TCF-7 is dispensable for anti-tumor response but required for persistent function of mature CD8 T cells

Rebecca Harris[1*], Mahinbanu Mammadli[1*], Shannon Hiner[1], Liye Suo[2], Qi Yang[3], Jyoti Misra Sen[4,5], Mobin Karimi[1ℜ]

[1]Department of Microbiology and Immunology, SUNY Upstate Medical University, Syracuse, NY 13210.

[2]Department of Pathology, SUNY Upstate Medical University, Syracuse, NY 13210.

[3]Department of Pediatrics, Rutgers Robert Wood Johnson Medical School
Rutgers Child Health Institute of New Jersey New Brunswick, NJ 08901

[4]National Institute on Aging-National Institutes of Health, 08C218, Biomedical Research Center, 251 Bayview Boulevard, Suite 100, Baltimore, MD 21224.

[5]Immunology Program, Department of Medicine, Johns Hopkins School of Medicine, Baltimore, MD 21224.

Rebecca Harris[1*] and Mahinbanu Mammadli[1*] equally contributed to this manuscript

To whom correspondence should be addressed ℜ:
Mobin Karimi
Assistant Professor of Immunology and Microbiology
SUNY Upstate Medical University,
766 Irving Ave Weiskotten Hall Suite 2281,
Syracuse, NY 13210
Office Phone: 315-464-2344
Laboratory Phone: 315-464-7652
Email: karimim@upstate.edu

Conflict of interest statement: The authors have declared that no conflict of interest exists

Abstract

T Cell Factor-1, encoded by *TCF-7,* is a transcription factor that plays an essential role

during T cell development and differentiation. In this manuscript we utilized a pre-clinical model

provided evidence that *TCF-7* is dispensable for the anti-tumor response, and that *TCF-7*

suppresses key transcriptional factors Eomes and T-bet and molecules responsible for peripheral

CD8 T cell cytolytic function. We discovered that *TCF-7* regulates NKG2D expression on naïve

and activated mouse CD8 T cells, and that peripheral CD8 T cells from *TCF-7* cKO utilize

NKG2D to clear tumor cells.

We also provide evidence that *TCF-7* regulates key signaling molecules, including LCK,

LAT, ITK, PLC-γ1, P65, ERKI/II, and JAK/STATs required for peripheral CD8 T cell persistent

function. Our data transcriptomic and protein data uncovered the mechanism of how *TCF-7*

impacting peripheral CD8 T cell inflammatory cytokine production, CD8 T cell activation, and

apoptosis. Our pre-clinical model showed that CD8 T cells from *TCF-7* cKO mice did not cause

GVHD, but effectively cleared primary tumor cells.

Introduction

64
T Cell Factor-1 (*TCF-7*), a major T cell developmental transcription factor, is involved in the

Wnt signaling pathway, and is critical for T cell development as well as activation (Escobar et

al., 2020; Ma et al., 2012). Dysfunction of the Wnt/β-catenin/*TCF-7* signaling pathway leads to

immune deficiency or autoimmunity (Shi et al., 2016). It is well known that *TCF-7* is involved in

the regulation of cell proliferation and survival during later T cell development (Kim et al.,

2020). In the absence of *TCF-7*, T cell development is completely blocked after early thymocyte

progenitor (ETP) cells, which suggests that *TCF-7* is also important in T cell lineage

specification (Germar et al., 2011; Weber et al., 2011). Several studies have shown that *TCF-7* is

critical for controlling viral infection (He et al., 2016; Im et al., 2016; Utzschneider et al., 2016).

Furthermore, *TCF-7* + CD8+ T cell have self-renewal capacity, while CD8+ T cells lacking

TCF-7 do not (Utzschneider *et al.*, 2016; Wu et al., 2016). Overwhelming evidence has

suggested that *TCF-7* is critical for CD8 T cell persistence and capacity to control viral infection

(Kurtulus et al., 2019; Miller et al., 2019; Siddiqui et al., 2019; Wu *et al.*, 2016). While

investigating the role of *TCF-7* in viral infection, single cell RNA sequencing data uncovered

that CD8 T cells expressing higher levels of *TCF-7* are quiescent and tissue resident, while CD8

T cells expressing higher levels of *TCF-7* in periphery are highly proliferative, and that *TCF-7*

negative T-bet^{hi} CD8 T cells are transitory effector cells. *TCF-7$^-$* Eomes^{hi} CD8 T cells are not

proliferative and express higher levels of immune checkpoint receptors, but maintain effector

function (Beltra et al., 2020; LaFleur et al., 2019; Zander et al., 2019). Studies have also reported

that *TCF-7+* CD8 T cells with stem-like abilities expresses low levels of PD-1 and Tim

checkpoint receptors, further providing evidence that *TCF-7* is required for CD8 T cell formation

and persistent function (Kurtulus *et al.*, 2019; Siddiqui *et al.*, 2019; Utzschneider *et al.*, 2016).

Studies have demonstrated that dysfunctional virus-specific CD8 T cells transplanted into naïve

mice give rise to *TCF-7*-negative CD8 T cells (Utzschneider et al., 2013). However, tumor-

specific CD8 T cells transferred to naïve mice might give rise to *TCF-7+* CD8 T cells,

demonstrating the differential role of *TCF-7* in viral infection and tumor (Schietinger et al.,

2016). These differences could be due to the differences between microenvironments in viral

infection and in tumors (Philip et al., 2017). Whether *TCF-7+* cells will give rise to *TCF-7+* or

TCF-7- cells will be dependent on internal and external signaling. Some studies have shown that

both the long (p45) and the short (p33) *TCF-7* isoform are expressed by CD8 T cells that will

give rise to stem-like CD8 T cells during viral infection, and it has been shown that the long

isoform of *TCF-7* is capable of promoting stem-like CD8 T cell formation during viral infection

by regulating genes like CD127, CXCR5 and cMyb (Chen et al., 2019). Currently, the role of

long and short *TCF-7* isoforms is largely unknown.

Studies of T cells as immunotherapy in both human and mice showed that for a superior

anti-tumor response, less differentiated cells are more favorable than a more differentiated subset

of CD8 T cells (Gattinoni et al., 2011; Im *et al.*, 2016; Lugli et al., 2013). Ideal CD8 T cells for

immunotherapy have been shown to exhibit stem-like abilities (such as those obtained by

inducing *ex vivo* cell growth with IL-17, IL-15, IL-21) and higher expression of *TCF-7*, Eomes,

and Bcl6 (Cieri et al., 2013; Cui et al., 2011). Thus, the suitability of *TCF-7* as a target for

immunotherapy to clear viral infection and cancer might also indicate considerable consequences

for autoimmune diseases.

To study the role of *TCF-7* in a clinically relevant model, we utilized allogeneic

hematopoietic stem cell transplantation (allo-HSCT). In allo-HSCT, mature peripheral donor T

cells found in the graft become alloactivated upon recognition of host HLA as non-self. These T

cell activities help to clear residual malignant cells, which is called the graft-versus-leukemia

effect (GVL) (Balassa et al., 2019; Giralt and Bishop, 2009; Hall and Shenoy, 2019). On the

other hand, alloactivated T cells also target healthy recipient tissues, an effect known as graft-

versus-host disease (GVHD) (Mavers and Bertaina, 2018). We used a unique mouse strain which

has deletion of *TCF-7* in mature T cells, rather than a global deletion (Xing et al., 2016). This

TCF-7 flox/flox x CD4cre mouse experiences deletion of *TCF-7* in all T cells at the double-

positive phase of development, when all T cells express CD4 (Wang et al., 2019b). This allows

us to overcome the severe T cell developmental defect that occurs in global *TCF-7* deletion, as

TCF-7 is critical for the double-negative stage of development (Yang et al., 2019).

Using a mouse model of GVHD and GVL following allo-HSCT, we were able to study all

of the major T cell functions, as well as phenotype, clinical outcomes, and gene expression, in a

single model. In this model, we transplanted CD8 T cells from either WT or *TCF-7* cKO mice

into irradiated BALB/c mice (H2K^b→H2K^d) (Mammadli et al., 2021a; Mammadli et al., 2021c;

Mammadli et al., 2021d). Using allogenic pre-clinical model, we have shown that CD8 T cells

from *TCF-7* cKO effectively clear tumor cells without inducing GVHD by producing

significantly less inflammatory cytokines as proinflammatory cytokines are con sidered the

hallmark of allo-immunity (Ju et al., 2005; Seif et al., 2017). Our data also uncovered that CD8 T

cells from *TCF-7* cKO mice cause significantly less tissue damage in GVHD target organs

(Bleakley et al., 2012; Breems and Lowenberg, 2005). Molecular analysis showed that CD8 T

cells from *TCF-7* cKO mice significantly showed reduction in key molecules required for CD8 T

cell persistent function (Germar *et al.*, 2011; Giralt and Bishop, 2009; Gounari and Khazaie,

2022). CD8 T cells from *TCF-7* cKO mice exhibited innate memory-like phenotype by

upregulating CD122, CD44, and effector and central memory phenotypes in mature CD8 T cells

critical for GVHD development (Dutt et al., 2011; Huang et al., 2013). We also uncovered that

CD8 T cells from *TCF-7* cKO mice significantly upregulated Eomes and T-bet, two downstream

transcription factors which are known to be involved in GVL (Mammadli *et al.*, 2021a; Weeks et

al., 2021; Zhou et al., 2010). Our data demonstrated that naïve CD8 T cells from *TCF-7* cKO

mice upregulated NK cell type 2 receptor (NKG2D). NKG2D, encoded by *Klrk1*, is an activating

cell surface receptor that is predominantly expressed on Natural killer cells (Larson et al., 2006;

Wensveen et al., 2018). While naïve human CD8$^+$ T cells express NKG2D, in mice CD8 T cells

only upregulate NKG2D upon activation (Hu et al., 2016; Maasho et al., 2005). Upregulation of

Granzyme B on CD8 T cells from *TCF-7* cKO mice was also observed. The loss of *TCF-7* also

led to upregulation of NK cell type 2 receptor (NKG2D). NKG2D, encoded by *Klrk1*, is an

activating cell surface receptor that is predominantly expressed on Natural killer cells (Larson *et*

al., 2006; Wensveen *et al.*, 2018). NKG2D is abundantly present on all NK cells, NKT cells, and

subsets of γδ T cells (Stojanovic et al., 2018). While naïve human CD8$^+$ T cells express NKG2D,

in mice CD8 T cells only upregulate NKG2D upon activation (Hu *et al.*, 2016; Maasho *et al.*,

2005). Upregulation of Granzyme B on CD8 T cells from *TCF-7* cKO mice was also observed.

Our molecular and animal data were confirmed by transcriptomic analysis.

Altogether, our data demonstrate that loss of *TCF-7* in mature murine CD8 T cells

enhanced Eomes and T-bet expression and reduced TCR-signaling, resulting in less severe

GVHD. Our data demonstrated that *TCF-7*-deficient CD8 T cells utilized NKG2D receptors to

kill tumor targets. These findings will have a considerable impact on developing strategies to

uncouple GVHD from GVL, and for developing therapeutic interventions for T cell-driven

autoimmune disorders.

Results

Loss of *TCF-7* in donor CD8 T cells reduced severity and persistence of GVHD symptoms,

increased survival from lethal GVHD, and retained anti-tumor capabilities for the GVL

effect. Most of the previous research on *TCF-7* utilized a global *TCF-7* knockout because the

primary focus was on *TCF-7's* role as a developmental factor (Gounari and Khazaie, 2022;

Weber *et al.*, 2011). However, we wanted to study the role of *TCF-7* in mature T cells. Global

loss of *TCF-7* results in minimal T cell production from the thymus, because *TCF-7* is critical for

DN stages of development (Johnson et al., 2018). Therefore, we obtained mice with a T cell-

specific knockout for *TCF-7* (Tcf7 flox/flox x CD4cre, called *TCF-7* cKO here (Xing *et al.*,

2016). This allowed us to study mature T cells that developed normally in the thymus, then lost

expression of *TCF-7* at the DP phase (Berga-Bolanos et al., 2015).

To determine whether *TCF-7* plays a role in mature alloactivated T cell regulation, which

is currently unknown, we used a mouse model of MHC-mismatched allo-HSCT leading to

GVHD and GVL. Briefly, BALB/c mice (MHC haplotype d) were lethally irradiated and

transplanted with wild-type (WT) bone marrow and C57Bl/6-background (MHC haplotype b)

donor CD8 T cells (Mammadli *et al.*, 2021c; Mammadli *et al.*, 2021d). The donor CD8 T cells

came from wild-type (WT), or *TCF-7*-deficient (*TCF-7* cKO) mice. Recipients were given

$1X10^6$ CD8 T cells and $10X10^6$ WT T cell-depleted bone marrow cells, as well as $2X10^5$

luciferase-expressing B-cell lymphoma (A-20) cells (Edinger et al., 2003a; Edinger et al., 2003b)

to assess GVL responses (Mammadli *et al.*, 2021a; Mammadli et al., 2021b; Mammadli *et al.*,

2021c; Mammadli *et al.*, 2021d). A20 cells are syngeneic to BALB/c mice and allogeneic to

C57BL/6 (B6) mice (Edinger *et al.*, 2003a; Edinger *et al.*, 2003b), meaning that the cells will not

be naturally rejected by the BALB/c hosts, but will be attacked by the transplanted donor T cells.

The MHC haplotype mismatch between host and donor cells drives alloactivation of donor T

cells, which in turn causes GVHD and GVL effects (Hoffmann et al., 2002). To examine disease

severity, progression, and recipient mouse survival, recipient mice were weighed and given a

clinical score three times per week following transplant, until about day 70 **(Fig. 1A-D)**. The

mice were scored based on six factors: skin integrity, fur texture, posture, activity level, weight

loss, and diarrhea (Cooke et al., 1996). Since the A20 cells express luciferase (called A20 luc)

(Mammadli *et al.*, 2021c), allowing us to track them by injecting D-luciferin into the recipient

mice and imaging them with an *in vivo* bioluminescence scanner (IVIS 50), the mice were

scanned one time per week with IVIS 50 until the end of the experiment **(Fig.1A, 1E).**

We found that mice who received WT donor CD8 T cells had a rapid increase in GVHD

severity, with a high score being reached by day 14, suggesting severe GVHD **(Fig. 1B)**. This

high score was maintained, suggesting persistent disease, and reached a high peak score at day

40 when the recipient mice died of disease burden **(Fig. 1B-D)**. WT-transplanted mice lost

weight initially and were unable to regain much weight (**Fig.1C**). In contrast, mice given *TCF-7*

cKO CD8 T cells had much better survival, lower peak and average clinical scores, minimal

disease burden, and a gain in weight following the initial weight loss period **(Fig.1B-D).** In

addition, the clinical scores for *TCF-7* cKO CD8 T cells transplanted mice quickly reduced to

near-control levels following peak score at day 10 **(Fig. 1B)**, suggesting that disease does not

persist in these mice. Therefore, loss of *TCF-7* in donor T cells led to reduced GVHD severity

and persistence, with improved survival **(Fig.1D).**

Regarding anti-tumor effects, we observed that over time, the group receiving only bone

marrow and tumor cells showed a large increase in tumor growth **(Fig. 1A, 1E)**, because no T

cells were present to control the tumor cells. In contrast, most mice given CD8 T cells from any

donor type along with the BM and A20 luc cells were able to clear the tumor cells by the end of

the experiment **(Fig. 1A, 1E)**. The GVL effect was maintained even in *TCF-7* cKO-transplanted

mice **(Fig. 1A, 1E)**. Altogether, these data show that *TCF-7* is dispensable for GVL effects, but

critical for GVHD. Therefore, loss of *TCF-7* in donor T cells provides a clinically optimal

phenotype, where GVHD severity is reduced but beneficial GVL effects are maintained.

Loss of *TCF-7* drives changes to mature CD8 T cell phenotype which are primarily cell-

extrinsic. It has been shown that loss of *TCF-7* in late stages of T cell development led to

impaired output of CD4 T cells, and redirection of CD4 T cells to a CD8 T cell fate(Steinke et

al., 2014). To determine whether loss of *TCF-7* affected mature donor T cell phenotype, we

performed flow cytometry phenotyping on CD8 T cells **(Fig. 2)**. First, we confirmed the loss of

TCF-7 expression in *TCF-7* cKO mice by flow cytometry **(Fig.2A).** Next, we examined whether

the loss of *TCF-7* also altered Eomesodermin (Eomes) and T-box transcription factor 21 (T-bet)

expression, both of which are downstream of *TCF-7* (Chen *et al.*, 2019). Some reports have

claimed that Eomes is activated by *TCF-7* (meaning loss of *TCF-7* reduces Eomes expression)

(Paley and Wherry, 2010). However, in our model of conditional *TCF-7* deletion, we found that

TCF-7 cKO CD8 T cells had increased expression of Eomes compared to WT CD8 T cells **(Fig.**

2B). Other reports have claimed that T-bet may be activated or not affected by *TCF-7* (Ma *et al.*,

2012), but we found that loss of *TCF-7* led to increased T-bet expression in CD8 T cells **(Fig.**

2C). This suggests that *TCF-7* normally suppresses the expression of Eomes and T-bet in mature

CD8 T cells.

Some reports have suggested that CD44[hi] T cells do not cause GVHD or cause less severe

GVHD (Dutt *et al.*, 2011). Therefore, we wanted to examine CD8 T cells from *TCF-7* cKO mice

for activation markers like CD44 and CD122. Our data showed that CD8 T cells from *TCF-7*

cKO mice exhibit increased expression of CD122 and CD44 **(Fig. 2D-E)**. Next, using CD62L

and CD44 markers, we identified three major memory subsets: central memory (CD44^{hi}

CD62L^{hi}), effector memory (CD44^{hi} CD62L^{low}), and naive (CD44^{low} CD62L^{hi}) cells **(Fig.2F)**.

TCF-7 cKO mice showed increased central memory CD8 T cell subsets and decreased naive

CD8 T cells **(Fig. 2F)**. Thus, loss of *TCF-7* results in a more memory-skewed phenotype for

CD8 T cells. Some reports have suggested that memory T cells delay induction of GVHD (Dutt

et al., 2011; Mammadli *et al.*, 2021c), so this phenotypic change may be beneficial (Nakajima et

al., 2021).

Changes to cell phenotype in a knock-out mouse may be cell-intrinsic (due directly to

gene deficiency within the cell) or cell-extrinsic (due to changes in the microenvironment from

gene deficiency) (Decman et al., 2010; Mammadli *et al.*, 2021d). To determine whether the

phenotypic effects we observed were cell-intrinsic or cell-extrinsic, we developed a chimeric

mouse model. Briefly, we mixed bone marrow from WT and *TCF-7* cKO mice at a 1:4

(WT:TCF) ratio for a total of 50X10^6 BM cells, then used this mixture to reconstitute lethally

irradiated Thy1.1 mice. We used a 1:4 ratio based on our previous published work (Mammadli et

al., 2020; Mammadli *et al.*, 2021d), to ensure survival of the KO cells (based on our initial

observations that *TCF-7* cKO T cells did not proliferate well in culture). At 9 weeks post-

transplant, blood was taken to ensure reconstitution and survival of both donor types in each

mouse. At 10 weeks, splenocytes were phenotyped by flow cytometry, with donor cells being

identified by H2K^b, CD45.1 (WT), and CD45.2 (*TCF-7* cKO) markers (Mammadli *et al.*, 2020).

First, we looked at the *TCF-7* expression in CD45.1+ (WT), and CD45.2+ (*TCF-7* cKO)

cells and confirmed that cells from *TCF-7* cKO mice did not express *TCF-7* in chimeric mice

(Supp.Fig.1A). We did see a statistically significant increase in T-bet expression in CD8 T cells

from *TCF-7* cKO donor cells compared to WT donor cells in chimeric mice, when we performed

a t-test (data not shown). However, when we compared the T-bet expression in chimeric versus

naïve CD8 T cells, we observed that T-bet expression in CD8 T cells from *TCF-7* cKO donor

mice was reduced to near-WT levels from elevated levels **(Supp. Fig. 1B).** This suggests that the

increased expression of T-bet seen in *TCF-7* cKO CD8 T cells from naive mice is a cell-extrinsic

effect.   Interestingly, in the chimeric mice we observed that Eomes and CD122 expression levels

in WT CD8 T cells were significantly increased to near-*TCF-7* cKO levels, suggesting that the

increase in Eomes and CD122 expression in CD8 T cells from *TCF-7* cKO mice is primarily

cell-intrinsic **(Supp.Fig.1C-D).**

Next, we examined the expression of CD44 and central memory phenotype in chimeric

mice. We observed that while the frequencies of these subsets were lower in *TCF-7* cKO-derived

CD8 T cells compared to WT-derived CD8 T cells in the chimera (opposite of the trend observed

in naive mice), this was because the frequencies of CD44 and CM phenotype in WT cells was

enhanced to the levels expressed by *TCF-7* cKO cells from naive mice **(Supp. Fig. 1E-F).**

These results suggest that the effects of *TCF-7* deficiency on CD44 and CM phenotype

expression in naïve mice could be primarily cell-intrinsic, with cell-extrinsic elements as well.

Interestingly, effector memory phenotype in the chimeric mice we observed that levels in WT

CD8 T cells were significantly increased to near-*TCF-7* cKO levels, suggesting that the increase

in effector memory phenotype in CD8 T cells from *TCF-7* cKO mice is primarily cell-intrinsic

(Supp. Fig. 1G). Finally, the naïve CD8 T cell population in the chimera coming from *TCF-7*

cKO bone marrow was significantly increased compared to CD8 T cells from WT bone marrow

and compared to naïve *TCF-7* cKO mice (**Supp. Fig. 1H**). This suggests that the effect on naive

CD8 T cells in *TCF-7* cKO mice could be either cell-intrinsic or cell-extrinsic. Altogether, these

data suggest that the phenotypic changes seen in *TCF-7* cKO may be primarily cell-intrinsic,

with some additional cell-extrinsic effects being present.

Loss of *TCF-7* alters cytotoxic mediator production in mature CD8 T cells. Our data

demonstrated that the loss of *TCF-7* increases Eomes and T-bet expression in mature CD8 T

cells (**Fig. 2B-C**). Considering that Eomes and T-bet have been reported to play a central role in

anti-tumor responses, we hypothesized that by upregulating Eomes and T-bet, CD8 T cells

lacking *TCF-7* can maintain cytotoxicity, and that *TCF-7* is not required for CD8 T cell-

mediated cytolytic function (Zhu et al., 2010). We anticipated that CD8 T cells from *TCF-7* cKO

mice may have attenuated TCR signaling, so we examined this and other activating receptors by

flow cytometry. It is also known that Eomes and T-bet overexpression increases NKG2D

expression in NK cells (Kiekens et al., 2021). Considering that loss of *TCF-7* in mature T cells

led to upregulation of Eomes and T-bet expression, we hypothesized that loss of *TCF-7* may also

lead to upregulation of NKG2D expression in CD8 T cells and enhance the anti-tumor response.

Natural killer group 2 member D (NKG2D) is constitutively expressed on mouse NK

cells, NKT cells and some other cells (Abel et al., 2018; Al Dulaimi et al., 2018), but does not

get expressed on naïve mouse CD8 T cells (Prajapati et al., 2018). Human CD8 T cells always

express NKG2D on their surface, but mouse CD8 T cells only express it upon activation

(Wensveen *et al.*, 2018) NKG2D is activated by NKG2D ligands (Raulet et al., 2013), and

NKG2D ligands are relatively restricted to malignant or transformed cells (Raulet, 2003; Raulet

et al., 2013). In order to determine whether loss of *TCF-7* affects NKG2D expression and anti-

tumor responses, we analyzed NKG2D expression in CD8 T cells. We measured NKG2D

expression by flow cytometry before and at different time points after CD3/CD28 activation

(Karimi et al., 2015). We found that CD8 T cells from *TCF-7* cKO mice had significantly

increased expression of NKG2D on the cells surface compared to CD8 T cells from WT mice,

before stimulation **(Fig.3A).** Next, we wanted to examine whether NKG2D expression was

further upregulated on CD8 T cells from *TCF-7* cKO mice compared to CD8 T cells from WT

mice after stimulation. CD8 T cells were cultured with 2.5ug/ml anti-CD3 and 2.5ug/ml anti-

CD28 antibodies for 24, 48, or 72 hours. These cultured cells were examined for NKG2D

expression by flow cytometry. We observed an increase in NKG2D expression on CD8 T cells

from both WT and *TCF-7* cKO mice in a time-dependent manner, and at all time points,

expression of NKG2D was higher for cells from *TCF-7* cKO mice **(Fig.3B).** There was no

difference in the viability of the cells or CD8 T cell numbers before or after the culture

(Supp.Fig.2A-B).

We also wanted to compare the Granzyme B expression in CD8+, NKG2D+ T cells from

TCF-7 cKO and WT mice (Chu et al., 2013). We did not observe any Granzyme B expression in

CD8 T cells before stimulation **(Fig.3C).** Only 24 hours after stimulation, we observed

Granzyme B expression in T cells from both strains, peaking at 48 hours post-stimulation with

no difference between strains of mice **(Fig.3C)**. After 72 hours post-stimulation, CD8 T cells

from WT mice had significantly reduced Granzyme B expression compared to *TCF-7* cKO CD8

T cells **(Fig.3C)**. We also confirmed total Granzyme B expression in CD8 T cells from *TCF-7*

cKO mice, in the presence and absence of CD3/CD28 stimulation, using Western blotting. Total

Granzyme B expression was upregulated in CD8 T cells from *TCF-7* cKO mice compared to WT

mice **(Fig.3D-E).** These data demonstrated that CD8 T cells from *TCF-7* cKO mice may

maintain anti-tumor responses by killing the target cells with an NKG2D-mediated mechanism,

and by persistent upregulation of Granzyme B expression (Liu et al., 2022; Wang et al., 2022).

Next, we wanted to examine the functional consequences of upregulation of NKG2D

expression on CD8 T cells from *TCF-7* cKO mice. We utilized an in *vitro* cytotoxicity assay,

where we used anti-NKG2D neutralizing antibody. We isolated CD8 T cells from WT and *TCF-*

7 cKO mice and cultured them for 48 hours with anti-CD3/anti-CD28 antibodies in order to

induce optimal NKG2D expression in CD8 T cells. CD8 T cells from *TCF-7* cKO and WT mice

were then cultured with tumor target A20 cells (Edinger *et al.*, 2003b) in a 40:1 ratio of tumor

cells to CD8 T cells, along with anti-NKG2D antibody or isotype control antibody for 4 hours.

We used the A20 cell line as a tumor target because it is known for expressing NKG2D ligands

including Rae1, H60, and MULT1(Karimi *et al.*, 2015; Nishimura et al., 2008). Triplicate wells

were averaged and percent lysis was calculated from the data using the following equation: %

specific lysis = 100 × (spontaneous death bioluminescence – test bioluminescence)/ (spontaneous

death bioluminescence – maximal killing bioluminescence) (Karimi et al., 2014).

Our data showed that the addition of anti-NKG2D antibody significantly reduced the

cytotoxicity of CD8 T cells from *TCF-7* cKO mice, whereas addition of isotype control had no

effect on cytotoxicity of the CD8 T cells from *TCF-7* cKO mice **(Fig.3F).** In contrast, the

addition of anti-NKG2D antibody (Karimi *et al.*, 2015) did not change cytotoxicity of the CD8 T

cells from WT mice **(Fig.3F).** These data further support the idea that *TCF-7* cKO CD8 T cells

maintain their anti-tumor activity through an NKG2D-mediated mechanism. Taking into account

that normal tissue does not express NKG2D ligands on the surface and that primarily malignant

and transformed cells upregulate these ligands, this could explain why CD8 T cells from *TCF-7*

cKO mice cause less GVHD but maintain their anti-tumor activity (Nishimura *et al.*, 2008).

Loss of *TCF-7* alters cytokine production, chemokine expression, and expression of exhaustion markers by mature CD8 T cells. We confirmed that CD8 T cells from *TCF-7* cKO mice mediate cytolytic function primarily through NKG2D. Next, we wanted to examine the mechanism behind why CD8 T cells from *TCF-7* cKO mice induce less GVHD. One of the hallmarks of GVHD is the release of pro-inflammatory cytokines by alloactivated donor T cells, eventually leading to cytokine storm (Lynch Kelly et al., 2015; Mohty et al., 2005). We examined whether loss of *TCF-7* in donor CD8 T cells led to changes in cytokine production, thereby affecting GVHD damage. We allotransplanted lethally irradiated BALB/c mice as described above. Recipient mice were transplanted with 1.5×10^6 WT or *TCF-7* cKO CD8 donor T cells. Recipients were sacrificed at day 7 post-transplant. Splenocytes were isolated and restimulated by 6 hours of culture with PBS (control) or anti-CD3/anti-CD28 (stimulation), along with Golgiplug. Afterwards, the cultured cells were stained with antibodies against H2K^b, CD3, CD4, CD8, TNF-α, and IFN-γ. Our data showed that production of TNF-α by donor CD8 T cells trended toward decreasing when *TCF-7* was lost **(Supp. Fig. 3A)**. In contrast, IFN-γ trended toward increasing upon loss of *TCF-7* in CD8 T cells **(Supp. Fig. 3B)**.

We also obtained serum from cardiac blood of recipient mice at day 7 post-transplant and tested it with a mouse Th cytokine ELISA panel (LEGENDplex kit from Biolegend) (Mammadli *et al.*, 2020; Mammadli *et al.*, 2021d). Levels of TNF-α and IFN-γ in serum of recipient mice given *TCF-7* cKO CD8 T cells were lower than in mice given WT CD8 T cells at day 7 **(Fig. 4A)**. In contrast, the serum levels of IL-2 in mice given *TCF-7* cKO CD8 T cells was higher than in mice given WT CD8 T cells at day 7 **(Fig. 4A)**. At day 14 post-transplant, the reduction in TNF-α and IFN-γ levels observed at day 7 for *TCF-7* cKO-transplanted mice was preserved **(Fig. 4B)**. We observed a trend towards decreased serum levels of IL-2 in mice given *TCF-7* cKO

CD8 T cells compared with mice given WT CD8 T cells at day 14 post-transplant, opposite of

the effect observed on day 7 post-transplant **(Fig.4B)**. This suggests that *TCF-7* cKO CD8 T

cells may be capable of IFN-γ and TNF-α production at the same level as WT CD8 T cells when

restimulated, but in reality, produce less IFN-γ and TNF-α than WT cells when allotransplanted.

In order for GVHD to persist, donor T cells must proliferate in secondary lymphoid

organs and target organs (Beilhack et al., 2005; Ferrara, 2014). Naive and effector T cells drive

GVHD, but they are short-lived and must be replaced to maintain an alloresponse (Jiang et al.,

2021; Jiang et al., 2014). Also, given that memory cells are increased among CD8 T cells when

TCF-7 is lost, we hypothesized that activation and/or exhaustion of these cells may also be

affected. Ki-67 (Blessin et al., 2021) is a marker of T cell activation and proliferation, and TOX

(Khan et al., 2019) and PD-1 are markers of exhaustion (Ahn et al., 2018). Therefore, we wanted

to determine the Ki-67, TOX, and PD-1 expression levels on WT and *TCF-7* cKO CD8 T cells *in*

vitro. We cultured splenocytes from either WT mice or *TCF-7* cKO mice *in vitro* with anti-CD3

and anti-CD28 antibodies for 24, 48, and 72 hours. We did not observe any difference in Ki-67

expression in cells that were not stimulated, but the CD8 T cells from *TCF-7* cKO mice that were

stimulated for 72 hours *in vitro* showed significant upregulation of Ki-67 expression (**Fig. 4 C-**

D), suggesting that CD8 T cells from *TCF-7* cKO mice could potentially proliferate more than

CD8 T cells from WT mice when restimulated**.** We also observed the same trend of increased

expression for PD-1 at 72 hours post-stimulation **(Fig.4E-F)**. There were no differences in

expression of TOX at any time points *in vitro* when CD8 T cells from *TCF-7* cKO mice were

compared to WT **(Supp.Fig.3C).**

Next, we checked the expression of these markers *in vivo* from donor cells that were allo-

transplanted in recipient Balb/c mice as described earlier. At day 7 post-transplant, splenocytes

were isolated and Ki-67, TOX, and PD-1 expression were detected by flow cytometry. We

observed that donor CD8+T cells from *TCF-7* cKO mice expressed more TOX compared to WT,

suggesting that donor T cells *TCF-7* cKO mice were more exhausted following

allotransplantation **(Fig.5G-H)**. We did not observe any statistically significant differences in Ki-

67 or PD-1 expression at day 7 post-transplant **(Supp.Fig.3D-E).** Taken together these data

suggest that CD8 T cells from *TCF-7* cKO mice could be more exhausted than WT CD8 T cells

both *in vivo* and *in vitro*.

One of the major functions of alloactivated T cells is migration from spleen to

GVHD target organs, including liver and small intestine(Beilhack *et al.*, 2005; Ferrara, 2014).

Expression of chemokines and chemokine receptors is a critical aspect of T cell migration to

target organs. To determine whether expression of these molecules was affected by loss of *TCF-*

7 in CD8 T cells, we FACS sorted pre- and post-transplanted donor CD8 T cells from WT or

TCF-7 cKO mice (spleen only for pre-transplant, spleen, or liver for post-transplant). We then

extracted RNA from the cells, converted it to cDNA, and performed qPCR using a 96-well

mouse chemokine/chemokine receptor plate (Thermo Fisher). As expected, expression of these

markers was generally upregulated in alloactivated cells. Expression of these markers was

generally higher in *TCF-7* cKO CD8 T cells from pre-transplant spleen compared to WT pre-

transplant spleen **(Supp. Fig. 4A)**, but these markers were generally downregulated in *TCF-7*

cKO CD8 T cells from post-transplant liver and spleen compared to WT cells **(Supp. Fig. 4B-**

C). Therefore, *TCF-7* controls expression of CD8 T cell chemokine/chemokine receptors.

Loss of *TCF-7* in donor CD8 T cells led to decreased damage to the GVHD target organs.

During GVHD, host tissues are damaged by the activity of alloactivated T cells. To determine

whether damage to target organs of GVHD (skin, liver, and small intestine) was altered by loss

of *TCF-7* in donor T cells, we collected organs from mice allotransplanted as described above

(Beilhack *et al.*, 2005; Mammadli *et al.*, 2021a; Mammadli *et al.*, 2021d; Weeks *et al.*, 2021). At

day 7 and day 21 post-transplant, we collected pieces of skin, small intestine, and liver from

recipient BALB/c mice. These organs were fixed, sectioned, stained with hematoxylin and eosin

(H&E), and analyzed by a pathologist (L.S.) **(Fig. 5)**. At day 7, *TCF-7* cKO mice showed

significantly less inflammatory infiltrates in all the organs. We observed much less inflammatory

infiltrates in the bile duct epithelium of the portal triad (black arrows showing the interlobular

bile ducts) in the liver of the *TCF-7* cKO-transplanted recipients compared with WT-transplanted

recipients **(Fig. 5A)**. In the small intestines, no apoptotic bodies were seen in the crypts of the

small intestine in the *TCF-7* cKO CD8 T cell-transplanted mice, while frequent apoptotic bodies

were present in the WT CD8 T cell-transplanted mice at day 7 post-transplantation (black

arrows) **(Fig. 5B)**. In the skin, a mild increase in inflammatory cells was observed in the dermis

of the WT CD8 T transplanted mice, while the dermis of the *TCF-7* cKO CD8 T transplanted

mice appears normal at day 7 post-transplantation **(Fig. 5C)**.

Again, at day 21 post-transplant, *TCF-7* cKO CD8 T cell transplanted mice showed

significant less inflammatory infiltrates in all the sectioned GVHD target organs. We observed

much less inflammatory infiltrates involving the bile duct epithelium of the portal triad (black

arrows showing the interlobular bile ducts) in the liver of the *TCF-7* cKO CD8 T cell-

transplanted mice compared with WT CD8 T cell-transplanted mice **(Fig. 5D)**. At day 21 post-

transplant, no apoptotic bodies were seen in the crypts of the small intestines of the *TCF-7* cKO

CD8 T cell transplanted mice, while few apoptotic bodies are present in the small intestines of

the WT CD8 T cell transplanted mice (black arrows) **(Fig. 5E)**. A marked increase in

inflammatory cells (red circle) and destruction of the adnexal glands was observed in the dermis

of the WT CD8 T cell-transplanted recipients, while the dermis of the *TCF-7* cKO CD8 T cell

transplanted recipients showed only a mild increase in dermal inflammatory cells, with

preservation of adnexal glands **(Fig. 5F)**. Altogether, these data suggest that *TCF-7* normally

promotes GVHD damage to healthy tissues and is indispensable for T cell-driven damage. Thus,

loss of *TCF-7* in donor T cells leads to reduced severity and persistence of GVHD over time.

***TCF-7* alters the transcriptomic signature of alloactivated T cells.** Given that the phenotype

and functions of donor T cells, as well as disease outcomes, were significantly altered by loss of

TCF-7 on donor cells, we sought to determine what specific gene changes occurred to support

this. We allotransplanted recipient BALB/c mice with $1X10^6$ donor CD3 T cells as above, and

FACS-sorted pre- and post-transplant WT or *TCF-7* cKO donor CD8 T cells, which were stored

in Trizol and transcriptionally profiled. When we analyzed the genetic profile of the pre-

transplanted CD8 T cells from *TCF-7* cKO and WT mice, we were not able to determine any

differentially expressed genes (DEGs, FDR<0.1). However, when we performed Gene Set

Enrichment analysis (GSEA) using the Hallmark pathways collection from Molecular Signatures

Database (MSigDB), we identified that cytokine signaling pathways like TNF-a Signaling via

NF-κβ and Interferon gamma response were enriched in pre-transplanted CD8 T cells from WT

mice compared with cells from *TCF-7* cKO mice **(Supp.Fig.5A-C)**. Meanwhile, a number of

pathways involved in cell cycle also were enriched in WT cells versus *TCF-7* cKO pre-

transplanted CD8 T cells, like the P53 pathway, G2M checkpoint, DNA repair, and Myc targets

pathways **(Supp.Fig.5A).** MTOR signaling, Allograft rejection, and Oxidative phosphorylation

pathways were also enriched in pre-transplant CD8 T cells from WT mice **(Supp.Fig.5A,6D-F),**

suggesting that loss of *TCF-7* alters the transcriptional profile of CD8 T cell towards decreased

cytokine release while also altering the cell cycle, leading to a lessening of the alloactivation

responses.

When we analyzed the post-transplanted CD8 T cells which were alloactivated *in vivo*, we

identified 2548 differentially expressed genes (DEGs; FDR<0.1) when comparing *TCF-7* cKO

cells to WT cells **(Fig. 6A-B).** A majority of the DEGs (2000 genes) were downregulated

(module 2 in heatmap) and only 548 genes (module 1 in heatmap) were upregulated in post-

transplant CD8 T cells from *TCF-7* cKO mice compared to WT mice **(Fig. 6A-B).** We analyzed

both up- and downregulated DEGs for the Gene Ontology (GO) enrichment analysis using

Functional Annotation Chart tools, selecting only the top 20 GO BP (Biological Process) terms

in the Database for Visualization and Integrative Discovery (DAVID)(Huang da et al., 2009;

Sherman et al., 2022). The analysis showed that the differentially expressed genes in post-

transplant CD8 T cells from *TCF-7* cKO mice compared to WT were involved in Cell Cycle,

Cell-cell adhesion, Cell division, Apoptotic process, Antigen processing and presentation via

MHC class I, TCR signaling, Regulation of NF-κB signaling, Metabolic process, and others

(Supp.Fig.6). Once we knew which processes these DEGs played a role in, we pulled out the top

genes that were altered for each GO BP term that we were interested in. When we looked at the

top 35 DEGs based on P-value that were altered in Cell cycle, we observed that a majority of

them were downregulated in *TCF-7* cKO compared to WT **(Fig.6C).** We also looked at the top

30 genes based on P-value that were altered in Apoptotic processes **(Fig.6D**), which also showed

that most of the genes were downregulated in CD8 T cells in *TCF-7* cKO mice. When we looked

at the top 30 genes that play a role in metabolic processes, we observed that only 1 gene was

upregulated, and the rest were downregulated in *in vivo* alloactivated CD8 T cells from *TCF-7*

cKO mice **(Fig. 6E).** While the upregulated genes in the NF-κB pathway were Fasl, Ubd, Chuk

and others, the downregulated genes were Rela, Irf3, Traf2, Mavs, Tradd, Nod1, Tnfrsf1a, and

Trim25 **(Fig.6F)**.

Gene Set Enrichment analysis (GSEA) using the Hallmark pathways identified that

signaling pathways like PI3K-AKT-MTOR Signaling and TNFA Signaling via NF-κβ were

enriched in post-transplanted CD8 T cell from WT mice compared to cells from TCF-7 cKO

mice **(Supp.Fig.7A-B)**. GSEA analysis using the C2 canonical pathways showed that a number

of cytokines signaling pathways involving IL-1, IL-2, IL-4, and IL-7 were enriched in post-

transplanted CD8 T cells from WT mice compared to cells from *TCF-7* cKO mice **(Fig.6G)**.

While Cell cycle pathways were enriched in CD8 T cells from *TCF-7* cKO mice compared to

cells from WT mice, Apoptosis and Cell adhesion pathways were enriched in post-transplanted

CD8 T cells from WT mice **(Fig.6H).** Meanwhile, a number of cells signaling pathways were

also enriched in WT CD8 T cells compared to *TCF-7* cKO cells, such as TCR signaling, Toll

like receptor signaling, Jak-Stat signaling, ERK-MAPK signaling, and NF-kB pathways **(Fig.6I).**

We also analyzed the genes that were altered in KEGG pathways, which revealed that

DEGs that were altered in *in vivo* alloactivated CD8 T cells from *TCF-7* cKO mice (compared to

WT) were involved in pathways like Cell cycle, DNA replication, Metabolic pathways, Natural

killer mediated cytotoxicity, TCR signaling, JAK-STAT signaling, Chemokine receptor

signaling, and others **(Fig.7A).** Specifically, Klrk1 gene for NKG2D on Natural killer mediated

cytotoxicity pathway were enriched in allo-activated *TCF-7* deficient CD8+ T cells compared to

WT CD8+ T cells. Clustering of genes that were affected in TCR signaling showed that while

AKT1, AKT2, Pik3r5, Zap70, LCK, Lat, PLCγ1, Pdcd1, Vav1, Rela, Mapk3, Nfkbia, and Nfatc1

were downregulated, Ifng and Ptprc were upregulated in post-transplanted CD8 T cells from

TCF-7 cKO mice **(Fig.7B).** The JAK/STAT signaling pathway is important for cytokine

production and for the response of T cells to cytokines. Analysis revealed that IL2RB, JAK3,

STAT5B, STAT3, STAT1, Cish, Il2rb, Socs3, and Socs1 genes were downregulated in the JAK-

STAT pathway for *TCF-7* cKO CD8 T cells compared to WT **(Fig.7C)**.

In order to confirm these changes in genes, we isolated CD8 T cells from WT and *TCF-7*

cKO mice and determined the baseline level protein expression of LCK, ZAP70, LAT, ITK,

PLCγ1, ERK1-2, Jak2, Jak3, Stat3, p65-rela, AKT, and actin in unstimulated and 10 minute-anti-

CD3/CD28-stimulated samples. Data from non-stimulated samples revealed that protein

expression of most of the markers was downregulated in CD8 T cells from *TCF-7* cKO mice

compared to WT mice; only ZAP70, JAK3 and AKT were not affected by loss of *TCF-7*

(Fig.7D, Supp.Fig.8A). We observed even more robust differences in samples that were

stimulated with anti-CD3/CD28 for 10 minutes, and again only ZAP70 and AKT were

unaffected by loss of *TCF-7* **(Fig.7E, Supp.Fig.8B).** Altogether, the data from RNA sequencing

analysis and Western blots of stimulated and unstimulated samples showed attenuation of TCR

signaling and many other pathways in CD8 T cells from *TCF-7* cKO mice. These results help to

explain why CD8 T cells from *TCF-7* cKO mice cannot induce GVHD as severely as CD8 T

cells from WT mice.

518
519
520
521

Discussion

T Cell Factor-1 (*TCF-7*) is a critical regulatory transcription factor in T cell development and functions (Zuniga-Pflucker, 2004). *TCF-7* is known to be important for T cell development, as well as activation in some contexts (Yu et al., 2010). *TCF-7* has been extensively studied in viral infection(Escobar *et al.*, 2020; He *et al.*, 2016; Im *et al.*, 2016; Kurtulus *et al.*, 2019; Miller *et al.*, 2019; Siddiqui *et al.*, 2019; Utzschneider *et al.*, 2016; Weber *et al.*, 2011; Wu *et al.*, 2016). However, the role of (*TCF-7*) in a mouse model of allogeneic transplant has not been investigated. Using a murine allogeneic transplant model, we have shown that CD8 T cells from *TCF-7* cKO effectively clear tumor cells without inducing GVHD by producing significantly less inflammatory cytokines (Ju *et al.*, 2005; Seif *et al.*, 2017). Our data also uncovered that CD8 T cells from *TCF-7* cKO mice cause significantly less tissue damage in GVHD target organs (Bleakley *et al.*, 2012; Breems and Lowenberg, 2005). Here, we show that loss of *TCF-7* alters a number of CD8 T cell functions, and while it is dispensable for anti-tumor responses, it is essential for host tissue damage, cytokine production and signaling, and gene expression, playing a role in a number of immunological and biological pathways during alloactivation. (Yu *et al.*, 2010). This murine model allows us to study T cell function, clinical outcomes, and gene expression all in one model. Here, we showed that *TCF-7*-deficiency alters the phenotype of CD8 T cells by upregulating CD44 and CD122. We and other have shown that innate memory-like CD8 T cells expressing CD12hi and CD44 hi, Eomes and T-bet ameliorate GVHD development (Huang et al., 2019; Karimi *et al.*, 2014; Mammadli *et al.*, 2020; Mammadli *et al.*, 2021d; Zheng et al., 2009). The role of central (CD44^{hi}, CD62L^{hi}) and effector (CD44^{hi}, CD62L^{low}) memory phenotypes has been investigated previously (Huang *et al.*, 2019; Zheng *et al.*, 2009). Our data demonstrated that *TCF-7* significantly impacts CD8 T cell central memory phenotypes, suggesting that *TCF-7* might be a regulator for T cell activation. While effector and

naive cells are known to cause severe GVHD, central memory cells are often associated with less

severe disease (Dutt *et al.*, 2011; Tugues et al., 2018; Zheng *et al.*, 2009). Our findings

suggesting that this phenotypic change in *TCF-7 cKO* cells may be beneficial for reducing

disease severity. Our experiments in mixed bone marrow chimeras showed that bone marrow-

derived CD8 T cells from WT and or *TCF-7* cKO mice developed in the same thymus have a

similar phenotype to each other, and a different phenotype than CD8 T cells from naïve WT or

TCF-7 cKO mice. We found that the upregulation of activation marker expression like Eomes,

CD122, and the effector memory phenotype is primarily cell-intrinsic, with changes to other

markers being either cell-extrinsic or primarily cell-intrinsic with other extrinsic effects.

Donor T cells are crucial for target organ injury in graft-versus-host disease (GVHD).

These alloactivated T cells proliferate, migrate to target organs (liver, skin, and small intestine),

and produce cytokines during GVHD(Bastien et al., 2012; Reddy and Ferrara, 2008; Villarroel et

al., 2014). Our data showed that tissue damage in liver, skin, and small intestine (all target

organs) was reduced by loss of *TCF-7* in donor CD8 T cells at all timepoints. This shows that

TCF-7 in donor T cells is required for GVHD damage and persistence over time.

Donor T cells eliminate tumor cells (GVL) but also cause graft-versus-host disease

(GVHD) (Bleakley *et al.*, 2012; Tugues *et al.*, 2018). Our data showed that CD8 T cells from

TCF-7 cKO mice were able to clear tumor without causing GVHD, suggesting that *TCF-7* is

dispensable for anti-tumor responses. Our data revealed that CD8 T cells from *TCF-7* cKO mice

mediate cytolytic function via NKG2D. We also confirmed this hypothesis by blocking the

surface NKG2D by anti-NKG2D antibody on CD8 T cells from *TCF-7* cKO and WT mice and

performing an *in vitro* cytotoxicity assay. While the anti-tumor response of TCF1-deficient CD8

T cell against A20 cells (which express the NKG2D ligands like Rae1, H60, and MULT1)

(Nishimura *et al.*, 2008) was diminished, cytotoxicity of WT CD8 T cells was not affected.

Upregulation of the *KLRK1* gene (also known as NKG2D) in alloactivated *TCF-7* deficient

CD8+ donor T cells further confirmed our hypothesis. Furthermore, the increase in Granzyme B

expression in *TCF-7* cKO CD8 T cells by flow cytometry and Western blot also provides

evidence as to why the anti-tumor response is preserved despite weakened TCR signaling in

TCF-7 cKO CD8 T cells (Presotto et al., 2017).

Once we had a clear mechanism for the GVL effect, we looked at functions that were

altered in CD8 T cells from *TCF-7* cKO mice that could produce the attenuated GVHD effect.

We discovered that the serum levels of cytokines like TNFα and IFNγ were decreased in *TCF-7*

deficient CD8 T cell-transplanted mice. Published data has shown the CD8 T cell lacking *TCF-7*

develop exhaustion while clearing viral infection(Gautam et al., 2019; Seo et al., 2019) While the

expression of TOX, an exhaustion marker, was not affected in freshly isolated or *in vitro*-

stimulated *TCF-7* deficient CD8 T cells, alloactivated *TCF-7*-deficient CD8 donor T cells

upregulated TOX at day 7 post-transplant (Scott et al., 2019). Another exhaustion marker, PD-1,

was upregulated after 72 hours in *in vitro*-stimulated CD8 T cells from TCF-7 cKO mice, but *in*

vivo expression of PD-1 in donor T cells was not increased (Wang et al., 2019a; Xu et al., 2019).

This could be explained by differences between alloactivation *in vivo* and TCR-mediated

activation *in vitro.* Ki-67, a proliferation marker, was also altered by loss of *TCF-7*, suggesting

that CD8 T cells from *TCF-7* cKO mice may proliferate more compared to WT CD8 T cells

(Sobecki et al., 2016). We also confirmed that CD8 T cells from *TCF-7* cKO mice cause less

tissue damage to the target organs, and loss of *TCF-7* alters chemokine receptor expression both

pre- and post-transplant, which could explain why these cells cause less severe GVHD with

increased survival of recipient mice (Ferrara, 2014).

These studies provide evidence for how *TCF-7* regulates the functions of peripheral T

cells. The phenotype caused by loss of *TCF-7* is clinically optimal, because it allows for

clearance of residual malignant cells while limiting the risk of life-threatening GVHD damage

(Guinan et al., 1999). This observation, coupled with the increase in exhaustion of *TCF-7* cKO

donor CD8 T cells, suggests that donor cells lacking *TCF-7* are highly activated and cytotoxic to

malignant cells early on following transplant, but quickly become exhausted, limiting GVHD

progression.

Finally, to identify the changes to the genetic program of donor T cells that occurred in the

absence of *TCF-7*, we performed RNA sequencing analysis on pre- and post-transplant donor T

cells. Even though we could not identify the Differentially Expressed Genes (DEGs) in pre-

transplanted T cells (which we attribute to technical problems), gene set enrichment analysis

revealed that loss of *TCF-7* alters the genetic signature of pre-transplanted CD8 T cells. A

number of signaling pathways involved in cytokine production and cell cycle were enriched in

pre-transplanted CD8 T cell from WT mice compared to cells from *TCF-7* cKO mice, suggesting

that loss of *TCF-7* alters the transcriptional profile of the CD8 T cell towards decreased cytokine

release, while altering the cell cycle in baseline and leading to the lessening of the allo-activation

response. Meanwhile, 2548 DEGs (FDR<0.1) were identified when comparing post-transplanted

CD8 T cells from *TCF-7* cKO mice and WT mice. Both GO Annotation analysis of DEGs and

Gene Set Enrichment analysis revealed that a number of pathways such as Cell cycle, DNA

replication, Metabolic pathways, TCR signaling, JAK-STAT signaling, and Chemokine receptor

signaling were enriched in post-transplanted CD8 T cells from WT mice compared with cells

from *TCF-7* cKO mice. Transcriptomic analysis also revealed that KLRK-1 gene for NKG2D

were upregulated in allo-activated *TCF-7* cKO CD8+ T cells compared to the WT CD8+T cells,

which confirming our findings in in-vitro cytotoxicity assay and flow cytometry data. These

findings suggest that loss of *TCF-7* leads to changes in the transcriptomic profile of the CD8 T

cells towards producing less cytokines and attenuating the T cell response, while increasing

cytotoxicity. Chemokine receptor pathways were also enriched in alloactivated WT CD8 T cells

which confirmed the qPCR analysis of chemokine receptors. The decrease in chemokine

receptors helps to explain why we observed less tissue damage in GVHD target organs after allo-

transplantation, and attenuated GVHD persistence, in mice given *TCF-7* cKO CD8 T cells. By

using Western blotting, we also confirmed changes in TCR and JAK/STAT signaling before and

after stimulation with anti-CD3/CD28, which helps to explain why we saw less serum cytokines

at day 7 and day 14 post-transplant in mice given *TCF-7* cKO CD8 T cells. This also helps to

explain why GVHD didn't persist in recipients given CD8 T cells which lack *TCF-7*.

Altogether, these data suggest that *TCF-7* is a major transcription factor that plays a role

in T cell development. Our work shows that *TCF-7* is dispensable for cytotoxic function of

mature alloactivated CD8 T cells but is indispensable for GVHD. TCR, JAK-STAT, and NF-

κB signaling as well as cytokine production, these findings will help to establish an

understanding of *TCF-7* as a critical factor in the GVHD/GVL regulatory network of CD8 T

cells.

Materials and Methods

Mouse Models. For transplant, the following female donor mice were used: B6-Ly5.1

(CD45.1+, "WT" or B6.SJL-Ptprca Pepcb/BoyCrl, 494 from Charles River), C57Bl/6J

(CD45.2+, "WT", 000664 from Jackson Laboratories), or Tcf7 flox x CD4cre (referred to here as

" *TCF-7 cKO*" (Ma *et al.*, 2012) , obtained from Dr. Jyoti Misra Sen at NIH by permission of Dr.

Howard Xue, and bred in-house),. These donor mice were age-matched to each other and to

recipients as closely as possible. BALB/c female mice (CR:028 from Charles River, age 6-8

weeks or older) were used as recipient mice for transplant experiments, and Thy1.1 mice

(B6.PL-Thy1a/CyJ, 000406 from Jackson Labs) were used as recipient mice for chimera

experiments.

Allotransplant and Tumor Models. BALB/c recipient mice were irradiated with two doses of

400 cGy of x-rays (total dose 800 cGy) and rested for at least 12 hours between doses. Mice

were also rested for 4 hours prior to transplantation. T cells (total CD3+ or CD8+) were

separated from WT and *TCF-7 cKO* spleens using CD90.2 or CD8 microbeads and LS columns

(Miltenyi, CD8: 130-117-044, CD90.2: 130-121-278, LS: 130-042-401). 1×10^6 donor cells

(unless otherwise mentioned) were injected IV into the tail vein in PBS, along with 10×10^6 WT

bone marrow cells. Bone marrow was T-cell depleted with CD90.2 MACS beads (130-121-278

from Miltenyi) and LD columns (130-042-901 from Miltenyi). For short-term experiments, at

day 7 post-transplant, recipient mice were euthanized and serum, spleen, lymph nodes, small

intestine, or liver were collected, depending on the experiment. For GVHD and GVL

experiments, recipient mice were also given 2×10^6 luciferase expressing B-cell lymphoma (A-

20) (Edinger *et al.*, 2003a). Recipient mice were weighed, given a clinical score, and imaged

using the IVIS 50 imaging system three times per week until day 70 or longer. Clinical scores

were composed of scores for skin integrity, fur texture, posture, activity, diarrhea, and weight

loss. Imaging was done by injecting recipients I.P. with D-luciferin to detect tumor cell

bioluminescence. To produce mixed bone marrow chimeras, Thy1.1 mice were lethally

irradiated and reconstituted with a 1:4 (WT: *TCF-7 cKO*) mixture of bone marrow cells (total

$50x10^6$ cells), then rested for 9 weeks. At 9 weeks, tail vein blood was collected and checked by

flow cytometry for CD45.1 and CD45.2 to ensure reconstitution with both donor cell types. At

10 weeks, mice were used for phenotyping experiments.

Flow Cytometry, Sorting, and Phenotyping. Splenocytes (or cells from other organs) were

obtained from WT or *TCF-7* cKO mice or recipient allotransplanted mice. Lymphocytes were

obtained and lysed with RBC Lysis Buffer (00-4333-57 from eBioscience) to remove red blood

cells if needed. Cells were then stained with extracellular markers for 30 min on ice in MACS

buffer (1X PBS with EDTA and 4g/L BSA). If intracellular markers were used, the cells were

then fixed and permeabilized using the Fix/Perm Concentrate and Fixation Diluent from FOXP3

Transcription Factor Staining Buffer Set (eBioscience cat. No. 00-5523-00). The cells were then

run on a BD LSR Fortessa cytometer and data were analyzed using FlowJo software v9

(Treestar). All antibodies were used at a 1:100 dilution. For FACS sorting, the same methods

were applied, and cells were run on a BD FACS Aria IIIu with cold-sorting blocks. Cells were

sorted into sorting media (10% FBS in RPMI) or Trizol, depending on the experiment.

Depending on the experiment, antibodies used were: anti-CD4 (FITC, PE, BV785, BV21), anti-

CD8 (FITC, PE, APC, PerCP, Pacific Blue, PE/Cy7), anti-CD3 (BV605 or APC/Cy7), anti-

H2Kb-Pacific Blue, anti-H2Kd-PE/Cy7, anti-CD122 (FITC or APC), anti-CD44 (APC or Pacific

Blue), anti-CD62L (APC/Cy7), anti-TNF-α-FITC, anti-IFN-γ-APC, anti-Eomes (AF488 or

PE/Cy7), anti-T-bet-BV421, anti-CD45.2-PE/Dazzle594, anti-CD45.1-APC, anti-Ki67 (PE or

BV421), anti-PD1-BV785, anti-CTLA4-PE, NKG2D-BV711, Granzyme B-PE/Cy7.

Histology. Recipient mice were allotransplanted as described, and organs were removed for

histology at day 7, and day 21 post-transplant. Spleen, liver, small intestine, and skin (from ear

and back) were fixed, sectioned, and stained with H&E at Cornell University (

https://www.vet.cornell.edu/animal-health-diagnostic-center/laboratories/anatomic-

pathology/services). A pathologist (L.S) analyzed the sections for T cell-induced damage.

Cytokine Restimulation. Recipient BALB/c mice were allotransplanted with 1.5×10^6 CD3

donor T cells and euthanized at day 7. Splenocytes were taken and cultured for 6 hours with

GolgiPlug (1:1000) and PBS (control) or anti-CD3 (1ug/mL)/anti-CD28 (2ug/mL) (TCR

stimulation) at 37 C and 7% CO2. After 6 hours of culture, the cells were stained for CD3, CD4,

CD8, $H2K^b$, TNF-α, and IFN-γ using the BD Cytokine Staining kit (BD Biosciences, 555028),

and run on a flow cytometer.

LegendPLEX Serum ELISA Assay. Serum from cardiac blood was collected from recipient

mice in the cytokine restimulation experiment. Serum was analyzed using the Biolegend

LEGENDPlex Assay Mouse Th Cytokine Panel kit (741043). This kit quantifies serum

concentrations of: IL-2 (T cell proliferation), IFN-γ and TNF-α (Th1 cells, inflammatory), IL-4,

IL-5, and IL-13 (Th2 cells), IL-10 (Treg cells, suppressive), IL-17A/F (Th17 cells), IL-21 (Tfh

cells), IL-22 (Th22 cells), IL-6 (acute/chronic inflammation/T cell survival factor), and IL-9

(Th2, Th17, iTreg, Th9 – skin/allergic/intestinal inflammation).

Western blot. Splenocytes from WT or *TCF-7* cKO donor mice were collected. CD8 T cells

were separated using CD8 MACS beads. CD8 T cells were either stimulated with 2.5ug/ml anti-

CD3 (Biolegend #100202) and anti-CD28 antibodies (Biolegend #102115) for 10 minutes or left

unstimulated. These cells were counted and lysed with RIPA Buffer (89900 from Thermo Fisher)

plus protease inhibitors (11697498001 from Millipore Sigma) and phosphatase inhibitors

(P5726-1ML and P0044-1ML from Millipore Sigma). The lysates were run on a Western blot

and probed for Perforin (Cell Signaling Technology #3693), Granzyme B (Cell Signaling

Technology #4275), LCK (Thermo Fisher PA5-34653), ZAP70(Cell Signaling Technology

#3165), LAT(Cell Signaling Technology # 45533), ITK (Thermo Fisher PA5-49363), PLCγ1

(Cell Signaling Technology #2822), ERK1-2 (Cell Signaling Technology #9107), JAK 2(Cell

Signaling Technology # 3230), JAK3 (Cell Signaling Technology #8863), STAT3 (Cell

Signaling Technology #9139), p65-Rela (Cell Signaling Technology #4764), AKT (Cell

Signaling Technology #9272), , and β-actin (Cell Signaling Technology #4970). All the western

blots repeated at least three times and one representative of each protein and quantification is

shown.

qPCR Analysis. To perform qPCR, BALB/c mice were allotransplanted as described ($1X10^6$

CD3 donor T cells). Pre-transplant donor cells and post-transplant (day 7) spleen and liver cells

from recipients were FACS-sorted to obtain CD8 donor cells. The cells were sorted into Trizol,

RNA was extracted using chloroform (https://www.nationwidechildrens.org/

Document/Get/93327), and eluted using the Qiagen RNEasy Mini kit (74104 from Qiagen).

Concentration was checked with a spectrophotometer, then RNA was converted to cDNA with

an Invitrogen Super Script IV First Strand synthesis System kit (18091050 from Invitrogen).

Final cDNA concentration was checked with a spectrophotometer, then cDNA was mixed with

Taqman Fast Advanced Master Mix (4444557 from Invitrogen) at a 10ng/μL cDNA

concentration. This master mix was added to premade 96 well TaqMan Array plates with

chemokine/chemokine receptor primers (Thermo Fisher, Mouse Chemokines & Receptors Array

plate, 4391524). qPCR was performed in a Quant Studio 3 thermocycler, and data were analyzed

using the Design and Analysis software v2.4 (provided by Thermo Fisher). Five separate

recipient mice were sorted, and cells were combined to make one sample for qPCR testing per

condition/organ.

NKG2D expression and NKG2D mediated cytotoxicity in CD8 T cells. To determine the

NKG2D expression in CD8 T cells, we obtained splenocytes from WT and *TCF-7* cKO mice and

stimulated T cells with 2.5ug/ml anti-CD3 (Biolegend #100202) and anti-CD28 antibodies

(Biolegend #102115) for 24, 48, or 72 hours in culture, or left them unstimulated. GolgiPlug

(1:1000) was added to stimulated samples for each time point, and samples were incubated at 37

C and 7% CO_2. After 6 hours of culture, the cells were stained with LIVE/DEAD Aqua and for

CD3, CD8, NKG2D, and Granzyme B using the BD Cytokine Staining kit (BD Biosciences,

555028), and run on a flow cytometer. To assess the NKG2D mediated cytotoxicity, we used

luciferase-expressing A20 cells as target cells as described earlier. Effector cells (MACS-sorted

CD8 T cells from *TCF-7* cKO or WT mice) were incubated in 2.5μg/ml anti-CD3 and anti-CD28

coated plates for 48 hours to induce optimal NKG2D expression. Then effector cells were added

at 40:1 effector-to-target ratios and incubated at 37°C for 4 hours with the A20 cells. Anti-

NKG2D antibody (10 µg/mL, Bio X Cell #BE0334) or rat IgG1 isotype control antibody (10

µg/mL, Bio X Cell #BE0334) was added and incubated for 30 minutes before washing and

plating. Triplicate wells were averaged and percent lysis was calculated from the data using the

following equation: % specific lysis = 100 × (spontaneous death bioluminescence – test

bioluminescence)/(spontaneous death bioluminescence – maximal killing bioluminescence).

Exhaustion/Activation Assay. To determine the *in vitro* exhaustion and activation of CD8 T

cells, we obtained splenocytes from WT and *TCF-7* cKO mice and either activated them with

2.5ug/ml anti-CD3 (Biolegend #100202) and anti-CD28 antibodies (Biolegend #102115) for 24,

48, or 72 hours in culture, or left them unstimulated, and stained for CD3, CD8, Ki-67, Tox, and

PD-1 markers. To assess exhaustion and activation of *in vivo* donor CD8 T cells, recipient mice

were allotransplanted as before ($1X10^6$ CD3 donor T cells) and euthanized at day 7.

Lymphocytes were obtained from spleen, and stained for CD3, CD4, CD8, H2K^b, TOX, Ki-67

and PD-1 markers.

DNA Extraction and PCR. Donor mice were genotyped using PCR on DNA extracted from ear

punches. At 4 weeks of age mice were ear punched, and DNA was extracted using the Accustart

II Mouse Genotyping kit (95135-500 from Quanta Biosciences). Standard PCR reaction

conditions and primer sequences from Jackson Laboratories were used for CD4cre. For Tcf7,

primer sequences and reaction conditions were obtained from Dr. Jyoti Misra Sen of NIH.

RNA Sequencing. Recipient mice were allotransplanted as before ($1\text{X}10^6$ CD3 donor T cells),

except that donor CD8 T cells were also FACS-sorted prior to transplant. A sample of sorted

donor cells was also saved for pre-transplanted RNA sequencing in Trizol. At day 7 post-

transplant, donor CD8 T cells were FACS-sorted back from recipient spleen of *TCF-7* cKO and

WT transplanted mice. The cells were all sorted into Trizol, then RNA was extracted and

prepped by the Molecular Analysis Core (SUNY Upstate,

https://www.upstate.edu/research/facilities/molecular-analysis.php). Paired end sequencing was

done with an Illumina NovaSeq 6000 system at the University at Buffalo Genomics Core

(http://ubnextgencore.buffalo.edu). For data analysis we used the statistical computing

environment R (v4.0.4), the Bioconductor suite of packages for R, and R studio (v1.4.1106). We

calculated the transcript abundance by performing pseudoalignment using the Kallisto (Bray et

al., 2016) (version 0.46.2). Calculated Transcript per million (TPM) values were normalized and

fitted to a linear model by empirical Bayes method with the Voom (Law et al., 2014) and Limma

(Ritchie et al., 2015) R packages to determine Differentially expressed genes – DEGs (FDR<0.1,

after controlling for multiple testing using the Benjamini-Hochberg method). DEG's were used

for hierarchical clustering and heatmap generation in R. Gene Ontology enrichment analysis was

conducted using either the Function Annotation Chart tools using only GO BP and KEGG terms

in the Database for Visualization and Integrative Discovery (Huang da et al., 2009a; b) (Huang

da *et al.*, 2009a) DAVID enrichment scores >1.3 are equivalent to a P value<0.05. For Gene set

enrichment analysis (GSEA) we used Hallmark and C2 gene set collections of Molecular

Signatures Database (MsigDB) and cluster Profiler package in R. Data will be deposited on the

Gene Expression Omnibus (GEO) database for public access

https://www.ncbi.nlm.nih.gov/geo/

The RNAseq experiment described here was performed as part of the experiment described in

other recent publications from our laboratory (Mammadli et al., 2021a; Mammadli et al., 2021b;

Mammadli et al., 2020; Mammadli et al., 2021c). Therefore, the data generated for WT pre- and

post-transplanted samples (CD4 and CD8) are the same as that shown in the papers mentioned,

but here, these data are compared to data for *Cat-Tg* mice(Mammadli *et al.*, 2021a).

Statistical Analysis. Unless otherwise noted in the figure legends, all numerical data are

presented as means and standard deviations with or without individual points. Analysis was done

in GraphPad Prism v7 or v9. Most data were analyzed with Student's t-test, one-way ANOVA,

or two-way ANOVA, with Tukey's multiple comparisons test for ANOVA methods, depending

on the number of groups. Kaplan-Meier survival analyses were done for survival experiments.

All tests were two-sided, and p-values less than or equal to ($\leq$) 0.05 were considered significant.

Transplant experiments used 3-5 mice per group, with at least two repeats. *Ex vivo* experiments

were done two to three times unless otherwise noted with at least three replicates per condition

each time. RNA seq was done once with three replicates per group and condition. qPCR was

done once with one sample per condition, and 5 mice were combined to make the one sample.

Western blots were done 3 times for unstimulated and 10 min anti-CD3/CD28 stimulated

samples, one experiment each is shown.

Study Approval. All animal studies were reviewed and approved by the IACUC at SUNY

Upstate Medical University. All procedures and experiments were performed according to these

approved protocols.

Author Contributions

Author Contributions

RH, MM, and MK designed and conducted experiments, analyzed data, and wrote the

manuscript. MK assisted with scientific/technical research design. MK, and JMS edited the

manuscript. MK, SH assisted with data collection. LS performed histology analyses. QY

provided technical and scientific advice and assisted with data analysis.

Acknowledgments

We thank all members of the Karimi lab for helpful discussions. We also thank Joel Wilmore for

help in flow cytometry analysis. This research was funded in part by a grant from the National

Blood Foundation Scholar Award to MK, the National Institutes of Health (NIH LRP #L6

MD0010106 and K22 (AI130182) to MK), and an Upstate Medical University Cancer Center

grant (1146249-1-75632) to MK.

JMS was supported by the Intramural Research Program of the National Institute of

Aging. We thank Dr. Howard Xue for permission to use *TCF-7* cKO mice. *TCF-7* flox/flox

mice were provided by Dr. Jyoti Misra Sen from NIH. RH was a PH. D student at SUNY

Upstate Medical University at the time the study was conducted from 2017-2021. A version of

this manuscript was previously included as a chapter in RH's dissertation.

Summary Figure: *TCF-7* is required for persistent function of CD8 T cells but dispensable

for anti-tumor response. Here we have utilized a novel mouse model that lacks *TCF-7*

specifically on CD8 T cells for an allogeneic transplant model. We uncovered a molecular

mechanism of how *TCF-7* regulates key signaling pathways at both transcriptomic and protein

levels. These key molecules included LCK, LAT, ITK, PLC-γ1, p65, ERK I/II, and JAK/STAT

signaling. Next, we showed that the lack of *TCF-7* impacted phenotype, proinflammatory

cytokine production, chemokine expression, T cell activation. We provided clinical evidence for

how these changes impact GVHD target organs (skin, small intestine, and liver). Finally, we

provided evidence that *TCF-7* regulates NKG2D expression on mouse naïve and activated CD8

T cells. We have shown that CD8 T cells from *TCF-7* cKO mice mediate cytolytic functions via

NKG2D.

Figure 1: Loss of *TCF-7* in donor CD8 T cells reduces severity and persistence of GVHD

symptoms. BALB/c recipient mice (MHC haplotype d) were lethally irradiated and

allotransplanted with 1×10^6 CD8 T cells from WT or *TCF-7* cKO donor mice (MHC haplotype

b), as well as 10×10^6 T cell-depleted bone marrow cells (BM) from WT mice (MHC haplotype

b). Recipient mice were also given 2×10^5 luciferase-expressing A20 tumor cells **(A)** The

recipient mice were imaged 1 time a week using IVIS50 for 70 days. Gross pictures of

representatives of recipients transplanted with CD8 T cells from *TCF-7* cKO and WT mice at

day 25 post-transplant are shown. **(B)** Recipient mice were also given a GVHD clinical score

three times per week until day 70, based on combined scores of fur texture, activity level, weight

loss, posture, skin integrity, and diarrhea. Mean and SD are plotted, analyzed by one-way

ANOVA. **(C)** Weight changes of the recipient mice also were tracked over the time course of

disease. **(D)** Survival for each group of recipient mice up to 70 days post-transplant, analyzed by

Kaplan-Meier survival analysis. **(E)** Quantification of bioluminescence of tumor growth. For all

graphs, * means p-value ≤ 0.05, ** means p-value ≤ 0.01, *** means p-value ≤ 0.001, and ****

 means p-value ≤ 0.0001 (*n* = 3 mice/group for BM alone: *n* = 4 experimental mice/group for all

other groups). Survival is a combination of two experiments. Note: Control mouse is one of the

mice from bone marrow only transplanted group used as a negative control for BLI.

Figure 2. Loss of *TCF-7* changes the mature CD8 T cell phenotype. Naive WT or *TCF-7*

cKO donor mice were euthanized and splenocytes were stained for flow cytometry phenotyping**.**

(A) Percent of CD8 T cells expressing *TCF-7* and quantified statistical analysis **(B)** Percent of

CD8 T cells expressing Eomes and quantified statistical analysis. **(C)** Percent of CD8 T cells

expressing T-bet and quantified statistical analysis. **(D)** Percent of CD8 T cells expressing

CD122 and quantified statistical analysis. **(E)** Percent of CD8 T cells expressing CD44 and

quantified statistical analysis. **(F)** Percent of CD8 T cells expressing central memory, effector

memory, or naive phenotypes and quantified statistical analysis. All data are plotted as individual

Figure 3: Loss of *TCF-7* alters cytotoxic mediator production by mature alloactivated CD8

T cells. Total splenocytes were isolated from *TCF-7* cKO or WT mice and either left

unstimulated or stimulated with anti-CD3/CD28 for 24, 48, or 72 hours in culture. GolgiPlug

(1:1000) was added to stimulated samples in each time point except 0-hour samples, and samples

were incubated at 37 C with 7% CO_2. After 6 hours of culture, the cells were stained with

LIVE/DEAD Aqua, and for CD3, CD8, NKG2D, and Granzyme B expression, as determined by

flow cytometry. **(A)** Percent of NKG2D expression in freshly isolated CD8 T cells from WT or

TCF-7 cKO mice and quantified statistical analysis. **(B)** Percent of NKG2D expression in 24, 48,

or 72 hour-stimulated CD8 T cells from WT and *TCF-7* cKO mice and quantified statistical

analysis. **(C)** Percent of Granzyme B expression in unstimulated or 24, 48, or 72 hour-stimulated

CD3+ CD8+ NKG2D+ T cells from WT and *TCF-7* cKO mice, and quantified statistical

analysis. **(D)** Granzyme B expression in unstimulated CD8 T cells from WT and *TCF-7* cKO

mice by Western blot, and bands' integrated density normalized to β-actin (quantified). **(E)**

Granzyme B expression in 10 minute-anti-CD3/CD28-stimulated CD8 T cells from WT and

TCF-7 cKO mice by Western blot, and bands' integrated density normalized to β-actin

(quantified) **(F)** To assess the NKG2D-mediated cytotoxicity, we used luciferase-expressing A20

cells as target cells. Effector cells (MACS-sorted CD8 T cells from *TCF-7* cKO or WT mice)

were incubated in 2.5μg/ml anti-CD3 and anti-CD28 coated plates for 48 hours to induce optimal

NKG2D expression. Then effector cells were added at 40:1 effector-to-target ratios and

incubated at 37°C for 4 hours with A20 cells. Anti-NKG2D antibody (10 μg/mL) or rat IgG1

isotype control antibody (10 μg/mL) was added and incubated for 30 minutes before washing

and plating. Triplicate wells were averaged and percent lysis was calculated from the data using

the following equation: % specific lysis = 100 × (spontaneous death bioluminescence – test

bioluminescence)/ (spontaneous death bioluminescence – maximal killing

bioluminescence)(Karimi et al., 2014a; Karimi et al., 2015; Karimi et al., 2014b). N=4 per group

with one rappresentative of 2 experiments shown. Western blot were repeated twice and

representative examples are shown. All data are shown as individual points with mean and SD,

and were analyzed with Student's t-test or one-way ANOVA (depending on groups). * means p-

value ≤ 0.05, ** means p-value ≤ 0.01, and *** means p-value ≤ 0.001.

Fig.4. *TCF-7* controls cytokine production and exhaustion of mature alloactivated CD8 T

cells. (A-B) Recipient mice were allotransplanted with $1.5X10^6$ WT or *TCF-7* cKO donor CD3 T

cells as before**.** At day 7 and day 14 post-transplant, serum was obtained from cardiac blood of

allotransplanted recipient mice and tested for various cytokines using a LEGENDplex ELISA

assay. **(A)** Serum levels (pg/mL) of TNF-α, IFN-γ, and IL-2 for WT and *TCF-7* cKO-

transplanted mice at day 7 post-transplant. **(B)** Serum levels (pg/mL) of TNF-α, IFN-γ, and IL-2

for WT and *TCF-7* cKO-transplanted mice at day 14 post-transplant. **(C-F)** Splenocytes from

TCF-7 cKO and WT mice were obtained and stimulated with anti-CD3/CD28 antibodies for

24hrs, 48hrs, or 72hrs in culture and stained for Ki-67 and PD-1 or were stained immediately

after isolation without stimulation. **(C)** Percent expression of Ki-67 in CD8 T cells after 72

hours of anti-CD3/CD28 stimulation in culture determined by flow cytometry. **(D)**

Quantification of the *in vitro* Ki-67 expression of CD8 T cells at different time points. **(E)**

Percent expression of PD-1 in CD8 T cells after 72 hours of anti-CD3/CD28 stimulation in

culture determined by flow cytometry. **(F)** Quantification of the *in vitro* PD-1 expression of CD8

T cells are different time points**. (G-H)** Balb/c mice were allotransplanted as before, with WT or

TCF-7 cKO CD8 donor T cells. On day 7, spleens were removed from recipients, processed to

isolate lymphocytes, and spleen-derived lymphocytes from allotransplanted recipient mice were

stained for CD3, CD8, H2K^b, and TOX to identify exhausted T cells. **(G)** Percent expression of

TOX in CD8 T cells after allotransplantation (day 7 post-transplant) as determined by flow

cytometry. **(H)** Quantification of the TOX expression in *in vivo* donor CD8 T cells. N=3-5 per

group for **A-B** with two experiments shown. N=4 per group for **C-F,** one representative of two

experiments shown. N=3-5 per group with one representative experiment shown for **G-H**. All

data are shown as individual points with mean and SD, and were analyzed with Student's t-test

or two-way ANOVA (depending on groups). * means p-value $\leq$ 0.05, ** means p-value $\leq$ 0.01,

and *** means p-value $\leq$ 0.001.

Fig.5. Loss of *TCF-7* in donor CD8 T cells decrease the damage to the GVHD target organs.
We collected organs from mice allotransplanted as described above. At day 7 and day 21 post-
transplant, organs were taken from recipient mice for histology analyses. Skin, liver, spleen, and
small intestine were sectioned, stained with H&E and analyzed by pathologist. Representative
sections for each organ per group and timepoint are shown. **(A)** H&E staining of the liver of the

recipient mice at day 7 post-transplant (black arrows showing the interlobular bile ducts and red

circle showing inflammatory infiltrates). **(B)** H&E staining of the small intestines of the recipient

mice at day 7 post-transplant (black arrows showing the crypts of the small intestine and red

circle showing apoptotic bodies). **(C)** H&E staining of the skin of the recipient mice at day 7

post-transplant (black arrows showing the dermis of the skin and red circle showing increase of

inflammatory cells) **(D)** H&E staining of the liver of the recipient mice at day 21 post-transplant

(black arrows showing the interlobular bile ducts and red circle showing inflammatory

infiltrates). **(E)** H&E staining of the small intestines of the recipient mice at day 21 post-

transplant (blacks arrows showing apoptotic bodies). **(F)** H&E staining of the skin of the

recipient mice at day 21 (black arrows showing dermis of skin and red circle showing increased

inflammatory cells with destructed adnexal glands).

Fig.6. Loss of *TCF-7* changes the genetic signature of alloactivated donor CD8 T cells.

CD8 T cells from WT and *TCF-7* cKO mice were FACS sorted either into Trizol as pre-

transplanted samples, or into 10% FBS-containing media for transplantation into recipient mice

as described previously. At day 7 post-transplant, donor CD8 T cells were FACS-sorted back

from recipient spleens of *TCF-7* cKO- and WT-transplanted mice. RNA was extracted and

prepped by the Molecular Analysis Core (SUNY Upstate). Paired end sequencing was done with

an Illumina NovaSeq 6000 system at the University at Buffalo Genomics Core. For data analysis,

we used the statistical computing environment R (v4.0.4), the Bioconductor suite of packages for

R, and RStudio (v1.4.1106). We calculated the transcript abundance by performing

pseudoalignment using Kallisto. **(A)** Hierarchical clustering of genes and samples, heatmap

illustrating the expression of the differentially expressing genes (DEG's; FDR<0.1) of post-

transplanted CD8 T cells *TCF-7* cKO compared to WT. **(B)** Table showing the number of up- or

downregulated DEGs in post-transplanted CD8 T cells (*TCF-7* cKO compared to WT). **(C)**

Heatmap showing top significant 35 differentially expressed genes that play a role in the Cell

Cycle pathway, identified using DAVID Functional Annotation Analysis and GO-BP terms in

post-transplanted CD8 T cells (*TCF-7* cKO compared to WT). **(D)** Heatmap showing top

significant 30 differentially expressed genes that play a role in the Apoptotic Processes pathway,

identified using DAVID Functional Annotation Analysis and GO-BP terms in post-transplanted

CD8 T cells (*TCF-7* cKO compared to WT). **(E)** Heatmap showing top significant 30

differentially expressed genes that play a role in the Metabolic Processes pathway, identified

using DAVID Functional Annotation Analysis and GO-BP terms in post-transplanted CD8 T

cells (*TCF-7* cKO compared to WT). **(F)** Heatmap showing top significant 35 differentially

expressed genes that play a role in the NF-κB pathway, identified using DAVID Functional

Annotation Analysis and GO-BP terms in post-transplanted CD8 T cells (*TCF-7* cKO compared

to WT). **(G)** Gene set enrichment analysis (GSEA) enrichment plots of Cytokine signaling

pathways, including IL-1, IL-2, IL-4, and IL-7 signaling pathways from WP terms that are

enriched in post-transplanted CD8 T cells from WT mice (versus *TCF-7* cKO). Negative

Enrichment Score is an indicator of downregulation, and positive Enrichment Score is an

indicator of upregulation of the genes in the post-transplanted CD8 T cells from *TCF-7* cKo

mice. DAVID enrichment scores >1.3 are equivalent to a P value<0.05 **(H)** GSEA enrichment

plots of G2 to M Cell Cycle and Cell Cycle S pathways that are enriched in *TCF-7* cKO mice,

and Integrin Mediated Cell Adhesion and Apoptosis pathways that are enriched in post-

transplanted CD8 T cells from WT mice. **(I)** GSEA enrichment plots of JAK-STAT signaling,

Toll like receptor signaling, T cell receptor signaling, ERK MAPK signaling, signaling by

Tyrosine kinases, Rankl - Rank mediated NF-kB signaling pathways that are enriched in post-

transplanted CD8 T cells from WT mice. Again, Negative Enrichment Score is an indicator of

downregulation, and positive Enrichment Score is an indicator of upregulation of the genes in the

post-transplanted CD8+ T cells from *TCF-7* cKo mice. DAVID enrichment scores >1.3 are

equivalent to a P value<0.05.

1010

Fig.7. Loss of *TCF-7* decrease TCR, JAK-STAT, and NF-kB signaling downstream. (A) A

bar plot showing KEGG (Kyoto Encyclopedia of Genes and Genomes) pathways identified by

David functional annotation analysis. **(B)** Heatmap showing the altered genes in TCR signaling

pathway from KEGG pathways**. (C)** Heatmap showing the altered genes in JAK-STAT signaling

pathway from KEGG pathways. **(D)** Western blot showing the protein expression levels of LCK,

ZAP70, LAT, ITK, PLCγ1, ERK1/2, JAK2, JAK3, STAT3, P65-RelA, AKT, and β-actin of

freshly isolated CD8 T cell lysates from *TCF-7* cKO and WT mice**. (E)** Western blot showing

the protein expression levels of LCK, ZAP70, LAT, ITK, PLCγ1, ERK1/2, JAK2, JAK3,

STAT3, P65-RelA, AKT, and β-actin in 10 minute-CD3/CD28-stimulated CD8 T cell lysates

from *TCF-7* cKO and WT mice. All the western blots repeated at least three and one

representative of each protein and quantification is shown.

1022

1023

1024 **Supplemental Figure Legends:**

1025

Supp.Fig.1. Related to Fig2. Loss of *TCF-7* drives changes to mature CD8 T cell phenotype

that are cell-intrinsic, with the possibility of extrinsic effects. Bone marrow chimeras were

developed by lethally irradiating Thy1.1 mice and reconstituting with a 1:4 (WT:*TCF-7* cKO)

mixture of bone marrow cells. Blood was tested at 9 weeks to ensure reconstitution with both

donor cell types, and splenocytes were used at 10 weeks for phenotyping by flow cytometry. **(A)**

Percentage of CD8 T cells from chimeric and naive mice expressing *TCF-7*. **(B)** Percentage of

CD8 T cells from chimeric and naive mice expressing T-bet. **(C)** Percentage of CD8 T cells from

chimeric and naive mice expressing Eomes. **(D)** Percentage of CD8 T cells from chimeric and

naive mice expressing CD122. **(E)** Percentage of CD8 T cells from chimeric and naive mice

expressing CD44. **(F)** Percentage of CD8 T cells from chimeric and naive mice expressing

central memory (CM) phenotype. **(G)** Percentage of CD8 T cells from chimeric and naive mice

expressing effector memory (EM) phenotype. **(H)** Percentage of CD8 T cells from chimeric and

naive mice expressing naïve phenotype. All data are plotted as individual points with mean and

SD, all were analyzed with one-way ANOVA, or Student's t-test (depending on groups). For all

graphs, * means p-value ≤ 0.05, *** means p-value ≤ 0.001, and **** means p-value ≤ 0.0001.

For naïve cells 3 different experiments combined (N=2-3 per group of mice) and for chimera cell

(N=5) with one experiment shown (done once).

1043

Supp.Fig.2 Related to Fig.3. Cell viability and CD8 + T cell percent in NKG2D induction *in*

vitro. Total splenocytes were isolated from *TCF-7* cKO or WT mice and either left unstimulated

or stimulated with anti-CD3/CD28 for 24, 48, or 72 hrs in culture. GolgiPlug (1:1000) was added

to stimulated samples for each time point except 0 hr samples, and samples were incubated at 37

C with 7% CO_2. After 6 hours of culture, the cells were stained for CD3, CD8, NKG2D, and

Granzyme B expression, as determined by flow cytometry. **(A)** Quantification of percent of live

cells (dead cells that were positive for LIVE/DEAD Aqua excluded) for different time points of

stimulation. **(B)** Quantification of CD8 T cell percentages for different time points of

stimulation. N=4 per group with one representative of 2 experiments shown. All data are shown

as individual points with mean and SD, and were analyzed with two-way ANOVA (depending

on groups). * means p-value ≤ 0.05, ** means p-value ≤ 0.01, and *** means p-value ≤ 0.001.

1056

Supp.Fig3. Related to Fig.5. *TCF-7* controls cytokine production and exhaustion of mature

alloactivated CD8 T cells. (A-B) Recipient mice were allotransplanted with $1.5X10^6$ WT or

TCF-7 cKO donor CD3 T cells as before. Splenocytes were taken at day 7 post-transplant,

restimulated by 6-hour culture with Golgi Plug and PBS (control) or anti-CD3/anti-CD28 (TCR

restim.), then stained for $H2K^b$, CD3, CD4, CD8, TNF-α, and IFN-γ. **(A)** TNF-α production and

(B) IFN-γ production by donor CD8 T cells, as measured by percent cytokine-positive donor

cells. **(C)** Splenocytes from *TCF-7* cKO and WT mice were obtained and either stimulated with

anti-CD3/CD28 antibodies for 24hrs, 48hrs, or 72hrs in culture and stained for TOX, or were

immediately stained after isolation without stimulation. Percent expression of TOX in CD8 T

cells after 72 hours of CD3/CD28 stimulation in culture, as determined by flow cytometry. **(D-E)**

Balb/c mice were allo-transplanted as before, with WT BM and WT or *TCF-7* cKO CD8 donor T

cells. On day 7, spleens were removed from recipients, processed to isolate lymphocytes, and

lymphocytes were stained for CD3, CD8, H2K[b], Ki-67, and PD-1 to identify proliferating and

exhausted T cells, respectively. **(D)** Quantification of the Ki-67 expression in *in vivo* donor CD8

T cells. (**E)** Quantification of the PD-1 expression in *in vivo* donor CD8 T cells. N=3-5 per group

for **A-B and D-E** with two experiments shown. N=4 per group for **C,** one representative of two

experiments shown. All data are shown as individual points with mean and SD, and were

analyzed with Student's t-test or two-way ANOVA (depending on groups). * means p-value ≤

0.05, ** means p-value ≤ 0.01, and *** means p-value ≤ 0.001.

1076

Supp.Fig.4. *TCF-7* **controls chemokine and chemokine receptor expression of donor CD8 T**

cells during alloactivation. As before, BALB/c mice were allotransplanted with BM and 1X10^6

donor CD3 T cells from WT or *TCF-7* cKO mice (not mixed). Donor CD8 T cells were FACS-

sorted from spleen of donor's pre-transplant, and from spleen and liver of recipients at day 7

post-transplant. Cells were sorted into Trizol, then RNA was extracted using chloroform and

converted to cDNA for qPCR analysis. cDNA was run on premade mouse chemokine/chemokine

receptor assay plates, and results are displayed as heatmaps. Scales are shown at right, with fold

change per gene compared to an 18S reference gene on each plate. White boxes with an "X"

represent signals too low to detect or otherwise unreadable due to technical limitation/error. **(A)**

Pre-transplant spleen donor CD8 T cells, **(B)** post-transplant spleen donor CD8 T cells, and **(C)**

post-transplant liver donor CD8 T cells for WT versus *TCF-7* cKO donors. N=5 mice into one

sample per condition**,** summary data shown.

1089

Supp.Fig.5: Loss of *TCF-7* **alters the enrichment of gene sets in pre-transplanted CD8+T**

cells. (A) Bar plot of the altered pathways indentified in Gene set enrichment analysis (GSEA)

by using Hallmark gene set collections of the Molecular Signatures Database (MSigDB) in pre-

transplanted CD8 T cells (*TCF-7* cKO compared to WT). The normalized enrichment score

(NES) for the pathway is defined as the peak score furthest from zero, with a negative NES

meaning enrichment in the WT group. **(B)** GSEA plot for the

"HALLMARK_TNFA_SIGNALING_VIA_NFKB" pathway comparing pre-transplanted CD8 T

cells from *TCF-7* cKO to WT cells. The running enrichment score (ES) for the pathway is

defined as the peak score furthest from zero, with a negative ES meaning enrichment in the WT

group. **(C)** GSEA plot for the "HALLMARK_INTERFERON_GAMMA_RESPONSE" pathway

comparing pre-transplanted CD8 T cells from *TCF*-7 cKO to WT cells. **(D)** GSEA plot for the

"HALLMARK_MTORC1_SIGNALING" pathway comparing pre-transplanted CD8 T cells

from *TCF-7* cKO to WT cells. **(E)** GSEA plot for the

"HALLMARK_ALLOGRAFT_REJECTION" pathway comparing pre-transplanted CD8 T cells

from *TCF-7* cKO to WT cells. **(F)** GSEA plot for the

"HALLMARK_OXIDATIVE_PHOSPHORYLATION" pathway comparing pre-transplanted

CD8 T cells from *TCF-7* cKO to WT cells.

Top 20 pathways in GO-BP

Term ID	Term	Count	%	P Value	Fold Enrichment	FDR
GO:0007049	Cell cycle	141	5.73870574	2.34E-14	1.906511069	1.19E-10
GO:0015031	Protein transport	134	5.45380545	3.42E-13	1.879194277	8.74E-10
GO:0098609	Cell-cell adhesion	55	2.23850224	6.46E-10	2.415958527	9.33E-07
GO:0051301	Cell division	88	3.58160358	7.31E-10	1.953438125	9.33E-07
GO:0007067	Mitotic nuclear division	69	2.80830281	5.24E-09	2.068035126	5.35E-06
GO:0006915	Apoptotic process	114	4.63980464	4.40E-08	1.660422406	3.75E-05
GO:0006810	Transport	291	11.8437118	1.39E-07	1.325968496	1.01E-04
GO:0006974	Cellular response to DNA damage stimulus	86	3.5002035	8.53E-07	1.699956273	5.31E-04
GO:0016192	Vesicle-mediated transport	52	2.11640212	9.36E-07	2.026806693	5.31E-04
GO:0006783	Heme biosynthetic process	12	0.48840049	2.25E-06	5.243439176	0.0011491
GO:0002479	Antigen processing and presentation of exogenous peptide antigen via MHC class I, TAP-dependent	15	0.61050061	2.85E-06	4.151056015	0.00132461
GO:0050852	T cell receptor signaling pathway	20	0.81400081	3.32E-06	3.255730208	0.00141253
GO:0006260	DNA replication	34	1.38380138	5.93E-06	2.294892756	0.00232917
GO:0002376	Immune system process	76	3.09320309	1.26E-05	1.647416486	0.00460664
GO:0043123	Positive regulation of I-kappaB kinase/NF-kappaB signaling	36	1.46520147	6.87E-05	2.005879416	0.02338981
GO:0008152	Metabolic process	84	3.41880342	1.21E-04	1.506214709	0.03848315
GO:0031663	Lipopolysaccharide-mediated signaling pathway	13	0.52910053	1.40E-04	3.481530851	0.04077673
GO:0006281	DNA repair	62	2.52340252	1.44E-04	1.618650773	0.04077673
GO:0008643	Carbohydrate transport	14	0.56980057	2.33E-04	3.141339687	0.06257663

1107

1108 **Supp.Fig.6. Related to Fig.7**. Top 20 GO-BP terms identified in DAVID functional annotation

1109 analysis of differentially expressed genes in post-transplanted samples. Count mean – number of

1110 genes involved in the Term, % - percentage of involved genes/ total genes in the Term.

1111

Supp.Fig.7. Loss of *TCF-7* alters the Gene Set Enrichment Analysis (GSEA) of post-

transplanted CD8+T cells. (A) GSEA plot for the

"HALLMARK_PI3K_AKT_MTOR_SIGNALING" pathway comparing post-transplanted CD8

T cells from *TCF-7* cKO to WT mice. The running enrichment score (ES) for the pathway is

defined as the peak score furthest from zero, with a negative ES meaning enrichment in the WT

group. **(B)** GSEA plot for the "HALLMARK_TNFA_SIGNALING_VIA_NFKB" pathway

comparing post-transplanted CD8 T cells from *TCF-7* cKO to WT mice. Again, the running

enrichment score (ES) for the pathway is defined as the peak score furthest from zero, with a

negative ES meaning enrichment in the WT group.

1121

Supp.Fig.8. Related to Fig.8. **Quantification of Western blot of TCR and JAK-STAT**

signaling. (**A**) Comparison of the quantified bands integrated density normalized to β-actin for

unstimulated CD8 T cells from *TCF-7* cKO and WT mice**. (B)** Comparison of the quantified

bands integrated density normalized to β-actin for 10-minute-anti-CD3/CD28-stimulated CD8 T

cells from *TCF-7* cKO and WT mice. All the western blots repeated at least three times and one

representative of each protein and quantification is shown.

1128

1129

1130

1131

1132

1133

1134

1135

References

Abel, A.M., Yang, C., Thakar, M.S., and Malarkannan, S. (2018). Natural Killer Cells:
Development, Maturation, and Clinical Utilization. Front Immunol *9*, 1869.
10.3389/fimmu.2018.01869.
Ahn, E., Araki, K., Hashimoto, M., Li, W., Riley, J.L., Cheung, J., Sharpe, A.H., Freeman, G.J.,
Irving, B.A., and Ahmed, R. (2018). Role of PD-1 during effector CD8 T cell differentiation. Proc
Natl Acad Sci U S A *115*, 4749-4754. 10.1073/pnas.1718217115.
Al Dulaimi, D., Klibi, J., Olivo Pimentel, V., Parietti, V., Allez, M., Toubert, A., and Benlagha, K.
(2018). Critical Contribution of NK Group 2 Member D Expressed on Invariant Natural Killer T
Cells in Concanavalin A-Induced Liver Hepatitis in Mice. Front Immunol *9*, 1052.
10.3389/fimmu.2018.01052.
Balassa, K., Danby, R., and Rocha, V. (2019). Haematopoietic stem cell transplants: principles
and indications. Br J Hosp Med (Lond) *80*, 33-39. 10.12968/hmed.2019.80.1.33.
Bastien, J.P., Roy, J., and Roy, D.C. (2012). Selective T-cell depletion for haplotype-mismatched
allogeneic stem cell transplantation. Seminars in oncology *39*, 674-682.
10.1053/j.seminoncol.2012.09.004.
Beilhack, A., Schulz, S., Baker, J., Beilhack, G.F., Wieland, C.B., Herman, E.I., Baker, E.M., Cao,
Y.A., Contag, C.H., and Negrin, R.S. (2005). In vivo analyses of early events in acute graft-versus-
host disease reveal sequential infiltration of T-cell subsets. Blood *106*, 1113-1122.
10.1182/blood-2005-02-0509.
Beltra, J.C., Manne, S., Abdel-Hakeem, M.S., Kurachi, M., Giles, J.R., Chen, Z., Casella, V., Ngiow,
S.F., Khan, O., Huang, Y.J., et al. (2020). Developmental Relationships of Four Exhausted CD8(+)
T Cell Subsets Reveals Underlying Transcriptional and Epigenetic Landscape Control
Mechanisms. Immunity *52*, 825-841 e828. 10.1016/j.immuni.2020.04.014.
Berga-Bolanos, R., Zhu, W.S., Steinke, F.C., Xue, H.H., and Sen, J.M. (2015). Cell-autonomous
requirement for TCF1 and LEF1 in the development of Natural Killer T cells. Mol Immunol *68*,
484-489. 10.1016/j.molimm.2015.09.017.
Bleakley, M., Turtle, C.J., and Riddell, S.R. (2012). Augmentation of anti-tumor immunity by
adoptive T-cell transfer after allogeneic hematopoietic stem cell transplantation. Expert Rev
Hematol *5*, 409-425. 10.1586/ehm.12.28.
Blessin, N.C., Li, W., Mandelkow, T., Jansen, H.L., Yang, C., Raedler, J.B., Simon, R., Buscheck, F.,
Dum, D., Luebke, A.M., et al. (2021). Prognostic role of proliferating CD8(+) cytotoxic Tcells in
human cancers. Cell Oncol (Dordr) *44*, 793-803. 10.1007/s13402-021-00601-4.
Bray, N.L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic RNA-seq
quantification. Nat Biotechnol *34*, 525-527. 10.1038/nbt.3519.
Breems, D.A., and Lowenberg, B. (2005). Autologous stem cell transplantation in the treatment
of adults with acute myeloid leukaemia. Br J Haematol *130*, 825-833. BJH5628 [pii]
10.1111/j.1365-2141.2005.05628.x.
Chen, Z., Ji, Z., Ngiow, S.F., Manne, S., Cai, Z., Huang, A.C., Johnson, J., Staupe, R.P., Bengsch, B.,
Xu, C., et al. (2019). *TCF-1*-Centered Transcriptional Network Drives an Effector versus
Exhausted CD8 T Cell-Fate Decision. Immunity *51*, 840-855 e845.
10.1016/j.immuni.2019.09.013.
Chu, T., Tyznik, A.J., Roepke, S., Berkley, A.M., Woodward-Davis, A., Pattacini, L., Bevan, M.J.,

Zehn, D., and Prlic, M. (2013). Bystander-activated memory CD8 T cells control early pathogen
load in an innate-like, NKG2D-dependent manner. Cell Rep *3*, 701-708.
10.1016/j.celrep.2013.02.020.
Cieri, N., Camisa, B., Cocchiarella, F., Forcato, M., Oliveira, G., Provasi, E., Bondanza, A.,
Bordignon, C., Peccatori, J., Ciceri, F., et al. (2013). IL-7 and IL-15 instruct the generation of
human memory stem T cells from naive precursors. Blood *121*, 573-584. 10.1182/blood-2012-
05-431718.
Cooke, K.R., Kobzik, L., Martin, T.R., Brewer, J., Delmonte, J., Jr., Crawford, J.M., and Ferrara, J.L.
(1996). An experimental model of idiopathic pneumonia syndrome after bone marrow
transplantation: I. The roles of minor H antigens and endotoxin. Blood *88*, 3230-3239.
Cui, W., Liu, Y., Weinstein, J.S., Craft, J., and Kaech, S.M. (2011). An interleukin-21-interleukin-
10-STAT3 pathway is critical for functional maturation of memory CD8+ T cells. Immunity *35*,
792-805. 10.1016/j.immuni.2011.09.017.
Decman, V., Laidlaw, B.J., Dimenna, L.J., Abdulla, S., Mozdzanowska, K., Erikson, J., Ertl, H.C.,
and Wherry, E.J. (2010). Cell-intrinsic defects in the proliferative response of antiviral memory
CD8 T cells in aged mice upon secondary infection. J Immunol *184*, 5151-5159.
10.4049/jimmunol.0902063.
Dutt, S., Baker, J., Kohrt, H.E., Kambham, N., Sanyal, M., Negrin, R.S., and Strober, S. (2011).
CD8+CD44(hi) but not CD4+CD44(hi) memory T cells mediate potent graft antilymphoma
activity without GVHD. Blood *117*, 3230-3239. 10.1182/blood-2010-10-312751.
Edinger, M., Hoffmann, P., Contag, C.H., and Negrin, R.S. (2003a). Evaluation of effector cell fate
and function by in vivo bioluminescence imaging. Methods *31*, 172-179. 10.1016/s1046-
2023(03)00127-0.
Edinger, M., Hoffmann, P., Ermann, J., Drago, K., Fathman, C.G., Strober, S., and Negrin, R.S.
(2003b). CD4+CD25+ regulatory T cells preserve graft-versus-tumor activity while inhibiting
graft-versus-host disease after bone marrow transplantation. Nat Med *9*, 1144-1150.
10.1038/nm915.
Escobar, G., Mangani, D., and Anderson, A.C. (2020). T cell factor 1: A master regulator of the T
cell response in disease. Sci Immunol *5*. 10.1126/sciimmunol.abb9726.
Ferrara, J.L. (2014). Blood and Marrow Transplant Clinical Trials Network: Progress since the
State of the Science Symposium 2007. Biology of blood and marrow transplantation : journal of
the American Society for Blood and Marrow Transplantation *20*, 149-153.
10.1016/j.bbmt.2013.11.006.
Gattinoni, L., Lugli, E., Ji, Y., Pos, Z., Paulos, C.M., Quigley, M.F., Almeida, J.R., Gostick, E., Yu, Z.,
Carpenito, C., et al. (2011). A human memory T cell subset with stem cell-like properties. Nat
Med *17*, 1290-1297. 10.1038/nm.2446.
Gautam, S., Fioravanti, J., Zhu, W., Le Gall, J.B., Brohawn, P., Lacey, N.E., Hu, J., Hocker, J.D.,
Hawk, N.V., Kapoor, V., et al. (2019). The transcription factor c-Myb regulates CD8(+) T cell
stemness and antitumor immunity. Nat Immunol *20*, 337-349. 10.1038/s41590-018-0311-z.
Germar, K., Dose, M., Konstantinou, T., Zhang, J., Wang, H., Lobry, C., Arnett, K.L., Blacklow,
S.C., Aifantis, I., Aster, J.C., and Gounari, F. (2011). T-cell factor 1 is a gatekeeper for T-cell
specification in response to Notch signaling. Proc Natl Acad Sci U S A *108*, 20060-20065.
10.1073/pnas.1110230108.
Giralt, S., and Bishop, M.R. (2009). Principles and overview of allogeneic hematopoietic stem

cell transplantation. Cancer Treat Res *144*, 1-21. 10.1007/978-0-387-78580-6_1.

Gounari, F., and Khazaie, K. (2022). TCF-1: a maverick in T cell development and function. Nat
Immunol. 10.1038/s41590-022-01194-2.

Guinan, E.C., Boussiotis, V.A., Neuberg, D., Brennan, L.L., Hirano, N., Nadler, L.M., and Gribben,
J.G. (1999). Transplantation of anergic histoincompatible bone marrow allografts. N Engl J Med
340, 1704-1714. 10.1056/NEJM199906033402202.

Hall, E., and Shenoy, S. (2019). Hematopoietic Stem Cell Transplantation: A Neonatal
Perspective. Neoreviews *20*, e336-e345. 10.1542/neo.20-6-e336.

He, R., Hou, S., Liu, C., Zhang, A., Bai, Q., Han, M., Yang, Y., Wei, G., Shen, T., Yang, X., et al.
(2016). Follicular CXCR5- expressing CD8(+) T cells curtail chronic viral infection. Nature *537*,
412-428. 10.1038/nature19317.

Hoffmann, P., Ermann, J., Edinger, M., Fathman, C.G., and Strober, S. (2002). Donor-type
CD4(+)CD25(+) regulatory T cells suppress lethal acute graft-versus-host disease after allogeneic
bone marrow transplantation. J Exp Med *196*, 389-399. 10.1084/jem.20020399.

Hu, J., Batth, I.S., Xia, X., and Li, S. (2016). Regulation of NKG2D(+)CD8(+) T-cell-mediated
antitumor immune surveillance: Identification of a novel CD28 activation-mediated, STAT3
phosphorylation-dependent mechanism. Oncoimmunology *5*, e1252012.
10.1080/2162402X.2016.1252012.

Huang da, W., Sherman, B.T., and Lempicki, R.A. (2009a). Bioinformatics enrichment tools:
paths toward the comprehensive functional analysis of large gene lists. Nucleic Acids Res *37*, 1-
13. 10.1093/nar/gkn923.

Huang da, W., Sherman, B.T., and Lempicki, R.A. (2009b). Systematic and integrative analysis of
large gene lists using DAVID bioinformatics resources. Nat Protoc *4*, 44-57.
10.1038/nprot.2008.211.

Huang, W., Hu, J., and August, A. (2013). Cutting edge: innate memory CD8+ T cells are distinct
from homeostatic expanded CD8+ T cells and rapidly respond to primary antigenic stimuli. J
Immunol *190*, 2490-2494. 10.4049/jimmunol.1202988.

Huang, W., Mo, W., Jiang, J., Chao, N.J., and Chen, B.J. (2019). Donor Allospecific CD44(high)
Central Memory T Cells Have Decreased Ability to Mediate Graft-vs.-Host Disease. Front
Immunol *10*, 624. 10.3389/fimmu.2019.00624.

Im, S.J., Hashimoto, M., Gerner, M.Y., Lee, J., Kissick, H.T., Burger, M.C., Shan, Q., Hale, J.S., Lee,
J., Nasti, T.H., et al. (2016). Defining CD8+ T cells that provide the proliferative burst after PD-1
therapy. Nature *537*, 417-421. 10.1038/nature19330.

Jiang, H., Fu, D., Bidgoli, A., and Paczesny, S. (2021). T Cell Subsets in Graft Versus Host Disease
and Graft Versus Tumor. Front Immunol *12*, 761448. 10.3389/fimmu.2021.761448.

Jiang, T., Piao, D., Zhu, A., and Jiang, H. (2014). Changes in T lymphocyte subsets in mice with
CT26 colon tumors after treatment with donor lymphocyte infusion. Tumour Biol *35*, 5599-
5605. 10.1007/s13277-014-1740-4.

Johnson, J.L., Georgakilas, G., Petrovic, J., Kurachi, M., Cai, S., Harly, C., Pear, W.S., Bhandoola,
A., Wherry, E.J., and Vahedi, G. (2018). Lineage-Determining Transcription Factor TCF-1 Initiates
the Epigenetic Identity of T Cells. Immunity *48*, 243-257 e210. 10.1016/j.immuni.2018.01.012.

Ju, X.P., Xu, B., Xiao, Z.P., Li, J.Y., Chen, L., Lu, S.Q., and Huang, Z.X. (2005). Cytokine expression
during acute graft-versus-host disease after allogeneic peripheral stem cell transplantation.
Bone Marrow Transplant *35*, 1179-1186. 10.1038/sj.bmt.1704972.

Karimi, M.A., Bryson, J.L., Richman, L.P., Fesnak, A.D., Leichner, T.M., Satake, A., Vonderheide,
R.H., Raulet, D.H., Reshef, R., and Kambayashi, T. (2015). NKG2D expression by CD8+ T cells
contributes to GVHD and GVT effects in a murine model of allogeneic HSCT. Blood *125*, 3655-
3663. 10.1182/blood-2015-02-629006.
Karimi, M.A., Lee, E., Bachmann, M.H., Salicioni, A.M., Behrens, E.M., Kambayashi, T., and
Baldwin, C.L. (2014). Measuring cytotoxicity by bioluminescence imaging outperforms the
standard chromium-51 release assay. PLoS One *9*, e89357. 10.1371/journal.pone.0089357.
Khan, O., Giles, J.R., McDonald, S., Manne, S., Ngiow, S.F., Patel, K.P., Werner, M.T., Huang, A.C.,
Alexander, K.A., Wu, J.E., et al. (2019). TOX transcriptionally and epigenetically programs CD8(+)
T cell exhaustion. Nature *571*, 211-218. 10.1038/s41586-019-1325-x.
Kiekens, L., Van Loocke, W., Taveirne, S., Wahlen, S., Persyn, E., Van Ammel, E., De Vos, Z.,
Matthys, P., Van Nieuwerburgh, F., Taghon, T., et al. (2021). T-BET and EOMES Accelerate and
Enhance Functional Differentiation of Human Natural Killer Cells. Front Immunol *12*, 732511.
10.3389/fimmu.2021.732511.
Kim, C., Jin, J., Weyand, C.M., and Goronzy, J.J. (2020). The Transcription Factor TCF1 in T Cell
Differentiation and Aging. Int J Mol Sci *21*. 10.3390/ijms21186497.
Kurtulus, S., Madi, A., Escobar, G., Klapholz, M., Nyman, J., Christian, E., Pawlak, M., Dionne, D.,
Xia, J., Rozenblatt-Rosen, O., et al. (2019). Checkpoint Blockade Immunotherapy Induces
Dynamic Changes in PD-1(-)CD8(+) Tumor-Infiltrating T Cells. Immunity *50*, 181-194 e186.
10.1016/j.immuni.2018.11.014.
LaFleur, M.W., Nguyen, T.H., Coxe, M.A., Miller, B.C., Yates, K.B., Gillis, J.E., Sen, D.R., Gaudiano,
E.F., Al Abosy, R., Freeman, G.J., et al. (2019). PTPN2 regulates the generation of exhausted
CD8(+) T cell subpopulations and restrains tumor immunity. Nat Immunol *20*, 1335-1347.
10.1038/s41590-019-0480-4.
Larson, J.H., Marron, B.M., Beever, J.E., Roe, B.A., and Lewin, H.A. (2006). Genomic organization
and evolution of the ULBP genes in cattle. BMC Genomics *7*, 227. 10.1186/1471-2164-7-227.
Law, C.W., Chen, Y., Shi, W., and Smyth, G.K. (2014). voom: Precision weights unlock linear
model analysis tools for RNA-seq read counts. Genome Biol *15*, R29. 10.1186/gb-2014-15-2-r29.
Liu, L., Chen, X., Jin, H.M., Zhao, S.S., Zhu, Y., Qian, S.X., and Wu, Y.J. (2022). [The Expression and
Function of NK Cells in Patients with Acute Myeloid Leukemia]. Zhongguo Shi Yan Xue Ye Xue Za
Zhi *30*, 49-55. 10.19746/j.cnki.issn.1009-2137.2022.01.009.
Lugli, E., Dominguez, M.H., Gattinoni, L., Chattopadhyay, P.K., Bolton, D.L., Song, K., Klatt, N.R.,
Brenchley, J.M., Vaccari, M., Gostick, E., et al. (2013). Superior T memory stem cell persistence
supports long-lived T cell memory. J Clin Invest *123*, 594-599. 10.1172/JCI66327.
Lynch Kelly, D., Lyon, D.E., Ameringer, S.A., and Elswick, R.K. (2015). Symptoms, Cytokines, and
Quality of Life in Patients Diagnosed with Chronic Graft-Versus-Host Disease Following
Allogeneic Hematopoietic Stem Cell Transplantation. Oncol Nurs Forum *42*, 265-275.
10.1188/15.ONF.265-275.
Ma, J., Wang, R., Fang, X., and Sun, Z. (2012). beta-catenin/TCF-1 pathway in T cell development
and differentiation. J Neuroimmune Pharmacol *7*, 750-762. 10.1007/s11481-012-9367-y.
Maasho, K., Opoku-Anane, J., Marusina, A.I., Coligan, J.E., and Borrego, F. (2005). NKG2D is a
costimulatory receptor for human naive CD8+ T cells. J Immunol *174*, 4480-4484.
10.4049/jimmunol.174.8.4480.
Mammadli, M., Harris, R., Mahmudlu, S., Verma, A., May, A., Dhawan, R., Waickman, A.T., Sen,

J.M., August, A., and Karimi, M. (2021a). Human Wnt/beta-Catenin Regulates Alloimmune Signaling during Allogeneic Transplantation. Cancers (Basel) *13*. 10.3390/cancers13153798.

Mammadli, M., Harris, R., Mahmudlu, S., Verma, A., May, A., Dhawan, R., Waickman, A.T., Sen, J.M., and Karimi, M. (2021b). Wnt/β-catenin regulates alloreactive T cells for the treatment of hematological malignancies. bioRxiv, 2021.2004.2012.439538. 10.1101/2021.04.12.439538.

Mammadli, M., Harris, R., Suo, L., May, A., Gentile, T., Waickman, A.T., Bah, A., August, A., Nurmemmedov, E., and Karimi, M. (2021c). Interleukin-2-inducible T-cell kinase (Itk) signaling regulates potent noncanonical regulatory T cells. Clin Transl Med *11*, e625. 10.1002/ctm2.625.

Mammadli, M., Huang, W., Harris, R., Sultana, A., Cheng, Y., Tong, W., Pu, J., Gentile, T., Dsouza, S., Yang, Q., et al. (2020). Targeting Interleukin-2-Inducible T-Cell Kinase (ITK) Differentiates GVL and GVHD in Allo-HSCT. Front Immunol *11*, 593863. 10.3389/fimmu.2020.593863.

Mammadli, M., Huang, W., Harris, R., Xiong, H., Weeks, S., May, A., Gentile, T., Henty-Ridilla, J., Waickman, A.T., August, A., et al. (2021d). Targeting SLP76:ITK interaction separates GVHD from GVL in allo-HSCT. iScience *24*, 102286. https://doi.org/10.1016/j.isci.2021.102286.

Mavers, M., and Bertaina, A. (2018). High-Risk Leukemia: Past, Present, and Future Role of NK Cells. J Immunol Res *2018*, 1586905. 10.1155/2018/1586905.

Miller, B.C., Sen, D.R., Al Abosy, R., Bi, K., Virkud, Y.V., LaFleur, M.W., Yates, K.B., Lako, A., Felt, K., Naik, G.S., et al. (2019). Author Correction: Subsets of exhausted CD8(+) T cells differentially mediate tumor control and respond to checkpoint blockade. Nat Immunol *20*, 1556. 10.1038/s41590-019-0528-5.

Mohty, M., Blaise, D., Faucher, C., Vey, N., Bouabdallah, R., Stoppa, A.M., Viret, F., Gravis, G., Olive, D., and Gaugler, B. (2005). Inflammatory cytokines and acute graft-versus-host disease after reduced-intensity conditioning allogeneic stem cell transplantation. Blood *106*, 4407-4411. 10.1182/blood-2005-07-2919.

Nakajima, Y., Chamoto, K., Oura, T., and Honjo, T. (2021). Critical role of the CD44(low)CD62L(low) CD8(+) T cell subset in restoring antitumor immunity in aged mice. Proc Natl Acad Sci U S A *118*. 10.1073/pnas.2103730118.

Nishimura, R., Baker, J., Beilhack, A., Zeiser, R., Olson, J.A., Sega, E.I., Karimi, M., and Negrin, R.S. (2008). In vivo trafficking and survival of cytokine-induced killer cells resulting in minimal GVHD with retention of antitumor activity. Blood *112*, 2563-2574. 10.1182/blood-2007-06-092817.

Paley, M.A., and Wherry, E.J. (2010). TCF-1 flips the switch on Eomes. Immunity *33*, 145-147. 10.1016/j.immuni.2010.08.008.

Philip, M., Fairchild, L., Sun, L., Horste, E.L., Camara, S., Shakiba, M., Scott, A.C., Viale, A., Lauer, P., Merghoub, T., et al. (2017). Chromatin states define tumour-specific T cell dysfunction and reprogramming. Nature *545*, 452-456. 10.1038/nature22367.

Prajapati, K., Perez, C., Rojas, L.B.P., Burke, B., and Guevara-Patino, J.A. (2018). Functions of NKG2D in CD8(+) T cells: an opportunity for immunotherapy. Cell Mol Immunol *15*, 470-479. 10.1038/cmi.2017.161.

Presotto, D., Erdes, E., Duong, M.N., Allard, M., Regamey, P.O., Quadroni, M., Doucey, M.A., Rufer, N., and Hebeisen, M. (2017). Fine-Tuning of Optimal TCR Signaling in Tumor-Redirected CD8 T Cells by Distinct TCR Affinity-Mediated Mechanisms. Front Immunol *8*, 1564. 10.3389/fimmu.2017.01564.

Raulet, D.H. (2003). Roles of the NKG2D immunoreceptor and its ligands. Nat Rev Immunol *3*,

781-790. 10.1038/nri1199.

Raulet, D.H., Gasser, S., Gowen, B.G., Deng, W., and Jung, H. (2013). Regulation of ligands for
the NKG2D activating receptor. Annu Rev Immunol *31*, 413-441. 10.1146/annurev-immunol-
032712-095951.

Reddy, P., and Ferrara, J.L.M. (2008). Mouse models of graft-versus-host disease. In StemBook.
10.3824/stembook.1.36.1.

Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015). limma
powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic
Acids Res *43*, e47. 10.1093/nar/gkv007.

Schietinger, A., Philip, M., Krisnawan, V.E., Chiu, E.Y., Delrow, J.J., Basom, R.S., Lauer, P.,
Brockstedt, D.G., Knoblaugh, S.E., Hammerling, G.J., et al. (2016). Tumor-Specific T Cell
Dysfunction Is a Dynamic Antigen-Driven Differentiation Program Initiated Early during
Tumorigenesis. Immunity *45*, 389-401. 10.1016/j.immuni.2016.07.011.

Scott, A.C., Dundar, F., Zumbo, P., Chandran, S.S., Klebanoff, C.A., Shakiba, M., Trivedi, P.,
Menocal, L., Appleby, H., Camara, S., et al. (2019). TOX is a critical regulator of tumour-specific T
cell differentiation. Nature *571*, 270-274. 10.1038/s41586-019-1324-y.

Seif, F., Khoshmirsafa, M., Aazami, H., Mohsenzadegan, M., Sedighi, G., and Bahar, M. (2017).
The role of JAK-STAT signaling pathway and its regulators in the fate of T helper cells. Cell
Commun Signal *15*, 23. 10.1186/s12964-017-0177-y.

Seo, H., Chen, J., Gonzalez-Avalos, E., Samaniego-Castruita, D., Das, A., Wang, Y.H., Lopez-
Moyado, I.F., Georges, R.O., Zhang, W., Onodera, A., et al. (2019). TOX and TOX2 transcription
factors cooperate with NR4A transcription factors to impose CD8(+) T cell exhaustion. Proc Natl
Acad Sci U S A *116*, 12410-12415. 10.1073/pnas.1905675116.

Shi, J., Chi, S., Xue, J., Yang, J., Li, F., and Liu, X. (2016). Emerging Role and Therapeutic
Implication of Wnt Signaling Pathways in Autoimmune Diseases. J Immunol Res *2016*, 9392132.
10.1155/2016/9392132.

Siddiqui, I., Schaeuble, K., Chennupati, V., Fuertes Marraco, S.A., Calderon-Copete, S., Pais
Ferreira, D., Carmona, S.J., Scarpellino, L., Gfeller, D., Pradervand, S., et al. (2019). Intratumoral
Tcf1(+)PD-1(+)CD8(+) T Cells with Stem-like Properties Promote Tumor Control in Response to
Vaccination and Checkpoint Blockade Immunotherapy. Immunity *50*, 195-211 e110.
10.1016/j.immuni.2018.12.021.

Sobecki, M., Mrouj, K., Camasses, A., Parisis, N., Nicolas, E., Lleres, D., Gerbe, F., Prieto, S.,
Krasinska, L., David, A., et al. (2016). The cell proliferation antigen Ki-67 organises
heterochromatin. Elife *5*, e13722. 10.7554/eLife.13722.

Steinke, F.C., Yu, S., Zhou, X., He, B., Yang, W., Zhou, B., Kawamoto, H., Zhu, J., Tan, K., and Xue,
H.H. (2014). TCF-1 and LEF-1 act upstream of Th-POK to promote the CD4(+) T cell fate and
interact with Runx3 to silence Cd4 in CD8(+) T cells. Nat Immunol *15*, 646-656. 10.1038/ni.2897.

Stojanovic, A., Correia, M.P., and Cerwenka, A. (2018). The NKG2D/NKG2DL Axis in the Crosstalk
Between Lymphoid and Myeloid Cells in Health and Disease. Front Immunol *9*, 827.
10.3389/fimmu.2018.00827.

Tugues, S., Amorim, A., Spath, S., Martin-Blondel, G., Schreiner, B., De Feo, D., Lutz, M.,
Guscetti, F., Apostolova, P., Haftmann, C., et al. (2018). Graft-versus-host disease, but not graft-
versus-leukemia immunity, is mediated by GM-CSF-licensed myeloid cells. Sci Transl Med *10*.
10.1126/scitranslmed.aat8410.

Utzschneider, D.T., Charmoy, M., Chennupati, V., Pousse, L., Ferreira, D.P., Calderon-Copete, S.,
Danilo, M., Alfei, F., Hofmann, M., Wieland, D., et al. (2016). T Cell Factor 1-Expressing Memory-
like CD8(+) T Cells Sustain the Immune Response to Chronic Viral Infections. Immunity *45*, 415-
427. 10.1016/j.immuni.2016.07.021.
Utzschneider, D.T., Legat, A., Fuertes Marraco, S.A., Carrie, L., Luescher, I., Speiser, D.E., and
Zehn, D. (2013). T cells maintain an exhausted phenotype after antigen withdrawal and
population reexpansion. Nat Immunol *14*, 603-610. 10.1038/ni.2606.
Villarroel, V.A., Okiyama, N., Tsuji, G., Linton, J.T., and Katz, S.I. (2014). CXCR3-mediated skin
homing of autoreactive CD8 T cells is a key determinant in murine graft-versus-host disease. J
Invest Dermatol *134*, 1552-1560. 10.1038/jid.2014.2.
Wang, S., Wu, Q., Chen, T., Su, R., Pan, C., Qian, J., Huang, H., Yin, S., Xie, H., Zhou, L., and
Zheng, S. (2022). Blocking CD47 promotes anti-tumor immunity through CD103+ dendritic cell-
NK cell axis in murine hepatocellular carcinoma model. J Hepatol. 10.1016/j.jhep.2022.03.011.
Wang, X., He, Q., Shen, H., Xia, A., Tian, W., Yu, W., and Sun, B. (2019a). TOX promotes the
exhaustion of antitumor CD8(+) T cells by preventing PD1 degradation in hepatocellular
carcinoma. J Hepatol *71*, 731-741. 10.1016/j.jhep.2019.05.015.
Wang, Y., Hu, J., Li, Y., Xiao, M., Wang, H., Tian, Q., Li, Z., Tang, J., Hu, L., Tan, Y., et al. (2019b).
The Transcription Factor TCF1 Preserves the Effector Function of Exhausted CD8 T Cells During
Chronic Viral Infection. Front Immunol *10*, 169. 10.3389/fimmu.2019.00169.
Weber, B.N., Chi, A.W., Chavez, A., Yashiro-Ohtani, Y., Yang, Q., Shestova, O., and Bhandoola, A.
(2011). A critical role for TCF-1 in T-lineage specification and differentiation. Nature *476*, 63-68.
10.1038/nature10279.
Weeks, S., Harris, R., and Karimi, M. (2021). Targeting ITK signaling for T cell-mediated diseases.
iScience *24*, 102842. 10.1016/j.isci.2021.102842.
Wensveen, F.M., Jelencic, V., and Polic, B. (2018). NKG2D: A Master Regulator of Immune Cell
Responsiveness. Front Immunol *9*, 441. 10.3389/fimmu.2018.00441.
Wu, T., Ji, Y., Moseman, E.A., Xu, H.C., Manglani, M., Kirby, M., Anderson, S.M., Handon, R.,
Kenyon, E., Elkahloun, A., et al. (2016). The TCF1-Bcl6 axis counteracts type I interferon to
repress exhaustion and maintain T cell stemness. Sci Immunol *1*. 10.1126/sciimmunol.aai8593.
Xing, S., Li, F., Zeng, Z., Zhao, Y., Yu, S., Shan, Q., Li, Y., Phillips, F.C., Maina, P.K., Qi, H.H., et al.
(2016). Tcf1 and Lef1 transcription factors establish CD8(+) T cell identity through intrinsic
HDAC activity. Nat Immunol *17*, 695-703. 10.1038/ni.3456.
Xu, W., Zhao, X., Wang, X., Feng, H., Gou, M., Jin, W., Wang, X., Liu, X., and Dong, C. (2019). The
Transcription Factor Tox2 Drives T Follicular Helper Cell Development via Regulating Chromatin
Accessibility. Immunity *51*, 826-839 e825. 10.1016/j.immuni.2019.10.006.
Yang, B.H., Wang, K., Wan, S., Liang, Y., Yuan, X., Dong, Y., Cho, S., Xu, W., Jepsen, K., Feng, G.S.,
et al. (2019). TCF1 and LEF1 Control Treg Competitive Survival and Tfr Development to Prevent
Autoimmune Diseases. Cell Rep *27*, 3629-3645 e3626. 10.1016/j.celrep.2019.05.061.
Yu, Q., Sharma, A., and Sen, J.M. (2010). TCF1 and beta-catenin regulate T cell development and
function. Immunol Res *47*, 45-55. 10.1007/s12026-009-8137-2.
Zander, R., Schauder, D., Xin, G., Nguyen, C., Wu, X., Zajac, A., and Cui, W. (2019). CD4(+) T Cell
Help Is Required for the Formation of a Cytolytic CD8(+) T Cell Subset that Protects against
Chronic Infection and Cancer. Immunity *51*, 1028-1042 e1024. 10.1016/j.immuni.2019.10.009.
Zheng, H., Matte-Martone, C., Jain, D., McNiff, J., and Shlomchik, W.D. (2009). Central memory

CD8+ T cells induce graft-versus-host disease and mediate graft-versus-leukemia. J Immunol *182*, 5938-5948. 10.4049/jimmunol.0802212.

Zhou, X., Yu, S., Zhao, D.M., Harty, J.T., Badovinac, V.P., and Xue, H.H. (2010). Differentiation and persistence of memory CD8(+) T cells depend on T cell factor 1. Immunity *33*, 229-240. 10.1016/j.immuni.2010.08.002.

Zhu, Y., Ju, S., Chen, E., Dai, S., Li, C., Morel, P., Liu, L., Zhang, X., and Lu, B. (2010). T-bet and eomesodermin are required for T cell-mediated antitumor immune responses. J Immunol *185*, 3174-3183. 10.4049/jimmunol.1000749.

Zuniga-Pflucker, J.C. (2004). T-cell development made simple. Nat Rev Immunol *4*, 67-72. 10.1038/nri1257.

Huang da, W., Sherman, B.T., and Lempicki, R.A. (2009). Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. Nat Protoc *4*, 44-57. 10.1038/nprot.2008.211.

Karimi, M.A., Aguilar, O., Zou, B., Bachmann, M.H., Carlyle, J.R., Baldwin, C.L., and Kambayashi, T. (2014a). A truncated human NKG2D splice isoform negatively regulates NKG2D-mediated function. J Immunol *193*, 2764-2771. 10.4049/jimmunol.1400920.

Karimi, M.A., Bryson, J.L., Richman, L.P., Fesnak, A.D., Leichner, T.M., Satake, A., Vonderheide, R.H., Raulet, D.H., Reshef, R., and Kambayashi, T. (2015). NKG2D expression by CD8+ T cells contributes to GVHD and GVT effects in a murine model of allogeneic HSCT. Blood *125*, 3655-3663. 10.1182/blood-2015-02-629006.

Karimi, M.A., Lee, E., Bachmann, M.H., Salicioni, A.M., Behrens, E.M., Kambayashi, T., and Baldwin, C.L. (2014b). Measuring cytotoxicity by bioluminescence imaging outperforms the standard chromium-51 release assay. PLoS One *9*, e89357. 10.1371/journal.pone.0089357.

Mammadli, M., Harris, R., Mahmudlu, S., Verma, A., May, A., Dhawan, R., Waickman, A.T., Sen, J.M., August, A., and Karimi, M. (2021a). Human Wnt/beta-Catenin Regulates Alloimmune Signaling during Allogeneic Transplantation. Cancers (Basel) *13*. 10.3390/cancers13153798.

Mammadli, M., Harris, R., Suo, L., May, A., Gentile, T., Waickman, A.T., Bah, A., August, A., Nurmemmedov, E., and Karimi, M. (2021b). Interleukin-2-inducible T-cell kinase (Itk) signaling regulates potent noncanonical regulatory T cells. Clin Transl Med *11*, e625. 10.1002/ctm2.625.

Mammadli, M., Huang, W., Harris, R., Sultana, A., Cheng, Y., Tong, W., Pu, J., Gentile, T., Dsouza, S., Yang, Q., et al. (2020). Targeting Interleukin-2-Inducible T-Cell Kinase (ITK) Differentiates GVL and GVHD in Allo-HSCT. Front Immunol *11*, 593863. 10.3389/fimmu.2020.593863.

Mammadli, M., Huang, W., Harris, R., Xiong, H., Weeks, S., May, A., Gentile, T., Henty-Ridilla, J., Waickman, A.T., August, A., et al. (2021c). Targeting SLP76:ITK interaction separates GVHD from GVL in allo-HSCT. iScience *24*, 102286. https://doi.org/10.1016/j.isci.2021.102286.

Sherman, B.T., Hao, M., Qiu, J., Jiao, X., Baseler, M.W., Lane, H.C., Imamichi, T., and Chang, W. (2022). DAVID: a web server for functional enrichment analysis and functional annotation of gene lists (2021 update). Nucleic Acids Res. 10.1093/nar/gkac194.

1483

Anti-Tumor Function

↓ TCR signaling

CD8

↑ NKG2D

LRP

Wnt

Frizzled

LAT ↓

↓ LCK   ↓ ITK

↓ PLC-γ

↓ ERK1/II

Dishevelled

Vav1   DAP10

GrB2

↓ JAK2

↓ JAK3

↓ STAT3

Required for
TCR signaling

↑ Eomes   ↑ T-bet

Required for
anti-tumor
responses

↓ p65   β-Catenin

TCF-1 ✗

T-bet Eomes

Target
Genes

Molecular Mechanisms

Gene expression

LCK, LAT, ITK, PLC-g, p65,
Erk1/II, JAK2, JAK3, STAT3

Significantly decreased

Cellular Mechanisms

Inflammatory
Cytokines

Activation

Apoptosis

Clinical Impact

Less damage to
GVHD target organs

Skin

SI

Liver

Anti-Tumor Response

Increased Eomes, T-bet, and surface NKG2D expression