Symptomatic and Neurotrophic Effects of GABAA Receptor Positive Allosteric Modulation in a Mouse Model of Chronic Stress

Ashley Bernardo[1], PhD; Philip Lee[2], BSc; Michael Marcotte[1], MSc; Md Yeunus Mian[3], MSc; Sepideh Rezvanian[3], MSc; Dishary Sharmin[3], MSc; Aleksandra Kovačević[4], Msc; Miroslav Savić[4], PhD; James M. Cook[3], PhD; Etienne Sibille[1,2,5*], PhD and Thomas D. Prevot[1,5*], PhD.

[1]Campbell Family Mental Health Research Institute of CAMH, Toronto, Canada

[2]Department of Pharmacology and Toxicology, University of Toronto, Toronto, Canada

[3]Department of Chemistry and Biochemistry, University of Wisconsin–Milwaukee, Milwaukee, USA

[4]Department of Pharmacology, Faculty of Pharmacy, University of Belgrade, Belgrade, Serbia

[5]Department of Psychiatry, University of Toronto, Toronto, Canada

*Corresponding Authors

Thomas D. Prevot, Ph.D., CAMH, 250 College Street, room 131, Toronto, ON M5T 1R8, Canada

Tel: 416-535-8501, ext. 33785; E-mail: thomas.prevot@camh.ca

Etienne Sibille, Ph.D. CAMH, 250 College Street, Room 134, Toronto, ON M5T 1R8, Canada

Tel: 416-535-8501, ext. 36571, E-mail: Etienne.sibille@camh.ca

Introduction to Discussion; 3919 words

Abstract:

Chronic stress is a risk factor for Major depressive disorder (MDD), and in rodents, it recapitulates human behavioral, cellular and molecular changes. In MDD and after chronic stress, neuronal dysfunctions and deficits in GABAergic signaling are observed and responsible for symptom severity. GABA signals predominantly through GABAA receptors (GABAA-R) composed of various subunit types that relate to downstream outcomes. Activity at α2-GABAA-Rs contributes to anxiolytic properties, α5-GABAA-Rs to cognitive functions, and α1-GABAA-Rs to sedation. Therefore, a therapy aiming at increasing α2- and α5-GABAA-Rs activity, but devoid of α1-GABAA-R activity, has potential to address several symptomologies of depression while avoiding side effects. This study investigated the activity profiles and behavioral efficacy of two molecules enantiomers of each other (GL-II-73 and GL-I-54), separately and as a racemic mixture (GL-RM), and potential disease-modifying effects on neuronal morphology. Results confirm GL-I-54 and GL-II-73 exert positive allosteric modulation at the α2-, α3-, α5-GABAA-Rs and α5-containing GABAA-Rs, respectively, and have anti-depressant and pro-cognitive effects independently. Using unpredictable chronic mild stress (UCMS) in male and female mice (n=12/group), we show that acute and chronic administration of GL-RM combined the anti-depressant and pro-cognitive effects of each enantiomer, although at lower doses avoiding sedation. Morphology studies showed reversal of spine density loss caused by UCMS after chronic GL-RM treatment at apical and basal dendrites of the PFC and CA1. Together, these results support using a racemic mixture with combined α2-, α3-, α5-GABAA-R profile to reverse chronic stress-induced mood symptoms, cognitive deficits, and with anti-stress neurotrophic effects.

Introduction

Chronic stress is a risk factor for psychiatric disorders including Major Depressive Disorder (MDD), displaying mood and cognitive symptoms [1-3]. Interventions to treat depression and other stress-related psychopathologies have focused on the monoaminergic system [4] with limited efficacy on mood symptoms, no efficacy on cognitive symptoms, and no effect on structural features of brain cells, highlighting unmet clinical needs. Recent studies have focused on the glutamatergic system [5], with promising effect on mood, cognition and cell structure [6, 7]. Similar efficacies were reported with compounds targeting the GABAergic system [8].

Mood and cognitive symptoms result from cellular dysfunctions, altered communication between cells and neuronal atrophy [9-13]. Transcranial magnetic stimulation in MDD patients shows reduced GABAergic function (i.e., cortical inhibition) [14, 15], contributing to impaired excitation/inhibition balance in MDD [16, 17]. Reduced GABA levels are reported in the occipital cortex [18-24], prefrontal cortex (PFC) [25] and anterior cingulate cortex [18, 25-27]. In the PFC, synaptic densities and expression of synaptic function-related genes are reduced [12, 28]. Chronic stress-related disorders [13, 29] and animal models report similar findings [30-32] and demonstrate critical links between prefrontal and hippocampal functions [33]. Chronic stress reduces dendritic length and spine density in the PFC and hippocampus (HPC), likely contributing to cognitive deficits. Of approved treatments, only the NMDA receptor antagonist, ketamine, has shown efficacy at increasing spine density and dendritic complexity in mice, along with antidepressant properties [34]. The neurotrophic effects of ketamine are hypothesized to act through BDNF-TrkB signaling while its fast-acting antidepressant properties are suggested to be initially mediated by activity on GABAergic neurons followed by long-term GABA and glutamate changes [35], highlighting a potential role for GABA in these therapeutic effects.

GABA signals through ionotropic GABAA receptors (GABAA-Rs) and metabotropic GABAB receptors. GABAA-Rs are pentameric ion channels, formed by a combination of 19

subunit subtypes [36]. Drugs acting on GABAA-Rs exist (benzodiazepines or imidazodiazepines),

but their use is limited because of side effects (sedation) due to broad activity at several receptor

subunits, including α1-GABAA-Rs [37-39]. Studies demonstrated α2-GABAA-Rs, strongly present

in the amygdala [40-42] and HPC [40, 43, 44], are linked to regulation of anxious and depressive

phenotypes [45-49]. The role of α3-GABAA-Rs are not fully characterized, but appear linked to

behavioral despair in mice [50]. Other studies link α5-GABAA-Rs with cognitive performances

[51-53], aligned with their preferential expression in cognitive processing regions (HPC and

mPFC) [40, 54]. Our group and others showed targeting α5-GABAA-Rs with a positive allosteric

modulator (PAM) reverses stress- [52] and age-[55, 56] related cognitive deficits, and chronically

reverses age-related neuronal atrophy [53]. Potentiation of α2- and α5-GABAA-Rs, while limiting

potentiation of α1-GABAA-R, is hypothesized to improve mood and cognitive symptoms while

reducing side effects [37, 57], altogether suggesting potential for treatment of symptomatic and

morphological alterations in chronic stress and MDD.

Previously, we showed the α5-PAM GL-II-73, has pro-cognitive, anxiolytic and anti-

depressant properties at 10mg/kg IP but not at 5mg/kg IP [52], and neurotrophic effects at

30mg/kg PO [56] in mice. However, it remained unknown if its enantiomer, GL-I-54, has similar

or complementary properties. Here, we investigated the similarities and/or complementarities of

receptor selectivity and behavioral efficacy profile of GL-I-54 and GL-II-73 in stressed mice. We

then investigated the symptomatic and neurotrophic potentials of a racemic mixture of both

compounds in a mouse model of chronic stress.

Materials and Methods

Details provided in the supplementary material.

Compounds

GL-I-54 and GL-II-73 were synthesized as in [58], with >99% chemical and optical purity. GL-RM

is a 1:1 mix of each compound.

Electrophysiology

Electrophysiological recordings used HEK-293T cell line transfected with full-length cDNAs for

human GABAA-R subtypes α(1/2/3/4/5)β3γ2 [59, 60]. PubMed Protein Sequence repository for

GABAA-R subunits reports the binding pocket formed by α1/2/3/5 and γ2 is 100% conserved

between mouse and human, predicting similar activity and selectivity across species. Current

recordings were performed in absence or presence (5µM) of GABA. GL-I-54 or GL-II-73 was

applied at 0.033-33.33µM concentrations.

Animals

Six independent cohorts of 8-week old C57BL/6 mice (50% female) were obtained from Military

Medical Academy (Belgrade, Serbia) or Jackson Laboratories (Stock#000664; MA, USA),

measuring: pharmacokinetics (#1; n=25), individual enantiomers on elevated plus maze and

forced swim test (#2; n=24), effect of GL-I-54 on Y-maze (#3; n=42), GL-I-54 side effects in the

rotarod (#4, n=18), and acute (#5; n=36) and chronic (#6; n=36) efficacy of GL-RM. Animals were

individually housed and maintained on a 12 hour light/dark cycle (7:00 ON, 19:00 OFF), with *ad*

libitum food and water.

Ethical Statement

107 All animal work was completed in accordance with Ontario Animals for Research Act (RSO 1990,

108 Chapter A.22), Canadian Council on Animal Care (CCAC), Ethical Council for the Protection of

109 Experimental Animals of the Ministry of Agriculture, Forestry and Water Management of the

110 Republic of Serbia, and was approved by the Institutes' Animal Care Committees.

Drug Administration

GL-I-54 (1, 3, 5, or 10mg/kg), GL-II-73 (5mg/kg) or GL-RM (10mg/kg) were diluted in vehicle

solution (85% distilled H_2O, 14% propylene glycol (Sigma Aldrich) and 1% Tween 80 (Sigma

Aldrich)) and administered intraperitoneal (IP), 30min before testing. For oral administration, GL-

RM was prepared in tap water (30mg/kg) and given through drinking water, considering 6mL

average daily fluid intake/animal.

GL-I-54 Pharmacokinetics

Mice were treated IP with 3mg/kg GL-I-54 and euthanized at different time points (5-720 min post-

injection) for brain and plasma quantification of ligand by ultraperformance LC-MS/MS (cassette

dosing) [61]. In vitro hydrolytic plasma stability of GL-I-54 was tested in vitro at 37°C, utilizing

blank mouse plasma spiked with GL-I-54 and internal standard, as in [62].

Plasma Protein and Brain Tissue Binding Studies

A rapid equilibrium dialysis assay determined the free fraction of GL-I-54 in mouse plasma and

brain tissue as in [63]. GL-I-54-free brain concentrations were calculated by multiplying the total

brain concentrations with the appropriate free fractions determined by rapid equilibrium dialysis.

Liver Microsomal Assay

Metabolic stability was tested in C57BL/6 Mouse liver microsomes at 2µM and 0.5mg/mL of matrix

concentration, as in [64]. NADPH solution, Microsomes/S9 and compounds were added to assay

plates. Plates were quenched at T0, 30, 60 and 120 min and assessed in refrigerated LC-MS/MS

autosampler. Intrinsic hepatic clearance calculations used 45mg microsomes/g of liver and 87.5g

liver/kg of body weight for mouse.

Stress Paradigms

Chronic restraint stress (CRS) was used with cohorts #2 and #3, consisting of placing mice in a

50mL Falcon™ Tube for 1hr twice daily for 1 week. Cohorts #5 and #6 were subjected to

Unpredictable Chronic Mild Stress (UCMS), using randomized mild stressors (twice daily) over 6

weeks. Weeks with behavioral testing applied one stressor per day after the behavioral testing (2

weeks). UCMS animals were housed in a separate room without environmental enrichment to

exacerbate effects of other stressors.

Behavioral Tests

Side effects were measured in the rotarod test in cohort #4, assessing locomotor coordination on

an accelerating rod. Latency to fall was measured over 6 trials. Mice in cohorts #2 were tested in

the elevated plus maze (EPM), measuring exploration of open arms for 10 min as a proxy of

anxiety [65], and in the forced swim test (FST), measuring time spent immobile for 6 min as a

proxy of antidepressant-like effect. In cohort #3, mice were exposed to CRS to induce a working

memory in the Y maze test, and the capacity of reversal of GL-I-54. In cohorts #5 and #6, mice

were subjected to UCMS, and each animal's coat state was scored (0=well-groomed, smooth

coat, 1=soiled coat or bald patches) weekly for seven coat regions, as in [66]. Behavioral screen

included: Anxiety-like behavior (EPM, open-field, OF; Novelty Suppressed Feeding, NSF; and the

Phenotyper Test [65], PT), anhedonia (Sucrose Consumption; SC), antidepressant-like activity

(FST) and cognitive function (Y-maze alternation task [52, 56, 67]).

Tissue Collection and Golgi Staining

Cohort #6 was euthanized by cervical dislocation 24hr after last behavioral test. Brains were

immersed in Golgi-Cox staining solution, and assigned a unique identifier (4 brains/group). Brains

were shipped to NeuroDigiTech (San Francisco CA, USA) for sectioning (100μm thickness),

mounting and blind quantification of basal and apical dendrites of PFC and CA1 pyramidal cells

using NeuroLucida v10 software. Neurons (n=6/animal) were analyzed for dendritic length, spine

number and spine density [68].

Statistics

Statistical analyses used GraphPad Prism 9. Electrophysiology data and compounds activity at

GABAA-R subunits were compared to 100% using t-tests. Other analyzes used two-way ANOVA

with concentration and subunit as co-factors, and Bonferroni post-hoc tests. Behavioral analyzes

used one-way or two-way ANOVA, and repeated measure ANOVA as relevant. Fishers PLSD

tests were used for post-hoc analyses. Spearman ranks correlation was used between

morphological features and behavioral outcomes. Statistical significance was considered $p<0.05$.

Data are presented as mean ± SEM.

Z-scores were calculated to assess consistency of anxiety-like and depressive-like behavioral

phenotypes across tests, referred to as z-emotionality using averaged z-scores of behavioral tests

as in [69]. Full calculations in supplementary.

Results

Statistical significances are provided in **Supplementary Tables**.

GL-I-54 and GL-II-73 exhibit different GABAA-R potentiation profiles but similar metabolic

stability

Both enantiomers were confirmed PAMs, not agonists (**Supplementary Fig. 1 and Fig. 1A-B**),

acting only in the presence of GABA. GL-I-54 significantly increased potentiation at α2-GABAA-

Rs (>0.03μM), α3-GABAA-Rs (>0.33μM), α1-GABAA-Rs (>1μM) and α5-GABAA-Rs (0.03, 0.33

and 3.33μM; **Fig. 1A**), with preferential activity at α3- and α2- over α1-, α4- and α5-GABAA-Rs

(Supplementary Table 1A).

GL-II-73 significantly increased potentiation at α5-GABAA-Rs (>1μM), α2-GABAA-Rs (3.33μM),

α3-GABAA-Rs (33.33μM) and α1-GABAA-Rs (33.33μM; **Fig. 1B**), with preferential activity at α5-

GABAA-Rs over α1-GABAA-Rs (>0.3μM), α2-GABAA-Rs (between 3.3 and 10μM), α3-GABAA-

Rs (between 1 and 10µM) and α_4-GABAA-Rs (>0.33µM) (**Supplementary Table 1B**), confirming

data previously published [52].

Pharmacokinetic profile, brain penetrance and stability

Pharmacokinetic profile of GL-I-54 dosed at 3mg/kg in mice is represented in **Fig. 1C**. Accounting

for the doses administered in respective studies, GL-I-54 and GL-II-73 exhibit comparable

pharmacokinetic patterns. As a representative example, 20min after the 3mg/kg dose, GL-I-54

achieved 311.75±6.72ng/g, while GL-II-73 reached 264.61±38.22ng/g (**Supplementary Fig. 2)**.

After appropriate adjustments (for the 5mg/kg dose, free fraction, brain tissue density and unit),

these close-to-maximum concentrations achievable in the mouse brain correspond to the 0.100-

0.330µM concentration range in electrophysiological recordings (calculation in **Supplementary**

Fig. 2). This implies that, in acute behavioral experiments with 5mg/kg dose, GL-I-54 likely

potentiates α_1, α_2, α_3 and α_5 GABAA-Rs, however, α_1 potentiation is mild. 5mg/kg of GL-II-73, on

the other hand, is fairly silent on all GABAA-Rs, with the potential to mildly activate α_2 and α_5

receptors; on α_1 and α_3 receptors, nevertheless, it behaves as a null modulator. Plasma and brain

free fractions of GL-I-54 were 22.57% and 11.49%, respectively, while, as previously reported for

GL-II-73 were 20.39% and 12.14% [52], respectively. Brain to plasma ratio (K_p) evaluates brain

penetrance (penetrant when K_p>0.04). GL-I-54 is brain penetrant (K_p=0.20), similarly to GL-II-73

(K_p=0.30), which is further supported by the ratio of unbound brain to unbound plasma ligand

concentrations values, $K_{p,uu}$ (0.10 vs 0.18, respectively). GL-I-54 displayed high in vitro metabolic

stability, although lower compared to GL-II-73; after 4hr of incubation in mouse plasma, the

fraction of remaining intact ligand was 78.41% (vs 98.88% of GL-II-73 [52]). In the liver microsomal

assay, GL-I-54 showed a 198min half-life with 70% remaining after 2hr. Intrinsic hepatic clearance

was 27.6mL/min/kg. GL-II-73 showed a longer half-life, 256min, with 75.6% remaining after 2hr.

Intrinsic hepatic clearance was 21.3mL/min/kg.

Acute administration of GL-I-54 displays antidepressant and procognitive effects in the chronic retrain stress model with locomotor side effects at higher dose

In mice subjected to CRS, anxiolytic and antidepressant properties of GL-I-54 (5mg/kg) and GL-II-73 (5mg/kg) were tested using the elevated plus maze (EPM), forced swim test (FST).Pro-cognitive properties were tested in the Y-maze (5mg/kg and 10mg/kg). After CRS, GL-I-54 and GL-II-73 had no effect on percent time in open arms of the EPM (**Fig. 1D**). In the FST, CRS increased time spent immobile, which was significantly reversed by GL-I-54 (**Fig. 1E**), suggesting antidepressant potential. In the Y-maze, CRS induced an alternation rate deficit that was reversed by GL-I-54 at 10mg/kg but not 5mg/kg (**Fig. 1F**), suggesting the capacity of reversing memory deficits. Locomotor side effects of GL-I-54 at 10mg/kg were tested in the rotarod (**Fig. 1G**). Decreasing latency to fall 60min after injection was found, suggesting sedation at this dose.

Acute administration of GL-RM displays antidepressant and procognitive effects in the UCMS model

Based on the selective effect on mood and no effect on working memory at 5mg/kg for GL-I-54 or GL-II-73 alone, we tested potential additive and low side effects using a 1:1 racemic mixture of 5mg/kg each, called GL-RM, in the UCMS model (**Fig. 2A**), with a final GL-RM dose of 10mg/kg. Weekly monitoring showed UCMS deteriorated coat state, not reversed by acute treatment, and weight gain remained unaffected (**Supplementary Fig. 3**).

In the EPM, there was no effect of UCMS or treatment on percent time in open arms (**Fig.2B**). There was a trend level effect of stress reducing the percent of entries into open arms but no effect of treatment (**Fig.2C**). NSF latency to approach found a significant interaction between UCMS and treatment, due to reduced latency in UCMS-GL-RM treated animals compared to Control-GL-RM and compared to UCMS-vehicle (**Fig.2D**), suggesting anxiolytic properties. Latency to bite was unaffected by UCMS or treatment (**Fig. 2E**). Phenotyper test and associated

measure of residual avoidance used in Prevot et al. [65] assess anxiety-like behavior weekly. The

residual avoidance score captures the time spent in the shelter, after a 1hr-light challenge,

measuring residual avoidance of the previously-lit zone. Prior to UCMS, there was no group

difference (**Supplementary Fig. 4A-B**). Upon UCMS, residual time spent in the shelter following

an acute light challenge was increased (**Fig. 2F-G**; **Supplementary Fig. 4C-G**). On week 6, acute

GL-RM treatment did not affect time in the shelter after the light challenge (**Fig. 2F**). Sex-

dependent analyses showed males spent more time in the shelter than females (data not shown).

Anhedonia-like and antidepressant-like behaviors were evaluated using the SC and FST tests,

respectively. At week 6, sucrose preference was significantly reduced by UCMS but unaffected

by GL-RM (**Fig. 2H**), suggesting a lack of effect on anhedonia. In the FST, UCMS increased time

immobile, which was reduced by GL-RM (**Fig. 2I**), suggesting antidepressant properties. Z-

scoring of behavioral emotionality confirmed the effect of UCMS, and a limited efficacy of acute

GL-RM treatment (**Fig. 2J**). In contrast, the Y-maze spatial working memory deficit induced by

UCMS was significantly reversed by acute GL-RM treatment (**Fig. 2K**).

Altogether, these results showed that acute administration of GL-RM has antidepressant and

procognitive potential in the UCMS model, while anxiety measures are inconclusive.

Chronic GL-RM conserves antidepressant and pro-cognitive properties in the UCMS

model.

The effects of chronic GL-RM administration were investigated in mice subjected to UCMS (**Fig.**

3A). Altered coat state and reduced weight gain with UCMS was significant from week 1-6, with

no effect on chronic GL-RM (**Supplementary Fig. 5**).

There was no effect of UCMS or treatment in the EPM, OF and NSF latency to approach (**Fig.**

3B-F). Latency to bite was decreased in UCMS-GL-RM mice compared to Control-GL-RM mice

(**Fig. 3G**). In the Phenotyper test (**Fig. 3H** and **Supplementary Fig. 6**), UCMS increased residual

avoidance in the shelter zone, while chronic GL-RM was trending to decrease it.

Sucrose preference was unaffected by UCMS, but chronic GL-RM increased preference (**Fig. 3I**),

suggesting improvement of anhedonia-like behavior. In the FST, the time spent immobile was

unaffected by UCMS or treatment (**Fig. 3J**). UCMS significantly increased z-emotionality, with a

trend level effect of chronic GL-RM reducing z-emotionality (**Fig. 3K**).

In the Y-maze, UCMS decreased percent alternation rate (Control-Vehicle vs UCMS-Vehicle) and

chronic GL-RM significantly increased percent alternation rate between UCMS-vehicle and

UCMS-GL-RM groups (**Fig. 3L**), confirming efficacy at reversing stress-induced memory deficits.

Results suggest that chronic GL-RM may contribute to reducing z-emotionality, with discrete

effects on individual mood symptoms, and confirm its effect at reversing UCMS-induced working

memory deficits.

Chronic GL-RM administration prevents neuronal morphology deficits induced by UCMS

We quantified morphological changes in apical and basal dendrites of PFC pyramidal neurons

(**Fig. 4A-C,** and **Supplementary Fig.7**). Dendritic length was not affected by UCMS or treatment,

but spine count and spine density were significantly lower in UCMS compared to control mice

(**Fig. 4D-E,** and **Supplementary Fig.7**). Such decreases were reversed by chronic GL-RM in

basal and apical segments, with spine density in apical PFC dendrites of GL-RM treated UCMS

animals being significantly higher than UCMS animals and not significantly different from controls

(**Fig. 4D**).

Similarly, significant UCMS effects on spine density and reversal by chronic GL-RM treatment

were observed in CA1 pyramidal neurons (**Fig. 4E,** and **Supplementary Fig. 8).**

Spine architecture in the PFC and CA1 are related to spatial working memory and emotionality

Spine densities in the PFC and CA1 were correlated to spatial working memory performance in the Y-maze (**Fig. 5 and Supplementary Table 5)**. In both regions, positive correlations were found at apical and basal segments, with lower alternation rates associated with lower spine densities in the PFC (**Fig. 5A-5B)** and the CA1 (**Fig. 5C**-**D**). No correlations were found for dendritic length or spine count in either region with Y-maze performance.

Correlations between morphological outcome and z-emotionality were calculated (**Supplementary Table 5)**. There were no significant correlations between z-emotionality and any morphological parameters in the PFC. In the CA1, increased dendritic length correlated with lower z-emotionality scores in basal and apical dendrites (**Fig. 5E-F**), with no correlations with spine count or spine density.

<u>Discussion</u>

This study is based on observations that patients with psychopathologies related to chronic stress exposure experience mood and cognitive deficits, while current treatments display moderate-to-no efficacy for mood and cognition [1-3, 9]. We investigated the efficacy of two enantiomers with activity at α3-, α2-, α5-GABAA-Rs (GL-I-54) and α5-GABAA-Rs (GL-II-73) separately, and in combination (GL-RM) at alleviating such symptoms and neuronal atrophy in CRS and UCMS models. GL-I-54 showed antidepressant-like effects, no anxiolytic nor cognitive effects at low dose (5mg/kg), and reverses working memory deficits at a higher dose (10mg/kg), while causing slight sedation. GL-II-73 previously showed no effect on working memory at doses lower than 10mg/kg [52]. Acute and chronic treatment of a racemix combination of both enantiomers (GL-RM) improved stress-induced cognitive deficits. Acute GL-RM showed anti-depressant potential but no anxiolytic effects. Finally, chronic GL-RM treatment reversed the

UCMS-induced reduction in spine densities in the PFC and CA1 of UCMS mice, with

morphological measures correlating with improvement of emotionality and working memory.

Racemic mixtures have therapeutic relevance by harnessing different enantiomer GABAA-

R specificities

Therapeutic development often favors pure substances over racemic mixtures due to superior

selectivity and reduced off-target activities [70, 71]. Racemic development becomes justified if

therapeutic benefits of each enantiomer are similar, complimentary or devoid of side-effects and

toxicity [72]. GL-RM harnessed the similar, yet unique selectivity profiles of GL-I-54 and GL-II-73

to target the GABAA-Rs known to be involved in anxiety, depression and cognition. GL-I-54

displays activity at α3-, α2-, α5-, α1-GABAA-Rs while GL-II-73 confirmed preferential activity at

α5-GABAA-Rs [52], likely reflecting different interactions of each compound's conformation in the

binding pocket.

Racemic mixtures are not uncommon in the treatment of MDD [73, 74]. Fluoxetine [73, 75],

citalopram [76] or ketamine [34] are clinically used racemic antidepressants [73, 77, 78]. Racemic

R,S-ketamine has more efficacy than non-racemic S-Ketamine [79]. Racemic therapies can

harness broader selectivity profiles to address multiple pathophysiological mechanisms but have

the added risk of off-target side effects. *In vivo,* considering estimated free concentrations in the

brain after a 5mg/kg dose, we found that concentration is slightly above 100nM. At this

concentration, GL-I-54 may act as a PAM at non-α1 GABAARs, while GL-II-73 would mainly have

only slight PAM activity at α2- and α5-GABAA-Rs, and be a null modulator at α1-GABAA-Rs.

When combined together, low concentrations of GL-II-73 may be effective as an "on-site softener"

of activity of GL-I-54, especially at α1-GABAA-Rs, limiting binding of GL-I-54 in the binding pocket

and therefore limiting negative effects observed when potentiating this subunit, further suggesting

GL-RM as an intervention for treatment of complete depression symptomatology.

Targeting α2-, α3-, α5-GABRA-Rs with GL-RM displays superior anti-depressant and pro-cognitive effects at lower concentrations than individual enantiomers.

Reconciling potentiation profiles with behavioral results, we compared GL-I-54 and GL-II-73 alone at 5mg/kg or 10mg/kg to GL-RM 10mg/kg (each enantiomer at 5mg/kg). At 5mg/kg, GL-I-54 has an antidepressant effect by reducing immobile time in the FST. Previously reported, GL-II-73 only reduced immobility time at 10 not 5mg/kg [52]. GL-RM reduced immobility time in the FST suggesting GL-RM retains antidepressant effects likely due to GL-I-54. Neither enantiomer shows clear anxiolytic effects alone or in GL-RM.

Interestingly, GL-RM combined the efficacy of both enantiomers for pro-cognitive effects. While neither enantiomer elicited pro-cognitive effects at 5mg/kg alone, their combined activity in GL-RM improved working memory deficits. This could be the results of an additive effect of the two compounds at GABAA-Rs, or a reduction of a putative amnestic effects caused by α1-GABAA-R null potentiation [80] with the use of a low GL-II-73 dose. Nevertheless, this finding demonstrates a clinical advantage for using GL-RM over either pure enantiomer at low concentrations, avoiding side effects and benefiting from combined profiles.

Chronic GL-RM administration has neurotrophic effects.

As in MDD, chronic stress leads to reduced synchronicity between the PFC and HPC, and dysfunctional information processing within cortical and hippocampal structures [81]. Dendritic length and spine density are reduced in the PFC and HPC of MDD patients and stressed mice [82], contributing to cognitive deficits. Our findings are consistent with other UCMS studies reporting neuronal atrophy and working memory deficits [83-85]. We previously showed that chronic administration of GL-II-73 alone remedies age-related spine loss within apical segments of pyramidal neurons, through α5-GABAA-R modulation confined to apical dendrites [56], because α5-GABAA-Rs are primarily located at apical dendrites [86-88] and GL-II-73 is

preferentially selective for α5-GABAA-Rs. With GL-RM, neurotrophic effects were extended to

basal segments in both PFC and HPC, suggesting a potential role of α2- and/or α3-GABAA-Rs in

this neurotrophic effect. Previously, α5- and α2-GABAA-Rs have been implicated in dendritic

outgrowth, spine maturation and synapse formation [89-91], thus their modulation by GL-RM may

be responsible for spine density restoration at apical and basal dendrites.

While showing spine count improvements, it remains unclear whether spines are prevented from

shrinking, or if they are generated *de novo*. Ketamine showed ability to overcome stress hormone-

induced spine loss by a combination of restored spines and *de novo* spines, with some preference

for *de novo* [92]. A similar combination may also be the case for GL-RM. Evidence that GL-RM

has an effect at a pathophysiological level (not only symptomatic levels) is substantially valuable

for translational efficacy within humans.

Limitations

We tested GL-RM on a single cognitive domain, exploring others could add more insight.

Regarding neurotrophic effects, we investigated the PFC and HPC for their regulation of stress-

related and depressive symptoms. It could be valuable to investigate how GL-RM modulates

connectivity in subnuclei of the amygdala, potentially decreasing the hyperactivity/increased

connectivity reported after chronic stress [93, 94]. We used chronic stress models because they

recapitulate behavioral and cellular changes observed in human depression [95-98]. While using

the most commonly reported stressors [99], variability in anxiety-related outcomes limited clear

conclusions on procedure and treatment, not common in behavioral studies [65, 100]. The

Phenotyper test measuring context anxiety [65] provided insight to anxiolytic properties of GL-RM

with better consistency. The discrepancy between EPM/NSF and Phenotyper results highlight

potential biases from hands-on behavioral assays versus automatized approaches [101].

To conclude, the therapeutic relevance of a targeted PAM approach at GABAA-Rs is

reinforced and the role of α5-GABAA-Rs in cognition is supported. At low-to-moderate dose, both

compounds, used together, show promising effects for the treatment of mood and cognitive

symptoms, as well as morphological changes in disorders such as MDD.

Funding and Disclosure

CAMH Internal Fund (Discovery Fund), awarded to TP at the time of study execution, and the

Campbell Family Mental Health Research Institute of CAMH. Chemistry synthesis funded by NIH

(DA-043204, R01NS076517) to JMC. The Ministry of Education, Science and Technological

Development, Republic of Serbia funded the pharmacokinetic study through Grant Agreement

with University of Belgrade-Faculty of Pharmacy No: 451-03-9/2021-14/200161. JMC, MS, ES

and TP are listed inventors on patents covering syntheses and use of the compounds. ES is

Founder of Damona Pharmaceuticals, a biopharma dedicated to bring the described compounds

to the clinic.

Acknowledgements

Authors thank Mehrab Ali and Netta Ussyshkin for support with administrative tasks throughout

the study. They thank CAMH animal facility staff for the caring for our animals over the study

duration. Authors further thank the members from Charles River Laboratories and NeuroDigitech

for their contribution to data generation. We also thank Milwaukee Institute for Drug Discovery

and University of Wisconsin-Milwaukee's Shimadzu Laboratory for Advanced and Applied

Analytical Chemistry for help with spectroscopy and National Science Foundation, Division of

Chemistry [CHE-1625735].

Author Contributions

TP and ES designed the study. YM, ZAK, DS and JMC synthesized the compounds tested in this

study. TP and PL performed the behavioral piece. AK and MS performed the pharmacokinetic

piece. Electrophysiology was outsourced to Charles River Laboratories. TP, AB and MM analyzed

the data. TP and AB wrote the manuscript.

References

1. MacQueen, G.M. and K.A. Memedovich, *Cognitive dysfunction in major depression and bipolar disorder: Assessment and treatment options.* Psychiatry and Clinical Neurosciences, 2017. **71**(1): p. 18-27.
2. Malhi, G.S. and J.J. Mann, *Depression.* The Lancet, 2018. **392**(10161): p. 2299-2312.
3. Gartlehner, G., et al., *Pharmacological and non-pharmacological treatments for major depressive disorder: review of systematic reviews.* BMJ Open, 2017. **7**(6): p. e014912.
4. Hirschfeld, R.M.A., *History and evolution of the monoamine hypothesis of depression.* The Journal of Clinical Psychiatry, 2000. **61**(Suppl6): p. 4-6.
5. Xia, C.-Y., et al., *Targeting the dysfunction of glutamate receptors for the development of novel antidepressants.* Pharmacology & Therapeutics, 2021. **226**: p. 107875.
6. Davis, M.T., et al., *Acute cognitive effects of single-dose intravenous ketamine in major depressive and posttraumatic stress disorder.* Translational Psychiatry, 2021. **11**(1): p. 205.
7. Gill, H., et al., *The Effects of Ketamine on Cognition in Treatment-Resistant Depression: A Systematic Review and Priority Avenues for Future Research.* Neurosci Biobehav Rev, 2021. **120**: p. 78-85.
8. Prevot, T. and E. Sibille, *Altered GABA-mediated information processing and cognitive dysfunctions in depression and other brain disorders.* Mol Psychiatry, 2021. **26**(1): p. 151-167.
9. Duman, C.H. and R.S. Duman, *Spine synapse remodeling in the pathophysiology and treatment of depression.* Neurosci Lett, 2015. **601**: p. 20-9.
10. Williams, L.M., *Precision psychiatry: a neural circuit taxonomy for depression and anxiety.* The Lancet Psychiatry, 2016. **3**(5): p. 472-480.
11. Banasr, M., J.M. Dwyer, and R.S. Duman, *Cell atrophy and loss in depression: reversal by antidepressant treatment.* Current opinion in cell biology, 2011. **23**(6): p. 730-737.
12. Kang, H.J., et al., *Decreased expression of synapse-related genes and loss of synapses in major depressive disorder.* Nat Med, 2012. **18**(9): p. 1413-7.
13. Holmes, S.E., et al., *Lower synaptic density is associated with depression severity and network alterations.* Nature Communications, 2019. **10**(1): p. 1529.
14. Lissemore, J.I., et al., *Reduced GABAergic cortical inhibition in aging and depression.* Neuropsychopharmacology, 2018. **43**(11): p. 2277-2284.
15. Lee, J.C., et al., *The role of transcranial magnetic stimulation in treatment-resistant depression: a review.* Curr Pharm Des, 2012. **18**(36): p. 5846-52.
16. Fogaca, M.V. and R.S. Duman, *Cortical GABAergic Dysfunction in Stress and Depression: New Insights for Therapeutic Interventions.* Front Cell Neurosci, 2019. **13**: p. 87.
17. Newton, D.F., et al., *Chapter 28 - Altered GABAergic Function, Cortical Microcircuitry, and Information Processing in Depression*, in *Neurobiology of Depression*, J. Quevedo, A.F. Carvalho, and C.A. Zarate, Editors. 2019, Academic Press. p. 315-329.
18. Bhagwagar, Z., et al., *Low GABA concentrations in occipital cortex and anterior cingulate cortex in medication-free, recovered depressed patients.* The international journal of neuropsychopharmacology / official scientific journal of the Collegium Internationale Neuropsychopharmacologicum (CINP), 2008. **11**(2): p. 255-60.
19. Goddard, A.W., et al., *Reductions in Occipital Cortex GABA Levels in Panic Disorder Detected With 1H-Magnetic Resonance Spectroscopy.* Archives of General Psychiatry, 2001. **58**(6): p. 556-556.
20. Price, R.B., et al., *Amino acid neurotransmitters assessed by proton magnetic resonance spectroscopy: relationship to treatment resistance in major depressive disorder.* Biol Psychiatry, 2009. **65**(9): p. 792-800.

21. Sanacora, G., et al., *Subtype-specific alterations of gamma-aminobutyric acid and glutamate in*
patients with major depression. Arch Gen Psychiatry, 2004. **61**(7): p. 705-13.
22. Sanacora, G., et al., *Increased occipital cortex GABA concentrations in depressed patients after*
therapy with selective serotonin reuptake inhibitors. Am J Psychiatry, 2002. **159**(4): p. 663-5.
23. Sanacora, G., et al., *Subtype-specific alterations of gamma-aminobutyric acid and glutamate in*
patients with major depression. Archives of general psychiatry, 2004. **61**(7): p. 705-13.
24. Sanacora, G., et al., *Reduced Cortical γ-Aminobutyric Acid Levels in Depressed Patients Determined*
by Proton Magnetic Resonance Spectroscopy. Archives of General Psychiatry, 1999. **56**(11): p.
1043-1043.
25. Hasler, G., et al., *Reduced prefrontal glutamate/glutamine and gamma-aminobutyric acid levels*
in major depression determined using proton magnetic resonance spectroscopy. Archives of
general psychiatry, 2007. **64**(2): p. 193-200.
26. Gabbay, V., et al., *Anterior Cingulate Cortexy-Aminobutyric Acid in Depressed Adolescents.*
Archives of General Psychiatry, 2012. **69**(2): p. 139-139.
27. Plante, D.T., et al., *Reduced gamma-aminobutyric acid in occipital and anterior cingulate cortices*
in primary insomnia: a link to major depressive disorder? Neuropsychopharmacology, 2012. **37**(6):
p. 1548-57.
28. Feyissa, A.M., et al., *Reduced levels of NR2A and NR2B subunits of NMDA receptor and PSD-95 in*
the prefrontal cortex in major depression. Progress in Neuro-Psychopharmacology and Biological
Psychiatry, 2009. **33**(1): p. 70-75.
29. Christoffel, D.J., S.A. Golden, and S.J. Russo, *Structural and synaptic plasticity in stress-related*
disorders. Reviews in the neurosciences, 2011. **22**(5): p. 535-549.
30. Izquierdo, A., C.L. Wellman, and A. Holmes, *Brief uncontrollable stress causes dendritic retraction*
in infralimbic cortex and resistance to fear extinction in mice. J Neurosci, 2006. **26**(21): p. 5733-8.
31. Liston, C., et al., *Stress-induced alterations in prefrontal cortical dendritic morphology predict*
selective impairments in perceptual attentional set-shifting. J Neurosci, 2006. **26**(30): p. 7870-4.
32. Shansky, R.M., et al., *Stress-Induced Dendritic Remodeling in the Prefrontal Cortex is Circuit*
Specific. Cerebral Cortex, 2009. **19**(10): p. 2479-2484.
33. Abdallah, C.G., et al., *Prefrontal cortical GABA abnormalities are associated with reduced*
hippocampal volume in major depressive disorder. European Neuropsychopharmacology, 2015.
25(8): p. 1082-1090.
34. Phoumthipphavong, V., et al., *Longitudinal Effects of Ketamine on Dendritic Architecture In Vivo*
in the Mouse Medial Frontal Cortex. eNeuro, 2016. **3**(2): p. ENEURO.0133-15.2016.
35. Gerhard, D.M., et al., *GABA interneurons are the cellular trigger for ketamine's rapid*
antidepressant actions. J Clin Invest, 2020. **130**(3): p. 1336-1349.
36. Mohler, H., *GABA(A) receptor diversity and pharmacology.* Cell Tissue Res, 2006. **326**(2): p. 505-
16.
37. Sigel, E. and M. Ernst, *The Benzodiazepine Binding Sites of GABAA Receptors.* Trends in
Pharmacological Sciences, 2018. **39**(7): p. 659-671.
38. Pariente, A., et al., *The Benzodiazepine–Dementia Disorders Link: Current State of Knowledge.* CNS
Drugs, 2016. **30**(1): p. 1-7.
39. Sieghart, W. and M.M. Savić, *International Union of Basic and Clinical Pharmacology. CVI: GABA(A)*
Receptor Subtype- and Function-selective Ligands: Key Issues in Translation to Humans. Pharmacol
Rev, 2018. **70**(4): p. 836-878.
40. Fritschy, J.M. and H. Mohler, *GABAA-receptor heterogeneity in the adult rat brain: differential*
regional and cellular distribution of seven major subunits. J Comp Neurol, 1995. **359**(1): p. 154-94.

41. Geracitano, R., et al., *Functional expression of the GABA(A) receptor α2 and α3 subunits at synapses between intercalated medial paracapsular neurons of mouse amygdala.* Front Neural Circuits, 2012. **6**: p. 32.

42. Stefanits, H., et al., *GABAA receptor subunits in the human amygdala and hippocampus: Immunohistochemical distribution of 7 subunits.* Journal of Comparative Neurology, 2018. **526**(2): p. 324-348.

43. Marowsky, A., J.M. Fritschy, and K.E. Vogt, *Functional mapping of GABA A receptor subtypes in the amygdala.* Eur J Neurosci, 2004. **20**(5): p. 1281-9.

44. Heldt, S.A. and K.J. Ressler, *Training-induced changes in the expression of GABAA-associated genes in the amygdala after the acquisition and extinction of Pavlovian fear.* Eur J Neurosci, 2007. **26**(12): p. 3631-44.

45. Dias, R., et al., *Evidence for a significant role of alpha 3-containing GABAA receptors in mediating the anxiolytic effects of benzodiazepines.* J Neurosci, 2005. **25**(46): p. 10682-8.

46. Möhler, H., *The GABA system in anxiety and depression and its therapeutic potential.* Neuropharmacology, 2012. **62**(1): p. 42-53.

47. Atack, J.R., *GABAA receptor alpha2/alpha3 subtype-selective modulators as potential nonsedating anxiolytics.* Curr Top Behav Neurosci, 2010. **2**: p. 331-60.

48. Engin, E., J. Liu, and U. Rudolph, *α2-containing GABA(A) receptors: a target for the development of novel treatment strategies for CNS disorders.* Pharmacology & therapeutics, 2012. **136**(2): p. 142-152.

49. Collins, I., et al., *3-Heteroaryl-2-pyridones: benzodiazepine site ligands with functional delectivity for alpha 2/alpha 3-subtypes of human GABA(A) receptor-ion channels.* J Med Chem, 2002. **45**(9): p. 1887-900.

50. Miller, B.H., et al., *Quantitative trait locus analysis identifies Gabra3 as a regulator of behavioral despair in mice.* Mammalian genome : official journal of the International Mammalian Genome Society, 2010. **21**(5-6): p. 247-257.

51. Koh, M.T., S. Rosenzweig-Lipson, and M. Gallagher, *Selective GABA(A) alpha5 positive allosteric modulators improve cognitive function in aged rats with memory impairment.* Neuropharmacology, 2013. **64**: p. 145-52.

52. Prevot, T., et al., *Novel Benzodiazepine-Like Ligands with Various Anxiolytic, Antidepressant, or Pro-Cognitive Profiles.* Molecular Neuropsychiatry, 2019. **5**(2): p. 84-97.

53. Prévot, T. and E. Sibille, *Altered GABA-mediated information processing and cognitive dysfunctions in depression and other brain disorders.* Mol Psychiatry, 2021. **26**(1): p. 151-167.

54. Sigurdsson, T. and S. Duvarci, *Hippocampal-Prefrontal Interactions in Cognition, Behavior and Psychiatric Disease.* Frontiers in systems neuroscience, 2016. **9**: p. 190-190.

55. Koh, M.T., S. Rosenzweig-Lipson, and M. Gallagher, *Selective GABA(A) α5 positive allosteric modulators improve cognitive function in aged rats with memory impairment.* Neuropharmacology, 2013. **64**(1): p. 145-152.

56. Prevot, T.D., et al., *Reversal of Age-Related Neuronal Atrophy by α5-GABAA Receptor Positive Allosteric Modulation.* Cerebral Cortex, 2020.

57. Rudolph, U., et al., *Benzodiazepine actions mediated by specific gamma-aminobutyric acid(A) receptor subtypes.* Nature, 1999. **401**(6755): p. 796-800.

58. Li, G., et al., *Synthesis of chiral GABA(A) receptor subtype selective ligands as potential agents to treat schizophrenia as well as depression.* Arkivoc, 2018. **2018**(4): p. 158-182.

59. Billakota, S., et al., *Personalized medicine: Vinpocetine to reverse effects of GABRB3 mutation.* Epilepsia, 2019. **60**(12): p. 2459-2465.

535   60.   Kirsch, G.E., et al., *Electrophysiology-Based Assays to Detect Subtype-Selective Modulation of*
536   *Human Nicotinic Acetylcholine Receptors.* ASSAY and Drug Development Technologies, 2016.
537   **14**(6): p. 333-344.
538   61.   Bayliss, M.K. and L.W. Frick, *High-throughput pharmacokinetics: cassette dosing.* Curr Opin Drug
539   Discov Devel, 1999. **2**(1): p. 20-5.
540   62.   Stamenic, T.T., et al., *Ester to amide substitution improves selectivity, efficacy and kinetic behavior*
541   *of a benzodiazepine positive modulator of GABAA receptors containing the alpha5 subunit.* Eur J
542   Pharmacol, 2016. **791**: p. 433-443.
543   63.   Obradovic, A., et al., *Sh-I-048A, an in vitro non-selective super-agonist at the benzodiazepine site*
544   *of GABAA receptors: the approximated activation of receptor subtypes may explain behavioral*
545   *effects.* Brain Res, 2014. **1554**: p. 36-48.
546   64.   Di, L., et al., *Optimization of a higher throughput microsomal stability screening assay for profiling*
547   *drug discovery candidates.* Journal of biomolecular screening, 2003. **8**(4): p. 453-462.
548   65.   Prevot, T.D., et al., *Residual avoidance: A new, consistent and repeatable readout of chronic stress-*
549   *induced conflict anxiety reversible by antidepressant treatment.* Neuropharmacology, 2019. **153**:
550   p. 98-110.
551   66.   Nollet, M., A.M. Le Guisquet, and C. Belzung, *Models of depression: unpredictable chronic mild*
552   *stress in mice.* Curr Protoc Pharmacol, 2013. **Chapter 5**: p. Unit 5.65.
553   67.   Faucher, P., et al., *Hippocampal Injections of Oligomeric Amyloid β-peptide (1-42) Induce Selective*
554   *Working Memory Deficits and Long-lasting Alterations of ERK Signaling Pathway.* Front Aging
555   Neurosci, 2015. **7**: p. 245.
556   68.   Wu, C.C., et al., *Selective vulnerability of dentate granule cells prior to amyloid deposition in PDAPP*
557   *mice: digital morphometric analyses.* Proc Natl Acad Sci U S A, 2004. **101**(18): p. 7141-6.
558   69.   Guilloux, J.-P., et al., *Integrated behavioral z-scoring increases the sensitivity and reliability of*
559   *behavioral phenotyping in mice: Relevance to emotionality and sex.* Journal of Neuroscience
560   Methods, 2011. **197**(1): p. 21-31.
561   70.   Sekhon, B.S., *Exploiting the power of stereochemistry in drugs: an overview of racemic and*
562   *enantiopure drugs.* J. Mod. Med. Chem, 2013. **1**: p. 10-36.
563   71.   Nguyen, L.A., H. He, and C. Pham-Huy, *Chiral drugs: an overview.* International journal of
564   biomedical science : IJBS, 2006. **2**(2): p. 85-100.
565   72.   Srinivas, N.R., R.H. Barbhaiya, and K.K. Midha, *Enantiomeric drug development: issues,*
566   *considerations, and regulatory requirements.* J Pharm Sci, 2001. **90**(9): p. 1205-15.
567   73.   Budău, M., et al., *Chirality of Modern Antidepressants: An Overview.* Adv Pharm Bull, 2017. **7**(4):
568   p. 495-500.
569   74.   Zarate, C.A., *Commentary on the Canadian Network for Mood and Anxiety Treatments (CANMAT)*
570   *Task Force Recommendations for the Use of Racemic Ketamine in Adults with Major Depressive*
571   *Disorder.* The Canadian Journal of Psychiatry. **0**(0): p. 07067437211004023.
572   75.   Henry, M.E., et al., *A comparison of brain and serum pharmacokinetics of R-fluoxetine and racemic*
573   *fluoxetine: A 19-F MRS study.* Neuropsychopharmacology, 2005. **30**(8): p. 1576-83.
574   76.   Gorman, J.M., A. Korotzer, and G. Su, *Efficacy comparison of escitalopram and citalopram in the*
575   *treatment of major depressive disorder: pooled analysis of placebo-controlled trials.* CNS Spectr,
576   2002. **7**(4 Suppl 1): p. 40-4.
577   77.   Perez-Caballero, L., et al., *Fluoxetine: a case history of its discovery and preclinical development.*
578   Expert Opin Drug Discov, 2014. **9**(5): p. 567-78.
579   78.   Bezchlibnyk-Butler, K., I. Aleksic, and S.H. Kennedy, *Citalopram--a review of pharmacological and*
580   *clinical effects.* J Psychiatry Neurosci, 2000. **25**(3): p. 241-54.

581   79.   Bahji, A., G.H. Vazquez, and C.A. Zarate, Jr., *Comparative efficacy of racemic ketamine and*
582   *esketamine for depression: A systematic review and meta-analysis.* J Affect Disord, 2021. **278**: p.
583   542-555.
584   80.   Möhler, H., *Pathophysiological aspects of diversity in neuronal inhibition: a new benzodiazepine*
585   *pharmacology.* Dialogues in clinical neuroscience, 2002. **4**(3): p. 261-269.
586   81.   Abbas, A.I., et al., *Somatostatin Interneurons Facilitate Hippocampal-Prefrontal Synchrony and*
587   *Prefrontal Spatial Encoding.* Neuron, 2018. **100**(4): p. 926-939.e3.
588   82.   Duman, C.H. and R.S. Duman, *Spine synapse remodeling in the pathophysiology and treatment of*
589   *depression.* Neuroscience letters, 2015. **601**: p. 20-29.
590   83.   Prévôt, T.D., et al., *sst(2)-receptor gene deletion exacerbates chronic stress-induced deficits:*
591   *Consequences for emotional and cognitive ageing.* Prog Neuropsychopharmacol Biol Psychiatry,
592   2018. **86**: p. 390-400.
593   84.   Gumuslu, E., et al., *The Antidepressant Agomelatine Improves Memory Deterioration and*
594   *Upregulates CREB and BDNF Gene Expression Levels in Unpredictable Chronic Mild Stress (UCMS)-*
595   *Exposed Mice.* Drug Target Insights, 2014. **8**: p. 11-21.
596   85.   Fee, C., et al., *Chronic Stress-induced Behaviors Correlate with Exacerbated Acute Stress-induced*
597   *Cingulate Cortex and Ventral Hippocampus Activation.* Neuroscience, 2020. **440**: p. 113-129.
598   86.   Hörtnagl, H., et al., *Patterns of mRNA and protein expression for 12 GABAA receptor subunits in*
599   *the mouse brain.* Neuroscience, 2013. **236**: p. 345-372.
600   87.   Schulz, J.M., et al., *Dendrite-targeting interneurons control synaptic NMDA-receptor activation via*
601   *nonlinear α5-GABA(A) receptors.* Nat Commun, 2018. **9**(1): p. 3576.
602   88.   Serwanski, D.R., et al., *Synaptic and nonsynaptic localization of GABAA receptors containing the*
603   *alpha5 subunit in the rat brain.* J Comp Neurol, 2006. **499**(3): p. 458-70.
604   89.   Duveau, V., et al., *Spatiotemporal specificity of GABAA receptor-mediated regulation of adult*
605   *hippocampal neurogenesis.* Eur J Neurosci, 2011. **34**(3): p. 362-73.
606   90.   Deprez, F., et al., *Partial inactivation of GABAA receptors containing the α5 subunit affects the*
607   *development of adult-born dentate gyrus granule cells.* Eur J Neurosci, 2016. **44**(5): p. 2258-71.
608   91.   Brady, M.L. and T.C. Jacob, *Synaptic localization of α5 GABA (A) receptors via gephyrin interaction*
609   *regulates dendritic outgrowth and spine maturation.* Dev Neurobiol, 2015. **75**(11): p. 1241-51.
610   92.   Moda-Sava, R.N., et al., *Sustained rescue of prefrontal circuit dysfunction by antidepressant-*
611   *induced spine formation.* Science, 2019. **364**(6436).
612   93.   Nikolova, Y.S., et al., *Shifting priorities: highly conserved behavioral and brain network adaptations*
613   *to chronic stress across species.* Transl Psychiatry, 2018. **8**(1): p. 26.
614   94.   Zhang, X., et al., *Stress-Induced Functional Alterations in Amygdala: Implications for*
615   *Neuropsychiatric Diseases.* Frontiers in neuroscience, 2018. **12**: p. 367-367.
616   95.   Luscher, B., Q. Shen, and N. Sahir, *The GABAergic deficit hypothesis of major depressive disorder.*
617   Mol Psychiatry, 2011. **16**(4): p. 383-406.
618   96.   Karolewicz, B., et al., *Reduced level of glutamic acid decarboxylase-67 kDa in the prefrontal cortex*
619   *in major depression.* Int J Neuropsychopharmacol, 2010. **13**(4): p. 411-20.
620   97.   Romeo, B., et al., *Meta-analysis of central and peripheral γ-aminobutyric acid levels in patients*
621   *with unipolar and bipolar depression.* J Psychiatry Neurosci, 2018. **43**(1): p. 58-66.
622   98.   Hasler, G., et al., *Reduced prefrontal glutamate/glutamine and gamma-aminobutyric acid levels*
623   *in major depression determined using proton magnetic resonance spectroscopy.* Arch Gen
624   Psychiatry, 2007. **64**(2): p. 193-200.
625   99.   Antoniuk, S., et al., *Chronic unpredictable mild stress for modeling depression in rodents: Meta-*
626   *analysis of model reliability.* Neuroscience & Biobehavioral Reviews, 2019. **99**: p. 101-116.
627   100.   Mineur, Y.S., C. Belzung, and W.E. Crusio, *Effects of unpredictable chronic mild stress on anxiety*
628   *and depression-like behavior in mice.* Behav Brain Res, 2006. **175**(1): p. 43-50.

629   101.   Aarts, E., et al., *The light spot test: Measuring anxiety in mice in an automated home-cage*
630   *environment.* Behav Brain Res, 2015. **294**: p. 123-30.

  Figure Legend

Figure 1. Electrophysiological, pharmacokinetic and behavioral profiles of GL-II-73 and GL-I-54

Electrophysiological recordings were obtained from HEK-293T cells transfected with full-length cDNA for human GABAA receptor subtypes α1β3γ2, α2β3γ2, α3β3γ2, α4β3γ2 or α5β3γ2, in presence of GL-I-54 (**A**) or GL-II-73 (**B**), and in the presence of GABA in the medium (5μM). Pharmacokinetic profile of GL-I-54 was also examined (**C**). Plasma, brain and free brain concentration-time profile of GL-I-54 after intraperitoneal cassette administration of 3 mg/kg dose in male C57BL/6 mice (n = 3 per time point). C_{max}, maximum concentration in plasma or brain; T_{max}, time of maximum concentration in plasma or brain; AUC_{0-720}, area under the plasma or brain concentration-time curve from 0 to 720 min; $AUC_{0-\infty}$, area under the plasma or brain concentration-time curve from 0 to extrapolated infinite time; $t_{1/2}$, elimination half-life from plasma or brain; β, elimination constant rate from plasma or brain; K_p, brain-to-plasma partition coefficient ($K_p = AUC_{0-\infty}$, brain/$AUC_{0-\infty}$, plasma); $K_{p,uu}$, ratio of unbound brain to unbound plasma drug concentrations ($K_{p,uu} = K_p \times$ unbound fraction in brain/unbound fraction in plasma). GL-I-54 and GL-II-73 were then tested in the elevated plus maze and forced swim test at the dose of 5mg/kg, in mice previously exposed to chronic restraint stress (CRS). Drugs were administered IP, 30min prior to testing. Percent time spent in the open arms (**D**) showed no difference between groups. Time immobile (**E**) showed more time spent immobile in the CRS-Vehicle group, and less time immobile in animals receiving GL-I-54, compared to CRS-Vehicle. GL-II-73 did not show an effect. GL-I-54 was tested in the Y-maze task, assessing working memory (**F**). Animals subjected to CRS and receiving vehicle showed a significant decrease in alternation rate, suggesting a working memory deficit. Animal subjected to CRS and receiving the highest dose of GL-I-54 showed significant increase in alternation rate, suggesting reversal of working memory deficits induced by CRS. Finally, independent mice were tested in the rotarod (**G**; N=5 Control Vehicle and N=6 GL-I-54). Mice were trained to maintain themselves on a rotating rod (rotarod) for 3 trials. Then, they were injected with GL-I-54 at 10 mg/kg, and tested 5 min, 20 min and 60 min past injection time. Latency to fall from the rod was recorded, and showed significant reductions in latency to fall in mice receiving GL-I-54. All values are represented as mean ± standard error of the mean. *p<0.05, **p<0.01, ***p<0.001 compared to 100 % in A-B) or to Control, Control/Vehicle in D-G. θp<0.05 compared to CRS-Vehicle.

Figure 2. Effect of acute treatment of GL-RM on anxiety, emotionality and working memory deficits in mice subjected to chronic stress

Male and female mice were tested at baseline in the Y-maze, the phenotyper and the sucrose consumption test prior to being subjected to 6 weeks of UCMS (**A**). Weekly, mice were tested in the phenotyper test, the sucrose consumption and their weight and coat state were measured. After 6 weeks of UCMS, acute injections were performed 30 minutes prior to behavioral testing. In the elevated plus maze, time spent (**B**) and entries (**C**) in the open arms were measured, with no significant effect of UCMS, treatment or interaction. In the novelty suppressed feeding test, latency to approach (**D**) and latency to bite (**E**) were assessed. Statistical analyses showed that acute GL-RM treatment reduces latency to approach in mice subjected to UCMS. In the phenotyper test (**F**), mice were placed in the box overnight, where a stressful stimulus was applied at 11pm, for 1hr. Time spent in the shelter zone showed that mice subjected to UCMS spent more time in the shelter than control mice. Calculating a residual avoidance score (**G**), statistical analyses showed a significant increase in residual avoidance in mice subjected to UCMS. In the sucrose consumption test (**H**), mice subjected to UCMS showed a significant decrease in preference to sucrose. In the forced swim test (**I**), mice subjected to UCMS showed increased immobility, while mice treated with GL-RM showed a reduction in immobility. Combining the individual score into a global z-score (**J**), statistical analyses confirmed a significant impact of UCMS, with reduced effect of GL-RM. Finally, mice were tested in the Y-maze (**K**), where statistical analyses showed altered alternation with UCMS, which is reversed by acute GL-RM treatment. $^t p<0.1$, *$p<0.05$, **$p<0.01$, ***$p<0.001$ effect of UCMS, #$p<0.05$, ##$p<0.01$ effect of GL-RM

Figure 3. Effect of chronic treatment of GL-RM on anxiety, emotionality and working memory deficits in mice subjected to chronic stress

Male and female mice were tested at baseline in the Y-maze, the phenotyper and the sucrose consumption test prior to being subjected to 6 weeks of UCMS (**A**). After 3 weeks of UCMS, chronic treatment with GL-RM in the drinking water was initiated, for a total of 4 weeks. Weekly, mice were tested in the phenotyper test, the sucrose consumption and their weight and coat state were measured. After 6 weeks of UCMS, and 3 weeks of treatment, mice were tested in the elevated plus maze. Time spent (**B**) and entries (**C**) in the open arms were measured, but did not show statistical differences. Mice were also tested in the open field for the time spent (**D**) and number of entries (**E**) in the inner zone. Again, statistical analyses did not reveal any effect of UCMS nor treatment. In the novelty suppressed feeding test, latency to approach (**F**) and latency

to bite (**G**) were assessed. Mice were tested in the Phenotyper weekly, and the residual avoidance
scores from week 3 to 6 were analyzed, since the treatment was onboard during these testing
periods (**H**). In the sucrose consumption test (**I**), mice receiving chronic GL-RM showed a
significant increase in preference to sucrose. In the forced swim test (**J**), statistical analyses did
not reveal significant differences between groups. Combining the individual score into a global z-
score (**K**), statistical analyses confirmed a significant impact of UCMS at increasing emotionality,
with a trend level effect of chronic GL-RM reducing emotionality. Finally, mice were tested in the
Y-maze (**L**), where statistical analyses showed altered alternation with UCMS, which is reversed
by acute GL-RM treatment. *p<0.05, **p<0.01, ***p<0.001 effect of UCMS, ^{f}p<0.1, #p<0.05,
###p<0.001 effect of GL-RM

Figure 4. Chronic treatment with GL-RM reverses chronic-stress induced spine density
reduction in the PFC and the CA1.
After completion of the behavioral screening, mice were euthanized and brains were stained with
Golgi-Cox solution. Pyramidal neurons (N=6 per mouse) from 4 mice per group (**A-C**) were
analyzed for dendritic length, spine counts and spine density. Basal and apical spine densities
were measured in the PFC (**D**) and the CA1 of the hippocampus (**E**). ANOVA in the basal and
apical segments revealed significant differences between groups, in both brain regions. This
difference was explained by a decrease in spine density in mice subjected to UCMS compared to
Control mice that was partially reversed by chronic treatment with GL-RM. *p<0.05, **p<0.01,
***p<0.001, ****p<0.0001 compared to "Control"; $$p<0.01, $$$p<0.001, $$$$p<0.0001
compared to "UCMS". Scale bar in (A-C) represents 50μm.

Figure 5. Morphological changes in the PFC and CA1 correlates with behavioral outcomes.

Morphological features from the PFC and the CA1 were correlated to behavioral outcomes using
the alternation rate (**A-D**) and the z-emotionality (**E-F**). Basal (**A**) and apical (**B**) spine density in
the PFC correlated positively with the alternation rate, where higher density was associated with
higher alternation rate, i.e. higher memory performances. Similar findings were reported when
looking at basal and apical spine density from the CA1 (**C-D**). Basal and Apical dendritic length in
the CA1 also correlated with z-emotionality, this time negatively (**E-F**), where higher dendritic
length was associated with lower z-emotionality, i.e. lower anxiety- and depressive-like behaviors.