1

Revisiting the role of *Toxoplasma gondii* ERK7 in the maintenance and stability

of the apical complex

Nicolas Dos Santos Pacheco[1*], Nicolò Tosetti[1*], Aarti Krishnan[1], Romuald Haase[1] & Dominique

Soldati-Favre[1#]

[1]Department of Microbiology and Molecular Medicine, Faculty of Medicine, University of Geneva.

1 Rue Michel-Servet, 1211 Geneva, Switzerland

Running title: *Revisiting the role of ERK7 in Toxoplasma gondii*

*These authors contributed equally to this work

#Address correspondence to Dominique Soldati-Favre: Dominique.Soldati-Favre@unige.ch

Abbreviations: apical cap protein AC; subpellicular microtubules, SPMTs; inner membrane

complex, IMC; Extracellular signal-regulated kinase 7, ERK7.

Keywords: Apicomplexa, *Toxoplasma gondii*, conoid, apical cap, subpellicular microtubules,

comparative proteomics, egress, motility, invasion, microneme secretion.

Abstract

Toxoplasma gondii ERK7 is known to contribute to the integrity of the apical complex and to be involved only in the final step of the conoid biogenesis. In the absence of ERK7, mature parasites lose their conoid complex and are unable to glide, invade or egress from host cells. In contrast to a previous report, we show here that depletion of ERK7 phenocopies the depletion of the apical cap proteins AC9 or AC10. The absence of ERK7 leads to the loss of the apical polar ring, the disorganization of the basket of subpellicular microtubules and an impairment in micronemes secretion. Ultra-expansion microscopy (U-ExM) coupled to NHS-Ester staining on intracellular parasites offers an unprecedented level of resolution and highlights the disorganization of the rhoptries as well as the dilated plasma membrane at the apical pole in the absence of ERK7. Comparative proteomics analysis of wild-type and ERK7 or AC9 depleted parasites led to the disappearance of known, predicted, as well as putative novel components of the apical complex. In contrast, the absence of ERK7 led to an accumulation of microneme proteins, resulting from the defect in exocytosis of the organelles.

Importance

The conoid is an enigmatic, dynamic organelle positioned at the apical tip of the coccidian subgroup of the Apicomplexa, close to the apical polar ring (APR) from which the subpellicular microtubules (SPTMs) emerge and at the site of microneme and rhoptry secretory organelles exocytosis. In *Toxoplasma gondii,* the conoid protrudes concomitantly to microneme secretion, during egress, motility and invasion. The conditional depletion of the apical cap structural protein AC9 or AC10 leads to a disorganization of SPTMs as well as the loss of APR and conoid that result in microneme secretion defect and block in motility, invasion and egress. We show here that depletion of the kinase ERK7 phenocopies completely AC9 and AC10 mutants. Moreover, the combination of ultrastructure expansion microscopy with an NHS ester staining revealed that ERK7 depleted parasites exhibit a dilated apical plasma membrane and a mis-positioning of the rhoptry organelles.

Introduction

The Apicomplexa phylum is defined by the so-called apical complex, a structure that harbors unique secretory organelles, termed rhoptries and micronemes, as well as membranous and cytoskeletal elements that critically contributes to parasite dissemination. The overall cytoskeleton is made of three interacting layers, conferring structure and stability to the parasite: the inner membrane complex (IMC), formed by a patchwork of flattened vesicles, the subpellicular network (SPN) composed of intermediate filaments-like proteins termed alveolins and the subpellicular microtubules (SPMTs). The most apical region of the IMC, called the apical cap, is formed by a single vesicle and was shown to participate to the apical complex stability (Back et al. 2020; O'Shaughnessy et al. 2020; Tosetti et al. 2020) Members of the coccidian subgroup possess an additional open-barrel shaped structure made of unique fibers of tubulin polymers termed the conoid (Hu et al. 2002), with two pre-conoidal rings (PCRs) at the top and two short intra-conoidal microtubules on the inside (Dos Santos Pacheco et al. 2020). The conoid lays between the apical polar ring (APR) which serves as microtubule organizing center (MTOC) for the spiraling SPMTs. The apical cap proteins 9 and 10 (AC9, AC10) were previously shown to be SPN resident proteins whose depletion resulted in the loss of the conoid and the APR, and disorganization of the SPMTs at the apical pole (Back *et al.* 2020; Tosetti *et al.* 2020). The conoid is the site of convergence for calcium and lipid-mediated signaling cascades that coordinate microneme secretion and actin polymerization, and is also the site of for the glideosome, the molecular machinery powering parasite gliding motility (Bisio & Soldati-Favre 2019). Strikingly AC9 and AC10 depleted parasites were severely impaired in microneme secretion and consequently unable to move, invade or egress from host cells. We postulated that the loss of the conoid and SPMTs disorganization, could result from the disappearance of the APR. Concordantly another study reported that the double knock-out of two APR proteins (APR1 and KinA) led to the fragmentation of the APR, partial detachment of the conoid and reduced microneme exocytosis (Leung et al. 2017). In contrast, depletion of the mitogen-activated protein kinase (MAPK) ERK7 reportedly cause the loss of conoid in mature

parasite without seemingly affecting microneme secretion nor APR integrity (O'Shaughnessy *et al.* 2020). Moreover, AC9 was assigned a dual role in localizing ERK7 at the apical cap and in regulating its kinase activity and substrate specificity (Back *et al.* 2020). Another protein, the conoidal ankyrin repeat-containing protein (CPH1) was also reported to contribute to the conoid integrity in extracellular parasites. CPH1 depleted parasites harbored shortened, partially collapsed conoids, again without seemingly impacting on microneme secretion (Long et al. 2017a).

Here, we have revisited the role of ERK7 and CPH1 in microneme secretion. Depletion of ERK7 phenocopies strictly the depletion AC9 or AC10, leading not only to the loss of the conoid but also the disappearance of the APR and a severe impairment in induced microneme secretion. A comparative proteomics analysis of wild-type and ERK7 or AC9 depleted parasites highlighted the loss of known, as well as novel candidate proteins composing the apical complex. Conversely the accumulation of microneme proteins reflected the severe defect in exocytosis of these secretory organelles.

Results

ERK7 is essential for the stability of the conoid complex and the organization of the subpellicular microtubules in mature parasites

Three mitogen-activated protein kinases (MAPKs) are encoded in the genome of *T. gondii* and conserved across the superphylum of the Alveolata, except for TgMAPK-like 1 which is missing in haemosporidia and piroplasmida orders (Lacey et al. 2007; Talevich et al. 2011); their respective localization and function in *T. gondii* and *Plasmodium* spp. are summarized in Figure S1A. Briefly, TgMAPK-like 1 and TgMAPK2 are both involved in different steps of parasite replication (Suvorova et al. 2015; Hu et al. 2020), while TgERK7 is involved in conoid biogenesis thus hampering parasites invasion, egress and dissemination (O'Shaughnessy *et al.* 2020). The *Plasmodium* spp. orthologues of TgERK7 (Pf/PbMAP-1) and TgMAPK2 (PF/PbMAP-2) are

dispensable in the asexual blood stage with only Pf/PbMAP-2 required for male gametogenesis

(Khan et al. 2005; Rangarajan et al. 2005; Tewari et al. 2005; Hitz et al. 2020).

TgERK7 was carboxy-terminally tagged with the mAID-HA (ERK7-mAID-HA) at the endogenous

locus and shown to be tightly downregulated upon addition of auxin (IAA) (Figure S1B) and

localized at the apical cap of both mature and daughter cells (Figure S1C). As previously reported,

parasites depleted in ERK7 fail to form lysis plaques in a monolayer of human foreskin fibroblasts

(HFFs) after 7 days of IAA treatment (Figure S1D). Given the close relationship reported between

AC9 and ERK7, the cytoskeletal integrity was checked by deoxycholate (DOC) extraction. As

observed in absence of AC9 or AC10, the cytoskeleton of parasites depleted in ERK7 is

disassembled (Figure 1A). In addition to the loss of conoid, the APR has disappeared, leading to a

wide apical opening, explaining the detached SPMTs visible by DOC extraction (Figure 1B).

Ultrastructure expansion microscopy (U-ExM) was applied here for the first time on intracellular

parasites to compare with observations made on extracellular parasites, using the ERK7-mAID-HA

strain further modified by Ty-epitope tagging of AC2, a marker of the alveolin network (ERK7-

mAID-HA / AC2-Ty). As previously shown with AC9 deleted parasites, the absence of ERK7 led

to the loss of both conoid and APR in mature parasites, while the nascent daughter cells still

possessed an intact apical complex (Figure 1C). Overall AC2 was not dramatically affected by

ERK7 depletion; however, occasionally, AC2 was either not present apically or was markedly

reduced in some parasites (Figure 1C). The same observation was made with another apical cap

protein, ISP1 (Figure S1E).

The apical polar ring is lost in mature ERK7 depleted parasites

To further confirm the loss of the APR, three APR markers were epitope tagged in the ERK7-

mAID-HA strain, namely RNG1, APR1 and KinA (Tran et al. 2010; Leung *et al.* 2017). RNG1, a

very late maker of parasite division, was not visible at the APR of mature parasites (Figure 2A and

2B; Figure S2A); whereas the protein was still detectable by western blot analysis (Figure 2A).

Remarkably, the remaining RNG1 signal is mis-localized in the absence of ERK7, accumulating

mainly at the posterior pole of the parasites (Figure 2A; Figure S2A). The two other APR markers

were lost exclusively in the mature parasite but present in nascent daughter cells, confirming ERK7

role occurring late in parasite division (Figure 2C and 2D). In contrast to RNG1, APR1 and KinA

are both degraded based on western blot analysis (Figure 2C and 2D).

Rhoptries are disorganized in mature ERK7 depleted parasites by NHS-Ester staining

coupled to U-ExM

Next, to gain further insight into the morphology of the ERK7 depleted parasites, we coupled U-

ExM newly adapted to *T. gondii* (Tosetti *et al.* 2020) with the fluorescent dye N-

hydroxysuccinimide ester (NHS-Ester) able to bind to primary amines on proteins. Remarkably, this

combination allows the observation of a wide range of anatomical features while approaching the

resolution of electron microscopy (EM) (M'Saad & Bewersdorf 2020). This technique allowed the

visualization a wide spectrum of recognizable ultrastructures (movie 1 and 2) with a non-exhaustive

list of such features presented in individual stacks (Figure S3). As expected, in ERK7 depleted

parasites the conoid was clearly absent from mature parasites while still present in forming daughter

cells (Figure 3A, movie 3 and 4). Strikingly the neck of the rhoptries also detected with anti-RON9

antibodies, are still apical but dispersed compared to their apical positioning in presence of ERK7.

Interestingly, the so-called "neck" of the rhoptries seems to not extend all the way up to the conoid.

This observation was also confirmed with an anti-RON4 antibody (data not shown). Furthermore

as previously observed by EM for AC9 depleted parasites (Tosetti *et al.* 2020), in the absence of

ERK7 the plasma membrane was dilated at the apical pole (Figure 3A).

ERK7 depleted parasites are defective in induced microneme secretion

The dilated plasma membrane was examined by staining for the major surface antigen (SAG1) and

both in AC9 and ERK7 depleted conditions, the integrity of the membrane appeared compromised

in some intracellular parasites (Figure 4A). This leads to an apparent releases of few micronemes

outside the parasite body (Figure 4B; Figure S4A and S4B), as previously reported by EM in the

absence of AC9 (Tosetti *et al.* 2020).

Although AC9 and ERK7 are both essential for parasite survival, depleted mutants are not affected

in intracellular growth and replication (Figure 4C). In contrast, both mutants are seemingly able to

lyse the parasitophorous vacuole membrane (PVM) but remained trapped inside the host cell even

after the host cell detached in floating "bubbles" suggesting at least a partial microneme secretion

defect (Figure 4C). While depletion of ERK7 was reported not to impact on microneme secretion, a

severe exocytosis defect was observed in absence of AC9 (Back *et al.* 2020; O'Shaughnessy *et al.*

2020; Tosetti *et al.* 2020). The microneme secretion used by O'Shaughnessy *et al,* was based on a

luciferase assay applied previously to CPH1-mAID (Long et al. 2017). Here the induced secretion

assay was monitored by detection of cleaved released MIC2 in the supernatant and quantified by

western blot. Under these conditions, ERK7-mAID-HA treated with IAA showed a complete block

in microneme secretion as reported for AC9 (Figure 4D). As control, the CPH1-mAID strain was

generated (Figure S4C and S4D) and shown to exhibit a milder defect in microneme secretion

indicating that the discrepancy can be explained only in part by the type of assay used (Figure S4E).

Comparative proteomics of wild-type and ERK7/AC9 depleted parasites identifies novel candidate components of the apical complex

Given the dramatic disappearance of the structure as well as the protein markers of the conoid and

APR in both AC9 and ERK7 depleted parasites, we performed a comparative proteomic analysis by

mass spectrometry to gain further knowledge about the composition of the apical complex. The

datasets analysis revealed a good overall coverage in comparison to the hyperLOPIT dataset

(3,832 *T*. *gondii* detected proteins) (Barylyuk et al. 2020), with a total of 3477 and 3483 proteins

detected in ERK7 and AC9 samples, respectively (Figure 5C). In agreement with previously

published data and proteomics datasets, numerous known apical complex proteins disappeared in

both AC9 and ERK7 depleted parasites (Figure 5A, 5B, 5D and Figure 6A). Out of the 17 most

significantly depleted proteins in both datasets, 14 were already localized at the apical pole of the

parasite (Figure 6A). Overall, the analysis points toward at least a dozen of plausible novel

candidate proteins potentially associated to the apical complex.

Comparative proteomics of wild-type and ERK7 and AC9 depleted parasites show an accumulation of microneme proteins and components of the IMC

Beside the loss of components of apical complex and conversely, other distinct sets of proteins

accumulate in ERK7 and AC9 depleted parasites, respectively (Figure 5A, 5B, 5D and Figure 6B).

In the absence of ERK7, known microneme proteins as well as proteins predicted by hyperLOPIT

(Barylyuk et al. 2020) to be associated with micronemes are significantly more abundant. This is

easily accounted for by the severe microneme exocytosis defect observed in the mutant parasites. In

AC9 depleted parasites, microneme proteins also accumulated however other proteins predicted to

be localized at the IMC or dense granules are overrepresented. Out of the 17 proteins enriched in

both datasets and not localized experimentally to date, 7 share a cell cycle transcription profile

common to known IMC proteins (Figure 6B and Figure S5). A comprehensive list of the

comparative proteomics analysis of ERK7 and AC9 mutants is presented in Dataset 1.

Discussion

AC9 is a component of the alveolin network proposed to serve as regulatory platform for ERK7

kinase activity (Back *et al.* 2020). In this study we have revisited the role of ERK7 in the integrity

of the apical complex of *T. gondii* in light of unmatching phenotypic consequences compared to

AC9 depletion (O'Shaughnessy *et al.* 2020; Tosetti *et al.* 2020). ERK7 depleted parasites

phenocopied AC9 and AC10 depletion except that APR was shown to be still present and

microneme secretion was not affected. Here a new ERK7-mAID-HA strain was generated and

confirmed to be severely affected in invasion and egress from infected cells. U-ExM analysis

proved that like AC9 and AC10, ERK7 is required to maintain the conoid stability but also to preserve the tethering of the SPMTs in mature parasites. In contrast to the previous report, the depletion of ERK7 leads the disappearance of the APR with the degradation of APR1 and KinA and the mislocalization of RNG1. As a late marker of the APR, RNG1 might not be a structural component *per se* of the APR but rather associated to the structure only in mature parasites. The absence of the APR might explain why this protein end up in the residual body in ERK7 depleted parasites. The absence of APR also explains the wide apical opening observed in U-ExM and why detached individual microtubules are observed in deoxycholate extraction experiments. The APR is likely lost upon parasite final maturation as it presumably serves as MTOC for the SPMTs.

The NHS-Ester staining coupled with U-ExM offers novel opportunities to study in detail the internal structures of the parasite in its vacuolar niche. For example, in the less well-studied parasites like *Cryptosporidium spp*. in which a very sparse number of antibodies have been raised for localization studies (Wilke et al. 2018). The NHS-Ester could be a staining of great value as it can virtually reveal all structures within the cell. Also, as shown in this study, some structures are very well stained with this method. The "ring structures" systematically observed presumably corresponds to the micropore or the ring structures described with GAPM2 and GAPM3, two resident proteins of the IMC (Nichols et al. 1994; Harding et al. 2019). The basal complex of the parasite appears extremely dense with this technique, probably highlighting a density of protein complexes present at the basal pole of the parasite. In addition, very small structures in dividing parasites like the centrocone and centromeres are surprisingly evident with the NHS-Ester staining while only EM could achieve such level of details (Francia et al. 2020). Further improvements of these methods could potentially help retaining membrane or lipid-bound epitope as well as reaching greater expansion ratio without the need of heavy instruments (Damstra et al. 2021).

The induced microneme secretion monitored here by excretory-secretory antigen detection via western blot provides radically opposing outcome for ERK7-mAID-HA compared to the published results based on the luciferase assay. The presence of the conoid seems absolutely required for

micronerme secretion as no processed MIC2 can be detected in supernatant from parasites depleted

in ERK7, AC9 or AC10 (Tosetti *et al.* 2020). Since O'Shaughnessy *et al*, used a very sensitive

assay based on MIC2 fused to luciferase reporter we also generated a CPH1-mAID mutant

previously analyzed by the same method to include in the analysis (Long *et al.* 2017a). The two

assays are rather concordant for CPH1 depleted parasites although a mild defect in microneme

secretion was not detected in the luciferase-based assay (Long et al. 2017). The accumulation of

further processing of MIC2, only visible by western blot-based assay, reflects a defect in gliding

motility due to the lack of posterior translocation of MIC2, and hence the over-representation of

MIC2 protein trimmed by the apical subtilisin-like serin protease SUB1 (Lagal et al. 2010). It is

plausible that the luciferase assay is too sensitive for the detection of minor defects microneme

exocytosis, yet ERK7 depleted parasites have a very strong defect. Given that a fraction of these

mutants was shown to exhibit a dilated apical plasma membrane, they might be more prone to lysis

and could perturb the assay if parasites are not carefully washed before the experiment. In a

conceptually distinct assay, O'Shaughnessy et al, also assessed microneme secretion by monitoring

mVenus diffusion from the PV to the host cell cytoplasm. This diffusion is indicative of a

permeabilization of the PVM by the action of the secreted microneme protein perforin PLP1

(Kafsack et al. 2009). AC9 and ERK7 depleted parasites are indeed able to lyse the PVM but

remained trapped inside the host cell suggesting that a low level of microneme exocytosis was

occurring in a few intracellular parasites but not sufficient to trigger complete egress and hence

resulted in parasites trapped in floating detached host cells. Alternatively the discrepancies

observed in ERK7 mutant with regard to the disappearance of the APR and ability to secrete

microneme could be explained by revertant parasites that fail to respond anymore to IAA as

mentioned by the authors (O'Shaughnessy *et al.* 2020).

Capitalizing on the loss of APR and conoid that resulted in degradation of several apical complex

protein in both ERK7/AC9 depleted strains, we embarked on a proteomics analysis to compare the

mutants to wild-type parasites and identify novel components of the affected structures. The

comparative analysis between each replicate revealed a striking quantitative reproducibility for the vast majority of proteins. Consequently, the subsets of proteins disappearing or accumulating in the samples emerged very clearly. A large set of proteins already known or predicted to be associated with the apical complex were detected while only a small number of potential new apical complex proteins were found, potentially reflecting some technical limitation of mass spectrometry in detecting small, low abundant or not easily digested proteins. Alternatively, the combination of approaches used so far to determine the repertoire of apical complex protein might be almost comprehensive.

Of relevance, known, as well as predicted microneme proteins are accumulating in ERK7 and AC9 deleted parasites, concordantly with their impairment in microneme exocytosis. This tendency was more prominent in the ERK7 depleted parasites since twice as many proteins were found significantly enriched in those parasites compared to the AC9 depleted ones (88 versus 41). This might be due to the intrinsic nature and role of the two proteins: while AC9 is a structural component of the alveolin network, ERK7 is an active kinase (Back *et al.* 2020). Some proteins enriched in the absence of ERK7 could be affected by the absence of kinase activity and not only be a consequence of the loss APR and conoid and the disassembly of SPMTs. In the absence of ERK7 or AC9, some proteins predicted by hyperLOPIT to localize at the dense granules, ER, Golgi, nucleus also accumulate (Barylyuk *et al.* 2020). The experimental assessment of their localization will be instrumental as some protein such as the product of TGGT1_254870, mainly predicted to localize at the nucleus was previously localized at the apical pole of the parasite (Long *et al.* 2017). Similarly, the product of TGGT1_243200, predicted to localize to the dense granules is associated to the IMC of nascent daughter cells (Chen *et al.* 2016).

Overall, with novel technologies such as U-ExM, NHS-Ester staining and comparative proteomics mass-spectrometry, a deeper look into the role of ERK7 was taken in this study. Our observations revealed that the loss of ERK7, a kinase likely phosphorylating several components (known and unknown) of the apical complex, indeed dislocates the conoid complex. The severe microneme

secretion defect also strengthens previous observations that exocytosis of these organelles, and as a

consequence, proper egress, gliding, motility and invasion of intracellular parasites, might not be

achieved without the presence of a structurally intact conoid.

Figure legends

Figure 1 - ERK7 depletion caused major cytoskeletal defects at the apical pole

(A) Extracellular parasites were extracted with deoxycholate (DOC) and placed on gelatin

coverslips. Depletion of ERK7 caused the collapse of the microtubular cytoskeleton as only single

microtubules can be visualized by IFA with α-tubulin antibody suggesting that the APR is no more

present in those mutants. **(B)** Ultrastructure expansion microscopy (U-ExM) highlighted the

absence of the APR and consequently an enlargement of the apical pole in ERK7 depleted parasites.

(C) U-ExM was applied in intracellular conditions with AC2 tagged in the ERK7 inducible strain.

Upon addition of IAA, the APR and conoid are missing exclusively in the mother cell and in some

severe cases of SPMTs disorganization the AC2 staining disappears from the apical cap too

(arrowhead). Scale bars = 2 mm.

Figure 2 - Loss of apical markers in ERK7 depleted parasites

(A) RNG1, a late marker of parasite division, failed to be incorporated in most of the parasites. APR

and RNG1 signals could be detected in parasite cytoplasm/residual body (asterisk). Western blot

analysis confirmed that RNG1 is not degraded. **(B)** Quantification of apical RNG1 in ERK7

depleted parasites. **(C)** and **(D)** APR markers, APR1 and KinA, are incorporated very early during

daughter cells formation and are exclusively lost in mature parasites following deletion of ERK7.

Western blot analysis of extracellular parasites (no or very few daughter cells present) treated with

IAA showed that both markers are degraded. Arrowheads indicate mother cell apex. Scale bars = 2

mm.

Figure 3 - NHS-ester coupled with U-ExM

(A) Maximum projection pictures of ERK7 untreated and treated parasites (non-dividing and dividing) from videos 1, 2, 3 and 4. White arrowheads indicate mother cells conoid (blue arrowhead: intraconoidal microtubules); yellow arrowheads highlight daughter cells conoid; red arrowheads indicate deformation of the apical plasma membrane. Scale bars = 2 mm.

Figure 4 - ERK7 depleted parasites are defective in PM integrity and microneme secretion

(A) In both AC9 and ERK7 depletion, some parasites per vacuoles showed a defect in plasma membrane integrity at the apical tip highlighted by SAG1 staining and **(B)** leaking of micronemes (MIC4 staining). **(C)** Large bright field images of AC9 and EKR7 parasites treated for 48h with IAA. Depleted parasites showed no defects in intracellular growth; however, parasites remained trapped inside the floating host cell remnant suggesting a microneme secretion impairment. **(D)** Depletion of ERK7 caused a dramatic failure in parasite to secrete microneme when stimulated with ethanol (EtOH). Anti-MIC2 antibodies were used for secretion (white arrow: full length MIC2; black arrow: secreted MIC2), anti-catalase (CAT) to assess parasites lysis and anti-dense granule 1 (GRA1) for constitutive secretion; both pellets and supernatants (SN) were analyzed. Scale bars = 2 μm.

Figure 5 - Comparative proteomics of ERK7 and AC9 depleted parasites

(A) and **(B)** Volcano plots showing all the proteins found in ERK7 (A) and AC9 (B) mAID parasites and their differential abundance in auxin-treated parasites vs. untreated parasites. The most significant changes are colored and grouped based on their known and predicted (LOPIT) localization. On the left (in green) are proteins that are depleted and that majorly localize to the apical complex. The proteins on the right (in magenta) are the one that are enriched and are found majorly within the micronemes. **(C)** Venn diagrams showing the significant changes in ERK7 and AC9 (proteins individually depleted or enriched) and the common proteins depleted or enriched in both ERK7 and AC9 auxin-treated parasites. **(D)** Predicted localization of enriched and depleted proteins in both ERK7 and AC9 according to hyperLOPIT.

Figure 6 – Statistically significant proteins depleted/enriched in both strains

(A) and **(B)** Heat map showing the most significant changes in both ERK7 and AC9 (auxin-treated) conditions. In green (A) are the proteins that are depleted and in magenta (B) are the proteins that are enriched. Predicted (LOPIT) localizations and previously localized proteins are also indicated. . (Hoff et al. 2001; Hu et al. 2006; de Leon et al. 2013; Tomita et al. 2013; Katris et al. 2014; Graindorge et al. 2016; Huynh & Carruthers 2016; Wall et al. 2016; Chen et al. 2017; Leung *et al.* 2017; Long *et al.* 2017a; Long et al. 2017b; Barylyuk *et al.* 2020; Tosetti *et al.* 2020)

Supplementary materials

Figure S1

(A) Mitogen-activated protein kinases (MAPKs) present in *T. gondii* and *Plasmodium* spp. **1** (Suvorova *et al.* 2015), **2** (O'Shaughnessy *et al.* 2020), **3** (Wierk et al. 2013), **4** (Dorin-Semblat et al. 2007), **5** (Fang et al. 2018), **6** (Hitz *et al.* 2020), **7** (Hu *et al.* 2020), **8** (Khan *et al.* 2005), **9** (Rangarajan *et al.* 2005), **10** (Tewari *et al.* 2005).

(B) ERK7 was C-terminally tagged with the mAID-HA construct and was tightly downregulated upon addition of auxin (IAA). **(C)** ERK7 localized at the apical cap of both mature and daughter cells. Like AC9 and AC10, ERK7 is among the earliest markers appearing during endodyogeny, before the daughter cytoskeletal basket. **(D)** ERK7 depleted parasites failed to form lysis plaques after 7 days. TIR1 represents the parental strain. **(E)** Localization of another apical cap protein (ISP1) was not majorly impacted by ERK7 degradation; nevertheless, the apical region of some parasites per vacuoles is clearly enlarged and some shows no or reduced ISP1 signal (arrowhead). Scale bars = 2 μm.

Figure S2

(A) Raw pictures and quantification of RNG1 in ERK7 inducible strain. Scale bars = 2 μm.

Figure S3

Individual stacks from videos 1, 3 and 4 highlighting the broad spectrum of NHS-ester staining for ultrastructural elements and organelles. On top left the stack number. **(A)** Ring structures observed on parasite surface (from video 1). **(B)** Apicoplast (from video 1)**. (C)** Rhoptry neck (from video 1). **(D)** Post-Golgi vesicles (from video 3). **(E)** Apical micronemes collar just below the conoid (from video 1). **(F)** Centrocone (white arrowhead) and centromere (yellow arrowhead) during parasite division (from video 4).

Figure S4

(A) and **(B)** Additional pictures of membrane defects in both AC9 and ERK7 and microneme leaking. **(C)** CPH1 was terminally tagged with the mAID-HA construct and was tightly downregulated upon addition of IAA by IFA. **(D)** CPH1 depleted parasites failed to form lysis plaques after 7 days. TIR1 represents the parental strain. **(E)** Depletion of CPH1 caused a moderate defect in microneme secretion when stimulated with ethanol (EtOH). Anti-MIC2 antibodies were used for secretion (white arrow: full length MIC2; black arrow: secreted MIC2) and anti-dense granule 1 (GRA1) for constitutive secretion; both pellets and supernatants (SN) were analyzed. Scale bars = 2 μm.

Figure S5

Expression profile of proteins with no "experimental localization" presented in the figure 6B. Out of 17 proteins, 7 of them (thick lines / shades of magenta) share a similar expression profile with a typical IMC protein, namely IMC1 (cyan). The 10 others, with different expression profiles, are presented with thin gray lines. Data is retrieved from the ToxoDB website (Transcriptomic data "T. gondii ME49 Cell Cycle Expression Profiles (RH)" from Behnke *et al*).

Video 1

ERK7 untreated parasite (-IAA), non-dividing cells.

Video 2

ERK7 untreated parasite (-IAA), dividing cells.

Video 3

ERK7 treated parasite (+IAA), non-dividing cells.

Video 4

ERK7 treated parasite (+IAA), dividing cells.

Material and methods

Generation of strains

T. gondii were grown in human foreskin fibroblasts (HFFs, American Type Culture Collection-CRL

1634) in Dulbecco's Modified Eagle's Medium (DMEM, Gibco) supplemented with 5% fetal calf

serum (FCS), 2 mM glutamine and 25 mg/ml gentamicin. Human Foreskin Fibroblasts (HFF) were

obtained from the ATCC (CRL-2088 Ref# CCD1072Sk). ERK7-mAID-HA (TGGT1_233010) was

obtained by transfecting a gRNA obtained by Q5 site-directed mutagenesis kit (New England

Biolabs) on the pSAG1::Cas9-U6::sgUPRT vector (Shen et al. 2014) with primer 5'-

CACGCCGCATTTGTTGACTGGTTTTAGAGCTAGAAATAGC-3' and a KOD PCR obtained

with F primer 5'-TTTCAGTCTGCGTCCAAGACATACAACAGCGCTAGCAAGGGCTCGGG-

3' and R primer 5'-

CGGCTCTTCTTTGACACAAAGCAAGAGTCTATACGACTCACTATAGGG-3' as described in

(Brown et al. 2018). Depletion of ERK7-mAID-HA was obtained with 500 mM of auxin (IAA)

(Brown et al. 2017). Knock-in of APR1, KinA and RNG1 were performed as previously described

(Tosetti *et al.* 2020). Cloning were performed in *E. coli* XL-1 Gold chemo-competent bacteria.

Freshly egressed *T. gondii* tachyzoites were transfected by electroporation and mycophenolic acid

(25 mg/mL) and xanthine (50 mg/mL) (for HXGPRT cassette) or pyrimethamine (1 mg/ml) (for

DHFR cassette) were employed to select resistant parasites.

Immunofluorescence assay

Parasites were grown in HFF cells with cover slips in 24-well plates for 24–30 hr before fixation

with either 4% PFA/0.05% glutaraldehyde (PFA/GA) or cold methanol and neutralized in 0.1M

glycine/PBS for 5 min. For SAG1 staining, parasites were permeabilized with 0.1% saponin instead

of Triton X-100 which was used for all other IFA. The following antibodies were used: rabbit anti-

IMC1 (Frenal et al. 2014), rabbit anti-HA (Sigma), mouse α-acetylated α-tubulin (6-11B-1; Santa

Cruz Biotechnology), mouse anti-ISP1 (Beck et al. 2010), rabbit anti-ARO (Mueller et al. 2013),

mouse anti-Ty, anti-MIC2 and anti-ROP2-4 (gifts from J-F Dubremetz, Montpellier), anti-tubulin

AA344 scFv-S11B (β-tubulin) and AA345 scFv-F2C (α-tubulin), secondary antibodies Alexa Fluor

405-, Alexa Fluor 488-, Alexa Fluor 594-conjugated goat a-mouse/a-rabbit. For western blot,

secondary peroxidase conjugated goat a-rabbit or mouse antibodies (Sigma) were used. Confocal

images were obtained with a Zeiss laser scanning confocal microscope (LSM700; objective

apochromat 63x/1.4 oil). Ultrastructure expansion microscopy (U-ExM) was performed as

previously described (Tosetti *et al.* 2020) and images taken with a Leica TCS SP8 STED with

lightning deconvolution. For intracellular conditions, the parasites were seeded on non-confluent

host cells in order to guarantee the good expansion of the sample. All U-ExM gels were carefully

measured to assure a minimum expansion ratio of 4X. The NHS-Ester staining (Thermofisher,

DyLight™ 488 NHS Ester) was used at 5μg/mL and incubated 1 hour in PBS. Confocal and

expansion images were processed with ImageJ and LAS X, respectively. Images were obtained at

the Bioimaging core facility of the Faculty of Medicine at University of Geneva.

Plaque assay

433   HFF were inoculated with fresh parasites and grown for 7 days with or without IAA. HFF were

434   then fixed with PAF/GA, washed once with PBS and stained with crystal violet

Microneme secretion

After ~48 hr of grown parasites were harvested and centrifuged at 1000g. Pellets were then washed

twice in intracellular buffer pre-warmed at 37°C (5 mM NaCl, 142 mM KCl, 1 mM $MgCl_2$, 2 mM

EGTA, 5.6 mM glucose and 25 mM HEPES, pH 7.2). Next, parasites were incubated at 37°C for 15

min in DMEM containing 2% of ethanol (EtOH). After induction, parasite were centrifuged at 1000

g for 5 min at 4˚C and supernatants (SN) collected in different tubes and centrifuged at higher speed

(2000 g) for 5 min at 4˚C to clean from parasite debris; pellets were washed once in PBS. Pellets

and SNs were analyzed using anti-MIC2, anti-catalase (CAT; parasite lysis control) and anti-dense

granule 1 (GRA1; constitutive secretion control) antibodies by western blot.

Deoxycholate extraction

445   Freshly egressed parasites (-/+ IAA) were deposited on top of poly-L-Lysine-coated coverslips and

446   treated with 10 mM deoxycholate (20 min at room temperature). Parasites were fixed with cold

447   methanol for 8 min and immunofluorescence performed with α-acetylated α-tubulin antibody.

Mass spectrometry

Sample preparation: Parasite pellets were resuspended in 100 μl of 1 % RapiGest Surfactant

(Waters) in 0.1M triethylammonium bicarbonate (TEAB) and 7.5 mM Dithioerythritol (DTE).

Samples were heated for 5 min at 95°C. Lysis was performed by sonication (6 x 30 sec.) at 70%

amplitude and 0.5 pulse. Samples were kept 30 sec. on ice between each cycle of sonication.

Samples were centrifuged for 5 min. at 16'000 x g. Protein concentration was measured by

Bradford assay and 50 μg of each sample was subjected to protein digestion as follow: sample

volume was adjusted to 100 μl with 0.1M TEAB to obtain a final concentration of rapigest 0.1%. 2

μl of Dithioerythritol (DTE) 50 mM in distilled water were added and the reduction was carried out

at 60°C for 1h. Alkylation was performed by adding 2 μl of iodoacetamide (400 mM in distilled

water) during 1 hour at room temperature in the dark. Overnight digestion was performed at 37 °C

with 5 μl of freshly prepared trypsin (Promega; 0.2 μg/μl in TEAB 0.1M). To remove RapiGest,

samples were acidified with TFA, heated at 37°C for 45 min. and centrifuged 10 min. at 14'000 g.

Supernatants were then desalted with a C18 microspin column (Harvard Apparatus, Holliston, MA,

USA) according to manufacturer's instructions, completely dried under speed-vacuum and stored at

-20°C.

ESI-LC-MSMS: Samples were diluted in 50 μl of loading buffer (5% CH3CN, 0.1% FA) and 2 μl were injected on column. LC-ESI-MS/MS was performed on an Orbitrap Fusion Lumos Tribrid mass

spectrometer (Thermo Fisher Scientific) equipped with an Easy nLC1200 liquid chromatography system (Thermo Fisher Scientific). Peptides were trapped on a Acclaim pepmap100, C18, 3μm, 75μm x 20mm nano trap-column (Thermo Fisher Scientific) and separated on a 75 μm x 500 mm, C18 ReproSil-Pur (Dr. Maisch GmBH), 1.9 μm, 100 Å, home-made column. The analytical separation was run for 180 min using a gradient of H2O/FA 99.9%/0.1% (solvent A) and CH3CN/FA 99.9%/0.1% (solvent B). The gradient was run from 5 % B to 28 % B in 160 min, then to 40% B in 20 min, then to 95%B in 10 min with a final stay of 20 min at 95 % B. Flow rate was of 250 nL/min an total run time was of 210 min. Data-dependent analysis (DDA) was performed with MS1 full scan at a resolution of 120'000 FWHM followed by as many subsequent MS2 scans on selected precursors as possible within 3 second maximum cycle time

Database search: Peak lists (MGF file format) were generated from raw data using the MS Convert conversion tool from ProteoWizard. The peaklist files were searched against the *Toxoplasma gondii GT1* database (ToxoDB, release 48, 8460 entries) combined with an in-house database of common contaminant using Mascot (Matrix Science, London, UK; version 2.5.1). Trypsin was selected as the enzyme, with one potential missed cleavage. Precursor ion tolerance was set to 10 ppm and fragment ion tolerance to 0.02 Da. Variable amino acid modifications were oxidized methionine and deaminated (NQ). Fixed amino acid modification was carbamidomethyl cysteine. The Mascot searches were validated using Scaffold 4.11.1 (Proteome Software). Peptide identifications were accepted if they could be established at greater than 91.0% probability to achieve an FDR less than 0.1% by the Scaffold Local FDR algorithm. Protein identifications were accepted if they could be

established at greater than 96.0% probability to achieve an FDR less than 1.0% and contained at least 2 identified peptides. Protein probabilities were assigned by the Protein Prophet algorithm (Nesvizhskii et al. 2003). Proteins that contained similar peptides and could not be differentiated based on MS/MS analysis alone were grouped to satisfy the principles of parsimony.

Comparative proteomics: Bioinformatics analysis: For the comparative proteomics analysis, the average of the three replicates (minus and plus auxin, IAA) was used to compute the false discovery rate (FDR). A cutoff of 0.05 on the FDR was applied to obtain the most significant hits. For the volcano plots, the log10 of the FDR (y-axis) and the log2 fold change (FC), comparing minus and plus auxin treated parasites (x-axis) was computed. A cutoff of 0.5 was applied to the log2FC to mark the significant changes. The plots were generated using R statistical software, and the heat map with the common significant changes (depleted and enriched proteins) in both ERK7 and AC9 plus auxin was generated using GraphPad Prism v9. All the raw data can be found in Supplementary Table S1.

Acknowledgements

We thank the Bioimaging Core Facility and the Proteomic Core Facility (University of Geneva – CMU) for their excellent technical support. We thank the Brochet lab and the Guichard/Hamel lab for their helpful discussion to improve the U-ExM technology. This work was supported by the Swiss National Science Foundation (SNSF) 310030_185325 to DSF.

References

1. Back P.S., O'Shaughnessy W.J., Moon A.S., Dewangan P.S., Hu X., Sha J., Wohlschlegel J.A., Bradley P.J. & Reese M.L. (2020) Ancient MAPK ERK7 is regulated by an unusual inhibitory scaffold required for *Toxoplasma* apical complex biogenesis. Proc Natl Acad Sci U S A 117, 12164-73.
2. Barylyuk K., Koreny L., Ke H., Butterworth S., Crook O.M., Lassadi I., Gupta V., Tromer E., Mourier T., Stevens T.J., Breckels L.M., Pain A., Lilley K.S. & Waller R.F. (2020) A Comprehensive Subcellular Atlas of the *Toxoplasma* Proteome via hyperLOPIT Provides Spatial Context for Protein Functions. Cell Host Microbe 28, 752-66 e9.

518   3.   Beck J.R., Rodriguez-Fernandez I.A., de Leon J.C., Huynh M.H., Carruthers V.B., Morrissette N.S. &
519   Bradley P.J. (2010) A novel family of *Toxoplasma* IMC proteins displays a hierarchical organization
520   and functions in coordinating parasite division. PLoS Pathog 6, e1001094.

521   4.   Bisio H. & Soldati-Favre D. (2019) Signaling Cascades Governing Entry into and Exit from Host Cells
522   by *Toxoplasma gondii*. Annu Rev Microbiol 73, 579-99.

523   5.   Brown K.M., Long S. & Sibley L.D. (2017) Plasma Membrane Association by N-Acylation Governs
524   PKG Function in *Toxoplasma gondii*. mBio 8.

525   6.   Brown K.M., Long S. & Sibley L.D. (2018) Conditional Knockdown of Proteins Using Auxin-inducible
526   Degron (AID) Fusions in *Toxoplasma gondii*. Bio Protoc 8.

527   7.   Chen A.L., Moon A.S., Bell H.N., Huang A.S., Vashisht A.A., Toh J.Y., Lin A.H., Nadipuram S.M., Kim
528   E.W., Choi C.P., Wohlschlegel J.A. & Bradley P.J. (2017) Novel insights into the composition and
529   function of the *Toxoplasma* IMC sutures. Cell Microbiol 19.

530   8.   Damstra H.G.J., Mohar B., Eddison M., Akhmanova A., Kapitein L.C. & Tillberg P.W. (2021)
531   Visualizing cellular and tissue ultrastructure using Ten-fold Robust Expansion Microscopy (TREx).
532   bioRxiv, 2021.02.03.428837.

533   9.   de Leon J.C., Scheumann N., Beatty W., Beck J.R., Tran J.Q., Yau C., Bradley P.J., Gull K., Wickstead
534   B. & Morrissette N.S. (2013) A SAS-6-like protein suggests that the *Toxoplasma* conoid complex
535   evolved from flagellar components. Eukaryot Cell 12, 1009-19.

536   10.   Dorin-Semblat D., Quashie N., Halbert J., Sicard A., Doerig C., Peat E., Ranford-Cartwright L. &
537   Doerig C. (2007) Functional characterization of both MAP kinases of the human malaria parasite
538   *Plasmodium falciparum* by reverse genetics. Mol Microbiol 65, 1170-80.

539   11.   Dos Santos Pacheco N., Tosetti N., Koreny L., Waller R.F. & Soldati-Favre D. (2020) Evolution,
540   Composition, Assembly, and Function of the Conoid in Apicomplexa. Trends Parasitol 36, 688-704.

541   12.   Fang H., Gomes A.R., Klages N., Pino P., Maco B., Walker E.M., Zenonos Z.A., Angrisano F., Baum J.,
542   Doerig C., Baker D.A., Billker O. & Brochet M. (2018) Epistasis studies reveal redundancy among
543   calcium-dependent protein kinases in motility and invasion of malaria parasites. Nat Commun 9,
544   4248.

545   13.   Francia M.E., Bhavsar S., Ting L.M., Croken M.M., Kim K., Dubremetz J.F. & Striepen B. (2020) A
546   Homolog of Structural Maintenance of Chromosome 1 Is a Persistent Centromeric Protein Which
547   Associates With Nuclear Pore Components in Toxoplasma gondii. Front Cell Infect Microbiol 10,
548   295.

549   14.   Frenal K., Marq J.B., Jacot D., Polonais V. & Soldati-Favre D. (2014) Plasticity between MyoC- and
550   MyoA-glideosomes: an example of functional compensation in *Toxoplasma gondii* invasion. PLoS
551   Pathog 10, e1004504.

552   15.   Graindorge A., Frenal K., Jacot D., Salamun J., Marq J.B. & Soldati-Favre D. (2016) The Conoid
553   Associated Motor MyoH Is Indispensable for *Toxoplasma gondii* Entry and Exit from Host Cells. PLoS
554   Pathog 12, e1005388.

555   16.   Harding C.R., Gow M., Kang J.H., Shortt E., Manalis S.R., Meissner M. & Lourido S. (2019) Alveolar
556   proteins stabilize cortical microtubules in Toxoplasma gondii. Nat Commun 10, 401.

557   17.   Hitz E., Balestra A.C., Brochet M. & Voss T.S. (2020) PfMAP-2 is essential for male gametogenesis in
558   the malaria parasite *Plasmodium falciparum*. Sci Rep 10, 11930.

559   18.   Hoff E.F., Cook S.H., Sherman G.D., Harper J.M., Ferguson D.J., Dubremetz J.F. & Carruthers V.B.
560   (2001) *Toxoplasma gondii*: molecular cloning and characterization of a novel 18-kDa secretory
561   antigen, TgMIC10. Exp Parasitol 97, 77-88.

562   19.   Hu K., Johnson J., Florens L., Fraunholz M., Suravajjala S., DiLullo C., Yates J., Roos D.S. & Murray
563   J.M. (2006) Cytoskeletal components of an invasion machine--the apical complex of *Toxoplasma*
564   *gondii*. PLoS Pathog 2, e13.

565   20.   Hu K., Roos D.S. & Murray J.M. (2002) A novel polymer of tubulin forms the conoid of *Toxoplasma*
566   *gondii*. Journal of Cell Biology 156, 1039-50.

567   21.   Hu X., O'Shaughnessy W.J., Beraki T.G. & Reese M.L. (2020) Loss of the Conserved Alveolate Kinase
568   MAPK2 Decouples *Toxoplasma* Cell Growth from Cell Division. mBio 11.

569   22.   Huynh M.H. & Carruthers V.B. (2016) A *Toxoplasma gondii* Ortholog of *Plasmodium* GAMA
570   Contributes to Parasite Attachment and Cell Invasion. mSphere 1.

571   23.   Kafsack B.F., Pena J.D., Coppens I., Ravindran S., Boothroyd J.C. & Carruthers V.B. (2009) Rapid
572   membrane disruption by a perforin-like protein facilitates parasite exit from host cells. Science 323,
573   530-3.

574   24.   Katris N.J., van Dooren G.G., McMillan P.J., Hanssen E., Tilley L. & Waller R.F. (2014) The apical
575   complex provides a regulated gateway for secretion of invasion factors in *Toxoplasma*. PLoS Pathog
576   10, e1004074.

577   25.   Khan S.M., Franke-Fayard B., Mair G.R., Lasonder E., Janse C.J., Mann M. & Waters A.P. (2005)
578   Proteome analysis of separated male and female gametocytes reveals novel sex-specific
579   *Plasmodium* biology. Cell 121, 675-87.

580   26.   Lacey M.R., Brumlik M.J., Yenni R.E., Burow M.E. & Curiel T.J. (2007) *Toxoplasma gondii* expresses
581   two mitogen-activated protein kinase genes that represent distinct protozoan subfamilies. J Mol
582   Evol 64, 4-14.

583   27.   Lagal V., Binder E.M., Huynh M.H., Kafsack B.F., Harris P.K., Diez R., Chen D., Cole R.N., Carruthers
584   V.B. & Kim K. (2010) *Toxoplasma gondii* protease TgSUB1 is required for cell surface processing of
585   micronemal adhesive complexes and efficient adhesion of tachyzoites. Cell Microbiol 12, 1792-808.

586   28.   Leung J.M., He Y., Zhang F., Hwang Y.C., Nagayasu E., Liu J., Murray J.M. & Hu K. (2017) Stability and
587   function of a putative microtubule-organizing center in the human parasite *Toxoplasma gondii*. Mol
588   Biol Cell 28, 1361-78.

589   29.   Long S., Anthony B., Drewry L.L. & Sibley L.D. (2017a) A conserved ankyrin repeat-containing
590   protein regulates conoid stability, motility and cell invasion in *Toxoplasma gondii*. Nat Commun 8,
591   2236.

592   30.   Long S., Brown K.M., Drewry L.L., Anthony B., Phan I.Q.H. & Sibley L.D. (2017b) Calmodulin-like
593   proteins localized to the conoid regulate motility and cell invasion by *Toxoplasma gondii*. PLoS
594   Pathog 13, e1006379.

595   31.   M'Saad O. & Bewersdorf J. (2020) Light microscopy of proteins in their ultrastructural context. Nat
596   Commun 11, 3850.

597   32.   Mueller C., Klages N., Jacot D., Santos J.M., Cabrera A., Gilberger T.W., Dubremetz J.F. & Soldati-
598   Favre D. (2013) The *Toxoplasma* protein ARO mediates the apical positioning of rhoptry organelles,
599   a prerequisite for host cell invasion. Cell Host Microbe 13, 289-301.

600   33.   Nesvizhskii A.I., Keller A., Kolker E. & Aebersold R. (2003) A Statistical Model for Identifying Proteins
601   by Tandem Mass Spectrometry. Anal. Chem. 75, 4646-58.

602   34.   Nichols B.A., Chiappino M.L. & Pavesio C.E.N. (1994) Endocytosis at the micropore ofToxoplasma
603   gondii. Parasitology Research 80, 91-8.

604   35.   O'Shaughnessy W.J., Hu X., Beraki T., McDougal M. & Reese M.L. (2020) Loss of a conserved MAPK
605   causes catastrophic failure in assembly of a specialized cilium-like structure in *Toxoplasma gondii*.
606   Mol Biol Cell 31, 881-8.

607   36.   Rangarajan R., Bei A.K., Jethwaney D., Maldonado P., Dorin D., Sultan A.A. & Doerig C. (2005) A
608   mitogen-activated protein kinase regulates male gametogenesis and transmission of the malaria
609   parasite *Plasmodium berghei*. EMBO Rep 6, 464-9.

610   37.   Shen B., Brown K.M., Lee T.D. & Sibley L.D. (2014) Efficient gene disruption in diverse strains of
611   *Toxoplasma gondii* using CRISPR/CAS9. mBio 5, e01114-14.

612   38.   Suvorova E.S., Francia M., Striepen B. & White M.W. (2015) A novel bipartite centrosome
613   coordinates the apicomplexan cell cycle. PLoS Biol 13, e1002093.

614   39.   Talevich E., Mirza A. & Kannan N. (2011) Structural and evolutionary divergence of eukaryotic
615   protein kinases in Apicomplexa. BMC Evol Biol 11, 321.

616   40.   Tewari R., Dorin D., Moon R., Doerig C. & Billker O. (2005) An atypical mitogen-activated protein
617   kinase controls cytokinesis and flagellar motility during male gamete formation in a malaria
618   parasite. Mol Microbiol 58, 1253-63.

619   41.   Tomita T., Bzik D.J., Ma Y.F., Fox B.A., Markillie L.M., Taylor R.C., Kim K. & Weiss L.M. (2013) The
620   *Toxoplasma gondii* cyst wall protein CST1 is critical for cyst wall integrity and promotes bradyzoite
621   persistence. PLoS Pathog 9, e1003823.

622   42.   Tosetti N., Dos Santos Pacheco N., Bertiaux E., Maco B., Bournonville L., Hamel V., Guichard P. &
623   Soldati-Favre D. (2020) Essential function of the alveolin network in the subpellicular microtubules
624   and conoid assembly in *Toxoplasma gondii*. Elife 9.

625   43.   Tran J.Q., de Leon J.C., Li C., Huynh M.H., Beatty W. & Morrissette N.S. (2010) RNG1 is a late marker
626   of the apical polar ring in *Toxoplasma gondii*. Cytoskeleton (Hoboken) 67, 586-98.
627   44.   Wall R.J., Roques M., Katris N.J., Koreny L., Stanway R.R., Brady D., Waller R.F. & Tewari R. (2016)
628   SAS6-like protein in *Plasmodium* indicates that conoid-associated apical complex proteins persist in
629   invasive stages within the mosquito vector. Sci Rep 6, 28604.
630   45.   Wierk J.K., Langbehn A., Kamper M., Richter S., Burda P.C., Heussler V.T. & Deschermeier C. (2013)
631   *Plasmodium berghei* MAPK1 displays differential and dynamic subcellular localizations during liver
632   stage development. PLoS One 8, e59755.
633   46.   Wilke G., Ravindran S., Funkhouser-Jones L., Barks J., Wang Q., VanDussen K.L., Stappenbeck T.S.,
634   Kuhlenschmidt T.B., Kuhlenschmidt M.S. & Sibley L.D. (2018) Monoclonal Antibodies to Intracellular
635   Stages of Cryptosporidium parvum Define Life Cycle Progression In Vitro. mSphere 3.

A ERK7-mAID-HA

-IAA

α-Tubulin | Merge

IMC1

Zoom

+IAA

α-Tubulin | Merge

IMC1

Zoom

DOC extracted parasites

B ERK7-mAID-HA

-IAA

α/β-Tubulin

α/β-Tubulin

+IAA

α/β-Tubulin

α/β-Tubulin

C ERK7-mAID-HA / AC2-Ty

-IAA

α/β-Tubulin | Ty | Merge | Zoom

α/β-Tubulin | Ty | Merge | Zoom

+IAA

α/β-Tubulin | Ty | Merge | Zoom

α/β-Tubulin | Ty | Merge | Zoom

A

Significant (FDR < 0.05) in both ERK7 and AC9 +IAA

	log$_2$ fold-change	Fitness score
	-4 -2 0 2	-5 -2 0 2

Gene name	Gene ID	DEPLETED ERK7	AC9	Ess CRISPR	LOPIT	Experimental localisation	Reference
-	TGGT1_222350	-4.22	-1.84	-1.31	apical 1	Conoid	(Long et al. 2017a) ; (Barylyuk et al. 2020) ; (Koreny et al. 2020)
-	TGGT1_258090	-4.05	-2.63	-1.34	apical 1	Conoid	(Long et al. 2017a) ; (Barylyuk et al. 2020) ; (Koreny et al. 2020)
CPH1	TGGT1_266630	-3.35	-2.39	-4.16	apical 1	Conoid	(Long et al. 2017a) ; (Barylyuk et al. 2020) ; (Koreny et al. 2020)
APR1	TGGT1_315510	-3.11	-1.47	-0.05	apical 1	APR	(Leung et al. 2017)
RNG2	TGGT1_244470	-2.84	-0.94	-4.21	apical 1	APR	(Tosetti et al. 2020) ; (Katris et al. 2014) ; (Koreny et al. 2020)
SAS6-L	TGGT1_301420	-2.29	-1.55	-1.62	apical 1	Conoid	(de Leon et al. 2013) ; (Wall et al. 2016) ; (Koreny et al. 2020)
CaM3	TGGT1_226040	-3.23	-2.39	-3.25	apical 2	Conoid	(Long et al. 2017b)
MyoH	TGGT1_243250	-2.90	-1.90	-3.94	apical 2	Conoid	(Graindorge et al. 2016)
MLC7	TGGT1_315780	-2.48	-1.85	-0.12	apical 2	Apical complex	(Graindorge et al. 2016)
CAM2	TGGT1_262010	-2.12	-1.42	-0.81	apical 2	Conoid	(Long et al. 2017b) ; (Hu et al. 2006)
CAM1	TGGT1_246930	-2.08	-1.22	1.09	apical 2	Conoid	(Long et al. 2017b) ; (Hu et al. 2006) ; (Graindorge et al. 2016)
MLC5	TGGT1_311260	-1.46	-1.07	-0.33	apical 2	Apical complex	(Graindorge et al. 2016)
CST1	TGGT1_264660	-0.80	-1.10	0.85	dense granules 1	Cyst wall	(Tomita et al. 2013)
-	TGGT1_246040	-2.71	-3.83	-3.29	nucleus - chromatin	-	-
-	TGGT1_315720	-1.87	-1.64	-0.39	nucleus - chromatin	-	-
-	TGGT1_254870	-1.26	-0.61	0.64	nucleus - chromatin	Apical complex	(Long et al. 2017a)
-	TGGT1_274120	-3.05	-2.49	0.64	unknown	Apical complex	(Long et al. 2017a)

B

Significant (FDR < 0.05) in both ERK7 and AC9 +IAA

Gene name	Gene ID	ENRICHED ERK7	AC9	Ess CRISPR	LOPIT	Experimental localisation	Reference
-	TGGT1_301480	0.84	0.60	n/a	dense granules	-	-
IMC29	TGGT1_243200	0.90	1.04	-3.95	dense granules	daughter cell IMC	(Chen et al. 2016)
-	TGGT1_250955	1.09	1.16	n/a	dense granules	-	-
-	TGGT1_238950	0.76	0.93	-3.70	ER	-	-
-	TGGT1_269340	0.89	0.79	-3.34	ER	-	-
-	TGGT1_290980	1.05	0.71	-0.62	ER	-	-
-	TGGT1_281440	0.91	0.66	-4.82	Golgi	-	-
-	TGGT1_262710	1.49	1.15	-2.62	Golgi	Golgi	(Barylyuk et al. 2020)
-	TGGT1_235690	1.12	0.61	1.12	IMC	-	-
MIC10	TGGT1_250710	0.70	0.61	2.38	micronemes	Micronemes	(Hoff et al. 2001)
-	TGGT1_221180	0.94	0.77	0.93	micronemes	Micronemes	(Barylyuk et al. 2020)
GAMA	TGGT1_243930	0.95	0.80	0.25	micronemes	Micronemes	(Huynh et al. 2016)
CLP1	TGGT1_293770	1.03	0.63	0.70	micronemes	Micronemes	ToxoDB comment
-	TGGT1_249300	1.12	0.70	2.16	micronemes	-	-
-	TGGT1_248950	0.82	0.59	-3.72	mitochondrion - membranes	-	-
-	TGGT1_320080	0.89	0.81	-0.52	nucleus - chromatin	-	-
-	TGGT1_203820	2.07	0.91	1.39	nucleus - chromatin	-	-
-	TGGT1_222370	1.70	1.31	-0.07	PM - peripheral 1	-	-
-	TGGT1_220900	1.27	0.74	-3.42	PM - peripheral 2	-	-
-	TGGT1_294640	0.68	1.19	-4.79	unknown	-	-
-	TGGT1_310220	0.87	0.90	-1.36	unknown	-	-
-	TGGT1_294610	1.27	1.27	-0.98	unknown	-	-
-	TGGT1_282070	1.34	1.18	-2.70	unknown	-	-