Microbial Communities Performing Hydrogen Solventogenic Metabolism of Volatile Fatty Acids

[1,3]Gustavo Mockaitis*, [1]Guillaume Bruant, [2]Eugenio Foresti, [2]Marcelo Zaiat, [1]Serge R. Guiot.

[1]Anaerobic Technologies and Bioprocess Control group, Energy, Mining and Environment research

center, National Research Council Canada, 6100 Royalmount Avenue, H4P 2R2, Montreal, QC,

Canada.

[2]Biological Processes Laboratory, Center for Research, Development and Innovation in Environmental

Engineering, São Carlos School of Engineering, University of São Paulo (EESC/USP), Av. João

Dagnone, 1100, Santa Angelina, São Carlos, São Paulo 13563-120, Brazil.

[3]Interdisciplinary Research Group on Biotechnology Applied to the Agriculture and the Environment,

School of Agricultural Engineering, University of Campinas (GBMA/FEAGRI/UNICAMP), 501

Cândido Rondon Avenue, CEP, 13.083-875, Campinas, SP, Brazil.

Author	ORCID
Gustavo Mockaitis	0000-0002-4231-1056
Guillaume Bruant	0000-0002-8769-2868
Eugenio Foresti	0000-0002-7097-8990
Marcelo Zaiat	0000-0001-7336-9093
Serge Roger Guiot	0000-0002-2532-0620

12 (*Corresponding Author, gusmock@unicamp.br)

1 Abstract

This work evaluates four different physicochemical pretreatments (acidic, thermal, acidic-thermal and thermal-acidic) on an anaerobic inoculum used for alcohol production from acetate and butyrate. All experiments were conducted in single batches using acetate and butyrate as substrates at 30 °C and with a pressurized headspace of pure H_2 at 2.15 atm (218.2 MPa). Thermal and acidic-thermal pretreatments lead to higher production of both ethanol and butanol. Mathematical modelling shows that the highest attainable concentrations of ethanol and butanol produced were 122 mg L^{-1} and 97 mg L^{-1} for the thermal pretreatment (after 17.5 days) and 87 mg L^{-1} and 143 mg L^{-1} for the acidic-thermal pretreatment (after 18.9 days). Acetate was produced in all assays. Thermodynamic data indicated that a high H_2 partial pressure favoured solventogenic metabolic pathways. Finally, sequencing data showed that both thermal and acidic-thermal pretreatments selected mainly the bacterial genera *Pseudomonas*, *Brevundimonas* and *Clostridium*. The acidic-thermal pretreatment selected a bacterial community more adapted to the conversion of acetate and butyrate into ethanol and butanol, respectively. Thermal-acidic pretreatment was unstable, showing significant variability between replicates. Acidic pretreatment showed the lowest alcohol production.

2 Keywords

Anaerobic digestion, inoculum pretreatment, dark fermentation, alcohols, anaerobic microbiology

3 Background

The constant increase of prices of fossil fuels and the extensive land requirements for crop cultures targeting ethanol production is forcing the market to consider substitute avenues. The development of bioprocesses using organic residues as raw materials could be an alternative for fuel production. Anaerobic processes can be used to produce alcohol through solventogenic processes [1] and volatile fatty acids (VFAs) and hydrogen (H_2) through acidogenic processes [2]. The conversion of any wastes

into VFAs and H_2 has an immediate commercial interest. Both propionic and butyric acids are raw materials of great interest, with many applications in various sectors, such as pharmaceutical and chemical industries [3]. H_2 can be considered a raw material for subsequent processes and as an energy carrier to feed fuel cells [4].

Although anaerobic acidogenic processes could be used for wastes valorization, downstream processing of the VFAs obtained through those processes, such as bioconversion into their corresponding alcohols using H_2 produced concomitantly, could have an even greater economic interest [5,6]. Ethanol and butanol produced through such processes can be used as drop-in liquid fuels, which have a higher market value per unit of energy. This two-steps approach (acidogenic followed by solventogenic processes) for ethanol and butanol production could both improve solvent production and reduce the toxicity linked to acidogenic processes products [6,7].

The most important parameters influencing anaerobic solventogenic fermentations include pH, organic acids, nutrient limitation, temperature, oxygen, and inoculum source [1,6,8]. Most of the studies performed on ethanol and butanol production through anaerobic solventogenic processes focussed on using sugars as carbon source and on using microbial pure cultures, with the most studied bacterial genera being *Thermoanaerobacter*, *Thermoanaerobacterium* and *Caloramator* under thermophilic conditions [9–13], and *Clostridium* [14–18].

Using mixed microbial cultures rather than pure cultures to produce alcohols from wastes through biological processes is of great interest. Mixed cultures increase process stability, improve the resistance to both toxicity and microbial contaminations, and bring higher substrate flexibility [19,20]. However, to date, there is still a lack of fundamental knowledge on mixed cultures used as inoculum, especially when H_2 is used as an electron donor for the conversion of organic acids through solventogenic processes. Characterization of microbial communities capable of performing such processes is thus preponderant.

Enhancement of microbial communities through pretreatment of the inoculum is an effective way to

induce changes in the communities to improve process performance. This approach has already been

successfully tested, but mainly for optimizing the acidogenic step in H_2 production through dark

fermentation [21–23]. Pretreatments used in those studies consisted of modifications of the pH and

temperature applied to the inoculum. Since microbial communities involved in acidogenic processes

could use H_2 as an electron donor to shift their metabolism to produce alcohols [1,5,24], the same

inoculum pretreatments could be applied to improve solventogenic processes.

The present work evaluated the effects of four pretreatments of a mixed microbial population (acidic,

thermal, acidic-thermal and thermal-acidic) on its capacity to convert VFAs into alcohols, using H_2 as

electron donor and an equimolar mixture of acetate and butyrate as carbon sources. The composition

and dynamics of the mixed microbial community were analyzed.

4   Materials and Methods

4.1   Medium

The carbon source was an equimolar mixture of acetate and butyrate (17 mmol L^{-1}, which correspond

to 1,000 and 1,476 mg L^{-1} of acetate and butyrate, respectively). The nutrient medium (micro and

macro) composition was prepared considering previous work on anaerobic digestion microbiology [25]

and is described in Table 1. The initial pH was adjusted to 6.0.

Table 1 – Nutrients composition of the medium [25].

Nutrient	Concentration	Nutrient	Concentration	Nutrient	Concentration
NH_4Cl	1 g·L^{-1}	$MnCl_2 \cdot 4H_2O$	50 µg·L^{-1}	$(NH_4)_6Mo_7O_{24} \cdot 4H_2O$	50 µg·L^{-1}
NaCl	100 mg·L^{-1}	$AlCl_3$	50 µg·L^{-1}	Pyridoxine Chloride	10 µg·L^{-1}
$MgCl_2 \cdot 6H_2O$	100 mg·L^{-1}	$CoCl_2 \cdot 6H_2O$	50 µg·L^{-1}	HCl Concentrated	1 µL·L^{-1}
$CaCl_2 \cdot 2H_2O$	50 mg·L^{-1}	$NiCl_2 \cdot 6H_2O$	92 µg·L^{-1}	$Na_2SeO_3 \cdot 5H_2O$	100 µg·L^{-1}

$K_2HPO_4 \cdot 3H_2O$	400 mg·L^{-1}	EDTA	500 µg·L^{-1}	Nicotinic acid	5 µg·L^{-1}
$FeCl_2 \cdot 4H_2O$	2 mg·L^{-1}	Biotin	2 µg·L^{-1}	Pantothenic acid	5 µg·L^{-1}
H_3BO_3	50 µg·L^{-1}	Riboflavin	5 µg·L^{-1}	B_{12} Vitamin	0.1 µg·L^{-1}
$ZnCl_2$	50 µg·L^{-1}	Thiamine	5 µg·L^{-1}	p-aminobenzoic acid	5 µg·L^{-1}
$CuCl_2 \cdot 2H_2O$	38 µg·L^{-1}	Folic acid	2 µg·L^{-1}	Thioctic acid	5 µg·L^{-1}

4.2   Raw Inoculum

The sludge used as raw inoculum in all assays was a primary digestate of the Carleton Corner Farms

(Marionville, ON, Canada – 45º 11' 14.0" N; 75° 21' 54.1" W) collected in May 2013. The sludge was

sieved three times using a 2 mm mesh sieve to eliminate all inert and heterogeneous lignocellulosic

materials. The total volatile solids (TVS) content of the sieved sludge was 41 ± 3 mg TVS·L^{-1}. The

sludge was centrifuged (SorvalTM RC 6 Plus, Thermo Inc.) for 40 min at 10k min^{-1} and at 5°C. The

supernatant was discarded, and the pellet was resuspended in a phosphate buffer (500 mg·L^{-1} PO_4^{3-}),

using a homogenizer and disperser (Ultraturrax^{TM} T25, IKA Inc.) for 10 min at 15k·min^{-1}. The sludge

was then sonicated to disaggregate possible granules and biofilms, using a sonicator (Vibra Cell^{TM}

VC130, Sonics Inc.) with 30W of power. This step was repeated four times, on ice, with a time/volume

dependence relation of 4 s·mL^{-1}, with 2 minutes interval between sonications. These steps were carried

out to ensure the homogeneity of the inoculum and to wash possible organic dissolved materials

present in the sludge, which could be used as an alternative carbon source during the process. The pH

was corrected to 6.0 using a 1.0 M solution of HCl under vigorous stirring. This processed sludge was

considered as the control inoculum.

4.3   Pretreatments

Four different physicochemical pretreatments – acidic, thermal, acidic-thermal, and thermal-acidic –

were performed on the control inoculum. Pretreated inocula were then compared to each other, using

the control inoculum as reference. All pretreated inocula were submitted to a starvation process to

reduce the length of the lag phase before inoculation. This step consisted of incubating the inoculum

for 72 hours at 30°C, under stirring of 50 min^{-1}. The control inoculum was submitted to the same

starvation process and presented a TVS content of 31 ± 0 mg $TVS \cdot L^{-1}$.

Acidic pretreatment consisted of decreasing the raw inoculum pH to 3.0 with a 12 M HCl solution

under continuous stirring, followed by an incubation of 24 hours at 30°C, with stirring of 50 min^{-1}. pH

was then increased to 6.0 using a 2 M NaOH solution under stirring, followed by an incubation of 24

hours at 30°C under stirring of 50 min^{-1}. In both pH decreasing and increasing inoculum, pH was

controlled every hour for the five first hours of incubation to assure its stability. The acidic pretreated

inoculum had a TVS of 29 ± 1 mg $TVS \cdot L^{-1}$.

The thermal pretreatment consisted of heating the control inoculum at 90°C for 20 minutes, under

stirring and using a water batch. The inoculum was then immediately transferred to an ice batch until it

reached room temperature (23°C). The thermally pretreated inoculum had a TVS of 31 ± 0 mg $TVS \cdot L^{-1}$.

The acidic-thermal and thermal-acidic pretreatments consisted of performing the two pretreatments

sequentially, as indicated by the pretreatment name. The second pretreatment was performed

immediately after the first one. The acidic-thermal pretreated inoculum had a TVS of 36 ± 3 mg

$TVS \cdot L^{-1}$ and thermal-acidic pretreated inoculum had a TVS of 32 ± 0 mg $TVS \cdot L^{-1}$.

4.4 Experimental Setup

Initial physicochemical parameters for each experiment (pretreatments and control) and initial

concentration of acetate, butyrate (before inoculation) and inoculum are shown in Table 2. Each

experiment was carried out in quintuplicates, using 538 ± 3 mL sealed glass bottles, with an initial

working volume of 110 mL. All the bottles were incubated upside down to avoid any diffusion of H_2

through the bottle's cap. All experiments were performed at 30°C under a 150 min^{-1} stirring. At the

beginning of each experiment, the headspace of each bottle was totally replaced by pure H_2 (99.99%) at

a pressure of 2.39 ± 0.08 atm (242.2 ± 7.9 MPa). Such headspace replacement was repeated after each

sampling to ensure constant pressure and composition all along with the experiment.

Table 2 – Initial conditions (prior to inoculation) for each assay.

Assay	pH	Acetate mg·L^{-1}	Butyrate mg·L^{-1}	Inoculum mg TVS·L^{-1}	Total H_2 mmol	Dissolved H_2[*] mmol·L^{-1}
Control	5.89	1153	1727	11.3 ± 0.2 (5)	37.9 ± 1.6 (5)	1.69
Acidic	5.94	1139	1708	11.0 ± 0.2 (5)	39.2 ± 0.2 (5)	1.74
Thermal	6.07	1143	1712	12.8 ± 3.7 (5)	40.2 ± 0.1 (5)	1.78
Acidic-thermal	5.84	1233	1846	12.7 ± 0.2 (5)	40.8 ± 0.2 (5)	1.81
Thermal-acidic	5.85	1213	1816	11.3 ± 0.1 (5)	39.7 ± 1.3 (5)	1.77

Values in parenthesis are the number of replicates. * Dissolved H_2 was estimated through Henry's Law [26,27].

Initial analyses of the liquid phase at the beginning of the essay confirmed the absence of significant

amounts of alcohols, mono or disaccharides, and fatty acids other than acetate and butyrate.

4.5   Physicochemical Analyses

pH monitoring was performed with a portable potentiometer (Accumet[TM] AP115, Fisher Scientific[TM]),

with a microprobe electrode (Accumet[TM] 55500-40, Cole Parmer[TM]), using the method 4500-H^+ B,

described by APHA [28]. TVS analyses were performed in quintuplicate, following method 2540-E,

accordingly to APHA [28]. The pressure inside the bottles was measured using a digital manometer

(DM8200, General Tools & Instruments[TM]) with a range pressure of 0 - 6,804 atm (0 – 689.5 MPa).

Dissolved CO_2 was measured through alkalinity determination by potentiometric titration [29,30].

Mono, disaccharides, and organic fatty acids were analyzed using a Waters[TM] HPLC, which consisted

of a pump (model 600) and an autosampler (model 717 Plus). The system was equipped with a

refractive index detector (model 2414) for mono and disaccharides analyses. Organics acids were

monitored from the same samples using the same equipment through a linked photodiode array detector

(model 2996). A Transgenomic[TM] ICSep IC-ION-300 (300 mm x 7.8 mm outer diameter) column was

used to separate all compounds and was operated at 35°C. The mobile phase was 0.01 N H_2SO_4 at 0.4

mL min^{-1} under an isocratic flow.

Alcohols (methanol, ethanol, acetone, 2-propanol, tert-butanol, n-propanol, sec-butanol, and n-butanol)

were measured on an Agilent[TM] 6890 gas chromatograph (GC) equipped with a flame ionization

detector [31].

100 µL gas samples, obtained using a gas-tight syringe (model 1750, Hamilton[TM]), were used for gas

composition (H_2, CO_2, CH_4 and N_2) measurements with a GC (HP 6890, Hewlett Packard[TM]) equipped

with a thermal conductivity detector (TCD) and an 11 m x 3.2 mm 60/80 mesh packed column

(Chromosorb[TM] 102, Supelco[TM]). The column temperature was held at 50 ºC for the entire run (4 min).

The carrier gas was argon. The injector and detector were maintained at 125°C and 150°C,

respectively.

4.6   16S rRNA gene Sequencing and microbial characterization

Bacterial 16S rRNA genes (V2 region) were amplified using the set of primers 16S-F343 IonA L01

(343-357; 5′ TACGGRAGGCAGCAG 3′) and 16S-R533 Ion P1 (516-533; 5′

ATTACCGCGGCTGCTGGC 3′) [32]. A sample-specific multiplex identifier was added to each

forward primer and an Ion Torrent adapter to each primer. DNA extraction was conducted [33]. DNA

was then purified [34,35]. Amplification reactions were performed in a final volume of 20 µL, which

contained 1 µL of DNA, 0.5 µM of each primer, 7.5 µL of RNAse free H_2O and 10 µL of 2X

HotStarTaq[TM] Plus Master Mix (HotStarTaq[TM] *Plus* Master Mix Kit, Qiagen, USA). PCR conditions

were an initial denaturation of 5 min at 95°C followed by 25 cycles of 30 s at 95°C, 30 s at 55°C, and

45 s at 72°C, with a final elongation step of 10 min at 72°C. PCR products were purified and quantified

using a QIAquick™ gel extraction kit (Qiagen, USA) and a Quant-iT PicoGreen™ double-stranded

DNA quantitation kit (Life Technologies Inc., USA) according to manufacturer's instructions. The

pooled amplicons were then sequenced using the Ion TorrentTM (Life Technologies Inc., USA)

sequencing platform with a 314 chip. Bacterial 16S rRNA gene sequences generated were and analyzed

using the ribosomal database project (RDP) classifier [36], using a bootstrap confidence cutoff of 50%,

as recommended by RDP classifier [37] for short sequences (less than 250 bp). Before analysis,

sequences shorter than 75 bp and sequences with unidentified bases (N) were removed.

4.7   Kinetics

Kinetic parameters for ethanol and butanol production were calculated using a modified Boltzmann

sigmoidal model, as shown by Equation 1. The model was modified to incorporate parameter r_{max} as

the maximum rate of the process (mg L^{-1} d^{-1}).

$$C(t) = C_{max} - \frac{C_{max}}{1 + e^{\left(\frac{4 \cdot r_{max}(t - t_m)}{C_{max}}\right)}} \qquad\qquad \text{Equation 1}$$

174   Where $C(t)$ is the function of concentration in respect of time (mg L^{-1}), t is time (d), C_{max} is the

175   maximum concentration reached (mg L^{-1}), and t_m is the value of time when $\frac{dC(t_m)}{dt} = r_{max}$ (d).

176   From some of the calculated parameters showed in Equation 1, is possible to calculate the length of

177   both lag (t_i, d) and exponential (t_e, d) phases, as depicted in Equation 3 and 4, respectively.

$$t_i = t_m - \frac{C_{max}}{2 \cdot r_{max}} \qquad\qquad \text{Equation 3}$$

$$t_e = \frac{C_{max}}{r_{max}}$$
<div align="right">Equation 4</div>

All fittings were performed using the software Microcal Origin Pro^{TM} 9.0, using a Levenberg-

Marquardt algorithm for fitting and initializing the equations parameters.

4.8   Metabolisms' Gibb's free energy

Gibb's free energy values were calculated for all pathways depicted in Figure 1, except for pathway 1,

for both initial (ΔG^I_r) and final (ΔG^F_r) conditions and were compared with the free energy for standard

conditions (ΔG^o_r). ΔG^o_r values for each reaction were estimated [38,39], considering 1 mol of each

product and reagent at STP. ΔG^I_r and ΔG^F_r values were calculated using the Nernst equation at 30 °C

and considered the concentrations of the metabolites presented in Table 4. To make all calculations

possible, compounds that were not detected through the experiments, such as butanol, propionate,

propanol, and ethanol, were assumed to be present, but at the same concentration of their respective

lower detection limit for each methodology. This represents 10^{-3} mmol L^{-1} for volatile acids and

alcohols and 10^{-1} mmol L^{-1} for CO_2.

5   Results and Discussion

5.1   Metabolic Model and Molar Balances

For all experiments, gas composition of the headspace was monitored prior to its replacement and H_2

pressure was constantly kept at 2.15 ± 0.10 atm. H_2 was the sole gas detected in the headspace in all

cases, except for the control experiment, which showed a methane production of 0.24 mmol (5.37 mL

at standard temperature and pressure (STP)). Initial and final concentrations of the main metabolites

(added or produced) and of dissolved H_2 and CO_2 are presented in Table 3. Dissolved H_2 was not

measured but estimated through Henry's law [26,27], and its concentration was considered constant,

due to its continuous replacement. Dissolved CO_2 was calculated from the alkalinity value measured at

the beginning of each assay. CO_2, O_2 and N_2 were never detected at the headspace. Acetone, lactate,

and methanol were only sporadically detected in the liquid phase, in traces concentrations (below 1 mg

L^{-1}). Neither propionate nor alcohols were detected at the beginning of each experiment.

Table 3 – Mean concentrations of all metabolites detected and of dissolved H_2 and CO_2.

Metabolite[a]	Concentration (mg L^{-1})									
	Control		Acidic		Thermal		Acidic-thermal		Thermal-acidic	
	Initial	Final	Initial	Final	Initial	Final	Initial	Final	Initial	Final
Acetate	1225 ± 1	1575 ± 40	1222 ± 3	1492 ± 24	1206 ± 2	1722 ± 120	1235 ± 3	1645 ± 44	1224 ± 2	1447 ± 26
Butyrate	1831 ± 3	1773 ± 24	1828 ± 1	1935 ± 14	1791 ± 4	1796 ± 17	1840 ± 5	1664 ± 25	1831 ± 4	1738 ± 58
Propionate	-	42 ± 12	-	35 ± 4	-	107 ± 25	-	52.8 ± 9.2	-	40.1 ± 5.4
Ethanol	-	40 ± 6	-	14 ± 2	-	104 ± 7	-	94 ± 9	-	33 ± 6
Propanol	-	-	-	0.3 ± 0.2	-	-	-	1.8 ± 0.1	-	1.0 ± 0.3
Butanol	-	49 ± 8	-	11 ± 2	-	83 ± 3	-	154 ± 14	-	61 ± 25
Dissolved H_2	1.6 ± 0.1		1.7 ± 0.0		1.6 ± 0.1		1.7 ± 0.1		1.7 ± 0.1	
Dissolved CO_2	3.07	-	2.40	-	3.30	-	1.56	-	2.19	-
Duration (d)	29.2		29.1		29.1		29.1		29.0	

[a]All values are expressed in mg L^{-1} except for dissolved H_2 and CO_2 expressed in mmol L^{-1}. Mean concentrations of all volatile acids and
alcohols was calculated considering 5 replicates. The amount of H_2 was kept constant in all experiments, there were no difference in its
initial   and   final   concentrations.   Dissolved H_2 mean   concentration   considered   50   replicates.

Acetate and butyrate concentrations along the process demonstrated acetate production in all cases.

Only minor variations of butyrate concentrations (slight decrease for the control and the acidic-thermal

and thermal-acidic pretreatments, low production for the acidic and the thermal pretreatments) were

observed. Such results suggested an alternative organic matter input in the system. This hypothesis is

consistent with the decrease of the concentration of total volatile solids (TVS) that was observed for all

conditions tested, as a direct effect of the different pretreatments on the biomass, which might have

resulted in partial microbial cell death. Such non-living cells constituted non-soluble organic matter

which was probably hydrolyzed and then consumed as a supplementary carbon source. Based on well-

known anaerobic acidogenic and solventogenic metabolisms [1,24], and considering the inoculum as

the only possible alternative source of organic matter, an alternative metabolic model including this

alternative contribution and considering the metabolites (added and produced) presented in Table 3,

was developed, as shown in Figure 1.

Figure 1 depicts butanol as the only product from lysed inoculum (pathway 1). Pathway 1 seems the

only way to explain butanol production maintaining the butyric acid concentration constant since it is

thermodynamically unfeasible to form butanol from acetate. Acetate could only be produced from

butyrate, considering the high concentration of hydrogen and lack of CO_2 in all assays, indicating that

acetate was produced through acetogenesis from butyrate (pathway 2) or homoacetogenesis (pathway

7) after initial dissolved CO_2 was consumed.

Metabolic Pathways

(1)	Organic matter in the inoculum to butyrate	
(2)	Acetogenesis from butyrate	$But^- + 4H_2O \rightarrow Ac^- + 2CO_2 + 6H_2$
(3)	Solventogenesis from butyrate to butanol	$But^- + H^+ + 2H_2 \rightarrow But(OH) + H_2O$
(4)	Acidogenesis from butyrate to propionate	$But^- + 2H_2O \rightarrow Prop^- + CO_2 + 3H_2$
(5)	Solventogenesis from propionate to propanol	$Prop^- + H^+ + 2H_2 \rightarrow Prop(OH) + H_2O$
(6)	Solventogenesis from acetate to ethanol	$Ac^- + H^+ + 2H_2 \rightarrow Eth(OH) + H_2O$
(7)	Homoacetogenesis	$2CO_2 + 4H_2 \rightarrow Ac^- + H^+ + 2H_2O$

Figure 1 – Alternative metabolic model proposed for the solventogenic process using butyrate and
acetate as substrates. Pathways in orange are exergonic; pathways in blue are endergonic (at 30 °C, pH
of 6.0, with a constant CO_2 concentration of 10^{-3} mmol L^{-1}, in initial conditions as depicted in Table 3).

The metabolic model depicted in Figure 1 was used to perform a molar balance of all carbon inputs and

outputs during the process (Table 4) and estimates the butyrate input from inoculum. All balances were

performed on a 1 L basis and derived from initial and final experimental values obtained in all

conditions (Table 3), except for the organic matter from the inoculum (pathway 1) which was

expressed in butyrate equivalent.

Table 4 – Molar balance of all metabolites detected in all metabolic pathways of each experiment.

Metabolite	Pathway[a]	Molar Balance (mmol)[b]				
		Control	Acidic	Thermal	Acidic-thermal	Thermal-acidic
Butyrate	Initial[*]	21.0	21.0	20.6	21.1	21.0
	Final[*]	20.4	22.2	20.6	19.1	20.0
	1	*+ 3.16*	*+ 3.56*	*+ 6.92*	*+ 4.73*	*+ 1.96*
	2	- 2.53	- 1.73	- 4.34	- 3.92	- 1.57
	3	- 0.66	- 0.15	- 1.12	- 2.08	- 0.82
	4	- 0.57	- 0.48	- 1.46	- 0.75	- 0.57
Butanol	3[*]	+ 0.66	+ 0.15	+ 1.12	+ 2.08	+ 0.82
Propionate	4*	+ 0.57	+ 0.48	+ 1.46	+ 0.75	+ 0.57
	5	-	-	-	- 0.03	- 0.02
Propanol	5[*]	-	-	-	+ 0.03	+ 0.02
Acetate	Initial[*]	20.7	20.7	20.4	20.9	20.7
	Final[*]	26.7	25.3	29.2	27.9	24.5
	2	+ 2.53	+ 1.73	+ 4.34	+ 3.92	+ 1.57
	6	- 0.87	- 0.30	- 2.26	- 2.04	- 0.72
	7	+ 4.35	+ 3.17	+ 6.72	+ 5.08	+ 2.95
Ethanol	6[*]	+ 0.87	+ 0.30	+ 2.26	+ 2.04	+ 0.72
Dissolved CO_2	Initial*	3.07	2.40	3.30	1.56	2.19
	2	+ 5.05	+ 3.46	+ 8.68	+ 7.85	+ 3.14
	4	+ 0.57	+ 0.48	+ 1.46	+ 0.75	+ 0.57
	7	- 8.69	- 6.34	- 13.4	- 10.2	- 5.90

236 [*]Values obtained experimentally (derived from initial and final concentrations as indicated in Table 3).
237 [a]Metabolic pathways described in Figure 1.
238 [b]Balance on a 1 L basis; "+" indicates a production; "-" indicates a consumption.
239

According to the mass balance shown in Table 4, CO_2 was produced by converting butyrate into

acetate (pathway 2) and by acidogenesis from butyrate to propionate (pathway 4). This CO_2 was then

consumed to form acetate through a homoacetogenic pathway (pathway 7). This hypothesis was

supported by the absence of gaseous CO_2 in all experiments, followed by an increase in acetate

concentration. This CO_2 absence may be linked to microbial communities more adapted to convert

butyrate and CO_2 into acetate (pathways 2 and 7, respectively) rather than to the conversion of acetate

into ethanol (pathway 6). This metabolic model also indicates that butyrate was mainly consumed to

produce acetate (pathway 2) in all studied conditions and the organic matter from the inoculum

(pathway 1) represented an external input since butyrate concentrations were almost constant in all

assays, although consumed to form butanol (pathway 3). Ethanol and butanol were produced (pathways

3 and 6) in all experiments. These pathways were more active in the experiments with thermal and

acidic-thermal pretreatments, as these conditions showed the highest ethanol and butanol production

among all other experiments. Although acidogenesis from butyrate to propionate (pathway 4) was

active in all conditions, conversion of propionate into propanol (pathway 5) was not an important

metabolic pathway. Propanol production was not expressive in neither experiments since it showed the

highest concentration of 1.82 mg L^{-1} in acidic-thermal pretreatment essay.

The successful closure of molar balances shown in Table 4, with a stoichiometrically balanced sum of

all inputs and outputs (considering the contribution of the inoculum), indicates that the metabolic

model proposed in Figure 1 accurately represents the solventogenic processes occurring in all studied

conditions. An energy balance based on the mass balance depicted in Table 4 was calculated to

estimate Gibb's free energy values for each pathway for both initial and final conditions (Figure 2).

Those values provide an overview of the thermodynamic feasibility of each pathway involved in the

solventogenic process and tend to validate the metabolism proposed in Figure 1.

Metabolic Pathway	ΔG^o_r	Metabolism Free Energy (kJ mol^{-1})									
		Control		Acidic		Thermal		Acidic-thermal		Thermal-acidic	
		ΔG^I_r	ΔG^F_r	ΔG^I_r	ΔG^F_r	ΔG^I_r	ΔG^F_r	ΔG^I_r	ΔG^F_r	ΔG^I_r	ΔG^F_r
2	109	234	-158	221	-162	237	-156	200	-156	217	-160
3	-20,6	-291	-129	-291	-168	-290	-116	-291	-99	-291	-123
4	55,7	-129	-170	-135	-177	-127	-147	-146	-163	-138	-171
5	-24,4	-47	-205	-47	-201	-47	-228	-47	-125	-47	-135
6	-20,5	-290	-129	-290	-154	-290	-107	-291	-109	-290	-131
7	-40	-67	339	-54	337	-70	341	-33	340	-50	337

-400	Exergonic	Endergonic	400

Figure 2 – Gibbs' free energy for initial (ΔG^I_r), final (ΔG^F_r) and standard (ΔG^o_r) conditions for each pathway (as defined in Figure 1) in each experiment. Greenish values (< 0) are exergonic and reddish values (> 0) are endergonic.

Energetic profiles could help to anticipate changes within metabolic pathways. As shown in Figure 2, the estimated ΔG^o_r values indicate that acetogenesis from butyrate and acidogenesis from butyrate to propionate (pathways 2 and 4, respectively) are theoretically thermodynamically unfeasible metabolisms. Nevertheless, these pathways became thermodynamically feasible due to the absence (or low concentration) of dissolved CO_2 at the course of each essay. Acetogenesis from butyrate (pathway 2) was likely inhibited at the beginning of each experiment due to the concentration of H_2 and CO_2. During the experiment, CO_2 was progressively consumed through homoacetogenesis (pathway 7), which rendered pathway 2 thermodynamically feasible. At low concentrations of CO_2 as found after the beginning of each essay, the homoacetogenic pathway might have become thermodynamically unfeasible. Gibb's free energy values found for these pathways could explain the production of acetate observed in all studied conditions, all along the process. Metabolic pathways 3 (solventogenesis from butyrate to butanol) and 6 (solventogenesis from acetate to ethanol) were thermodynamically feasible

281 in all studied conditions all along the process, explaining the observed production of such alcohols.

282 Metabolic pathway 4 (acidogenesis from butyrate to propionate) was unfeasible at the ending of each

283 condition, showing that propionate production was stationed in a low concentration at some point of the

284 experiments.

285
Figure 3 – Evolution of the standard Gibbs' free energy (ΔG^o_r) in function of the partial pressure of H_2.
Evolution for each metabolic pathway (as defined in Figure 1), at 30 °C, and with a standard
concentration of 1.0 mol L^{-1} for each reagent other than H_2. Shadowed area indicates
thermodynamically feasible reactions.

As described previously, H_2 has a significant role as a co-substrate in the anaerobic process energetics

[40,41]. As shown in Figure 3, solventogenesis of butanol from butyrate (pathway 3) is less sensitive to

low pp_{H_2} than solventogenesis of ethanol and propanol, respectively from acetate (pathway 6) and

propionate (pathway 5). By extrapolation, equilibrium ($\Delta G^o_r = 0$) in pathway 3 would be achieved at a

pp_{H_2} of $4 \cdot 10^{-4}$ atm (43.7 Pa), implying that solventogenesis of butanol could theoretically be carried

out at very low pp_{H_2}. It is also possible to infer from Figure 3 that at pp_{H_2} higher than 0.62 atm

(62.5 MPa), acidogenesis of propionate (pathway 4) and acetogenesis of acetate (pathway 2) from

butyrate will stop, shutting down the production of acetate, propionate and H_2. In addition to low CO_2

concentrations, high pp_{H_2} thus represents an important factor that renders such process

thermodynamically feasible and favours solventogenesis from VFAs.

5.2   Alcohols and VFAs metabolisms

As shown in Figure 4, the highest concentrations of alcohols were obtained for both thermal and acidic-

thermal pretreatments, with the best rate observed for the acidic-thermal pretreatment, especially for

butanol production. On the opposite, the lowest alcohol production was observed for the acidic

pretreatment. Acetate and butyrate concentrations along the essays confirmed the contribution of the

inoculum as an important source of organic matter as the substrate for alcohol production, since in all

conditions tested, acetate concentration increased, and butyrate concentration increased or only slightly

decreased. These results reinforced the hypothesis described in Figure 1 and Table 4 of an acetate

production through acetogenesis of butyrate (pathway 2) and homoacetogenesis (pathway 7), and

preferentially through pathway 2 due to low concentrations of CO_2 at the beginning of all assays.

Figure 4 – Kinetics of VFAs (acetate and butyrate) and alcohols (ethanol and butanol) productions for each studied condition. Red dotted- and blue dashed-lines represent the modified Boltzmann model fittings for ethanol and butanol concentrations, respectively.

Figure 5 – Behaviour of propionate concentration for each studied condition.

As shown in Figure 5, propionate was produced in all conditions tested. No significant difference was

observed in the concentrations produced (42.4 ± 10.0 mg L^{-1}), except for the thermal pretreatment

essay, for which the level of production was 2.5 times higher than in all other conditions. As presented

in Table 3, propionate was then consumed and converted into propanol (pathway 5, Figure 1) for the

acidic, acidic-thermal and thermal-acidic pretreatments. Despite the favourable thermodynamics of this

reaction (Figure 2), propanol was only produced in trace amounts. Oxidation of propionate into acetate

was considered unlikely since such reaction is energetically unfeasible in standard conditions ($\Delta G^{o}_{r} =$

53.3 kJ mol^{-1}), and that, as previously described [39], at high H_2 concentrations ΔG_r values are

increased.

Table 5 presents the parameters of the modified Boltzmann model fitting the experimental data from

Figure 4. Those parameters are related to the kinetics of alcohol production and compare efficiencies

between all pretreatments. A high correspondence was obtained between replicates for all conditions

tested, apart from the thermal-acidic pretreatment. The correlation coefficient was very low in that

condition, only reaching 0.5 for ethanol and 0.4 for butanol, indicating that the process was

unpredictable and could not be reproduced. Due to its instability and unpredictability, the thermal-

acidic pretreatment was thus no longer considered for analyses.

Table 5 – Parameters of the modified Boltzmann model fitting for ethanol and butanol production.

	Parameters	Pretreatment of the inoculum				
		Control	Acidic	Thermal	Acidic-thermal	Thermal-acidic
Ethanol	C_{max} (mg L^{-1})	38.5 ± 1.5	13.6 ± 0.7	122 ± 10	87.3 ± 1.9	27.6 ± 2.8
	t_m (d)	12.7 ± 0.6	10.4 ± 0.8	21.0 ± 1.1	6.9 ± 0.3	6.1 ± 1.0
	r_{max} (mg L^{-1} d^{-1})	1.9 ± 0.0	0.7 ± 0.0	7.0 ± 0.48	9.4 ± 0.7	3.1 ± 1.2
	R^2	0.96	0.89	0.94	0.95	0.50
	t_i (d)	2.7	0.9	12	2.3	1.6
	t_e (d)	20	19	18	9.3	9.0
Butanol	C_{max} (mg L^{-1})	50.7 ± 2.3	10.9 ± 0.5	96.7 ± 6.8	143 ± 2	53.8 ± 5.4
	t_m (d)	17.6 ± 0.7	14.2 ± 0.8	21.1 ± 1.0	5.3 ± 0.2	5.1 ± 0.8
	r_{max} (mg L^{-1} d^{-1})	2.4 ± 0.0	0.5 ± 0.0	5.5 ± 0.5	25 ± 2	26 ± 23
	R^2	0.97	0.94	0.96	0.96	0.40
	t_i (d)	7.0	2.9	12	2.4	4.1
	t_e (d)	21	23	18	6	2

C_{max}, maximum concentration; t_m, time when maximum production rate is achieved; r_{max}, maximum production rate; R^2, correlation coefficient; t_i and t_e, initial and ending time of exponential growth phase.

As shown in Figure 4 and Table 5, both alcohol production and maximum production rates (r_{max}) were

improved for thermal and acidic-thermal pretreatments. Although the highest ethanol production was

observed for the thermal pretreatment, the best results were obtained for the acidic-thermal

pretreatment, which allowed the best butanol production and an increase of 4.5 times of the ethanol and

of 10.2 times of the butanol maximum production rate. In opposition, a decrease in alcohol production

was observed for the acidic pretreatment. Such results clearly indicate that an acidic-thermal

pretreatment of the inoculum is a promising approach for designing more efficient and smaller-sized

bioreactors.

Length of the lag phase (t_i) and duration of the bacterial exponential growth phase (t_e) were evaluated

for each pretreatment and compared to the control by considering alcohol production curves as growth-

associated curves (Table 3). The length of the lag phase is related to the time required for a bioreactor

to initiate its process (start-up) and achieve higher rates of alcohol production. For both ethanol and

butanol production, the shortest lag phases were observed for the acidic and acidic-thermal

pretreatments. In both pretreatments, the lag phase was shorter than found in the control essay. The

most extended lag phase was observed for the thermal pretreatment, which was considerably greater for

both ethanol and butanol production than observed in control. Such results indicate that the thermal

pretreatment had the highest impact on the inoculum, while acidic and acidic-thermal pretreatments

selected microbial communities which were the best adapted to solventogenic processes. Variations in

the duration of the bacterial exponential growth phases (t_e) were also observed (Table 5). A shorter

exponential growth phase is representative of a faster process. However, this parameter must be

evaluated concomitantly with the maximum rates of production (r_{max}). For example, a short t_e

occurring at low r_{max} indicates a process occurring with low efficiency. Among all the conditions

tested, low t_e values and high r_{max} values were obtained for the acidic-thermal pretreatment for

ethanol and butanol production. Taken all together, results obtained for all conditions tested indicate

that the acidic-thermal pretreatment was the best approach to generate a rapid and efficient process.

363
Figure 6 – Evolution of pH along the process for each studied condition.

pH was initially set to a value of 5.92 ± 0.09 for all essays, and their value increased during the

experiments (Figure 6). This rise in pH values likely reflects the consumption of H2 as an electron

donor to form alcohols since solventogenic pathways (3, 5 and 6) require the consumption of H+ and

alcohols show a low ionization on water. Such a pH increase occurred to a lesser extent in all

conditions, including an inoculum acidic pretreatment (acidic, acidic-thermal and thermal-acidic

pretreatments). This difference might be explained by a more drastic initial pH drop resulting from the

acidic addition on these pretreatments, probably lowering the buffering capacity of the inoculum. This

hypothesis is reinforced by the concentrations of dissolved CO_2, which were higher for both control

and thermal pretreatment than acidic and acidic-thermal pretreatments (Table 3).

375
Figure 7 – Evolution of the total volatile solids (TVS) along the process for each studied condition.

A decrease in total volatile solids (TVS) was also observed for all conditions tested (Figure 7). This

decrease occurred probably due to a direct effect of the different pretreatments on the biomass,

resulting in partial microbial cell death. Such non-living cells constituted non-soluble organic matter

which was probably hydrolyzed and then consumed as a carbon source, contributing to the organic

matter input proposed in pathway 1 of Figure 1. The TVS concentration stopped decreasing and started

to increase after 6.3 days in the acidic-thermal pretreatment slightly. Such evolution is probably linked

to the higher values of   observed for this pretreatment (Table 5), which reflect an increase in the

biomass growth's rate, and consequently, the TVS concentration.

5.3 Bacterial communities

Based on the results presented above and focusing on alcohol production, both thermal and acidic-

thermal pretreatments experiments were submitted to microbial community analyses. Bacterial

populations were thus characterized and monitored all along with those two processes. Thirteen phyla

were detected, namely *Firmicutes*, *Proteobacteria*, *Bacteroidetes*, *Actinobacteria*, *Synergistetes*,

Cyanobacteria, *Tenericutes*, *Spirochaetes*, *Deinococcus-Thermus*, *Fibrobacteres*, *Verrucomicrobia*,

Chloroflexi and *Nitrospirae*. Among those, three (*Firmicutes*, *Proteobacteria* and *Bacteroidetes*)

represented up to 94.6% of the total OTUs detected in each sample for both pretreatments (Figure 8).

Firmicutes was the most abundant phylum detected in the initial non-treated inoculum (corresponding

to 66.6% of the detected OTUs). Both pretreatments generated a shift in the bacterial population by

stimulating the development of *Proteobacteria* and strongly reducing the number of *Firmicutes*.

Bacterial populations evolved differently along the process for the two pretreatments. In the thermal

pretreatment experiment (Figure 8A), a decrease of *Proteobacteria* was observed concomitantly with

an increase in *Bacteroidetes* and *Firmicutes*, the latest becoming the dominant phylum at the end of the

process. This shift started after 200 hours of experiment, which correspond to the early beginning of the

bacterial exponential growth phase (Figure 4C and Table 5). In opposition, in the acidic-thermal

pretreatment experiment (Figure 8B), relative stability was observed for the bacterial population, with

Proteobacteria remaining the main phylum all along the process ($72.0 \pm 4.99\%$ of the detected OTUs)

and *Firmicutes* constantly being the second phylum of importance ($24.0 \pm 5.10\%$ of the detected

OTUs). *Bacteroidetes* increased during the process, before starting to decrease after 500h of

experiment, to reach their initial level at the end of the experiment.

407
Figure 8 – Evolution of the three most representative bacterial phyla for the thermal (A) and acidic-
thermal (B) pretreatments. Dashed lines represent OTUs associated to those same phyla found in the
inoculum without pretreatment.

Deeper phylogenetic analyses were performed down to the genus level to infer the potential metabolic

pathway(s) that could be associated with the enhancement of alcohol production observed for both

thermal and acidic-thermal pretreatments (Figure 9). Substantial differences were noticed between the

bacterial population from the initial inoculum (before any pretreatment) and those from the thermal and

acidic-thermal pretreated inocula.

417
Figure 9 – Evolution of the bacterial population at the genus taxonomic level for thermal and acidic-
thermal pretreatments, and comparison with the initial bacterial population (I) for the 31 most detected
genera.

A total of 317 genera were detected in all samples tested, but only 31 of them, representing up to 85%

of the bacterial population, were considered for further analysis (Figure 9). One of the most noticeable

changes consisted of an increase of *Pseudomonas* in pretreatment experiments compared to the initial

population. Such increase was observed as soon as the first hours of experiments. *Pseudomonas* then

remained a dominant genus of the bacterial population, despite a slight decrease observed for the

thermal pretreatment in the second half of the process. Several *Pseudomonas* species have been

genetically well-characterized and have been shown to possess the genetic components for both ethanol

(from pyruvate) and butanol (from glycerol and pyruvate) production [42]. The prevalence of

Pseudomonas is likely linked with the higher levels of alcohol production observed in thermal and

acidic-thermal pretreatments. Some *Pseudomonas* species also possess genes responsible for the

degradation of ethanol into acetyl-CoA [42]. A pathway related to ethanol degradation might explain

better butanol production compared to ethanol, observed for the acidic-thermal pretreatment. Among

the other noticeable results, two bacterial genera, namely *Acinetobacter* and *Paenibacilus*, which were

not detected in the initial population, appeared to be positively affected by both pretreatments. It is

being reported [43,44] that *Acinetobacter* strains are related to alcohol consumption as they can express

alcohol dehydrogenase (ADH) to convert ethanol into acetate, reverting the solventogenic pathway 6.

As shown in Figure 2, this reversed pathway probably not occurred during the essays since in all

experiments, in their beginning and ending, ΔG_r for pathway 6 were thermodynamically feasible.

Conversely, ADH is also a molecule that is strongly related to bacterial quorum sensing with a key role

in biofilm formation [44,45]. The growth of *Acinetobacter* was more stimulated in the thermal

pretreatment, so this quorum sense mechanism could be linked with the best performance of this

pretreatment to produce alcohols, considering that the presence of ethanol could have stimulated the

expression of ADH, although the alcohol degradation metabolism was probably shut down. Acidic-

thermal pretreatment preferentially stimulated the growth of *Paenibacilus*, which is directly related to

alcohol production [46].

In addition to those changes, several other bacterial genera evolved differently depending on the

pretreatment applied (Figure 9). The thermal pretreatment appeared to stimulate the growth of

Brevundimonas, *Bacteroides*, *Butyrivibrio*, *Bacillus*, *Alcaligenes*, *Eubacterium*, *Clavibacter*,

Psychrobacter, *Serratia* and *Microbulbifer*. The latter three being even exclusively detected in this

pretreatment. On the opposite, the acidic-thermal pretreatment appeared to improve the growth of

Tissierella, *Novosphingobium*, *Sedimentibacter* and *Yersinia*. The latter one being exclusively detected

in this pretreatment.

The genus *Brevudimonas* started to increase in the thermal pretreatment after 8.3 days, coinciding with

the very beginning of the exponential growth phase (Figure 4 and Table 5). Members belonging to this

genus have already been shown to possess metabolic pathways related to the acidogenesis of alcohols

[47]. High H_2 partial pressures applied during the process might have rendered solventogenesis

thermodynamically favourable through the reverse same metabolic pathway (Figure 3). The genus

Yersinia started to increase in the acidic-thermal pretreatment after approximately 4.2 days, coinciding

with the early exponential growth phase (Figure 4 and Table 5). Members belonging to this genus have

already been shown to possess metabolic pathways to convert pyruvate into butanol and ethanol [48].

The genus *Sedimentibacter* is one of the predominant genus found in the fermentation of Baijiu, and it

could be directly related to alcohol production [49]. It is noticeable that the genus *Clostridium*, which is

composed of a high number of known alcohol producers [15,50–52], was detected for both

pretreatments all along the process, without any significant variations. The genus *Clostridium* usually

produces alcohol by converting glycerol and pyruvate into ethanol and butanol. Thus, this genus

probably has an essential role in producing ethanol and butanol in both thermal and acidic-thermal

pretreatments. There is no reporting on alcohol-producing metabolism for all other genera found within

the experiments.

Other results were observed at higher taxonomic levels, notably decreasing microorganisms from the

Peptostreptococcaceae family and the *Epsilonproteobacteria* class, in both pretreatments.

Peptostreptococcaceae family represented almost half (45% of total bacterial OTUs) found in the

initial inoculum (before any pretreatment), strongly decreased its presence to $5.4 \pm 2.2\%$ in thermal and

$9.2 \pm 3.1\%$ in acidic-thermal pretreatments. *Epsilonproteobacteria* class represented up to 6% of total

bacterial OTUs in the initial inoculum but significantly decreased to less than 0.3% of total bacterial

OTUs in both thermal and acidic-thermal pretreatments. Such results indicate that neither bacteria from

the *Peptostreptococcaceae* family nor *Epsilonproteobacteria* class played a significant role in the

processes studied.

6   Conclusions

Characterization of the initial microbial population showed the presence of bacteria possessing

metabolic pathways for alcohol production. Both thermal and acidic-thermal pretreatments were able to

select the best adapted bacterial communities to produce ethanol and butanol. The significant decrease

of members belonging to the family *Peptostreptococaceae* coupled to an increase of *Pseudomonas*

appeared to be related to the best performance of ethanol and butanol production. The acidic-thermal

pretreatment generated the best results, reaching a concentration of 87 mg L^{-1} of ethanol and of 143 mg

L^{-1} of butanol after 240 hours of processing. The thermal pretreatment achieved the highest ethanol

production (122 mg L^{-1}) but at a much slower rate (after 710 hours). The two other pretreatments

studied showed either instability and inconsistency for the thermal-acidic pretreatment or very low

alcohol production for the acidic pretreatment. Among the VFAs added to the medium, butyrate was

used solely as the substrate, acetate being produced along the process. The mass balance study

highlighted another substrate input, which was probably originating from the inoculum.

Thermodynamic data indicated that homoacetogenesis was a critical pathway to produce acetate from

dissolved CO_2 and H_2. Finally, H_2 partial pressure was a preponderant factor for solventogenesis,

enabling alcohol production and inhibiting its consumption.

7   List of Abbreviations Used

| 495 | **7** | **List of Abbreviations Used** |

STP   Standard temperature and pressure (0° C and 1 atm);

ppH$_2$ Partial H$_2$ pressure (atm);

R^2 Levenberg-Marquardt algorithm correlation coefficient;

TVS   Total volatile solids (mg L^{-1});

OTU   Operational taxonomic unit;

HPLC   High performance liquid chromatography;

PCR   Polymerase chain reaction;

RDP   Ribosomal data project;

VFA   Volatile fatty acid;

But$^-$ Butyrate;

Prop$^-$ Propionate;

Ac$^-$ Acetate;

But(OH)   Butanol;

Prop(OH)   Propanol;

Eth(OH)   Ethanol;

$C(t)$ Concentration in respect of time (mg L^{-1});

t Time (h);

t_i Initial time for exponential growth phase (h);

t_e Ending time for exponential growth phase (h);

t_m Time in which maximum production rate (r_{max}) is achieved (h);

r_{max} Maximum production rate (mg L^{-1} h^{-1});

C_{max} Maximum concentration (mg L^{-1});

ΔG_r Variation of Gibbs' free energy at a given condition (kJ mol^{-1});

ΔG_r° Variation of Gibbs' free energy at standard condition (kJ mol^{-1});

ΔG_r^{F} Variation of Gibbs' free energy at final experiment condition (kJ mol^{-1});

ΔG_r^{I} Variation of Gibbs' free energy at initial experiment condition (kJ mol^{-1}).

8   Competing Interests

The authors declare that they have no competing interests.

9   Authors' Contribution

All authors have planned the experiments. GM performed all the experiments, data treatment, statistics, and mathematical modeling. GB treated and analysed all sequencing data. All authors have analysed and discussed the results. GM have written the paper. All authors have read, reviewed, and approved the final manuscript. Experiments were conducted in Montréal branch of National Research Council of Canada from January to July of 2014.

10   Authors' Informations

GM is professor of sanitation and applied microbiology of the Agricultural Engineering College at Campinas State University (FEAGRI/UNICAMP). This research was developed as an invited researcher at National Research Council of Canada (NRC) from October/2013 to October/2014, under a post-doctoral program in biofuels in the São Carlos Engineering School at University of São Paulo (EESC/USP).

11   Acknowledgements

This work was supported by FAPESP – Fundação de Amparo a Pesquisa do Estado de São Paulo (processes 2010/18.463-9 and 2013/18.172-2 – G. Mockaitis and 2009/15.984-0 – M. Zaiat), and the NRC – National Research Council of Canada (project A1-004645 – S.R. Guiot).

The authors thankfully acknowledge Marie-Josée Lévesque, Christine Maynard and Sylvie Sanschagrin for their assistance with the biomolecular techniques (DNA extraction, purification and PCR) and sequencing (through Ion Torrent[TM]); and also, Stephane Deschamps and Alain Corriveau for their valuable contribution with physicochemical analysis (HPLC and gas chromatography of alcohols).

12 References

[1] Jones DT, Woods DR. Acetone-butanol fermentation revisited. Microbiol Rev 1986;50:484–524.

[2] Das D, Veziroglu T. Advances in biological hydrogen production processes. Int J Hydrogen Energy 2008;33:6046–57. https://doi.org/10.1016/j.ijhydene.2008.07.098.

[3] Baumann I, Westermann P. Microbial Production of Short Chain Fatty Acids from Lignocellulosic Biomass: Current Processes and Market. Biomed Res Int 2016;2016. https://doi.org/10.1155/2016/8469357.

[4] Levin D. Biohydrogen production: prospects and limitations to practical application. Int J Hydrogen Energy 2004;29:173–85. https://doi.org/10.1016/S0360-3199(03)00094-6.

[5] Steinbusch KJJ, Hamelers HVM, Buisman CJN. Alcohol production through volatile fatty acids reduction with hydrogen as electron donor by mixed cultures. Water Res 2008;42:4059–66. https://doi.org/10.1016/j.watres.2008.05.032.

[6] Zverlov V V, Berezina O, Velikodvorskaya G a, Schwarz WH. Bacterial acetone and butanol production by industrial fermentation in the Soviet Union: use of hydrolyzed agricultural waste for biorefinery. Appl Microbiol Biotechnol 2006;71:587–97. https://doi.org/10.1007/s00253-006-0445-z.

[7] Agler MT, Wrenn B a, Zinder SH, Angenent LT. Waste to bioproduct conversion with undefined mixed cultures: the carboxylate platform. Trends Biotechnol 2011;29:70–8. https://doi.org/10.1016/j.tibtech.2010.11.006.

[8] Dürre P. New insights and novel developments in clostridial acetone/butanol/isopropanol fermentation. Appl Microbiol Biotechnol 1998:639–48.

[9] Brynjarsdottir H, Wawiernia B, Orlygsson J. Ethanol Production from Sugars and Complex Biomass by Thermoanaerobacter AK 5☐: The Effect of Electron-Scavenging Systems on End-Product Formation. Energy and Fuels 2012;26:4568–74.

[10] Jessen JE, Orlygsson J. Production of ethanol from sugars and lignocellulosic biomass by Thermoanaerobacter J1 isolated from a hot spring in Iceland. J Biomed Biotechnol 2012;2012:186982. https://doi.org/10.1155/2012/186982.

[11] Almarsdottir AR, Sigurbjornsdottir MA, Orlygsson J. Effect of various factors on ethanol yields from lignocellulosic biomass by Thermoanaerobacterium AK☐☐. Biotechnol Bioeng 2012;109:686–94. https://doi.org/10.1002/bit.24346.

[12] Crespo CF, Badshah M, Alvarez MT, Mattiasson B. Ethanol production by continuous fermentation of D-(+)-cellobiose, D-(+)-xylose and sugarcane bagasse hydrolysate using the thermoanaerobe Caloramator boliviensis. Bioresour Technol 2012;103:186–91. https://doi.org/10.1016/j.biortech.2011.10.020.

[13] Xu L, Tschirner U. Improved ethanol production from various carbohydrates through anaerobic thermophilic co-culture. Bioresour Technol 2011;102:10065–71. https://doi.org/10.1016/j.biortech.2011.08.067.

[14] Mes-Hartree M, Saddler J. Butanol production of Clostridium acetobutylicum grown on sugars found in hemicellulose hydrolysates. Biotechnol Lett 1982;4:247–52.

[15] Maddox I. Production of ethanol and n-butanol from hexose/pentose mixtures using consecutive fermentations with Saccharomyces cerevisiae and Clostridium acetobutylicum. Biotechnol Lett

1982;4:23–8.

[16] Ounine K, Petitdemange H, Raval G, Gay R. Acetone-butanol production from pentoses by
Clostridium acetobutylicum. Biotechnol Lett 1983;5:605–10.

[17] Li Z, Shi Z, Li X. Models construction for acetone-butanol-ethanol fermentations with
acetate/butyrate consecutively feeding by graph theory. Bioresour Technol 2014;159:320–6.
https://doi.org/10.1016/j.biortech.2014.02.095.

[18] Kumar M, Goyal Y, Sarkar A, Gayen K. Comparative economic assessment of ABE
fermentation based on cellulosic and non-cellulosic feedstocks. Appl Energy 2012;93:193–204.
https://doi.org/10.1016/j.apenergy.2011.12.079.

[19] Kleerebezem R, van Loosdrecht MCM. Mixed culture biotechnology for bioenergy production.
Curr Opin Biotechnol 2007;18:207–12. https://doi.org/10.1016/j.copbio.2007.05.001.

[20] Puig S, Coma M, Monclús H, van Loosdrecht MCM, Colprim J, Balaguer MD. Selection
between alcohols and volatile fatty acids as external carbon sources for EBPR. Water Res
2008;42:557–66. https://doi.org/10.1016/j.watres.2007.07.050.

[21] O-Thong S, Prasertsan P, Birkeland N-K. Evaluation of methods for preparing hydrogen-
producing seed inocula under thermophilic condition by process performance and microbial
community   analysis.   Bioresour   Technol   2009;100:909–18.
https://doi.org/10.1016/j.biortech.2008.07.036.

[22] Luo G, Karakashev D, Xie L, Zhou Q, Angelidaki I. Long-term effect of inoculum pretreatment
on fermentative hydrogen production by repeated batch cultivations: homoacetogenesis and
methanogenesis as competitors to hydrogen production. Biotechnol Bioeng 2011;108:1816–27.
https://doi.org/10.1002/bit.23122.

[23] Pendyala B, Chaganti SR, Lalman J a., Shanmugam SR, Heath DD, Lau PCK. Pretreating mixed
anaerobic communities from different sources: Correlating the hydrogen yield with hydrogenase
activity   and   microbial   diversity.   Int   J   Hydrogen   Energy   2012;37:12175–86.
https://doi.org/10.1016/j.ijhydene.2012.05.105.

[24] Kumar M, Gayen K, Saini S. Role of extracellular cues to trigger the metabolic phase shifting
from acidogenesis to solventogenesis in Clostridium acetobutylicum. Bioresour Technol
2013;138:55–62. https://doi.org/10.1016/j.biortech.2013.03.159.

[25] Angelidaki I, Petersen SP, Ahring BK. Applied Microbiolog . v Effects of lipids on thermophilic
anaerobic digestion and reduction of lipid inhibition upon addition of bentonite 1990:469–72.

[26] Sander R. Compilation of Henry ' s Law Constants for Inorganic and Organic Species of
Potential Importance in Environmental Chemistry 1999.

[27] Lide DR, Frederikse HPR, editors. CRC Handbook of Chemistry and Physics. 76th ed. Boca
Ratón, FL, USA: CRC Press Inc.; 1995.

[28] APHA. Standard methods for the examination of water and wastewater. 21st ed. Washington
DC: 2005.

[29] Dilallo R, Albertson O. Volatile Acids by Direct Titration. J Water Pollut Control Fed
1961;33:356–65.

[30] Ripley L, Boyle W, Converse J. Improved alkalimetric monitoring for anaerobic digestion of
high-strength wastes. J Water Pollut Control Fed 1986;58:406–11.

[31] Guiot SR, Cimpoia R, Carayon G. Potential of wastewater-treating anaerobic granules for
biomethanation   of   synthesis   gas.   Environ   Sci   Technol   2011;45:2006–12.

https://doi.org/10.1021/es102728m.

[32]   Muyzer G, Dewaal EC, Uitterlinden AG. Profiling of complex microbial populations by
desnaturing gradient gel electrophoresis analysis ofpolymerase chain reaction amplified genes
coding for 16S ribosomal RNA. Appl Environ Microbiol 1993;59:695–700.

[33]   Griffiths RI, Whiteley a S, O'Donnell a G, Bailey MJ. Rapid method for coextraction of DNA
and RNA from natural environments for analysis of ribosomal DNA- and rRNA-based microbial
community composition. Appl Environ Microbiol 2000;66:5488–91.

[34]   Lévesque MJ, La Boissière S, Thomas JC, Beaudet R, Villemur R. Rapid method for detecting
Desulfitobacterium frappieri strain PCP-1 in soil by the polymerase chain reaction. Appl
Microbiol Biotechnol 1997;47:719–25.

[35]   Berthelet M, Whyte LG, Greer CW. Rapid, direct extraction of DNA from soils for PCR
analysis using polyvinylpolypyrrolidone spin columns. FEMS Microbiol Lett 1996;138:17–22.

[36]   Wang Q, Garrity GM, Tiedje JM, Cole JR. Naive Bayesian classifier for rapid assignment of
rRNA sequences into the new bacterial taxonomy. Appl Environ Microbiol 2007;73:5261–7.
https://doi.org/10.1128/AEM.00062-07.

[37]   Claesson MJ, O'Sullivan O, Wang Q, Nikkilä J, Marchesi JR, Smidt H, et al. Comparative
analysis of pyrosequencing and a phylogenetic microarray for exploring microbial community
structures   in   the   human   distal   intestine.   PLoS   One   2009;4:e6669.
https://doi.org/10.1371/journal.pone.0006669.

[38]   Mavrovouniotis ML. Group contributions for estimating standard gibbs energies of formation of
biochemical   compounds   in   aqueous   solution.   Biotechnol   Bioeng   1990;36:1070–82.
https://doi.org/10.1002/bit.260361013.

[39]   Mavrovouniotis ML. Errata Group Contributions for Estimating Standard Gibbs Energies of
Formation of Biochemical Compounds in Aqueous Solution 1991;38:803–4.

[40]   Harper SR, Pohland FG. Recent Developments in Hydrogen Management During Anaerobic
Biological Wastewater Treatment. Biotechnol Bioeng 1985;28:585–602.

[41]   Speece RE. Anaerobic biotechnology for industrial wastewater. Vanderbilt University; 1996.

[42]   Caspi R, Billington R, Fulcher CA, Keseler IM, Kothari A, Krummenacker M, et al. The
MetaCyc database of metabolic pathways and enzymes. Nucleic Acids Res 2018;46:D633–9.
https://doi.org/10.1093/nar/gkx935.

[43]   John J, Saranathan R, Adigopula LN, Thamodharan V, Singh SP, Pragna Lakshmi T, et al. The
quorum sensing molecule N-acyl homoserine lactone produced by Acinetobacter baumannii
displays   antibacterial   and   anticancer   properties.   Biofouling   2016;32:1029–47.
https://doi.org/10.1080/08927014.2016.1221946.

[44]   Zhang K, Yang X, Yang J, Qiao X, Li F, Liu X, et al. Alcohol dehydrogenase modulates quorum
sensing in biofilm formations of Acinetobacter baumannii. Microb Pathog 2020;148:104451.
https://doi.org/10.1016/j.micpath.2020.104451.

[45]   Subhadra B, Hwan Oh M, Hee Choi C. Quorum sensing in Acinetobacter: with
special emphasis on antibiotic resistance, biofilm formation and quorum quenching. AIMS
Microbiol 2016;2:27–41. https://doi.org/10.3934/microbiol.2016.1.27.

[46]   Grady EN, MacDonald J, Liu L, Richman A, Yuan ZC. Current knowledge and perspectives of
Paenibacillus: A review. Microb Cell Fact 2016;15:203. https://doi.org/10.1186/s12934-016-
0603-7.

[47] Leifson E, Hugh R. A new type of polar monotrichous flagellation. J Gen Microbiol 1954;10:68–70. https://doi.org/10.1099/00221287-10-1-68.

[48] Caspi R, Altman T, Dale JM, Dreher K, Fulcher CA, Gilham F, et al. The MetaCyc database of metabolic pathways and enzymes and the BioCyc collection of pathway/genome databases. Nucleic Acids Res 2010;38:D473-9. https://doi.org/10.1093/nar/gkp875.

[49] Xu J, Sun L, Xing X, Sun Z, Gu H, Lu X, et al. Culturing Bacteria From Fermentation Pit Muds of Baijiu With Culturomics and Amplicon-Based Metagenomic Approaches. Front Microbiol 2020;11:1223. https://doi.org/10.3389/fmicb.2020.01223.

[50] Tracy BP, Jones SW, Fast AG, Indurthi DC, Papoutsakis ET. Clostridia: the importance of their exceptional substrate and metabolite diversity for biofuel and biorefinery applications. Curr Opin Biotechnol 2012;23:364–81. https://doi.org/10.1016/j.copbio.2011.10.008.

[51] Ueki A, Hirono T, Sato E, Mitani A, Ueki K. Ethanol and amylase production by a newly isolated Clostridium sp. World J Microbiol … 1991;7:385–93.

[52] Bruant G, Lévesque M-J, Peter C, Guiot SR, Masson L. Genomic analysis of carbon monoxide utilization and butanol production by Clostridium carboxidivorans strain P7. PLoS One 2010;5:e13033. https://doi.org/10.1371/journal.pone.0013033.

13   One-Sentence Summary

The inoculum pretreatment acid and thermal improved significantly the production of both ethanol and butanol, through an anaerobic solventogenic process, using acetate and butyrate as carbon source and H_2 as electron donor.