Sexual selection affects the evolution of physiological and life history traits

Martin D. Garlovsky[1+], Luke Holman[2+], Andrew L. Brooks[1] and Rhonda R. Snook[3]

[1]Department of Animal and Plant Sciences, University of Sheffield, UK

[2]School of Applied Sciences, Edinburgh Napier University, UK

[3]Department of Zoology, Stockholm University, Sweden

[+]These authors contributed equally to this work

Corresponding author: martingarlovsky@gmail.com

ABSTRACT

Sexual selection and sexual conflict are expected to affect all aspects of the phenotype, not only traits that are directly involved in reproduction. Here, we show coordinated evolution of multiple physiological and life history traits in response to long-term experimental manipulation of the mating system in populations of *Drosophila pseudoobscura*. Development time was extended under polyandry relative to monogamy in both sexes, potentially due to higher investment in traits linked to sexual selection and sexual conflict. Individuals (especially males) evolving under polyandry had higher metabolic rates and locomotor activity than those evolving under monogamy. Polyandry individuals also invested more in metabolites associated with increased endurance capacity and efficient energy metabolism and regulation, namely lipid and glycogen. Finally, polyandry males were less desiccation- and starvation-resistant than monogamy males, suggesting trade-offs between resistance and sexually selected traits. Our results provide experimental evidence that mating systems can impose selection that influences the evolution of non-sexual phenotypes such as development, activity, metabolism, and nutrient homeostasis.

KEYWORDS: Sexual selection, polyandry, metabolism, life history evolution, trade-offs, experimental evolution.

INTRODUCTION

Natural and sexual selection often act differently in males and females, which have different routes to evolutionary fitness as a result of anisogamy [1,2], though independent evolution of the sexes is partly constrained due to their shared genome [3]. Recent work has increasingly highlighted that sexual selection affects all aspects of the phenotype, and not only classic sexually-selected traits such as male ornaments [4]. This is because sexually-selected traits often depend on overall condition [5], which in turn depends on fundamental aspects of organisms such as their development, physiology, life history and metabolism [6,7].

Sexual selection often favours energetically costly bouts of sustained locomotor activity, e.g. during mate searching, courtship, or the 'harassment and resistance' behaviours that typify interlocus sexual conflict [8–12]. Moreover, postcopulatory sexual selection can select for males that produce metabolically expensive ejaculates [13,14]. We therefore expect sexual selection and sexual conflict to favour physiological adaptations that augment the efficiency of metabolism and respiration [11,15]. Sexual selection is also hypothesised to affect the evolution of development and life history, potentially in a sex-specific way [16], for instance by selecting for a 'live fast, die young' strategy in species where competition for mates is intense [17,18], or selecting for the reallocation of limiting resources away from growth and somatic maintenance [10,19–21]. Indeed, the conspicuous trade-offs involved in the production of sexually-selected traits served as the original motivation for the theory of sexual selection [1].

In addition to these links between sexually selected traits and other phenotypes, the picture is further complicated by genetic correlations between the sexes. Male and female traits have a shared genetic basis, such that selection on males results in a (frequently maladaptive) evolutionary response in females, and *vice versa* e.g. [18,22–29]. Therefore, sexual selection on males not only results in evolution of the males' life history, physiology etc., but also those of females; similarly, selection on female reproductive traits can influence the life history of males.

Here, we investigate how the long-term experimental manipulation of sexual selection and sexual conflict alters the evolution of physiological and life history traits in a fruit fly,

Drosophila pseudoobscura. One set of replicate populations (n = 4, hereafter 'lines')
experienced enforced monogamy (M), eliminating sexual selection and sexual conflict, and
another set of lines (n = 4) experienced elevated polyandry (E), where 6 males were housed
with 1 female to facilitate inter- and intra- sexual selection, likely at elevated levels compared
to natural populations [30]. We have previously found divergence between the E and M lines
in a number of traits under sexual selection. For example, E males produce more abundant
and complex chemical signals [31], perform a faster and more vigorous courtship song
[12,32,33], court and mate more frequently [34], are more competitive in mating encounters
[35], and have larger glands for producing seminal fluid [36]. Additionally, coevolutionary
patterns have been found, such that E males are more harmful but E females more resilient
to such harm [36,37], and females prefer courtship song of males from their own treatment
[33]. These evolved phenotypic differences are accompanied by divergence in gene
expression [38,39] and concerted genetic differences between treatments [40]. In the present
study, we tested whether the mating system similarly caused life history and metabolic traits
to evolve, which we tested by measuring respiration, macro-metabolites, and developmental
traits.

METHODS

Detailed methods, including establishment and maintenance of the experimental evolution lines, can be found in the online supplementary material. In brief, in the enforced monogamy or 'M' treatment, the adult population was housed in groups of two (one male, one female) with a family size of 60 vials. In the elevated polyandry or 'E' treatment, each group of adults comprised six males and one female, with a family size of 40 vials. In each generation, unmated males and females were housed in 'interaction vials' for 5 days, before being transferred to 'oviposition vials' (OVs) for a further 5 days. Flies were kept at 22°C on a 12:12 light:dark cycle on standard cornmeal-agar-molasses media with added live yeast. Importantly, effective population size was standardized across the treatments so divergent treatment responses represent response to selection rather than differential genetic drift [41].

Experimental individuals

Prior to the experiments described below, flies from each line were placed in a 'common garden' to minimise non-genetic differences between lines. Newly-eclosed individuals were collected *en masse* from the OVs within each replicate population. A random sample of these flies were allowed to mate and oviposit for two days. From these eggs, we set up controlled density vials (CDVs) by placing 100 first instar larvae into food vials. Unmated flies eclosing from CDVs were collected and stored in same-sex food vials until 3-5 days old. Excluding the development time experiment, each measurement in our study was performed on a 'triad' of three age-matched flies (due to practical constraints resulting from the small size of individual flies).

Juvenile development time

We measured juvenile development time at generations 180, 179, 178 and 176 for replicates 1-4, respectively. For each replicate of the E and M treatments, we seeded 6 CDVs (see above) on three consecutive days (i.e., 600 larvae per replicate population per seeding day = 14,400 larvae). On the day of eclosion, emerging flies were CO_2 anaesthetised and killed in ethanol. We continued collecting until no individuals eclosed for two consecutive days. We subsequently counted the number of adult males and females emerging each day from each vial.

We used the length of wing vein IV as a proxy for body size [37] by measuring a random

subsample of individuals (n = 15 per sex per replicate per seeding day). We removed the left

wing from flies preserved in ethanol and mounted wings on a microscope slide in a drop of

phosphate-buffered saline and dried at room temperature overnight. We imaged wings using

a Motic camera and Motic Images Plus 2.0 software (Motic Asia, Hong Kong). We measured

wing vein IV length using ImageJ software [42]. Image files were anonymised prior to

measurement.

Desiccation and starvation resistance

We measured desiccation and starvation resistance at generations 199, 198, 197 and 195 for

replicates 1-4, respectively. Triads (n = 7-10) were housed in 8-dram plastic vials stoppered

with cotton balls and covered with Parafilm®. For the desiccation resistance assay, vials

contained no food and between the cotton and Parafilm® we placed a packet of silica gel

beads. For the starvation resistance assay, vials contained an agar solution that provided

moisture but no food. Vials were checked every 2 hours and any deaths recorded until all flies

perished. Flies were scored as dead if they were not able to right themselves or no movement

was observed.

Respirometry

We measured metabolic rates at generations 196, 195, 194 and 192 for replicates 1-4,

respectively. Triads (n = 3) were weighed to the nearest 0.1 mg (Sartorius Genius ME 235P-

OCE) before transfer to a respirometry chamber (a glass cylinder; 17mm × 70 mm). We

measured the amount of CO_2 produced and O_2 consumed as a proxy for metabolic rate using

a Sable Systems (Las Vegas, NV, USA) respirometry system [43]. Each respirometry chamber

was placed in an activity detector (AD-2, Sable Systems) connected to a data acquisition

interface (Quick-DAQ, National Instruments, Coleman Technologies, Newton Square, US),

which uses reflective infrared-light technology to provide a precise and continuous measure

of locomotor activity of the subjects.

Metabolite extractions

We measured metabolite composition at generations 199, 198, 197 and 195 for replicates 1-

4, respectively. Triads (n = 3 per sex) were weighed to the nearest 1µg (METTLER TOLEDO®

UMX2 ultra-microbalance) and flash frozen in liquid nitrogen. Triads were then placed in a

0.35ml glass vial insert (SUPELCO Analytical®) of known weight, dried at 55ºC overnight and

re-weighed to obtain dry weight. We subsequently processed each sample to acquire

measurements of lipids, soluble carbohydrates, soluble protein, glycogen and chitin.

Statistical analysis

All statistical analyses were performed in R version 4.0.3 [44]. We fit models using Bayesian

approaches in *brms* [45], and specified conservative, regularising priors on the fixed and

random effects. Complete code and description of models are provided in the code repository

(https://lukeholman.github.io/exp_evol_respiration/). In brief, we analysed development

time and survival using survival analysis. For the respirometry and metabolite data we used

structural equation models (SEM), motivated by the causal relationships that we hypothesise

to exist between the variables, given the experimental design and our biological

understanding of the system (Fig. S1). For the SEMs, after fitting the model, we calculated the

treatment effect size (Cohen's *d*) for each response variable, both with and without the

moderators (dry weight and/or activity). We also calculated the difference in treatment effect

size between the sexes, to test for a treatment × sex interaction. All models included replicate

line as a random intercept to reflect the experimental design of the selection experiment (i.e.

where n = 8) and selection treatment as a random slope to allow lines to vary in their response

to selection [46]. All models also included vial identity as a random intercept, except for wing

vein length where we pooled individuals across vials prior to measurements.

RESULTS

Juvenile development time

Juvenile development time differed significantly between the E and M treatments (Cox

proportional hazards model; PP = 0.013, where PP is the posterior probability the effect is of

the opposite direction, similar to a 1-tailed p-value) and between the sexes (PP < 0.001). E

treatment flies had a reduced hazard (Hazard ratio = 0.45; 95% confidence intervals [CI] = 0.27

– 0.85), i.e., development took longer in the E treatment in both sexes. Males took longer to

eclose than females (Hazard ratio = 0.84; 95% CI = 0.80 – 0.89) (Fig. 1A), as usual in this

species. There was a significant treatment × sex interaction (PP = 0.001), such that the sex

difference in development time was greater in the E treatment, and the treatment affected

males more strongly than females.

As expected, females were larger than males (mean wing vein IV length (μm); M females:

2324 ± 4.80, n = 152; E females: 2335 ± 5.76, n = 118; M males: 2099 ± 4.93, n = 154; E males:

2114 ± 5.96, n = 127). We found no statistically significant effect of sexual selection treatment

(PP = 0.317) or the treatment × sex interaction (PP = 0.384).

Desiccation and starvation resistance

We found a significant treatment × sex interaction for both desiccation and starvation

resistance (desiccation: PP = 0.024; starvation: PP = 0.018). E males survived 0.91 (95% CI =

0.83 – 0.99), and 0.88 (95% CI = 0.80 – 0.97), times as long as M males under desiccation and

starvation resistance, respectively. However, the treatment effect was opposite in females: E

females survived 1.06 (95% CI = 0.87 – 1.29), and 1.09 (95% CI = 0.88 – 1.35), times longer

than M females under desiccation and starvation resistance, respectively. In short, M males

had better survival than E males, while E females had better survival than M females.

Figure 1: Kaplan-Meier plots for (a) juvenile development time, (b) desiccation resistance, and
(c) starvation resistance. '+' indicates right censored individuals. Monogamy (M) = pink and
light blue, Elevated polyandry (E) = red and dark blue; Males = dashed lines, Females = solid
lines. Shaded areas show confidence intervals.

Respirometry

Beginning with the mediator variables, females were heavier than males (PP < 0.001; Table

S1), and E females were non-significantly heavier than M females (Fig. 2A; Mean dry weight

per triad (mg): M females: 0.40 ± 0.02; E females: 0.47 ± 0.04; M males: 0.29 ± 0.02; E males:

0.29 ± 0.02; n = 12 each). The treatment (PP = 0.078) and treatment × sex (PP = 0.099) terms

were not significantly related to body mass. Activity level showed a sex × cycle interaction (PP

< 0.031), indicating that activity levels declined over cycles in males but not females. This sex

difference in the rate of decline was stronger in M treatment than E treatment, though the 3-

way interaction was not statistically significant (PP = 0.072).

There was a strong correlation between activity level and O_2 consumption (PP < 0.001; Fig.

S2), and O_2 consumption declined across cycles (PP < 0.001). Body weight did not correlate

with O_2 consumption (PP = 0.47; Fig. S2). Calculating the posterior estimate of the difference

in treatment means for each sex and cycle revealed that O_2 consumption was higher in the E

treatment (indicated by positive effect sizes in Fig. 2B), especially in males. This effect was

considerably attenuated when we statistically adjusted for differences in activity level

between treatments and sexes, indicating that the evolved difference in activity between the

E and M treatments was largely responsible for the treatment effect on O_2 consumption.

Controlling for the evolved differences in body mass did not change the effect size, illustrating

that the (modest) changes in body mass between the E and M treatments did not explain the

evolved difference in O₂ consumption.

The respiratory quotient (RQ) did not differ detectably between the E and M treatments,

whether or not one controlled for the mediator variables (Fig. S3). The grand mean RQ = 0.90

(± 0.02) indicated flies used a mixture of carbohydrates, protein, and lipids as metabolic

substrate.

Figure 2: Effects of sex and selection treatment on metabolic rates and mediator variables. A)
Posterior estimates of the means for mediator variables (body mass and activity). Note that
females are larger than males, and that E females are somewhat larger than M females. B)
Posterior estimates of effect sizes of selection treatment (E minus M) on metabolic rate
(measured as volume of O2 consumed) controlling for activity, body mass, or neither
mediator variable. The x-axes show the posterior estimate of standardised effect size
(Cohen's d), i.e., a value of 1 would mean that the E treatment has a mean that is larger by 1
standard deviation. The horizontal bars show the 66% and 95% quantiles and the median of
the posterior distribution. All response variables have been mean-centred and divided by the
overall standard deviation (such that the dashed line at zero marks the mean across sexes,
treatments and cycles). For brevity, only the first cycle is shown for activity and metabolic
rate; see online supplementary material for all cycles.

Metabolite composition

As in the respiration experiment, females were heavier than males and E flies were heavier

than M flies (Dry weight (mg): M females: 0.56 ± 0.02; E females: 0.64 ± 0.17; M males: 0.33

± 0.01; E males: 0.35 ± 0.01; n = 12 each; Table S2). There was a significant sex × treatment

interaction (P = 0.023), such that the E treatment positively affected body mass more strongly

in females. Furthermore, dry weight was significantly correlated with lipid and chitin

concentration (Fig. S4; Table S2), and therefore could act as a mediator for some of the effect

of sex and treatment on these metabolites.

We also found differences in metabolite composition between the E and M treatments and

between sexes (Fig. 3). E treatment flies contained a higher concentration of lipids and

glycogen than M flies, while M flies had relatively more carbohydrates and chitin; the

proportion of protein did not differ between treatments. These treatment differences

reached statistical significance in males for carbohydrate, glycogen, and chitin, and in females

for lipid. When we controlled for the evolved differences in dry weight between the E and M

treatments, only the carbohydrate and glycogen differences between E and M males

remained statistically significant; however, the actual change in effect size when controlling

for dry weight was very small, indicating that dry weight was not an especially important

mediator of the evolved changes in metabolite composition (compare Fig. 3 and Fig. S5).

<u>Figure 3.</u> Posterior estimates of the treatment effect size for both sexes, for each of the five
metabolites. A positive value indicates that the mean metabolite concentration is higher in
the E treatment than the M treatment, while a negative value denotes M > E. A strongly
supported treatment effect is implied by the majority of the posterior lying to one side of
zero. The vertical bars show the 66% and 95% quantiles and the median of the posterior
distribution. This plot was created using posterior predictions of the means that were not
adjusted for differences in dry weight between treatments.

Finally, we investigated the treatment × sex interaction term by calculating the posterior

difference in the treatment effect size between sexes. Though none of the results reached

statistical significance, there was some indication that the decrease in carbohydrate

concentration (in E relative to M) was greater in males than females, that the increase in

glycogen was greater in males (PP = 0.091), and that the sexes showed different changes in

protein concentration (PP = 0.11) (Fig. S6).

DISCUSSION

We found that experimental manipulation of sexual selection and sexual conflict caused

several physiological and life history traits to evolve. First, flies from the elevated polyandry

(E) treatment took longer to develop into adults than their monogamous (M) counterparts

and the difference in development time between the sexes was greater in the E lines. Second,

E treatment flies of both sexes had faster metabolic rates, largely due to E line flies being

more active than M line flies. Third, macro-metabolite composition has diverged between

sexual selection treatments, such that E females had a higher lipid content than M females, E

males higher glycogen content than M males, and M males had higher chitin and sugar

content than E males. Fourth, E males were less resistant to desiccation and starvation than

were M males, whereas E females were more resistant than M females. Finally, the difference

between the sexes in the response to sexual selection treatment was greater evolving under

elevated polyandry for development time and stress resistance.

E males produce a faster and more vigorous courtship song [12,33], more cuticular

hydrocarbons [31], and larger accessory glands [36] than M males. These sexually selected

traits may be energetically costly to produce and maintain, demanding greater metabolic

activity [14,15,21]. Locomotor activity is beneficial for male fitness but costly to females in *D.*

melanogaster and has a significant intersexual genetic correlation [22]. Therefore, sexual

selection favouring increased activity and metabolic rate in males may generate sexual

conflict by increasing activity and metabolic rate above optimal levels for females. We found

no difference in the respiratory quotient, which indicates metabolic substrate use, despite

evolved differences in activity and metabolic rate, suggesting E flies have evolved

compensatory physiological mechanisms to offset the metabolic costs of increased energy

expenditure [47].

We found that E males had higher glycogen concentrations than M males. Carbohydrates are

the main fuel used during intense aerobic activities, such as flight [48] and courtship [49], and

glycogen provides the main source of trehalose marshalled during intense activity [50]. Sexual

selection favouring endurance capacity may also increase lipid respiration and fat storage

[11]. However, despite male courtship song being an endurance contest [12], in males we

found no difference between treatments in lipid concentration. Limited time and resources

may demand strategic resource allocation decisions. We did find higher lipid concentrations

in E females, possibly reflecting selection for endurance capacity, as E females are courted

more frequently [33,37]. No metabolites showed a significant treatment × sex interaction,

indicating the response to sexual selection treatment did not differ between the sexes,

potentially highlighting a strong intersexual genetic correlation constraining divergence. Our

results suggest that heightened sexual selection and sexual conflict favours storage and

regulation of metabolites to meet increased metabolic demands [11,12,15,33,36], whereas

relaxed sexual selection may instead enable investment in other components of fitness.

E males were less tolerant of desiccation and starvation than M males despite their greater

glycogen content, which buffers against these stresses [51]. Previous work artificially selecting

on desiccation tolerance in *D. melanogaster* found selected lines had lower metabolic rates

[52]; our experiment imposed different selective pressures, but found the same negative

correlation between metabolic rate and resistance. This suggests a trade-off in males, where

sexual selection favours investment in current reproduction at a cost to later-life survival,

whereas the relaxation of sexual selection favours increased investment in longevity [9,10].

In contrast, E females showed no such trade-off, as E females were similarly stress resistant

as M females, despite investing more in fecundity [37,38]. The elevated lipid concentrations

we found in E females could mitigate such a trade-off, because lipids provide protection

against starvation alongside being a major component of eggs [53]. Thus, the sexes have

altered investment in macro-metabolite composition under sexual selection, which may allow

differential investment in sexually selected traits and sexual conflict persistence and

resistance traits. The subsequent fitness consequences during stress have different outcomes

on the sexes. This suggests sex specific selection in the adult stage which could further

generate sexual conflict over shared traits, particularly during the juvenile stage [16].

Finally, E flies of both sexes took significantly longer to reach adulthood than flies evolving

under monogamy, perhaps due to pleiotropic costs of increased investment in traits involved

in sexual selection or sexual conflict [21,54,55]. Moreover, because development time is

probably genetically correlated across sexes [23,24], the fact that both sexes evolved longer

development might reflect intralocus sexual conflict; for example, E males might have evolved

to invest more in sexual traits, slowing their development and causing a similar delay in E

females because of pleiotropy. This would be similar to the results of Harano et al. [56], who

selected for male beetles with larger mandibles (a sexually-selected trait), which caused

maladaptive morphological evolution in females despite females lacking the male trait that

was under selection. Conversely, longer development might benefit females, e.g. by enabling

increased resource acquisition and allocation towards reproduction or resistance to male

harm [34,37,38]. The sex difference in development time was also greater in the E lines,

possibly resulting from evolution towards the (shorter) female optimum under monogamy.

To conclude, we have experimentally shown that sexual selection and sexual conflict affects

not only directly selected traits but those affecting whole organismal performance.

Differences in activity levels and metabolic rate coincided with differential investment in

macro-metabolites, reflecting the energetic demands associated with the elevation or

removal of sexual selection. Males evolving under elevated polyandry suffered reduced

survival, potentially trading off current vs. future reproduction. However, the same was not

true of females, perhaps due to selection favouring endurance capacity in females to offset

the costs of frequent male courtship. The physiological and life history traits we measured

likely have a shared genetic basis, and thus may be subject to intralocus sexual conflict. The

magnitude of any sexual dimorphism did not differ between treatments for the physiological

traits we measured (macro-metabolites and metabolic rate), whereas life history traits

(development time and stress resistance) were more dimorphic in populations evolving under

elevated polyandry, suggesting tighter intersexual genetic correlations over physiological

traits. Overall, our findings highlight that sexual selection results in coordinated evolution of

fundamental physiological and non-reproductive life history traits, implicating a broad role of

sexual selection in the evolution of life history strategies.

ACKNOWLEDGEMENTS

We would like to thank Göran Arnqvist and Zorana K. Novicic who kindly collected data on metabolic rates. Andrew Beckerman, Dylan Childs, John Jackson and Björn Rogell provided helpful discussion about statistical analysis and Henry Barton helped relabel images for blind analysis. Many members of the Snook lab contributed to running the experimental sexual selection lines over the years.

FUNDING

This work was funded by the National Environment Research Council (NERC) through the Adapting to the Challenges of a Changing Environment Doctoral Training Partnership (ACCE DTP; NE/L002450/1). Previous funding for the lines came from: the US National Science Foundation (DEB-0093149) and NERC grants (NE/B504065/1, NE/D003741/1, NE/I014632/1).

REFERENCES

1. Darwin C. 1871 *The Descent of Man, and Selection in Relation to Sex*. REV-Revised. Princeton University Press. See https://www.jstor.org/stable/j.ctt19zbz6c.

2. Kokko H, Jennions MD. 2008 Parental investment, sexual selection and sex ratios. *J. Evol. Biol.* **21**, 919–948. (doi:https://doi.org/10.1111/j.1420-9101.2008.01540.x)

3. Lande R. 1980 Sexual Dimorphism, Sexual Selection, and Adaptation in Polygenic Characters. *Evolution* **34**, 292–305. (doi:10.2307/2407393)

4. Cally JG, Stuart-Fox D, Holman L. 2019 Meta-analytic evidence that sexual selection improves population fitness. *Nat. Commun.* **10**, 2017. (doi:10.1038/s41467-019-10074-7)

5. Rowe L, Houle D. 1996 The lek paradox and the capture of genetic variance by condition dependent traits. *Proc. R. Soc. Lond. B Biol. Sci.* **263**, 1415–1421.

6. Lailvaux SP, Irschick DJ. 2006 A functional perspective on sexual selection: insights and future prospects. *Anim. Behav.* **72**, 263–273. (doi:10.1016/j.anbehav.2006.02.003)

7. Orr TJ, Garland T. 2017 Complex reproductive traits and whole-organism performance. *Integr. Comp. Biol.* **57**, 407–422. (doi:10.1093/icb/icx052)

8. Watson PJ, Stallmann RR, Arnqvist G. 1998 Sexual conflict and the energetic costs of mating and mate choice in water striders. *Am. Nat.* **151**, 46–58. (doi:10.1086/286101)

9. Kotiaho JS. 2001 Costs of sexual traits: a mismatch between theoretical considerations and empirical evidence. *Biol. Rev.* **76**, 365–376. (doi:10.1017/S1464793101005711)

10. Hunt J, Brooks R, Jennions MD, Smith MJ, Bentsen CL, Bussière LF. 2004 High-quality
male field crickets invest heavily in sexual display but die young. *Nature* **432**, 1024–
1027. (doi:10.1038/nature03084)

11. Gyulavári HA, Therry L, Dévai G, Stoks R. 2014 Sexual selection on flight endurance,
flight-related morphology and physiology in a scrambling damselfly. *Evol. Ecol.* **28**, 639–
654. (doi:10.1007/s10682-014-9703-1)

12. Debelle A, Courtiol A, Ritchie MG, Snook RR. 2017 Mate choice intensifies motor
signalling in *Drosophila*. *Anim. Behav.* **133**, 169–187.
(doi:10.1016/j.anbehav.2017.09.014)

13. Linklater JR, Wertheim B, Wigby S, Chapman T. 2007 Ejaculate depletion patterns evolve
in response to experimental manipulation of sex ratio in *Drosophila melanogaster*.
Evolution **61**, 2027–2034.

14. Immonen E, Rönn J, Watson C, Berger D, Arnqvist G. 2016 Complex mitonuclear
interactions and metabolic costs of mating in male seed beetles. *J. Evol. Biol.* **29**, 360–
370. (doi:10.1111/jeb.12789)

15. Montooth KL, Marden JH, Clark AG. 2003 Mapping determinants of variation in energy
metabolism, respiration and flight in *Drosophila*. *Genetics* **165**, 623–635.

16. Badyaev AV. 2002 Growing apart: an ontogenetic perspective on the evolution of sexual
size dimorphism. *Trends Ecol. Evol.* **17**, 369–378. (doi:10.1016/S0169-5347(02)02569-7)

17. Hollis B, Keller L, Kawecki TJ. 2017 Sexual selection shapes development and maturation
rates in *Drosophila*. *Evolution* **71**, 304–314. (doi:10.1111/evo.13115)

18. Hämäläinen A, Immonen E, Tarka M, Schuett W. 2018 Evolution of sex-specific pace-of-
life syndromes: causes and consequences. *Behav. Ecol. Sociobiol.* **72**, 50.
(doi:10.1007/s00265-018-2466-x)

19. Pitnick S, Markow TA, Spicer GS. 1995 Delayed male maturity is a cost of producing large
sperm in *Drosophila*. *Proc. Natl. Acad. Sci.* **92**, 10614–10618.

20. Emlen DJ, Warren IA, Johns A, Dworkin I, Lavine LC. 2012 A mechanism of extreme
growth and reliable signaling in sexually selected ornaments and weapons. *Science* **337**,
860–864. (doi:10.1126/science.1224286)

21. Berson JD, Zuk M, Simmons LW. 2019 Natural and sexual selection on cuticular
hydrocarbons: a quantitative genetic analysis. *Proc. R. Soc. B Biol. Sci.* **286**, 20190677.
(doi:10.1098/rspb.2019.0677)

22. Long TAF, Rice WR. 2007 Adult locomotory activity mediates intralocus sexual conflict in
a laboratory-adapted population of *Drosophila melanogaster*. *Proc. R. Soc. B Biol. Sci.*
274, 3105–3112. (doi:10.1098/rspb.2007.1140)

23. Lewis Z, Wedell N, Hunt J. 2011 Evidence for Strong Intralocus Sexual Conflict in the
Indian Meal Moth, Plodia Interpunctella. *Evolution* **65**, 2085–2097.
(doi:https://doi.org/10.1111/j.1558-5646.2011.01267.x)

24. Berger D, Berg EC, Widegren W, Arnqvist G, Maklakov AA. 2014 Multivariate intralocus
sexual conflict in seed beetles. *Evolution* **68**, 3457–3469. (doi:10.1111/evo.12528)

25. Jensen K, McClure C, Priest NK, Hunt J. 2015 Sex-specific effects of protein and
carbohydrate intake on reproduction but not lifespan in <i>Drosophila
melanogaster<\i>. *Aging Cell* **14**, 605–615. (doi:10.1111/acel.12333)

26. Arnqvist G, Stojković B, Rönn JL, Immonen E. 2017 The pace-of-life: A sex-specific link
between metabolic rate and life history in bean beetles. *Funct. Ecol.* **31**, 2299–2309.
(doi:10.1111/1365-2435.12927)

27. Holman L, Jacomb F. 2017 The effects of stress and sex on selection, genetic covariance,
and the evolutionary response. *J. Evol. Biol.* **30**, 1898–1909.
(doi:https://doi.org/10.1111/jeb.13149)

28. Immonen E, Hämäläinen A, Schuett W, Tarka M. 2018 Evolution of sex-specific pace-of-
life syndromes: genetic architecture and physiological mechanisms. *Behav. Ecol.
Sociobiol.* **72**, 60. (doi:10.1007/s00265-018-2462-1)

29. Videlier M, Rundle HD, Careau V. 2019 Sex-Specific Among-Individual Covariation in
Locomotor Activity and Resting Metabolic Rate in *Drosophila melanogaster*. *Am. Nat.*
194, E164–E176. (doi:10.1086/705678)

30. Anderson WW. 1974 Frequent multiple insemination in a natural population of
Drosophila pseudoobscura. *Am. Nat.* **108**, 709–711.

31. Hunt J, Snook RR, Mitchell C, Crudgington HS, Moore AJ. 2012 Sexual selection and
experimental evolution of chemical signals in *Drosophila pseudoobscura*. *J. Evol. Biol.* **25**,
2232–2241. (doi:10.1111/j.1420-9101.2012.02603.x)

32. Snook RR, Robertson A, Crudgington HS, Ritchie MG. 2005 Experimental manipulation of
sexual selection and the evolution of courtship song in *Drosophila pseudoobscura*.
Behav. Genet. **35**, 245–255. (doi:10.1007/s10519-005-3217-0)

33. Debelle A, Ritchie MG, Snook RR. 2014 Evolution of divergent female mating preference
in response to experimental sexual selection. *Evolution* **68**, 2524–2533.
(doi:10.1111/evo.12473)

34. Crudgington HS, Fellows S, Snook RR. 2010 Increased opportunity for sexual conflict
promotes harmful males with elevated courtship frequencies. *J. Evol. Biol.* **23**, 440–446.
(doi:10.1111/j.1420-9101.2009.01907.x)

35. Debelle A, Ritchie MG, Snook RR. 2016 Sexual selection and assortative mating: an
experimental test. *J. Evol. Biol.* **29**, 1307–1316. (doi:10.1111/jeb.12855)

36. Crudgington HS, Fellows S, Badcock NS, Snook RR. 2009 Experimental manipulation of
sexual selection promotes greater male mating capacity but does not alter sperm
investment. *Evolution* **63**, 926–938. (doi:10.1111/j.1558-5646.2008.00601.x)

37. Crudgington HS, Beckerman AP, Brüstle L, Green K, Snook RR. 2005 Experimental
removal and elevation of sexual selection: does sexual selection generate manipulative
males and resistant females? *Am. Nat.* **165**, S72–S87. (doi:10.1086/429353)

38. Immonen E, Snook RR, Ritchie MG. 2014 Mating system variation drives rapid evolution
of the female transcriptome in *Drosophila pseudoobscura*. *Ecol. Evol.* **4**, 2186–2201.
(doi:10.1002/ece3.1098)

39. Veltsos P, Fang Y, Cossins AR, Snook RR, Ritchie MG. 2017 Mating system manipulation
and the evolution of sex-biased gene expression in *Drosophila*. *Nat. Commun.* **8**, 2072.
(doi:10.1038/s41467-017-02232-6)

40. Wiberg RAW, Veltsos P, Snook RR, Ritchie MG. 2021 Experimental evolution supports
signatures of sexual selection in genomic divergence. *bioRxiv* , 2020.09.07.285650.
(doi:10.1101/2020.09.07.285650)

41. Snook RR, Briistle L, Slate J. 2009 A test and review of the role of effective population
size on experimental sexual selection patterns. *Evolution* **63**, 1923–1933.

42. Schneider CA, Rasband WS, Eliceiri KW. 2012 NIH Image to ImageJ: 25 years of image
analysis. *Nat. Methods* **9**, 671–675. (doi:10.1038/nmeth.2089)

43. Lighton JRB. 2008 *Measuring metabolic rates a manual for scientists*. 1 edition. Oxford ;
New York: Oxford University Press.

44. R Core Team. 2020 *R: A language and environment for statistical computing*. Vienna,
Austria: R Foundation for Statistical Computing. See http://www.R-project.org.

45. Bürkner P-C. 2017 brms: An R package for Bayesian multilevel models using Stan. *J. Stat.*
Softw. **80**, 1–28. (doi:doi:10.18637/jss.v080.i01)

46. Schielzeth H, Forstmeier W. 2009 Conclusions beyond support: overconfident estimates
in mixed models. *Behav. Ecol.* **20**, 416–420. (doi:10.1093/beheco/arn145)

47. Husak JF, Swallow JG. 2011 Compensatory traits and the evolution of male ornaments.
Behaviour **148**, 1–29. (doi:10.1163/000579510X541265)

48. Wigglesworth VB. 1949 The utilization of reserve substances in Drosophila during flight.
J. Exp. Biol. **26**, 150–163.

49. Bertram SM, Thomson IR, Auguste B, Dawson JW, Darveau C-A. 2011 Variation in cricket
acoustic mate attraction signalling explained by body morphology and metabolic
differences. *Anim. Behav.* **82**, 1255–1261. (doi:10.1016/j.anbehav.2011.08.021)

50. Becker A, Schlöder P, Steele JE, Wegener G. 1996 The regulation of trehalose
metabolism in insects. *Experientia* **52**, 433–439. (doi:10.1007/BF01919312)

51. Marron MT, Markow TA, Kain KJ, Gibbs AG. 2003 Effects of starvation and desiccation
on energy metabolism in desert and mesic *Drosophila*. *J. Insect Physiol.* **49**, 261–270.
(doi:10.1016/S0022-1910(02)00287-1)

52. Hoffmann AA, Parsons PA. 1989 Selection for Increased Desiccation Resistance in
Drosophila Melanogaster: Additive Genetic Control and Correlated Responses for Other
Stresses. *Genetics* **122**, 837–845.

53. Chippindale AK, Chu TJF, Rose MR. 1996 Complex trade-offs and the evolution of
starvation resistance in *Drosophila melanogaster*. *Evolution* **50**, 753–766.
(doi:10.2307/2410848)

54. Zera AJ, Harshman LG. 2001 The physiology of life history trade-offs in animals. *Annu.*
Rev. Ecol. Syst. **32**, 95–126. (doi:10.1146/annurev.ecolsys.32.081501.114006)

55. Simmons LW, Lüpold S, Fitzpatrick JL. 2017 Evolutionary trade-off between secondary
sexual traits and ejaculates. *Trends Ecol. Evol.* **32**, 964–976.
(doi:10.1016/j.tree.2017.09.011)

56. Harano T, Okada K, Nakayama S, Miyatake T, Hosken DJ. 2010 Intralocus Sexual Conflict
Unresolved by Sex-Limited Trait Expression. *Curr. Biol.* **20**, 2036–2039.
(doi:10.1016/j.cub.2010.10.023)

583