Experimental sexual selection affects the evolution of physiological and life history traits

Running title: Sexual selection affects physiology and life history

Martin D. Garlovsky[1,2+], Luke Holman[3+], Andrew L. Brooks[1], Zorana K. Novicic[4] and Rhonda R. Snook[5]

[1]Department of Animal and Plant Sciences, University of Sheffield, UK

[2]Current address: Applied Zoology, Faculty Biology, Technische Universität Dresden, Germany

[3]School of Applied Sciences, Edinburgh Napier University, UK

[4]Animal Ecology, Department of Ecology and Genetics, Evolutionary Biology Center, Uppsala University, Uppsala, Sweden.

[5]Department of Zoology, Stockholm University, Sweden

[+]These authors contributed equally to this work

Corresponding author: martingarlovsky@gmail.com

ORCiD: MDG: 0000-0002-3426-4341, LH: 0000-0002-7268-2173, ALB: 0000-0002-2556-5249, ZKN: 0000-0001-5157-3229, RRS: 0000-0003-1852-1448.

ACKNOWLEDGEMENTS

We would like to thank Göran Arnqvist who kindly offered the use of his laboratory to measure metabolic rates. Andrew Beckerman, Dylan Childs, John Jackson and Björn Rogell provided helpful discussion about statistical analysis and Henry Barton helped relabel images for blind analysis. Many members of the Snook lab contributed to running the experimental sexual selection lines over the years. This work was funded by the National Environment Research Council (NERC) through the Adapting to the Challenges of a Changing Environment Doctoral Training Partnership (ACCE DTP; NE/L002450/1). Previous funding for the lines came from: the US National Science Foundation (DEB-0093149) and NERC grants (NE/B504065/1, NE/D003741/1, NE/I014632/1).

CONTRIBUTIONS

RRS designed the experiments. ALB, MDG, ZKN, and RRS collected the data. MDG and LH performed statistical analyses. MDG, LH and RRS wrote the manuscript. All authors agreed to the final version of the manuscript.

CONFLICT OF INTERESTS

The authors have no conflict of interest to declare

ABSTRACT

Sexual selection and sexual conflict are expected to affect all aspects of the phenotype, not only traits that are directly involved in reproduction. Here, we show coordinated evolution of multiple physiological and life history traits in response to long-term experimental manipulation of the mating system in populations of *Drosophila pseudoobscura*. Development time was extended under polyandry relative to monogamy in both sexes, potentially due to higher investment in traits linked to sexual selection and sexual conflict. Individuals (especially males) evolving under polyandry had higher metabolic rates and locomotor activity than those evolving under monogamy. Polyandry individuals also invested more in metabolites associated with increased endurance capacity and efficient energy metabolism and regulation, namely lipid and glycogen. Finally, polyandry males were less desiccation- and starvation- resistant than monogamy males, suggesting trade-offs between resistance and sexually selected traits. Our results provide experimental evidence that mating systems can impose selection that influences the evolution of non-sexual phenotypes such as development, activity, metabolism, and nutrient homeostasis.

KEYWORDS: Sexual selection, polyandry, metabolism, physiology, life history evolution, trade-offs, experimental evolution.

INTRODUCTION

Natural and sexual selection often act differently in males and females, which have different routes to evolutionary fitness as a result of anisogamy (Darwin, 1871; Kokko & Jennions, 2008). However, independent evolution of the sexes is partly constrained due to their shared genome (Lande, 1980). Recent work has increasingly highlighted that sexual selection affects all aspects of the phenotype, and not only classic sexually-selected traits such as male ornaments (reviewed in Cally *et al.*, 2019). This is because sexually-selected traits often depend on overall condition (Rowe & Houle, 1996), which in turn depends on underlying aspects of organisms such as their development, physiology, life history and metabolism (Lailvaux & Irschick, 2006; Orr & Garland, 2017).

Sexual selection often favours energetically costly bouts of sustained locomotor activity, e.g. during mate searching, courtship, or the 'harassment and resistance' behaviours that typify interlocus sexual conflict (Watson *et al.*, 1998; Kotiaho, 2001; Hunt *et al.*, 2004; Gyulavári *et al.*, 2014; Debelle *et al.*, 2017). Moreover, postcopulatory sexual selection can select for males that produce metabolically expensive ejaculates (Linklater *et al.*, 2007; Immonen *et al.*, 2016). We therefore expect sexual selection and sexual conflict to favour physiological adaptations that augment the efficiency of metabolism and respiration (Montooth *et al.*, 2003; Gyulavári *et al.*, 2014). Sexual selection is also hypothesised to affect the evolution of development and life history, potentially in a sex-specific way (Badyaev, 2002; Stångberg *et al.*, 2020). For instance, in populations experiencing heightened sexual selection, male *Drosophila melanogaster* evolved increased activity and courtship frequency and shorter lifespan (Nandy *et al.*, 2013), and in a separate experiment, males experiencing heightened sexual selection evolved faster development time, whereas females developed faster under monogamy (Hollis *et al.*, 2017). Likewise, selection for early life reproduction in *Acanthoscelides obtectus* beetles favoured higher metabolic rate in males (Arnqvist *et al.*, 2017), and in *Callosobruchus maculatus* beetles, females (but not males) evolved under polygamy had higher mortality and aging rates than females evolved under monogamy (Maklakov *et al.*, 2007). Thus, selection may cause trade-offs between different aspects of fitness, for instance, favouring a 'live fast, die young' strategy in species where competition for mates is intense (Nandy *et al.*, 2013; Hollis *et al.*, 2017; Hämäläinen *et al.*, 2018) or selecting for the reallocation of limiting resources away from growth and somatic maintenance (Pitnick *et al.*, 1995; Hunt *et al.*, 2004; Emlen *et al.*, 2012; Berson *et al.*, 2019). Indeed, the conspicuous fitness trade-offs associated with sexually-selected traits was a key motivation for the theory of sexual selection (Darwin, 1871).

In addition to these links between sexually selected traits and other phenotypes, the picture is further complicated by genetic correlations between the sexes. Male and female traits have a shared genetic basis, such that selection on males results in a (frequently maladaptive) evolutionary response in females, and *vice versa* e.g. (Long & Rice, 2007; Harano *et al.*, 2010; Lewis *et al.*, 2011; Berger *et al.*, 2014; Jensen *et al.*, 2015; Arnqvist *et al.*, 2017; Holman &

Jacomb, 2017; Hämäläinen *et al.*, 2018; Immonen *et al.*, 2018; Videlier *et al.*, 2019). Therefore,
sexual selection on males produces an evolutionary response in female, as well as male, life
history traits and vice versa. For instance, male-limited selection for short life span in *C.*
maculatus caused a correlated response in females (Berger *et al.*, 2014), and in *D.*
melanogaster, male-limited evolution selected for increased activity in both sexes, to the
detriment of female fitness (Long & Rice, 2007). While these studies have measured the
response of individual or several key physiological or life history traits to the strength of sexual
selection, in this paper, we investigate the effect of variation in the strength of sexual
selection on multiple inter-related life history and physiological traits to capture a broader
range of consequences of sexual selection.
134
We use the fruit fly, *Drosophila pseudoobscura*, subjected to either experimentally enforced
monogamy (M) that eliminated sexual selection and sexual conflict, or to elevated polyandry
(E), where 6 males were housed with 1 female to facilitate inter- and intra- sexual selection,
likely at elevated levels compared to natural populations (Anderson, 1974). For each sexual
selection treatment, there were four replicate populations (hereafter 'lines'). Previous studies
have found divergence between the E and M lines in several traits important during episodes
of sexual selection. For example, E males produce more abundant and complex chemical
signals (Hunt *et al.*, 2012), perform a faster and more vigorous courtship song (Snook *et al.*,
2005; Debelle *et al.*, 2014, 2017), court and mate more frequently (Crudgington *et al.*, 2010),
are more competitive in mating encounters (Debelle *et al.*, 2016), and have larger glands for
producing seminal fluid (Crudgington *et al.*, 2009). Additionally, coevolutionary patterns have
been found, such that E males are more harmful but E females more resilient to such harm
(Crudgington *et al.*, 2005, 2009), and females prefer courtship song of males from their own
treatment (Debelle *et al.*, 2014). These evolved phenotypic differences are accompanied by
divergence in gene expression (Immonen *et al.*, 2014; Veltsos *et al.*, 2017) and concerted
genetic differences between treatments (Wiberg *et al.*, 2021).
151
Given these evolved differences in traits directly associated with reproduction, we tested
whether the mating system similarly caused divergence in life history and physiological
phenotypes that likely contribute to reproductive success, namely metabolic rate, macro-
metabolite content, development time, and stress resistance. Assuming that these traits
contribute to reproductive success, we predict that they may have diverged between the E
and M lines, reflecting the different behaviours, life history decisions, and physiological states
that maximise fitness under the different selective regimes. Due to the genetic non-
independence of the sexes, the multiple differences in selection on both sexes selection in
the E and M treatments, and the genetic correlations between the variables examined, the
direction of evolutionary change is difficult to predict, making our study exploratory in nature.
However, we predict higher activity levels and metabolic rates in the E treatment in both
sexes, due to the elevated importance of courtship, harassment, and resistance behaviours.
As such, elevated metabolism might require altered resource allocation, affecting

macrometabolite composition, development time, and the ability to resist stressors. Shorter
development may be favoured by sexual selection (Hollis *et al.*, 2017), however the effects of
sexual selection on macrometabolite profile is unknown *a priori*.
168
METHODS
Establishment and maintenance of experimental evolution lines
Details of the establishment and maintenance of the experimental evolution lines has been
previously described (Crudgington *et al.*, 2005). Briefly, the ancestral population was
established from 50 wild-caught, inseminated female *Drosophila pseudoobscura* from
Tucson, Arizona in 2001; the descendants of these females were used to establish four
replicate populations (termed lines) for each sexual selection treatment. In the enforced
monogamy or 'M' treatment, the adult population was housed in groups of two (one male,
one female); this protocol reduces the opportunity for sexual selection and sexual conflict. In
the elevated polyandry or 'E' treatment, each group of adults comprised six males and one
female. Each M line contained a greater number of groups (n = 80) than each E line (n = 40),
to offset the smaller group size and thereby minimise differences between treatments in the
autosomal effective population size (Snook *et al.*, 2009). In each generation, unmated males
and females were housed in 'interaction vials' (IVs) for 5 days, before being transferred to
'oviposition vials' (OVs) for a further 5 days; using both IVs and OVs reduces larval competition
and provides more opportunity for episodes of pre- and post-copulatory sexual selection. To
facilitate selection favouring the most productive groups (and the most competitive males,
for the E treatment), offspring from all groups within each line were gathered and mixed in a
single container, and a random sample of all the offspring was used to set up the next
generation. Flies were kept at 22ºC on a 12:12 light:dark cycle on standard cornmeal-agar-
molasses media with added live yeast.
190
Experimental individuals
Prior to the experiments described below, flies from each line were taken out of selection and
placed in a 'common garden' to minimise non-genetic differences between lines. Newly-
eclosed individuals were collected *en masse* from the OVs within each line. A random sample
of these flies were allowed to mate and oviposit for two days. From these eggs, we set up
controlled density vials (CDVs) by placing 100 first instar larvae into food vials. Unmated flies
eclosing from CDVs were collected and stored in same-sex food vials until 3-5 days old.
Excluding the development time experiment each measurement in our study was performed
on a 'triad' of three age-matched, same sex flies within each line, due to practical constraints
resulting from the small size of individual flies.
201
Juvenile development time
We measured juvenile development time at generations 180, 179, 178 and 176 for lines 1-4,
respectively. For each replicate of the E and M treatments, we seeded 6 CDVs on three
consecutive days (i.e., 600 larvae per replicate population per seeding day = 14,400 larvae).

Vials were checked daily for new eclosees, and flies were CO_2 anaesthetised and killed in
ethanol. We continued collecting until no individuals eclosed for two consecutive days. We
subsequently counted the number of adult males and females emerging each day from each
vial.
210
We used the length of wing vein four as a proxy for body size (Crudgington *et al.*, 2005) by
measuring a random subsample of individuals (n = 15 per sex per replicate per seeding day).
We removed the left wing from flies preserved in ethanol and mounted wings on a
microscope slide in a drop of phosphate-buffered saline and dried at room temperature
overnight. We imaged wings using a Motic camera and Motic Images Plus 2.0 software (Motic
Asia, Hong Kong). We measured wing vein four length using ImageJ software (Schneider *et
al.*, 2012). Image files were anonymised prior to measurement.
218

Desiccation and starvation resistance
We measured desiccation and starvation resistance at generations 199, 198, 197 and 195 for
lines 1-4, respectively. Triads (n = 7-10) were housed in 8-dram plastic vials stoppered with
cotton balls and covered with Parafilm®. For the desiccation resistance assay, vials contained
no food and between the cotton and Parafilm® we placed a packet of silica gel beads. For the
starvation resistance assay, vials contained an agar solution that provided moisture but no
food. Vials were checked every 2 hours and any deaths recorded until all flies perished. Flies
were scored as dead if they were not able to right themselves or no movement was observed.
227

Respirometry
We measured metabolic rates at generations 196, 195, 194 and 192 for lines 1-4, respectively.
Triads (n = 3) were weighed to the nearest 0.1 mg (Sartorius Genius ME 235P-OCE) before
transfer to a respirometry chamber (a glass cylinder; 17mm × 70 mm). Metabolic rate was
measured using a Sable Systems (Las Vegas, NV, USA) respirometry system (Lighton, 2008).
This system pumps air at a precise flow rate through a sealed chamber containing the
organisms undergoing measurement. Downstream gas analysers measure two response
variables: the amount of CO_2 produced and O_2 consumed. Briefly, the respirometry system
was set up in stop-flow mode (Lighton, 2008), in which each chamber was sealed for 60 min
and then flushed for 2.5 min. Each cycle (through all 24 chambers) lasted for 62.5 minutes,
and each triad of flies was recorded over consecutive cycles, giving four readings of CO_2 and
O_2 flux in each individual chamber. The first recording was discarded as a wash-out, while the
other three were used for analyses. Each respirometry chamber was placed in an activity
detector (AD-2, Sable Systems) connected to a data acquisition interface (Quick-DAQ,
National Instruments, Coleman Technologies, Newton Square, US), which uses reflective
infrared-light technology to provide a precise and continuous measure of locomotor activity
of the subjects; one activity measurement per triad per cycle was recorded. One of the 24
chambers was left empty and used as a baseline to control for any drift of the gas analysers
during each session (washed out twice in each cycle). Thus, each observation consisted of

three consecutive readings of the amount of CO_2 produced and O_2 consumed during 62.5
minutes by a triad of flies, under dark conditions, with a known weight and total amount of
activity performed. We also calculated the respiratory quotient (RQ; i.e., VCO_2 / VO_2), which
describes the metabolic substrate used for respiration, where a value of 0.7 indicates pure
fatty acid oxidation, 1.0 indicates pure carbohydrate oxidation, and intermediate values
indicate mixed substrate or protein-based oxidation

Metabolite extractions
We measured metabolite composition at generations 199, 198, 197 and 195 for lines 1-4,
respectively. Triads (n = 3) were weighed to the nearest 1μg (METTLER TOLEDO® UMX2 ultra-
microbalance) and flash frozen in liquid nitrogen. Triads were then placed in a 0.35ml glass
vial insert (SUPELCO Analytical®) of known weight, dried at 55ºC overnight, and re-weighed
to obtain dry weight. We subsequently processed each sample to acquire measurements of
lipids, soluble carbohydrates, soluble protein, glycogen, and chitin (see online supplementary
material).

Statistical analysis
All statistical analyses were performed in R version 4.0.3 (R Core Team, 2020). We fit models
using Bayesian approaches in *brms* (Bürkner, 2017), and specified conservative, regularising
priors on the fixed and random effects. Complete code and description of models are
provided in the code repository (https://anonymous.4open.science/r/exp_evol_respiration-
2647). In brief, we analysed development time and survival using survival analysis. For the
respirometry and metabolite data we used structural equation models (SEM), motivated by
the causal relationships that we hypothesise to exist between the variables, given the
experimental design and our biological understanding of the system (Fig. S1). For the
metabolite data, we began by expressing the abundance of each metabolite as a proportion
of dry weight and scaled each response variable to have mean zero and unit standard
deviation. For the SEMs, after fitting the model, we calculated the treatment effect size
(Cohen's *d*) for each response variable, both with and without the moderators (dry weight
and/or activity). We also calculated the difference in treatment effect size between the sexes,
to test for a treatment × sex interaction. All models included replicate line as a random
intercept to reflect the experimental design of the selection experiment (i.e. where n = 8). We
also performed analyses with selection treatment fitted as a random slope to allow lines to
vary in their response to selection (Schielzeth & Forstmeier, 2009), however results were
qualitatively identical, and thus we present only the results of models without random slopes
in the Results. All models also included vial identity as a random intercept, except for wing
vein length where we measured a random sample of individuals across multiple vials.

RESULTS

Juvenile development time

Juvenile development time differed significantly between the E and M treatments (Cox proportional hazards model; PP = 0.008, where PP is the posterior probability the effect is of the opposite direction, similar to a 1-tailed p-value) and between the sexes (PP < 0.001). E treatment flies had a reduced hazard (Hazard ratio = 0.44; 95% confidence intervals [CI] = 0.26 − 0.81), i.e., development took longer in the E treatment in both sexes. Males took longer to eclose than females (Hazard ratio = 0.84; 95% CI = 0.80 − 0.88) (Fig. 1A), as usual in this species. There was a significant treatment × sex interaction (PP = 0.007), such that the sex difference in development time was greater in the E treatment, and the treatment affected males more strongly than females.

As expected, females were larger than males (mean wing vein IV length (μm); M females: 2324 ± 4.80, n = 152; E females: 2335 ± 5.76, n = 118; M males: 2099 ± 4.93, n = 154; E males: 2114 ± 5.96, n = 127). We found no statistically significant effect of sexual selection treatment (PP = 0.312) or the treatment × sex interaction (PP = 0.382).

Desiccation and starvation resistance

We found a significant treatment × sex interaction for both desiccation and starvation resistance (desiccation: PP = 0.024; starvation: PP = 0.018). E males survived 0.91 (95% CI = 0.83 − 0.99), and 0.88 (95% CI = 0.80 − 0.97), times as long as M males under desiccation and starvation resistance, respectively (Fig. 1B-C). However, the treatment effect was opposite in females: E females survived 1.06 (95% CI = 0.87 − 1.29), and 1.09 (95% CI = 0.88 − 1.35), times longer than M females under desiccation and starvation resistance, respectively. In short, M males survived longer than E males, while E females survived longer than M females.

Figure 1: Kaplan-Meier plots for (a) juvenile development time, (b) desiccation resistance, and (c) starvation resistance. '+' indicates right censored individuals. Monogamy (M) = pink and light blue, Elevated polyandry (E) = red and dark blue; Males = dashed lines, Females = solid lines. Shaded areas show confidence intervals.

Respirometry

Beginning with the mediator variables, females were heavier than males (PP < 0.001; Table
S1), and E females were non-significantly heavier than M females (Fig. 2A; Mean dry weight
per triad (mg): M females: 0.40 ± 0.02; E females: 0.47 ± 0.04; M males: 0.29 ± 0.02; E males:
0.29 ± 0.02; n = 12 each). The treatment (PP = 0.075) and treatment × sex (PP = 0.105) terms
were not significantly related to body mass. There was a significant effect of sexual selection
treatment on activity level (PP < 0.001), with E flies more active than M flies (Fig. 2A). Activity
level also showed a sex × cycle interaction (PP < 0.026), indicating that activity levels declined
over cycles in males but not females. This sex difference in the rate of decline was stronger in
M treatment than E treatment, though the 3-way interaction was not statistically significant
(PP = 0.066).

There was a strong correlation between activity level and O_2 consumption (PP < 0.001; Fig.
S2), and O_2 consumption declined across cycles (PP < 0.001). Body weight did not correlate
with O_2 consumption (PP = 0.119; Fig. S2). Calculating the posterior estimate of the difference
in treatment means for each sex and cycle revealed that O_2 consumption was higher in the E
treatment (indicated by positive effect sizes in Fig. 2B), especially in males. This effect was
considerably attenuated when we statistically adjusted for differences in activity level
between treatments and sexes, indicating that the evolved difference in activity between the
E and M treatments was largely responsible for the treatment effect on O_2 consumption.
Controlling for the evolved differences in body mass did not change the effect size, illustrating
that the (modest) changes in body mass between the E and M treatments did not explain the
evolved difference in O_2 consumption.

The respiratory quotient (RQ) did not differ detectably between the E and M treatments,
whether or not one controlled for the mediator variables (Fig. S3). The grand mean RQ = 0.90
(± 0.02) indicated flies used a mixture of carbohydrates, protein, and lipids as metabolic
substrate.

349
Figure 2: Effects of sex and selection treatment on metabolic rates and mediator variables. A)
Posterior estimates of the means for mediator variables (body mass and activity). Note that
females are larger than males, and that E females are somewhat larger than M females. B)

Posterior estimates of effect sizes of selection treatment (E minus M) on metabolic rate
(measured as volume of O2 consumed) controlling for activity, body mass, or neither
mediator variable. The x-axes show the posterior estimate of standardised effect size
(Cohen's d), i.e., a value of 1 would mean that the E treatment has a mean that is larger by 1
standard deviation. The horizontal bars show the 66% and 95% quantiles and the median of
the posterior distribution. All response variables have been mean-centred and divided by the
overall standard deviation (such that the dashed line at zero marks the mean across sexes,
treatments, and cycles). For brevity, only the first cycle is shown for activity and metabolic
rate; see online supplementary material for all cycles.

Metabolite composition

As in the respiration experiment, females were heavier than males and E flies were heavier
than M flies (Dry weight (mg): M females: 0.56 ± 0.02; E females: 0.64 ± 0.17; M males: 0.33
± 0.01; E males: 0.35 ± 0.01; n = 12 each; Table S2). There was a significant treatment × sex
interaction (P = 0.038), such that the E treatment positively affected body mass more strongly
in females. Furthermore, dry weight was significantly correlated with lipid and chitin content
(Fig. S4; Table S2), and therefore could act as a mediator for some of the effect of sex and
treatment on these metabolites.

We also found differences in metabolite composition between the E and M treatments and
between sexes (Fig. 3). E treatment flies contained more lipids and glycogen than M flies,
while M flies had relatively more carbohydrates and chitin; the amount of protein did not
differ between treatments. These treatment differences reached statistical significance in
males for carbohydrate, glycogen, and chitin, and in females for lipid. When we controlled for
the evolved differences in dry weight between the E and M treatments, only the carbohydrate
and glycogen differences between E and M males remained statistically significant; however,
the actual change in effect size when controlling for dry weight was very small, indicating that
dry weight was not an especially important mediator of the evolved changes in metabolite
composition (compare Fig. 3 and Fig. S5).

<u>Figure 3.</u> Posterior estimates of the treatment effect size for both sexes, for each of the five
metabolites. A positive value indicates that the mean metabolite content is higher in the E
treatment than the M treatment, while a negative value denotes M > E. A strongly supported
treatment effect is implied by most of the posterior lying to one side of zero. The vertical bars
show the 66% and 95% quantiles and the median of the posterior distribution. This plot was
created using posterior predictions of the means that were not adjusted for differences in dry
weight between treatments.
391
392 Finally, we investigated the treatment × sex interaction term by calculating the posterior
393 difference in the treatment effect size between sexes. Though none of the results reached
394 statistical significance, there was some indication that the decrease in carbohydrate content
395 (in E relative to M) was greater in males than females (PP = 0.092), and that the sexes showed
396 different changes in protein content (PP = 0.11) (Fig. S6).
397

<u>DISCUSSION</u>
We found that experimental manipulation of sexual selection and sexual conflict caused
several physiological and life history traits to evolve. First, flies from the elevated polyandry
(E) treatment took longer to develop into adults than their monogamous (M) counterparts
and the difference in development time between the sexes was greater in the E lines. Second,
E treatment flies of both sexes had faster metabolic rates, largely due to E line flies being
more active than M line flies. Third, macro-metabolite composition has diverged between
sexual selection treatments, such that E females had a greater lipid content than M females,
E males greater glycogen content than M males, and E males less chitin and sugar content
than M males. Finally, E males were less resistant to desiccation and starvation than were M
males, whereas E females were more resistant than M females.
409

E males produce a faster and more vigorous courtship song (Debelle *et al.*, 2014, 2017), more
cuticular hydrocarbons (Hunt *et al.*, 2012), and larger accessory glands (Crudgington *et al.*,
2009) than M males. These sexually selected traits may be energetically costly to produce and
maintain, demanding greater metabolic activity (Montooth *et al.*, 2003; Immonen *et al.*, 2016;
Berson *et al.*, 2019). Metabolic rate showed a sex-specific response to selection for early life
reproduction in bean beetles (Arnqvist *et al.*, 2017). However, locomotor activity and
metabolic rate have a significant intersexual genetic correlation in *D. melanogaster* (Long &
Rice, 2007; Nandy *et al.*, 2013; Videlier *et al.*, 2021). Therefore, sexual selection favouring
increased activity and metabolic rate in males may generate sexual conflict by increasing
activity and metabolic rate above optimal levels for females (Hämäläinen *et al.*, 2018). We
found no difference between treatments or sexes in the respiratory quotient (RQ), which
describes metabolic substrate use, despite evolved differences in activity and metabolic rate.
The lack of evolution of RQ may suggest that E flies have evolved compensatory physiological
mechanisms to offset the metabolic costs of increased energy expenditure (Husak & Swallow,
2011).

We found that E males had higher glycogen content than M males. Carbohydrates are the main fuel used during intense aerobic activities, such as flight (Wigglesworth, 1949) and courtship (Bertram *et al.*, 2011), and glycogen provides the main source of trehalose marshalled during intense activity (Becker *et al.*, 1996). Sexual selection favouring endurance capacity may also increase lipid respiration and fat storage (Gyulavári *et al.*, 2014). However, despite male courtship song being an endurance contest (Debelle *et al.*, 2017), in males we found no difference between treatments in lipid content. Limited time and resources may demand strategic resource allocation decisions. We did find higher lipid content in E females, possibly reflecting selection for endurance capacity, as E females are courted more frequently (Crudgington *et al.*, 2005; Debelle *et al.*, 2014). No metabolites showed a significant treatment × sex interaction, indicating the response to sexual selection treatment did not differ between the sexes, potentially highlighting a strong intersexual genetic correlation constraining divergence of physiological traits (Videlier *et al.*, 2021; Wittman *et al.*, 2021). Our results suggest that heightened sexual selection and sexual conflict favours storage and regulation of metabolites to meet increased metabolic demands (Montooth *et al.*, 2003; Crudgington *et al.*, 2009; Debelle *et al.*, 2014, 2017; Gyulavári *et al.*, 2014), whereas relaxed sexual selection may instead enable investment in other components of fitness.

E males were less tolerant of desiccation and starvation than M males despite their greater glycogen content, which buffers against these stresses (Marron *et al.*, 2003). Previous work artificially selecting on desiccation tolerance in *D. melanogaster* found selected lines had lower metabolic rates (Hoffmann & Parsons, 1989); our experiment imposed different selective pressures but found the same negative correlation between metabolic rate and stress resistance. The E lines have greater cuticular hydrocarbon (CH) content but there is no difference in the abundance of long-chain CHs (which offer greater desiccation resistance) between treatments (Hunt *et al.*, 2012). That E males are less stress resistant despite having more macromolecules which buffer against such stressors suggests a trade-off, where heightened sexual selection favours investment in current reproduction at a cost to later-life survival, while the relaxation of sexual selection favours increased investment in longevity (Kotiaho, 2001; Hunt *et al.*, 2004; Nandy *et al.*, 2013). In contrast, E females showed no such trade-off, as E females were as stress resistant as M females, despite investing more in fecundity (Crudgington *et al.*, 2005; Immonen *et al.*, 2014). The greater lipid content we found in E females could mitigate such a trade-off, because lipids provide protection against starvation alongside being a major component of eggs (Chippindale *et al.*, 1996). Thus, the sexes have altered investment in macro-metabolite composition under sexual selection, which may allow differential investment in sexually selected traits and sexual conflict persistence and resistance traits. The subsequent fitness consequences during stress have different outcomes on the sexes. This suggests sex-specific selection in the adult stage which could further generate sexual conflict over shared traits, particularly during the juvenile stage (Badyaev, 2002).

Related to the juvenile stage, in both sexes, E flies took significantly longer to reach adulthood
than flies evolving under monogamy, perhaps due to pleiotropic costs of increased
investment in traits involved in sexual selection or sexual conflict (Zera & Harshman, 2001;
Simmons *et al.*, 2017; Berson *et al.*, 2019). Moreover, because development time is probably
genetically correlated across sexes (Lewis *et al.*, 2011; Berger *et al.*, 2014), the fact that both
sexes evolved longer development might reflect intralocus sexual conflict. For example, E
males might have evolved to invest more in sexual traits, slowing their development and
causing a similar delay in E females because of pleiotropy. This would be similar to the results
of Harano et al. (Harano *et al.*, 2010), who selected for male beetles with larger mandibles (a
sexually-selected trait), which caused maladaptive morphological evolution in females
despite females lacking the male trait that was under selection (Harano *et al.*, 2010).
Conversely, longer development might benefit females, e.g. by enabling increased resource
acquisition and allocation towards reproduction or resistance to male harm (Crudgington *et*
al., 2005, 2010; Immonen *et al.*, 2014). The sex difference in development time was also
greater in the E lines, possibly resulting from evolution towards the (shorter) female optimum
under monogamy.
482
To conclude, we have shown that experimental manipulation of sexual selection and sexual
conflict caused the evolution of several traits that are not directly involved in sexual
interactions. Differences in activity levels and metabolic rate coincided with differential
investment in macro-metabolites, reflecting the energetic demands associated with the
elevation or removal of sexual selection. Males evolving under elevated polyandry suffered
reduced survival, potentially trading off current vs. future reproduction. However, the same
was not true of females, perhaps due to selection favouring endurance capacity in females to
offset the costs of frequent male courtship. The physiological and life history traits we
measured likely have a shared genetic basis, and thus may be subject to intralocus sexual
conflict. The magnitude of any sexual dimorphism did not differ between treatments for the
physiological traits we measured (macro-metabolites and metabolic rate), whereas life
history traits (development time and stress resistance) were more dimorphic in populations
evolving under elevated polyandry, suggesting tighter intersexual genetic correlations over
physiological traits. Overall, our findings highlight that sexual selection results in coordinated
evolution of fundamental physiological and non-reproductive life history traits, implicating
sexual selection as an important factor in the evolution of life history strategies.
499
500   **DATA AVAILABILITY STATEMENT**
501   Data and code are available from the Dryad digital repository:
502   https://doi.org/10.5061/dryad.9cnp5hqhk, and GitHub:
503   https://lukeholman.github.io/exp_evol_respiration/
504
505
506

REFERENCES

Anderson, W.W. 1974. Frequent multiple insemination in a natural population of *Drosophila*
pseudoobscura. *Am. Nat.* **108**: 709–711.

Arnqvist, G., Stojković, B., Rönn, J.L. & Immonen, E. 2017. The pace-of-life: A sex-specific link
between metabolic rate and life history in bean beetles. *Funct. Ecol.* **31**: 2299–2309.

Badyaev, A.V. 2002. Growing apart: an ontogenetic perspective on the evolution of sexual
size dimorphism. *Trends Ecol. Evol.* **17**: 369–378.

Becker, A., Schlöder, P., Steele, J.E. & Wegener, G. 1996. The regulation of trehalose
metabolism in insects. *Experientia* **52**: 433–439.

Berger, D., Berg, E.C., Widegren, W., Arnqvist, G. & Maklakov, A.A. 2014. Multivariate
intralocus sexual conflict in seed beetles. *Evolution* **68**: 3457–3469.

Berson, J.D., Zuk, M. & Simmons, L.W. 2019. Natural and sexual selection on cuticular
hydrocarbons: a quantitative genetic analysis. *Proc. R. Soc. B Biol. Sci.* **286**:
20190677.

Bertram, S.M., Thomson, I.R., Auguste, B., Dawson, J.W. & Darveau, C.-A. 2011. Variation in
cricket acoustic mate attraction signalling explained by body morphology and
metabolic differences. *Anim. Behav.* **82**: 1255–1261.

Bürkner, P.-C. 2017. brms: An R package for Bayesian multilevel models using Stan. *J. Stat.*
Softw. **80**: 1–28.

Cally, J.G., Stuart-Fox, D. & Holman, L. 2019. Meta-analytic evidence that sexual selection
improves population fitness. *Nat. Commun.* **10**: 2017.

Chippindale, A.K., Chu, T.J.F. & Rose, M.R. 1996. Complex trade-offs and the evolution of
starvation resistance in *Drosophila melanogaster*. *Evolution* **50**: 753–766.

Crudgington, H.S., Beckerman, A.P., Brüstle, L., Green, K. & Snook, R.R. 2005. Experimental
removal and elevation of sexual selection: does sexual selection generate
manipulative males and resistant females? *Am. Nat.* **165**: S72–S87.

Crudgington, H.S., Fellows, S., Badcock, N.S. & Snook, R.R. 2009. Experimental manipulation
of sexual selection promotes greater male mating capacity but does not alter sperm
investment. *Evolution* **63**: 926–938.

Crudgington, H.S., Fellows, S. & Snook, R.R. 2010. Increased opportunity for sexual conflict
promotes harmful males with elevated courtship frequencies. *J. Evol. Biol.* **23**: 440–
446.

Darwin, C. 1871. *The Descent of Man, and Selection in Relation to Sex*, REV-Revised.
Princeton University Press.

Debelle, A., Courtiol, A., Ritchie, M.G. & Snook, R.R. 2017. Mate choice intensifies motor
signalling in *Drosophila*. *Anim. Behav.* **133**: 169–187.

Debelle, A., Ritchie, M.G. & Snook, R.R. 2014. Evolution of divergent female mating
preference in response to experimental sexual selection. *Evolution* **68**: 2524–2533.

Debelle, A., Ritchie, M.G. & Snook, R.R. 2016. Sexual selection and assortative mating: an
experimental test. *J. Evol. Biol.* **29**: 1307–1316.

Emlen, D.J., Warren, I.A., Johns, A., Dworkin, I. & Lavine, L.C. 2012. A mechanism of extreme
growth and reliable signaling in sexually selected ornaments and weapons. *Science*
337: 860–864.

Grieshop, K., Maurizio, P.L., Arnqvist, G. & Berger, D. 2021. Selection in males purges the
mutation load on female fitness. *Evol. Lett.* **n/a**.

Gyulavári, H.A., Therry, L., Dévai, G. & Stoks, R. 2014. Sexual selection on flight endurance,
flight-related morphology and physiology in a scrambling damselfly. *Evol. Ecol.* **28**:
639–654.

Hämäläinen, A., Immonen, E., Tarka, M. & Schuett, W. 2018. Evolution of sex-specific pace-
of-life syndromes: causes and consequences. *Behav. Ecol. Sociobiol.* **72**: 50.

Harano, T., Okada, K., Nakayama, S., Miyatake, T. & Hosken, D.J. 2010. Intralocus Sexual
Conflict Unresolved by Sex-Limited Trait Expression. *Curr. Biol.* **20**: 2036–2039.
Elsevier.

Hoffmann, A.A. & Parsons, P.A. 1989. Selection for Increased Desiccation Resistance in
Drosophila Melanogaster: Additive Genetic Control and Correlated Responses for
Other Stresses. *Genetics* **122**: 837–845.

Hollis, B., Keller, L. & Kawecki, T.J. 2017. Sexual selection shapes development and
maturation rates in *Drosophila*. *Evolution* **71**: 304–314.

Holman, L. & Jacomb, F. 2017. The effects of stress and sex on selection, genetic covariance,
and the evolutionary response. *J. Evol. Biol.* **30**: 1898–1909.

Hunt, J., Brooks, R., Jennions, M.D., Smith, M.J., Bentsen, C.L. & Bussière, L.F. 2004. High-
quality male field crickets invest heavily in sexual display but die young. *Nature* **432**:
1024–1027.

Hunt, J., Snook, R.R., Mitchell, C., Crudgington, H.S. & Moore, A.J. 2012. Sexual selection and
experimental evolution of chemical signals in *Drosophila pseudoobscura*. *J. Evol. Biol.*
25: 2232–2241.

Husak, J.F. & Swallow, J.G. 2011. Compensatory traits and the evolution of male ornaments.
Behaviour **148**: 1–29.

Immonen, E., Hämäläinen, A., Schuett, W. & Tarka, M. 2018. Evolution of sex-specific pace-
of-life syndromes: genetic architecture and physiological mechanisms. *Behav. Ecol.*
Sociobiol. **72**: 60.

Immonen, E., Rönn, J., Watson, C., Berger, D. & Arnqvist, G. 2016. Complex mitonuclear
interactions and metabolic costs of mating in male seed beetles. *J. Evol. Biol.* **29**:
360–370.

Immonen, E., Snook, R.R. & Ritchie, M.G. 2014. Mating system variation drives rapid
evolution of the female transcriptome in *Drosophila pseudoobscura*. *Ecol. Evol.* **4**:
2186–2201.

Jensen, K., McClure, C., Priest, N.K. & Hunt, J. 2015. Sex-specific effects of protein and
carbohydrate intake on reproduction but not lifespan in <i>Drosophila
melanogaster<\i>. *Aging Cell* **14**: 605–615.

Kokko, H. & Jennions, M.D. 2008. Parental investment, sexual selection and sex ratios. *J.*
Evol. Biol. **21**: 919–948.

Kotiaho, J.S. 2001. Costs of sexual traits: a mismatch between theoretical considerations
and empirical evidence. *Biol. Rev.* **76**: 365–376.

Lailvaux, S.P. & Irschick, D.J. 2006. A functional perspective on sexual selection: insights and
future prospects. *Anim. Behav.* **72**: 263–273.

Lande, R. 1980. Sexual Dimorphism, Sexual Selection, and Adaptation in Polygenic
Characters. *Evolution* **34**: 292–305.

Lewis, Z., Wedell, N. & Hunt, J. 2011. Evidence for Strong Intralocus Sexual Conflict in the
Indian Meal Moth, Plodia InterpuncTELla. *Evolution* **65**: 2085–2097.

Lighton, J.R.B. 2008. *Measuring metabolic rates a manual for scientists*, 1 edition. Oxford
University Press, Oxford ; New York.

Linklater, J.R., Wertheim, B., Wigby, S. & Chapman, T. 2007. Ejaculate depletion patterns
evolve in response to experimental manipulation of sex ratio in *Drosophila*
melanogaster. *Evolution* **61**: 2027–2034.

Long, T.A.F. & Rice, W.R. 2007. Adult locomotory activity mediates intralocus sexual conflict
in a laboratory-adapted population of *Drosophila melanogaster*. *Proc. R. Soc. B Biol.*
Sci. **274**: 3105–3112.

Maklakov, A.A., Fricke, C. & Arnqvist, G. 2007. Sexual selection affects lifespan and aging in
the seed beetle. *Aging Cell* **6**: 739–744.

Marron, M.T., Markow, T.A., Kain, K.J. & Gibbs, A.G. 2003. Effects of starvation and
desiccation on energy metabolism in desert and mesic *Drosophila*. *J. Insect Physiol.*
49: 261–270.

Montooth, K.L., Marden, J.H. & Clark, A.G. 2003. Mapping determinants of variation in
energy metabolism, respiration and flight in *Drosophila*. *Genetics* **165**: 623–635.

Nandy, B., Gupta, V., Sen, S., Udaykumar, N., Samant, M.A., Ali, S.Z., *et al.* 2013. Evolution of
mate-harm, longevity and behaviour in male fruit flies subjected to different levels of
interlocus conflict. *BMC Evol. Biol.* **13**: 212.

Orr, T.J. & Garland, T. 2017. Complex reproductive traits and whole-organism performance.
Integr. Comp. Biol. **57**: 407–422.

Pitnick, S., Markow, T.A. & Spicer, G.S. 1995. Delayed male maturity is a cost of producing
large sperm in *Drosophila*. *Proc. Natl. Acad. Sci.* **92**: 10614–10618.

R Core Team. 2020. *R: A language and environment for statistical computing*. R Foundation
for Statistical Computing, Vienna, Austria.

Rowe, L. & Houle, D. 1996. The lek paradox and the capture of genetic variance by condition
dependent traits. *Proc. R. Soc. Lond. B Biol. Sci.* **263**: 1415–1421.

Schielzeth, H. & Forstmeier, W. 2009. Conclusions beyond support: overconfident estimates
in mixed models. *Behav. Ecol.* **20**: 416–420.

Schneider, C.A., Rasband, W.S. & Eliceiri, K.W. 2012. NIH Image to ImageJ: 25 years of image
analysis. *Nat. Methods* **9**: 671–675.

Simmons, L.W., Lüpold, S. & Fitzpatrick, J.L. 2017. Evolutionary trade-off between secondary
sexual traits and ejaculates. *Trends Ecol. Evol.* **32**: 964–976.

Snook, R.R., Briistle, L. & Slate, J. 2009. A test and review of the role of effective population
size on experimental sexual selection patterns. *Evolution* **63**: 1923–1933.

Snook, R.R., Robertson, A., Crudgington, H.S. & Ritchie, M.G. 2005. Experimental
manipulation of sexual selection and the evolution of courtship song in *Drosophila*
pseudoobscura. *Behav. Genet.* **35**: 245–255.

Stångberg, J., Immonen, E., Moreno, P.P. & Bolund, E. 2020. Experimentally induced
intrasexual mating competition and sex-specific evolution in female and male
nematodes. *J. Evol. Biol.* **33**: 1677–1688.

Veltsos, P., Fang, Y., Cossins, A.R., Snook, R.R. & Ritchie, M.G. 2017. Mating system
manipulation and the evolution of sex-biased gene expression in *Drosophila*. *Nat.*
Commun. **8**: 2072.

Videlier, M., Rundle, H.D. & Careau, V. 2019. Sex-specific among-individual covariation in
locomotor activity and resting metabolic rate in *Drosophila melanogaster*. *Am. Nat.*
194: E164–E176.

Videlier, M., Rundle, H.D. & Careau, V. 2021. Sex-specific genetic (co)variances of standard
metabolic rate, body mass and locomotor activity in Drosophila melanogaster. *J.*
Evol. Biol. **n/a**.

Watson, P.J., Stallmann, R.R. & Arnqvist, G. 1998. Sexual conflict and the energetic costs of
mating and mate choice in water striders. *Am. Nat.* **151**: 46–58.

Wiberg, R.A.W., Veltsos, P., Snook, R.R. & Ritchie, M.G. 2021. Experimental evolution
supports signatures of sexual selection in genomic divergence. *Evol. Lett.* **n/a**.

Wigglesworth, V.B. 1949. The utilization of reserve substances in Drosophila during flight. *J.*
Exp. Biol. **26**: 150–163.

Wittman, T.N., Robinson, C.D., McGlothlin, J.W. & Cox, R.M. 2021. Hormonal pleiotropy
structures genetic covariance. *Evol. Lett.* **n/a**.

Zera, A.J. & Harshman, L.G. 2001. The physiology of life history trade-offs in animals. *Annu.*
Rev. Ecol. Syst. **32**: 95–126.