Shining a light on Laurentian Great Lakes cisco (*Coregonus artedi*): how ice coverage may

impact embryonic development

Taylor R. Stewart[a,b*], Mark R. Vinson[c], and Jason D. Stockwell[b]

[a] Department of Biology, University of Vermont, 109 Carrigen Drive, Burlington, VT 05405,

USA

[b] Rubenstein Ecosystem Science Laboratory, University of Vermont, 3 College Street,

Burlington, VT 05401, USA

[c] U.S. Geological Survey, Lake Superior Biological Station, 2800 Lake Shore Drive East,

Ashland, WI 54806, USA

*Corresponding Author: Taylor R. Stewart, Department of Biology, Rubenstein Ecosystem

Science Laboratory, University of Vermont, 3 College St, Burlington, VT 05401, USA.

Email: taylor.stewart@uvm.edu

Abstract:

Changes in winter conditions, such as decreased ice coverage and duration, have been observed in the Laurentian Great Lakes over the past 20+ years and hypothetically linked to low *Coregonus* spp. survival to age-1. Most cisco (*Coregonus artedi*) populations are autumn spawners whose embryos incubate under ice throughout the winter. The quantity and quality of light during winter is regulated by ice and snow coverage, and light has been shown to affect embryo survival and development in some teleosts. We experimentally evaluated how cisco embryos from lakes Superior and Ontario responded to three light treatments that represented day-light intensity under 0-10, 40-60, and 90-100% ice coverage. Embryonic response measures included two developmental factors (embryo survival and incubation period) and two morphological traits (length-at-hatch and yolk-sac volume). Embryo survival was highest at the medium light treatment and decreased at high and low treatments for both populations suggesting cisco may be adapted to withstand some light exposure from inter-annual variability in ice coverage. Light intensity had no overall effect on length of incubation. Length-at-hatch decreased with increasing light in Lake Superior, but had no effect in Lake Ontario. Yolk-sac volume was positively correlated with increasing light in Lake Superior and negatively correlated in Lake Ontario. Contrasting responses in embryo development between lakes suggests differences in populations' flexibility to light. These results provide a step towards better understanding the recent high variability observed in coregonine recruitment and may help predict what the future of this species may look like under current climate trends.

Keywords: Coregonus; cisco; climate change; embryo incubation; ice coverage; light intensity

Introduction:

Freshwater whitefishes, Salmonidae Coregoninae (hereafter coregonines) play important economic (Ebener et al., 2008) and ecological (Lynch et al., 2010; Nyberg et al., 2001; Stockwell et al., 2014) roles throughout the northern hemisphere, but populations have declined over the past century (Eshenroder et al., 2016). Historical coregonine declines were attributed to overfishing, invasive species, habitat alterations, and competition (Anneville et al., 2009; Rosinski et al., 2020; Stockwell et al., 2009). More recently, coregonine populations worldwide have experienced declines due to highly variable and low survival to age-1 (Lepak et al., 2017; Nyberg et al., 2001; Parks and Rypel, 2018) which have been associated with climate-induced changes in early-life stage environments (Nyberg et al., 2001). However, an underlying mechanism between changing lake environments and coregonine year-class strength has yet to be established.

Year-class strength in most fish species, including coregonines, is thought to be established prior to the end of the first season of growth (Cushing, 1990; Hjort, 1914; Karjalainen et al., 2015). Most coregonines are autumn spawners whose embryos incubate under ice throughout the winter (Karjalainen et al., 2000; Stockwell et al., 2009). Embryos are static, which leaves them vulnerable to predation (Stockwell et al., 2014) and unable to evade detrimental changes in winter environmental conditions (Pepin, 1991). Changes in winter conditions, such as decreased ice coverage and duration, that have been observed over the past 20+ years (Austin and Colman, 2007; O'Reilly et al., 2015; Sharma et al., 2019), could alter developmental rates, embryo survival, and time of hatching (Karjalainen et al., 2015). Potential mechanisms by which ice coverage influences coregonine embryonic development include reduction of physical wave action (Austin and Colman, 2007; Nguyen et al., 2017; Walter et al., 2006; Wang et al., 2010), more stable winter and spring water temperatures (Magnuson et al., 1997; Winslow et al., 2017), and the amount of sunlight reaching the lake bottom (Bolsenga and Vanderploeg, 1992; Hampton et al., 2015).

Sunlight (i.e., photoperiod) is the most consistent abiotic factor in nature (Ruchin, 2020) and can regulate fish development phenology, behavior, and physiology (Ruchin, 2007; Villamizar et al., 2011). The length of photoperiods characterize circadian rhythms and ensure that biological

processes are synchronized with the environment (Gaston et al., 2013; Marchesan et al., 2005;

Ruchin, 2020). In seasonally ice-covered lakes, winter lake light levels are regulated by ice

coverage and snow depth (Bolsenga and Vanderploeg, 1992; Hampton et al., 2015). Ice can

reduce light transmittance to 62% under clear ice, and to ≤ 10% under snow covered ice

(Bolsenga and Vanderploeg, 1992).

Salmonid embryos incubated under elevated light levels had higher mortality and deformity

rates, slower formation of cartilaginous skeletal elements, decreased time to hatching, and

smaller size-at-age; with accelerated development after organogenesis (Chernyaev, 2007; Eisler,

1961, 1958; Kwain, 1975; MacCrimmon and Kwain, 1969). However, other teleosts (e.g., turbot

Scophthalmus maximus, Atlantic halibut *Hippoglossus hippoglossus*, brown-marbled grouper

Epinephelus fuscoguttatus) exhibit opposite responses, or no response, to light illumination

(Iglesias et al., 1995; Mangor☐Jensen and Waiwood, 1995; Ruchin, 2020; Seth et al., 2014). To

our knowledge, no previous work has examined the effects of light on coregonine embryos from

North America.

We experimentally evaluated how cisco (*Coregonus artedi*) embryos responded to different

photoperiod intensities, as a proxy for different ice coverages. We hypothesized that exposure to

elevated light intensity (a proxy for low ice coverage) decreases embryo survival and accelerates

embryogenesis, resulting in earlier hatching, larger yolk-sacs, and shorter lengths-at-hatch. Our

objective was to identify the extent to which light influences cisco embryo survival, incubation

duration, and length and yolk-sac volume at hatching. If our hypothesis is supported, we would

expect populations adapted to lower light levels (high ice coverage) to experience a greater

magnitude of change as light intensity increases.

Methods:

Ethics

All work described here was approved for ethical animal care under University of Vermont's

Institutional Animal Care and Use Committee (Protocol # PROTO202000021).

Study Species and Locations

Mature cisco were collected from the Apostle Islands, Lake Superior (46.85°, -90.55°) and

Chaumont Bay, Lake Ontario (44.05°, -76.20°) in December 2019. Lake Superior cisco were

collected at an open lake location at depths between 15 and 50 m. Lake Ontario cisco were

collected in a shallow, protected bay on rocky shoals at depths between 2 to 5 m. Historical

(1973-2020) ice conditions over the sampled spawning locations varied between lakes with the

shallower, more protected Lake Ontario spawning site having more consistent ice coverage

between January and March than the deeper, open location in Lake Superior (Figure 1). The

different spawning habitats provide a contrast in light levels that coregonine embryos from each

population would naturally experience because maximum light availability decreases with depth

(Fleming-Lehtinen and Laamanen, 2012; Preisendorfer, 1986; Ramus et al., 1976; Secchi, 1864)

and winter light availability is further restricted by ice and snow conditions.

Crossing Design and Fertilization

The design is fully described in Stewart et al., (2021). Briefly, gametes were stripped from 12

females and 16 males from each lake and artificially fertilized to create 48 families from each

lake. Reconstructed fresh water was used during fertilizations and rearing (OECD ISO

6341:2012) to standardize the chemical properties of the water used between lakes. Embryos

were transported to the University of Vermont in coolers by shipping overnight for Lake

Superior samples and driven the same-day for Lake Ontario samples. A temperature logger

recorded air temperature inside the cooler during transport (Lake Superior: mean (SD) = 2.80°C

(0.21); Lake Ontario: mean (SD) = 3.28°C (0.37)). Total length, mass, and egg diameter were

collected from the spawned adults. Fertilization success was determined by assessing 10

haphazardly selected embryos under microscopy (Oberlercher and Wanzenböck, 2016). If

fertilization was low (<30%), the family was removed from the experiment.

Rearing Conditions

Embryos were individually distributed into 24-well cell culture microplates and incubated in 2

ml of reconstructed fresh water (Stewart et al., 2021). A total of 36 embryos were used for each

Lake Ontario and Lake Superior cisco family. Families were randomly distributed across three

microplates (i.e., 12 eggs per family per microplate resulting in two families per 24-well

microplate).

Microplates from each population were incubated under three light treatments (0.6-6.2 μmol m^{-2}

s^{-1}) that represented day-light intensity under 0-10 (low), 40-60 (medium), and 90-100 % (high)

ice coverage (Table 1), and followed mean weekly photoperiods with gradual sunrise and sunset

transitions. Light intensities for each treatment were chosen to mimic *in situ* winter, lakebed light

measurements that were previously recorded with a photometer (JFE Advantech Co., Ltd.

DEFI2-L) from Lake Superior (46.97°, -90.99°) at 10 m of water in 2016-17. Remote-sensing ice

data (U.S. National Ice Center; usicecenter.gov) were used to quantify the daily percentage of ice

coverage above the light sensor (Figure 2). Embryos were incubated at a constant target water

temperature of 4.0°C in a climate-controlled chamber (Conviron® E8; Table 2). Forced airflow

was used in the climate-controlled chamber to ensure equal air circulation around the microplates

and opaque, plastic sheeting was used to separate light treatments. Microplates were covered

with transparent lids to minimize evaporation and rotated (i.e., orientation and position within the

incubator) weekly. Water temperature and light intensity were recorded hourly with loggers

(HOBO® Water Temperature Pro v2 and JFE Advantech Co., Ltd. DEFI2-L) and daily mean

values calculated (Table 1). During the hatch period, microplates were checked on a three-day

cycle for newly hatched embryos. All hatched embryos were photographed ventrally (Nikon®

D5600 and Nikon® AF-S DX 18-55mm lens) and then immediately preserved in 95% ethanol.

Egg size at fertilization, total length-at-hatch, and post-hatching yolk-sac axes were measured

from photographed images using Olympus® LCmicro.

Developmental and Morphological Traits

Embryo survival was estimated as the percent of embryos surviving between eye-up and post-

hatch stages. Incubation period was assessed with two variables: the number of days from

fertilization to hatching (days post-fertilization; DPF) and the sum of the degree-days

(accumulated degree-days; ADD). Total length-at-hatch (LAH; mm) and yolk-sac volume (YSV;

mm^3) were measured from five individuals per family at, or as close as possible to, 50% hatching

for each family. Yolk-sac volume was calculated assuming the shape of an ellipse (Blaxter,

1963):

$$YSV = \frac{\pi}{6}ab^2$$

where a = length of the yolk sac (mm) and b = height of the yolk sac (mm).

Statistical Analyses

Embryo survival was analyzed as a binomial response variable, with incubation period, length-

at-hatch, and yolk-sac volume at hatching as continuous response variables. Because embryos

were raised independently, the replication unit in the statistical models is the individual embryo

and the design was unbalanced due to different levels of embryo mortality. All non-proportional

data were visually checked for approximate normality using histograms and Q-Q plots. A cubic

transformation was applied to LAH and a cubic root transformation was applied to DPF, ADD,

and YSV to normalize the distributions. Embryo survival was analyzed with binomial

generalized linear mixed-effects models, and variables with distributions not strongly deviating

from normal (i.e., incubation period, LAH, and YSV) were analyzed with restricted maximum

likelihood linear mixed-effects models with the *lme4* package v.1.1-26 (Bates et al., 2015).

Population and incubation light treatment were included as fixed effects and female, male,

female x male, and fertilization block as random effects. All traits and possible interactions were

examined with backward, stepwise effect-selection and the maximal model for each trait selected

using the *buildmer* package v.1.7.1 (Voeten, 2020). The significance for population, species,

incubation temperature, interaction effects, and any random-effects selected were determined

using a likelihood ratio test between the maximal model and reduced models with the model

effect of interest removed.

To enable population comparisons, the response to temperature for each trait was standardized to

what we assumed was the optimal light treatment - the low light treatment (Table 1). For each

trait, the within-family mean was calculated for each light treatment and the percent change from

the optimal light intensity estimated. Standard error was calculated as the among-family

variation in percent change.

All analyses were performed in R version 4.0.4 (R Core Team, 2021).

Results:

Spawning Adult and Egg Measurements

Lake Superior spawning adults ranged from 326-503 mm (total length mean (SD) = 412.5 (40.8)

mm) and 298.9-970.0 g (fresh mass mean (SD) = 589.1 (171.4) g), and were larger in total length

and fresh mass than Lake Ontario adults which ranged from 321-425 mm (mean (SD) = 372.5

(25.3) mm) and 280.5-795.8 g (mean (SD) = 496.6 (126.4) g). Egg diameter was larger in Lake

Ontario (mean (SD) = 2.30 (0.08) mm) than Lake Superior (mean (SD) = 2.14 (0.12) mm).

Developmental and Morphological Traits

Incubation period (both DPF and ADD) and YSV had significant interaction effects between

population and light treatments (maximum $P = 0.008$; Table 3). The interaction effects precluded

any interpretation of main effects for incubation period and YSV, but did suggest contrasting

norms of reaction between populations. Below we describe the interaction effects for incubation

period and YSV, and the population main effects and light treatment pairwise comparisons for

embryo survival and LAH. All random effects (i.e., female, male, and female x male) were

significant (maximum $P = 0.009$) except female for embryo survival, male for embryo survival

and YSV, and female x male for embryo survival and LAH (Table 3). All statistical model

results can be found in Table 3.

Embryo Survival

Embryo survival was highest for both populations at the medium light treatment, but lowest at

the low light treatment for Lake Ontario and at the high light treatment for Lake Superior (Figure

3). Light and population main effects were significant. Only Lake Ontario pairwise light

treatment comparisons with the low light treatment were significant (Low - Medium $P < 0.001$;

Low - High $P < 0.001$). All pairwise light treatment comparisons for Lake Superior were not

significant (minimum $P = 0.089$). Embryo survival was higher for Lake Ontario at the high

(98.4%) and medium (99.6%) light treatments than for Lake Superior (85.3 and 89.3%,

respectively) but there was no difference between populations at the low light treatment (0.9%;

Figure 3).

Incubation Period

The number of days between fertilization and hatching was highest for Lake Ontario at the low

light treatment (115.47 days) and for Lake Superior at the high light treatment (101.22 days;

Figure 3). Lake Ontario cisco had a decrease in DPF from the low light to the high light

treatments (-0.7%), while Lake Superior had an increase from the low light to the high light

treatments (1.9%; Figure 3). Incubation period (DPF) was longer for Lake Ontario than Lake

Superior across all light treatments (mean (SD) difference = 13.9 (0.8) days).

The effect of light depended on population because the difference in ADD between populations

was less pronounced at the high light treatment (difference = 60.8 ADD), while ADD was higher

for Lake Ontario at the low and medium light treatments (497.7 and 485.9 ADD, respectively)

than Lake Superior (427.5 and 420.8 ADD, respectively; Figure 3). Lake Ontario ADD had a

negative response from the low to high light treatments (-2.5%), while ADD for Lake Superior

did not change from the low to high light treatments (0.05%; Figure 3).

Length-at-Hatch

Light was not a component returned in the stepwise-selected model, but the population main

effect between Lake Ontario and Lake Superior was significant ($P < 0.001$; Table 3). Lake

Ontario had a higher LAH than Lake Superior across all light treatments (Figure 4). Length-at-

hatch decreased with increasing light by 3.2 and 0.2% in Lake Superior and Lake Ontario

populations, respectively (Figure 4).

Yolk-sac Volume

Yolk-sac volume had a different response to light intensity between populations (Figure 4). The

effect of light depended on population because the difference in YSV between populations was

less pronounced at the low light treatment (0.22 mm^3), while YSV was lower for Lake Ontario at

the high and medium light treatments (0.35 and 0.37 mm^3, respectively) than Lake Superior

(0.67 and 0.63 mm^3, respectively; Figure 4). YSV increased from the low to high light treatments

(15.3%) in Lake Superior and decreased from the low to high light treatments (-5.5%) in Lake

Ontario (Figure 4).

Discussion:

Developmental and morphological traits for Lake Superior and Lake Ontario cisco populations demonstrated similar and contrasting reaction norms to incubation light intensity. First, we found different responses to light intensity in embryo survival. Second, increasing light intensity had minimal impact on incubation periods (DPF and ADD) for both populations. Lastly, each population responded differently to light for LAH and YSV. Consequently, cisco from lakes Superior and Ontario are likely to have different responses to changes in ice coverage and subsequent light conditions.

Our hypothesis that embryo survival would be highest at the lowest light treatment was not supported. Embryo survival was highest for both populations at the medium light levels, suggesting that populations may be adapted to withstand some light exposure from high inter-annual variability in ice coverage. This result was contradictory to that observed in two Pacific salmonid species (chinook salmon *Oncorhynchus tshawytscha* and rainbow trout *Oncorhynchus mykiss*) for which embryo survival was highest at the lowest light exposures evaluated (0.04 μmol m^{-2} s^{-1}; Eisler, 1961, 1958; Kwain, 1975). Lake Ontario cisco had a sharper decrease in survival than Lake Superior cisco at the low light treatment. The difference was surprising because average historical ice coverage over the Lake Ontario spawning location is higher than the Lake Superior spawning location, and thus low light conditions are more likely to occur for Lake Ontario cisco embryos. However, the Lake Ontario cisco spawning location is shallow (< 5 m) and would have high light intensity with little or no ice coverage. Higher variability in winter illuminance may allow the population of Lake Ontario cisco sampled to have greater flexibility to higher light conditions than deeper spawning cisco sampled from Lake Superior.

Our hypothesis that elevated light intensity would accelerate embryogenesis was not supported. The greatest difference in incubation periods was between populations, and was likely due to differences in embryo size, as larger embryos (i.e., Lake Ontario cisco) require more time to develop (Hodson and Blunt, 1986; Kamler, 2008). Previous studies of other salmonid species (European whitefish *Coregonus lavaretus*, chinook salmon, rainbow trout) found increasing light intensity decreased the length of incubation (Chernyaev, 2007; Eisler, 1958; Kwain, 1975).

In contrast to incubation period, LAH and YSV responded to the light treatment and matched our

hypotheses, but responses differed between populations. Lake Ontario cisco exhibited minimal

change in LAH as light increased, but YSV decreased, suggesting that light intensity increased

the metabolic demand of embryos and diverted energy away from somatic growth. In

comparison, Lake Superior cisco showed a trade-off between LAH and YSV. A negative

relationship between LAH and YSV is a common finding in fish temperature incubation studies

(Blaxter, 1991; Karjalainen et al., 2015; Stewart et al., 2021), but the relationship is usually

found alongside a change in incubation period as basal metabolic demand consumes yolk as a

function of the length of incubation. We found that light influenced incubation periods similarly

among light treatments; therefore, the trade-off between LAH and YSV in Lake Superior cisco

suggests decreased rates of yolk conversion into somatic tissue occurred as light intensity

increased. This suggests future decreases in ice coverage and subsequent increases in embryonic

light exposure may result in smaller, less-robust larvae, which may explain the lack of observed

survival of Lake Superior cisco and other coregonines to age-1 over the past 20 years (Lepak et

al., 2017; Stockwell et al., 2009). The reasons underlying differences between Lakes Ontario and

Superior remain unknown. However, the contrasting responses in LAH and YSV between

populations suggests that embryogenesis for each population has different levels of plasticity to

light.

Embryo development is sensitive to environmental conditions, which can greatly influence life-

history trajectories, performances, and reproductive success (Colby and Brooke, 1970;

Karjalainen et al., 2016; Luczynski, 1991). We did not quantify developmental stages, except eye

pigmentation, so specific life-stage developmental rates are unknown. Changes in the frequency

of light (i.e., periodicity) can have adverse effects on fish embryos after yolk plug closure and

first vertebrate trunk segment formation (Abdel-Rahim et al., 2019; Chernyaev, 2007, 1993;

Ruchin, 2020). Fluctuating light cycles (e.g., 6:6h light:dark) and constant light (e.g., 24h light)

accelerated the rate of embryonic development compared to 'normal' photoperiods (e.g., 12:12h

light:dark; Chernyaev, 2007, 1993; John and Hasler, 1956; Ruchin, 2020). Photoperiod

disruptions can inhibit the pineal organ and melatonin synthesis, which is critical to regulate and

synchronize diurnal and seasonal biological rhythms (Delgado et al., 1987; Ekstrzöm and Meissl,

1997; Falcón et al., 2010; Roberts, 1978). The role photoperiod and the endocrine system plays

in embryo development and phenology remains unknown for coregonines. Further studies that

examine the impact of changing light intensities and photoperiods throughout incubations (e.g.,

decreased or no light during winter from ice coverage and increased light intensity and

periodicity during spring ice-out) will help determine the fine-scale influence light and

photoperiod may have on specific development stages (i.e., hatching), hormone regulation, and

organ, tissue, and skeletal formation.

Sunlight intensity, albedo, and attenuation are strongly influenced by the angle of the sunlight,

which is determined by season and latitude (Forsythe et al., 1995; Goldberg and Klein, 1977).

Latitude and sun angle are negatively correlated and this negative relationship is strongest at the

winter solstice in the northern hemisphere (Goldberg and Klein, 1977; Wielgolaski and Inouye,

2003). Lake Ontario is at a lower latitude and thus experiences a higher sun angle than Lake

Superior, which results in a more intense and longer period of daylight. Our light treatments

were calculated from light sensors deployed only in Lake Superior; thus, the experimental light

intensity treatments for Lake Ontario cisco may not have captured an accurate light environment

representation. Under-ice light data from more lakes, depths, and habitats would add to our

understanding of cisco embryo light environments and improve the authenticity of experimental

treatments. Additionally, comparing populations from high latitude lakes which experience

decreased winter sunlight would provide an additional contrast for local adaptation and plasticity

across geographic regions.

Turbidity also contributes to light attenuation, and spring ice-out and river discharge can

drastically increase the presence of suspended particulates and light absorption (Shao et al.,

2019). Hydrological responses to climate change indicate earlier and protracted winter/spring

runoff and higher runoff volume (Blahušiaková et al., 2020; Cutforth et al., 1999; Shen et al.,

2018). Seasonal runoff, including snowmelt pulses, often drive high nutrient loads and primary

productivity in temperate lakes (Isles et al., 2017; Rosenberg and Schroth, 2017). Runoff

entering ice-covered lakes is expected to suspend near the ice surface, rather than mixing into the

water column (Cortés et al., 2017; Yang et al., 2020), and therefore, have implications on when

nutrients are used by primary producers and the onset of spring plankton blooms (Sommer et al.,

2012). If the timing of spring plankton blooms changes as a result of earlier and protracted

winter/spring nutrient loads, the potential mis-match between interacting species may produce
bottom-up consequences (Rogers et al., 2020). Our results showed changing light intensities did
not affect cisco hatch dates; thus, the ability of cisco to match optimal spring nursery conditions
may be weakened if coregonines do not respond to changing ice conditions similarly to the
plankton community (Cushing, 1990; Myers et al., 2015). The proximity of spawning and
nursery grounds to shoreline and river outlets would likely impact the synchrony between
coregonine larvae and planktonic prey.
353
Many fish species are iteroparous and in some species individuals repeatedly use the same
spawning location (Marsden et al., 1995; Skjæraasen et al., 2011; Thorrold et al., 2001). The
question of what constrains the choice of a spawning location cannot be separated from the
question of what constrains early-life development and survival (Ciannelli et al., 2015; Iles and
Sinclair, 1982; Petitgas et al., 2012; Sinclair and Iles, 1989). Embryo survival is largely
determined by incubation habitat (e.g., water temperature, light exposure, oxygen availability,
protection from predators), thus, selective pressure is focused on 'correct' and adaptive choices of
spawning sites by the parents. The amount of spawning plasticity (e.g., spawning site selection,
fidelity to spawning sites, spawning time) among populations could serve as an indicator for the
level of evolutionary constraints for offspring (Ciannelli et al., 2015). For example, Atlantic
herring (*Clupea harengus*) exhibit a wide range of reproductive strategies across diverse
geographical locations, but have limited spawning site plasticity because embryo survival is
dependent on substrate and vegetation (Petitgas et al., 2012). Coregonines are considered to be
behaviorally and developmentally plastic and do not appear to be constrained by a spawning
habitat type (Karjalainen et al., 2015; Muir et al., 2013; Paufve, 2019); however, our
understanding of coregonine reproductive behavior and spawning-site selection is limited. The
selection of deeper or shallower spawning locations would provide a gradient in environment
conditions (e.g., light, temperature) depending on population-specific habitat requirements, and
both suitable nearshore and offshore spawning habitats are likely to be present in each sampled
lake (Goodyear, 1982; Paufve, 2019).
374
375   The existence of varying trait responses between populations raises questions concerning causal
376   mechanisms. Genomic studies can aid our understanding by determining what functional

377 pathways could be up or down-regulated due to light energy. Any potential changes in metabolic
378 or catabolic genes from light will enhance trait analyses and allow further partitioning of the
379 effects of light from other energy demanding environmental variables (e.g., temperature).
380
Conclusion:
Given the extensive degree of developmental plasticity in coregonines, propagation has been
proposed as a practical way to reintroduce native species from lakes with extirpated or reduced
population levels (Bronte et al., 2017; Zimmerman and Krueger, 2009). A key uncertainty to
maximizing restoration efforts is whether managers should prioritize survival in propagation
methods by mimicking natural environmental conditions (Bronte et al., 2017). Our study
highlights the potential role of winter light conditions, the influence of light intensity on cisco
embryo development, and the impact changing ice regimes may have on cisco survival and
recruitment in the wild. We did not identify a consistent directional reaction between and within
the two cisco populations to increasing light, and light is likely to have a differential effect on a
number of physiological and biochemical processes. Large-scale, cross-lake propagation and
reintroduction efforts are likely to be limited by the ability to match cisco phenotypes and
optimal incubation conditions. Our results provide a step towards better understanding the recent
high variability observed in coregonine recruitment and may help predict what the future of this
species may look like under current climate trends.
396

Acknowledgments:
We thank the staff at the Wisconsin Department of Natural Resources Bayfield Fisheries Field
Station, United States Geological Survey (USGS) Tunison Laboratory of Aquatic Science, and
New York State Department of Environmental Conservation Cape Vincent Fisheries Station for
conducting field collections of spawning adults. The staff at Apostle Islands National Lakeshore
(U.S. National Park Service) conducted sensor deployment and retrieval. We also thank Rachel
Taylor, Dan Yule, and Caroline Rosinski for help with fertilizations and experiment
maintenance. [NAME] provided the USGS solicited review that strengthened the manuscript, as
did anonymous peer reviewers and Stockwell and Marsden lab members. This work was
supported by the USGS [grant number G17AC00042]. Any use of trade, product, or firm names
is for descriptive purposes only and does not imply endorsement by the U.S. Government.

References:

Abdel-Rahim, M.M., Lotfy, A.M., Aly, H.A., Sallam, G.R., Toutou, M.M., 2019. Effects of light source, photoperiod, and intensity on technical and economic performance of meagre, Argyrosomus regius, on intensive land-based farms. Aquac. Aquarium, Conserv. Legis. 12, 1531–1545.

Anneville, O., Souissi, S., Molinero, J.C., Gerdeaux, D., 2009. Influences of human activity and climate on the stock☐recruitment dynamics of whitefish, Coregonus lavaretus, in Lake Geneva. Fish. Manag. Ecol. 16, 492–500. https://doi.org/10.1111/j.1365-2400.2009.00703.x

Austin, J.A., Colman, S.M., 2007. Lake Superior summer water temperatures are increasing more rapidly than regional temperatures: A positive ice-albedo feedback. Geophys. Res. Lett. 34, 1–5. https://doi.org/10.1029/2006GL029021

Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models Using lme4. J. Stat. Softw. 67, 1–48.

Blahušiaková, A., Matoušková, M., Jenicek, M., Ledvinka, O., Kliment, Z., Podolinská, J., Snopková, Z., 2020. Snow and climate trends and their impact on seasonal runoff and hydrological drought types in selected mountain catchments in Central Europe. Hydrol. Sci. J. 65, 2083–2096. https://doi.org/10.1080/02626667.2020.1784900

Blaxter, J.H.S., 1991. The effect of temperature on larval fishes. Netherlands J. Zool. 42, 336–357. https://doi.org/10.1163/156854291X00379

Blaxter, J.H.S., 1963. The influence of egg size on herring larvae (Clupea harengus L). J. du Cons. / Cons. Perm. Int. pour l'Exploration la Mer 28, 211–240. https://doi.org/10.1093/icesjms/28.2.211

Bolsenga, S.J., Vanderploeg, H.A., 1992. Estimating photosynthetically available radiation into open and ice-covered freshwater lakes from surface characteristics; a high transmittance case study. Hydrobiologia 243–244, 95–104. https://doi.org/10.1007/BF00007024

Bronte, C.R., Bunnell, D.B., David, S.R., Gordon, R., Gorsky, D., Millard, M.J., Read, J., Stein, R.A., Vaccaro, L., 2017. Report from the workshop on coregonine restoration science. US Geological Survey. https://doi.org/10.3133/ofr20171081

Chernyaev, Z.A., 2007. Factors and possible mechanisms causing changes in the rate of embryonic development of bony fish (with reference to Coregonidae). J. Ichthyol. 47, 494–503. https://doi.org/10.1134/S003294520707003X

Chernyaev, Z.A., 1993. The Impact of Light Factor on the Embryonic Development of
Coregonids. Izv. Akad. Nauk. Ser. Biol 64–73.

Ciannelli, L., Bailey, K., Olsen, E.M., 2015. Evolutionary and ecological constraints of fish
spawning habitats. ICES J. Mar. Sci. 72, 285–296. https://doi.org/10.1093/icesjms/fsu145

Colby, P.J., Brooke, L.T., 1970. Survival and development of lake herring (Coregonus artedii)
eggs at various incubation temperatures. Biol. Coregonid Fishes 417–428.

Cortés, A., MacIntyre, S., Sadro, S., 2017. Flowpath and retention of snowmelt in an
ice⬜covered arctic lake. Limnol. Oceanogr. 62, 2023–2044.
https://doi.org/10.1002/lno.10549

Cushing, D.H., 1990. Plankton production and year-class strength in fish populations: An update
of the match/mismatch hypothesis. Adv. Mar. Biol. 26, 249–293.
https://doi.org/10.1016/S0065-2881(08)60202-3

Cutforth, H.W., McConkey, B.G., Woodvine, R.J., Smith, D.G., Jefferson, P.G., Akinremi, O.O.,
1999. Climate change in the semiarid prairie of southwestern Saskatchewan: Late winter–
early spring. Can. J. Plant Sci. 79, 343–350. https://doi.org/10.4141/P98-137

Delgado, M.J., Gutiérrez, P., Alonso-Bedate, M., 1987. Melatonin and photoperiod alter growth
and larval development in Xenopus laevis tadpoles. Comp. Biochem. Physiol. Part A
Physiol. 86, 417–421. https://doi.org/10.1016/0300-9629(87)90517-2

Ebener, M.P., Stockwell, J.D., Yule, D.L., Gorman, O.T., Hrabik, T.R., Kinnunen, R.E., Mattes,
W.P., Oyadomari, J.K., Schreiner, D.R., Geving, S., Scribner, K., Schram, S.T., Seider,
M.J., Sitar, S.P., 2008. Status of cisco (Coregonus artedi) in Lake Superior during 1970-
2006 and management and research considerations. Ann Arbor, Michigan Gt. Lakes Fish.
Comm. Lake Super. Tech. Rep. 1, 126.

Eisler, R., 1961. Effects of visible radiation on salmonoid embryos and larvae. Growth 25, 281–
346.

Eisler, R., 1958. Some effects of artificial light on salmon eggs and larvae. Trans. Am. Fish. Soc.
87, 151–162. https://doi.org/10.1577/1548-8659(1957)87[151:SEOALO]2.0.CO;2

Ekstrzöm, P., Meissl, H., 1997. The pineal organ of teleost fishes. Rev. Fish Biol. Fish. 7, 199–
284. https://doi.org/10.1023/A:1018483627058

Eshenroder, R.L., Vecsei, P., Gorman, O.T., Yule, D.L., Pratt, T.C., Mandrak, N.E., Bunnell,
D.B., Muir, A.M., 2016. Ciscoes (Coregonus, subgenus Leucichthys) of the Laurentian

Great Lakes and Lake Nipigon, Canadian Journal of Fisheries and Aquatic Sciences. Great

Lakes Fishery Commission.

Falcón, J., Migaud, H., Munoz-Cueto, J.A., Carrillo, M., 2010. Current knowledge on the

melatonin system in teleost fish. Gen. Comp. Endocrinol. 165, 469–482.

https://doi.org/10.1016/j.ygcen.2009.04.026

Fleming-Lehtinen, V., Laamanen, M., 2012. Long-term changes in Secchi depth and the role of

phytoplankton in explaining light attenuation in the Baltic Sea. Estuar. Coast. Shelf Sci.

102, 1–10. https://doi.org/10.1016/j.ecss.2012.02.015

Forsythe, W.C., Rykiel Jr, E.J., Stahl, R.S., Wu, H., Schoolfield, R.M., 1995. A model

comparison for daylength as a function of latitude and day of year. Ecol. Modell. 80, 87–95.

https://doi.org/10.1016/0304-3800(94)00034-F

Gaston, K.J., Bennie, J., Davies, T.W., Hopkins, J., 2013. The ecological impacts of nighttime

light pollution: a mechanistic appraisal. Biol. Rev. 88, 912–927.

https://doi.org/10.1111/brv.12036

Goldberg, B., Klein, W.H., 1977. Variations in the spectral distribution of daylight at various

geographical locations on the earth's surface. Sol. Energy 19, 3–13.

https://doi.org/10.1016/0038-092X(77)90083-4

Goodyear, C.D., 1982. Atlas of the spawning and nursery areas of Great Lake fishes. US Fish

and Wildlife Service.

Hampton, S.E., Moore, M. V, Ozersky, T., Stanley, E.H., Polashenski, C.M., Galloway, A.W.E.,

2015. Heating up a cold subject: prospects for under-ice plankton research in lakes. J.

Plankton Res. 37, 277–284. https://doi.org/10.1093/plankt/fbv002

Hjort, J., 1914. Fluctuations in the great fisheries of Northern Europe, in: Rapports et Procés-

Verbaux. ICES, pp. 1–228.

Hodson, P. V, Blunt, B.R., 1986. The effect of time from hatch on the yolk conversion efficiency

of rainbow trout, Salmo gairdneri. J. Fish Biol. 29, 37–46. https://doi.org/10.1111/j.1095-

8649.1986.tb04924.x

Iglesias, J., Rodríguez-Ojea, G., Peleteiro, J.B., 1995. Effect of light and temperature on the

development of turbot eggs (Scophthalmus maximus L.), in: ICES Marine Science

Symposia. Copenhagen, Denmark: International Council for the Exploration of the Sea,

1991-, pp. 40–44.

Iles, T.D., Sinclair, M., 1982. Atlantic herring: stock discreteness and abundance. Science (80-.).

215, 627–633. https://doi.org/10.1126/science.215.4533.627

Isles, P.D.F., Xu, Y., Stockwell, J.D., Schroth, A.W., 2017. Climate-driven changes in energy

and mass inputs systematically alter nutrient concentration and stoichiometry in deep and

shallow regions of Lake Champlain. Biogeochemistry 133, 201–217.

https://doi.org/10.1007/s10533-017-0327-8

John, K.R., Hasler, A.D., 1956. Observations on Some Factors Affecting the Hatching of Eggs

and the in Lake Mendota , Wisconsin. Limnol. Oceanogr. 1, 176–194.

https://doi.org/10.4319/lo.1956.1.3.0176

Kamler, E., 2008. Resource allocation in yolk-feeding fish. Rev. Fish Biol. Fish. 18, 143–200.

https://doi.org/10.1007/s11160-007-9070-x

Karjalainen, J., Auvinen, H., Helminen, H., Marjomäki, T.J., Niva, T., Sarvala, J., Viljanen, M.,

2000. Unpredictability of fish recruitment - interannual variation in YOY abundance. J. Fish

Biol. https://doi.org/DOI 10.1006/jfbi.1999.1206

Karjalainen, J., Jokinen, L., Keskinen, T., Marjomäki, T.J., 2016. Environmental and genetic

effects on larval hatching time in two coregonids. Hydrobiologia 780, 135–143.

https://doi.org/10.1007/s10750-016-2807-6

Karjalainen, J., Keskinen, T., Pulkkanen, M., Marjomäki, T.J., 2015. Climate change alters the

egg development dynamics in cold-water adapted coregonids. Environ. Biol. Fishes 98,

979–991. https://doi.org/10.1007/s10641-014-0331-y

Kwain, W.-H., 1975. Embryonic development, early growth and meristic variation in rainbow

trout (Salmo gairdneri) exposed to combinations of light intensity and temperature. J. Fish.

Res. Board Canada 32, 397–402. https://doi.org/10.1139/f75-046

Lepak, T.A., Ogle, D.H., Vinson, M.R., 2017. Age, year-class strength variability, and partial

age validation of Kiyis from Lake Superior. North Am. J. Fish. Manag. 37, 1151–1160.

https://doi.org/10.1080/02755947.2017.1350222

Luczynski, M., 1991. Temperature requirements for growth and survival of larval vendace,

Coregonus albula (L.). J. Fish Biol. 38, 29–35. https://doi.org/10.1111/j.1095-

8649.1991.tb03088.x

Lynch, A.J., Taylor, W.W., Smith, K.D., 2010. The influence of changing climate on the ecology

and management of selected Laurentian Great Lakes fisheries. J. Fish Biol. 77, 1964–1982.

https://doi.org/10.1111/j.1095-8649.2010.02759.x

MacCrimmon, H.R., Kwain, W.-H., 1969. Influence of light on early development and meristic

characters in the rainbow trout, Salmo gairdneri Richardson. Can. J. Zool. 47, 631–637.

https://doi.org/10.1139/z69-108

Magnuson, J.J., Webster, K.E., Assel, R.A., Bowser, C.J., Dillon, P.J., Eaton, J.G., Evans, H.E.,

Fee, E.J., Hall, R.I., Mortsch, L.R., 1997. Potential effects of climate changes on aquatic

systems: Laurentian Great Lakes and Precambrian Shield Region. Hydrol. Process. 11, 825–

871. https://doi.org/10.1002/(SICI)1099-1085(19970630)11:8<825::AID-

HYP509>3.0.CO;2-G

Mangor☐Jensen, A., Waiwood, K.G., 1995. The effect of light exposure on buoyancy of halibut

eggs. J. Fish Biol. 47, 18–25. https://doi.org/10.1111/j.1095-8649.1995.tb01869.x

Marchesan, M., Spoto, M., Verginella, L., Ferrero, E.A., 2005. Behavioural effects of artificial

light on fish species of commercial interest. Fish. Res. 73, 171–185.

https://doi.org/10.1016/j.fishres.2004.12.009

Marsden, J.E., Casselman, J.M., Edsall, T.A., Elliott, R.F., Fitzsimons, J.D., Horns, W.H.,

Manny, B.A., McAughey, S.C., Sly, P.G., Swanson, B.L., 1995. Lake trout spawning

habitat in the Great Lakes—a review of current knowledge. J. Great Lakes Res. 21, 487–

497. https://doi.org/10.1016/S0380-1330(95)71120-0

Muir, A.M., Vecsei, P., Pratt, T.C., Krueger, C.C., Power, M., Reist, J.D., 2013. Ontogenetic

shifts in morphology and resource use of cisco Coregonus artedi. J. Fish Biol. 82, 600–617.

https://doi.org/10.1111/jfb.12016

Myers, J.T., Yule, D.L., Jones, M.L., Ahrenstorff, T.D., Hrabik, T.R., Claramunt, R.M., Ebener,

M.P., Berglund, E.K., 2015. Spatial synchrony in cisco recruitment. Fish. Res. 165, 11–21.

https://doi.org/10.1016/j.fishres.2014.12.014

Nguyen, T.D., Hawley, N., Phanikumar, M.S., 2017. Ice cover, winter circulation, and exchange

in Saginaw Bay and Lake Huron. Limnol. Oceanogr. 62, 376–393.

https://doi.org/10.1002/lno.10431

Nyberg, P., Bergstrand, E., Degerman, E., Enderlein, O., 2001. Recruitment of pelagic fish in an

unstable climate: studies in Sweden's four largest lakes. Ambio 30, 559–564.

https://doi.org/10.1579/0044-7447-30.8.559

O'Reilly, C.M., Rowley, R.J., Schneider, P., Lenters, J.D., Mcintyre, P.B., Kraemer, B.M., 2015.

Rapid and highly variable warming of lake surface waters around the globe. Geophys. Res.

Lett. 42, 1–9. https://doi.org/10.1002/2015GL066235

Oberlercher, T.M., Wanzenböck, J., 2016. Impact of electric fishing on egg survival of whitefish,

Coregonus lavaretus. Fish. Manag. Ecol. 23, 540–547. https://doi.org/10.1111/fme.12197

Parks, T.P., Rypel, A.L., 2018. Predator–prey dynamics mediate long-term production trends of

cisco (Coregonus artedi) in a northern Wisconsin lake. Can. J. Fish. Aquat. Sci. 75, 1969–

1976. https://doi.org/10.1139/cjfas-2017-0302

Paufve, M.R., 2019. Diversity in spawning habitat across Great Lakes Cisco populations

[Master's thesis]. Cornell University.

Pepin, P., 1991. Effect of temperature and size on development, mortality, and survival rates of

the pelagic early life history stages of marine fish. Can. J. Fish. Aquat. Sci. 48, 503–518.

https://doi.org/10.1139/f91-065

Petitgas, P., Alheit, J., Peck, M.A., Raab, K., Irigoien, X., Huret, M., Van Der Kooij, J.,

Pohlmann, T., Wagner, C., Zarraonaindia, I., 2012. Anchovy population expansion in the

North Sea. Mar. Ecol. Prog. Ser. 444, 1–13. https://doi.org/10.3354/meps09451

Preisendorfer, R.W., 1986. Secchi disk science: Visual optics of natural waters 1. Limnol.

Oceanogr. 31, 909–926. https://doi.org/10.4319/lo.1986.31.5.0909

R Core Team, 2021. R: A Language and Environment for Statistical Computing.

Ramus, J., Beale, S.I., Mauzerall, D., Howard, K.L., 1976. Changes in photosynthetic pigment

concentration in seaweeds as a function of water depth. Mar. Biol. 37, 223–229.

https://doi.org/10.1007/BF00387607

Roberts, A., 1978. Pineal eye and behaviour in Xenopus tadpoles. Nature 273, 774–775.

https://doi.org/10.1038/273774a0

Rogers, T.L., Munch, S.B., Stewart, S.D., Palkovacs, E.P., Giron☐Nava, A., Matsuzaki, S.S.,

Symons, C.C., 2020. Trophic control changes with season and nutrient loading in lakes.

Ecol. Lett. 23, 1287–1297. https://doi.org/10.1111/ele.13532

Rosenberg, B.D., Schroth, A.W., 2017. Coupling of reactive riverine phosphorus and iron

species during hot transport moments: impacts of land cover and seasonality.

Biogeochemistry 132, 103–122. https://doi.org/10.1007/s10533-016-0290-9

Rosinski, C.L., Vinson, M.R., Yule, D.L., 2020. Niche Partitioning among Native Ciscoes and

Nonnative Rainbow Smelt in Lake Superior. Trans. Am. Fish. Soc. 149, 184–203.

https://doi.org/10.1002/tafs.10219

Ruchin, A.B., 2020. Effect of illumination on fish and amphibian: development, growth,

physiological and biochemical processes. Rev. Aquac. 13, 567–600.

https://doi.org/10.1111/raq.12487

Ruchin, A.B., 2007. Effect of photoperiod on growth, physiologica and hematological indices of

juvenile Siberian sturgeon Acipenser baerii. Biol. Bull. 34, 583–589.

https://doi.org/10.1134/S1062359007060088

Secchi, P.A., 1864. Relazione delle esperienze fatte a bordo della pontificia pirocorvetta

Imacolata Concezione per determinare la trasparenza del mare. Mem. del PA Secchi. Nuovo

Cim. G. Fis. Chim. e Stor. Nat. Ottobre 1864, Publ. 1865 20, 205–237.

Seth, S.N.M., Nai, H.T., Rosli, M.K., Saad, S., Noor, N.M., Yukinori, M., 2014. Egg hatching

rates of brown-marbled grouper, Epinephelus fuscoguttatus under different light

wavelengths and intensities. MJS 33, 150–154. https://doi.org/10.22452/mjs.vol33no2.3

Shao, T., Wang, T., Liang, X., Li, L., 2019. Seasonal dynamics of light absorption by suspended

particulate matter and CDOM in highly turbid inland rivers on the Loess Plateau, China.

River Res. Appl. 35, 905–917. https://doi.org/10.1002/rra.3493

Sharma, S., Blagrave, K., Magnuson, J.J., O'Reilly, C.M., Oliver, S., Batt, R.D., Magee, M.R.,

Straile, D., Weyhenmeyer, G.A., Winslow, L.A., 2019. Widespread loss of lake ice around

the Northern Hemisphere in a warming world. Nat. Clim. Chang. 9, 227.

https://doi.org/10.1038/s41558-018-0393-5

Shen, Y.-J., Shen, Y., Fink, M., Kralisch, S., Chen, Y., Brenning, A., 2018. Trends and

variability in streamflow and snowmelt runoff timing in the southern Tianshan Mountains.

J. Hydrol. 557, 173–181. https://doi.org/10.1016/j.jhydrol.2017.12.035

Sinclair, M., Iles, T.D., 1989. Population regulation and speciation in the oceans. ICES J. Mar.

Sci. 45, 165–175. https://doi.org/10.1093/icesjms/45.2.165

Skjæraasen, J.E., Meager, J.J., Karlsen, Ø., Hutchings, J.A., Fernö, A., 2011. Extreme spawning-

site fidelity in Atlantic cod. ICES J. Mar. Sci. 68, 1472–1477.

https://doi.org/10.1093/icesjms/fsr055

Sommer, U., Adrian, R., De Senerpont Domis, L., Elser, J.J., Gaedke, U., Ibelings, B., Jeppesen,

E., Lürling, M., Molinero, J.C., Mooij, W.M., 2012. Beyond the Plankton Ecology Group

(PEG) model: mechanisms driving plankton succession. Annu. Rev. Ecol. Evol. Syst. 43,

429–448. https://doi.org/10.1146/annurev-ecolsys-110411-160251

Stewart, T.R., Mäkinen, M., Goulon, C., Guillard, J., Marjomäki, T.J., Lasne, E., Karjalainen, J.,

Stockwell, J.D., 2021. Influence of warming temperatures on coregonine embryogenesis

within and among species. Hydrobiologia *In review*.

https://doi.org/10.1101/2021.02.13.431107

Stockwell, J.D., Ebener, M.P., Black, J.A., Gorman, O.T., Hrabik, T.R., Kinnunen, R.E., Mattes,

W.P., Oyadomari, J.K., Schram, S.T., Schreiner, D.R., Seider, M.J., Sitar, S.P., Yule, D.L.,

2009. A Synthesis of Cisco Recovery in Lake Superior: Implications for Native Fish

Rehabilitation in the Laurentian Great Lakes. North Am. J. Fish. Manag. 29, 626–652.

https://doi.org/10.1577/M08-002.1

Stockwell, J.D., Yule, D.L., Hrabik, T.R., Sierszen, M.E., Isaac, E.J., 2014. Habitat coupling in a

large lake system: Delivery of an energy subsidy by an offshore planktivore to the nearshore

zone of Lake Superior. Freshw. Biol. 59, 1197–1212. https://doi.org/10.1111/fwb.12340

Thorrold, S.R., Latkoczy, C., Swart, P.K., Jones, C.M., 2001. Natal homing in a marine fish

metapopulation. Science 291, 297–299. https://doi.org/10.1126/science.291.5502.297

Villamizar, N., Blanco-Vives, B., Migaud, H., Davie, A., Carboni, S., Sanchez-Vazquez, F.J.,

Sánchez-Vázquez, F.J., 2011. Effects of light during early larval development of some

aquacultured teleosts: a review. Aquaculture 315, 86–94.

https://doi.org/10.1016/j.aquaculture.2010.10.036

Voeten, C.C., 2020. buildmer: Stepwise Elimination and Term Reordering for Mixed-Effects

Regression.

Walter, B., Cavalieri, D.J., Thornhill, K.L., Gasiewski, A.J., 2006. Aircraft measurements of heat

fluxes over wind-driven coastal polynyas in the Bering Sea. IEEE Trans. Geosci. Remote

Sens. 44, 3118–3134.

Wang, J., Hu, H., Schwab, D., Leshkevich, G., Beletsky, D., Hawley, N., Clites, A., 2010.

Development of the Great Lakes ice-circulation model (GLIM): application to Lake Erie in

2003–2004. J. Great Lakes Res. 36, 425–436. https://doi.org/10.1016/j.jglr.2010.04.002

Wielgolaski, F.E., Inouye, D.W., 2003. High latitude climates, in: D, S.M. (Ed.), Phenology: An

Integrative Environmental Science. Springer, pp. 175–194. https://doi.org/10.1007/978-94-

007-0632-3_12

Winslow, L.A., Read, J.S., Hansen, G.J.A., Rose, K.C., Robertson, D.M., 2017. Seasonality of

change: Summer warming rates do not fully represent effects of climate change on lake

temperatures. Limnol. Oceanogr. 62, 2168–2178. https://doi.org/10.1002/lno.10557

Yang, B., Wells, M.G., Li, J., Young, J., 2020. Mixing, stratification, and plankton under

lake☐ice during winter in a large lake: Implications for spring dissolved oxygen levels.

Limnol. Oceanogr. 65, 2713–2729. https://doi.org/10.1002/lno.11543

Zimmerman, M.S., Krueger, C.C., 2009. An Ecosystem Perspective on Re-establishing Native

Deepwater Fishes in the Laurentian Great Lakes. North Am. J. Fish. Manag. 29, 1352–

1371. https://doi.org/10.1577/M08-194.1

Tables:

Table 1. Mean daily $\pm$ SD light intensity (μmol m^{-2} s^{-1}) for three ice coverage classes from Lake

Superior and corresponding laboratory experimental light conditions.

Location	Ice Coverage (Light Treatment)		
	> 90% (Low)	40-60% (Medium)	< 10% (High)
Lake Superior	2.0 ± 1.1	3.4 ± 2.5	5.5 ± 5.9
Laboratory	0.6 ± 0.1	3.9 ± 1.9	6.2 ± 1.0

671   Table 2. Mean daily ± SD water temperatures (°C) during embryo incubations from each light

672   treatment for Lakes Superior and Ontario.

	Light Treatment		
Lake	High	Medium	Low
Superior	4.3 ± 0.2	4.3 ± 0.3	4.3 ± 0.3
Ontario	4.2 ± 0.3	4.3 ± 0.3	4.4 ± 0.4

675 Table 3. Likelihood ratio test output for each model selected for embryo survival (%), incubation

676 period (number of days post-fertilization (DPF) and accumulated degree days (°C; ADD)),

677 length-at-hatch (mm), and yolk-sac volume (mm^3) from Lakes Superior and Ontario cisco

678 (*Coregonus artedi*). pop indicates population. The full model that was selected is bolded for each

679 trait.

Trait	Model	Effect Tested	df	χ^2	p-value
Embryo Survival	**light + pop**				
	pop	light	2	181.92	< 0.001
	light	pop	1	95.00	< 0.001
Incubation Period (DPF)	**light + pop + light:pop + female:male + female + male**				
	pop + female:male + female + male	light	2	10.80	0.005
	light + female:male + female + male	pop	1	3,023.89	< 0.001
	light + pop + female:male + female + male	light:pop	2	9.66	0.008
	light + pop + light:pop + female + male	female:male	1	79.91	< 0.001
	light + pop + light:pop + female:male + male	female	1	25.29	< 0.001
	light + pop + light:pop + female:male + female	male	1	10.80	0.001
Incubation Period (ADD)	**light + pop + light:pop + female:male + female + male**				
	pop + female:male + female + male	light	2	51.72	< 0.001
	light + female:male + female + male	pop	1	3,092.41	< 0.001
	light + pop + female:male + female + male	light:pop	2	13.23	0.001
	light + pop + light:pop + female + male	female:male	1	79.99	< 0.001
	light + pop + light:pop + female:male + male	female	1	25.25	< 0.001
	light + pop + light:pop + female:male + female	male	1	10.75	< 0.001
Length-at-Hatch	**pop + female + male**				
	female + male	pop	1	373.34	< 0.001
	pop + male	female	1	100.97	< 0.001
	pop + female	male	1	11.37	< 0.001
Yolk-sac Volume	**light + pop + light:pop + female:male + female**				
	pop + female:male + female	light	2	1.96	0.376
	light + female:male + female	pop	1	712.18	< 0.001
	light + pop + female:male + female	light:pop	2	19.04	< 0.001
	light + pop + light:pop + female	female:male	1	6.52	< 0.001
	light + pop + light:pop + female:male	female	1	38.94	< 0.001

 Figure captions:

Figure 1. Histogram of annual mean ice coverage between 1-Jan and 15-Mar from 1973-2020 for

the sampling location in Lake Superior (top) and Lake Ontario (bottom). Error bars represent the

interquartile range and point indicates median. Ice coverage data was obtained from the U.S.

National Ice Center (usicecenter.gov/).

Figure 2. Daily ice coverage (%; blue line) and light intensity (μmol m^{-2} s^{-1}; gray line) based on

light sensors set at 10 m depth off Sand Island, Lake Superior. Ice coverage data above the

sensor was obtained from the U.S. National Ice Center (usicecenter.gov/).

Figure 3. Mean embryo survival (%) and incubation period (number of days post-fertilization

(DPF) and accumulated degree days (°C; ADD)) at each incubation light treatment (left) and

standardized responses to assumed optimal light conditions (i.e., low) within each population (%;

right) from Lake Superior and Lake Ontario cisco (*Coregonus artedi*). Error bars indicate

standard error.

Figure 4. Mean length-at-hatch (mm; LAH) and yolk-sac volume (mm^3; YSV) at each incubation

light treatment (left) and standardized responses to assumed optimal light conditions (i.e., low)

within each population (%; right) from Lake Superior and Lake Ontario cisco (*Coregonus*

artedi). Error bars indicate standard error.