Effects of domestication on neophobia: A comparison between the domesticated Bengalese

finch (*Lonchura striata* var. *domestica*) and its wild ancestor, the white-rumped munia

(*Lonchura striata*)

Kenta Suzuki[a,c,d,e] *, Maki Ikebuchi[c,d,e], Hiroko Kagawa[b,c,d,e], Taku Koike[c], Kazuo Okanoya[b,c,d,e]

[a] *Faculty of Health Sciences, Nihon Institute of Medical Science, Moroyama-machi 350-0435,*

JAPAN

[b] *Department of Life Sciences, Graduate School of Arts and Sciences, The University of Tokyo,*

Meguro-ku, Tokyo 153-8902, JAPAN

[c] *Laboratory for Biolinguistics, RIKEN Brain Science Institute (BSI), Wako, Saitama 351-0198,*

JAPAN

[d] *Okanoya Emotional Information Project, Exploratory Research for Advanced Technology*

(ERATO), Japan Science and Technology Agency (JST), Wako, Saitama 351-0198, JAPAN

[e] *Cognition and Behavior Joint Research Laboratory, RIKEN Center for Brain Science (CBS), Wako,*

Saitama 351-0198, JAPAN

*Corresponding author: Faculty of Health Sciences, Nihon Institute of Medical Science, 1276

Shimogawara, Moroyama-machi, Iruma-gun, Saitama 350-0435, JAPAN. Tel.: +81-49-294-9000,

Fax: +81-49-294-9009.

E-mail address: kentszk@gmail.com (K. Suzuki).

ABSTRACT

Bengalese finches (*Lonchura striata* var. *domestica*) have more complex song traits than their wild

ancestors, white-rumped munias (*Lonchura striata*). Domesticated finches are likely able to allocate

more resources to song development rather than allocating resources to mechanisms aimed at coping

with predation, which are no longer needed under domesticated conditions. Here, we aimed to

examine the effects of changes in selection pressure due to domestication on the behaviour of

Bengalese finches and to contemplate the possible evolutionary mechanisms underlying these

changes. To do so, we compared neophobic responses to novel-object conditions as an assessment of

reactions to potential predators. We studied groups of Bengalese finches and white-rumped munias

and found that Bengalese finches were more likely to eat the food provided to them under

novel-object conditions. Bengalese finches had a shorter latency time to eat, and this latency time

was less affected by the novel object in the case of Bengalese finches compared to white-rumped

munias. Therefore, Bengalese finches have reduced neophobic responses due to domestication. The

behavioural strategies of white-rumped munias appear to be more suitable for natural environments,

which include unpredictable risks, whereas Bengalese finches have likely adapted their behaviour to

the conditions of artificial selection.

Keywords: Bengalese finch (*Lonchura striata* var. *domestica*); Domestication; Neophobia;

Neophobic response; White-rumped munia (*Lonchura striata*)

Abbreviations

BF, Bengalese finch; WRM, white-rumped munia.

1. Introduction

The Bengalese finch (BF, *Lonchura striata* var. *domestica*) is a domesticated variety of the

wild white-rumped munia (WRM, *Lonchura striata*), which was imported from China to Japan

approximately 250 years ago (Washio, 1996; Okanoya, 2004a; Svanberg, 2008). BFs sing

phonologically and syntactically complex songs, as opposed to the stereotypical simple songs sung

by WRMs (Honda and Okanoya, 1999; Okanoya, 2004a, b). BFs were domesticated for their

plumage colour morphs, reproductive efficiency, parental behaviour, and ability to foster other bird

species (Taka-Tsukasa, 1917), but the complex songs of BF have not been selected for in an artificial

manner (Washio, 1996). Complex song as a high-quality sexual trait is believed to have evolved

through the domestication process in BFs (Okanoya, 2004a, b). WRMs are thought to experience

strong natural selection pressures in wild environments. Conversely, domesticated BFs experience

safe and resource-rich conditions under human-controlled rearing conditions (i.e., no environmental

perturbations, no predation, abundant food, and low risk of parasitism and injury). Thus, we

hypothesized that the relaxation of natural selection pressures and the presence of artificial selection

57   may have led BFs to allocate fewer resources to behaviours associated with efforts for survival and

58   more to efforts associated with reproduction.

59   Domestication results in species being removed from many natural selection pressures;

however, domesticated species are exposed to artificial selection pressures exerted by their captive

environments and by humans (Fox, 1968; Boice, 1973; Ratner and Boice, 1975; Price, 1984). This

change in selection pressure results in changes in physiological and behavioural traits (Hale, 1969;

Price, 1984, Künzl and Sachser, 1999; Lepage et al., 2000). Domesticated species have reduced

behavioural expressions related to ontogenetic survival, such as fear responses (Desforges and

Wood-Gush, 1975; Schütz et al., 2001; Campler et al., 2009), and have enhanced behavioural traits

that are not directly related to survival, such as sexual and reproductive behaviours (Künzl and

Sachser, 1999). Therefore, these behavioural changes have been a major target of domestication

effects.

69   In natural environments, birds must be vigilant of unpredictable risks such as predation

when foraging or eating; this is particularly important while approaching novel conditions such as

novel places, objects, and food items. Neophobia is the aversion that an animal displays while

approaching novel conditions (Greenberg, 2003). In birds, the most well-established behavioural

responses to novel conditions (neophobic responses) are the reactions to potential predators

(Greenberg, 1990). In the wild, the presence of unpredictable risks is considered to enhance induced

neophobia (Brown et al., 2020). Domesticated BFs are likely able to allocate more resources to

reproduction (e.g., song development) as a trade-off for a reduction in the behaviours and strategies

aimed at coping with predation (e.g., neophobic responses). In the present study, to examine the

effects of changes in selection pressure due to domestication on the behaviour of BFs, and to

contemplate the evolutionary mechanisms underlying these effects, we compared neophobic

responses to novel conditions in BFs and their wild ancestors, WRMs.

2. Materials and methods

2.1 Birds used in this study

Seventy BFs and 58 WRMs were used in the experiment. All the birds were sexually

mature. BFs (n = 10) were bought from local suppliers and others (n = 60) were bred in our

laboratory. WRMs were captured in the wild in Taiwan. These captured birds were reared for more

than a year in our laboratory before being used in this study. Birds were housed in a group of 8 (one

group of WRM) or 10 per cage resulting in us using seven groups of BFs and six groups of WRMs

for our experiments. Sex ratios of the birds that made up the groups were different for each cage

(BF-1 and WRM-1: all females, BF-2, 3 and WRM-2: all males, BF-4, 5 and WRM-3, 4: 50% of

each sex, BF-6, 7 and WRM-5, 6: 6 males and 4 females). Birds were housed in stainless steel cages

($370 \times 415 \times 440$ mm) within an animal-rearing room at RIKEN Brain Science Institute (BSI) and

given finch seed mixture, shell grit, and vitamin-enhanced water *ad libitum*. The light cycle was kept

constant at 14 h light and 10 h dark. The room was maintained at an ambient temperature of

approximately 25 °C, with a humidity of 50 %. The birds were acclimatized for at least four months

to the animal-rearing room before performing the experiments. All experimental procedures and the

housing conditions of the birds were approved by the Animal Experiments Committee at RIKEN

(#H20-2-231, #H22-2-217), and all the birds were cared for in accordance with the Institutional

Guidelines of RIKEN for experiments using animals.

2.2 Novel-object experiment

Before the experiment, each group of birds was transferred from their home cage in the

animal-rearing room to a testing cage (with the same dimensions and characteristics as the home

cage) in a sound-proof box. To avoid the effects of isolation stress, experiments were conducted

using the same groups of birds that were housed together during keeping. The test cage was

equipped with two wooden perches and with food and water. Experiments were conducted under two

conditions: the control (non-object) condition and the novel-object condition. Experiments were

carried out under the different conditions on different days. All groups were first examined under the

control (non-object) condition to avoid the effects of disturbance by the novel-object. To eliminate

the influence of the degree of acclimation to the soundproof box and the testing cage, the number of

days between the two conditions (the control and the novel-object condition) was randomly set

between 1 d and 7 d in both BFs and WRMs. New food and water were provided at the beginning of

the experiments. Under the novel-object condition, a novel object (a small toy dog, approximately 15

mm × 15 mm × 20 mm) was placed on the food. The behaviour of birds was recorded for 60 min

using a video camera. We quantified the number of groups of birds that ate food during the

observation period (at least one bird in the group) and the latency time (the time that passed before

first birds started eating the food) under both conditions. If the birds did not approach or eat the food

during the test period, the latency time of the birds was set at the maximum time (3600 s). All groups

were used only once under both conditions. All tests were conducted between 13:00 and 15:00 to

avoid other factors that could affect our results, such as diurnal changes in activity, hunger, and

hormone levels.

2.3 Statistical analyses

The numbers of groups of BFs and WRMs that ate the food during the observation period

were compared with Fisher's exact test. The latency times to start eating the food are represented as

the mean ± standard error of the mean (SEM). The latency times in BFs and WRMs were analysed

using a repeated measures two-way analysis of variance (ANOVA) (birds × conditions) with

post-hoc Bonferroni/Dunn tests. The difference in the effect of the novel object on the latency times

(calculated as the latency times under the novel-object conditions - the latency times under control

conditions) between BFs and WRMs were analysed by an unpaired *t*-test with a Welch's correction.

Values of $p < 0.05$ were considered significant. We used three different statistical software packages

for analyses. Statistical analyses were performed using R statistical software (version 4.03, R

Foundation for Statistical Computing, Vienna, Austria) for Fisher's extract test, Stat View software

(version 5, SAS Institute Inc., Cary, NC, USA) for repeated measures 2-way ANOVA with post-hoc

Bonferroni/Dunn tests, and GraphPad Prism software (version 4, GraphPad Software Inc., San

Diego, CA, USA) for *t*-tests with a Welch's correction.

3. Results

3.1 Responses of BFs and WRMs to novel objects

We examined the effects of domestication on behavioural responses to novel objects in

BFs and WRMs. Under control conditions, all groups of birds (both BFs and WRMs) ate the food at

some point during the observation time (BFs: 7/7, 100 %; WRMs: 6/6, 100 %). Thus, there were no

differences between BFs and WRMs ($p = 1.00$). However, under the novel-object conditions, six of

the seven groups of BFs (6/7, 85.7 %) ate the food at some point during the observation time,

whereas none of the WRMs groups ate the food (0/6, 0 %). Therefore, there were significant

differences between BFs and WRMs ($p < 0.01$). In BFs, there were no significant differences in the

number of groups that ate the food between the control and novel-object conditions ($p > 0.90$).

However, WRMs were significantly less likely to eat the food under the novel-object conditions ($p <$

0.01).

3.2 Latency times in BFs and WRMs to start eating food under control and novel-object conditions

152   The latency times to start eating the food were compared between BFs and WRMs (Fig.

1). Under the control conditions, the latency times were 25.6 ± 9.0 s for BFs and 159.0 ± 46.7 s for

WRMs. Under the novel-object conditions, the latency times were 1204.9 ± 524.9 s for BFs, but

none of the WRMs approached the food within the observation time (3600.0 ± 0.0 s). The latency

times was significantly affected by the birds (BF or WRM, $F(1, 11) = 16.9$, $p < 0.01$), conditions

(novel object or control, $F(1, 11) = 69.4$, $p < 0.0001$), and interactions (birds × conditions, $F(1, 11)$

$= 12.7$, $p < 0.01$). Latency times to eat were significantly shorter in BFs than in WRMs regardless of

the presence or absence of novel objects (control conditions: $p < 0.05$, novel-object conditions: $p <$

0.01). In both species, the latency times were extended by the presence of the novel object (BFs: $p <$

0.05, WRMs: $p < 0.0001$), and the extended times were significantly longer in WRMs than those in

BFs (BFs: 1379.3 ± 530.7 s, WRM: 3441.0 ± 46.7 s; $t = 3.87$, $p < 0.01$).

4. Discussion

We compared responses to novel conditions in BFs and their wild ancestor, WRMs, and

examined the effects of changes in selection pressure due to domestication. Our findings indicate

that these two birds responded differently to novel-object conditions. BFs had significantly lower

neophobia levels (more groups of BFs ate the food, and BFs had shorter latency times to eat under

novel-object conditions) than those of their wild ancestors, the WRMs. The novel objects extended

the latency times to eat in both birds, but latency times were significantly longer in WRMs than

those in BFs. Therefore, it is likely that BFs have reduced neophobic responses due to domestication.

Under control conditions, the proportion of the groups that ate the food was the same in BFs and

WRMs. However, BFs ate the food sooner than WRMs. It is probable that the WRMs were more

cautious about the experimental situation (the soundproof box is different from the WRMs' normal

environment). In a previous report, male WRMs had higher faecal corticosterone concentrations than

those in male BFs (Suzuki et al., 2012; 2014a, b). Therefore, WRMs are considered to have a higher

level of baseline vigilance than that in BFs. This strain difference with respect to vigilance seems to

have been further increased under novel-object conditions.

Our results are similar to those of a previous study that found that domesticated ducks

(*Anas platyrhynchos* var. *domesticus*) had lower levels of neophobia than those of wild mallards

(*Anas platyrhynchos*) (Desforges and Wood-Gush, 1975). In addition, laboratory mice (*Mus*

musculus) had lower levels of neophobia than those in wild mice (Meddock and Osborn, 1968). In a

comparison of dogs (*Canis lupus familiaris*) and wolves (*Canis lupus*), dogs had lower levels of

neophobia than those in wolves, but dogs had an overall lower interest in novel objects (Moretti et

al., 2015). In the present study, BFs showed a fear response to novel objects and were considered to

have no diminished interest in novel objects.

This study was conducted using groups of birds, and, as such, there may have been a social

effect. Because it was difficult to identify individuals in this study, the analysis was conducted as a

group-by-group comparison, and it was therefore not possible to take into account the differences in

the responses of each individual. Wolves and dogs spend more time approaching a novel object in

groups than as individuals, so risk sharing may increase vigilance (Moretti et al., 2015). In contrast,

in the zebra finches (*Taeniopygia guttata*), existence of conspecifics can modulate (decrease) the

responses to stressors (e.g., novel stimuli), known as "social buffering" (Coleman and Mellgren,

1994; Emmerson and Spencer, 2017). Responses to novel objects may also be influenced by the

identities of the group members. In the Gouldian finches (*Chloebia gouldiae*), shy birds took more

risks when they were paired with bold partners, and bold birds took fewer risks when they were

paired with shy partners (King et al., 2015). Since this experiment was conducted using groups, there

may have been an effect of risk sharing or social buffering. Therefore, in experiments conducted

using groups, individual differences may be reduced, and more of the characteristics of the groups

may be expressed. Conversely, in experiments conducted using individuals, there is a high

possibility that individual differences will be noticeable. Future studies on BFs and WRMs should

take into consideration the reactions of each individual bird to novel-object conditions. In a previous

study, we compared the tonic immobility response, which can be used as a measure of fear

responses, between individual BFs and WRMs and found that BFs had lower fear responses (Suzuki

et al., 2013). Therefore, we believe that the fear response is lower in BFs than that in WRMs, even

on an individual basis.

In this study, the effects of sex ratio and environmental or rearing conditions (bought from

local supplier, wild-caught, or bred in the laboratory) were ruled out because the groups in the cages

were neither separated by sex nor by environmental or rearing conditions. Moreover, previous

studies have shown that sex and environmental or rearing conditions of the white-rumped munia

(captive-born, bought from a supplier, or captured) did not affect the fear response (tonic immobility

reactions) (Suzuki et al., 2013). Additionally, the conditions under which Bengalese finches and

white-rumped munias were bred or reared did not affect the corticosterone levels, which is known to

affect fear responses (Suzuki et al., 2012). All the munias used in this experiment were wild-caught;

however, differences in rearing conditions have not been an issue in previous experiments.

Therefore, the magnitude of the impact of these factors is small, but it must be considered in future

research.

We hypothesized that BFs may allocate more resources to reproduction (e.g., song

development) as a trade-off for a reduction in the behavioural strategies aimed at coping with

predation (e.g., neophobic responses). The behavioural strategies of WRMs appear to be more

suitable for natural, wild environments, which include unpredictable risks (e.g., predation risk),

whereas BFs have likely adapted their behaviour to the conditions of artificial selection. Singing is

important for reproduction but can increase susceptibility to predators. Birds are able to respond to

increased predation risk by reducing singing, and it has been found that in areas with increased

perceived predation risk, fewer songs are produced per bird (Abbey-Lee et al., 2016). Continuous

auditory feedback is necessary to maintain the complex songs of the BF (Okanoya and Yamaguchi,

1997), and domesticated BFs might be able to allocate more time to singing and song maintenance

due to the reduced need for neophobic responses. Songs incur a cost in their development and

maintenance linked to corticosterone levels (Gil and Gahr, 2002; Okanoya, 2004a, b; Wada et al.,

2008; MacDougall-Shackleton et al., 2009). Administration of corticosterone during the

developmental period enhances the fear response (including the neophobic response) but reduces

song complexity (Spencer et al., 2003; Buchanan et al., 2004; Emmerson and Spencer, 2017).

Chronic stress and corticosterone treatment enhance dendritic expansion in neurons of the

basolateral amygdala and increase anxiety as a function of that brain region (Vyas et al., 2002; Vyas

and Chattarji, 2004; Mitra et al., 2005; Mitra and Sapolsky, 2008). In contrast, chronic stress and

corticosterone administration in juvenile and adult birds reduce the volume of song nuclei and the

numbers of new and matured neurons in the HVC (Buchanan et al., 2004; Newman et al., 2010).

Both the amygdala and the song nuclei have two receptors (mineralocorticoid and glucocorticoid

receptors) for corticosterone, and different levels of their expression might cause different

site-specific effects (Matsunaga et al., 2011; Suzuki et al., 2011; Suzuki et al., 2014a). Therefore,

low levels of corticosterone in BFs (Suzuki et al., 2012; Suzuki et al., 2014b) are considered to be

required for the development of low neophobic responses and complex songs.

In summary, our results suggest that the domestication process led to differences in the

responses of BFs and WRMs to novel conditions. The behavioural strategies of WRMs seem to be

suitable for the natural environment, which includes unpredictable risks, whereas BFs might have

adapted their behaviours to the conditions of captivity and artificial selection. BFs are likely able to

allocate the resources that would be needed in the wild to cope with predators to song development,

due to domestication. In addition, there were variations in latency times to eat under novel object

conditions in BFs, which indicate individual differences in neophobic responses. Although this study

was a group-by-group examination, future studies would need to focus on individual differences and

examine the trade-off between personality traits (such as neophobia) and reproductive traits (such as

song complexity). This is the first study to evaluate neophobia in Bengalese finches and

white-rumped munias, and also explore the evolutionary mechanisms of behavioural changes in

domesticated animals, including the evolution of complex songs.

Acknowledgements

We thank members of the Biolinguistics team (RIKEN BSI, Saitama, Japan) for animal care. This study was supported by RIKEN BSI and Japan science and technology agency (JST) - the exploratory research for advanced technology (ERATO) research funding.

Declarations of interest

None

Author contributions

Kenta Suzuki: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Visualization, Writing - original draft; **Maki Ikebuchi:** Conceptualization; **Hiroko Kagawa:** Conceptualization, Writing - review & editing; **Taku Koike:** Data curation, Investigation; **Kazuo Okanoya:** Conceptualization, Funding acquisition, Supervision, Writing - review & editing.

Funding

This work was supported by JSPS KAKENHI [grant numbers 15K14581, 17H06380, 20H00105] to KO.

References

Abbey-Lee, R.N., Kaiser, A., Mouchet, A., Dingemanse, N.J., 2016. Immediate and carry-over effects of perceived predation risk on communication behavior in wild birds. Behav. Ecol. 27(3), 708–716. https://doi.org/10.1093/beheco/arv210.

Boice, R., 1973. Domestication. Psychol. Bull. 80, 215–230. https://doi.org/10.1037/h00348.

Brown, G.E., Demers, E.E., Goldman, J.A., Singh, A., Chivers, D.P., Ferrari, M.C., 2020. Unpredictable risk enhances induced neophobia in northern red-bellied dace. Anim. Behav. 168, 121–127. https://doi.org/10.1016/j.anbehav.2020.08.012.

Buchanan, K.L., Leitner, S., Spencer, K.A., Goldsmith, A.R. and Catchpole, C.K., 2004. Developmental stress selectively affects the song control nucleus HVC in the zebra finch. Proc. R. Soc. B 271, 2381–2386. https://doi.org/10.1098/rspb.2004.2874.

Campler, M., Jöngren, M., Jensen, P., 2009. Fearfulness in red junglefowl and domesticated White Leghorn chickens. Behav. Processes 81, 39–43. https://doi.org/10.1016/j.beproc.2008.12.018.

Coleman, S. L., Mellgren, R. L., 1994. Neophobia when feeding alone or in flocks in zebra finches,

Taeniopygia guttata. Anim. Behav. 48(4), 903–907. https://doi.org/10.1006/anbe.1994.1315.

Desforges, M.F., Wood-Gush, D.G.M., 1975. A behavioural comparison of domestic and mallard

ducks.   Habituation   and   flight   reactions.   Anim.   Behav.   23,   692–697.

https://doi.org/10.1016/0003-3472(75)90145-1.

Emmerson, M.G., Spencer, K.A., 2017. Long-term effects of adolescent stress on neophobic

behaviors in zebra finches are modulated by social context when in adulthood. Horm. Behav. 90,

48–55. https://doi.org/10.1016/j.yhbeh.2017.02.004.

Fox, M.W., 1968. The influence of domestication upon behavior of animals. in: Fox, M.W. (Ed),

Abnormal behaviour in animals. Saunders, Philadelphia, London, pp. 64–76.

Gil, D., Gahr, M., 2002. The honesty of bird song: multiple constraints for multiple traits. Trends

Ecol. Evol. 17, 133–141. https://doi.org/10.1016/S0169-5347(02)02410-2.

Greenberg, R.S., 1990. Ecological plasticity, neophobia, and resource use in birds. Stud. Avian Biol.

13, 431–437.

Greenberg, R.S., 2003. The role of neophobia and neophilia in the development of innovative

behaviour of birds. in: Reader, S.N. and Laland, K.N. (Eds), Animal innovation. Oxford university

press, NY, pp. 175–196.

Hale, E.B., 1969. Domestication and the evolution of behaviour. in: Hafez, E.S.E. (Ed) The

Behaviour of Domestic Animals, 2nd Edition. Baillière, Tindall and Cassell, London, pp. 22–42.

Honda, E., Okanoya, K., 1999. Acoustical and syntactical comparisons between songs of the

white-backed munia (*Lonchura striata*) and its domesticated strain, the Bengalese finch (*Lonchura*

striata var. *domestica*). Zool. Sci. 16, 319–326. https://doi.org/10.2108/zsj.16.319.

King, A.J., Williams, L.J., Mettke-Hofmann, C., 2015. The effects of social conformity on Gouldian

finch personality. Anim. Behav. *99*, 25–31. https://doi.org/10.1016/j.anbehav.2014.10.016.

Künzl, C., Sachser, N., 1999. The behavioral endocrinology of domestication: A comparison

between the domestic guinea pig (*Cavia aperea*f.*porcellus*) and its wild ancestor, the cavy (*Cavia*

aperea). Horm. Behav. 35, 28–37. https://doi.org/10.1006/hbeh.1998.1493.

Lepage, O., Øverli, Ø., Petersson, E., Järvi, T., Winberg, S., 2000. Differential stress coping in wild

and domesticated sea trout. Brain Behav. Evol. 56, 259–268. https://doi.org/10.1159/000047209.

MacDougall-Shackleton, S.A., Dindia, L., Newman, A.E., Potvin, D.A., Stewart, K.A.,

MacDougall-Shackleton, E.A., 2009. Stress, song and survival in sparrows. Biol. Lett. 5, 746–748.

https://doi.org/10.1098/rsbl.2009.0382.

Matsunaga, E., Suzuki, K., Kobayashi, T., Okanoya, K., 2011. Comparative analysis of

mineralocorticoid receptor expression among vocal learners (Bengalese finch and budgerigar) and

non-vocal learners (quail and ring dove) has implications for the evolution of avian vocal learning.

Dev. Growth Differ. 53, 961–970. https://doi.org/10.1111/j.1440-169X.2011.01302.x.

Meddock, T.D., Osborn, D.R., 1968. Neophobia in wild and laboratory mice. Psychonom. Sci. 12(5),

223–223. https://doi.org/10.3758/BF03331280.

Mitra, R., Jadhav, S., McEwen, B.S., Vyas, A., Chattarji, S., 2005. Stress duration modulates the

spatiotemporal patterns of spine formation in the basolateral amygdala. Proc. Natl. Acad. Sci. USA

102, 9371–9376. https://doi.org/10.1073/pnas.0504011102.

Mitra, R., Sapolsky, R.M., 2008. Acute corticosterone treatment is sufficient to induce anxiety and

amygdaloid dendritic hypertrophy. Proc. Natl. Acad. Sci. USA 105, 5573–5578.

https://doi.org/10.1073/pnas.0705615105.

Moretti, L., Hentrup, M., Kotrschal, K., Range, F., 2015. The influence of relationships on

neophobia and exploration in wolves and dogs. Anim. Behav. 107, 159–173.

https://doi.org/10.1016/j.anbehav.2015.06.008.

Newman, A.E., MacDougall-Shackleton, S.A., An, Y.S., Kriengwatana, B., Soma, K.K., 2010.

Corticosterone and dehydroepiandrosterone have opposing effects on adult neuroplasticity in the

avian song control system. J. Comp. Neurol. 518, 3662–3678. https://doi.org/10.1002/cne.22395.

Okanoya, K., 2004a. Song syntax in Bengalese finches: proximate and ultimate analyses. Adv. Stud.

Behav. 34, 297–346. https://doi.org/10.1016/S0065-3454(04)34008-8.

Okanoya, K., 2004b. The Bengalese finch: a window on the behavioral neurobiology of birdsong

syntax. Ann. N. Y. Acad. Sci. 1016, 724–735. https://doi.org/10.1196/annals.1298.026.

Okanoya, K., Yamaguchi, A., 1997. Adult Bengalese finches (*Lonchura striata* var. *domestica*)

require real□time auditory feedback to produce normal song syntax. J. neurobiol. 33(4), 343–356.

https://doi.org/10.1002/(SICI)1097-4695(199710)33:4<343::AID-NEU1>3.0.CO;2-A.

Price, E.O., 1984. Behavioral aspects of animal domestication. Q. Rev. Biol. 59, 1–32.

https://doi.org/10.1086/413673.

Ratner, S.C., Boice, R., 1975. Effects of domestication on behaviour. in: Hafez, E.S.E. (ed), The

behaviour of domestic animals, third ed., Williams & Wilkins, Baltimore, pp. 3–19.

Schütz, K.E., Forkman, B., Jensen, P., 2001. Domestication effects on foraging strategy, social

behaviour and different fear responses: a comparison between the red junglefowl (Gallus gallus) and

a modern layer strain. Appl. Anim. Behav. Sci. 74(1), 1–14.

https://doi.org/10.1016/S0168-1591(01)00156-3.

Spencer, K.A., Buchanan, K.L., Goldsmith, A.R., Catchpole, C.K., 2003. Song as an honest signal of

developmental stress in the zebra finch (*Taeniopygia guttata*). Horm. Behav. 44, 132–139.

https://doi.org/10.1016/S0018-506X(03)00124-7.

Suzuki, K., Ikebuchi, M., Bischof, H. J., Okanoya, K., 2014a. Behavioral and neural trade-offs

between song complexity and stress reaction in a wild and a domesticated finch strain. Neurosci.

Biobehav. Rev. 46, 547–556. https://doi.org/10.1016/j.neubiorev.2014.07.011.

Suzuki, K., Ikebuchi, M., Okanoya, K., 2013. The impact of domestication on fearfulness: a

comparison of tonic immobility reactions in wild and domesticated finches. Behav. Processes. 100,

58–63. https://doi.org/10.1016/j.beproc.2013.08.004.

Suzuki, K., Matsunaga, E., Kobayashi, T., Okanoya, K., 2011. Expression pattern of

mineralocorticoid and glucocorticoid receptors in Bengalese finch (*Lonchura striatta* var. *domestica*)

brain suggest the relation between stress hormone and the song system development. Neuroscience

194, 72–83. https://doi.org/10.1016/j.neuroscience.2011.07.073.

Suzuki, K., Matsunaga, E., Yamada, H., Kobayashi, T., Okanoya, K., 2014b. Complex song

development and stress hormone levels in the Bengalese finch. Avian Biol. Res. 7(1), 10–17.

https://doi.org/10.3184/175815514X13903270812502.

Suzuki, K., Yamada, H., Kobayashi, T., Okanoya, K., 2012. Decreased fecal corticosterone levels

due to domestication: a comparison between the white-backed munia (*Lonchura striata*) and its

domesticated strain, the Bengalese finch (*Lonchura striata* var. *domestica*) with a suggestion for

complex song evolution. J. Exp. Zool. A 317, 561–570. https://doi.org/10.1002/jez.1748.

Svanberg, I., 2008. Towards a cultural history of the Bengalese Finch (*Lonchura domestica*). Der

Zoologische Garten 77(5–6), 334–344. https://doi.org/10.1016/j.zoolgart.2008.05.003.

Taka-Tsukasa, N., 1917. Companion Birds. Shoka-bo, Tokyo (in Japanese).

Vyas, A., Chattarji, S., 2004. Modulation of different states of anxiety-like behavior by chronic

stress. Behav. Neurosci. 118, 1450–1454. https://doi.org/10.1037/0735-7044.118.6.1450.

Vyas, A., Mitra, R., Shankaranarayana Rao, B.S., Chattarji, S., 2002. Chronic stress induces

contrasting patterns of dendritic remodeling in hippocampal and amygdaloid neurons. J. Neurosci.

22, 6810–6818. https://doi.org/10.1523/JNEUROSCI.22-15-06810.2002.

Wada, H., Salvante, K.G., Stables, C., Wagner, E., Williams, T.D., Breuner, C.W., 2008.

Adrenocortical responses in zebra finches (*Taeniopygia guttata*): individual variation, repeatability,

and relationship to phenotypic quality. Horm. Behav. 53, 472–480.

https://doi.org/10.1016/j.yhbeh.2007.11.018.

Washio, K., 1996. Chasing the mystery of the Bengalese finch. Modern Literature Inc., Tokyo. (in

Japanese)

Fig. 1. Latency times (in seconds) to start eating food in Bengalese finches (BFs, *Lonchura striata*

var. *domestica*) and white-rumped munias (WRMs, *Lonchura. striata*). Bars indicate means, and

vertical lines indicate standard errors of mean (SEM). BFs had significantly shorter latency times to

start eating the provided food than those WRMs under both conditions (control and exposure to a

novel object, i.e., a small dog toy). Latency times were extended under novel-object conditions in

both BFs and WRMs. Different letters indicate *significant differences between* groups (a vs b: *p*

< 0.05, A vs B: *p* < 0.01, a vs A: *p* < 0.05, b vs B: *p* < 0.0001).