A year in the life of the Eastern Mediterranean:

Monthly dynamics of phytoplankton and bacterioplankton in an ultra-oligotrophic sea

Tom Reich[1], Tal Ben-Ezra[1,2], Natalya Belkin[3], Anat Tsemel[1,2], Dikla Aharonovich[1], Dalit

Roth-Rosenberg[1], Shira Givati[1], Or Bialik[4], Barak Herut[3,4], Ilana Berman-Frank[1], Miguel

Frada[5,6], Michael D. Krom[1,2], Yoav Lehahn[4], Eyal Rahav[3]* and Daniel Sher[1]*

[1]Department of Marine Biology, Leon H. Charney School of Marine Sciences, University of

Haifa, Israel; [2]Morris Kahn Marine Research Station, Leon H. Charney School of Marine

Science, University of Haifa, Haifa, 3498838, Israel. [3]Israel Oceanographic and Limnological

Research, Haifa, Israel. [4]Department of Maritime Geosciences, Leon H. Charney School of

Marine Sciences, University of Haifa, Israel. [5]The Interuniversity Institute for Marine

Sciences in Eilat, POB 469, 88103 Eilat, Israel. [6]Dept. of Ecology, Evolution and Behavior –

Alexander Silberman Institute of Life Sciences, Hebrew University of Jerusalem, Jerusalem

91904, Israel

*Corresponding authors:

dsher@univ.haifa.ac.il

eyal.rahav@ocean.org.il

Keywords: Eastern Mediterranean, Levantine Basin, seasonal dynamics, primary

productivity, bacterial productivity, phytoplankton, *Prochlorococcus*, *Synechococcus*, pico-

eukaryotes.

Highlights

• Bacterioplankton dynamics were assessed monthly in the Eastern Mediterranean Sea

• Small-sized picophytoplankton numerically dominated the phytoplankton community

• Seasonal phytoplankton dynamics are similar to BATS and Red Sea, but not to HOT

• Annual primary productivity is among the lowest in the world's oceans

• Bacterial to primary production ratio is higher than most oligotrophic seas

Abstract

The Eastern Mediterranean Sea (EMS) is a poorly studied ultra-oligotrophic marine environment, dominated by small-size phyto- and bacterioplankton. Here, we describe the dynamics of a single annual cycle (2018-19) of phyto- and bacterioplankton (abundances, pigments and productivity) in relation to the physical and chemical conditions in the photic water column at an offshore EMS site (Station THEMO-2, ~1,500m depth, 50km offshore). We show that phytoplankton biomass (as chlorophyll a), primary and bacterial productivity differed between the mixed winter (January-April) and the thermally stratified (May-December) periods. *Prochlorococcus* and *Synechococcus* numerically dominated the picophytoplankton populations, with each clade revealing different temporal and depth changes indicative to them, while pico-eukaryotes (primarily haptophytes) were less abundant, yet likely contributed significant biomass. Estimated primary productivity (~32 gC m^{-2} y^{-1}) was lower compared with other well-studied oligotrophic locations, including the north Atlantic and Pacific (BATS and HOT observatories), the western Mediterranean (DYFAMED observatory) and the Red Sea, and was on-par with the ultra-oligotrophic South Pacific Gyre. In contrast, integrated bacterial production (~11 gC m^{-2} y^{-1}) was similar to other oligotrophic locations. Phytoplankton seasonal dynamics were similar to those at BATS and the Red Sea, suggesting an observable effect of winter mixing in this ultra-oligotrophic location. These results highlight the ultra-oligotrophic conditions in the EMS and provide, for the first time in this region, a full-year baseline and context to ocean observatories in the region.

1 Introduction

Convective mixing of the water column is one of the main mechanisms responsible for transport of nutrients to the photic zone of oligotrophic oceans and seas, often resulting in increased phytoplankton biomass and activity (Behrenfeld, 2010). This process usually occurs during wintertime upon the progressive cooling of the sea surface, although other mechanisms may also deliver nutrients to the mixed layer such as physical upwelling along shorelines, frontal systems and gyres (Anabalón et al., 2016), nutrient runoff from rivers (Jickells, 1998), and atmospheric deposition (Guieu et al., 2014). Stratification is established during springtime, nutrients gradually become depleted, which, together with other processes such as increased predation, lead to a decline in algal biomass and productivity (Behrenfeld and Boss, 2014). This cycle has been extensively studied over decades in the major oligotrophic gyres (e.g. HOT and BATS) (Steinberg et al., 2001), as well in other locations including the Western Mediterranean and Red Seas (Genin et al., 2018; Marty et al., 2002; Marty and Chiavérini, 2010), and is considered a fundamental driving force of marine ecosystem structure and function.

The offshore water of the Eastern Mediterranean Sea are considered ultra-oligotrophic (Berman-Frank and Rahav, 2012; Siokou-Frangou et al., 2010). The oligotrophic nature of this system, and especially of the Eastern Mediterranean Sea (EMS), is mainly driven by its general anti-estuarine circulation (Pinardi and Masetti, 2000). Additionally, the relatively stable density stratification throughout most of the year (winter mixing rarely exceeds the 200 m depth, (D'Ortenzio et al., 2005) has been suggested to result in a very low supply of deep water nutrients to the euphotic zone (Hazan et al., 2018). Finally, modern riverine inputs into the EMS are extremely low, especially since the Nile was dammed in the 1960s (Krom et al., 2014). The abovementioned conditions makes the EMS, and specifically the easternmost

Levantine basin, among the warmest, saltiest and least productive waters in the world (Ozer

et al., 2017).

Despite the importance of the EMS coastline in providing ecosystem services to over 70

million people e.g. (Peled et al., 2018), it lacks the continued high resolution records

available from other oligotrophic regions such as the North Atlantic (the Bermuda Atlantic

Time Series, BATS), North Pacific (The Hawaii Ocean Time-Series station ALOHA, HOT),

the Western Mediterranean e.g. station DYFAMED (Marty et al., 2002; Marty and

Chiavérini, 2010) and, to a lesser extent, the Red Sea (Shaked and Genin, 2017). Specifically,

detailed phytoplankton and bacterioplankton time-series are missing, and are important in

order to both understand the current system and the ways it may be impacted by local,

regional and global change (e.g. (Marty and Chiavérini, 2010). Thus, studying the dynamics

of the microorganisms at the base of the marine food-web over time are of great ecological

importance.

Studies from the early 1980's showed a clear seasonal cycle of chlorophyll a (a proxy of algal

biomass) and primary production in surface waters of both neritic and pelagic locations, with

the changes attributed primarily to picophytoplankton (Azov, 1986; Berman et al., 1984;

Kimor and Wood, 1975; Yacobi et al., 1995). More recent studies of the EMS show similar

seasonal trends using remote-sensing of surface chlorophyll a (Rosenberg et al., 2020) and in

measurements of cell numbers, production and dinitrogen fixation in coastal waters (Rahav et

al., 2018; Raveh et al., 2015). However, to the best of our knowledge, no detailed temporal

(monthly) measurements have been presented of microbial processes and phytoplankton

community structure at the offshore EMS waters. In this study, we followed the composition

and activity of phytoplankton and bacterioplankton from a pelagic location (station THEMO-

2) in the offshore EMS over a full year at monthly temporal resolution, and at high depth

resolution (~20 samples, ~12 of them across the photic zone). These measurements were

performed as part of the SoMMoS (Southeastern Mediterranean Monthly cruise Series)

campaign, which compared an open-ocean station with one at the edge of the continental

shelf (Figure 1A). Additional studies from this cruise series will focus on the carbonate

system (Juntao et al. *in prep*), nutrient dynamics (Ben Ezra et al., 2021), coccolithophore

dynamics (Keuter et al. *in prep*) and a detailed comparison of the offshore and coastal

stations (Krom et al. *in prep*). Together, these studies provide a baseline allowing improved

interpretation for future research from the two long-term ocean observatories recently

established in the EMS: the DeepLev (Katz et al., 2020) and THEMO (Diamant et al., 2020),

as well as comparison to long-term monitoring activities (Ozer et al., 2017; Rahav et al.,

2019; Sisma-Ventura et al., 2021).

Figure 1: Overview of the SoMMoS sampling locations context. A) A map of the Levantine Basin

(EMS) showing the location of the THEMO-2 sampling site (the focus of this study), as well as of

additional observatories and locations of long-term studies discussed here: THEMO-1, DeepLev (Katz

et al., 2020), N-1200 (Rosenberg et al., 2020), TB1000 (Yogev et al., 2011) and the Haifa section

transect cruises (Ozer et al., 2017). At the bottom right corner an inset of the EMS with a black frame

indicating the study site. B) Changes in satellite-derived sea surface temperature (orange) and sea

surface chlorophyll a (green) from the 1 km^2 region surrounding THEMO-2 between 2017-2020. Data

were extracted from the Copernicus Marine Environment Monitoring Service (CMEMS,

https://marine.copernicus.eu/). The period of this study is shaded.

2 Materials and Methods

2.1 Cruises and sample collection

Water samples were collected as part of the SoMMoS project during twelve cruises, from

February 2018 until January 2019. Each cruise took samples at two stations: THEMO2, an

open ocean station at a depth of ~1500 m (~50 Km from the coast, 32.820 N, 34.380 E), and

THEMO1 which is positioned at the edge of the continental shelf at ~125m depth (~10 Km

from the coast, 33.040 N, 34.950 E). THEMO2 was sampled from 12:00-22:00 (local time),

while THEMO1 was sampled during the night, typically 00:00-02:00 (local time). Here, we

will present information from data collected at the offshore THEMO2 station. Samples were

collected using a 12-bottle rosette with 8 L Niskin bottles. Samples were collected at 20-24

depths across the entire water column (11-12 bottles between the surface and 200 m, which

we define here as the photic zone). Sampling depths were selected based on real-time data of

Conductivity, Temperature, Depth (CTD) profiler (Seabird 19 Plus) from the down-cast

before each sample collection in the up-cast, an oxygen optode (on some cruises) and a

fluorescence meter (Turner designs, Cyclops-7). The continuous data was processed using the

Sea-Bird data conversion software, and minimized using bin averaging. One bin of data lines

was defined as a change of 1 decibar (db) between each bin. The first two meters of

measurements were compiled together to account for sensors and CTD pump adjustment time

and rosette depth while at sea surface. Each cast allocated water for analysis in the following

order to give priority to the more 'sensitive' parameters: dissolved methane, pH, alkalinity,

dissolved inorganic carbon, inorganic nutrients, total and dissolved organic carbon, cell

count, primary and bacterial production, DNA and algal pigment markers. Pre-filtered

inorganic nutrient samples were analyzed fresh (unfrozen) the day after the cruise, using a

SEAL AA-3 autoanalyzer system, and are described in detail elsewhere (Ben-Ezra et al.,

2021). A summary of all of the currently-available measurements can be found in

Supplementary Table S1, and the BCO-DMO (acronym: SoMMoS) and ISRAMAR

(https://isramar.ocean.org.il/isramar2009/) databases. Mixed layer depth (MLD) was

calculated using a temperature difference of $\Delta 0.3$ °C (Mena et al., 2019). Calculations based

on a density difference of $\Delta 0.15$ kg/m^3 yielded similar results. During several months

(February-April 2018 and January 2019), the density plots revealed a progressive increase in

density without a clear pycnocline but with multiple "bumps", indicative of water column

instability down to below 200 m (Supplementary Figure S1). At these times, the MLD

calculations based on a defined difference in temperature or salinity from the surface

preclude a robust estimate of the mixed layer depth, as they may underestimate the actual

values. Based on these calculations, we divide the study period into 'a generally mixed

period' during January-April (winter/spring), and a 'stratified period' during May-December

(summer/autumn, Table 1). Estimates based on the vertical distribution of inorganic nutrient

concentrations suggest the mixing period may have begun as early as November 2018 (Ben-

Ezra et al., 2021). We note that the monthly sampling resolution likely to 'miss' short-lived

deep mixing events, as can be observed from mooring operations (e.g. (Gunn et al., 2020).

2.2 Bacterial and primary productivity

Heterotrophic prokaryotic productivity (hereafter referred to as bacterial productivity, BP)

was estimated using the ^{3}H-leucine incorporation method (Simon and Azam, 1989).

Triplicate 1.7 ml of ocean water were taken from each sampled depth and incubated with a

7:1 mixture of 'cold' leucine and 'hot' ^{3}H-leucine (final concentration 100 nmol leucine L^{-1})

for 4 h at room temperature in the dark immediately after sampling. Preliminary experiments

show that this was a saturating level of leucine in the offshore water of the SE Mediterranean

Sea. After incubation, incorporation was terminated by adding 100 µl trichloroacetic acid

(TCA). As a negative control for non-specific binding, another set of triplicates were sampled

from a surface layer and treated with TCA immediately after the addition of the radioactive

tracer. At the end of each cruise, the samples were processes using the micro-centrifugation

protocol and 1ml scintillation cocktail (ULTIMA-GOLD) was added to all samples before

counted using TRI-CARB 2100 TR (PACKARD) scintillation counter. A conversion factor

of 3 kg C per mole of leucine incorporated and an isotopic dilution of 2.0 were used to

calculate the C incorporated (Simon and Azam, 1989).

Net daily photosynthetic carbon fixation rates were estimated using the ^{14}C incorporation

method (Nielsen, 1952), with several modifications (Hazan et al., 2018). Triplicate 50 ml

samples were taken from each depth within the photic zone and from one aphotic depth using

sterile vials and kept at surface light and temperature conditions. The 'dark' sample served as

blank and was kept under the same temperature as the 'light' samples. Radioactive spiking

was done at ~08:00 AM the day following of the cruise in order to start a 24h incubation for

all samples (including those collected at THEMO-1 station, not shown) at the same time.

Early work by Letelier and colleagues (1996) at station HOT showed that prolonged on-deck

incubations, similarly to the protocol used in this study, may result in underestimated PP rates

as it cannot precisely mimic the temperature and illumination levels in-situ. Our preliminary

tests concur with this conclusion and found that ashore incubations underestimate PP rates by

up to ~20% compared to incubations onto a mooring rope tied to the ship (Figure S2).

Samples were spiked with 50µl (5 µCi) of NaH$^{14}CO_3$ tracer and were incubated for 24 h

under 3 light regimes: surface illumination (samples from the upper mixing depths), 50%

illumination (samples from below the mixing depth to the DCM) and ~1% illumination

(samples from the DCM and below). Shading was performed using neutral density nets, thus

changing light intensity but not spectral properties. Water samples were then filtered through

GF/F filters (0.7 μm nominal pore size, 25 mm diameter) using low vacuum pressure (< 50

mmHg) and rinsed 3 times with filtered sea water. Filters from each sample were then put in

scintillation vials where 50 μl of 32%HCl solution was immediately added in order to remove

excess ^{14}C-bicarbonate and kept overnight for incubation. After incubation 5 mL scintillation

cocktail (ULTIMA-GOLD) was added to the samples and counted using TRI-CARB 2100

TR (PACKARD) scintillation counter. Three random aliquots were counted immediately

after the addition of the radiotracer (without incubation) with ethanolamine to serve as added

activity measurements.

2.3 Picophytoplankton abundance using flow-cytometry

Triplicates water samples (1.5 ml) were collected from each sampling depth, put in cryo-vials

(Nunc), and supplemented with 7.5 μl 25% glutaraldehyde (Sigma). Vials were incubated in

the dark for 10 min, flash-frozen in liquid nitrogen, and stored in -80 ºC freezer. Before

analysis, samples were thawed in the dark at room temperature. Each sample was run twice

on a BD Canto II flow-cytometer with 2 μm diameter fluorescent beads (Polysciences,

Warminster, PA, USA) as a size and fluorescence standard. In the first run three types of

phytoplankton cells were identified based on their natural auto-fluorescence:

Prochlorococcus, *Synechococcus* and picoeukaryotes. Cells were differentiated based on cell

chlorophyll (Ex482nm/Em676nm, PerCP channel) and phycoerythrin fluorescence

(Ex564nm/Em574nm), and by the size of cell (forward scatter). Before the second FCM, run

samples were stained with SYBR Green I (Molecular Probes/ ThermoFisher) to enable

counting followed by detection at Ex494nm/Em520nm (FITC channel). This provided counts

of the total bacterial population (phytoplankton + heterotrophic bacteria and archaea) as well

as a distinction between cells with High or Low DNA content (not shown). Data were

processed using FlowJo software. Flow rates were determined several times during each

running session by weighing tubes with double-distilled water, and counts of the standard

beads were used to verify a consistent flow rate.

2.4 Algal pigment markers

Eight litters of seawater were collected from all photic sample depths and one from a dark

depth (depth varies between cruises). Water was filtered onto GF/F filters (0.7 μm nominal

pore size, 47mm diameter, Waters) using a peristaltic pump until either all 8 L were filtered

or the filter became blocked, in which case the volume filtered was recorded. Filters were

placed in cryo-vials and flash frozen in liquid nitrogen until they could be stored in a -80 ºC

freezer. Pigments were extracted in 1ml 100% methanol for 3 h at room temperature and

clarified using syringe filters (Acrodisc CR, 13 mm, 0.2 μm PTFE membranes, Pall Life

Sciences). Total chlorophyll was measured spectrophotometrically using a NanoDrop 2000c

(Thermo Sciences) at 632, 652, 665 and 695 nm, and the concentration of chlorophyll a was

calculated (Ritchie 2008). Ultra high-pressure Liquid Chromatography (UPLC) was

performed on an ACQUITY UPLC system (Waters) equipped with a photodiode array

detector. A C8 column (1.7 μm particle size, 2.1 mm internal diameter, 50 mm column

length, ACQUITY UPLC BEH, 186002877) was used. The chromatography method was

adapted for UPLC from the LOV method (Hooker et al., 2005). Samples were preheated to

30 °C and column to 50 °C before each run. Running buffers were a 70:30 mixture of

methanol and 0.5M ammonium acetate (buffer A) and 100% methanol (buffer B). The

program consisted of an isocratic run using a 80:20 mixture of buffers A:B for 2min,

followed by a linear gradient to 50:50 for 7 minutes and an increase to 100% solvent B. The

flow rate was 0.5ml/min. Pigment standards from DHI (Denmark) were used to identify the

UPLC peaks (chlorophyll a, divinyl-chlorophyll a, chlorophyll b, chlorophyll c2, zeaxanthin,

beta-carotene, diatoxanthin, fucoxanthin, peridinin, 19'- butanoyloxyfucoxanthin and 19'-

hexanoyloxyfucoxanthin). Due to potential degradation of the pigment standards, we present

the total chlorophyll measured spectrophotometrically and the pigment ratios within each

UPLC run.

3 Results

3.1 Physical and chemical properties of the water column

Between January 2016 and December 2019, a clear pattern was observed in satellite-derived

sea surface temperature at the THEMO-2 location (Figure 1B), which was mirrored during

the monthly cruise measurements (February 2018-January 2019, Figure 2A). The measured

in-situ sea surface temperatures increased by $> 11°C$ from the winter minimum of ~17.9 °C to

the summer maximum of 29.1 °C (Figure 2A, Supplementary Figure S3, Table 1). Sea

surface salinity also increased from a winter minimum of ~39.3 psu to a summer maximum

of 39.8 psu (Figure 2B, Supplementary Figure S3). Both temperature and salinity minima

were higher than the climatological minima (~15.2 °C and 38.9 psu, measured between 2002-

2020 (Herut et al., 2020), suggesting that the sampling period represents a relatively warm

and salty year. The temporal changes in sea surface temperature and salinity led to

differences in the water density profiles (Figure S1), indicative of stratification of the upper

water layer between May and December 2018 (Mixed Layer Depth = 15-49 m) and mixed

between February-April 2018 and January 2019 (MLD not determined, see materials and

methods) (Table 1). Inorganic nutrient (NO_3+NO_2) concentrations began to increase in the

mixed layer already during November, suggesting that the stratification had started to erode

earlier than observed based on the density profiles, possibly due to short-term mixing events

(e.g. (Gunn et al., 2020).

Dissolved oxygen concentrations were at or above 100% saturation throughout the whole

photic layer (0-200 m), ranging from ~180 µmol L^{-1} to ~240 µmol L^{-1} (Figure 2C). Where,

soluble reactive phosphorus (SRP) concentrations at the surface water were at or close to the

limit of detection ~0.006 µM, (~6 nM) throughout the year, whereas nitrate+nitrite (NOx)

were close to the detection limits from August to November (limit of detection 0.013 µM) but

reached 0.3-0.5 µM during the mixed period (Table 1, and see Ben Ezra et al., 2021 for more

information).

Table 1: Median and range of major measured oceanographic parameters during the mixed

and stratified periods. BDL – Below detection limit. ND – not determined (see materials and

methods and Supplementary Figure S1 for details). Depth-integrated values are from 0-200m.

Parameter	Mixed period (Jan-Apr)	Thermally Stratified period (May-Dec)
SST (°C)	18.6 (17.9-19.6)	24.2 (22.1-29.1)
Mixed layer depth (m)	ND	26.5 (15-49)
DCM depth (m)	105 (67-125)	100 (100-135)
Surface (10 m) PO_4 [μM]	BDL ($\leq$0.006)	BDL ($\leq$0.006)
Surface (10 m) NOx [μM]	0.3 (0.3-0.6)	0.1 (0.03-0.5)
Surface Si [μM]	0.8 (0.8-1)	0.7 (0.6-1)
Chl a [mg m^{-2}]	14.6 (8.6-17.8)	8.8 (5.4-24.6)
Synechococcus [cells m^{-2}]	1×10^{12} (5.8×10^{11}-1.6×10^{12})	5×10^{11} (5×10^{10}-8.5×10^{11})
Prochlorococcus [cells m^{-2}]	7.1×10^{11} (2.5×10^{11}-2.2×10^{12})	2.3×10^{12} (1×10^{12}-3.2×10^{12})
Pico-eukaryotes [cells m^{-2}]	1.7×10^{11} (BDL-7.67×10^{11})	8.6×10^9 (7.2×10^7-2.3×10^{11})
Heterotrophic prokaryotes [cells m^{-2}]	6.6×10^{13} (6×10^{13}-1.03×10^{14})	5.4×10^{13} (3.1×10^{13}-1.1×10^{14})
PP [mgC m^{-2} d^{-1}]	113 (69-142)	60 (48-111)
BP [mgC m^{-2} d^{-1}]	54 (32-71)	18 (7-27)

3.2 Chlorophyll a, Primary Productivity and Bacterial Productivity

While satellite-derived sea surface chlorophyll *a* showed a clear seasonal cycle over three years, with lower levels during the stratified period (Figure 1B), the entire photic zone displayed more complex evolution in depth and chlorophyll concentrations (Figure 3A, B). A prominent Deep Chlorophyll Maximum (DCM) was observed year-round, ranging from depths of 60 m (February 2019) to 120 m (April 2019). The concentration of total chlorophyll a at the DCM was always higher than that measured at the surface, and ranged between ~0.15 μg L^{-1} during February-July, decreasing to ~0.07 μg L^{-1} during August-November. These values are within the range or somewhat lower than those measured in other studies in the EMS (~0.09-0.42 μg L^{-1}) (Christaki et al., 2001; Yacobi et al., 1995), and are ~50-80% lower than at BATS (Steinberg et al., 2001). Due primarily to the changes in chlorophyll a at the DCM, the integrated chlorophyll did not follow the same temporal variability as observed with the sea-surface satellites measurements (Figure 1B).

Figure 3: Seasonal changes in depth-resolved (left panels) and depth-integrated (right panels) chlorophyll *a* (A), primary productivity (B) and bacterial productivity (C). Black dots in panels A-C represent sampling points from each cruise. See Supplementary Figure S4 for the full depth profile of BP to ~1,400m.

Primary productivity (PP) was highest at the surface most of the year (1-3 mg C m^{-3} d^{-1}),

declining with depth, with no observable maximum during most of the year at the DCM

(0.08-0.47 mg C m^{-3} d^{-1}, Figure 3B). This is consistent with the chlorophyll maximum in

ultra-oligotrophic oceans being decoupled from the PP maximum (Lazzari et al., 2012). An

exception to this decoupling was observed during December 2019 and January 2020, where a

peak of PP was observed at around 75 m (0.93-0.96 mg C m^{-3} d^{-1}), slightly above the DCM

(Figure 3B). The observed values at the surface are within the range or somewhat lower than

previously measured values in the EMS (typically around ~2 mg C m^{-3} day^{-1}, but with values

as high as ~18 mg C m^{-3} day^{-1} recorded), (Hazan et al., 2018; Rahav et al., 2013). The

integrated PP values ranged between 69-142 mg C m^{-2} d^{-1} during the mixing period, and from

48 to 113 mg C m^{-2} d^{-1} during the stratified months (Figure 3B). These values are within the

ranges observed in other studies of this region (reviewed in (Berman-Frank and Rahav, 2012;

Siokou-Frangou et al., 2010) and discussed below). Our measurements result in an annual PP

of ~32 gC m^{-2}, which is lower by ~50% than most estimates from the EMS (Boldrin et al.,

2002; Psarra et al., 2000). These annual estimates highlight the ultra-oligotrophic

characteristics of the easternmost Levantine Basin.

During most of the year BP was highest at the surface (~0.2-0.5 mg C m^{-3} d^{-1}, Figure 3C,

Supplementary Figure S4), within the range of previous observations at this area (Rahav et

al., 2019; Tanaka et al., 2007; Van Wambeke et al., 2000). However, BP differed from PP in

several aspects. First, a significant increase in BP was observed during April, with the highest

rate observed at ~110 m where the DCM was detected. Indeed, a smaller secondary peak in

BP was observed at a depth corresponding to the DCM throughout the year (Figure 3C).

Second, surface BP increased during summer from a minimum in May to a maximum in

September-October, before decreasing again. Third, during January 2019 a large increase was

observed in BP, which was more-or-less homogenously distributed throughout the water

column down to ~200 m. Depth-integrated BP ranged from a maximum of ~70 mg C m^{-2} d^{-1}

during April 2018 and January 2019 to a minimum of ~20 mg C m^{-2} d^{-1} during May-

December 2018, within the range of previous measurements ~10-45 mg C m^{-2} day^{-1} (e.g.

(Christaki et al., 2011; Robarts et al., 1996; Van Wambeke et al., 2000).The integrated PP to

BP ratio also differed between the mixed and stratified periods. Where, during the mixed

season the PP:BP ratio was ~3:1, while during the stratified period it reached ~5:1.

3.3 Phytoplankton abundance and specific functional groups

Picocyanobacteria (*Prochlorococcus* and *Synechococcus*) were the most abundant

picophytoplankton cells throughout the year (Figure 4, Supplementary Figure S5).

Prochlorococcus was more abundant during the stratified period, and predominated below

the mixed layer (Figure 4A). Based on the ratio of divinyll chlorophyll a to total chlorophyll a

(Figure 5A), *Prochlorococcus* contributed up to ~45% of the total phytoplankton biomass at

the DCM during late fall (Figure 5A). *Synechococcus* was more abundant during the mixed

period and in the surface waters (Figure 4B). Pico-eukaryotes were present at up to 10^4 cells

ml^{-1} during February-June (Figure 4C), but there was no clear group-specific signal in the

tested photosynthetic pigments during this period (Figure 5B-D), and thus the specific pico-

eukaryotic groups present at this time could not be identified. In contrast, the presence of 19-

hexanoyloxyfucoxanthin and 19-butanoyloxyfucoxanthin (19-Hex+19-But), primarily during

August to November (Figure 5B), suggests the presence of haptophytes, and this was

corroborated with microscopic identification of coccolithophores (see a more detailed

discussion in Keuter et al*, in prep*). Finally, peridinin and fucoxanthin peaked during January,

suggesting the presence of dinoflagellates and diatoms, respectively (Figure 5C,D).

Figure 4: Annual dynamics of monthly measured picophytoplankton and heterotrophic bacterial abundance: *Prochlorococcus* (A), *Synechococcus* (B), PicoEukaryotes (C) and heterotrophic bacteria (D) at the THEMO2 station. Black dots represent sample point from each cruise. No data were available from the March cruise for total microbial counts. Note the differences in the color scale for each plot. Dashed line represents the DCM. See Supplementary Figure S5 for the full depth profiles to ~1,400m.

Throughout the year, total prokaryotic microbial counts (Sybr-stained cells, thus including

cyanobacteria, heterotrophic bacteria and archaea) ranged between 3×10^5 to 6×10^5 cells ml^{-1},

with higher values observed during February and May (Figure 3SB). Heterotrophic bacteria

were much more abundant than the combined phototrophs, the latter forming 2.4-8.5% of the

total Sybr-stained cell counts (Figure 4D). Below the photic layer, cell counts were typically

lower, ranging from 1.9×10^4-4.5×10^5 cells ml^{-1} (Supplementary Figure S5).

Figure 5: Monthly changes in the ratios of major accessory pigments to Total chlorophyll *a*. A) divinyl chlorophyll *a* (DVChl-a). B) Combined 19-hexanoyloxyfucoxanthin and 19-butanoyloxyfucoxanthin (19-Hex+19-But). C) Fucoxanthin (Fuco). D) Peridinin. Black dots represent sampling depths. Dashed line represents the DCM.

4 Discussion

4.1 How oligotrophic is the EMS compared to other oligotrophic oceans?

The EMS has previously been suggested to be one of the most oligotrophic marine systems

on Earth, including a claim for the "deepest Secchi-depth world record" - 53 m during

summertime (Berman et al., 1984). A comparison of published values of PP, including those

measured in this study, supports this notion, with the median integrated PP being ~66% lower

than Bermuda, the Red Sea and the Western Mediterranean Sea (WMS), and ~80% lower

than station ALOHA (HOT) (Figure 6A). Our estimates of PP during the SoMMoS cruises

are among the lowest in the EMS (Figure 6A), yielding an annual PP of ~32-39 gC m^{-2} y^{-1},

although similar values have previously been reported (Dugdale and Wilkerson, 1988). The

annual PP during the SoMMoS cruises is approximately half of that reported above the

continental slope of the Cretan Sea, ~59 gC m^{-2} y^{-1}, (Psarra et al., 2000), as well as in other

more western locations in the EMS, ~62 gC m^{-2} y^{-1} (Boldrin et al., 2002). It should be noted,

though, that our incubation approach do not precisely mimic the *in-situ* illumination and

temperature at the time of sampling, which, based on our previous work, may underestimate

the actual PP rates by up to ~20% (Figure S2). Under these circumstances, the annual PP we

report may reach ~39 gC m^{-2} y^{-1}, approximately ~33% lower than previously reported from

the EMS (Psarra et al., 2000; Boldrin et al., 2002). Moreover, comparison between different

studies that used similar, yet not identical, methodology for PP estimates, should be done

with care. For example, in the comparison presented in Figure 6A many of the studies

measured PP from dawn to dusk while in this study we used longer incubations which

account for the whole day (net PP). These differences may also, partly, account for the

changes between the different oceanic locations, yet even with these caveats taken into

account the PP in the EMS is still lower than other oceanic regions (Figure 6A).

The values of integrated chlorophyll (Table 1, Figure 2A) and phytoplankton cell counts are

also somewhat lower than previously-published measurements from the same region (Figure

4, Figure 7) (Berman et al., 1984; Robarts et al., 1996). Both the temperature and salinity

minima during the SoMMoS cruise series were higher than the 2002-2020 climatological

average (Herut et al., 2020), and thus we currently cannot determine whether the lower PP

values measured during this annual cruise series are due to inter-annual variability, or

whether these are due to the possible underestimation of the PP values discussed above

(section 2.2). Nevertheless, it is clear that the EMS is one of the most oligotrophic locations

on Earth, as defined using primary production, and that key seasonal dynamics are similar

between studies performed a decade apart (compare Figures 7D, E).

In addition to low PP, the EMS exhibits low sinking fluxes of particulate organic carbon

(POC). The POC flux measured from the bottom of the photic layer (180 m) at the DeepLev

mooring station, which is in the same region as the THEMO-2 site (Figure 1A, ~10Km north

of our study site, Lat: 32.820 N; 34.380 E), is ~0.2 gC m^{-2} during the summer months (May-

September 2018) and ~0.8 gC m^{-2} per year (Alkalay et al., 2020). This POC flux to the

aphotic layer is less than 2% of the PP measured during summertime (~12 gC m^{-2} in 5

months) or 2.5% from the annual PP rates (~32 gC m^{-2} y^{-1}). The fraction of PP exported as

particulate matter in the EMS is therefore lower than that reported in other oligotrophic

systems; i.e., HOT and BATS stations ~5% (Karl and Church, 2014; Steinberg et al., 2001).

This low contribution provides an additional biogeochemical implication of the extreme

oligotrophic conditions of the EMS. It also points to that the dissolved organic carbon

fraction derived from PP is rapidly recycled in the photic layer by heterotrophic/mixotrophic

bacteria as often observed in LNLC environments, as reviewed in (Santinelli, 2015). We

note, however, that estimates of export production based on nitrate+nitrate loss from the

photic zone, calculated from the same data, are significantly higher (172 mmol N m^{-2} year^{-1},

corresponding to ~13.6g C m^{-2} year^{-1} , presented in a companion paper, Ben Ezra et al.,

2021). The reason for this discrepancy is unclear, but may be due to over-estimation of the

export production rates (e.g. if part of the nitrate+nitrite was taken up by heterotrophic

organisms) and/or underestimation of the sinking carbon flux from sediment traps, and/or

non-Redfield C:N conversion factors that may prevail in the EMS. Moreover, such an

underestimation may be due to sinking flux from organisms too large to be caught in the

sediment traps such as fish or jellyfish (Edelist et al., 2020).

In contrast to PP, integrated BP in the EMS was higher than estimates at BATS and HOT,

and similar to the Red Sea (Figure 6B, which compares only measurements obtained using

the [3]H-leucine incorporation method). Yet, values obtained in the South Pacific Gyre, 43-61

mg C m^{-2} d^{-1}, (Berman-Frank et al., 2016) do come close to our measurements in the EMS.

Thus, differences in PP between oligotrophic regions are not necessarily coupled to BP

measurements. Previous studies, including from HOT and BATS, have shown that BP and PP

are decoupled across multiple time-scales (e.g hours-months) (Viviani and Church, 2017).

Similarly, PP and BP were uncoupled in the cyclonic Rhodes Gyre and the anti-cyclonic

Cyprus Eddy of the EMS (Rahav et al., 2013) and also in our study when BP increased in

surface waters between June and November, while PP rates declined (Figure 3). One

potential reason for this discrepancy is that PP is often measured on filtered samples

(particulate PP), and thus does not include the fraction of dissolved organic carbon (DOC)

fixed through PP that is released from the cells due to exudation or lysis (Viviani et al.,

2015).

When phytoplankton have sufficient carbon and light for photosynthesis yet nutrients such as

P and N are limiting, photosynthesis is uncoupled from growth and high concentrations of

DOC are often released to the environment (Berman-Frank and Dubinsky, 1999). Previous

studies from natural samples and from cultures of the numerically-dominant phytoplankton in

the EMS, *Prochlorococcus* and *Synechococcus*, suggest a potentially high but variable

fraction of released DOC (typically 20-70% of PP, with one study suggesting >90% in lab

cultures, (Roth-Rosenberg et al., 2020; Viviani et al., 2015). It is currently unclear whether

autochthonous production (that is, local PP) is sufficient to support the growth requirements

of heterotrophic bacteria, given what is known about the growth efficiency of bacteria. A

study from the Cretan Sea suggesting that this is possible if the release of DOC from

phytoplankton exceeds 40% of PP (Anderson and Turley, 2003).

The EMS is affected by coastal intrusions of relatively chlorophyll-rich waters (Efrati et al.,

2013), which could contribute also additional DOC, yet the importance of these processes in

determining BP and PP requires additional studies, as they occur on temporal and spatial

scales not captured by the SoMMoS cruise series. Such studies are also needed to understand

the relative contribution of dissolved compared to particulate carbon to export processes in

oligotrophic regions.

An additional explanation for the uncoupling between PP and BP was previously suggested

following an *in-situ* phosphorus addition experiment at an anti-cyclonic eddy in the EMS,

where orthophosphate addition led to chlorophyll *a* decrease while heterotrophic biomass and

activity increased (Thingstad et al., 2005). The authors postulated two possible scenarios to

explain their observation; 1) fast growing bacteria out-competed phytoplankton for the added

phosphorus. Then, the accumulated heterotrophic biomass was quickly channeled toward

larger consumers (the 'bypass hypothesis'). 2) luxury uptake of P, mainly by heterotrophic

bacteria (and to a lesser extent by small-size picophytoplankton), formed a phosphorus-rich

diet for grazers, resulting in increased egg production (the 'tunneling hypothesis').

Additionally, a scenario where an increase in temperature drives increased growth by

heterotrophic bacteria (Luna et al., 2012), resulting in a draw-down of inorganic nutrients,

could explain the increased BP in surface waters during summer, when no increase in PP was

observed (Figure 3).

Figure 6: **Comparative literature analysis of the integrated primary and bacterial productivity** (PP and BP, respectively) in the EMS and other well-studied oligotrophic

locations (Bianchi et al., 1999; Bonnet et al., 2011; Casotti et al., 2003; Christaki et al., 2011; Decembrini et al., 2009; Fernández et al., 1994; Gasol et al., 1998; Gaudy et al., 2003; Hazan et al., 2018; Ignatiades et al., 2002; Lemée et al., 2002; Lohrenz et al., 1988; Morán and Estrada, 2001; Moutin and Raimbault, 2002; Pedrós-Alió et al., 1999; Rahav et al., 2019; Eyal Rahav et al., 2013; Robarts et al., 1996; Siokou-Frangou et al., 2002; Van Wambeke et al., 2004; Vidussi et al., 2000; Wambeke et al., 2002; Zervoudaki et al., 2007). Box-Whisker plots show the interquartile range (25th–75th percentile) of the data set. The horizontal line within the box represents the median value. The red dashed line in the EMS boxes shows the median values for the SoMMoS cruises described here. The letters above the box-plots represent significant differences (ANOVA, $p < 0.05$) for mean values between sampling sites. The BP compilation includes only measurements obtained using the ^{3}H-leucine incorporation method. Note the different Y axis.

4.2 Who are the main phytoplankton in the EMS?

During the thermally mixed period, *Synechococcus* were more abundant than

Prochlorococcus at the upper ~100 m (Figures 4). This period was characterized by

measurable N at the surface while P concentrations were still extremely low – possibly

causing P limitation (December-July, Table 1, Ben Ezra et al., 2021). In contrast, the

extreme surface nutrient scarcity during the stratified period, when both N and P were below

the limit of detection and were potentially co-limiting (Ben Ezra et al., 2021), likely

contributed to the dominance of the small-size cyanobacterium *Prochlorococcus* over the

somewhat larger-cell *Synechococcus* (Figures 4). It is generally accepted that, as the nutrient

concentrations decrease (i.e., the water becomes more oligotrophic), so does the average size

of phytoplankton (Margalef and Kinne, 1997). Additionally, while most *Synechococcus*

genomes encode the genes required for nitrate uptake and utilization, this trait is found only

in a subset of the *Prochlroococcus* genomes (Berube et al., 2018), and single-cell analysis

suggests that more *Synechococcus* cells in nature utilize nitrate compared to *Prochlorococcus*

(Berthelot et al., 2019). Thus, the type of nutrient limitation, P or N+P, (Krom et al., 1991;

Thingstad et al., 2005; Zohary et al., 2005) may also affect the temporal dynamics of these

two clades in the water column.

The abundances of these two cyanobacterial groups also differed with depth. *Synechococcus*

was more abundant in the surface waters than *Prochlorococcus* (Fig 4). This was especially

evident from May to September when *Prochlorococcus* cells could not be identified in the

surface layer using either flow cytometry or pigment analysis (Figures 4 and 5). While the

low *Prochlorococcus* abundance at the surface could be partly due to the limit of detection of

the flow cytometer used, it is supported by the lack of observed DVChl-a (Figure 5a) and by

previous genetic analyses (Rosenberg et al., 2020). It has been suggested that

Prochlorococcus' unique photosynthetic pigments (DVChl-a and DVChl-b) provide a

competitive advantage for these organisms at deeper layers of the water column, leading to a

niche separation between these two pico-cyanobacteria (Moore et al., 1995). The observed

niche (depth) separation between *Prochlorococcus* and *Synechococcus* has been previously

observed including in the first description of *Prochlorococcus* from the North Atlantic and

North Pacific (Chisholm et al., 1988), and the Eastern South Pacific (BIOSOPE cruise,

(Huang et al., 2015)). However, this is not a universal observation. At the HOT station in the

North Pacific, *Prochlorococcus* and *Synechococcus* often show similar depth distributions,

both being more abundant at the surface, with *Prochlorococcus* more abundant numerically

and in terms of biomass (van den Engh et al., 2017). Similarly, in the Indian Ocean during

May-June 2003 *Prochlorococcus* dominated the surface waters whereas *Synechococcus* were

much less abundant (Huang et al., 2015). Indeed, a global analysis of >35,000 quantitative

measurements of *Prochlorococcus* and *Synechococcus* found that *Prochlorococcus* were

480 often more abundant than *Synechococcus* at high PAR and temperature values representative

481 of surface waters in tropical waters (Flombaum et al., 2013). Thus, the reasons why

482 *Prochlorococcus* and *Synechococcus* partitioned by both depth and season in the EMS, and

483 why this is not necessarily observed elsewhere, remain unclear. Other factors probably also

484 impact distribution, as was suggested for surface populations of *Prochlorococcus* that

485 appeared to be negatively affected by atmospheric deposition through biological (e.g.,

486 airborne viruses or other biological agents that infect the cells) and chemical (e.g., trace-

487 metals toxicity) processes (Rahav et al., 2020).

488 While the pico-cyanobacteria *Prochlorococcus* and *Synechococcus* were the most

489 numerically abundant phytoplankton at THEMO-2, when biomass is examined the

490 contribution of picoeukaryotes can be as high as that of the two pico-cyanobacteria, or even

491 higher (Supplementary Figure S6, and see below). A previous study of the diversity of pico-

492 phytoplankton in the EMS showed that pico-eukaryotes were not very common in terms of

493 their DNA sequences or flow cytometry signals, but were often dominant in terms of RNA

494 sequences, one potential proxy for photosynthetic activity (Man-Aharonovich et al., 2010). In

495 that study, pico-eukaryotes were suggested to comprise up to 60% of the photosynthetic

496 picoplankton biomass in surface waters, and were mainly composed of Haptophytes

497 (primarily *Prymnesiophytes*) and *Stramenopiles* (Man-Aharonovich et al., 2010). In

498 agreement with these observations, the photosynthetic pigment 19'-hexanoyloxyfucoxanthin

499 (19'-hex), which is considered (together with 19'-butanoyloxyfucoxanthin, or 19'-but) to be a

500 diagnostic pigment for *prymnesiophytes*, was found year-round in the water column, but was

501 more abundant relative to the total phytoplankton biomass (i.e. in relation to total

502 chlorophyll) during August-October (stratified period). Thus, both molecular evidence (Man-

503 Aharonovich et al., 2010) and biochemical evidence (pigment analysis) point to the

504 importance of prymnesiophytes in the EMS. A companion study from the SoMMoS cruises

provides an overview of seasonal dynamic and impact of calcified haptophytes

(coccolithophores) during this yearly survey (Keuter et al, *in prep*).

Interestingly, we often observed 19'-hex around or below the DCM, including at depths

where light intensities are very low and photosynthesis may not provide enough carbon or

energy to support growth (Fig 5). Recent studies have unveiled that haptophytes, including

coccolithophores, are mixotrophic - capable of acquiring prey by phagotrophy or organic

compounds by osmotrophy in addition to photosynthesis (e.g. (Avrahami and Frada, 2020;

Godrijan et al., 2020; Tillmann, 1998). Indeed, haptophytes can contribute to large fractions

of total bacteriovory in marine settings (Frias-Lopez et al., 2009; Unrein et al., 2014). Further

analysis of the DNA samples collected on the SoMMoS cruises may help identify

mixotrophy within the phytoplankton clades found at the base of the photic layer (e.g.

(Yelton et al., 2016) and help illuminate the metabolic capacities required to support

phytoplankton growth in this environment.

Pigments associated with diatoms and dinoflagellates, which are often dominant

phytoplankton groups (Andersen et al., 1996), were observed at relatively high concentrations

only during January 2019 (Figure 5C, D). At this time DVChl-*a* measurement were below the

detection limit in the water column, suggesting a succession from pico-cyanobacteria to

larger eukaryotic organisms, which occurred relatively rapidly - over several weeks. This

occurred concomitantly with a drop in surface temperature (Figure 2A) and thus a break in

stratification, an increase in nitrate+nitrite (Ben Ezra et al., 2021), and a shift in

coccolithophore population towards r-selected species such as *Emiliania huxleyi* which are

indicative for a higher nutrient regime (Keuter et al*, in prep*). While chlorophyll-*a*

concentrations increased both at the surface (including by satellite sensing, Figure 1B) and at

the DCM (Figure 3A), no concomitant increase was measured in PP (Figure 3B). Yet, a

major increase was observed in BP (Figure 3C). As described in detail in the materials and

methods, our measurements of PP likely underestimate productivity at depth, and the

pigments were observed primarily below the mixed layer. It is possible that the

phytoplankton bloom actually occurred several days before our cruise, and that the pigments

observed are from stressed or dead phytoplankton. In this case, the major increase in BP

observed at the time may represent heterotrophic bacteria utilizing dissolved and particulate

organic carbon produced during the bloom.

Figure 7: Integrated cell count time series of *Prochlorococcus* (green), *Synechococcus* (orange) and picoeukaryotes (gray) at stations HOT (A), BATS (B), the Northern Red Sea (C), and the EMS (D, E – this study and (Yogev et al., 2011) respectively). Abundance measurements were carried out using flow cytometry and are presented in a log scale. Data for stations HOT and BATS were compiled from (Malmstrom et al., 2010) and from the Northern Red Sea from the Eilat National Monitoring Program (Shaked and Genin, 2017).

4.3 Phytoplankton composition and seasonality in the EMS in comparison to HOT, BATS and the Red Sea

The north Atlantic and Pacific gyres, the northern Red Sea and the EMS are all considered Low-Nutrient Low-Chlorophyll marine systems, with phytoplankton numerically dominated by pico-cyanobacteria throughout most of the year. Nevertheless, there are differences in the compositions and the seasonal dynamics of the different phytoplankton clades among these LNLC location (Fig. 7, Sup Fig. 5). Specifically, BATS, the Red Sea and the EMS all show seasonal succession patterns in the abundance of the main phytoplankton groups, although there are subtle differences between these sites: i) The period when integrated *Synechococcus* numbers are higher than *Prochlorococcus* is consistently longer in the Red Sea and EMS (3-4 months) compared to BATS (1-2 months); ii) *Prochlorococcus* reached higher absolute cell abundances during the stratified period at BATS ($\sim 10^{13}$ cells m^{-2}) compared to the Northern Red Sea or the EMS ($\sim 2.5 \times 10^{11}$ cells m^{-2}); iii) In terms of calculated biomass, the Red Sea is dominated by pico-eukaryotes for much of the year, whereas in BATS and the EMS the contribution of *Prochlorococcus*, *Synechococcus* and pico-eukaryotes to the total biomass is similar, and the dominant groups change over the year (Supporting Fig S6).

The absolute nutrient concentrations and the identity of the limiting nutrient(s) are different between these locations. For example, BATS is typically considered P limited, whereas the

EMS is potentially P limited during winter and co-limited by N and P during summer

(Zohary et al., 2005). There may also be differences in the length of the mixed period, its

intensity, or the composition of the mixed water (e.g. deep water in the EMS have an N:P

ratio of ~28:1, compared to the canonical 16:1 Redfield ratio (Redfield, 1934). Future work

may use the observed differences in phytoplankton dynamics as a sensitive "readout" of the

system, enabling a better understanding of the oceanographic processes underlying the

differences between BATS, the Red Sea and the EMS.

In contrast to BATS, the Red Sea and the EMS, seasonality at HOT in is much less

pronounced (Figure 7). *Prochlorococcus* are about 2-orders of magnitude more abundant than

Synechococcus and pico-eukaryotes, and are also the most dominant in terms of biomass (Sup

Fig. 5). The drivers of phytoplankton temporal dynamics at HOT are thus likely

fundamentally different compared to BATS, the Red Sea and the EMS. It has been suggested

that a-periodic events such as mesoscale eddies, Fe deposition from winds, and biological

processes such as dinitrogen fixation, may be major drivers of phytoplankton dynamics at

HOT (Karl and Church, 2014) and less so in the EMS (Berman-Frank and Rahav, 2012;

Yogev et al., 2011).

5 Closing remarks

The year of data presented here show that the EMS is one of the most oligotrophic regions in

the global oceans, as estimated based on PP (32 mg C m^{-2} year $^{-1}$), chlorophyll a (median ~10

mg m^{-2}), and dominance of small-size picophytoplankton with overall low abundances

compared to other LNLC regions (in some cases ~2 orders of magnitude lower). In contrast,

BP in the EMS is higher than at HOT and BATS, and similar to the Red and Western

Mediterranean seas. This suggests that the system is overall heterotrophic, and that different

factors might limit phytoplankton and heterotrophic bacteria (e.g. (Thingstad et al., 2005).

Despite these observations, there is a clear seasonal cycle in the phytoplankton populations

and their activities, reminiscent of BATS and the Northern Red Sea (but not of HOT),

suggesting that mixing (and potentially other seasonal processes such as dust deposition)

drive similar processes across many, but not all, LNLC regions.

Our results also highlight important knowledge gaps regarding the EMS. For example, we do

not know to what extent the bacterial populations or their physiological traits (e.g. bacterial

carbon demand or growth efficiency) differ seasonally or vertically in the EMS, as well as

compared to other LNLC locations. Detailed studies are also lacking of higher trophic levels

in this region (dominant micro-grazer species, diversity, biomass, grazing rates, etc.). These

are critical for understanding ecosystem dynamics including top-down vs. bottom-up

regulations etc. Biogeochemical characterization of the EMS will also benefit from more

explicit characterization of carbon-per-cell estimates, used to derive the biomass

contributions from flow-cytometry counts (i.e. Supplementary Figure S6) can vary depending

on taxonomy and physiological state (Kirchman, 2013). Future studies of these aspects,

together with other data emerging from the SoMMoS cruise series and the ocean

observatories in the EMS, should provide the needed background for governments and other

stakeholders to employ science-based environmental policies in this rapidly changing region.

Acknowledgements

We thank the captains and crew of the R/V Mediterranean Explorer (EcoOcean) and R/V

Bat-Galim (IOLR), Oshra Yosef, Elad Rachmilovitz, Guy Sisma-Ventura for help with the

sampling. The SoMMoS ship-time was funded by the Leon H. Charney School of Marine

Sciences with help from EcoOcean and IOLR. This study was supported by grant

RGP0020/2016 from the Human Frontiers Science Program (to DS), by grant number

1635070/2016532 from the NSF-BSF program in Oceanography (NSFOCE-BSF, to DS), and

partly by the Israel Science Foundation (grants number 1211/17 to ER and BH and 996/08 to

I. B-F and B.H). This study is in partial fulfilment of the M.Sc. thesis of Tom Reich

(University of Haifa).

Author Contributions

MDK, YL, ER and DS initiated and designed study; TR, TBE, NB, DRR, OB, MDK, ER and

DS collected samples; TR, TBE, NB and AT analyzed samples with help from MDK, DA,

DRR and SG; TR, ER and DS wrote the manuscript with input from all co-authors.

References

Alkalay, R., Zlatkin, O., Katz, T., Herut, B., Halicz, L., Berman-Frank, I., Weinstein, Y.,

2020. Carbon export and drivers in the southeastern Levantine Basin. Deep. Res. Part II

Top. Stud. Oceanogr. 171, 104713. https://doi.org/10.1016/j.dsr2.2019.104713

Anabalón, V., Morales, C.E., González, H.E., Menschel, E., Schneider, W., Hormazabal, S.,

Valencia, L., Escribano, R., 2016. Micro-phytoplankton community structure in the

coastal upwelling zone off Concepción (central Chile): Annual and inter-annual

fluctuations in a highly dynamic environment. Prog. Oceanogr. 149, 174–188.

https://doi.org/10.1016/j.pocean.2016.10.011

Andersen, R.A., Bidigare, R.R., Keller, M.D., Latasa, M., 1996. A comparison of HPLC

pigment signatures and electron microscopic observations for oligotrophic waters of the

North Atlantic and Pacific Oceans. Deep. Res. Part II Top. Stud. Oceanogr. 43, 517–

537. https://doi.org/10.1016/0967-0645(95)00095-x

Anderson, T.R., Turley, C.M., 2003. Low bacterial growth efficiency in the oligotrophic

eastern Mediterranean Sea: A modelling analysis. J. Plankton Res. 25, 1011–1019.

https://doi.org/10.1093/plankt/25.9.1011

Avrahami, Y., Frada, M.J., 2020. Detection of Phagotrophy in the Marine Phytoplankton

Group of the Coccolithophores (Calcihaptophycidae, Haptophyta) During Nutrient-

replete and Phosphate-limited Growth. J. Phycol. 56, 1103–1108.

https://doi.org/10.1111/jpy.12997

Azov, Y., 1986. Seasonal patterns of phytoplankton productivity and abundance in nearshore

oligotrophic waters of the Levant Basin (Mediterranean). J. Plankton Res. 8, 41–53.

https://doi.org/10.1093/plankt/8.1.41

Behrenfeld, M.J., 2010. Abandoning sverdrup's critical depth hypothesis on phytoplankton

blooms. Ecology 91, 977–989. https://doi.org/10.1890/09-1207.1

Behrenfeld, M.J., Boss, E.S., 2014. Resurrecting the ecological underpinnings of ocean

plankton blooms. Ann. Rev. Mar. Sci. 6, 167–194. https://doi.org/10.1146/annurev-

marine-052913-021325

Berman-Frank, I., Dubinsky, Z., 1999. Balanced Growth in Aquatic Plants: Myth or

Reality?<subtitle>Phytoplankton use the imbalance between carbon assimilation and

biomass production to their strategic advantage</subtitle>. Bioscience 49, 29–37.

https://doi.org/10.1525/bisi.1999.49.1.29

Berman-Frank, I., Rahav, E., 2012. Dinitrogen fixation as a source for new production in the

Mediterranean Sea: A review. Life Mediterr. Sea A Look Habitat Chang. 199–226.

Berman-Frank, I., Spungin, D., Rahav, E., Van Wambeke, F., Turk-Kubo, K., Moutin, T.,

2016. Dynamics of transparent exopolymer particles (TEP) during the VAHINE

mesocosm experiment in the New Caledonian lagoon. Biogeosciences 13, 3793–3805.

https://doi.org/10.5194/bg-13-3793-2016

Berman, T., TOWNSEND, D., EL SAYED, S., TREES, C., AZOV, Y., 1984. Optical

transparency, chlorophyll and primary productivity in the eastern Mediterranean near the

Israeli coast. Oceanol. acta 7, 367–372.

Berthelot, H., Duhamel, S., L'Helguen, S., Maguer, J.F., Wang, S., Cetinić, I., Cassar, N.,

2019. NanoSIMS single cell analyses reveal the contrasting nitrogen sources for small

phytoplankton. ISME J. 13, 651–662. https://doi.org/10.1038/s41396-018-0285-8

Berube, P.M., Rasmussen, A., Braakman, R., Stepanauskas, R., Chisholm, S.W., 2018.

Emergence of trait variability through the lens of nitrogen assimilation in

Prochlorococcus. bioRxiv. https://doi.org/10.1101/383927

Bianchi, F., Boldrin, A., Civitarese, G., Del Negro, P., Giordani, P., Malaguti, A., Socal, G.,

Rabitti, S., Turchetto, M.M., 1999. Biogenic particulate matter and primary productivity

in the Southern Adriatic and Northern Ionian seas, in: 4th MTP-Workshop MATER. pp.

120–121.

Boldrin, A., Miserocchi, S., Rabitti, S., Turchetto, M.M., Balboni, V., Socal, G., 2002.

Particulate matter in the southern Adriatic and Ionian Sea: Characterisation and

downward fluxes. J. Mar. Syst. 33–34, 389–410. https://doi.org/10.1016/S0924-

7963(02)00068-4

Bonnet, S., Grosso, O., Moutin, T., 2011. Planktonic dinitrogen fixation along a longitudinal

gradient across the Mediterranean Sea during the stratified period (BOUM cruise).

Biogeosciences 8, 2257–2267. https://doi.org/10.5194/bg-8-2257-2011

Campbell, L., 2001. Flow cytometric analysis of autotrophic picoplankton. Methods

Microbiol. 30, 317–343.

Casotti, R., Landolfi, A., Brunet, C., D'Ortenzio, F., Mangoni, O., Ribera d'Alcalà, M.,

Denis, M., 2003. Composition and dynamics of the phytoplankton of the Ionian Sea

(eastern Mediterranean). J. Geophys. Res. Ocean. 108.

https://doi.org/10.1029/2002jc001541

Chisholm, S.W., Olson, R.J., Zettler, E.R., Goericke, R., Waterbury, J.B., Welschmeyer,

N.A., 1988. A novel free-living prochlorophyte abundant in the oceanic euphotic zone.

Nature 334, 340–343. https://doi.org/10.1038/334340a0

Christaki, U., Giannakourou, A., Van Wambeke, F., Grégori, G., 2001. Nanoflagellate

predation on auto- and heterotrophic picoplankton in the oligotrophic Mediterranean

Sea. J. Plankton Res. 23, 1297–1310. https://doi.org/10.1093/plankt/23.11.1297

Christaki, U., Van Wambeke, F., Lefevre, D., Lagaria, A., Prieur, L., Pujo-Pay, M.,

Grattepanche, J.D., Colombet, J., Psarra, S., Dolan, J.R., Sime-Ngando, T., Conan, P.,

Weinbauer, M.G., Moutin, T., 2011. Microbial food webs and metabolic state across

oligotrophic waters of the Mediterranean Sea during summer. Biogeosciences 8, 1839–

1852. https://doi.org/10.5194/bg-8-1839-2011

D'Ortenzio, F., Iudicone, D., de Boyer Montegut, C., Testor, P., Antoine, D., Marullo, S.,

Santoleri, R., Madec, G., 2005. Seasonal variability of the mixed layer depth in the

Mediterranean Sea as derived from in situ profiles. Geophys. Res. Lett. 32, 1–4.

https://doi.org/10.1029/2005GL022463

Decembrini, F., Caroppo, C., Azzaro, M., 2009. Size structure and production of

phytoplankton community and carbon pathways channelling in the Southern Tyrrhenian

Sea (Western Mediterranean). Deep. Res. Part II Top. Stud. Oceanogr. 56, 687–699.

https://doi.org/10.1016/j.dsr2.2008.07.022

Diamant, R., Shachar, I., Makovsky, Y., Ferreira, B.M., Cruz, N.A., 2020. Cross-sensor

quality assurance for marine observatories. Remote Sens. 12, 1–16.

https://doi.org/10.3390/rs12213470

Dugdale, R.C., Wilkerson, F.P., 1988. Nutrient sources and primary production in the Eastern

Mediterranean. Océanographie pélagique méditerranéenne 179–184.

Edelist, D., Guy-Haim, T., Kuplik, Z., Zuckerman, N., Nemoy, P., Angel, D.L., 2020.

Phenological shift in swarming patterns of Rhopilema nomadica in the Eastern

Mediterranean Sea. J. Plankton Res. 42, 211–219.

https://doi.org/10.1093/plankt/fbaa008

Efrati, S., Lehahn, Y., Rahav, E., Kress, N., Herut, B., Gertman, I., Goldman, R., Ozer, T.,

Lazar, M., Heifetz, E., 2013. Intrusion of coastal waters into the pelagic eastern

Mediterranean: In situ and satellite-based characterization. Biogeosciences 10, 3349–

3357. https://doi.org/10.5194/bg-10-3349-2013

Fernández, M., Bianchi, M., Van Wambeke, F., 1994. Bacterial biomass, heterotrophic

production and utilization of dissolved organic matter photosynthetically produced in the

Almeria-Oran front. J. Mar. Syst. 5, 313–325. https://doi.org/10.1016/0924-

7963(94)90053-1

Flombaum, P., Gallegos, J.L., Gordillo, R.A., Rincón, J., Zabala, L.L., Jiao, N., Karl, D.M.,

Li, W.K.W., Lomas, M.W., Veneziano, D., Vera, C.S., Vrugt, J.A., Martiny, A.C., 2013.

Present and future global distributions of the marine Cyanobacteria Prochlorococcus and

Synechococcus. Proc. Natl. Acad. Sci. U. S. A. 110, 9824–9829.

https://doi.org/10.1073/pnas.1307701110

Frias-Lopez, J., Thompson, A., Waldbauer, J., Chisholm, S.W., 2009. Use of stable isotope-

labelled cells to identify active grazers of picocyanobacteria in ocean surface waters.

Environ. Microbiol. 11, 512–525. https://doi.org/10.1111/j.1462-2920.2008.01793.x

Gasol, J.M., Doval, M.D., Pinhassi, J., Calderón-Paz, J.I., Guixa-Boixareu, N., Vaqué, D.,

Pedrós-Alió, C., 1998. Diel variations in bacterial heterotrophic activity and growth in

the northwestern Mediterranean Sea. Mar. Ecol. Prog. Ser. 164, 107–124.

https://doi.org/10.3354/meps164107

Gaudy, R., Youssara, F., Diaz, F., Raimbault, P., 2003. Biomass, metabolism and nutrition of

zooplankton in the Gulf of Lions (NW Mediterranean). Oceanol. Acta 26, 357–372.

https://doi.org/10.1016/S0399-1784(03)00016-1

Genin, A., Zarubin, M., Lindemann, Y., 2018. Deep mixing, stratification and spring

phytoplankton bloom in the northern Red Sea, in: 2018 Ocean Sciences Meeting. AGU.

Godrijan, J., Drapeau, D., Balch, W.M., 2020. Mixotrophic uptake of organic compounds by

coccolithophores. Limnol. Oceanogr. 65, 1410–1421. https://doi.org/10.1002/lno.11396

Guieu, C., Aumont, O., Paytan, A., Bopp, L., Law, C.S., Mahowald, N., Achterberg, E.P.,

Marañón, E., Salihoglu, B., Crise, A., Wagener, T., Herut, B., Desboeufs, K.,

Kanakidou, M., Olgun, N., Peters, F., Pulido-Villena, E., Tovar-Sanchez, A., Völker, C.,

2014. The significance of the episodic nature of atmospheric deposition to Low Nutrient

Low Chlorophyll regions. Global Biogeochem. Cycles 28, 1179–1198.

https://doi.org/10.1002/2014GB004852

Gunn, K.L., Beal, L.M., Elipot, S., McMonigal, K., Houk, A., 2020. Mixing of subtropical,

central, and intermediate waters driven by shifting and pulsing of the agulhas current. J.

Phys. Oceanogr. 50, 3545–3560. https://doi.org/10.1175/JPO-D-20-0093.1

Hazan, O., Silverman, J., Sisma-Ventura, G., Ozer, T., Gertman, I., Shoham-Frider, E., Kress,

N., Rahav, E., 2018. Mesopelagic prokaryotes alter surface phytoplankton production

during simulated deep mixing experiments in Eastern Mediterranean Sea waters. Front.

Mar. Sci. 5, 1–11. https://doi.org/10.3389/fmars.2018.00001

Herut, B., 2015. The National Monitoring Program of Israel's Mediterranean waters –

Scientific Report for 2013/14, IOLR Report H21/2015. IOLR.

Hooker, S.B., Van Heukelem, L., Thomas, C.S., Claustre, H., Ras, J., Barlow, R., Sessions,

H., Schlüter, L., Perl, J., Trees, C., Stuart, V., Head, E., Clementson, L., Fishwick, J.,

Llewellyn, C., Aiken, J., 2005. Second SeaWiFS HPLC analysis round-robin experiment

(SeaHARRE-2), NASA Technical Memorandum. National Aeronautics and Space

Administration, Goddard Space Flight Center.

Huang, S., Zhang, S., Jiao, N., Chen, F., 2015. Marine cyanophages demonstrate

biogeographic patterns throughout the global ocean. Appl. Environ. Microbiol. 81, 441–

452. https://doi.org/10.1128/AEM.02483-14

Ignatiades, L., Psarra, S., Zervakis, V., Pagou, K., Souvermezoglou, E., Assimakopoulou, G.,

Gotsis-Skretas, O., 2002. Phytoplankton size-based dynamics in the Aegean Sea

(Eastern Mediterranean). J. Mar. Syst. 36, 11–28. https://doi.org/10.1016/S0924-

7963(02)00132-X

Jickells, T.D., 1998. Nutrient biogeochemistry of the coastal zone. Science (80-.). 281, 217–

222. https://doi.org/10.1126/science.281.5374.217

Karl, D.M., Church, M.J., 2014. Microbial oceanography and the Hawaii Ocean Time-series

programme. Nat. Rev. Microbiol. 12, 699–713. https://doi.org/10.1038/nrmicro3333

Katz, T., Weinstein, Y., Alkalay, R., Biton, E., Toledo, Y., Lazar, A., Zlatkin, O., Soffer, R.,

Rahav, E., Sisma-Ventura, G., Bar, T., Ozer, T., Gildor, H., Almogi-Labin, A., Kanari,

M., Berman-Frank, I., Herut, B., 2020. The first deep-sea mooring station in the eastern

Levantine basin (DeepLev), outline and insights into regional sedimentological

processes. Deep. Res. Part II Top. Stud. Oceanogr. 171, 104663.

https://doi.org/10.1016/j.dsr2.2019.104663

Kimor, B., Wood, E.J.F., 1975. A plankton study in the eastern Mediterranean Sea. Mar.

Biol. 29, 321–333. https://doi.org/10.1007/BF00388852

Kirchman, D.L., 2013. Degradation of organic material. Process. Microb. Ecol. 79–98.

https://doi.org/10.1093/acprof:oso/9780199586936.003.0005

Krom, M., Kress, N., Berman-Frank, I., Rahav, E., 2014. Past, present and future patterns in

the nutrient chemistry of the eastern mediterranean, in: The Mediterranean Sea: Its

History and Present Challenges. Springer, pp. 49–68. https://doi.org/10.1007/978-94-

007-6704-1_4

Krom, M.D., Kress, N., Brenner, S., Gordon, L.I., 1991. Phosphorus limitation of primary

productivity in the eastern Mediterranean Sea. Limnol. Oceanogr. 36, 424–432.

https://doi.org/10.4319/lo.1991.36.3.0424

Lazzari, P., Solidoro, C., Ibello, V., Salon, S., Teruzzi, A., Béranger, K., Colella, S., Crise,

A., 2012. Seasonal and inter-annual variability of plankton chlorophyll and primary

production in the Mediterranean Sea: A modelling approach. Biogeosciences 9, 217–

233. https://doi.org/10.5194/bg-9-217-2012

Lemée, R., Rochelle-Newall, E., Van Wambeke, F., Pizay, M.-D., Rinaldi, P., Gattuso, J.-P.,

2002. Seasonal variation of bacterial production, respiration and growth efficiency in the

open NW Mediterranean Sea. Aquat. Microb. Ecol. 29, 227–237.

Letelier, R.M., Dore, J.E., Winn, C.D., Karl, D.M., 1996. Seasonal and interannual variations

in photosynthetic carbon assimilation at station ALOHA. Deep. Res. Part II Top. Stud.

Oceanogr. 43, 467–490. https://doi.org/10.1016/0967-0645(96)00006-9

Lohrenz, S.E., Wiesenburg, D.A., DePalma, I.P., Johnson, K.S., Gustafson Jr, D.E., 1988.

Interrelationships among primary production, chlorophyll, and environmental conditions

in frontal regions of the western Mediterranean Sea. Deep Sea Res. Part A. Oceanogr.

Res. Pap. 35, 793–810.

Luna, G.M., Bianchelli, S., Decembrini, F., De Domenico, E., Danovaro, R., Dell'Anno, A.,

2012. The dark portion of the Mediterranean Sea is a bioreactor of organic matter

cycling. Global Biogeochem. Cycles 26.

Malmstrom, R.R., Coe, A., Kettler, G.C., Martiny, A.C., Frias-Lopez, J., Zinser, E.R.,

Chisholm, S.W., 2010. Temporal dynamics of Prochlorococcus ecotypes in the Atlantic

and pacific oceans. ISME J. 4, 1252–1264. https://doi.org/10.1038/ismej.2010.60

Man-Aharonovich, D., Philosof, A., Kirkup, B.C., Le Gall, F., Yogev, T., Berman-Frank, I.,

Polz, M.F., Vaulot, D., Béjà, O., 2010. Diversity of active marine picoeukaryotes in the

Eastern Mediterranean Sea unveiled using photosystem-II psbA transcripts. ISME J. 4,

1044–1052. https://doi.org/10.1038/ismej.2010.25

Margalef, R., Kinne, O., 1997. Excellence in Ecology. Germany: Ecology Institute.

Marty, J.-C.C., Chiavérini, J., Pizay, M.-D.D., Avril, B., 2002. Seasonal and interannual

dynamics of nutrients and phytoplankton pigments in the western Mediterranean Sea at

the DYFAMED time-series station (1991–1999). Deep Sea Res. Part II Top. Stud.

Oceanogr. 49, 1965–1985. https://doi.org/10.1016/S0967-0645(02)00022-X

Marty, J.C., Chiavérini, J., 2010. Hydrological changes in the Ligurian Sea (NW

Mediterranean, DYFAMED site) during 1995-2007 and biogeochemical consequences.

Biogeosciences 7.

Mena, C., Reglero, P., Hidalgo, M., Sintes, E., Santiago, R., Martín, M., Moyà, G., Balbín,

R., 2019. Phytoplankton community structure is driven by stratification in the

oligotrophic mediterranean sea. Front. Microbiol. 10.

https://doi.org/10.3389/fmicb.2019.01698

Moore, L.R., Goericke, R., Chisholm, S.W., 1995. Comparative physiology of

Synechococcus and Prochlorococcus: influence of light and temperature on growth,

pigments, fluorescence and absorptive properties. Mar. Ecol. Prog. Ser. 259–275.

Morán, X.A.G., Estrada, M., 2001. Short-term variability of photosynthetic parameters and

particulate and dissolved primary production in the Alboran Sea (SW Mediterranean).

Mar. Ecol. Prog. Ser. 212, 53–67.

Moutin, T., Raimbault, P., 2002. Primary production, carbon export and nutrients availability

in western and eastern Mediterranean Sea in early summer 1996 (MINOS cruise). J.

Mar. Syst. 33, 273–288.

Nielsen, E.S., 1952. The use of radio-active carbon (c14) for measuring organic production in

the sea. ICES J. Mar. Sci. 18, 117–140. https://doi.org/10.1093/icesjms/18.2.117

Ozer, T., Gertman, I., Kress, N., Silverman, J., Herut, B., 2017. Interannual thermohaline

(1979–2014) and nutrient (2002–2014) dynamics in the Levantine surface and

intermediate water masses, SE Mediterranean Sea. Glob. Planet. Change 151, 60–67.

https://doi.org/10.1016/j.gloplacha.2016.04.001

Pedrós-Alió, C., Calderón-Paz, J.-I., Guixa-Boixereu, N., Estrada, M., Gasol, J.M., 1999.

Bacterioplankton and phytoplankton biomass and production during summer

stratification in the northwestern Mediterranean Sea. Deep Sea Res. Part I Oceanogr.

Res. Pap. 46, 985–1019.

Peled, Y., Zemah Shamir, S., Shechter, M., Rahav, E., Israel, A., 2018. A new perspective on

valuating marine climate regulation: The Israeli Mediterranean as a case study. Ecosyst.

837   Serv. 29, 83–90. https://doi.org/10.1016/j.ecoser.2017.12.001

838 Pinardi, N., Masetti, E., 2000. Variability of the large scale general circulation of the

839   Mediterranean Sea from observations and modelling: a review. Palaeogeogr.

840   Palaeoclimatol. Palaeoecol. 158, 153–173. https://doi.org/10.1016/S0031-

841   0182(00)00048-1

842 Psarra, S., Tselepides, A., Ignatiades, L., 2000. Primary productivity in the oligotrophic

843   Cretan Sea (NE Mediterranean): seasonal and interannual variability. Prog. Oceanogr.

844   46, 187–204.

845 Rahav, E., Belkin, N., Paytan, A., Herut, B., 2018. Phytoplankton and bacterial response to

846   desert dust deposition in the coastal waters of the southeastern Mediterranean Sea: A

847   four-year in situ survey. Atmosphere (Basel). 9. https://doi.org/10.3390/atmos9080305

848 Rahav, E, Herut, B., Levi, A., Mulholland, M.R., Berman-Frank, I., 2013. Springtime

849   contribution of dinitrogen fixation to primary production across the Mediterranean Sea.

850   Ocean Sci. 9, 489–498.

851 Rahav, Eyal, Herut, B., Stambler, N., Bar-Zeev, E., Mulholland, M.R., Berman-Frank, I.,

852   2013. Uncoupling between dinitrogen fixation and primary productivity in the eastern

853   Mediterranean Sea. J. Geophys. Res. Biogeosciences 118, 195–202.

854 Rahav, E., Paytan, A., Mescioglu, E., Bar-Zeev, E., Martínez Ruiz, F., Xian, P., Herut, B.,

855   2020. Bio-Aerosols Negatively Affect Prochlorococcus in Oligotrophic Aerosol-Rich

856   Marine Regions. Atmosphere (Basel). 11, 540.

857 Rahav, E., Silverman, J., Raveh, O., Hazan, O., Rubin-Blum, M., Zeri, C., Gogou, A., Kralj,

858   M., Pavlidou, A., Kress, N., 2019. The deep water of Eastern Mediterranean Sea is a

859   hotspot for bacterial activity. Deep. Res. Part II Top. Stud. Oceanogr. 164, 135–143.

860   https://doi.org/10.1016/j.dsr2.2019.03.004

861 Raveh, O., David, N., Rilov, G., Rahav, E., 2015. The temporal dynamics of coastal

phytoplankton and bacterioplankton in the eastern mediterranean sea. PLoS One 10,

e0140690. https://doi.org/10.1371/journal.pone.0140690

Redfield, A.C., 1934. On the proportions of organic derivatives in sea water and their relation

to the composition of plankton. University Press of Liverpool Liverpool.

Robarts, R.D., Zohary, T., Waiser, M.J., Yacobi, Y.Z., 1996. Bacterial abundance, biomass,

and production in relation to phytoplankton biomass in the levantine basin of the

southeastern Mediterranean Sea. Mar. Ecol. Prog. Ser. 137, 273–281.

https://doi.org/10.3354/meps137273

Rosenberg, D.R., Haber, M., Goldford, J., Lalzar, M., Aharoonovich, D., Al-Ashhab, A.,

Lehahn, Y., Krom, M.D., Segrè, D., Steindler, L., Sher, D., 2020. Particle-associated

and free-living bacterial communities in an oligotrophic sea are affected by different

environmental and anthropogenic factors. bioRxiv 2020.04.20.051391.

https://doi.org/10.1101/2020.04.20.051391

Roth-Rosenberg, D., Aharonovich, D., Luzzatto-Knaan, T., Vogts, A., Zoccarato, L.,

Eigemann, F., Nago, N., Grossart, H.P., Voss, M., Sher, D., 2020. Prochlorococcus cells

rely on microbial interactions rather than on chlorotic resting stages to survive long-term

nutrient starvation. MBio 11, 1–13. https://doi.org/10.1128/mBio.01846-20

Santinelli, C., 2015. DOC in the Mediterranean Sea, in: Biogeochemistry of Marine

Dissolved Organic Matter. Elsevier, pp. 579–608.

Shaked, Y., Genin, A., 2017. Eilat IUI monitoring NMP Report 2017.

Simon, M., Azam, F., 1989. Protein content and protein synthesis rates of planktonic marine

bacteria. Mar. Ecol. Prog. Ser. 51, 201–213. https://doi.org/10.3354/meps051201

Siokou-Frangou, I., Bianchi, M., Christaki, U., Christou, E.D., Giannakourou, A., Gotsis, O.,

Ignatiades, L., Pagou, K., Pitta, P., Psarra, S., 2002. Carbon flow in the planktonic food

web along a gradient of oligotrophy in the Aegean Sea (Mediterranean Sea). J. Mar.

Syst. 33, 335–353.

Siokou-Frangou, I., Christaki, U., Mazzocchi, M.G., Montresor, M., Ribera D'Alcala, M.,

Vaque, D., Zingone, A., 2010. Plankton in the open mediterranean Sea: A review.

Biogeosciences 7, 1543–1586. https://doi.org/10.5194/bg-7-1543-2010

Sisma-Ventura, G., Kress, N., Silverman, J., Gertner, Y., Ozer, T., Biton, E., Lazar, A.,

Gertman, I., Rahav, E., Herut, B., 2021. Post-eastern Mediterranean Transient Oxygen

Decline in the Deep Waters of the Southeast Mediterranean Sea Supports Weakening of

Ventilation Rates. Front. Mar. Sci. 7. https://doi.org/10.3389/fmars.2020.598686

Steinberg, D.K., Carlson, C.A., Bates, N.R., Johnson, R.J., Michaels, A.F., Knap, A.H., 2001.

Overview of the US JGOFS Bermuda Atlantic Time-series Study (BATS): a decade-

scale look at ocean biology and biogeochemistry. Deep Sea Res. Part II Top. Stud.

Oceanogr. 48, 1405–1447. https://doi.org/10.1016/S0967-0645(00)00148-X

Tanaka, T., Zohary, T., Krom, M.D., Law, C.S., Pitta, P., Psarra, S., Rassoulzadegan, F.,

Thingstad, T.F., Tselepides, A., Woodward, E.M.S., Flaten, G.A.F., Skjoldal, E.F.,

Zodiatis, G., 2007. Microbial community structure and function in the Levantine Basin

of the eastern Mediterranean. Deep Sea Res. Part I Oceanogr. Res. Pap. 54, 1721–1743.

https://doi.org/10.1016/j.dsr.2007.06.008

Thingstad, T.F., Krom, M.D., Mantoura, R.F.C., Flaten, G.A.F., Groom, S., Herut, B., Kress,

N., Law, C.S., Pasternak, A., Pitta, P., 2005. Nature of phosphorus limitation in the

ultraoligotrophic eastern Mediterranean. Science (80-.). 309, 1068–1071.

Tillmann, U., 1998. Phagotrophy by a plastidic haptophyte, Prymnesium patelliferum. Aquat.

Microb. Ecol. 14, 155–160.

Unrein, F., Gasol, J.M., Not, F., Forn, I., Massana, R., 2014. Mixotrophic haptophytes are

key bacterial grazers in oligotrophic coastal waters. ISME J. 8, 164–176.

van den Engh, G.J., Doggett, J.K., Thompson, A.W., Doblin, M.A., Gimpel, C.N.G., Karl,

D.M., 2017. Dynamics of Prochlorococcus and Synechococcus at station ALOHA

revealed through flow cytometry and high-resolution vertical sampling. Front. Mar. Sci.

4, 1–14. https://doi.org/10.3389/fmars.2017.00359

Van Wambeke, F., Christaki, U., Bianchi, M., Psarra, S., Tselepides, A., 2000. Heterotrophic

bacterial production in the Cretan Sea (NE Mediterranean). Prog. Oceanogr. 46, 205–

216.

Van Wambeke, F., Lefèvre, D., Prieur, L., Sempéré, R., Bianchi, M., Oubelkheir, K.,

Bruyant, F., 2004. Distribution of microbial biomass, production, respiration, dissolved

organic carbon and factors controlling bacterial production across a geostrophic front

(Almeria-Oran, SW Mediterranean Sea). Mar. Ecol. Prog. Ser. 269, 1–15.

Vidussi, F., Marty, J.-C.C., Chiavérini, J., 2000. Phytoplankton pigment variations during the

transition from spring bloom to oligotrophy in the northwestern Mediterranean Sea.

Deep Sea Res. Part I Oceanogr. Res. Pap. 47, 423–445. https://doi.org/10.1016/S0967-

0637(99)00097-7

Viviani, D.A., Church, M.J., 2017. Decoupling between bacterial production and primary

production over multiple time scales in the North Pacific Subtropical Gyre. Deep Sea

Res. Part I Oceanogr. Res. Pap. 121, 132–142.

Viviani, D.A., Karl, D.M., Church, M.J., 2015. Variability in photosynthetic production of

dissolved and particulate organic carbon in the North Pacific Subtropical Gyre. Front.

Mar. Sci. 2, 73.

Wambeke, F., Christaki, U., Giannakourou, A., Moutin, T., Souvemerzoglou, K., 2002.

Longitudinal and vertical trends of bacterial limitation by phosphorus and carbon in the

Mediterranean Sea. Microb. Ecol. 43, 119–133.

Yacobi, Y.Z., Zohary, T., Kress, N., Hecht, A., Robarts, R.D., Waiser, M., Wood, A.M., Li,

W.K.W., 1995. Chlorophyll distribution throughout the southeastern Mediterranean in

relation to the physical structure of the water mass. J. Mar. Syst. 6, 179–190.

https://doi.org/10.1016/0924-7963(94)00028-A

Yelton, A.P., Acinas, S.G., Sunagawa, S., Bork, P., Pedrós-Alió, C., Chisholm, S.W., 2016.

Global genetic capacity for mixotrophy in marine picocyanobacteria. ISME J. 10, 2946–

2957. https://doi.org/10.1038/ismej.2016.64

Yogev, T., Rahav, E., Bar-Zeev, E., Man-Aharonovich, D., Stambler, N., Kress, N., Béjà, O.,

Mulholland, M.R., Herut, B., Berman-Frank, I., 2011. Is dinitrogen fixation significant

in the Levantine Basin, east Mediterranean Sea? Environ. Microbiol. 13, 854–871.

Zervoudaki, S., Christou, E.D., Nielsen, T.G., Siokou-Frangou, I., Assimakopoulou, G.,

Giannakourou, A., Maar, M., Pagou, K., Krasakopoulou, E., Christaki, U., 2007. The

importance of small-sized copepods in a frontal area of the Aegean Sea. J. Plankton Res.

29, 317–338.

Zohary, T., Herut, B., Krom, M.D., Mantoura, R.F.C., Pitta, P., Psarra, S., Rassoulzadegan,

F., Stambler, N., Tanaka, T., Thingstad, T.F., 2005. P-limited bacteria but N and P co-

limited phytoplankton in the Eastern Mediterranean—a microcosm experiment. Deep

Sea Res. Part II Top. Stud. Oceanogr. 52, 3011–3023.