Title - Negative effects of agricultural intensification on the food provisioning rate of a declining aerial insectivore

List of Authors - Daniel R. Garrett[1,2], Fanie Pelletier[1], Dany Garant[1], and Marc Bélisle[1]

[1]Département de biologie, Université de Sherbrooke, 2500 Boulevard de l'Université, Sherbrooke, Québec J1K2R1 Canada

[2]daniel.garrett@usherbrooke.ca

Abstract

The historical rise of intensive agricultural practices is hypothesized to be related to declines of

grassland and aerial insectivorous birds. Drivers of declines may also influence the overall

abundance and spatial distribution of insects within agricultural landscapes. Subsequently, the

food provisioning rate of birds breeding within more agro-intensive landscapes may be impacted.

Lower provisioning rates in agro-intensive landscapes may lead to reduced growth rate, body

condition or fledging success of nestlings but also to diminished body condition of food

provisioning adults. Results from a previous study supported this hypothesis as the fledging

success and proxies of nestling body condition were lowest for an aerial insectivore breeding in

more agro intensive landscapes. Of the multiple hypotheses put forward to explain these

correlations, one mechanism may act through variation in food provisioning rates. In this study,

we expounded on this hypothesis using data derived from the aforementioned study system and

assessed if provisioning rates to nestlings and food provisioning behavior of adults varied across

a gradient of agricultural intensification in a declining aerial insectivore, the Tree Swallow

(*Tachycineta bicolor*). We found that the hourly provisioning rate was lower in agro-intensive

landscapes, and yet travel distances were longest within less agro-intensive landscapes. Our

results highlight that, in order to maximize long term average gain rates, Tree Swallows breeding

within agro-intensive landscapes must forage with greater intensity, perhaps at a cost to

themselves, or else costs will transfer to growing broods. Our work provides further evidence

that agricultural intensification on the breeding grounds can contribute to the declines of aerial

insectivores in part through a trophic pathway.

Introduction

Populations of both grassland and farmland birds have severely declined since the 1970s (Stanton et al. 2018, Rosenberg et al. 2019, Bowler et al. 2019). These groups include many aerial insectivores, a foraging guild presenting steeper population declines than most other avian guilds (Nebel et al. 2010, Michel et al. 2016). Concurrent changes in landscape composition and avian population indices have led to the hypothesis that these declines are driven, at least partly, by the increasing use of intensive agricultural practices (Benton et al. 2003, Hallmann et al. 2014, Stanton et al. 2018, Spiller and Dettmers 2019). Indeed, agricultural practices increasing agricultural yield per unit area of cultivated land, such as row cropping and the use of both synthetic fertilizers and pesticides, have risen dramatically in the past decades (Tscharntke et al. 2005, DiBartolomeis et al. 2019, Malaj et al. 2020).

Although many aerial insectivores share a common foraging strategy and diet, the ecological diversity of this guild invalidates many of the hypothesized drivers of farmland bird declines (Stanton et al. 2018, Spiller and Dettmers 2019). For instance, increased mechanization, including the frequency of forage harvesting, results in the significant nest destruction of ground nesting farmland birds (Tews et al. 2013, Stanton et al. 2018). Yet, most aerial insectivores do not nest on the ground and thus many hypothesized mechanisms underlying their declines involve a trophic pathway. One of those mechanisms postulates that intensive agricultural practices have reduced the overall abundance and changed the species composition or nutritive value of their prey (Stanton et al. 2018, Twining et al. 2018, Spiller and Dettmers 2019). Many factors of agricultural intensification can indeed affect the abundance, species assemblages, distributions, and trophic interactions of insects (Raven and Wagner 2021, Wagner et al. 2021).

Therefore, intensive agricultural practices have the potential to affect the fitness of insectivorous

birds through food limitation.

Previous studies have indeed observed that agricultural intensification, including habitats

impactful to insect communities and abundances, correlate with changes in dietary composition

(Nocera et al. 2012, English et al. 2017, Bellavance et al. 2018), reductions in annual fitness

proxies (Benton et al. 2002, Ghilain and Bélisle 2008, Garrett et al. 2021a), and altered

population growth (Stanton et al. 2018) of several aerial insectivore species. Studies linking prey

availability to aerial insectivore breeding success have, however, often failed to observe such

relationships (Dawson and Bortolotti 2000, Dunn et al. 2011, Imlay et al. 2017, McClenaghan et

al. 2019). A lack of association between food availability and fitness proxies is potentially

explained by breeding individuals compensating for reduced food availability or quality by

altering foraging effort (Stephens et al. 2007). For altricial birds, food-provisioning is considered

one of the most costly behaviors as individuals must meet the increasing metabolic demands of

their brood while faring for themselves (Drent and Daan 1980, Wiersma 2005, Schifferli et al.

2014). Moreover, aerial insectivores are mostly short-lived species and thus life history theory

suggests impediments to breeding will have significant impacts on lifetime reproductive success

(Stearns 1992, Sæther and Bakke 2000, Berzins et al. 2020). Therefore, if compensation of

reduced prey availability is through increased foraging effort, then it potentially comes at a cost

to an individual's own survival and future reproduction (Stearns 1992, Reznick et al. 2000,

Harrison et al. 2011).

The large-scale homogenization of landscapes resulting from agricultural intensification (Benton

et al. 2003) may thus alter foraging behavior and exacerbate the stress experienced by food-

provisioning adults (Hinsley 2000, Olsson and Bolin 2014). Assuming that food-provisioning

birds are maximizing their total daily food delivery, and if food patches are indeed of lower

quality and/or more sparsely distributed within intensively cultivated landscapes, then central-

place foraging theory predicts lower gain and nest visitation rates within these landscapes

(Orians and Pearson 1979). Therefore, parents may optimize foraging strategies, potentially by

increasing foraging efforts, to compensate for lowered food provisioning rates and overcome the

consequences of nesting within such agro-intensive landscapes (Ydenberg 2007, Olsson et al.

2008). Empirical evidence for such elevated foraging workloads, often despite lower nest

visitation rates, includes foraging at greater distances and reduced nest attendance in landscapes

dominated by agriculture (Bruun and Smith 2003, Poulin et al. 2010, Catry et al. 2014, Stanton et

al. 2016, Staggenborg et al. 2017, Evens et al. 2018). Furthermore, birds whose central places are

located within more intensely cultivated landscapes present several conditions potentially related

to increased foraging costs (Evens et al. 2018), including greater oxidative stress (Stanton et al.

2017, Evens et al. 2018) and lower immunocompetency (Belant et al. 2000, Pigeon et al. 2013).

It is therefore imperative to assess the influence of landscape context on the capacity of breeding

individuals to compensate for poor foraging conditions and the resultant workload.

A recent 11-year study in southern Québec, Canada suggested that landscapes dominated by

more arable row crop monocultures (e.g. corn (*Zea mays*) or soybean (*Glycine max*)) offer

significantly less aerial insectivore prey than ones dominated by less agriculturally intensive

habitats such as forage crops, grasslands, and dairy farms (Garrett et al. 2021a). Furthermore, in

this same study, and upon controlling for local prey availability, several indices of Tree Swallow

(*Tachycineta bicolor*) breeding success remained associated with an agricultural gradient

(Garrett et al. 2021a). Breeding attempts within landscapes dominated by intensively managed

row crops resulted in reduced fledging success, longer nestling periods, and shorter fledging

wing lengths. If these agro-intensive habitats do indeed reduce the functional connectivity of

landscapes by reducing the amount and distribution of suitable foraging patches, then

provisioning central-place foragers in these landscapes potentially expend more time or energy

traveling between food patches and exploiting them, with the additional cost of having to settle

for patches of lower quality (Hinsley 2000, Bélisle 2005, Olsson and Bolin 2014, Evens et al.

2018). Moreover, under such conditions, provisioning individuals may need to devote more time

to self-feeding in order to maintain or increase their workload (Martins and Wright 1993,

Markman et al. 2002, Ydenberg 2007). Therefore, one hypothesized mechanism explaining the

results of Garrett et al. (2021a) is that swallows breeding in agro-intensive landscapes provide

food at lower rates with the consequence that nestlings experience slower growth and potentially

increased mortality.

Presented here are the results of a study evaluating the strength of evidence for this hypothesis

using the same study system as Garrett et al. (2021a). Our main objective was to evaluate how a

gradient of agricultural intensification influenced the provisioning rate and travel distances of

food provisioning female Tree Swallows. Specifically, we evaluated a series of hypotheses

grounded in central-place foraging theory (Olsson et al. 2008, Olsson and Bolin 2014).

Provisioning models can address the constraints and mechanisms underlying the food

provisioning behavior of Tree Swallows, notably because they take into account the time parents

may devote to self-maintenance (Ydenberg 2007). However, a simple central-place foraging

model, such as the marginal value theorem, likely provides the most parsimonious conceptual

framework to make sensible qualitative predictions about provisioning rates. We hence firstly

assumed the duration of each foraging bout is composed of the travel to, between, and within

foraging patches, and that the biomass of food collected while exploiting a patch increases, yet at

a decreasing rate with patch residency time (Olsson et al. 2008). Despite the uncertainty around

which fitness currency provisioning altricial birds attempt to maximize, the fact that predation

risk is likely negligible for adult foraging swallows and that agricultural intensification can

reduce prey availability, if they are to maximize total daily food delivery, provisioning swallows

may mostly be constrained by time, rather than physiology (Schoener 1971, McNamara and

Houston 1997, Ydenberg 2007, Houston and McNamara 2014). We therefore further assumed

food bolus biomass is positively related to its energy content, that traveling and foraging costs

are constant per time unit, and that nestling rearing females are attempting to maximize their

average long-term net rate of energy gains per time unit. Therefore, females whose nests are

within landscapes composed of more dispersed or poorer food patches, as expected in landscapes

dominated by agro-intensive cultures, will experience longer travel times (Appendix S1: Fig.

S1A and C) or lower instantaneous gain rates while exploiting patches (Appendix S1: Fig. S1B

and C), and thereby show lower food provisioning rates. Finally, shorter travel times can

counterbalance slower gain rates within food patches, resulting in different food provisioning

rates and yet similar net gain rates between contrasting landscapes (Appendix S1: Fig. 1D).

Central-place foragers should be willing to incur greater time costs to reach and exploit food

patches of a given quality in landscapes composed of more dispersed or poorer food patches

(Olsson and Bolin 2014). We therefore tested if foraging distances would be furthest within more

agro-intensive landscapes, indicating greater travel time and costs in such landscapes. We also

tested the expectation that these effects would be at least partly alleviated with increasing local

prey availability (i.e. prey availability close to the nest), but exacerbated by increasing food-

demand from the brood (e.g., larger and older broods; Martins and Wright 1993, Markman et al.

2002, Geary et al. 2020).

Methods

Study sites and nest box visits

Data were collected between 2011 and 2016 within a nest box system placed across 40 farms

throughout a gradient of agricultural intensification in southern Québec, Canada (Fig. 1, but see

Ghilain and Bélisle 2008 for details). The western portion of the study area is composed

primarily of large-scale row cropping, resulting in expanses denuded of forest cover and

dominated by corn and soybean monocultures, and to a lesser extent, other cereals (principally

wheat, *Triticum spp.*) (Bélanger and Grenier 2002, Ruiz and Domon 2009). This focus on

monocultures has led to the near ubiquitous use of fertilizers, herbicides, insecticides and

fungicides, as well as increased mechanization and removal of both surface waters and wetlands

in this region (Jobin et al. 1996). These agro-intensive landscapes transition to ones composed

primarily of pastures and forage crops (e.g. hay, alfalfa (*Medicago sativa*) and clover (*Trifolium*

spp.)) in the eastern parts of the study system and are interspersed within large areas of forest

cover and a systematic drop in pesticide detections (Poisson et al. 2021).

Ten nest boxes were placed on the field margins of each farm (total N=400) and have been

monitored every two days throughout each Tree Swallow breeding season since 2006 (Porlier et

al. 2009). The number of eggs and nestlings were counted during each nest box visit, and thus

the age and phenology of each brood is known within two days of accuracy. We captured > 99%

of adult females (during incubation) and >80% of males (during the brooding period) of each

breeding attempt. Adults were banded with an aluminum US Geological Survey band containing

a unique identification code and a series of morpho-metrics and samples were collected (Ghilain

and Bélisle 2008, Lessard et al. 2014).

Food provisioning

Food provisioning was characterized via two rates defining the amount and frequency of food

being delivered to growing broods: the total biomass of delivered food boluses per hour (bolus

biomass; mg/hr) and the number of food boluses delivered per hour (bolus delivery). These rates

were estimated during food-provisioning sessions conducted between 2011 and 2016. Sessions

consisted of two consecutive 30-min blocks in which collars were applied to each nestling within

broods aged 6, 8, 10, and 12 days old (Bellavance et al. 2018). Collars were composed of a

rubber band slid through a 5-mm long feeding tube and adjusted like a bolo-tie to capture food

boluses in the crop yet loose enough to not restrict respiration. This method has often been used

to document the diet of Tree Swallows, and has not been found to impact nestling survival

(McCarty and Winkler 1999, Johnson and Lombardo 2000, Smits et al. 2005, Bellavance et al.

2018). At the end of each block, boluses were carefully removed via forceps and the nest was

checked for discarded boluses. Boluses were then placed in sterile 50-ml conical tubes and

immediately placed in a portable freezer (< 12 hours) until being stored at -80°C. At the end of

each session, collars were removed and the starting time, the number of days past 1 January

(Ordinal date) and the presence of precipitation was recorded. The maximum temperature during

sessions was also noted at the end of each breeding season via iButtons (model DS1922L;

Embedded Data Systems, Lawrenceburg, Kentucky, USA) attached to the underside of a single

nest box on each farm that recorded hourly temperatures.

In a tandem study occurring between 2013 and 2016, the food boluses collected during food

provisioning sessions were analyzed for the presence of the principal active agents of many

pesticides (Haroune et al. 2015, Poisson et al. 2021). The methods of this study required

recording the wet mass of food boluses (± 1 mg). We subsequently used this measure to estimate

the total biomass of food boluses delivered during the one-hour food provisioning sessions.

Foraging distance

The total distance an individual travels does not necessarily provide information on how far it

ventured from its nest box. For central-place foragers, this distance will be related to the rate or

effort to return to their nest box, and thus to the rate of food provisioning. Therefore, considering

both the distance an individual travels and how far it ventured from its nest box will be more

indicative of the possible foraging costs incurred than either value by itself. Consequently, we

used GPS locations of nestling rearing females (nestling age 4-12 days) to estimate their total

hourly distance traveled (DT) and mean hourly distance from their nest box (DNB). These

metrics were assumed to be indicative of the amount of travel undergone by each individual and

therefore to reflect the amount of energy expended while food provisioning. We used hourly

travel distances to have a comparable time scale to the hourly rates estimated by the food-

provisioning sessions. Specifically, we fitted PinPoint-10 GPS tags (Lotek Wireless Inc,

Newmarket, Ontario, Canada) to 43 nestling rearing females across 14 farms in 2019 (Fig. 1,

white points). Chosen farms represented landscapes dominated by either forage and pasture

fields or the principal row-crop monocultures of our study system (i.e., corn, soybean and other

cereals), with minimal tree cover (Appendix S2: Table S1). Tags weighed between 0.9 g and

1.2 g, including the harness (Appendix S2: Fig. S1), and were fitted over the synsacrum using leg

loops (Rappole and Tipton 1991). We preprogrammed tags so GPS fixes ("SWIFT" fixes,

nominal horizontal accuracy ± 10 m) were taken every 15 min between 5:00 and 20:00 local time

(N = 60 locations) the day after deployment, allowing for a habituation period. This schedule

was the most intensive that we could use to cover the majority of the daily food provisioning

period given the battery-life (McCarty 2002, Rose 2009). Because of the amount of time elapsed

between each GPS location, we did not intend for these locations to depict the actual path that

individuals traveled throughout the day, and instead assumed that they provide enough

information to derive a proxy of the total distance an individual may have traveled. We assessed

the validity of this assumption by comparing GPS locations collected every 15 minutes to those

collected every 5 minutes for two females (Appendix S2 Fig S2 and Fig S3). We observed that

the total travel distance logically increases, and yet no general pattern was missed in terms of the

distance traveled or relative frequency distribution of distances between GPS locations and an

individual's nest box. However, recording locations at this frequency severely impacted the

longevity of the tag, resulting in the acquisition of GPS locations from only half the day when

compared to sampling every 15 minutes. Given that sampling every 15 minutes did not result in

the loss of major foraging forays, we are confident that this degree of sampling is the most

ethically possible given technological constraints. The day after data collection, individuals were

caught, and their tag removed. Once data were extracted, movements of each individual were

derived by connecting GPS locations, in chronological order, and then calculating the distance

along each segment of the resulting path using both the sp (Bivand et al. 2013) and rgeos

(Bivand and Rundel 2019) packages in the R interface (R Core Team 2019). We estimated DT

by summating the length of each segment between each point during each one-hour sampling

period. We estimated DNB by first calculating the distance between the nest and each point, and

then calculated the mean distance of these points within each one-hour block (Appendix S2:

Table S2 and Fig. S4).

Landscape characterization

Landscape characterization started with delineating habitats, including agricultural parcels, within 500 m of each nest box using orthophotos (scale 1:40 000) in QGIS (QGIS 2020) at the end of each breeding season between 2011 to 2019. This spatial scale hosts close to 80% of nestling rearing females' daily time budget. We then characterized each habitat parcel in situ, defining agricultural habitats by the cultures grown. We then reclassified habitats into one of four higher order categories represented by forest, corn and soybean, forage fields (including pastures, hay fields, alfalfa, and clover), and cereals (principally wheat). Higher order categories represent the hypothesized habitats Tree Swallows use as indicators during nest-site selection, and the principal habitats composing landscapes throughout this system (Rendell and Robertson 1990, Courtois 2020, Winkler et al. 2020).

We calculated the mean percent cover of each higher order category across all nest boxes on each farm for each breeding season, resulting in 240 separate landscapes (i.e., 40 farms x 6 years). To obtain an integrative measure of the percent cover of all higher order habitats, defined as the landscape context, we used a robust principal components analysis (PCA) for compositional data (Filzmoser et al. 2009) to assign "site scores" to each of the farms during each year. Site scores were defined by the values along the first two components of the resulting compositional PCA as fitted using the robCompositions package in R (Templ et al. 2011). The first component (Comp.1) explained 79.8% of the variance in landscape context and was correlated positively with corn and soybean cultures and negatively with both forest cover and forage fields (Fig. 2A). The second component (Comp.2) explained 16.3% of the variance and was positively correlated with forest cover, negatively with both forage and cereal covers. Site scores were assigned to each food-provisioning session throughout the study.

The study of foraging distance occurred separately and on a subset of farms than those used to

estimate food provisioning (Fig. 1). We thus repeated the landscape characterization used in the

food-provisioning study, yet for the subset of farms hosting the foraging distance study. The

resulting PCA was similar to that of the food provisioning sessions. The first component

explained 89.5% of the variance in landscape context and correlated positively with both corn

and soybean cultures, and negatively with both forest and forage cover (Fig. 2B). The second

component explained 10.8% of the variance and correlated positively with forest cover as well as

with both corn and soybean cultures, and negatively with forage and cereal cultures.

For ease of interpretation, all statistical predictions and interpretations were made in reference to

five landscape context scenarios. These scenarios represent the predominant types of landscape

contexts experienced across the agricultural gradient and are defined by the site scores of five

farms (1-5) identified in Fig. 2A and B. The scenarios for the food-provisioning sessions are: (1)

ones dominated by corn or soybean with 8.8% forest cover (corn/soybean with forest); (2) ones

denuded of forest cover and dominated by only corn or soybean (corn/soybean); (3) ones

denuded of forest cover and dominated by a mixture of corn, soybean and cereals (mixed row

crops); (4) ones with 9.0% forest cover and dominated by forage crops (forage); and (5)

landscapes with 69.0% forest cover and few cultivated parcels typically devoted to forage crops

(agro-forested). Although site score values differ between the PCA derived for the food-

provisioning and foraging distance studies, they are qualitatively similar and the above landscape

context scenarios apply to both.

The distance of GPS locations from their respective nest boxes, greater than 0.5 km from nest

boxes (~20% of locations), was positively related to the percent cover of water bodies (e.g.

rivers, wetlands and ponds) around a point (Appendix S2: Fig. S5 and Fig. S6), supporting the

contention that these features are important (Elgin et al. 2020, Berzins et al. 2020). These

habitats are however rare within our system and were considered as potential confounding

factors of the observed foraging responses. Moreover, we hypothesized that the influence of

water features located far from a nest may depend on their local availability (i.e., close to the

nest). We therefore undertook a modeling endeavor aimed at identifying the spatial extent and

the type of water features most influential to the rate of food provisioning (Appendix S3: Fig.

S1).

Due to several model convergence problems, we found numerical stability was greatest when

treating water features as open bodies of water (hitherto referred to as surface water). A local

maximum of AICc weight (w) was observed when including the percent cover of surface water

calculated within 1 km (local percent cover) and 10 km (regional percent cover) from the nest

including their interaction. These model terms were included in all models for all foraging proxy.

Surface water data were derived from vector layers acquired from the Canadian National Hydro

Network data sets (NHN 2020) and Ducks unlimited (Ducks Unlimited, 2020). All calculations

of percent cover were made using the sf (Pebesma 2018) R package.

Local prey availability

Local prey availability was represented by the dry biomass of Diptera found within insect

samples from two separate insect traps placed on each of the 40 farms between 2006 and 2016.

We focused on dry biomass of Diptera (using the number of Diptera provides qualitatively

similar results, analyses not shown, see also Rioux Paquette et al. 2013, Garrett et al 2021)

because they represent a majority of the insects found in the diets of nestling Tree Swallows

(McCarty and Winkler 1999, Johnson and Lombardo 2000, Mengelkoch et al. 2004, Twining et

al. 2018). Within our study system, 74% of the biomass and 77% of the insects found in food

boluses delivered to nestlings are Diptera (Rioux Paquette et al. 2013, Bellavance et al. 2018,

Garrett et al. 2021a). The focus of this work being on the availability of food during the nestling

period, we processed insect samples collected between 1 June and 15 July, representing the

nestling period of over 96% of first breeding attempts. Diptera within each insect sample were

removed, counted, and then placed in an oven at 50°C for over 24 hours to ensure no further

changes in mass. Sample dry mass was then recorded ($\pm$ 0.0001 g) without delay.

To obtain prey estimates more reflective of the farm vs. local scale of each trap, and avoid

punctuated events such as insect swarms, we chose to use daily model predictions of Diptera

biomass from generalized additive models (GAMs) in which raw values were regressed against

the ordinal date of sample collection for each farm and year using a Gamma distribution and log

link function. GAMs were fitted as a tensor product smoother using the mgcv package in R

(Wood 2017) with an identical degree of smoothness (k=10). Model predictions are the

accumulated biomass of Diptera over a roughly 48-hour period occurring over a 3-day window

(i.e., from the time the trap was reset on day one, the entirety of day 2, and until the content of

the trap was collected on day 3). As we wanted estimates of Diptera biomass on the day of the

food provisioning session, prey availability is represented by the predicted Diptera biomass on

the day following the food provisioning session.

Statistical methods

314   *Food provisioning*

Bolus biomass was a continuous right-skewed (conditional) response containing zeros and was

consequently modeled with generalized linear mixed effect models (GLMM) using a tweedie

distribution and a log link function (Lo and Andrews 2015). We modeled bolus delivery with

GLMM using a negative binomial distribution. In order to account for differences in sampling

effort (both 30-min blocks were completed in 86% of sessions), we included whether sessions

lasted half or an entire hour as an offset (i.e., $\log(0.5)$ or $\log(1)$, respectively) in both sets of

analyses. We further included year, farm, and brood IDs as random factors in order to account

for the data's hierarchical structure (i.e., broods nested within farms nested within years).

We took an information theoretic and multimodel approach to evaluate links between landscape

context and rates of food provisioning by comparing a set of 12 models representing biologically

relevant alternative hypotheses (Table 1 and Appendix S2: Table S3). Except for a Null model

containing only random effects, we started with a Base model including potential confounding

variables. This model contained two weather variables. First, the mean maximum temperature, as

insects require a threshold temperature before being active (Taylor 1963, Grüebler et al. 2008),

and the upper limit of foraging swallow behavior is potentially thermally constrained (Tapper et

al. 2020). Second, the presence of precipitation, as we expected precipitation to act as a

hindrance to foraging (Cox et al. 2019), and because food provisioning sessions did not last long

enough to estimate amount of precipitation. The base model also included the age of the female

(second year (SY) or after second year (ASY); Hussell 1983), time at the start of the

provisioning session as a second-order polynomial, the percent cover of surface water within

1 km and 10 km and their interaction (Elgin et al. 2020), and a proxy of the brood's food demand

(Geary et al. 2020). This brood demand proxy was represented by the multiplication of the brood

size with the brood age. We did not expect foraging rates to respond linearly with this proxy, as

the functional response likely reaches a plateau whereby food provisioning parents can no longer

increase their efforts. We therefore treated this variable as a third-order polynomial. All model

terms within Base were included in all subsequent models of the candidate set. Our principal

model of interest was one in which sites scores and their interaction were added to assess

landscape context effects (Base + Land). Because potential negative effects of foraging within

agro-intensive landscapes may be alleviated with increasing prey availability closer to nests, we

also considered models with only local prey availability (Base + Food), one with both prey

availability and site scores (Base + Food + Land), and one also including the interaction between

prey availability and Comp.1 (Base + Food*Land). We further hypothesized provisioning rates

may be influenced by brood demand and the availability of prime foraging habitat and included

four models with such relationships. These included one in which the effect of land cover and

local prey availability varied with the demand proxy (Base + Food + Land * Demand and Base +

Land + Food * Demand) or the local (1 km) percent cover of surface water (Base + Land + Food

* LSW and Base + Food + Land * LSW).

352   *Foraging distance*

Two response variables characterized the foraging distances of females, the total hourly distance

traveled (DT) and the mean hourly distance an individual was from its nest box (DNB). These

distances were highly correlated (r=0.8), yet considering both was ultimately more informative

of foraging behavior. These response variables were modeled using linear mixed effects models

(LMM). Both response variables were log-transformed to ensure normality of residuals. We

included the individual and farm IDs as random effects to account for the hierarchical structure

of the dataset (i.e., locations nested within individuals nested within farms).

We applied an information theoretic and multimodel approach to a set of six models to evaluate

hypotheses about foraging distances (Table 1 and Appendix S2: Table S3). We started with a

Null model only including random effects and a Base model including potential confounding

variables influential to foraging distances (McCarty 2002, Dreelin et al. 2018, Elgin et al. 2020).

Confounding factors included the brood demand proxy as a third-order polynomial, as well as

the percent cover of surface water within 1 km and 10 km and their interaction. We further

included proxies of local prey availability, including the ordinal date and the maximum observed

temperature during each one-hour sampling period as a second-order polynomial (Rioux

Paquette et al. 2013, Bellavance et al. 2018, Garrett et al. 2021a). Though the principal model of

interests was one including site scores and their interaction (Base + Land), we compared another

three models hypothesizing that the effects of foraging within agro-intensive landscapes may

either be alleviated or exacerbated by confounding variables. We thus considered models also

including an interaction term between Comp.1 and each potential confounder, namely brood

demand (Base + Land*Demand), local (1 km) percent cover of surface water (Base +

Land*LSW), and local prey availability (Base + Land*Food).

In all analyses, the effects of key individual variables, including interactions, were estimated via

multi-model inference whereby predictions were calculated by model-averaging with shrinkage

and shown with their 95% unconditional confidence intervals (Burnham and Anderson 2002).

All quantitative covariates were z-transformed and all analyses were performed in R version

3.6.2 (R Core Team 2019) using the glmmTMB (Brooks et al. 2017) and AICcmodavg

(Mazerolle 2020) packages. Model validation was performed following Zuur et al. (2009) using

the DHARMa package (Hartig 2020).

Results

Food provisioning

Bolus biomass

We measured the total bolus biomass delivered during 741 food provisioning sessions from 420 broods, 92% of which consisted of both 30-min blocks. Overall mean bolus biomass was 284.5 mg per hour (SD = 344.9), ranging from 208.2 mg to 319.2 mg across years (Appendix S2: Fig. S7). The Null and Base models were not supported by the data, contrary to models including information on landscape context (Appendix S2: Table S3). Compared to forage landscapes (#4 in Fig. 3B and Appendix S2: Fig. S8B), bolus biomass was 54.0% lower within agro-forested landscapes (#5), 42.8% lower in mixed row-crop landscapes (#3), and 32.8% lower within corn/soybean landscapes (#2). Unlike for bolus delivery rate (see below), we did not find that the effect of corn/soybean cover was dependent upon forest cover (#1). We found that bolus biomass was unrelated to local prey availability as measured (Table 2 and Appendix S2: Fig. S9 and S10). We further found bolus biomass increased almost linearly with brood demand (Appendix S2: Fig. S10 and S11), and was negatively correlated with the local percent cover of surface water and was greater for ASY females (Appendix S2: Fig. S12). Lastly, bolus biomass increased with both the start time and the maximum temperature of the food provisioning session, until plateauing at 14:12 and 25°C, respectively (Table 2 and Appendix S2: Fig. S10).

Bolus delivery

We conducted 1,208 food provisioning sessions from 662 different broods. Overall mean bolus delivery was 2.00 boluses per hour (SD = 2.05), ranging from 1.69 to 2.25 across years (Appendix S2: Fig. S7). While the Null and Base models were not supported by the data, several

alternative models received support (Appendix S2: Table S3). We observed substantial landscape

effects (Table 2, Fig. 3A). Compared to forage landscapes (#4 in Fig. 3A), bolus delivery rate

was 46.2% lower within agro-forested landscapes (#5), 34.5% lower in mixed row-crop

landscapes (#3), and 19.3% lower within corn/soybean landscapes (#2). The effect of

corn/soybean cover was however highly dependent upon forest cover as the presence of remnant

woodlots in landscapes highly dominated by these crops (#1) was associated with a bolus

delivery rate 11.2% lower than within forage landscapes (#4). We found that bolus delivery rate

was not related to local food availability (Appendix S2: Fig. S9 and S13). We further found that

bolus delivery rate was positively associated with brood demand (third-order) and both the start

time and the maximum temperature of the food provisioning session, until plateauing at 14:59

and 26°C, respectively (Appendix S2: Fig. S11 and S13). It also negatively correlated with the

local percent cover of surface water (Appendix S2: Fig. S14).

Foraging distance

We successfully deployed and recovered 43 GPS tags. Of these tags, 40 successfully recorded all

60 attempted locations, resulting in 2,516 swift-fixes and 631 hourly estimates of foraging

distances (Appendix S2: Table S2 and Fig. S4). The overall mean distance of each location from

an individual's nest box was 420 m (SD = 930) and 465 m (SD = 814) between each consecutive

fix. Six females recorded locations over 5 km from their nest box with a maximum distance of

9.5 km.

424   *Hourly distance traveled*

Overall mean DT was 1,857 m (SD=2,365, Appendix S2: Fig. S7). A single most predictive and

parsimonious model occurred when considering landscape context along with an interaction

between Comp.1 and local percent cover of surface water (Table 2 and Appendix S2: Table S3).

DT was longest within agro-forested landscapes (#5 in Fig. 3C) and was 75.6% further when

compared to forage landscapes (#4). Compared to forage landscapes, distances were 2.8%

shorter within mixed row-crop landscapes (#3), and yet 52.7% shorter within corn/soybean

landscapes (#2). The effect of corn/soybean cover was however highly dependent upon forest

cover, as the presence of remnant woodlots in landscapes highly dominated by these crops (#1)

was associated with a DT 34.2% shorter than within forage landscapes (#4). DT increased with

brood demand until plateauing at mean demand values (Appendix S2: Fig. S11 and S15). DT

was further associated in a quadratic fashion with both the ordinal date and maximum

temperature, peaking on ordinal date 183 (2 July) and at 26.1°C (Appendix S2: Fig. S15).

Finally, DT was shorter with increasing regional cover of water (Table S2, Appendix S2: Fig.

S16).

439   *Hourly mean distance from nest box*

Overall mean DNB was 421 m (SD=719, Appendix S2: Fig. S7). While no single model could

be considered the most predictive or parsimonious, models taking landscape context into account

were clearly better supported (Table 2 and Appendix S2: Table S3). DNB was furthest within

agro-forested landscapes (#5 in Fig. 3D) and was 65.5% further when compared to forage

landscapes (#4). Compared to forage landscapes, DNB was 34.3% further within mixed row-

crop landscapes (#3), and yet 37.4% closer within corn/soybean landscapes (#2). The effect of

corn/soybean cover was however highly dependent upon forest cover as the presence of remnant

woodlots in landscapes highly dominated by these crops (#1) was associated with DNB being

24.4% closer than within forage landscapes (#4). DNB increased significantly until plateauing at

mean values of brood demand (Appendix S2: Fig. S11 and Fig. S17). DNB was associated in a

quadratic fashion with both the ordinal date and maximum temperature, peaking on ordinal date

184 (3 July) and at 26.1°C (Appendix S2: Fig. S17). Finally, DNB was shorter with increasing

regional cover of water (Table S2, Appendix S2: Fig. S18).

Discussion

This study aimed at evaluating a hypothesized mechanism that negatively correlated the fitness

of an aerial insectivore with breeding in landscapes increasingly composed of more agriculturally

intensive monocultures (e.g., corn and soybean). This mechanism hypothesized that foragers in

agro-intensive landscapes expend more time or energy reaching food patches and traveling

between them, with the additional cost of settling for patches of lower quality, the combination

of which would result in lower food provisioning rates (Appendix S1: Fig. 1). In more

agriculturally dominated landscapes, we found that female Tree Swallows provided food to their

nestlings at lower rates within increasingly agro-intensive landscapes. The hourly rate of delivery

of both bolus biomass and bolus number were indeed lowest within more agro-intensive

landscapes and greatest within forage landscapes. Female Tree Swallows traveled most and

furthest from their nest within forest-dominated landscapes, and unexpectedly less and closer to

their nest in agro-intensive compared to forage landscapes.

The relationship between the rate at which boluses were delivered and landscape context

resembled that at which prey biomass was provided, suggesting that the biomass of individual

boluses was similar throughout the agricultural intensification gradient studied (McCarty 2002).

This outcome was corroborated by a supplemental analysis (Appendix S4: Table S1 and S2).

Furthermore, coupled with the result that females traveled furthest within predominantly forested

landscapes and to a lesser extent in forage landscapes, the relatively low variance in the biomass

of individual boluses suggests that forage landscapes result in marginally greater travel times.

Yet the increased frequency of food bolus delivery implies that foraging in forage landscapes

likely lead to more rapid loading curves, and greater gain rates, than that of agro-intensive

landscapes (Fig. 4). Such results stand, even if we assume Tree Swallows from this system do

not experience a decelerating gain curve while exploiting food patches, and instead "sift"

through the air column until reaching an individual-dependent fixed loading capacity (Appendix

S2. Fig S2).

It should be stated that our results were focused on correlations between landscape composition

and food provisioning rates and not on how these rates are correlated to measures of avian

fitness. Therefore, conclusions from these results should recognize the need to better establish

causal pathways between provisioning rates and both nestling and adult body condition, as well

as nestling survival. However, these factors are intimately linked, and despite our work being

correlative, they are ultimately proxies of some underlying mechanisms warranting future work.

For example, lower food provisioning rates within more agro-intensive landscapes may need to

be compensated by parents elongating the entire foraging period of a given day, foraging more

intensely when conditions allow, forgoing recuperation or foraging for themselves (Martins and

Wright 1993, Markman et al. 2002, Ydenberg 2007, Schifferli et al. 2014). In all cases, increases

in foraging effort may come at a cost to themselves through reduced body condition and thus

potentially impactful to their future reproductive success (Stearns 1992, Reznick et al. 2000,

Harrison et al. 2011, Lagrange 2015), as increases in effort may result in foraging parents being

in too poor of body condition for molting prior to migration (Jenni-Eiermann and Jenni 1996,

Rubolini et al. 2002). Furthermore, if parents cannot compensate for lowered food provisioning

rates, then broods may incur the cost through reduced growth rates or elevated mortality (Naef-

Daenzer et al. 1999, Tremblay et al. 2004). In fact, evidence suggest food provisioning Tree

Swallows as well as their broods already incur these types of costs. For example, parents

breeding in more agro-intensive landscapes spend less time in the nest (Lamoureux 2010,

Stanton et al. 2016), express reduced body condition (Pigeon et al. 2013, Stanton et al. 2017),

and both reduced fledging success and nestling growth (Houle et al. 2020, Garrett et al. 2021a).

Some of these natal effects are of concern as they may be carried over into later life stages

(Harrison et al. 2011).

One fundamental characteristic of our gradient of agricultural intensification, and as an

extension, most agricultural landscapes, is a loss of forest cover (Fischer and Lindenmayer

2007). Within our study area, landscapes dominated by a mixture of forest cover and agricultural

fields resulted in the lowest food provisioning rates and furthest foraging distances. The foraging

habitat of Tree Swallows, as well as of most swallow species, is frequently recognized as being

open terrestrial habitats or over large water surfaces (Evans et al. 2007, Boynton et al. 2020,

Elgin et al. 2020). Thus, forest cover may reduce landscape functional connectivity for these

species, at least with respect to foraging (Bélisle 2005, Olsson and Bolin 2014, Evens et al.

2018). In this way, as forest cover is reduced, the cover of foraging habitats, close to central

places and conducive to Tree Swallows, increases. In fact, second to forage landscapes, we found

provisioning rates were superior within agro-intensive landscapes with remnant forest cover.

Residual woodlots within these agro-intensive landscapes add to the tree lines separating fields

to increase the length of "forest" edges, where insect abundance tends to be greatest (Jokimäki et

al. 1998, Evans et al. 2003, Boetzl et al. 2020). The fledging success of Tree Swallows from the

system presented here is greater in agro-intensive landscapes containing residual woodlots than

ones denuded of forest cover (Garrett et al. 2021a). It may therefore be hypothesized that the

elevated insect abundance associated with these residual woodlots lead to greater provisioning

rates and thus greater fledging success in this landscape context.

It should be noted that during the years covered by this study, areas intensively-managed for the

growing of corn and soybean harbored elevated levels of pesticides in their surface waters,

notably several herbicides (e.g. glyphosate, atrazine, S-metolachlor, imazethapyr) and

neonicotinoid insecticides [e.g. clothianidin, thiacloprid, thiametoxam (Giroux 2019, Montiel-

León et al. 2019)]. For instance, a previous study found that, on average, exposure to active

pesticide agents within the food boluses was greatest within these landscapes (Poisson et al.

2021). The consumption or prolonged exposure to such chemicals can influence the neurological,

endocrine and immunological systems of non-target animals, including vertebrates, via "skin"

contact, breathing or consumption of food or water (Mayne et al. 2005, Mineau et al. 2013,

Lopez-Antia et al. 2015, Pisa et al. 2015). Importantly, neurotoxic insecticides can induce

anorexia, alter orientation and influence motor functions required for flight in birds (Eng et al.

2017, 2019). Foraging within agro-intensive landscapes increases exposure of foragers to these

chemicals, likely reducing their motivation and ability to forage, and thus potentially resulting in

decreased provisioning rates.

Finally, foraging is a metabolically expensive behavior, made worse when foraging bouts are

during the nestling periods (Bryant and Tatner 2008, Yap et al. 2017). Therefore, incoming food

need not only be frequent, but also nutritionally rich (Twining et al. 2018). The diversity and

abundance of insects found within food boluses of this system suggests that the nutritive value of

food varies across the agricultural gradient (Bellavance et al. 2018). Most notably, insects with

aquatic life stages (principally Ephemeroptera) are more abundant within the food boluses

retrieved from forage landscapes than from those dominated by more agro-intensive cultures.

These insects are greater in highly unsaturated omega-3 fatty acids (HUFA; (Twining et al.

2018). Nestlings with diets rich in HUFA can grow faster, gain a better body condition and have

greater immunocompetence, leading to elevated brood survival and fledging success (Twining et

al. 2018). Such effect can logically be extended to foraging adults. Subsequently, the energetic

losses of foraging may be more easily replenished within forage landscapes.

To our surprise, we found that both measures of food provisioning were not related to our

measure of local prey availability. Our measure of food availability was based on sampling over

a two-day period prior to nest box visits, while our sampling of food provisioning occurs over a

one-hour period. We believe that, despite food availability being linked to the fitness of Tree

Swallows in this system (Garrett et al. 2021a), the differing temporal scales of sampling likely

hinders our ability to confidently evaluate how local food availability is related to food

provisioning. Instead, we suggest future work explicitly focus on sampling aerial insects during

food provisioning sessions, as has been used in other similar studies (Grüebler et al. 2008,

Stanton et al. 2016, Elgin et al. 2020).

Global climate change results in increased global temperatures and predicts increases in both the

frequency and intensity of extreme weather events, such as prolonged periods of low or high

temperatures or precipitation (Rahmstorf and Coumou 2011, Wuebbles et al. 2014). Such

weather events may drastically alter the availability of insects as a food resource due to their

thermally dependent nature (Grüebler et al. 2008, Shipley et al. 2020, Garrett et al. 2021a).

Furthermore, recent evidence suggests Tree Swallows may already be foraging at some upper

thermal threshold (Tapper et al. 2020). Our results, combined with other pieces of evidence,

suggest food provisioning adults may also be foraging at some upper energetic limit inasmuch as

proxies of body condition are lower where food provisioning rates are lower (Pigeon et al. 2013,

Stanton et al. 2016, 2017). It is thus imperative that future work focuses not only on how

agricultural practices or global climate change independently impact declining avian species, but

also on how these factors may interact with one another. This importance is due to the ability of

growing broods or breeding parents to manage extreme weather is potentially lower within more

agro-intensive landscapes (Grue et al. 1997, Evans et al. 2003, Eng et al. 2019, Tapper et al.

2020, Garrett et al. 2021b). Finally, our results provide evidence that aerial insectivore declines

may be related to agricultural intensification through a trophic pathway.

571 Acknowledgments

We are indebted to the farm owners who kindly accepted to partake in our long-term study since

2004. We greatly appreciate the time and effort of the reviewers. Sincere thanks to Josiane Côté

Audet and the many graduate students, field and lab assistants who helped collect, enter, and

proof data throughout the years, notably with respect to insect sample processing. This work was

conducted under the approval of the animal care committee of the Université de Sherbrooke and

was financially supported by Natural Sciences and Engineering Research Council of Canada

discovery grants to FP, DG and MB, two team research grants from the Fonds de recherche du

Québec—Nature et technologies to FP, DG and MB, by the Canada Research Chairs program to

FP and MB, as well as by the Canadian Foundation for Innovation to FP, DG and MB, the

Canadian Wildlife Service of Environment and Climate Change Canada and the Université de

Sherbrooke.

literature cited

Agriculture and Agri-Food Canada (AAFC). 2018. Annual Crop Inventory.
https://open.canada.ca/data/en/dataset/.

Bélanger, L., and M. Grenier. 2002. Agriculture intensification and forest fragmentation in the
St. Lawrence valley, Québec, Canada. Landscape Ecology 17:495–507.

Belant, J. L., L. A. Tyson, and T. W. Seamans. 2000. Wildlife Management: Cropping to
Manage or Managing to Crop? Wildlife Society Bulletin 28:774–779.

Bélisle, M. 2005. Measuring landscape connectivity: the challenge of behavioral landscape
ecology. Ecology 86:1988–1995.

Bellavance, V., M. Bélisle, J. Savage, F. Pelletier, and D. Garant. 2018. Influence of agricultural
intensification on prey availability and nestling diet in Tree Swallows (Tachycineta bicolor).
Canadian Journal of Zoology 96:1053–1065.

Benton, T. G., D. M. Bryant, L. Cole, and H. Q. P. Crick. 2002. Linking agricultural practice to
insect and bird populations: a historical study over three decades. Journal of Applied
Ecology 39:673–687.

Benton, T. G., J. A. Vickery, and J. D. Wilson. 2003. Farmland biodiversity: is habitat
heterogeneity the key? Trends in Ecology & Evolution 18:182–188.

Berzins, L. L., R. D. Dawson, C. A. Morrissey, and R. G. Clark. 2020. The relative contribution
of individual quality and changing climate as drivers of lifetime reproductive success in a
short-lived avian species. Scientific Reports 10:1–12.

Bivand, R. S., E. J. Pebesma, and V. Gomez-Rubio. 2013. Applied spatial data analysis with R,
Second edition. Springer.

Bivand, R. S., and C. Rundel. 2019. Package 'rgeos' - Interface to Geometry Engine - Open
Source ('GEOS').

Boetzl, F. A., M. Schuele, J. Krauss, and I. Steffan☐Dewenter. 2020. Pest control potential of
adjacent agri☐environment schemes varies with crop type and is shaped by landscape
context and within☐field position. Journal of Applied Ecology 57:1482–1493.

Bowler, D. E., H. Heldbjerg, A. D. Fox, M. Jong, and K. Böhning☐Gaese. 2019. Long☐term
declines of European insectivorous bird populations and potential causes. Conservation
Biology 33:1120–1130.

Boynton, C. K., N. A. Mahony, and T. D. Williams. 2020. Barn Swallow (Hirundo rustica)
fledglings use crop habitat more frequently in relation to its availability than pasture and
other habitat types. The Condor 122:duz067.

Brooks, M. E., K. Kristensen, K. J. van Benthem, A. Magnusson, C. W. Berg, A. Nielsen, H. J.
Skaug, M. Mächler, and B. M. Bolker. 2017. glmmTMB balances speed and flexibility
among packages for zero-inflated generalized linear mixed modeling. R Journal.

Bruun, M., and H. G. Smith. 2003. Landscape composition affects habitat use and foraging flight
distances in breeding European starlings. Biological Conservation 114:179–187.

Bryant, D. M., and P. Tatner. 2008. Intraspecies variation in avian energy expenditure: correlates
and constraints. Ibis 133:236–245.

Burnham, K. P., and D. R. Anderson. 2002. Model Selection and Multimodel Inference. second.

Springer.

Canada, N. R. 2020. National Hydro Network - NHN - GeoBase Series.

626   https://www.nrcan.gc.ca/science-and-data/science-and-research/earth-

627   sciences/geography/topographic-information/geobase-surface-water-program-

628   geeau/national-hydrographic-network/21361.

629   Catry, I., A. M. A. Franco, and F. Moreira. 2014. Easy but ephemeral food: exploring the trade-

630   offs of agricultural practices in the foraging decisions of Lesser Kestrels on farmland. Bird

631   Study 61:447–456.

632   Courtois, È. 2020. Sélection De Nichoir Chez L'hirondelle Bicolore (Tachycineta Bicolor) En

633   Milieux Agricoles□: Une Trappe Écologique?

634   Cox, A. R., R. J. Robertson, Á. Z. Lendvai, K. Everitt, and F. Bonier. 2019. Rainy springs linked

635   to poor nestling growth in a declining avian aerial insectivore (Tachycineta bicolor).

636   Proceedings of the Royal Society B: Biological Sciences 286:20190018.

637   Dawson, R. D., and G. R. Bortolotti. 2000. Reproductive Success of American Kestrels: The

638   Role of Prey Abundance and Weather. The Condor 102:814–822.

639   DiBartolomeis, M., S. Kegley, P. Mineau, R. Radford, and K. Klein. 2019. An assessment of

640   acute insecticide toxicity loading (AITL) of chemical pesticides used on agricultural land in

641   the United States. PLOS ONE 14:e0220029.

642   Dreelin, R. A., J. R. Shipley, and D. W. Winkler. 2018. Flight Behavior of Individual Aerial

643   Insectivores Revealed by Novel Altitudinal Dataloggers. Frontiers in Ecology and

644   Evolution 6:1–7.

Drent, R. H., and S. Daan. 1980. The Prudent Parent: Energetic Adjustments in Avian Breeding.

Ardea 68:225–252.

Ducks   Unlimited.   2020.   Ducks   unlimited   GIS   mapping   applications.

https://www.ducks.ca/initiatives/gis-mapping-applications/.

Dunn, P. O., D. W. Winkler, L. a. Whittingham, S. J. Hannon, and R. J. Robertson. 2011. A test

of the mismatch hypothesis: How is timing of reproduction related to food abundance in an

aerial insectivore? Ecology 92:450–461.

Elgin, A. S., R. G. Clark, and C. A. Morrissey. 2020. Tree Swallow selection for wetlands in

agricultural landscapes predicted by central-place foraging theory. The Condor 122:1–12.

Eng, M. L., B. J. M. Stutchbury, and C. A. Morrissey. 2017. Imidacloprid and chlorpyrifos

insecticides impair migratory ability in a seed-eating songbird. Scientific Reports 7:1–9.

Eng, M. L., B. J. M. Stutchbury, and C. A. Morrissey. 2019. A neonicotinoid insecticide reduces

fueling and delays migration in songbirds. Science 365:1177–1180.

English, P. A., J. J. Nocera, B. A. Pond, and D. J. Green. 2017. Habitat and food supply across

multiple spatial scales influence the distribution and abundance of a nocturnal aerial

insectivore. Landscape Ecology 32:343–359.

Evans, K. L., R. B. Bradbury, and J. D. Wilson. 2003. Selection of hedgerows by Swallows

Hirundo rustica foraging on farmland: the influence of local habitat and weather. Bird Study

50:8–14.

Evans, K. L., J. D. Wilson, and R. B. Bradbury. 2007. Effects of crop type and aerial invertebrate

abundance on foraging barn swallows Hirundo rustica. Agriculture, Ecosystems &

Environment 122:267–273.

Evens, R., N. Beenaerts, T. Neyens, N. Witters, K. Smeets, and T. Artois. 2018. Proximity of

breeding and foraging areas affects foraging effort of a crepuscular, insectivorous bird.

Scientific Reports 8:1–11.

Filzmoser, P., K. Hron, and C. Reimann. 2009. Principal component analysis for compositional

data with outliers. Environmetrics 20:621–632.

Fischer, J., and D. B. Lindenmayer. 2007. Landscape modification and habitat fragmentation: a

synthesis. Global Ecology and Biogeography 16:265–280.

Garrett, D. R., F. Pelletier, D. Garant, and M. Bélisle. 2021a. Combined influence of food

availability and agricultural intensification on a declining aerial insectivore.

bioRxiv:10.1101/2021.02.02.427782.

Garrett, D. R., F. Pelletier, D. Garant, and M. Bélisle. 2021b. Interacting effects of cold snaps,

rain, and agriculture on the fledging success of a declining aerial insectivore. bioRxiv:1–33.

Geary, B., P. L. Leberg, K. M. Purcell, S. T. Walter, and J. Karubian. 2020. Breeding Brown

Pelicans Improve Foraging Performance as Energetic Needs Rise. Scientific Reports 10:1–

9.

Ghilain, A., and M. Bélisle. 2008. Breeding success of tree swallows along a gradient of

agricultural intensification. Ecological Applications 18:1140–1154.

Giroux, I. 2019. Présence de pesticides dans l'eau au Québec : Portrait et tendances dans les

zones de maïs et de soya – 2015 à 2017, Québec, ministère de l'Environnement et de la

Lutte contre les changements climatiques, Direction générale du suivi de l'état de l'environ.

Grue, C. E., P. L. Gibert, and M. E. Seeley. 1997. Neurophysiological and Behavioral Changes

in Non-Target Wildlife Exposed to Organophosphate and Carbamate Pesticides:

Thermoregulation, Food Consumption, and Reproduction. American Zoologist 37:369–388.

Grüebler, M. U., M. Morand, and B. Naef-Daenzer. 2008. A predictive model of the density of

airborne insects in agricultural environments. Agriculture, Ecosystems & Environment

123:75–80.

Hallmann, C. A., R. P. B. Foppen, C. A. M. van Turnhout, H. de Kroon, and E. Jongejans. 2014.

Declines in insectivorous birds are associated with high neonicotinoid concentrations.

Nature 511:341–343.

Haroune, L., R. Cassoulet, M.-P. Lafontaine, M. Bélisle, D. Garant, F. Pelletier, H. Cabana, and

J.-P. Bellenger. 2015. Liquid chromatography-tandem mass spectrometry determination for

multiclass pesticides from insect samples by microwave-assisted solvent extraction

followed by a salt-out effect and micro-dispersion purification. Analytica Chimica Acta

891:160–170.

Harrison, X. A., J. D. Blount, R. Inger, D. R. Norris, and S. Bearhop. 2011. Carry-over effects as

drivers of fitness differences in animals. Journal of Animal Ecology 80:4–18.

Hartig, F. 2020. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)

Regression Models. R package version 0.2.0.

Hinsley, S. A. 2000. The costs of multiple patch use by birds. Landscape Ecology 15:765–775.

Houle, C., F. Pelletier, M. Bélisle, and D. Garant. 2020. Impacts of environmental heterogeneity

on natural selection in a wild bird population. Evolution 74:1142–1154.

Houston, A. I., and J. M. McNamara. 2014. Foraging currencies, metabolism and behavioural

routines. Journal of Animal Ecology 83:30–40.

Hussell, D. J. T. 1983. Age and Plumage Color in Female Tree Swallows. Journal of Field

Ornithology 54:312–318.

Imlay, T. L., H. A. R. Mann, and M. L. Leonard. 2017. No effect of insect abundance on nestling

survival or mass for three aerial insectivores. Avian Conservation and Ecology

12:10.5751/ACE-01092–120219.

Jenni-Eiermann, S., and L. Jenni. 1996. Metabolic Differences Between the Postbreeding,

Moulting and Migratory Periods in Feeding and Fasting Passerine Birds. Functional

Ecology 10:62–72.

Jobin, B., J.-L. DesGranges, and C. Boutin. 1996. Population trends in selected species of

farmland birds in relation to recent developments in agriculture in the St. Lawrence Valley.

Agriculture, Ecosystems & Environment 57:103–116.

Johnson, M. E., and M. P. Lombardo. 2000. Nestling Tree Swallow (Tachycineta bicolor) Diets

in an Upland Old Field in Western Michigan. The American Midland Naturalist 144:216–

219.

Jokimäki, J., E. Huhta, J. Itämies, and P. Rahko. 1998. Distribution of arthropods in relation to

forest patch size, edge, and stand characteristics. Canadian Journal of Forest Research

28:1068–1072.

Lagrange, P. 2015. Déterminants De La Survie Et De La Dispersion De Reproduction Par Une

Approche Capture-Marquage-Recapture Chez L'Hirondelle Bicolore Au Québec.

Lamoureux, S. 2010. Impact de l'intensification agricole sur l'effort parental, la croissance et la

survie des oisillons chez l'hirondelle bicolore (Tachycineta bicolor). Université de

Sherbrooke.

Lessard, A., A. Bourret, M. Bélisle, F. Pelletier, and D. Garant. 2014. Individual and

environmental determinants of reproductive success in male tree swallow (Tachycineta

bicolor). Behavioral Ecology and Sociobiology 68:733–742.

Lo, S., and S. Andrews. 2015. To transform or not to transform: using generalized linear mixed

models to analyse reaction time data. Frontiers in Psychology 6:1–16.

Lopez-Antia, A., M. E. Ortiz-Santaliestra, F. Mougeot, and R. Mateo. 2015. Imidacloprid-treated

seed ingestion has lethal effect on adult partridges and reduces both breeding investment

and offspring immunity. Environmental Research 136:97–107.

Malaj, E., L. Freistadt, and C. A. Morrissey. 2020. Spatio-Temporal Patterns of Crops and

Agrochemicals in Canada Over 35 Years. Frontiers in Environmental Science 8:1–12.

Markman, S., B. Pinshow, and J. Wright. 2002. The manipulation of food resources reveals sex–

specific trade–offs between parental self-feeding and offspring care. Proceedings of the

Royal Society of London. Series B: Biological Sciences 269:1931–1938.

Martins, T. L. F., and J. Wright. 1993. Cost of reproduction and allocation of food between

parent and young in the swift (Apus apus). Behavioral Ecology 4:213–223.

Mayne, G. J., C. A. Bishop, P. A. Martin, H. J. Boermans, and B. Hunter. 2005. Thyroid

Function in Nestling Tree Swallows and Eastern Bluebirds Exposed to Non-Persistent

Pesticides and p, p-DDE in Apple Orchards of Southern Ontario, Canada. Ecotoxicology

14:381–396.

Mazerolle, M. J. 2020. AICcmodavg: model selection and multimodel inference based on

(Q)AIC(c).

McCarty, J. P. 2002. The number of visits to the nest by parents is an accurate measure of food

delivered to nestlings in Tree Swallows. Journal of Field Ornithology 73:9–14.

McCarty, J. P., and D. W. Winkler. 1999. Foraging Ecology and Diet Selectivity of Tree

Swallows Feeding Nestlings. The Condor 101:246–254.

McClenaghan, B., K. C. R. Kerr, and E. Nol. 2019. Does prey availability affect the reproductive

performance of Barn Swallows (Hirundo rustica) breeding in Ontario, Canada? Canadian

Journal of Zoology 97:979–987.

McNamara, J. M., and A. I. Houston. 1997. Currencies For Foraging Based on Energetic Gain.

The American Naturalist 150:603–617.

Mengelkoch, J. M., G. J. Niemi, and R. R. Regal. 2004. Diet of the nestling tree swallow. The

Condor 106:423–429.

Michel, N. L., A. C. Smith, R. G. Clark, C. A. Morrissey, and K. A. Hobson. 2016. Differences

in spatial synchrony and interspecific concordance inform guild-level population trends for

aerial insectivorous birds. Ecography 39:774–786.

Mineau, P., C. Palmer, J. Melorose, R. Perroy, S. Careas, D. R. Tobergte, and S. Curtis. 2013.

Summary for Policymakers. Pages 1–30 *in* Intergovernmental Panel on Climate Change,

editor. Climate Change 2013 - The Physical Science Basis. Cambridge University Press.

Montiel-León, J. M., G. Munoz, S. Vo Duy, D. T. Do, M. Vaudreuil, K. Goeury, F. Guillemette,

M. Amyot, and S. Sauvé. 2019. Widespread occurrence and spatial distribution of

glyphosate, atrazine, and neonicotinoids pesticides in the St. Lawrence and tributary rivers.

Environmental Pollution 250:29–39.

Naef-Daenzer, B., L. F. Keller, B. N. Daenzer, and L. F. Keller. 1999. The foraging performance

of great and blue tits (Parus major and P. caeruleus) in relation to caterpillar development,

and its consequences for nestling growth and fledging weight. Journal of Animal Ecology

68:708–718.

Nebel, S., A. Mills, J. D. McCracken, and P. D. Taylor. 2010. Declines of Aerial Insectivores in

North America Follow a Geographic Gradient. Avian Conservation and Ecology

5:10.5751/ACE-00391–050201.

Nocera, J. J., J. M. Blais, D. V. Beresford, L. K. Finity, C. Grooms, L. E. Kimpe, K. Kyser, N.

Michelutti, M. W. Reudink, and J. P. Smol. 2012. Historical pesticide applications

coincided with an altered diet of aerially foraging insectivorous chimney swifts.

Proceedings of the Royal Society B: Biological Sciences 279:3114–3120.

Olsson, O., and A. Bolin. 2014. A model for habitat selection and species distribution derived

from central place foraging theory. Oecologia 175:537–548.

Olsson, O., J. S. Brown, and K. L. Helf. 2008. A guide to central place effects in foraging.

Theoretical Population Biology 74:22–33.

Orians, G. H., and E. Pearson. 1979. On the theory of central place foraging. Analysis of

Ecological Systems:154–1177.

Pebesma, E. J. 2018. Simple Features for R: Standardized Support for Spatial Vector Data. The

R Journal 10:439.

Pigeon, G., R. Baeta, M. Bélisle, D. Garant, and F. Pelletier. 2013. Effects of agricultural
intensification and temperature on immune response to phytohemagglutinin in Tree
Swallows (Tachycineta bicolor). Canadian Journal of Zoology 91:56–63.

Pisa, L. W., V. Amaral-Rogers, L. P. Belzunces, J. M. Bonmatin, C. A. Downs, D. Goulson, D.
P. Kreutzweiser, C. Krupke, M. Liess, M. McField, C. A. Morrissey, D. A. Noome, J.
Settele, N. Simon-Delso, J. D. Stark, J. P. Van der Sluijs, H. Van Dyck, and M. Wiemers.
2015. Effects of neonicotinoids and fipronil on non-target invertebrates. Environmental
Science and Pollution Research 22:68–102.

Poisson, M., D. R. Garrett, A. Sigouin, M. Bélisle, D. Garant, L. Haroune, J. Bellenger, and F.
Pelletier. 2021. Assessing pesticides exposure effects on the reproductive performance of a
declining aerial insectivore. Ecological Applications 0:1–13.

Porlier, M., M. Bélisle, and D. Garant. 2009. Non-random distribution of individual genetic
diversity along an environmental gradient. Philosophical Transactions of the Royal Society
B: Biological Sciences 364:1543–1554.

Poulin, B., G. Lefebvre, and L. Paz. 2010. Red flag for green spray: adverse trophic effects of Bti
on breeding birds. Journal of Applied Ecology 47:884–889.

QGIS. 2020. QGIS Geographic Information System. QGIS Association. http://www.qgis.org.

R Core Team. 2019. R: A language and environment for statistical computing.

Rahmstorf, S., and D. Coumou. 2011. Increase of extreme events in a warming world.
Proceedings of the National Academy of Sciences 108:17905–17909.

Rappole, J. H., and A. R. Tipton. 1991. New harness design for attachment of radio transmitters

to small passerines. Journal of field ornithology 62:335–337.

Raven, P. H., and D. L. Wagner. 2021. Agricultural intensification and climate change are

rapidly decreasing insect biodiversity. Proceedings of the National Academy of Sciences

118:10.1073/pnas.2002548117.

Rendell, W. B., and R. J. Robertson. 1990. Influence of Forest Edge on Nest-Site Selection by

Tree Swallows. Wilson Journal of Ornithology 102:634–644.

Reznick, D., L. Nunney, and A. Tessier. 2000. Big houses, big cars, superfleas and the costs of

reproduction. Trends in Ecology & Evolution 15:421–425.

Rioux Paquette, S., D. Garant, F. Pelletier, and M. Bélisle. 2013. Seasonal patterns in Tree

Swallow prey (Diptera) abundance are affected by agricultural intensification. Ecological

Applications 23:122–133.

Rose, A. P. 2009. Temporal and Individual Variation in Offspring Provisioning by Tree

Swallows: A New Method of Automated Nest Attendance Monitoring. PLoS ONE

4:10.1371/journal.pone.0004111.

Rosenberg, K. V., A. M. Dokter, P. J. Blancher, J. R. Sauer, A. C. Smith, P. A. Smith, J. C.

Stanton, A. Panjabi, L. Helft, M. Parr, and P. P. Marra. 2019. Decline of the North

American avifauna. Science 366:120–124.

Rubolini, D., A. Massi, and F. Spina. 2002. Replacement of body feathers is associated with low

pre-migratory energy stores in a long-distance migratory bird, the barn swallow (Hirundo

rustica). Journal of Zoology 258:S0952836902001590.

Ruiz, J., and G. Domon. 2009. Analysis of landscape pattern change trajectories within areas of

intensive agricultural use: case study in a watershed of southern Québec, Canada.

Landscape Ecology 24:419–432.

Sæther, B.-E., and O. Bakke. 2000. Avian Life History Variation and Contribution of

Demographic Traits to the Population Growth Rate. Ecology 81:642–653.

Schifferli, L., M. U. Grüebler, H. A. J. Meijer, G. H. Visser, and B. Naef-Daenzer. 2014. Barn

Swallow Hirundo rustica parents work harder when foraging conditions are good. Ibis

156:777–787.

Schoener, T. W. 1971. Theory of Feeding Strategies. Annual Review of Ecology and

Systematics 2:369–404.

Shipley, J. R., C. W. Twining, C. C. Taff, M. N. Vitousek, A. Flack, and D. W. Winkler. 2020.

Birds advancing lay dates with warming springs face greater risk of chick mortality.

Proceedings of the National Academy of Sciences 117:25590–25594.

Smits, J. E. G., G. R. Bortolotti, M. Sebastian, and J. J. H. Ciborowski. 2005. Spatial, temporal,

and dietary determinants of organic contaminants in nestling tree swallows in point Pelee

national park, Ontario, Canada. Environmental Toxicology and Chemistry 24:3159–3165.

Spiller, K. J., and R. Dettmers. 2019. Evidence for multiple drivers of aerial insectivore declines

in North America. The Condor 121:1–13.

Staggenborg, J., H. M. Schaefer, C. Stange, B. Naef-Daenzer, and M. U. Grüebler. 2017. Time

and travelling costs during chick-rearing in relation to habitat quality in Little Owls Athene

noctua. Ibis 159:519–531.

Stanton, R., R. G. Clark, and C. A. Morrissey. 2017. Intensive agriculture and insect prey

availability influence oxidative status and return rates of an aerial insectivore. Ecosphere

8:10.1002/ecs2.1746.

Stanton, R., C. A. Morrissey, and R. G. Clark. 2016. Tree Swallow (Tachycineta bicolor)

foraging responses to agricultural land use and abundance of insect prey. Canadian Journal

of Zoology 94:637–642.

Stanton, R., C. A. Morrissey, and R. G. Clark. 2018. Analysis of trends and agricultural drivers

of farmland bird declines in North America: A review. Agriculture, Ecosystems &

Environment 254:244–254.

Stearns, S. C. 1992. The evolution of life histories. Oxford University Press.

Stephens, D. W., J. S. Brown, and R. C. Ydenberg. 2007. Foraging: behavior and ecology.

University of Chicago Press.

Tapper, S., J. J. Nocera, and G. Burness. 2020. Experimental evidence that hyperthermia limits

offspring provisioning in a temperate-breeding bird. Royal Society Open Science

7:10.1098/rsos.201589.

Taylor, L. R. 1963. Analysis of the Effect of Temperature on Insects in Flight. The Journal of

Animal Ecology 32:99–117.

Templ, M., K. Hron, and P. Filzmoser. 2011. robCompositions: An R-package for Robust

Statistical Analysis of Compositional Data. Pages 341–355 Compositional Data Analysis.

John Wiley & Sons, Ltd.

Tews, J., D. G. Bert, and P. Mineau. 2013. Estimated Mortality of Selected Migratory Bird

Species from Mowing and Other Mechanical Operations in Canadian Agriculture. Avian

Conservation and Ecology 8:10.5751/ACE-00559–080208.

Tremblay, I., D. W. Thomas, J. Blondel, P. Perret, and M. M. Lambrechts. 2004. The effect of

habitat quality on foraging patterns, provisioning rate and nestling growth in Corsican Blue

Tits Parus caeruleus. Ibis 147:17–24.

Tscharntke, T., A. M. Klein, A. Kruess, I. Steffan☐Dewenter, and C. Thies. 2005. Landscape

perspectives on agricultural intensification and biodiversity - ecosystem service

management. Ecology Letters 8:857–874.

Twining, C. W., J. R. Shipley, and D. W. Winkler. 2018. Aquatic insects rich in omega-3 fatty

acids drive breeding success in a widespread bird. Ecology Letters 21:1812–1820.

Wagner, D. L., E. M. Grames, M. L. Forister, M. R. Berenbaum, and D. Stopak. 2021. Insect

decline in the Anthropocene: Death by a thousand cuts. Proceedings of the National

Academy of Sciences 118:10.1073/pnas.2023989118.

Wiersma, P. 2005. Metabolic adjustments to increasing foraging costs of starlings in a closed

economy. Journal of Experimental Biology 208:4099–4108.

Winkler, D. W., K. K. Hallinger, D. R. Ardia, R. J. Robertson, B. J. M. Stutchbury, and R. R.

Cohen. 2020. Tree Swallow (Tachycineta bicolor). Page *in* A. F. Poole, editor. Birds of the

World. Cornell Lab of Ornithology.

Wood, S. N. 2017. Generalized additive models: an introduction with R. CRC press.

Wuebbles, D. J., K. Kunkel, M. Wehner, and Z. Zobel. 2014. Severe Weather in United States

Under a Changing Climate. Eos, Transactions American Geophysical Union 95:149–150.

897 Yap, K. N., O. R. Kim, K. C. Harris, and T. D. Williams. 2017. Physiological effects of

898   increased foraging effort in a small passerine. The Journal of Experimental Biology

899   220:4282–4291.

900 Ydenberg, R. C. 2007. Provisioning. Pages 273–303 *in* D. W. Stephens, J. S. Brown, and R. C.

901   Ydenberg, editors. Foraging: behavior and ecology. University of Chicago Press.

902 Zuur, A. F., E. N. Ieno, N. Walker, A. A. Saveliev, and G. M. Smith. 2009. Mixed effects models

903   and extensions in ecology with R. Page Smart Society. Springer New York.

Tables

Table 1: Mean ± SD of covariates used to model response variables. Covariates for the hourly number of boluses and total bolus biomass delivered to nestlings were representative of the hourly food provisioning sessions conducted between 2011 and 2016. Covariates of foraging distances were calculated for the hourly interval's representative of the hourly GPS locations. Dashes imply the covariate was not used in the modeling of the response variable. Covariate groups identify which covariates were present within models found in Table 1.

Covariate	Acronym	Unit	Covariate group	Biomass	Number	Foraging distance
Starting time	ST	Time	Base	262.31 ± 150.16	251.57 ± 153.55	-
Maximum temperature	TMAX	°C	Base/Food	26.20 ± 4.08	25.54 ± 3.94	20.81 ± 5.11
Brood demand proxy	DM	Brood size * Brood age	Base/Demand	40.50 ± 13.89	39.14 ± 12.8	28.9 ± 10.59
Local surface water (1 km)	LSW	%	Base/ LSW	0.80 ± 1.43	0.87 ± 1.58	1.52 ± 1.73
Regional surface water (10 km)	RSW	%	Base	1.58 ± 1.46	1.56 ± 1.42	2.67 ± 1.93
Prey availability (Diptera biomass)	PA	g	Food	0.02 ± 0.01	0.02 ± 0.01	-
Site score Comp.1	Comp.1	-	Land	0.34 ± 1.73	0.42 ± 1.72	0.09 ± 2.42
Site score Comp.2	Comp.2	-	Land	-0.04 ± 0.88	-0.01 ± 0.86	-0.25 ± 1.24
Ordinal day	OD	days	Base	-	-	170.64 ± 7.06

Table 2: Standardized coefficient estimates and their 95% confidence intervals included in the top model of each response variable. See Table 1 for definitions of covariates and Appendix S2: Table S3 for outcome of model selection. Coefficients in bold indicate confidence intervals not overlapping zero. Numbers next to acronym refer to the order of the polynomial term.

Covariate	Biomass	Delivery	Travel	Distance
Comp.1	**0.04** **(0.00, 0.08)**	0.02 (-0.04, 0.07)	**-0.11** **(-0.16, -0.05)**	-0.08 (-0.16, 0.01)
Comp.1:Comp.2	**0.08** **(0.03, 0.13)**	**0.09** **(0.02, 0.16)**	**-0.17** **(-0.25, -0.09)**	**-0.15** **(-0.25, -0.05)**
Comp.2	**-0.11** **(-0.20, -0.02)**	**-0.15** **(-0.27, -0.02)**	0.05 (-0.06, 0.16)	0.03 (-0.15, 0.20)
Female Age (SY)	0.09 (-0.09, 0.26)	0.13 (-0.13, 0.40)	-	-
DM1	**5.09** **(2.62, 7.55)**	**9.78** **(7.09, 12.47)**	**3.12** **(0.41, 5.82)**	3.21 (-0.92, 7.33)
DM2	**-6.09** **(-8.58, -3.60)**	**-3.34** **(-6.11, -0.57)**	**-3.02** **(-5.81, -0.23)**	**-4.76** **(-8.81, -0.71)**
DM3	**3.72** **(1.15, 6.29)**	1.07 (-1.64, 3.78)	-1.41 (-1.45, 4.26)	1.45 (-3.02, 5.91)
DM1:Comp.1	0.47 (-0.90, 1.85)	-	-	-
DM2:Comp.1	**1.65** **(0.17, 3.13)**	-	-	-
DM3:Comp.1	-0.65 (-2.29, 0.98)	-	-	-
OD1	-	-	**10.80** **(7.32, 14.28)**	**12.50** **(7.03, 17.97)**
OD2	-	-	**-5.62** **(-8.57, -2.67)**	**-6.28** **(-10.90, -1.66)**
ST1	**2.93** **(0.53, 5.33)**	2.53 (-0.19, 5.26)	-	-
ST2	**-2.49** **(-4.90, -0.09)**	**-3.02** **(-5.61, -0.43)**	-	-
TMAX1	0.87 (-1.62, 3.36)	0.13 (-2.58, 2.83)	**8.79** **(6.34, 11.24)**	**8.24** **(5.53, 10.96)**
TMAX2	**-5.77** **(-8.21, -3.32)**	**-3.99** **(-6.75, -1.23)**	**-5.35** **(-7.73, -2.96)**	**-4.61** **(-7.23, -1.99)**
Rain (Yes)	-0.17 (-0.44, 0.10)	-0.18 (-0.52, 0.16)	-	-
LSW	**-0.10** **(-0.17, -0.03)**	**-0.12** **(-0.22, -0.02)**	0.17 (-0.02, 0.37)	0.06 (-0.22, 0.35)
LSW:Comp.1	-	-	**-0.09** **(-0.17, -0.02)**	-
LSW:RSW	0.03 (-0.07, 0.14)	0.12 (-0.03, 0.28)	**-0.29** **(-0.52, -0.07)**	-0.29 (-0.62, 0.05)
RSW	0.06 (-0.02, 0.14)	0.05 (-0.06, 0.16)	**-0.26** **(-0.44, -0.09)**	**-0.31** **(-0.58, -0.04)**

Figures

Figure 1: Map of study system. Each point represents the location of a farm, filled white

points are farms where GPS were applied to 43 females. Underlying image represents the

agricultural gradient derived from the Annual crop inventory (AAFC 2013). Light yellow

represents the more agro-intensive areas while light green represents less agro-intensive areas.

Figure 2: Results of the robust PCA for compositional data including the site scores of the

five landscape context scenarios for the food provisioning sessions (A) and foraging distance

studies (B). Points in the background represent the site scores assigned to each food provisioning

session. The arrows lengths are eigenvectors representing the direction and magnitude of

correlation between habitats. The five landscape scenarios are: ones dominated by corn or

soybean with 8.8% forest cover (Corn/soybean with forest, 1), ones denuded of forest cover and

dominated by only corn or soybean (Corn/Soybean,2), ones denuded of forest cover and

dominated by a mixture of corn, soybean and cereals (mixed row crop, 3), ones minimizing

forest cover (9.0%) and dominated by forage crops (Forage, 4), and agricultural landscapes with

69.0% forest cover (Agro-forested, 5)

Figure 3:   Response surfaces of the principal response variables against the first two components deriving the landscape context. Response surface are the two-dimensional unconditional predictions of each response variable over the observed range of values along the first and second component values of the PCA defining landscape context found in Fig 3. Deeper blue represents lower predicted values while yellow represents greater values. Predicted hourly bolus delivery rate (A), hourly bolus biomass (B) delivered to nestlings. Predicted hourly distance traveled (C) and hourly distance from nest box (D). Numbers in each panel are landscape context scenarios and represent the predominant types of landscape contexts experienced across the agricultural gradient and correspond to the points in numbers in Fig. 2.

Figure 4: Reasoning underlying our hypothesis that food patches within forage landscapes allow for a more rapid accumulation of food resources than those found in more agro-intensive landscapes. Assuming that food provisioning Tree Swallow females foraged according to the marginal value theorem, and given we found (1) that the individual bolus biomass was similar across the agricultural intensification gradient, (2) that travel distances were greater in forage than in agro-intense landscapes, then only a lower gain curve in agro-intense landscapes is compatible with our findings that hourly bolus biomass and hourly number of boluses delivered were higher in forage than in agro-intense landscapes. Gain rate is graphically represented by the slope of the dashed line connecting mean travel time to the tangent of the gain curve.