1 **Title:**   **Mugs   and   Plants: Object   Semantic   Knowledge   Alters**
2   **Perceptual Processing with Behavioral Ramifications**

5 **Authors:**   Dick Dubbelde and Sarah Shomstein

7 **Author affiliation:**   Department of Psychological and Brain Sciences

8   The George Washington University

9
10
11
Corresponding author:

Dick Dubbelde
Department of Psychology
The George Washington University
2125 G St. NW
Washington, DC 20052
E-mail: dubbelde@gwu.edu
19
20

Abstract

Neural processing of objects with action associations is thought to recruit dorsal visual regions

more than objects without such associations. We hypothesized that because the dorsal and

ventral visual pathways have differing proportions of magno- and parvo-cellular input, there

should be behavioral differences in perceptual tasks between manipulable and non-manipulable

objects. This hypothesis was tested using gap detection task, suited to the spatial resolution of the

ventral parvocellular processing, and object flicker discrimination task, suited to the temporal

resolution of the dorsal magnocellular processing. Directly predicted from the cellular

composition of each pathway, a non-manipulable object advantage was observed in tasks relying

on spatial resolution, and a manipulable object advantage in temporal discrimination. We also

show that these relative advantages are modulated by either reducing object recognition through

inversion or by suppressing magnocellular processing using red light. These results establish

perceptual differences between objects dependent on prior knowledge and experience.

34
35

Object perception and recognition underlies effective engagement with our complex

environment. When looking at a potted plant, for example, precise access to long term semantic

memory may be needed to facilitate a more detailed identification of the plant to know how often

it needs to be watered. However, when reaching for a mug to have your morning coffee, detailed

identification may be less important (e.g., which one of your 10 favorite mugs it is), but what is

more pertinent is computing a motor plan to bring the mug to your mouth without spilling. The

visual object processing necessary to identify plants and to drink from mugs is understood to

progress along two distinct but interacting neural processing pathways, each of which is

understood to subserve different end goals (Mishkin & Ungerleider, 1982; Kravitz et al., 2011).

While the hypothesis of biased processing across the two pathways is widely accepted, it is

unclear whether there are behavioral ramifications that are directly related to object processing

being biased to different pathways. Here, we provide evidence that object semantic knowledge

evokes differences in object perception based on the recruited regions across the dorsal or ventral

streams. Importantly, we gain insight into neural processing by using purely behavioral methods

that are custom tailored to the response properties of the neurons in both the dorsal (higher

temporal resolution) and ventral pathways (higher spatial resolution).

The ventral pathway, commonly referred to as the 'what' pathway, is characterized by object

feature selectivity and projects anteriorly from the occipital visual areas toward the hippocampus

and the inferior temporal cortex (Kravitz et al., 2013). Damage to cortical regions along the

ventral pathway results in visual agnosia, evidenced by difficulties in object recognition but

preserved motor interactions with objects (Goodale & Milner, 1992; 1994; Karnath et al., 2009).

Patients with ventral pathway damage but an intact dorsal pathway continue to manipulate,

move, and reach for objects without much difficulty, while their ability to identify objects is

greatly impaired (James et al., 2003). The dorsal pathway, in turn, colloquially known as the

'where' or 'how' pathway, courses from the occipital visual areas into the parietal cortex. The

dorsal pathway is commonly understood to subserve object processing for the purposes of

reaching, grasping, and manipulation. While both pathways contribute to visual perception, the

demands of perceiving a specific object can differentially engage the two pathways such that the

dorsal pathway provides greater contribution for visual processing of objects that are directly

relevant for object manipulation (e.g., a hammer, a plunger, or a mug). Evidence for dynamic

recruitment of processing from one pathway more than the other has been garnered from a wide

range of techniques and paradigms. For example, while presenting participants with objects of

various categories, including tools, places, animals, and faces, an increase in dorsal pathway

processing, specifically in the left ventral premotor and left posterior parietal cortices, has been

observed exclusively in response to tool presentation (Chao & Martin, 2000). Similar results

which evidence a dorsal bias for manipulable objects have been found in functional magnetic

resonance imaging studies (Noppeney et al., 2006a; Mahon et al., 2007, Almeida et al., 2013,

Chen et al., 2018) and with various other paradigms including interocular suppression (Fang and

He, 2005) and continuous flash suppression (Almeida et al., 2008; Almeida et al., 2010).

While neuroimaging and neurophysiological evidence has been convincing in showing

selectivity for objects in various areas across the two streams, the impact that differential

processing across the two pathways has on object perception and subsequent behavior has not

been characterized. Hypotheses regarding how dynamic recruitment of the dorsal and ventral

pathways differentially influences perception and behavior are based on the anatomy and

physiology of the two visual pathways. Considering cellular physiology, there is an asymmetry

in innervation of the two pathways that overlaps with the separate magnocellular and

parvocellular processing pathways identified in anatomical architecture of visual processing

(Maunsell et al., 1990; Baizer et al., 1991; Ferrera et al., 1992). The asymmetry in cellular

composition across the two pathways endows each pathway with different response properties in

accordance with cell stimulus preferences within the magno- and parvo-cellular pathways. The

magno- and parvo-cellular pathways originate from different types of ganglion cells within the

retina and course separately through different layers of the lateral geniculate nucleus (Leventhal

et al., 1981; Perry et al., 1984), innervating separate layers of V1 (Blasdel & Lund, 1983). From

V1, the parvocellular pathway can be followed into area V2, through area V4, and then into the

inferior parietal cortex while the magnocellular pathway can be traced into different regions of

area V2, through areas V3d and MT, and then into the posterior parietal cortices (DeYoe & van

Essen, 1988; Livingstone & Hubel, 1988).

Differentiation in cellular innervation of the two processing pathways leads to different

response properties that convey information of different spatial and temporal resolutions. The

magnocellular channel is derived from the parasol ganglion cells with relatively large receptive

fields, spanning large regions of the retina (Nassi & Callaway, 2009). The cells of the

magnocellular channel are more myelinated, and have less neuronal convergence than the

parvocellular channel (Maunsell et al., 1999a). The features of the parasol retinal ganglion cells

enable the dorsally biased magnocellular pathway to encode information with a higher temporal

resolution than that of the parvocellular channel (Pokorny and Smith, 1997; but see Maunsell et

al., 1999b). Conversely, the parvocellular channel is derived from the midget retinal ganglion

cells, which receive input primarily from cone receptors and have smaller receptive fields (Nassi

& Callaway, 2009). The features of the midget retinal ganglion cells enable the ventrally biased

parvocellular channel to encode information with a higher spatial resolution than that of the

magnocellular channel (Derrington & Lennie, 1984; Leonova et al., 2003; McAnany &

Alexander, 2008).

Due to the asymmetric contributions of the magno- and parvo-cellular pathways to the

dorsal and ventral pathway processing pathways, we hypothesize that semantic knowledge of

object utility (manipulable or non-manipulable) determines processing bias to either dorsal or

ventral pathway processing and thus has consequences for perceptual processing. Thus, if

semantic knowledge of an object's utility determines to which of the two pathways the bias will

accrue, then: (1) manipulable objects, that elicit a higher degree of magnocellularly biased dorsal

processing, should be processed with higher temporal resolution; and (2) non-manipulable

objects, that rely more on parvocellularly biased ventral processing, should be processed with

higher spatial resolution. Across five experiments, this hypothesis is tested by comparing

processing of temporal and spatial resolution in two object groups: manipulable and non-

manipulable objects. Each group consisted of ten line-drawings of objects which were carefully

controlled for low-level perceptual differences (width of the gap line, luminance, size, and aspect

ratio which a measure of elongation; see Methods for details). Due to the fact that our hypotheses

are predicated on the idea that action associations bias processing across the two streams, the

objects were chosen based on how often they are manipulated with the hands when they are

encountered rather than by their association to any particular semantic grouping. The

manipulable object set included objects that are typically manipulated by the hands when they

are encountered, such as a hammer, a plunger, and a mug (Fig. 1A). The non-manipulable object

set included objects that are rarely manipulated with the hands when they are encountered, such

as a fire hydrant, a picture frame, and a potted plant (Fig. 1A).

A	Fire Hydrant	Picture Frame	Window	Toilet	Candle	Garbage Can	Water Fountain	Potted Plant	Fan	Lamp
	0.010°	0.006°	0.008°	0.006°	0.010°	0.005°	0.018°	0.013°	0.006°	0.007°
	Snow Shovel	Handsaw	Plunger	Screw-driver	Hammer	Wrench	Knife	Bottle Opener	Spatula	Mug
	0.007°	0.011°	0.007°	0.012°	0.010°	0.007°	0.010°	0.008°	0.015°	0.010°

B	Fire Hydrant	Picture Frame	Window	Toilet	Candle	Garbage Can	Water Fountain	Potted Plant	Fan	Lamp
	0.010°	0.013°	0.012°	0.009°	0.009°	0.007°	0.011°	0.012°	0.007°	0.008°
	Snow Shovel	Handsaw	Plunger	Screw-driver	Hammer	Wrench	Knife	Bottle Opener	Spatula	Mug
	0.016°	0.018°	0.007°	0.014°	0.008°	0.009°	0.009°	0.009°	0.009°	0.013°

Figure 1 *Stimulus set used in the experimental paradigms. (A) Line drawings of manipulable and non-manipulable objects used in Exp. 1. (B) Pictures of real-world manipulable and non-manipulable objects used in the second control manipulation for Exp. 1. The numbers beneath each set of objects correspond to the size of the spatial gap titrated to each object.*

Experiment 1 employed two paradigms, each designed to take advantage of the unique stimulus response properties of the two visual processing pathways and their corresponding cellular composition. To measure ventral pathway engagement a gap detection task was employed. Each object appeared to the left or the right of fixation, and would either have an uninterrupted bottom contour or have a small spatial gap in the center of the bottom line of the object (Fig. 2A). Object group was counterbalanced such that manipulable objects were seen as commonly as non-manipulable objects, and objects within each group were counterbalanced such that all objects were seen an equal number of times. The size of the gap was titrated, in a separate procedure for each object individually in an independent group of participants, to a specific threshold making the task difficult but not impossible (See Methods for details). Participants' task was to report the presence or absence of the gap. If object semantic knowledge

determines the route of processing, then the presence of a spatial gap on non-manipulable objects

will be detected with higher sensitivity as compared to manipulable objects because processing

will be biased to ventral pathway's parvocellular processing.

To measure dorsal pathway engagement, a blink detection task was used. Similar to the

spatial gap detection task, participants detected the presence of a short blink. During a blink trial,

the object was presented for 80 ms, then removed for the short duration of 16 ms, then

redisplayed for additional 48 ms. The duration of the blink was titrated with a staircase

procedure, with the shortest titrations limited by the lower bound of the monitor's refresh rate.

Thus, shortest blink duration was set to 16 ms. During a non-blink trial, the objects remained on

the screen for the total duration of 144 ms. Similar to the spatial gap paradigm, participants

responded with a button press to indicate whether a blink was detected. If object semantic

knowledge determines the route of processing, then the presence of a temporal gap during the

presentation of manipulable objects will be detected with higher sensitivity as compared to non-

manipulable objects because processing will be biased to magnocellularly-dominated dorsal

pathway. For both tasks, participants were given feedback on incorrect trials. Each participant

performed both tasks (3 blocks each), in a counterbalanced order, across six blocks.

A

B

C

Object Group x Task Type
p = 0.024

Object Group x Task Type
p = 0.015

Gap   Blink   Gap   Blink

Upside-up   Upside-down

Figure 2. The paradigm and results for Experiments 1 and 2. Participants maintained fixation on the center cross. An object appeared to the left or right of fixation. A) For the spatial-gap paradigm, participants indicated by keypress if the bottom line of the presented object contained a gap. B) for the temporal paradigm, participants were to indicate whether the object flickered. C) Results for Experiment 1 (n=26) and 2 (n=30). For Experiment 1 (C, left), there was a significant main effect of task type (F(1, 25) = 50.953, p < 0.001, $\eta_p^2 = 0.671$) such that d' sensitivity was higher for blinks than for gaps ($M_{gaps} = 1.93$; $M_{blinks} = 2.97$); and a significant two-way interaction between object group and task type (F(1, 25) = 6.772, p = 0.015, $\eta_p^2 = 0.213$), driven by the difference between object groups in the gap condition (F(1, 25) = 9.888, p = 0.004, $\eta_p^2 = 0.283$, with non-manipulable objects having a higher d' for gaps than manipulable objects ($M_{non-manipulable} = 2.021$; $M_{manipulable} = 1.831$). For Experiment 2 (C, right), there was a significant main effect of object group (F(1, 25) = 5.860, p = 0.023, $\eta_p^2 = 0.190$) such that d' was higher for non-manipulable objects than for manipulable objects ($M_{non-manipulable} = 2.466$; $M_{manipulable} = 2.340$) (Fig. 2C, right). There was also a significant main effect of task type (F(1, 25) = 82.298, p < 0.001, $\eta_p^2 = 0.767$) such that d' was higher for blinks than for gaps ($M_{gaps} = 1.782$; $M_{blinks} = 3.024$). There was no significant interaction between object group and task type (F(1, 25) = 0.508, p = 0.483, $\eta_p^2 = 0.020$). Additionally, across the two experiments, with orientation as a between subjects factor, there was a significant three-way interaction between object group, task type, and orientation (F(1, 50) = 5.381, p = 0.024, $\eta_p^2 = 0.097$) such that orientation significantly reduced the effect for the inverted objects.

The spatial and temporal paradigms used d' as a measure of perceptual sensitivity (Fig.

2). A two-way repeated-measures analysis of variance (ANOVA) was conducted on d' with

object group (manipulable, non-manipulable) and stimulus type (gap, blink) as within-subject

variables (Fig. 2C, left). The ANOVA revealed no significant main effect of object group (*F*(1,

25) = 2.342, *p* = 0.138, $\eta_p^2 = 0.086$), and a significant main effect of task type (*F*(1, 25) =

50.953, *p* < 0.001, $\eta_p^2 = 0.671$) such that d' sensitivity was higher for blinks than for gaps (M_{gaps}

= 1.93; M_{blinks} = 2.97). Importantly, and consistent with the hypothesis of differential

engagement of two pathways depending on object utility, a significant two-way interaction

between object group and task type ($F(1, 25) = 6.772$, $p = 0.015$, $\eta_p^2 = 0.213$) was observed,

driven by the difference between object groups in the gap condition ($F(1, 25) = 9.888$, $p = 0.004$,

$\eta_p^2 = 0.283$), with non-manipulable objects having a higher d' for gaps than manipulable objects

($M_{non-manipulable} = 2.021$; $M_{manipulable} = 1.831$). The interaction effect and the driving main effect are

consistent with the prediction that non-manipulable objects, given reliance on ventral pathway

processing, should yield higher sensitivity in the detection of spatial gaps than manipulable

objects. Notably, the expected higher sensitivity to temporal gaps in the manipulable object set

was not supported. Given the high sensitivity for the temporal task type (d' $= 2.941$ for non-

manipulable objects and d' $= 2.993$ for manipulable objects), the lack of difference between the

object groups in the blink condition was likely due to the temporal task being overly easy (i.e.,

ceiling effect; see Exp. 4 for a manipulation that avoids ceiling effects).

While Experiment 1 provides evidence for biased engagement of the ventral pathway for

non-manipulable object perception during a temporal task, it could be argued that despite careful

low-level feature controls (e.g., luminance, size, elongation) an uncontrolled low-level difference

between manipulable and non-manipulable objects is responsible for driving the manipulable vs.

non-manipulable advantage in the spatial gap task. In order to rule out this possibility and,

simultaneously, to further probe the hypothesis that semantic knowledge of object utility

(manipulable or non-manipulable) biases the pathway that will ultimately process the object, we

conducted two control experiments. In addition to serving as a low-level control and a further

test of the hypothesis, these experiments provide internal replications of the reported finding (see

Open Science Collaboration, 2015).

In the first control manipulation, the same paradigm and objects from Experiment 1 were

used with the manipulation of inverting objects (upside-down). The inversion preserved the low-

level features of each object while impairing participants' ability to rapidly recognize and access

their semantic knowledge of the objects' function. Inversion has been found to interfere with

recognition of faces and objects (Diamond and Carey, 1986). Following from the hypothesis that

semantic knowledge of object's utility drives the perceptual difference between manipulable and

non-manipulable objects, we predict that object inversion should reduce the difference between

manipulable and non-manipulable objects. A two-way repeated-measures ANOVA was

conducted on d' with object group (manipulable, non-manipulable) and task type (gap, blink) as

within-subject variables. The ANOVA revealed a significant main effect of object group ($F(1,$

$25) = 5.860$, $p = 0.023$, $\eta_p^2 = 0.190$) such that d' was higher for non-manipulable objects than for

manipulable objects ($M_{non-manipulable} = 2.466$; $M_{manipulable} = 2.340$) (Fig. 2C, right). There was also

a significant main effect of task type ($F(1, 25) = 82.298$, $p < 0.001$, $\eta_p^2 = 0.767$) such that d' was

higher for blinks than for gaps ($M_{gaps} = 1.782$; $M_{blinks} = 3.024$). As predicted, there was no

significant interaction between object group and task type ($F(1, 25) = 0.508$, $p = 0.483$, $\eta_p^2 =$

0.020). In order to statistically demonstrate that results of the inversion experiment are indeed

different from those observed in Experiment 1, data were subjected to a between-subject

ANOVA with object orientation (upright, inverted) as a between subjects variable and object

group and task type as within subject factors. The ANOVA revealed a significant three-way

interaction between object group, task type, and orientation ($F(1, 50) = 5.381$, $p = 0.024$, $\eta_p^2 =$

0.097) such that, as predicted, orientation significantly reduced the effect for the inverted objects.

In the second control manipulation, line drawings were replaced by real-world images of

corresponding objects, such that a line drawing of a candle was replaced with a picture of a

candle, etc. (Fig. 1B). The prediction remained the same as in the original experiment, if object

semantic knowledge determines which visual processing pathway object processing is biased to

then non-manipulable objects will be biased to the ventral pathway leading to higher sensitivity

(as measured by d') in the spatial gap task, replicating the pattern of performance observed in

Experiment 1. A two-way repeated-measures ANOVA was conducted on d' with object group

(manipulable, non-manipulable) and task type (gap, blink) as within-subject variables. The

ANOVA revealed a significant main effect of object group ($F(1, 29) = 28.211$, $p < 0.001$, $\eta_p^2 =$

0.493) with non-manipulable objects having a higher sensitivity than manipulable objects (M_{non-}

$_{manipulable} = 2.361$, $M_{manipulable} = 2.046$) (Fig. 3). There was also a significant main effect of task

type ($F(1, 29) = 8.413$, $p = 0.007$, $\eta_p^2 = 0.225$) with blinks having a higher average sensitivity

than gaps ($M_{blinks} = 2.409$, $M_{gaps} = 1.998$). Importantly, a two-way interaction between object

group and task type was also significant ($F(1, 29) = 13.061$, $p = 0.001$, $\eta_p^2 = 0.311$), driven by a

simple main effect in the gap condition ($F(1, 29) = 46.424$, $p < 0.001$, $\eta_p^2 = 0.616$) such that non-

manipulable objects had a higher d' for gaps than did the manipulable objects ($M_{non-manipulable} =$

2.238; $M_{manipulable} = 1.758$). In order to statistically demonstrate that results observed using real

objects are indistinguishable from the results observed using the line-drawings in Experiment 1,

the data were subjected to a between-subject ANOVA with stimulus type (line drawings, real

object images) as a between subjects variable and object group and task type as within subject

factors. The ANOVA revealed no significant three-way interaction between object group, task

type, and stimulus type ($F(1, 54) = 0.454$, $p = 0.503$, $\eta_p^2 = 0.000159$). These results replicate

Experiment 1, providing strong additional support for the hypothesis that the perceptual

differences between manipulable objects and non-manipulable objects is due to the semantic

content of the objects rather than low-level features.

Figure 3. *Results from the second control manipulation of experiment one in which real object images were used instead of line drawing images. There was a significant main effect of object group ($F(1, 29) = 28.211$, $p < 0.001$, $\eta_p^2 = 0.493$) with non-manipulable objects having a higher sensitivity than manipulable objects ($M_{non-manipulable} = 2.361$, $M_{manipulable} = 2.046$) (Fig. 3). There was also a significant main effect of task type ($F(1, 29) = 8.413$, $p = 0.007$, $\eta_p^2 = 0.225$) with blinks having a higher average than gaps ($M_{blinks} = 2.409$, $M_{gaps} = 1.998$). The two-way interaction between object group and task type was also significant ($F(1, 29) = 13.061$, $p = 0.001$, $\eta_p^2 = 0.311$), driven by a simple main effect in the gap condition ($F(1, 29) = 46.424$, $p < 0.001$, $\eta_p^2 = 0.616$) such that non-manipulable objects had a higher d' for gaps than did the manipulable objects ($M_{non-manipulable} = 2.238$; $M_{manipulable} = 1.758$).*

235  While the first three experiments yielded strong supporting evidence that gap detection is

236 better on non-manipulable objects than manipulable objects, perhaps resulting from bias toward

237 the ventral pathway, manipulable objects failed to elicit an advantage in temporal resolution,

238 failing to provide evidence of a dorsal pathway bias for manipulable objects. One explanation for

239 the null effect is that the temporal gap detection task was too easy (ceiling effect), evidenced by

240 high d' values in the temporal task. To increase task difficulty, and move away from the ceiling

241 performance, the blink paradigm was re-designed to a discrimination task[1]. Objects were

242 presented for 80 ms, removed from the screen for either a 16 ms or 48 ms blink, and redisplayed

243 for 48 ms (Fig. 4). Participants' task was to indicate whether the blink duration was short or long.

244 The prediction was that manipulable objects should have a performance benefit over non-

245 manipulable objects due to the increased magnocellular input, and therefore temporal resolution,

246 of the dorsal pathway. Due to the difficulty of the task (mean accuracy 77%), evidenced by low

247 d' values, analyses were conducted on RT. A two-way repeated-measures ANOVA was

---

[1] We thank Ed Awh, from the University of Chicago, for the helpful suggestion.

248   conducted on RT with object group (manipulable, non-manipulable) and blink duration (short,

249   long) as within-subject variables. Only the RTs for correct responses were analyzed.

A

B

Object Group x Task Type

$p = 0.006$

The ANOVA revealed no significant main effect of object group ($F(1, 24) = 4.080, p =$

$0.055, \eta_p^2 = 0.145$) (Fig. 4). There was a significant main effect of blink duration ($F(1, 24) =$

$4.850 \ p = 0.038 \ \eta_p^2 = 0.168$) such that the short duration elicited a higher RTs than did the long

duration ($M_{short} = 642.68$ ms; $M_{long} = 621.997$ ms). The two-way interaction between object

group and blink duration was significant ($F(1, 24) = 9.011, p = 0.006, \eta_p^2 = 0.273$) driven by a

simple main effect in the short blink condition ($F(1, 24) = 11.710, p = 0.002, \eta_p^2 = 0.328$) such

that manipulable objects had a lower RT for short blinks than did the non-manipulable objects

($M_{manipulable} = 633.150; M_{non-manipulable} = 652.210$). In accuracy, no effect was observed for object

group ($F(1, 24) = 3.666, p = 0.068, \eta_p^2 = 0.133$), blink duration ($F(1, 24) = 1.509, p = 0.231, \eta_p^2$

= 0.059), nor the interaction ($F(1, 24) = 2.000$, $p = 0.170$, $\eta_p^2 = 0.077$), suggesting that all

significant effects were absorbed by the response time measure and no speed-accuracy tradeoffs

were observed, indicating consistency with response time results. These results, specifically the

simple main effect in the short gap condition with manipulable objects having a lower RT than

non-manipulable objects, supports our hypothesis that manipulable object processing would

elicit higher temporal resolution than non-manipulable object processing. The lack of a

difference in the long duration condition is likely due to a ceiling effect, similar to the lack of

temporal effect in Experiment 1.

The last test of our hypotheses is derived from the neurophysiological differences

between the dorsal and ventral pathways. It was reasoned that if the differences in spatial gap

sensitivity for non-manipulable objects are mechanistically derived from the magnocellular and

parvocellular dichotomy of the two visual processing pathways, then the effect should be

modulated through manipulation of processing within the pathways. Ambient red light has been

shown to suppress activity in the magnocellular processing channel because of the large number

of visually responsive cells with red inhibitory surrounds in their receptive fields (Wiesel &

Hubel, 1966). Therefore, presenting objects with a red background should suppress the

contribution from the parietal magnocellular pathway, whereas a background of any other color

should have no impact on processing. To test this prediction, the spatial resolution paradigm

from Experiment 1 was used with the manipulation of background color varying between green

or red (Fig. 5A). Due to the suppression of the magnocellular channel by red light, it was

predicted that the color of the background should modulate the perceptual difference in spatial

gap detected observed in Experiment 1. Namely, suppressing the contribution of the dorsal

pathway should exacerbate the difference between manipulable and non-manipulable objects. A

two-way repeated-measures ANOVA was conducted on d' with object group (manipulable, non-

manipulable) and background color (green, red) as within-subject variables. The ANOVA

revealed a significant main effect of object group ($F(1, 25) = 17.171$, $p < 0.001$, $\eta_p^2 = 0.407$)

with non-manipulable objects having a higher average than manipulable objects ($M_{\text{non-manipulable}} =$

2.25, $M_{\text{manipulable}} = 2.02$) (Fig. 5B). There was no significant main effect of background color

($F(1, 25) = 2.138$, $p = 0.156$, $\eta_p^2 = 0.079$) but, as predicted, there was a significant two-way

interaction between object group and background color ($F(1, 25) = 4.444$, $p = 0.045$, $\eta_p^2 = 0.151$)

such that the effect was increased with the red background. It was hypothesized that if the

perceptual differences between manipulable and non-manipulable objects are mechanistically

derived from the magno- and parvo-cellular processing in the two visual pathways, then

suppression of the magnocellular processing with red light should modulate the effect. The

results of Experiment 5 support our hypothesis by demonstrating an increase of the perceptual

difference between manipulable and non-manipulable objects with red light.

Figure 5. The paradigm and results of experiment 5. (A) Participants maintained fixation on the center cross. An object then appeared to the left or right of fixation. As in Experiment 1, participants were asked to detect the presence of small gap in the bottom outline of the object. The background color (red or green) was manipulated. (B) There was a significant main effect of object group ($F(1, 25) = 17.171$, $p < 0.001$, $\eta_p^2 = 0.407$) with non-manipulable objects having a higher average than manipulable objects ($M_{\text{non-manipulable}} = 2.25$, $M_{\text{manipulable}} = 2.02$). There was no significant main effect of background color ($F(1, 25) = 2.138$, $p = 0.156$, $\eta_p^2 = 0.079$) but there was a significant two-way interaction between object group and background color ($F(1, 25) = 4.444$, $p = 0.045$, $\eta_p^2 = 0.151$).

We hypothesized that manipulable and non-manipulable objects, due to differential

recruitment of the two visual pathways, would elicit perceptual differences stemming from the

characteristic spatial and temporal resolutions associated with the magno- and parvo-cellular

inputs. In five experiments we provide strong evidence in support of behavioral consequences

driven by physiological and anatomical differences of the two visual pathways, and argue that

semantic knowledge of object utility guides processing along a particular pathway. If an object

has a strong action association, its processing will be largely determined by activity in the dorsal

pathway which is not found with objects that lack strong action association (Chao & Martin,

2000; Noppeney et al., 2006; Mahon et al., 2007). The increased level of activity in the parietal

regions endows the perception of action-associated objects with greater access to the information

encoded in the magnocellular channel which preferentially courses through the dorsal pathway.

Without the enhanced dorsal pathway activity evoked by action associations, object

representation is more dependent on the ventral pathway, which has a higher ratio of

parvocellular channel information than does the dorsal pathway (Maunsell et al., 1990; Baizer et

al., 1991; Ferrera et al., 1992). Based on differential magno- and parvo-cellular input to the two

pathways, we predicted that the perception of objects with strong action associations results in an

increased temporal resolution and that the perception of objects without such associations results

in an increased spatial resolution. In the experiments reported here, we offer evidence that

objects are perceived with different spatial and temporal resolutions depending on object

semantic knowledge (namely, whether the object is manipulable or non-manipulable). In two

follow-up control experiments, further evidence was provided in support of the hypothesis that

the difference between manipulable and non-manipulable objects in the spatial task was driven

by the semantic knowledge of the objects rather than possible low-level visual features. Namely,

it was observed that the effect is curtailed by impeding the access of object semantic knowledge

through inversion and it was observed that the perceptual differences between manipulable and

non-manipulable objects replicated with a separate set of more realistic object images (with a

different set of low-level object properties). Lastly, to test if the differing relative proportions of

magno- and parvo-cellular input are responsible for the perceptual differences that were observed

in our first four experiments, ambient red light was used to suppress activity of the magnocellular

channel. It was observed that this increased the perceptual difference in the spatial task between

manipulable and non-manipulable objects.

Results are strengthened by a set of various low-level controls imposed on the stimulus

set. Manipulable and non-manipulable object groups were controlled for size, hue, saturation,

visual clutter, and image value. Additionally, object groups were controlled for the degree of

elongation difference (i.e., aspect ratio) as this factor has been shown to drive some amount of

dorsal visual pathway selective for tools, which are a subset of manipulable object (Chen et al.,

2018).

Based on the evidence provided, we argue that semantic knowledge of the manipulability

of objects, as defined by strong associations with an action appropriate for the item, generates the

perceptual differences between manipulable and non-manipulable objects. Previous studies have

shown that manipulable objects evoke a larger degree of dorsal pathway processing than do non-

manipulable objects (Chao & Martin, 2000; Almeida et al., 2013). However, the origin of this

processing difference and consequent perceptual differences between manipulable and non-

manipulable objects is poorly understood.

One possibility by which objects imbue semantic content related to manipulation is

implied manipulability. A potentially provocative investigation suggested that hand proximity (a

proxy for manipulability), leads to differences similar to the perceptual differences between

manipulable and non-manipulable objects (Gozli et al., 2012). Implied manipulability was

induced by varying the positions of participants' hands such that hands near the display implied

manipulability, while hands far from the display implied low manipulability. Importantly,

participants were presented with a simple circle shape, rather than manipulable or non-

manipulable objects. It was observed that when the participants' hands were positioned near the

experimental stimuli, participants' sensitivity to temporal blinks was heightened and when

participants' hands were far from the stimuli sensitivity to spatial gaps was heightened. In effect,

the placement of the participants' hand in this paradigm shifted the perception of the stimuli to

resemble either the non-manipulable objects, if the hands were far from the stimuli, or the

manipulable objects, if the hands were near the stimuli. It may be that when an object is learned

to be a manipulable, the neural representation of that object becomes associated with the near

hand effect reported by Gozli et al., (2012).

A possible mechanism linking the near-hand effect reported by Gozli et al. (2012) and the

perceptual difference between manipulable and non-manipulable objects derive from the bimodal

cells in the anterior parietal cortex (Graziano & Gross, 1993). The bimodal cells have a

somatosensory receptive field covering a part of the hand and they have a visual receptive field

corresponding to a part of the visual field near their associated hand area. When an object is

shown, if that object does not have a strong action association, then the activity of the bimodal

cells is unaffected. After the organism has learned to use the object with their hands, the bimodal

cells will respond solely to the visual presentation of the object even in the absence of

somatosensory input (Zhou & Fuster, 2000). During object training, it has been shown that the

anterior parietal regions which contain the bimodal cells upregulate brain-derived neurotrophic

factor (BDNF), a protein which encourages the development of new synapses (Ishibashi et al.,

2002). The upregulation of BDNF may be what changes the circuit structure around the bimodal

cells to enable purely visual activation without the corresponding somatosensory input. Thus, the

bimodal cells and their ability to become solely visually activated may be a driver of the known

object selectivity of the dorsal processing pathway (Vaziri-Pashkam & Xu, 2017; Kastner et al.,

2017; Freud et al., 2016) and lead to the dorsal pathway processing bias evoked by manipulable

objects (Chao & Martin, 2000). The object-specific neural activity in the dorsal processing

pathway, divorced from its need for somatosensory input, and largely derived from

magnocellular input, could then be read out during an identification task with high temporal

resolution, leading to heightened temporal precision for manipulable objects.

Taken together, our results demonstrate that object semantic knowledge determines the

processing bias of the object (via dorsal or ventral bias) and evokes subsequent behavioral

repercussions for perception and for action. Our work underscores the need for careful

consideration of object semantic knowledge and its subsequent possible bias to either dorsal or

ventral pathway processing when object images are used not only in psychological research, but

in applied settings such as display and product designs, environmental design, and in the design

of various cognitive assistants.

Methods

Participants. All participants were recruited from The George Washington University participant pool, gave informed consent according to the George Washington University's institutional review board (IRB), were naïve to the purpose of the experiment, and all reported normal or corrected-to-normal vision. In experiment 4, which utilized color stimuli no participant reported colorblindness.

For experiment 1, 26 undergraduate students (19 female; average age: 19.2; 6 left-handed) were recruited. No participant was removed from the analyses. For experiment 2, 27 undergraduate students were recruited. Twenty-six participants' data (14 female; average age: 19.0; 1 left-handed) were analyzed after one participant was cut for below chance accuracy in one of the conditions. For experiment 3, 33 undergraduate students were recruited. Thirty participants' data (26 female; average age: 19.07, 3 left-handed) were analyzed after eight participants were cut for below chance accuracy in at least one of the conditions. For experiment 4, 34 undergraduate students were recruited. Twenty-five participants' data (14 female; average age: 19.0; 1 left-handed) were analyzed after six participants were cut for not finishing the experiment and nine participants were cut for having less than chance accuracy in at least one of the conditions. For experiment 5, 26 undergraduate students were recruited. Twenty-five participants' data (18 female; average age: 19.5, 0 left-handed) were analyzed after one participant was cut for chance accuracy in all conditions.

Apparatus and Stimuli. All experiments were presented on a 24" Acer GN246 HL monitor with a refresh rate of 144 Hz, positioned at a distance of 60 cm from the viewer in a dark room. The experiment was generated and presented using PsychoPy v1.82.

The stimuli in Experiments 1, 2, 4, and 5 consisted of line drawings of real-world, everyday

objects drawn both from The Noun Project, an online repository of object icons and clip art

(https://thenounproject.com). The object stimuli consisted of 10 line-drawings of manipulable

objects, and 10 line-drawings of non-manipulable objects. The manipulable objects were as

follows: snow shovel, handsaw, plunger, screwdriver, hammer, wrench, knife, bottle opener,

spatula, and mug. The non-manipulable objects were: fire hydrant, picture frame, window, toilet,

candle, garbage can, water fountain, potted plant, fan, and lamp. The stimuli were displayed as

large as possible in a 4° x 4° area and all stimuli were presented in black (HSV = 0, 0, 0) on a dark

gray background (HSV = 0, 0, 50). All objects are displayed in Figure 1.

Objects were controlled for low-level differences by calculating the mean luminance, size,

and aspect ratio (i.e., measure of elongation) for each object. There was no mean luminance

difference between object groups, an independent samples t-test ($t(18) = -0.960$, $p = 0.350$). There

was no significant difference in mean size (t-test ($t(18) = 1.043$, $p = 0.311$), or aspect ratio ($t(18)$

$= 1.209$, $p = 0.242$). The width of the bottom line was controlled for each object on which the gap

appeared. There was no significant difference between manipulable and non-manipulable objects

in bottom line width in an independent samples t-test ($t(18) = 0.922$, $p = 0.369$).

For Experiment 3, the stimuli consisted of images of real-world, everyday objects drawn

from Google image search and manipulated in GIMP 2.10.4. The object stimuli consisted of 10

images of the same manipulable objects, and 10 images of the same non-manipulable objects used

in Experiments 1,2, 4, and 5. The stimuli were displayed as large as possible in a 4° x 4° area and

all stimuli were presented in color on a dark gray background (HSV = 0, 0, 50). All objects are

displayed in Figure 1.

Additionally, colored objects used in Experiment 3 were similarly controlled for low-level

differences by calculating the mean size, hue, saturation, value, aspect ratio, and visual clutter. In

analyzing mean size, there was no significant difference between manipulable and non-

manipulable objects in an independent samples t-test ($t(18) = -1.027$, $p = 0.318$). Additionally,

there were no significant differences for hue ($t(18) = -0.309$, $p = 0.761$), saturation ($t(18) = -0.031$,

$p = 0.976$), value ($t(18) = 0.032$, $p = 0.975$), aspect ratio ($t(18) = -0.094$, $p = 0.927$), or visual

clutter ($t(18) = 0.616$, $p = 0.545$). Additionally, we controlled the width of the bottom line of each

object on which the gap would appear. There was no significant difference between manipulable

and non-manipulable objects in bottom line width in an independent samples t-test ($t(18) = 0.557$,

$p = 0.584$).

Design and Procedure. In Experiment 1, each trial began with a single central fixation point,

which subtended a visual angle of 1° x 1°. The central fixation point was rendered in white (HSV

= 0, 0, 100). After 1000 ms of fixation presentation, a single object line drawing would appear 4°

to the left or right of the fixation point. The side of presentation was counterbalanced such that

each participant saw an equal number of stimuli on the left and on the right. The stimuli were

displayed as large as possible within a 4° x 4° area and all stimuli were presented in color on a

dark gray background (HSV = 0, 0, 50). All objects are displayed in Figure 1.

In one half of the experiment, the object appeared with or without a spatial gap in the center

of the bottom line of the object. The objects had an equal probability of having a spatial gap or not

having a spatial gap. The objects appeared for 100 ms and participants were to report the presence

of the spatial gap by pressing the right control button (present) or the left control button (absent)

on the keyboard. Feedback was presented on incorrect trials only.

453   A staircase procedure was used to calibrate the size of the gap to each object to ensure that

454 the gap was equally perceptible across each object regardless of each objects' individual

455 characteristics (Pelli and Bex, 2013). Sixteen undergraduate students (13 female; average age: 18.8;

456 4 left-handed) from the George Washington University participated in exchange for course credit.

457 In each trial the object would be presented with or without the spatial gap which would begin with

458 a size of $0.025°$ visual angle. If the gap was detected correctly for two consecutive trials, the gap

459 decreased by one $0.005°$ step. If the gap was missed, or a false alarm was made to the absence of

460 a gap, for two consecutive trials, the gap was increased by one $0.005°$ step. The gap was calibrated

461 until twenty trials had been completed or until the staircase reversed direction (2 correct followed

462 by 2 incorrect or vice versa) three times. The gap size for each object is displayed in figure 1.

463   In the other half of the experiment, the object appeared with or without a temporal blink.

464 The objects had an equal probability of having a temporal blink or not having a temporal blink.

465 The blinks were 16 ms long. The object first appeared for 96 ms, followed by the temporal blink,

466 and the object then appeared a second time for 32 ms. Participants were to report the presence of

467 the temporal blink by pressing the right control button and absence of the temporal blink by

468 pressing the left control button on the keyboard. Feedback was presented on incorrect trials only.

469   The presentation of stimulus type was counterbalanced across participants such that half of

470 the participants were first presented with spatial gaps and half of the participants were presented

471 with temporal blinks first. Participants completed a total of 720 trials broken into 6 blocks, 360

472 trials of spatial gaps and 360 trials of temporal blinks.

473   Experiment 2 was identical to Experiment 1 except each stimulus was presented upside

474 down. The spatial gap was presented in the same place as it was in experiment 1 relative to the

475 object.

476   Experiment 3 was identical to experiment 1, except more realistic images were used instead

477 of line drawings. Similarly to the line drawing stimuli, a staircase procedure was used to calibrate

478 the size of the gap to each object to ensure that the gap would be equally perceptible across each

479 object regardless of each objects' individual characteristics (Pelli and Bex, 2013). Twenty three

480 undergraduate students (18 female; average age: 18.9; 4 left-handed) from the George Washington

481 University participated exchange for course credit. In each trial the object would be presented with

482 or without the spatial gap which would begin with a size of 0.025° visual angle. If the gap was

483 detected correctly for two consecutive trials, the gap would be made smaller by one 0.005° step.

484 If the gap was not detected, or a false alarm was made to the absence of a gap, for two consecutive

485 trials, then the gap would be made larger by one 0.005° step. Calibration was done until twenty

486 trials had been completed or until the staircase reversed direction (2 correct followed by 2 incorrect

487 or vice versa) three times. The gap size for each object is displayed in figure 1.

488   Experiment 4 was similar in trial structure to the temporal blink condition of Experiment

489 1, except instead of temporal blink detection each trial had a temporal blink that was either short

490 or long in duration. The objects had an equal probability of having a short or a long temporal

491 blink. In the short blink condition, the object first appeared for 96 ms, followed by a 16 ms

492 temporal blink, and then the object appeared a second time for 32 ms. In the long blink

493 condition, the object first appeared for 64 ms, followed by a 48 ms temporal blink, and then the

494 object was presented for a second time for 32 ms. Participants were to report a presence of a

495 short temporal blink by pressing the 'c' key and a long temporal blink by pressing the 'm' key on

496 the keyboard. Feedback was presented exclusively on incorrect trials. Participants completed a

497 total of 720 trials broken into 6 blocks, 360 trials of short blink duration and 360 trials of long

498 blink duration.

Experiment 5, was identical to the spatial gap procedure of Experiment 1 except the

background color was either green (HSV = 110, 30, 90) or red (HSV = 5, 30, 90). The background

color was counterbalanced across participants such that half of the participants would first be

presented with the green background and half of the participants would first be presented with the

red background. Participants completed a total of 720 trials broken into 6 blocks, 360 trials of

green backgrounds and 360 trials of red backgrounds.

Data Analyses. Data for each experiment were analyzed using repeated-measures analysis of

variance with an α set to 0.05. Two-sample t-test were also employed where indicated in the text

and were two-tailed. Anticipatory response times faster than 250 ms and any response times

longer than 2000 ms were removed from analysis (2.67, 2.4, 5.59, 12.1, and 2.95% of the data

was removed from experiments 1, 2, 3, 4, and 5 respectively). Mean response times for each

condition in each experiment were calculated for each subject. For each analysis, all

assumptions, including normality and equality of variances, were formally tested and confirmed.

Code availability. The custom Python programs generated for this study are available from the

corresponding authors on reasonable request.

Data availability. The data that support the findings of this study are available from the

corresponding author upon reasonable request.

References

Almeida, J., Mahon, B. Z., Nakayama, K., & Caramazza, A. (2008). Unconscious processing

dissociates along categorical lines. *Proceedings of the National Academy of Sciences of the*

United States of America, *105*(39), 15214–15218. https://doi.org/10.1073/pnas.0805867105

Almeida, J., Mahon, B. Z., & Caramazza, A. (2010). The role of the dorsal visual processing

stream in tool identification. *Psychological Science: A Journal of the American*

Psychological Society / APS, *21*(6), 772–778. https://doi.org/10.1177/0956797610371343

Almeida, J., Fintzi, A. R., & Mahon, B. Z. (2013). Tool manipulation knowledge is retrieved by

way of the ventral visual object processing pathway. *Cortex*, *49*(9), 2334–2344.

https://doi.org/10.1016/j.cortex.2013.05.004

Baizer, J. S., Ungerleider, L. G., & Desimone, R. (1991). Organization of visual inputs to the

inferior temporal and posterior parietal cortex in macaques. *Journal of Neuroscience*, *11*(1),

168–190. https://doi.org/10.1523/jneurosci.11-01-00168.1991

Blasdel, G. G., & Lund, J. S. (1983). Termination of Afferent Axons in Macaque Striate Cortex.

Journal of Neuroscience, *3*(7), 1389–1413.

Chao, L. L., & Martin, A. (2000). Representation of manipulable man-made objects in the dorsal

stream. *NeuroImage*, *12*(4), 478–484. https://doi.org/10.1006/nimg.2000.0635

Chen, J., Snow, J. C., Culham, J. C., & Goodale, M. A. (2018). What Role Does "Elongation"

Play in "Tool-Specific" Activation and Connectivity in the Dorsal and Ventral Visual

Streams? *Cerebral Cortex (New York, N.Y., 1991)*, *28*(4), 1117–1131.

https://doi.org/10.1093/cercor/bhx017

Collaboration, O. S. (2015). Estimating the reproducibility of psychological science. *Science*,

349(6251). https://doi.org/10.1126/science.aac4716

Derrington, A. M., & Lennie, P. (1984). Spatial and temporal contrast sensitivities of neurons in

lateral geniculate nucleus of macaque. *The Journal of Physiology*, *357*(1), 219–240.

https://doi.org/10.1113/jphysiol.1984.sp015498

DeYoe, E. A., & van Essen, D. C. (1988). Concurrent processing streams in monkey visual

cortex. *Trends in Neurosciences*, *11*(5), 219–226. https://doi.org/10.1016/0166-

2236(88)90130-0

Diamond, R., & Carey, S. (1986). Why Faces Are and Are Not Special. An Effect of Expertise.

Journal of Experimental Psychology: General, *115*(2), 107–117.

https://doi.org/10.1037/0096-3445.115.2.107

Fang, F., & He, S. (2005). Cortical responses to invisible objects in the human dorsal and ventral

pathways. *Nature Neuroscience*, *8*(10), 1380–1385. https://doi.org/10.1038/nn1537

Ferrera, V. P., Nealey, T. A., & Maunsell, J. H. R. (1992). Mixed parvocellular and

magnocellular geniculate signals in visual area V4. *Nature*, *358*(6389), 756–758.

https://doi.org/10.1038/358756a0

Freud, E., Plaut, D. C., & Behrmann, M. (2016). 'What' Is Happening in the Dorsal Visual

Pathway. *Trends in Cognitive Sciences*, *20*(10), 773–784.

https://doi.org/10.1016/j.tics.2016.08.003

Gozli, D. G., West, G. L., & Pratt, J. (2012). Hand position alters vision by biasing processing

through different visual pathways. *Cognition*, *124*(2), 244–250.

https://doi.org/10.1016/j.cognition.2012.04.008

Graziano, M. S. A., & Gross, C. G. (1993). A bimodal map of space: somatosensory receptive

fields in the macaque putamen with corresponding visual receptive fields. *Experimental*

Brain Research, *97*(1), 96–109. https://doi.org/10.1007/BF00228820

Ishibashi, H., Hihara, S., Takahashi, M., Heike, T., Yokota, T., & Iriki, A. (2002). Tool-use

learning selectively induces expression of brain-derived neurotrophic factor, its receptor

trkB, and neurotrophin 3 in the intraparietal multisensory cortex of monkeys. *Cognitive*

Brain Research, *14*(1), 3–9. https://doi.org/10.1016/S0926-6410(02)00056-3

James, T. W., Culham, J., Humphrey, G. K., Milner, A. D., & Goodale, M. A. (2003). Ventral

occipital lesions impair object recognition but not object-directed grasping: An fMRI study.

Brain, *126*(11), 2463–2475. https://doi.org/10.1093/brain/awg248

Karnath, H. O., Rüter, J., Mandler, A., & Himmelbach, M. (2009). The anatomy of object

recognition - Visual form agnosia caused by medial occipitotemporal stroke. *Journal of*

Neuroscience, *29*(18), 5854–5862. https://doi.org/10.1523/JNEUROSCI.5192-08.2009

Kastner, S., Chen, Q., Jeong, S. K., & Mruczek, R. E. B. (2017). A brief comparative review of

primate posterior parietal cortex: a novel hypothesis on the human toolmaker.

Neuropsychologia, *105*, 123–134.

https://doi.org/10.1016/j.neuropsychologia.2017.01.034.A

Kravitz, D. J., Saleem, K. S., Baker, C. I., & Mishkin, M. (2011). A new neural framework for

visuospatial processing. *Nature Reviews Neuroscience*, *12*(4), 217–230.

https://doi.org/10.1038/nrn3008

Kravitz, D. J., Saleem, K. S., Baker, C. I., Ungerleider, L. G., & Mishkin, M. (2013). The ventral

visual pathway: An expanded neural framework for the processing of object quality. *Trends*

in Cognitive Sciences, *17*(1), 26–49. https://doi.org/10.1016/j.tics.2012.10.011

Leonova, A., Pokorny, J., & Smith, V. C. (2003). Spatial frequency processing in inferred PC-

and MC-pathways. *Vision Research*, *43*(20), 2133–2139. https://doi.org/10.1016/S0042-

6989(03)00333-X

Leventhal, A. G., Rodieck, R. W., & Dreher, B. (1981). Retinal ganglion cell classes in the old

world monkey: Morphology and central projections. *Science*, *213*(4512), 1142–1144.

https://doi.org/10.1126/science.7268423

Livingstone, M. S., & Hubel, D. H. (1988). Do the relative mapping densities of the magno- and

parvocellular systems vary with eccentricity? *Journal of Neuroscience*, *8*(11), 4334–4339.

https://doi.org/10.1523/jneurosci.08-11-04334.1988

Mahon, B. Z., Milleville, S. C., Negri, G. A. L., Rumiati, R. I., Caramazza, A., & Martin, A.

(2007). Action-Related Properties Shape Object Representations in the Ventral Stream.

Neuron, *55*(3), 507–520. https://doi.org/10.1016/j.neuron.2007.07.011

Maunsell, J. H. R., Ghose, G. M., Assad, J. A., McAdams, C. J., Boudreau, C. E., & Noerager,

B. D. (1999a). Visual response latencies of magnocellular and parvocellular LGN neurons

in macaque monkeys. *Visual Neuroscience*, *16*(1), 1–14.

https://doi.org/10.1017/S0952523899156177

Maunsell, J. H. R., Ghose, G. M., Assad, J. A., McAdams, C. J., Boudreau, C. E., & Noerager,

B. D. (1999b). Visual response latencies of magnocellular and parvocellular LGN neurons

in macaque monkeys. *Visual Neuroscience*, *16*(1), 1–14.

https://doi.org/10.1017/S0952523899156177

Maunsell, J. H. R., Nealey, T. A., & DePriest, D. D. (1990). Magnocellular and parvocellular

contributions to responses in the middle temporal visual area (MT) of the macaque monkey.

Journal of Neuroscience, *10*(10), 3323–3334. https://doi.org/10.1523/jneurosci.10-10-

03323.1990

McAnany, J. J., & Alexander, K. R. (2008). Spatial frequencies used in Landolt C orientation

judgments: Relation to inferred magnocellular and parvocellular pathways. *Vision*

Research, *48*(26), 2615–2624. https://doi.org/10.1016/j.visres.2008.02.012

Mishkin, M., & Ungerleider, L. G. (1982). Contribution of striate inputs to the visuospatial

functions of parieto-preoccipital cortex in monkeys. *Behavioural Brain Research*, *6*(1), 57–

77. https://doi.org/10.1016/0166-4328(82)90081-X

Nassi, J. J., & Callaway, E. M. (2009). Parallel processing strategies of the primate visual

system. *Nature Reviews Neuroscience*, *10*(5), 360–372. https://doi.org/10.1038/nrn2619

Noppeney, U., Price, C. J., Penny, W. D., & Friston, K. J. (2006a). Two distinct neural

mechanisms for category-selective responses. *Cerebral Cortex*, *16*(3), 437–445.

https://doi.org/10.1093/cercor/bhi123

Noppeney, U., Price, C. J., Penny, W. D., & Friston, K. J. (2006b). Two distinct neural

mechanisms for category-selective responses. *Cerebral Cortex*, *16*(3), 437–445.

https://doi.org/10.1093/cercor/bhi123

Perry, V. H., Oehler, R., & Cowey, A. (1984). Retinal ganglion cells that project to the dorsal

lateral geniculate nucleus in the macaque monkey. *Neuroscience*, *12*(4), 1101–1123.

https://doi.org/10.1016/0306-4522(84)90006-X

Pokorny, J., & Smith, V. C. (1997). Psychophysical signatures associated with magnocellular

and parvocellular pathway contrast gain: comment. *Journal of the Optical Society of*

America A, *14*(9), 2477–2486. https://doi.org/10.1364/josaa.15.002440

Vaziri-Pashkam, M., & Xu, Y. (2017). Goal-directed visual processing differentially impacts

human ventral and dorsal visual representations. *Journal of Neuroscience*, *37*(36), 8767–

8782. https://doi.org/10.1523/JNEUROSCI.3392-16.2017

Wiesel, T. N., & Hubel, D. H. (1966). Spatial and chromatic interactions in the lateral geniculate

body of the rhesus monkey. *Journal of Neurophysiology*, *29*(6), 1115–1156.

https://doi.org/10.1152/jn.1966.29.6.1115

Zhou, Y. di, & Fuster, J. M. (2000). Visuo-tactile cross-modal associations in cortical

somatosensory cells. *Proceedings of the National Academy of Sciences of the United States*

of America, *97*(17), 9777–9782. https://doi.org/10.1073/pnas.97.17.9777

Acknowledgements

This work was supported by a National Science Foundation grant no. BCS-1921415 and BCS-2022572 to S.S. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. We thank D. Kravitz and M. Behrmann for insightful comments on an earlier version of this manuscript. Special thanks go to E. Awh for suggesting a temporal manipulation adopted for Experiment 4.