Evolved changes in DNA methylation in response to the sustained loss of parental care in the burying beetle

Mashoodh R[1]*, Sarkies P[2,3], Westoby J[4], Kilner RM[1]

[1]Department of Zoology, University of Cambridge, United Kingdom

[2]MRC London Institute of Medical Sciences, London, UK

[3]Institute of Clinical Sciences, Imperial College London, UK

[4]Department of Genetics, University of Cambridge, United Kingdom

*Corresponding author

Email: rm786@cam.ac.uk

Abstract

Levels of parental care critically influence the development environment with the capacity to impact the growth, survival, physiology, and behaviour of offspring. Plastic changes in DNA methylation have been hypothesised to modulate gene expression responses to parental environments. Moreover, these effects can be inherited and so may affect the process of adaptive evolution. In this study, using experimental evolution, we investigated how plastic changes in DNA methylation induced by the loss of parental care have evolved in a biparental insect (*Nicrophorus vespilloides*) using experimental evolution. We show that removal of care in a single generation is associated with changes in gene expression in stress-related pathways in 1st instar larvae. However, in larvae that have adapted to the loss of parental care after being deprived of care for 30 generations, gene expression is shifted from stress-related gene expression towards growth and brain development pathways. We found that changes in gene body methylation arose both as a direct response to the loss of parental care and stochastically as populations diverged. Overall, our results suggest that a complex interplay between transcription and DNA methylation shapes the molecular adaptation to environmental change.

Introduction

There has been increasing interest in the idea that epigenetic mechanisms could participate in evolutionary change. Epigenetic mechanisms are the biochemical marks (*e.g.*, DNA methylation, histone modifications) that ultimately determine how accessible DNA is to factors within the cell which, in turn, can regulate gene expression (Holliday, 2006; Jaenisch & Bird, 2003). These processes have been shown to be sensitive to the environment and represent a mechanism through which plasticity can be achieved during development. A hypothesised role for epigenetics in adaptation and evolution is driven primarily from recent work in plants and animals (*e.g.*, *Arabidopsis*, *C. elegans*, mouse) demonstrating the capacity for transgenerational epigenetic inheritance *via* epigenetic modifications. These studies have shown that chronic challenges with stress, dietary changes or chemicals/drugs induce heritable changes in epigenetic marks in the germline that are transmitted to subsequent generations (Curley et al., 2011; Miska & Ferguson-Smith, 2016; Sarkies, 2020). The capacity for such inheritance and the degree to which it persists is highly dependent on the organism and the nature of the epigenetic change in question. Nevertheless, these observations raise the possibility that epigenetic mechanisms could buffer and/or modulate the pace of evolutionary change under changing environments. Indeed, theoretical models have supported this idea (Badyaev & Uller, 2009; Bondurianský & Day, 2018; Miska & Ferguson-Smith, 2016).

Although these ideas have generated much enthusiasm (and controversy), there is nevertheless limited empirical evidence to support the idea that epigenetic marks are induced by environments and persist in a way that drives long-term adaptive evolutionary change. There have been many observations of population-level differences in epigenetic marks between locally adapted wild populations in a range of diverse taxa (Dixon et al., 2018; Dubin et al., 2015; Heckwolf et al., 2020; Liew et al., 2020; Skinner et al., 2014) which are often correlated with local abiotic environmental conditions (*e.g.*, salinity, temperature, latitude). The accumulation of methylation changes is suggestive of a potential role in adaptation to changing environments. However, there are a number of pathways through which epigenetic marks could participate in adaptative processes and contribute to the emergence of population-level changes. For example, epigenetic marks could be a by-product of transcriptional states or genetic sequence change (Dixon et al., 2018; Dubin et al., 2015; Hu et al., 2021). In this scenario epigenetic modifications are effectors and/or markers rather than drivers of the adaptive change. Taken together, epigenetic modifications could result from a complex interplay between plastic, genetic and stochastic processes acting in the short- and/or long-term (Sarkies, 2020). A critical first step towards understanding the sources of epigenetic variation is to characterise the extent to which epigenetic changes respond to an initial change in the environment, and the degree to which

such changes might persist and be correlated with phenotypic changes (*i.e.*, gene
expression) as populations adapt to a changing environment.
Here, we used experimental evolution to examine how patterns of gene expression
and DNA methylation are induced and evolve in populations of burying beetles as they adapt
to the removal of post-hatching parental care (*i.e.,* a change in the social environment). In
natural populations of this locally abundant insect, burying beetle parents raise their young
on a carrion nest (formed from a small dead animal, such as a mouse or songbird). There is
continuous variation in the level of parental care provisioned (Jarrett, Evans, et al., 2018). At
one extreme, parents tend to their offspring throughout their development (including feeding
with regurgitated pre-digested carrion) whereas at the other, parents abandon their offspring,
leaving them to fend for themselves without any post-hatching care (Eggert et al., 1998).
Burying beetles are highly responsive to the levels of parental care they receive within a
single generation (Kilner et al., 2015). We have exploited this natural variation in care to
establish two types of experimentally evolving populations in the laboratory (each replicated
twice), which vary only in the family environment that larvae experience during development,
and where the same family environment is created for successive generations within
populations. In Full Care populations (FC_{POP}), parents remain with their young throughout
development; whereas in No Care populations (NC_{POP}), parents are removed just before
their offspring hatch (see Schrader et al., 2017; for more details on the experimental setup).
Previous work has found that that NC_{POP} populations rapidly adapt to a life without parental
care such that by generation 13 the proportion of successful broods matches that of the
FC_{POP} populations (Schrader et al., 2015, 2017). This adaptation to the loss of parental care
is associated with a number of changes in morphology and behaviour. For example, NC
offspring have evolved larger mandibles which likely helps with feeding on the carcass in the
absence of parents (Jarrett, Evans, et al., 2018), hatch more synchronously (Jarrett, Rebar,
et al., 2018) and cooperate more with each other (Rebar et al., 2020).
DNA-Me is a highly conserved epigenetic mark across plants and animals. This
involves the covalent addition of methyl groups to CG dinucleotides (CpG) within DNA. DNA
methylation (mCpG) is added by DNA methyltransferases (DNMTs 1-3; *e.g.*, for
maintenance during cell division and *de novo*) and, in mammals, can be actively removed by
ten-eleven translocation enzymes (TETs) (Law & Jacobsen, 2010). Although parental care
(and its loss) has been associated with plasticity in both DNA-Me and gene expression
across a number of species (Arsenault et al., 2018; Franklin et al., 2010; Nyman et al., 2020;
Weaver et al., 2006), its role in insect DNA-Me and gene expression plasticity is still poorly
understood. Burying beetles have a functioning DNA-Me system with levels of methylation
that are comparable to other holometabolous insects and possess single copies of DNMT1-3
(Bewick et al., 2016; Cunningham et al., 2015; Lewis et al., 2020). These enzymes are

highly conserved across multiple taxa and methylation occurs in actively transcribed genes suggesting important, yet still poorly understood, roles for DNA-Me in the species that carry this epigenetic modification (Feng et al., 2010; Schmitz et al., 2019).

In this study, we first characterized the changes in gene expression might underpin adaptation to the NC environment in developing larvae. To distinguish between evolved and environmentally-induced (plastic) changes, we used a common garden approach where offspring were exposed to the reciprocal parental environment (*i.e.,* FC_{POP} had parents removed to create a NC environment or parents remained post-hatching in NC_{POP} to create a FC environment). We then asked if changes in DNA methylation are correlated with adaptive divergence (see Figure 1a for experimental design summary).

Results

Differential gene expression in response to the sustained removal of care

To quantitatively characterise genome-wide variation in larval gene expression between our populations, we conducted a discriminant analysis of principal components (DAPC) using all expressed genes (n=12,772) within samples. The removal of care was significantly associated with a shift in gene expression (environment posterior mean = 4.22, 95% CI = [3.35, 5.15], p_{MCMC} < 0.001; Figure 1b) and this response was true whether individuals were derived from the FC or NC populations, as there was no effect of population (population posterior mean = 0.27, 95% CI = [-0.63, 1.20], p_{MCMC} = 0.576) nor an interaction between population of origin and the current environment (posterior mean = -7.91, 95% CI = [-2.12, 0.40], p_{MCMC} = 0.210). Further, each replicate responded to the care treatment in a similar way, since there was no effect of block on gene expression plasticity (posterior mean = 0.03, 95% CI = [-0.6, 0.72], p_{MCMC} = 0.940). Taken together, these results suggest that while there were large shifts in gene expression in response to the absence of parental care in the current environment, evolved changes in gene expression are likely to be more subtle and/or involve smaller subsets of genes.

To investigate these differences in more detail we performed differential expression analyses using $FC_{POP}FC_{ENV}$ as the reference population to estimate the number of differentially expressed genes (DEGs) across conditions. We found that 657 genes were differentially expressed when parental care was removed for the first time in the FC_{POP} (n=657; all $\log_2$ fold changes > 1; Figure 1c). By contrast, in the NC_{POP}, which had experienced a NC_{ENV} for the previous 29 generations, 385 genes were differentially expressed which is a significant reduction in response (X^2(3, n=12772) = 21558, p<2 x 10^{-16}). Interestingly, only 20 genes were differentially expressed when NC_{POP} parents were allowed to care for their offspring for the first time in 30 generations, suggesting that gene

expression was not constrained by their exposure to divergent selection pressures during
these generations.
There was significant overlap in identity of the genes being differentially expressed in
$FC_{POP}NC_{ENV}$ compared to the $NC_{POP}NC_{ENV}$ ($\log_2$(OR)=122.4265, $p<2,2 \times 10^{-16}$), with many
more changes being lost (n=354) than being gained (n=82; Figure 1d). To test whether this
reflected changes in the overall strength of response by individual genes to a NC
environment within each population, we extracted overlapping DEGs (even those of small
effect) and correlated the magnitude of response at each individual gene. This is the
equivalent of comparing reaction norms on a genome-wide scale (See Figure 1e inset). We
predicted that if the magnitude of change in response to a NC environment was similar
across populations then we would expect fold-changes to be highly correlated resulting in a
slope close to 1. We found that genes that were differentially expressed in response to a
NC_{ENV} in both populations were positively correlated, with no evidence that genes switched
direction of expression after evolving without parental care (slope = 0.489, t(3103)=91.875,
$p<2.2 \times 10^{-16}$; Figure 1e). However, the slope of this correlation was significantly less than a
slope of 1 (F(2,3103)=9176.4, $p<2.2 \times 10^{-16}$). Therefore, while there were similar changes
elicited by a NC_{ENV} across both populations, the evolved NC populations expressed those
genes at lower levels on average than the FC_{POP}.
We extracted significant gene ontology (GO) terms from each contrast and collapsed
across significant, but highly redundant terms, to look for overarching patterns. This analysis
revealed that changes in gene expression primarily occurred in three broad categories of
genes: 1) stress and its cellular response, 2) immune function and 3) growth and
development (see Figure 1f for examples of GOs falling within these categories). Taken
together, these results suggest that whilst the initial response to the removal of care is
associated with an upregulated stress response and increased immune defense, individuals
selected under a NC regime express fewer genes associated with these two categories.
These changes are accompanied by enhanced expression of genes associated with
physiological and neurobiological development (Figure 1g). Moreover, further examination of
gene expression changes that emerge after 30 generations of exposure to a NC
environment reveal an over-representation in the neurotransmission and neuropeptide
categories (*e.g.,* neuropeptide Y, orexin, and glutamate). For full list of GO categories see
Supplementary Tables.
174
CpG methylation in the burying beetle
We first sought to characterise the methylomes of first instar burying beetle larvae.
We found average CpG methylation across the whole genome to be between 0.7-0.8% for
all samples. Average levels of methylation were significantly higher in gene bodies (3.64 ±

8.60) compared to coding sequences (CDS; 2.72 ± 10.8), introns (0.952 ± 4.44), repeat/TEs

(0.523 ± 1.96) and upstream regions (5' UTR; 0.659 ± 3.22; all FDR-corrected Wilcoxon p

values < 2.2×10^{-16}; Figure 2a). Moreover, we found that gene body methylation was weakly

but positively correlated with gene expression ($t(1,11920)=36.34$, $p<2.2 \times 10^{-16}$, $R^2 = 0.10$).

When we classified genes into expression state categories (silent, low, medium or highly

expressed; Supp Figure 1a), we found that this correlation was mainly driven by expression

states that were classified as medium or low. Medium and high levels of expression did not

have significantly different methylation values from each other (Figure 2b, Supp Figure 1b).

These findings are consistent with previous reports of methylation residing primarily in gene

bodies for this species and the presence of higher methylation levels in active genes in

plants and animals (Cunningham et al., 2015; Feng et al., 2010; Lewis et al., 2020).

Next, we determined whether the removal of care resulted in methylation changes in

specific genomic features. We looked for enrichment of differentially methylated CpGs over

the expected distribution of methylated CpGs across genomic features (Figure 2b). A CpG

was classified as being methylated if the logit probability of it being methylated was greater

than the bisulfite conversion error rate in at least 1 sample (see Methods). The density of

methylated CpGs was higher in gene bodies (Figure 2c; $X^2(5, n=7400) = 14786$, $p<2.2 \times 10^{-16}$)

and we found differentially methylated CpGs (in both environmental and evolved contrasts)

to be significantly enriched in genes and, specifically, introns (FDR-corrected p's<0.05;

Figure 1c) but not in any other annotated features tested. We, therefore, chose to focus on

gene body methylation for subsequent differential methylation analyses.

Differential gene body methylation in response to the sustained removal of care

We performed differential methylation analysis across gene bodies to compare (1)

DNA-Me between populations (evolved changes; sustained exposure to a NC_{ENV} over 30

generations) with (2) DNA-Me induced by a single generation of exposure to a NC_{ENV} within

the FC_{POP} (proxy for initial response) and (3) DNA-Me within the NC_{POP} when exposed to a

FC_{ENV} (population-specific responses to NC_{ENV}). As has been previously reported, not all

genes in the burying beetle are methylated (Cunningham et al., 2015; Lewis et al., 2020).

Therefore, we clustered average weighted methylation values (calculated across the entire

gene body), identified two distinct distributions of methylation states within genes (see

Methods) and focussed our analyses on those that were classified as methylated (Supp

Figure 2a, n = 4431; 39% of genes for which methylation was present for 85% of samples).

These genes were also more highly expressed (W = 9161933, $p<2.2 \times 10^{-16}$, Supp Figure 2b)

confirming that this procedure extracted functionally-relevant methylated genes.

We first compared the initial environmental response to a NC environment (within the

FC_{POP}) to the response 30 generations later (evolved differences). Removal of care in

response to one generation of a NC_{ENV} resulted in 1486 differentially methylated genes,

whereas we found 1530 differences when comparing the evolved response to the sustained

loss of care (all genes FDR-corrected $p<0.01$). We also assessed the overall false discovery

rate empirically by randomly shuffling samples between conditions 1000 times and

extracting the number of differentially methylated genes (DMGs) after FDR correction at

each shuffle. The number of DMGs detected in each comparison far exceeded the number

that would be expected at random (p_{FDR} = 0 for both comparisons; Figure 3a). Moreover, the

response to the removal of care was highly similar within each population (overlap of 741

DMGs; $log_2(OR)=1.8$, $p<2.2x10^{-16}$), though a high proportion of these changes remained

specific to populations (Supp Figure 3). Taken together, this suggests that epigenetic

variation can arise from environmental, population genetic background as well as an

interaction between the two.

Differential gene body methylation and transcriptional change

Of the DMGs that were altered in response to a single generation of exposure to NC,

51% were significantly associated with differences in gene expression (all DEGs FDR-

corrected $p<0.01$; $log_2(OR)=1.21$, $p=0.003$). However, no significant association with gene

expression was identified for responses measured after 30 generations of exposure to NC

($log_2(OR)=1.16$, p=0.27). GO term enrichment analysis revealed that genes differentially

methylated after exposure to one generation of NC were enriched in stress and growth

reduction pathways (Figure 3d). Differentially methylated genes in the evolved response to a

NC_{ENV} were enriched in biological processes related to growth and brain development

(Figure 3e; see Supplementary Tables for a full list of GO terms).

Given the high degree of overlap in DMGs induced by the environment compared to

the evolved response to the NC_{ENV} 48%; Figure 3b), we hypothesised that the latter set of

changes might reflect environmentally-induced changes that are no longer coupled with

expression (i.e., methylation reflects ancestral rather than current transcriptional state). We,

therefore, tested the overlapping DMGs against changes in gene expression induced by a

single generation of exposure to NC. We found that overlapping DMGs are associated with

the initial environmental response ($log_2(OR)=1.17$, $p<0.05$) but not current changes in gene

expression ($log_2(OR)=1.04$, $p=n.s$).

To further investigate the idea that differential methylation can occur in evolved

responses without associated differential gene expression, we highlighted differentially

methylated genes on density plots describing the relationship between expression (FKPM)

and average methylation for all genes. We identified a cluster of genes that was persistently

differentially methylated but had lost differential expression after 30 generations of sustained

exposure to NC (Figure 3c). Differential methylation in this cluster (outlined in Figure 3c) is

associated with differential gene expression in populations exposed to NC for one
generation ($\log_2(OR)$=1.91, p = 0.01) but not in the evolved response ($\log_2(OR)$=0,
p=4.279e-05). Also differentially methylated at this cluster are genes from individuals in the
NC_{POP} that were exposed back to the FC_{ENV} for a single generation suggesting they are
environmentally labile (*i.e.*, not inherited) (Supp Figure 4). We performed a GO analysis on
this cluster and found these genes are almost exclusively enriched in ribosome biogenesis,
protein translation and transport processes (see Figure 3f and Supplementary Tables).
We then asked if there was a possibility that environmental responses could become
fixed evolved differences. In other words, DMGs that were initially responsive to the
environment but no longer environmentally sensitive. We reasoned that such changes would
not be reversed when NC_{POP} were provided with a $FC_{ENV.}$ There were 273 DMGs (18% of all
evolved DMGs) that were differentially methylated in the evolved but not reverted by
exposure of the NC_{POP} to a FC_{ENV}. While these genes were enriched in defense, cellular and
RNA metabolic processes these were not significantly associated with gene expression
changes in larvae ($\log_2(OR)$=0.97, p=ns). This raises the possibility that some heritable
methylation changes could become fixed during adaptive divergence, though the functional
consequences of such changes remain unclear.
270
271   **Discussion**
A role for epigenetic processes in mediating some of the molecular events underlying
adaptive divergence is the subject of intense interest and debate (Bonduriansky & Day,
2018; Charlesworth et al., 2017; Laland et al., 2014; Sarkies, 2020). Debate persists
because empirical studies testing the relationship between early environmentally-induced
epigenetic responses and the subsequent adaptive evolution of such modifications to
changing environments have been rare. How and when such changes are adaptive would
depend on the persistence, heritability and function of epigenetic modifications during
adaptation. Here, using experimental evolution in a wild-caught population of a subsocial
insect, the burying beetle, we describe how gene expression and genome-wide patterns of
DNA methylation diverge as beetles adapt to the experimentally imposed loss of parental
care in the lab. We show that (1) DNA methylation and gene expression is highly sensitive to
the loss of parental care, in both the initial and the evolved response (30 generations later),
(2) The relationship between gene expression and DNA methylation at methylated genes
shifts as populations diverge and (3) There is more variation in DNA methylation between
populations (evolved change) and in population-specific responses to the environment than
is reflected in gene expression profiles. Taken together, these data suggest that

environmental, genetic and stochastic processes contribute to the accumulation of DNA
methylation differences during adaptive divergence.
290
Evolved changes in gene expression
Levels of parental care can shape the development of a number of physiological and
behavioural traits that persist throughout the lifespan, including key fitness-related traits such
as lifespan, fecundity and survival (Gustafsson & Sutherland, 1988; Kilner et al., 2015;
Steiger, 2013). Conversely, the removal of parental care has been shown to be associated
with disruptions to behavioural development *via* changes in gene expression (Franklin et al.,
2010; Weaver et al., 2006). Although much of the molecular level data comes from rodent
model species, there is evidence for similar changes across primates, fishes and even a
subsocial bee suggesting that there might be suites of conserved molecular mediators
(Arsenault et al., 2018; Massart et al., 2014; Nyman et al., 2020). Therefore, it is not
surprising that the experience of a NC_{ENV} in burying beetles is the major source of variation
in larval gene expression particularly in stress response pathways that disrupt growth and
development. Adaptation to the loss of parental care in this species appears to be
associated with the evolution of a stress-tolerant phenotype because we observe (1)
reduced responsiveness of genes in the NC_{POP} to the NC_{ENV} (in terms of levels and number)
and (2) the appearance of a small number of changes that may compensate for stress-
related gene expression. This is consistent with the idea that adaptation to stressful
environments and/or challenges involves fine tuning the balance between stress- and
growth-related gene expression (López-Maury et al., 2008).
We further inspected the genes that were differentially expressed in the NC_{POP} but
not FC_{POP} in response to the NC_{ENV} (*i.e.*, new DEGs). One striking change was the high
expression of a cytochrome P450 gene which appears to be a homolog to the Drosophila
Cyp6a20 gene, the deletion of which has been associated with higher levels of aggression
and reduced sociality (Wang et al., 2008). Also, of particular interest was the differential
expression of a number of odorant binding proteins. These proteins are odour sensing
proteins that have been shown to be involved in a number of behavioural phenotypes. One
such highly expressed gene was a homolog to Drosophila *Obp69a* which is involved in
social responsiveness to social experience as well as starvation (Bentzur et al., 2018;
Farhadian et al., 2012). Changes in the expression of both sets of genes was particularly
interesting given that previous studies have shown that adaptation to the loss of parental
care is associated with a greater level of coordination among larvae (Jarrett, Rebar, et al.,
2018; Schrader et al., 2017) and a shift from sibling conflict to cooperation (Rebar et al.,
2020).
324

Sources of variation in DNA methylation

In rodent model systems, changes in gene expression in developing offspring in response to the loss of parental care have been shown to be, at least in part, mediated by changes in DNA methylation, most notably at promoter regions (Franklin et al., 2010; Massart et al., 2014; Suderman et al., 2012). Here we show that gene body methylation of developing larvae, though less widespread than vertebrate methylation, is similarly sensitive to the loss of parental care in a subsocial insect and that some of these changes are associated with differential gene expression. We further show patterns of DNA methylation diverge as populations adapt to the recurrent loss of parental care over multiple generations. Some of this change is simply induced afresh, each time the population is exposed to an NC environment. We know this because allowing parents to care for NC_{POP} offspring reversed some of these population-specific differences in DNA methylation. Nevertheless, some of the changes in DNA methylation that were induced originally by the environment did appear to become fixed over time in the NC_{POP}. However, while a significant portion of DNA methylation resembles the initial response to the loss of care, many of the changes in DNA methylation that we observed were not explained solely by the removal of care. Some epigenetic changes are short-lived whereas others appear to persist suggesting that there might be functional differences in the timing and duration of such changes throughout the course of adaptation. These findings do not rule out the possibility of adaptive transgenerational inheritance of environmentally-induced methylation but they do suggest that population differences in epigenetic variation could emerge through multiple parallel processes.

Population differences in DNA-Me could be due, at least partially, to genetic variation that emerges through drift and/or selection. For example, in sticklebacks, some but not all population-level variation in DNA-Me is correlated with plasticity-induced by differences in water salinity (Heckwolf et al., 2020). Some of these methylation changes have been linked to *cis* or *trans*-acting loci which show evidence of local adaptation (Hu et al., 2021). Similarly, population level changes in gene body methylation that appear in corals inhabiting different quality environments are associated with higher genetic differentiation at those loci (Dixon et al., 2018). Another explanation is that population differences in DNA methylation could have arisen randomly (*i.e.*, epimutations) and then been selected by the environment. This could be made more possible in insects by the reported high heritability of arthropod gene body methylation (Liew et al., 2020; Yagound et al., 2020). Indeed, theoretical models have shown that stochastic but heritable epigenetic variation can be adaptive under changing environments (Feinberg & Irizarry, 2010). In plants, there is evidence for both stochastic, genetic and environmental sources of DNA methylation in explaining local adaptation (Becker et al., 2011; Dubin et al., 2015; Kawakatsu et al., 2016). This further highlights the

diverse mechanisms through which these changes in DNA methylation could emerge.
Though we were able to distinguish between environmentally-induced and evolved changes
in DNA methylation, this lays the groundwork for distinguishing between other (*e.g.*, genetic,
epimutational) sources of variation and its functional role.

Of particular note is that while a single exposure to the loss of care leads to an
association between DNA methylation and gene expression, no such relationship is
apparent after populations had diverged and the NC_{POP} had become adapted to a life without
parental care. Indeed, over this timeframe, some methylation changes become uncoupled
from differential gene expression, which suggests that adaptation to a NC_{ENV} might involve
buffering against or overriding environmentally-induced changes in DNA methylation *via*
other mechanisms. Our analysis of GO terms suggests that differentially methylated genes
between the environmental and evolved conditions broadly reflect the same functional
categories involved in divergent gene expression patterns. So while the initial response to
the removal of care involves differential methylation in stress-related pathways, evolved
differential methylation is enriched in processes associated with growth and brain
development though this was not always associated with gene expression at the
developmental point at which we measured it. DNA methylation in insects, including the
burying beetle, appears to be stable across the lifespan (Cunningham et al., 2019; Yagound
et al., 2020). This raises the possibility that DNA methylation may be a product of ancestral
transcriptional states. An alternative explanation is that these marks may prime
transcriptional responses for activation at different developmental timepoints. What is clear
is that while DNA methylation may not always reflect current transcriptional states, it
accumulates as populations adapt and diverge and appears to be a marker for past
environmental experience.
386
Conclusions
There is accumulating evidence that populations accrue epigenetic variation that is
correlated with local abiotic conditions, as they become divergently and locally adapted.
These findings, coupled with work showing the transgenerational inheritance of
environmentally-induced modifications (primarily in isogenic lab animals and plants) has
provoked speculation that epigenetic modifications could play a key causal role in
adaptation. An implicit, yet mostly untested, hypothesis here is that epigenetic modifications
are inherited, and sustain adaptive divergence through a form of "plasticity-led" evolution.
Here, we have taken an experimental approach to test this hypothesis. By manipulating
environmental conditions (*i.e.*, the loss of parental care) we have found that a significant
amount of variation in DNA methylation can be induced and this is associated with
transcriptional activity. However, after several generations of exposure to the same

environmental cues, we find that population-level variation and transcription is not solely

explained by the environmental differences. These indicate that additional transcriptional

and stochastic processes contribute to the accumulation of differential DNA methylation as

populations adapt and diverge. In short, although it is still possible that epigenetic variation

contributes to adaptation, existing formulations of how this might work in practice are too

simplistic to account for the experimental results we obtained.

Methods

Breeding design & Experimental Evolution

We analysed experimental populations of *Nicrophorus vespilloides* that had been evolving

under different regimes of parental care, and which were founded from a genetically diverse

founding population generated by interbreeding beetles from multiple wild populations

across Cambridgeshire. These populations are described in detail in Schrader et al., 2017.

and comprise a total of 4 populations, two blocks (Block 1 and Block 2; separated by 1

week) containing two populations evolving with (FC_{POP}) or without parental care (NCpop).

On the 29th generation, seventeen days after their emergence as adults, when individuals

were sexually mature, we paired 15 males and females within each population (N=30 pairs

in total). Each pair was placed in a separate breeding box. Each pair was placed in a

separate breeding box with moist soil and a thawed carcass (10-12g). We then placed each

breeding box in a cupboard, and allowed parents to prepare the carcass and for the female

to lay the clutch of eggs. After 53h, populations were split such that both parents were either

removed (in keeping with the procedure experienced by the NC_{POP}) or left in the breeding

box. This produced offspring from both populations in their evolved condition ($FC_{POP}FC_{ENV}$ &

$NC_{POP}NC_{ENV}$) as well as their reciprocal environments ($FC_{POP}NC_{ENV}$, $NC_{POP}FC_{ENV}$). At 80h

post-pairing (approx. 10h post-hatching), first-instar larvae were collected from surviving

families. Only families that with brood sizes between 20-30 larvae were used for subsequent

RNA-sequencing analyses.

Larval tissue dissection, RNA extraction & Sequencing

For each family, RNA from four heads of first instar larvae were pooled and extracted using

Trizol (Invitrogen). Total RNA quality was checked using the BioAnalyzer System (Agilent),

and yield was quantified using a Qubit RNA Assay Kit (Thermo Fisher). PolyA-selected RNA

libraries were constructed and sequenced (150bp paired-end) at a depth of 30x by

Novogene (Hong Kong). The resulting sample group sizes for gene expression analysis

were: 5-6 libraries for each group within each block (total of 46 libraries).

RNA Sequencing & Read Mapping

Reads were trimmed using TrimGalore (0.5.0; https://github.com/FelixKrueger/TrimGalore)
to remove adaptor sequences, perform quality trimming and discard low-quality reads.
Reads were mapped and quantified using a custom pipeline using HiSat2 (2.1.0) and
Stringtie (2.0.3; Pertea et al., 2016). Transcript abundance estimation was based on
counting reads aligned to the *N. vespilloides* reference transcriptome (NCBI Refseq
Assembly: GCF_001412225.1; Cunningham et al., 2015). Lowly expressed genes (those
with less than 15 counts in more than 90% of samples) were filtered from raw counts table
leaving a total of 12,772 expressed genes in the dataset to be analysed.

DAPC using adegnet and MCMCglmm

To compare overall patterns of gene expression plasticity, we performed a discriminant
analysis of principal components (DAPC), which linearizes principal components within a
dataset into a single discriminant function, enabling the comparison of overall levels of
variation between groups. Gene counts were normalized and log-transformed using a
regularized log transform in DESeq2 (version 1.26.0; Love et al., 2014) and a discriminant
function was built by defining each populations response to their reciprocal environments
using adegenet (version 2.1.3; Jombart, 2008). To compare differences in gene expression
plasticity between populations we used the MCMCglmm package (version 2.29; Hadfield,
2010) to model DAPC score as a function of population background and current
environment (and an interaction between the two) using a default prior that assumes a
normal posterior distribution with large variances for the fixed effects and a weakly
informative (flat) prior.

Differential expression analysis

Differential expression was analysed using DESeq2 using custom scripts in R. Log2-fold
change estimates for each differentially expressed gene (DEG) were shrunk to generate
more conservative estimates of effect size using the ashr shrinkage estimator (Stephens,
2016). Moreover, to focus on genes that are differentially expressed at higher thresholds we
considered a gene to be significant differentially expressed only if it had increased by a 2-
fold change (*i.e.,* $Log_2FoldChange >= 1$). We included the separate blocks as a covariate in
all the analyses to increase power but also account for minor fluctuations in gene expression
across the blocks of replicate populations.

Functional Annotation

Functional enrichment analyses were conducted using the topGO R package (version
2.38.1; Alexa & Rahnenfuhrer, 2016) to identify over-representation of particular functional

groups within the DEGs in response to the removal of care as well as the evolved response

to the removal of care, based on GO classifications using Fisher's exact test. GO terms were

annotated to the *N. vespilloides* genome using the BLAST2GO (version 5.1.1) workflow to

assign homologs to the *Drosophila* non-redundant protein databases (Sun et al., 2020). To

increase functional predictions of *N. vespilloides* genes these annotations were

supplemented with GO term assignments based on ortholog searches within Arthropods

using eggNOG-mapper (Huerta-Cepas et al., 2017) and OrthoDB (v10.1; Kriventseva et al.,

2019).

Bisulfite Sequencing & Read Mapping

For each family, DNA from heads of first instar larvae were pooled and extracted using the

Qiagen DNEasy Blood & Tissues kit (Qiagen). Total DNA quality was checked using the

BioAnalyzer System (Agilent), and yield was quantified using a Qubit DNA HS Assay Kit

(Thermo Fisher). DNA methylation libraries were constructed and sequenced (150bp paired-

end) at a depth of 30x by Novogene (Hong Kong). The resulting sample group sizes for

gene expression analysis were: 3-4 libraries for each group (pooled from the same families

used for RNA-sequencing): $FC_{POP}FC_{ENV}$, $FC_{POP}NC_{ENV}$, $NC_{POP}NC_{ENV}$, $NC_{POP}FC_{ENV}$ (n=3-

4/group; total of 14 libraries). Following initial QC (using FastQC v0.11.9), reads were

trimmed using TrimGalore (0.6.4) to remove adaptor sequences and poor-quality reads.

Bisulfite conversion efficiency across all samples was estimated by Novogene and was

between 99.14-99.50% for all samples. Bisulfite-converted reads obtained from each library

were mapped to the *N. Vespilloides* genome using Bismark (v0.22.3; Krueger & Andrews,

2011) and were quantified following deduplication of reads. Mapping rates for samples to the

N. vespilloides reference genome were 65.45% ± 5.48, which after deduplication gave an

average coverage of 17.78x ± 1.79 (mean ± SD).

Differential Methylation Analysis

For differential methylation analysis of individual CpGs we first tested each site in every

sample to determine sites that were significantly methylated using a binomial test. We then

tested for differential methylation across sites using a weighted binomial glm (*p*-values were

adjusted using Benjamini-Hochperg corrections for multiple testing). We then mapped CpGs

to their feature using our custom genome annotation (see below for details) and bedtools

(Quinlan & Hall, 2010). For analyses of gene body methylation, we collapsed across all CpG

sites within each genic region (using bedtools) to compute the average weighted methylation

value for each gene for each sample. Genes were clustered into methylated and

unmethylated clusters using mixture models using the mixtools R package. To determine

which genes were differentially methylated we used a weighted binomial GLM (base R stats

package) and adjusted for multiple testing using the Benjamini-Hochberg procedure. Genes

were considered differentially methylated if they had an adjusted $p<0.01$. Fishers' exact tests

for enrichment were run using the base R stats packages.

Genome Annotation

To annotate exons in each genome we used existing annotations, excluding genes that were

split across multiple contigs. To annotate regions which may contain promoters or

enhancers (5' UTR), we took 1,000 bases upstream of each gene, excluding genes where

this exceeded the contig start or end point. We annotated introns based on the position of

exons, excluding genes that were split across multiple contigs. To annotate TEs, we used

RepeatModeller (v2.0.1) to generate a model of TEs for the *N. vespilloides* genome, and

then RepeatMasker (v4.1.0) to annotate TEs based on the model for that genome.

Data Access

All raw sequencing data generated have been submitted to the NCBI Gene Expression

Omnibus (GEO) under accession number (TBC).

References

Arsenault, S. V., Hunt, B. G., & Rehan, S. M. (2018). The effect of maternal care on gene expression and DNA methylation in a subsocial bee. *Nature Communications*, *9*(1), 3468. https://doi.org/10.1038/s41467-018-05903-0

Badyaev, A. V., & Uller, T. (2009). Parental effects in ecology and evolution: Mechanisms, processes and implications. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *364*(1520), 1169–1177. https://doi.org/10.1098/rstb.2008.0302

Becker, C., Hagmann, J., Müller, J., Koenig, D., Stegle, O., Borgwardt, K., & Weigel, D. (2011). Spontaneous epigenetic variation in the Arabidopsis thaliana methylome. *Nature*, *480*(7376), 245–249. https://doi.org/10.1038/nature10555

Bentzur, A., Shmueli, A., Omesi, L., Ryvkin, J., Knapp, J.-M., Parnas, M., Davis, F. P., & Shohat-Ophir, G. (2018). Odorant binding protein 69a connects social interaction to modulation of social responsiveness in Drosophila. *PLOS Genetics*, *14*(4), e1007328. https://doi.org/10.1371/journal.pgen.1007328

Bewick, A. J., Vogel, K. J., Moore, A. J., & Schmitz, R. J. (2016). Evolution of DNA Methylation across Insects. *Molecular Biology and Evolution*, msw264. https://doi.org/10.1093/molbev/msw264

Bonduriansky, R., & Day, T. (2018). *Extended heredity: A new understanding of inheritance and evolution*. Princeton University Press.

Charlesworth, D., Barton, N. H., & Charlesworth, B. (2017). The sources of adaptive variation. *Proceedings of the Royal Society B: Biological Sciences*, *284*(1855), 20162864. https://doi.org/10.1098/rspb.2016.2864

Cunningham, C. B., Ji, L., McKinney, E. C., Benowitz, K. M., Schmitz, R. J., & Moore, A. J. (2019). Changes of gene expression but not cytosine methylation are associated with male parental care reflecting behavioural state, social context and individual flexibility. *The Journal of Experimental Biology*, *222*(2), jeb188649. https://doi.org/10.1242/jeb.188649

Cunningham, C. B., Ji, L., Wiberg, R. A. W., Shelton, J., McKinney, E. C., Parker, D. J., Meagher, R. B., Benowitz, K. M., Roy-Zokan, E. M., Ritchie, M. G., Brown, S. J., Schmitz, R. J., & Moore, A. J. (2015). The Genome and Methylome of a Beetle with Complex Social Behavior, *Nicrophorus vespilloides* (Coleoptera: Silphidae). *Genome Biology and Evolution*, *7*(12), 3383–3396. https://doi.org/10.1093/gbe/evv194

Curley, J. P., Mashoodh, R., & Champagne, F. A. (2011). Epigenetics and the origins of paternal effects. *Hormones and Behavior*, *59*(3), 306–314. https://doi.org/10.1016/j.yhbeh.2010.06.018

Dixon, G., Liao, Y., Bay, L. K., & Matz, M. V. (2018). Role of gene body methylation in acclimatization and adaptation in a basal metazoan. *Proceedings of the National Academy of Sciences*, *115*(52), 13342–13346. https://doi.org/10.1073/pnas.1813749115

Dubin, M. J., Zhang, P., Meng, D., Remigereau, M.-S., Osborne, E. J., Paolo Casale, F., Drewe, P., Kahles, A., Jean, G., Vilhjálmsson, B., Jagoda, J., Irez, S., Voronin, V., Song, Q., Long, Q., Rätsch, G., Stegle, O., Clark, R. M., & Nordborg, M. (2015). DNA methylation in Arabidopsis has a genetic basis and shows evidence of local adaptation. *ELife*, *4*, e05255. https://doi.org/10.7554/eLife.05255

Eggert, A.-K., Reinking, M., & Müller, J. K. (1998). Parental care improves offspring survival and growth in burying beetles. *Animal Behaviour*, *55*(1), 97–107. https://doi.org/10.1006/anbe.1997.0588

Farhadian, S. F., Suárez-Fariñas, M., Cho, C. E., Pellegrino, M., & Vosshall, L. B. (2012). Post-fasting olfactory, transcriptional, and feeding responses in Drosophila. *Physiology & Behavior*, *105*(2), 544–553. https://doi.org/10.1016/j.physbeh.2011.09.007

Feinberg, A. P., & Irizarry, R. A. (2010). Stochastic epigenetic variation as a driving force of development, evolutionary adaptation, and disease. *Proceedings of the National*

| 578 | *Academy of Sciences*, *107*(suppl_1), 1757–1764. |
| 579 | https://doi.org/10.1073/pnas.0906183107 |

Feng, S., Cokus, S. J., Zhang, X., Chen, P.-Y., Bostick, M., Goll, M. G., Hetzel, J., Jain, J.,
Strauss, S. H., Halpern, M. E., Ukomadu, C., Sadler, K. C., Pradhan, S., Pellegrini,
M., & Jacobsen, S. E. (2010). Conservation and divergence of methylation patterning
in plants and animals. *Proceedings of the National Academy of Sciences*, *107*(19),
8689–8694. https://doi.org/10.1073/pnas.1002720107

Franklin, T. B., Russig, H., Weiss, I. C., Gräff, J., Linder, N., Michalon, A., Vizi, S., &
Mansuy, I. M. (2010). Epigenetic Transmission of the Impact of Early Stress Across
Generations. *Biological Psychiatry*, *68*(5), 408–415.
https://doi.org/10.1016/j.biopsych.2010.05.036

Gustafsson, L., & Sutherland, W. J. (1988). The costs of reproduction in the collared
flycatcher Ficedula albicollis. *Nature*, *335*(6193), 813–815.
https://doi.org/10.1038/335813a0

Hadfield, J. D. (2010). MCMC Methods for Multi-Response Generalized Linear Mixed
Models: The **MCMCglmm** *R* Package. *Journal of Statistical Software*, *33*(2).
https://doi.org/10.18637/jss.v033.i02

Heckwolf, M. J., Meyer, B. S., Häsler, R., Höppner, M. P., Eizaguirre, C., & Reusch, T. B. H.
(2020). Two different epigenetic information channels in wild three-spined
sticklebacks are involved in salinity adaptation. *Science Advances*, *6*(12), eaaz1138.
https://doi.org/10.1126/sciadv.aaz1138

Holliday, R. (2006). Epigenetics: A historical overview. *Epigenetics*, *1*(2), 76–80.
https://doi.org/10.4161/epi.1.2.2762

Hu, J., Wuitchik, S. J. S., Barry, T. N., Jamniczky, H. A., Rogers, S. M., & Barrett, R. D. H.
(2021). Heritability of DNA methylation in threespine stickleback (*Gasterosteus
aculeatus*). *Genetics*, *217*(1), 1–15. https://doi.org/10.1093/genetics/iyab001

Huerta-Cepas, J., Forslund, K., Coelho, L. P., Szklarczyk, D., Jensen, L. J., von Mering, C.,
& Bork, P. (2017). Fast Genome-Wide Functional Annotation through Orthology
Assignment by eggNOG-Mapper. *Molecular Biology and Evolution*, *34*(8), 2115–
2122. https://doi.org/10.1093/molbev/msx148

Jaenisch, R., & Bird, A. (2003). Epigenetic regulation of gene expression: How the genome
integrates intrinsic and environmental signals. *Nature Genetics*, *33*(S3), 245–254.
https://doi.org/10.1038/ng1089

Jarrett, B. J. M., Evans, E., Haynes, H. B., Leaf, M. R., Rebar, D., Duarte, A., Schrader, M.,
& Kilner, R. M. (2018). A sustained change in the supply of parental care causes
adaptive evolution of offspring morphology. *Nature Communications*, *9*(1), 3987.
https://doi.org/10.1038/s41467-018-06513-6

Jarrett, B. J. M., Rebar, D., Haynes, H. B., Leaf, M. R., Halliwell, C., Kemp, R., & Kilner, R.
M. (2018). Adaptive evolution of synchronous egg-hatching in compensation for the
loss of parental care. *Proceedings of the Royal Society B: Biological Sciences*,
285(1885), 20181452. https://doi.org/10.1098/rspb.2018.1452

Jombart, T. (2008). adegenet: A R package for the multivariate analysis of genetic markers.
Bioinformatics, *24*(11), 1403–1405. https://doi.org/10.1093/bioinformatics/btn129

Kawakatsu, T., Huang, S. C., Jupe, F., Sasaki, E., Schmitz, R. J., Urich, M. A., Castanon,
R., Nery, J. R., Barragan, C., He, Y., Chen, H., Dubin, M., Lee, C.-R., Wang, C.,
Bemm, F., Becker, C., O'Neil, R., O'Malley, R. C., Quarless, D. X., … Zhou, X.
(2016). Epigenomic Diversity in a Global Collection of Arabidopsis thaliana
Accessions. *Cell*, *166*(2), 492–505. https://doi.org/10.1016/j.cell.2016.06.044

Kilner, R. M., Boncoraglio, G., Henshaw, J. M., Jarrett, B. J. M., De Gasperin, O., Attisano,
A., & Kokko, H. (2015). *Parental effects alter the adaptive value of an adult
behavioural trait*. *4*, e07340. https://doi.org/10.7554/eLife.07340.001

Kriventseva, E. V., Kuznetsov, D., Tegenfeldt, F., Manni, M., Dias, R., Simão, F. A., &
Zdobnov, E. M. (2019). OrthoDB v10: Sampling the diversity of animal, plant, fungal,
protist, bacterial and viral genomes for evolutionary and functional annotations of

orthologs. *Nucleic Acids Research*, *47*(D1), D807–D811.
https://doi.org/10.1093/nar/gky1053

Krueger, F., & Andrews, S. R. (2011). Bismark: A flexible aligner and methylation caller for
Bisulfite-Seq applications. *Bioinformatics*, *27*(11), 1571–1572.
https://doi.org/10.1093/bioinformatics/btr167

Laland, K., Uller, T., Feldman, M., Sterelny, K., Müller, G. B., Moczek, A., Jablonka, E.,
Odling-Smee, J., Wray, G. A., Hoekstra, H. E., Futuyma, D. J., Lenski, R. E.,
Mackay, T. F. C., Schluter, D., & Strassmann, J. E. (2014). Does evolutionary theory
need a rethink? *Nature*, *514*(7521), 161–164. https://doi.org/10.1038/514161a

Law, J. A., & Jacobsen, S. E. (2010). Establishing, maintaining and modifying DNA
methylation patterns in plants and animals. *Nature Reviews Genetics*, *11*(3), 204–
220. https://doi.org/10.1038/nrg2719

Lewis, S. H., Ross, L., Bain, S. A., Pahita, E., Smith, S. A., Cordaux, R., Miska, E. A.,
Lenhard, B., Jiggins, F. M., & Sarkies, P. (2020). ------Widespread conservation and
lineage-specific diversification of genome-wide DNA methylation patterns across
arthropods. *PLOS Genetics*, *16*(6), e1008864.
https://doi.org/10.1371/journal.pgen.1008864

Liew, Y. J., Howells, E. J., Wang, X., Michell, C. T., Burt, J. A., Idaghdour, Y., & Aranda, M.
(2020). Intergenerational epigenetic inheritance in reef-building corals. *Nature
Climate Change*, *10*(3), 254–259. https://doi.org/10.1038/s41558-019-0687-2

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550.
https://doi.org/10.1186/s13059-014-0550-8

Massart, R., Suderman, M., Provencal, N., Yi, C., Bennett, A. J., Suomi, S., & Szyf, M.
(2014). Hydroxymethylation and DNA methylation profiles in the prefrontal cortex of
the non-human primate rhesus macaque and the impact of maternal deprivation on
hydroxymethylation. *Neuroscience*, *268*, 139–148.
https://doi.org/10.1016/j.neuroscience.2014.03.021

Miska, E. A., & Ferguson-Smith, A. C. (2016). Transgenerational inheritance: Models and
mechanisms of non-DNA sequence-based inheritance. *Science*, *354*(6308), 59–63.
https://doi.org/10.1126/science.aaf4945

Nyman, C., Hebert, F. O., Bessert-Nettelbeck, M., Aubin-Horth, N., & Taborsky, B. (2020).
Transcriptomic signatures of social experience during early development in a highly
social cichlid fish. *Molecular Ecology*, *29*(3), 610–623.
https://doi.org/10.1111/mec.15335

Pertea, M., Kim, D., Pertea, G. M., Leek, J. T., & Salzberg, S. L. (2016). Transcript-level
expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown.
Nature Protocols, *11*(9), 1650–1667. https://doi.org/10.1038/nprot.2016.095

Quinlan, A. R., & Hall, I. M. (2010). BEDTools: A flexible suite of utilities for comparing
genomic features. *Bioinformatics*, *26*(6), 841–842.
https://doi.org/10.1093/bioinformatics/btq033

Rebar, D., Bailey, N. W., Jarrett, B. J. M., & Kilner, R. M. (2020). An evolutionary switch from
sibling rivalry to sibling cooperation, caused by a sustained loss of parental care.
Proceedings of the National Academy of Sciences, *117*(5), 2544–2550.
https://doi.org/10.1073/pnas.1911677117

Sarkies, P. (2020). Molecular mechanisms of epigenetic inheritance: Possible evolutionary
implications. *Seminars in Cell & Developmental Biology*, *97*, 106–115.
https://doi.org/10.1016/j.semcdb.2019.06.005

Schmitz, R. J., Lewis, Z. A., & Goll, M. G. (2019). DNA Methylation: Shared and Divergent
Features across Eukaryotes. *Trends in Genetics*, *35*(11), 818–827.
https://doi.org/10.1016/j.tig.2019.07.007

Schrader, M., Jarrett, B. J. M., & Kilner, R. M. (2015). Using Experimental Evolution to Study
Adaptations for Life within the Family. *The American Naturalist*, *185*(5), 610–619.
https://doi.org/10.1086/680500

Schrader, M., Jarrett, B. J. M., Rebar, D., & Kilner, R. M. (2017). Adaptation to a novel family
environment involves both apparent and cryptic phenotypic changes. *Proceedings of*
the Royal Society B: Biological Sciences, *284*(1862), 20171295.
https://doi.org/10.1098/rspb.2017.1295
Skinner, M. K., Gurerrero-Bosagna, C., Haque, M. M., Nilsson, E. E., Koop, J. A. H., Knutie,
S. A., & Clayton, D. H. (2014). Epigenetics and the Evolution of Darwin's Finches.
Genome Biology and Evolution, *6*(8), 1972–1989. https://doi.org/10.1093/gbe/evu158
Steiger, S. (2013). Bigger mothers are better mothers: Disentangling size-related prenatal
and postnatal maternal effects. *Proceedings of the Royal Society B: Biological*
Sciences, *280*(1766), 20131225. https://doi.org/10.1098/rspb.2013.1225
Stephens, M. (2016). False discovery rates: A new deal. *Biostatistics*, kxw041.
https://doi.org/10.1093/biostatistics/kxw041
Suderman, M., McGowan, P. O., Sasaki, A., Huang, T. C. T., Hallett, M. T., Meaney, M. J.,
Turecki, G., & Szyf, M. (2012). Conserved epigenetic sensitivity to early life
experience in the rat and human hippocampus. *Proceedings of the National*
Academy of Sciences, *109*(Supplement_2), 17266–17272.
https://doi.org/10.1073/pnas.1121260109
Sun, S., Catherall, A. M., Pascoal, S., Jarrett, B. J. M., Miller, S. E., Sheehan, M. J., & Kilner,
R. M. (2020). Rapid local adaptation linked with phenotypic plasticity. *Evolution*
Letters, *4*(4), 345–359. https://doi.org/10.1002/evl3.176
Wang, L., Dankert, H., Perona, P., & Anderson, D. J. (2008). A common genetic target for
environmental and heritable influences on aggressiveness in Drosophila.
Proceedings of the National Academy of Sciences, *105*(15), 5657–5663.
https://doi.org/10.1073/pnas.0801327105
Weaver, I. C. G., Meaney, M. J., & Szyf, M. (2006). Maternal care effects on the
hippocampal transcriptome and anxiety-mediated behaviors in the offspring that are
reversible in adulthood. *Proceedings of the National Academy of Sciences*, *103*(9),
3480–3485. https://doi.org/10.1073/pnas.0507526103
Yagound, B., Remnant, E. J., Buchmann, G., & Oldroyd, B. P. (2020). Intergenerational
transfer of DNA methylation marks in the honey bee. *Proceedings of the National*
Academy of Sciences, *117*(51), 32519–32527.
https://doi.org/10.1073/pnas.2017094117
718

719 **Figure 1**

720
Figure 1. a) Summary of sampling from experimentally evolving populations and sequencing strategy.
Larval heads from Full Care (FC_{POP}) and No Care (NC_{POP}) populations in native ($FC_{POP}FC_{ENV}$;
$NC_{POP}NC_{ENV}$) and reciprocal environments ($FC_{POP}NC_{ENV}$; $NC_{POP}FC_{ENV}$) were sampled and used to
measure gene expression (RNA-Seq) and DNA methylation (whole-genome bisulfite sequencing;
WGBS). **b)** Density plots of discriminant function values for each family indicating that variation in
gene expression is primarily due to the current social environment. **c)** Number of differentially
expressed genes (DEGs) relative to the $FC_{POP}FC_{ENV}$ group. **d)** Venn diagram indicating degree of
overlap between plastic and evolved DEGs. **e)** Scatterplot showing the correlation between log-fold
changes ($log_2\Delta$) in the response to a NC_{ENV} within each population. **f)** Gene ontology (GO; biological
processes) enrichment for plastic response to a NC_{ENV}. **g)** GO terms for "new" DEGs following
adaptation. Only GO terms with FDR-corrected $p<0.05$ are shown. Numbers indicate number of
genes associated with that category.

Figure 2

Figure 2. a) Ridge plots showing the distribution of percent methylated CpG levels (mCpG/CpG %) in different genomic features: 1000bp upstream of genes (5' UTR), repeat or transposable elements (TE), introns, coding sequences (CDS) and across the entire gene body (gene). Median mCpG values are indicated by vertical dotted lines. **b)** Average mCpG levels across gene bodies (± 1kb) of genes categories by transcriptional activity: from inactive (0) to genes showing low (1), intermediate (2) and high (3) levels of expression. Plotted regions include 1kb upstream and downstream of transcription start sites (TSS) and transcription termination sites (TTS), respectively for each gene **c)** Bar plot indicating the distribution of methylated CpGs across gene features. **d)** Enrichment scores (odds-ratio; $\log_2(OR)$) for differentially methylated CpGs across different gene features. *Fishers' FDR-corrected p<0.05*.

747 **Figure 3**

748
Figure 3. a) Histogram of number of differentially methylated genes (DMGs) passing FDR-corrected
$p<0.01$ when samples are shuffled randomly. Vertical lines indicate number of DMGs found between
plastic (red) and evolved contrasts (blue). **b)** Venn diagram indicating degree of overlap between
plasticity and evolved DMGs. **c)** The relationship between gene expression (FKPM) and percent CpG
methylation (mCpG/CpG) for all genes analysed for plasticity (top) and evolved (bottom) contrasts.
Colours indicated differentially expressed genes (DEGs; blue) and DMGs (purple) for each contrast.
Gene ontology (GO; biological processes) enrichment for plastic response DMGs **(e)** Evolved DMGs
(f) and differentially methylated but not differentially expressed in the Evolved contrast **(g)**. Only GO
terms with FDR-corrected $p<0.05$ are shown. Numbers indicate number of genes associated with that
category.