Drying of virus-containing aerosol particles: Modelling effects of droplet origin and composition.

Michael C. Jarvis

School of Chemistry, Glasgow University, Glasgow G12 8QQ, Scotland, UK.

Email: michael.jarvis@glasgow.ac.uk

____

Abstract

Virus-containing aerosol droplets emitted by breathing, speech or coughing dry rapidly to

equilibrium with ambient relative humidity (RH), increasing in solute concentration with effects on

virus survival and decreasing in diameter with effects on sedimentation and respiratory uptake. In

simulations starting from drying data on mixed NaCl/KCl aerosols, the supersaturated salt

concentrations were shown to reach 20-25M at the efflorescence RH of 40-55%, depending on the

K:Na ratio. These salt concentrations may inactivate some viruses. The dependence on K:Na ratio

implies that the evaporation curves differ between aerosols derived from saliva and from airway

surfaces. Differences in drying behaviour are consequently predicted between breathing, speech and

coughing emissions and between droplet size fractions within these. The direct effect of liquid

droplet size through the Kelvin term was shown to be smaller and restricted to breath emissions.

Comparative simulations starting from osmotic pressure measurements on airway surface liquid

showed that salts are the primary determinants of drying equilibria down to the efflorescence RH,

and macromolecules at lower RH.

Keywords: evaporation; SARS-CoV-2; HRV-16; water activity; virus inactivation; Kohler theory;

efflorescence

____

Introduction

Some respiratory viruses can be transmitted in aerosol form, as well as in larger droplets and surface deposits[1]. Aerosols are conventionally defined as droplets or particles less than 5-10 μm in diameter that, according to Stokes' Law, remain suspended in still air for minutes or longer[2]. It has been argued[3,4] that the size range should be extended to 50-100 μm because turbulence, either in a cough jet[5] or due to draughts[6] or convection[7], keeps larger particles airborne for longer than is predicted by Stokes' Law.

Despite initial doubts, it is now quite widely accepted that certain viruses including SARS-CoV-2 are transmitted in aerosols, particularly from asymptomatic subjects[8,9]. Aerosol transmission is most likely in enclosed spaces such as schools, public buildings and transportation[10,11]. Aerosols can transmit viruses from person to person with transiently inadequate social distancing[4], but they can also build up, over minutes to hours, throughout the air in an enclosed space so that the risk of infection depends on the duration of emission and exposure, not on distancing[12]. In these circumstances the risk depends on Wells-Riley dynamics[13] and is reduced by ventilation with fresh or filtered air and by anything that decreases the viable lifetime of the virus[11,12].

The viable lifetime of airborne viruses varies. SARS-CoV-2 remains viable indoors for minutes to hours[14-17]. The rate of inactivation is enhanced by sunlight[18,19] and increases with temperature[19,20]. There is also a significant effect of humidity[14,19], but the nature and mechanism of the humidity effect are still poorly understood.

The effect of humidity on virus inactivation also differs between viruses. For some viruses inactivation is faster at intermediate levels of relative humidity, around 40%, than at high or very low humidity[21]. It has been suggested that the common cold virus HRV-16, which follows this pattern, loses activity in the supersaturated salt solution when dried from the high RH of emission to just above the precipitation (efflorescence) zone of the salts present, roughly 40% RH[22]. In contrast, for SARS-CoV-2 in aerosol form, the limited data available suggest that the rate of inactivation is low at all RH values tested up to about 50% and increases at higher RH[19]. The rate of inactivation of surface deposits of SARS[23] and SARS-CoV-2[24,25] is lower than in aerosols but likewise rises with increasing humidity[26]. This implies that the SARS viruses are more resistant than, for example, HRV-16 to very high salt concentrations, and that whatever its mechanism the inactivation of SARS-CoV-2 is dependent on available water.

Free water is present in aerosol droplets and surface deposits above the efflorescence RH of the solutes present. Water availability, expressed as the water activity a_w, increases with the ambient RH but is also modulated by the Kelvin effect in small aerosol droplets.

These observations suggest that that the drying process for aerosol-sized droplets deserves closer examination. The same is true for surface deposits, particularly as it is not known why viral viability is enhanced in that form[16,25]. The drying of aerosols when emitted into ambient air has been quite extensively studied[27-29] and modelled[27,28,30,31], often with the aim of predicting when droplets, initially large enough to sediment in still air, will shrink enough to remain suspended. In the biomedical literature, a droplet that has dried to equilibrium with the ambient air is called a droplet nucleus[1]. Depending on the solids present and the moisture that their hygroscopicity retains, a droplet nucleus may consist of a very concentrated solution, a polycrystalline salt precipitate, a protein gel or amorphous solid, or a combination of these phases; presenting very different environments in which viruses may be inactivated[21]. For example, thermal denaturation depends on mobile water[32]. Also polycrystalline or other solids may refract or absorb daylight, which is known to inactivate SARS-CoV-2[19].

Much of the published experimentation on droplet drying (cited by[27]) has made use of simplified analogues of the respiratory fluids in which viruses are emitted by breathing, speaking, singing, coughing or sneezing. Sometimes just NaCl solutions have been used. The detailed salt composition of natural aerosols has a profound effect on their drying behaviour[33]. Metzger et al.[34] give an accessible description of the underlying physics (Kohler theory) as an appendix: their nomenclature is adopted here. In respiratory droplets, proteins and glycoproteins have been recognised to contribute volume to the dried droplet nuclei, but little attention has been paid to

other ways in which these polymers might influence the drying process[27,35].

Because of the very high salt and polymer concentrations that can be reached when biological aerosols dry at low RH, classical colligative relationships like Raoult's Law become increasingly unsatisfactory approximations and the relevant physical chemistry becomes necessarily more empirical. In these circumstances direct experimental measurements using real biological fluids may be more informative than theoretical prediction[27]. However, experimentation on aerosols is technically challenging[27,36,37] and during the pandemic time is short.

This paper describes simulations of the effects of some of the main variables in the composition of virus-containing aerosols on drying equilibria. In view of the above comments, these simulations do not aim at quantitative descriptions of complex, real-life bioaerosols, but may serve to suggest some simplifying assumptions and to prioritise variables that deserve experimental investigation.

Results

Origins and composition of emitted droplets

Aerosols and larger droplets emitted during breathing[38,39], speech[37,40], coughing[39] and other activities[41] originate by aerodynamic disruption of the mucosal lining[42] in different zones in the respiratory tract[36,43,44], leading to different droplet size distributions[45,46]. The principal zones in which droplets are generated are the bronchioli (modal droplet diameter 1-2 μm) the laryngeal region (modal droplet diameter 1-2 μm), and the oral cavity and nasal region (modal droplet diameter >100 μm)[46]. Multiple sites of origin lead to bimodal, trimodal or broad continuous ranges of droplet diameter for each mode of emission[46]. With droplet diameters covering several orders of magnitude, caution is needed in the interpretation of modal figures because they may be derived by several experimental methods with differing size limitations[42,46] and because number-weighted and volume-weighted distributions are very different: volume-weighted distributions are more relevant to viral load[35].

Droplets emitted in normal breathing originate primarily from the bronchial zone and have diameters in the submicron to μm range[46,47], while droplets emitted in talking or coughing are derived partly from the laryngeal and oral zones, with a preponderance of larger particles[36,39,46]. The viral load of the mucosa in each region[48,49] varies with disease progression[50] and between individuals[39,40].

The drying of emitted droplets depends on their ionic and polymer composition. It has not been well recognised that the composition of the droplets differs according to their site of origin[27], and therefore also differs with droplet diameter.

A key function of the airway lining throughout the respiratory tract is to sustain hydration[51] and freedom of motion for the underlying cilia[52], defects in hydration leading to disorders such as cystic fibrosis[53]. Equilibrium hydration depends on osmolytes in a very similar way to water retention by emitted droplets[51], although the RH within the respiratory system is much higher. The cation composition of the airway surface liquid is dominated by sodium, with $Na^+:K^+$ molar ratios typically around 4:1[53-55]: the precision is lower for K^+ due to the difficulty of sampling without cellular damage[56]. The principal anion is Cl^-, with a much smaller amount of HCO_3^-[53,54]. In health the total osmolarity is approximately 300 mM[54], increasing from the lower respiratory tract to the nasal region[53] and increasing substantially in conditions such as chronic bronchitis[51]. Fluid harvested from human bronchial epithelial (HBE) cell cultures is rather similar in ionic composition to native airway fluids but with lower protein content[55].

In contrast, saliva has much lower ion concentrations and osmolarity[57]. In the normal (resting) state, the total osmolarity averages 50 mM and the main cation is K^+ with a $K^+:Na^+$ ratio of about 3:1[58]. The main anion is Cl^-. On stimulation, water secretion is driven by an increase in Na^+ and Cl^- concentrations. The $Na^+:K^+$ ratio is therefore variable and can exceed unity[58]. The large difference in solute concentrations between saliva and airway surface fluids means that sputum varies in composition between these extremes[55]. Similarly, emitted droplets are predicted to have an overall $K^+:Na^+$ ratio that depends on the saliva contribution and is highest for speech[46], whereas the aerosol droplets emitted by breathing originate mainly from the lining of the lower respiratory tract and are dominated by Na^+. In emissions of mixed origin, large droplets[46] are likely to be dominated

by K$^+$ and small droplets[46] by Na$^+$, leading to differences in their evaporation equilibria.

Simulated effects of Na$^+$:K$^+$ ratio

KCl is less hygroscopic than NaCl[59]. Therefore, neglecting the direct (Kelvin) effect of droplet size, large droplets with KCl as the predominant salt would be predicted to reach their efflorescence point at higher RH than small droplets with NaCl as the predominant salt.

Figure 1. Efflorescence relative humidity of KCl : NaCl mixtures in aerosol form, calculated from data in[59].

However, the behaviour of salt mixtures is not necessarily additive. Li et al.[59] measured droplet sizes of mixed KCl : NaCl aerosols as they became larger with increasing RH and smaller with decreasing RH, using a microscopy technique after impaction. Figure 1, calculated from their experimental data[59], shows that in aerosol mixtures of KCl and NaCl, with no other solutes present, the relationship between efflorescence RH and K:Na ratio is non-linear, with a minimum efflorescence RH at about K mol fraction 0.4. Thus each salt tends to keep the other in supersaturated solution until both precipitate together at the efflorescence RH.

Figure 2 shows that In the absence of other solutes, mixtures of NaCl and KCl can reach supersaturated concentrations up to 20 M or more. Such high concentrations have been suggested to reduce survival of susceptible viruses[22]. Somewhat higher maximal salt concentrations are reached when the major cation is potassium compared to sodium, with the maximum at higher RH. When the RH fell below the efflorescence point each droplet contracted abruptly to an irregular solid that

remained constant in size down to RH = 5%[59], from which it was assumed that the liquid phase disappeared at the efflorescence RH. The mixture with K mol fraction 0.2 is representative of the principal ion content of airway lining fluid[53,54] emitted mainly as small (<10 μm) droplets[46]. The mixture with K mol fraction 0.8 is representative of the principal ion content of saliva[58] emitted mainly as larger (>10 μm) droplets[46]. Ions other than Na$^+$ and K$^+$ are also present, of course, and their contribution to drying equilibria could be calculated[33] if comprehensive consensus values for their concentrations were available.

Figure 2. Simulated supersaturated total salt concentration in aerosols of KCl / NaCl mixtures as a function of decreasing relative humidity, calculated from[59]. Salt concentrations increased as the droplets dried until they reached the efflorescence RH (arrowed) at which point the droplets dried abruptly.

Droplet diameter: the Kelvin effect

The extent to which droplets dry at any RH is also influenced directly by their size due to the Kelvin effect. In Kohler theory, the increased surface curvature of small droplets leads to enhanced evaporation at a given RH, according to the diameter-dependent Kelvin term K_e in the expression for their drying equilibrium[34].

Droplet diameter, μm

○ 0.01   ○ 0.02   ○ 0.05   ○ 0.1   ○ 0.2   ○ 0.5

Figure 3. Simulated effect of droplet diameter, through inclusion of the Kelvin term, on the equilibrium salt concentration during drying of aerosol droplets with K mol fraction = 0.2 as in Figure 2.

Figure 3 shows that the effect of including the Kelvin term in the simulation is to displace the whole curve to higher RH. The Kelvin effect starts to become significant only at droplet diameters below about 0.1 μm. In calculating the Kelvin term it is generally assumed that the droplet is wholly liquid and is spherical[34]. When a mixture of irregular solid and liquid phases is present, the local radius at protuberances may be less than calculated and the effect augmented.

The magnitude of the Kelvin effect depends also on σ_{sol}, the surface tension, which in the simulation above was assumed constant and equivalent to that of water. However, surfactants reduce σ_{sol},[60] and are present in airway surface liquids[61], influencing their fragmentation into aerosols[42]. Surfactant proteins[61] are best known from the lungs but are detectable elsewhere[62] and are accompanied by deacylated phosphatidyl choline[63]. The effect of surfactants on biogenic droplet drying has not been quantified but it can be assumed that they reduce the magnitude of the Kelvin term, to an uncertain and possibly large extent[60].

Macromolecular composition: proteins and mucin glycoproteins

Figure 2 simulates the drying of NaCl/KCl mixtures with no other solutes present. However, virus-containing aerosols contain larger amounts of proteins and glycoproteins[55,58], which increase in concentration as the droplets dry until they constitute most of the mass of the droplet nucleus. It is often assumed that these macromolecules have no influence on water activity[28], an assumption that might not be valid when there is more protein than remaining water.

In the intact airway lining, mucin glycoproteins have been stated to play a role in maintaining hydration[51] and have been studied with that function in mind[55]. Reduced water content and increased viscosity are well-known factors in cystic fibrosis and other pathological conditions[51,55]. A substantial rheological change attributed to gelation has been observed when the solids content and mucin content of normal airway fluid are doubled[55], which would occur at RH > 90% during droplet drying. It would then follow that during much of the supersaturation part of the aerosol drying curve the mucin fraction, and perhaps other proteins, are in the gel state. Osmotic relations of polymer gels are difficult to handle, although for simplification it is often assumed that only low-molecular species – free salts and the counterions associated with any charges on the polymer – contribute to lowering water activity[64]. This assumption may not hold at high polymer concentrations or for very flexible polymers that undergo vigorous segmental motion.

In the case of airway fluid, these conceptual problems have been circumvented by direct measurement of the polymer-associated osmotic pressure using a membrane permeable to salts that are not associated with the polymer[51,52]. An empirical relationship of osmotic pressure to polymeric solids content (proteins plus glycoproteins) of airway mucus, above and below the sol-gel transition, was derived from the measurements of osmotic pressure in ref[52].

$$\Pi = kc[1+(c/c^*)^{(\beta-1)m}]^{1/m}$$

Where c is the mucus polymeric solids concentration (Kg/L), $c^* = 0.081$ Kg/L, $m = 3$, $\beta = 2.21$ and $k = 14.4$ KPa L/Kg.

Water activity a_w, and hence equilibrium relative humidity, can be connected to osmotic pressure using an appropriate form of the Van t'Hoff relationship[65]:

Osmotic pressure $\Pi = -RT\ln(a_w)/V_w$

Thus $a_w = \exp(-\Pi V_w /RT)$

Where RH is expressed as a fraction, R is the gas constant, T is absolute temperature and V_w is the molar volume of water.

Figure 4. Simulated reduction in water activity a_w, i.e. (1- a_w), during drying of airway surface liquid by the influence of salt, calculated from RH data for a NaCl/KCl mixture with initial total concentration 300 mM and K^+:Na^+ molar ratio 0.2 (see Figure 2); and by the influence of the macromolecular (protein and glycoprotein) fraction, calculated from the osmotic pressure measurements of ref[52]. Note that the (1- a_w) scale is logarithmic.

Figure 4 compares (a) the reduction in water activity due to the salts, calculated (without including the Kelvin term) from the RH data of Li et al.[59], with (b) the reduction in water activity due to proteins and glycoproteins calculated from the osmotic pressure data of Button et al[52] when solutions with ionic and protein/glycoprotein content representative of airway fluid were dried as far as equilibrium with 71% RH. Drying to this extent requires extrapolation of the osmotic pressure data somewhat beyond the measured protein concentration range[52]. Within the concentration range shown, the contribution of the protein and glycoprotein fraction to the reduction in water activity is two orders of

magnitude lower than the contribution of the salt mixture, and so would for many purposes be negligible. Extrapolation further into the concentration range where the mucin components gel would be unsafe, bearing in mind the complexity of the osmotic properties of biomacromolecule gels.

To explore what might happen at lower RH, Figure 5 shows the remaining moisture relative to the mass of protein and glycoprotein (approximated as total solids – (NaCl + KCl)) for compositions representative of airway fluid[55] (0.2 mol fraction K, 40 g/L total solids) and saliva[58] (0.8 mol fraction K, 5 g/L total solids). The extent of drying was calculated from the salt content only, neglecting the hygroscopic effect of the polymer fraction. The simulated moisture curves for solutions representing airway fluid and saliva were similar and remained above 30% until the efflorescence point was reached. In general proteins at less than 20% moisture form hydrated solids, with the water in adsorbed form retained quite strongly to low RH and no separate liquid phase[66]. This would be expected to happen below the efflorescence point.

Figure 5. Residual moisture content of the protein and glycoprotein fraction for solutions simulating airway surface liquid (Initial solids 40 g/L; Initial salt 0.15 mol/L; K mol fraction = 0.2) and saliva (Initial solids 5 g/L; Initial salt 0.027 mol/L; K mol fraction = 0.8) during drying.

Discussion

Supersaturation of salts in aerosol particles means that very high salt concentrations can potentially be reached in equilibrium with a rather narrow mid-range band of RH values (Figure 2). It should be noted than anything that can nucleate crystallisation, such as particulate solids or contact with a surface, would truncate the maximal salt concentration at higher RH. As the RH falls past the efflorescence point, NaCl / KCl mixtures precipitate abruptly (Figure 2), although a more gradual decrease in dissolved salt concentration at low RH is possible with more complex mixtures[33]. Such high salt concentrations inactivate some viruses[22], although SARS-CoV-2 appears to be relatively resistant[19]. Its survival is enhanced when droplets dry onto metal surfaces[24], for reasons that are not clear. Water activity in surface deposits depends on ambient RH in the same way as in aerosol droplets larger than the threshold of the Kelvin effect. Some fractionation of droplet components may occur during surface drying[67], and there may then be radial variation in salt and protein content within the dried deposit. There is a potential parallel in the stability of viruses adsorbed on atmospheric particulates[35].

The nature of the cations present in virus-containing emitted droplets deserves closer attention, since the ionic composition of saliva[58] differs from that of airway fluids[55]. The low salt content and high proportion of KCl in saliva droplets means that their maximal salt content is reached at higher RH than Na-rich airway fluids. Although a saliva-rich cough[46] and a sneeze[41] emit droplets with rather similar initial size ranges, their drying behaviour and droplet nucleus diameter are predicted to differ.

The dependence of drying equilibrium on droplet size through the Kelvin effect becomes noticeable when the liquid droplet diameter is less than 0.1 μm (Figure 3). Diameter distributions on emission by normal breathing include large numbers of droplets <0.1 μm[46]. However, the volume-weighted diameter distributions, which are more relevant to viral load, include only a small fraction of <0.1 μm droplets, even for breathing[35]. Also, the magnitude of the Kelvin effect is reduced to an unknown extent[60] by the presence of pulmonary and other surfactants. Thus for many purposes it is a reasonable assumption to neglect the Kelvin term

in drying simulations. This prediction has practical consequences. Variations in drying equilibria with droplet size are more likely to arise from differences in site of origin and composition, rather than from the Kelvin effect. More usefully, experiments on the hygroscopic properties of bulk biological fluids, e.g. measurements of osmotic pressure[51,52], are relevant to the behaviour of aerosols so that in suitable cases, it may be possible to obtain informative experimental data without the technical challenges inherent in aerosol generation and measurement.

However it may be inadvisable to neglect the Kelvin effect for partially dried particles or surface deposits where polycrystalline salts are present, because liquid films may have small local radii overlying crystal vertices and negative (inward) radii in interstices, leading to variation in local water activity.

In the drying range up to about 20% solids content, the hygroscopic effect of proteins and of glycoproteins such as mucins is predicted to be negligible in comparison with the effect of the salts present (Figure 4). Below the efflorescence RH the macromolecular fraction is likely to be essentially an amorphous solid holding any remaining water quite strongly (Figure 5). In the 40% - 80% solids range the mucin fraction at least is likely to form a gel, within which water relations are difficult to predict[64]. Other, salted-out proteins are likely to be interspersed with the mucin glycoproteins and protein aggregates may act as nucleation points for salt precipitation above the efflorescence RH. In this region of complex physical chemistry, experiments on real biological fluids[27] may be a better guide to the behaviour of the system than available theory.

Conclusions

The drying equilibria of aerosol and larger droplets containing infectious viruses are determined principally by the salt composition of the droplets. The salts present in saliva are K^+-dominated, whereas the more concentrated salts present in airway surface liquids are Na^+-dominated and, in the absence of other solutes, precipitate (effloresce) at lower RH. These differences mean that droplets emitted by breathing, speech, coughing and sneezing differ in drying behaviour

according to their site of origin and that in emissions with multiple sites of origin, drying behaviour differs between small and large droplets.

Methods

Modelling mixed NaCl/KCl solutions

The published efflorescence equilibria[59] in the form of measured droplet area ratios R_a with the constant droplet area at RH <30% set as unity, were converted to volume ratios $R_v = R_a^{3/2}$. This approach was justified by a close match (+/- <1%) with the efflorescence and deliquescence RH measured at bulk scale[59]. For RH below the efflorescence point $R_v = R_0$. For RH above the efflorescence point R_v is equivalent to the growth factor as conventionally defined[34]. R_0 was converted to mass using a solid density ρ_{solid} interpolated between the densities of KCl (1980 Kgm^{-3}) and NaCl (2176 Kgm^{-3}) according to the molar ratio, with a correction factor of 0.6 for void volume in the polycrystalline salt deposits. Solution concentrations were then calculated as $R_0 . \rho_{solid}(R_v - 1)$ approximating the liquid density as 1000 Kg m^{-3}.

The Kelvin term in the predicted drying equilibrium was calculated as a function of droplet diameter using the relationship

RH = $a_w . K_e = a_w . \exp(4 M_w \sigma_{sol}/(RT \rho_w D_{wet})$

Where M_w is the molar mass of water, ρ_w is its density and σ_{sol} is the surface tension of the solution[34].

Modelling water activity in macromolecular solutions

The water activity of protein/glycoprotein solutions without salt was calculated from published osmotic pressure data[52] using the following form of the Van t'Hoff relationship[65]:

Osmotic pressure $\Pi = -RT\ln(a_w)/V_w$

Rearranging, $a_w = \exp(-\Pi V_w /RT)$

Where RH is expressed as a fraction, R is the gas constant, T is absolute temperature and V_w is the molar volume of water.

All simulations were carried out in Microsoft Excel. The .xlsx files are available from the author on request.

Conflict of interest statement The author declares no conflict of interest.

Funding statement The preparation of this review was not supported by any funding body.

References

Pohlker, M. L. *et al.* Respiratory aerosols and droplets in the transmission of infectious diseases. *ArXiv*, 1-46, doi:arXiv:2103.01188 (2021).

Fenelly, K. P. Particle sizes of infectious aerosols: implications for infection control. *Lancet Respiratory Medicine*, doi:10.1016/ S2213-2600(20)30323-4 (2020).

Tang, J. W. The effect of environmental parameters on the survival of airborne infectious agents. *Journal of the Royal Society Interface* **6**, S737-S746, doi:10.1098/rsif.2009.0227.focus (2009).

Tang, J. W. *et al.* Dismantling myths on the airborne transmission of severe acute respiratory syndrome coronavirus (SARS-CoV-2). *The Journal of Hospital Infection*, doi:10.1016/j.jhin.2020.12.022 (2021).

Bourouiba, L., Dehandschoewercker, E. & Bush, J. W. M. Violent expiratory events: on coughing and sneezing. *Journal of Fluid Mechanics* **745**, 537-563, doi:10.1017/jfm.2014.88 (2014).

Choi, J. I. & Edwards, J. R. Large-eddy simulation of human-induced contaminant transport in room compartments. *Indoor Air* **22**, 77-87, doi:10.1111/j.1600-0668.2011.00741.x (2012).

Sun, S., Li, J. & Han, J. How human thermal plume influences near-human transport of respiratory droplets and airborne particles: a review. *Environmental Chemistry Letters*, doi:10.1007/s10311-020-01178-4 (2021).

Dancer, S. J. *et al.* Putting a balance on the aerosolization debate around SARS-CoV-2. *Journal of Hospital Infection* **105**, 569-570, doi:10.1016/j.jhin.2020.05.014 (2020).

Meyerowitz, E. A., Richterman, A., Gandhi, R. T. & Sax, P. E. Transmission of SARS-CoV-2: A review of viral, host, and environmental factors. *Annals of Internal Medicine* **174**, 69-79, doi:10.7326/m20-5008 (2021).

Lipinski, T., Ahmad, D., Serey, N. & Jouhara, H. Review of ventilation strategies to reduce the risk of disease transmission in high occupancy buildings. *International Journal of Thermofluids* **7**, doi:10.1016/j.ijft.2020.100045 (2020).

CIBSE: Emerging from Lockdown: Ventilation Guidance. (2020) https://www.cibse.org/ coronavirus-covid-19/emerging-from-lockdown#1

Sun, C. & Zhai, Z. The efficacy of social distance and ventilation effectiveness in preventing COVID-19 transmission. *Sustainable Cities and Society* **62**, 102390-102390, doi:10.1016/j.scs.2020.102390 (2020).

Pavilonis, B., Ierardi, A. M., Levine, L., Mirer, F. & Kelvin, E. A. Estimating aerosol transmission risk of SARS-CoV-2 in New York City public schools during reopening. *Environmental Research* **195**, 110805-110805, doi:10.1016/j.envres.2021.110805 (2021).

Smither, S. J., Eastaugh, L. S., Findlay, J. S. & Lever, M. S. Experimental aerosol survival of SARS-CoV-2 in artificial saliva and tissue culture media at medium and high humidity. *Emerging Microbes & Infections* **9**, 1415-1417, doi:10.1080/22221751.2020.1777906 (2020).

Ren, S.-Y. *et al.* Stability and infectivity of coronaviruses in inanimate environments. *World Journal of Clinical Cases* **8**, 1391-1399, doi:10.12998/wjcc.v8.i8.1391 (2020).

van Doremalen, N. *et al.* Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. *New England Journal of Medicine* **382**, 1564-1567, doi:10.1056/NEJMc2004973 (2020).

Fears, A. C. *et al.* Persistence of Severe Acute Respiratory Syndrome Coronavirus 2 in aerosol suspensions. *Emerging Infectious Diseases* **26**, doi:10.3201/eid2609.201806 (2020).

Schuit, M. *et al.* Airborne SARS-CoV-2 is rapidly inactivated by simulated sunlight. *The Journal of Infectious Diseases* **Published online ahead of print**, doi:10.1093/infdis/jiaa334 (2020).

Dabisch, P. *et al.* The influence of temperature, humidity, and simulated sunlight on the infectivity of SARS-CoV-2 in aerosols. *Aerosol Science and Technology* **55**, 142-153, doi:10.1080/02786826.2020.1829536 (2021).

Chin, A. W. H. *et al.* Stability of SARS-CoV-2 in different environmental conditions. *Lancet Microbe* **1**, e10, doi:10.1016/S2666-5247(20)30003-3 (2020).

Lin, K., Schulte, C. R. & Marr, L. C. Survival of MS2 and phi 6 viruses in droplets as a function of relative humidity, pH, and salt, protein, and surfactant concentrations. *Plos One* **15**, doi:10.1371/journal.pone.0243505 (2020).

Niazi, S. *et al.* Susceptibility of an airborne common cold virus to relative humidity. *Environmental Science and Technology* **55**, 499-508, doi:https://dx.doi.org/10.1021/acs.est.0c06197 (2021).

Chan, K. H. M. P., J.S.Lam, S.Y.Poon, L.L.M.Yuen, K.Y.Seto, W.H. The effects of temperature and relative humidity on the viability of the SARS coronavirus. *Advances in Virology*, doi:10.1155/2011/734690 (2011).

Biryukov, J. *et al.* Increasing temperature and relative humidity accelerates inactivation of SARS-CoV-2 on surfaces. *mSphere* **5**, doi:10.1128/mSphere.00441-20 (2020).

Bueckert, M., Gupta, R., Gupta, A., Garg, M. & Mazumder, A. Infectivity of SARS-CoV-2 and other coronaviruses on dry surfaces: potential for indirect transmission. *Materials* **13**, doi:10.3390/ma13225211 (2020).

Marques, M. & Domingo, J. L. Contamination of inert surfaces by SARS-CoV-2: Persistence, stability and infectivity. A review. *Environmental Research* **193**, doi:10.1016/j.envres.2020.110559 (2021).

Lieber, C., Melekidis, S., Koch, R. & Bauer, H.-J. Insights into the evaporation characteristics of saliva droplets and aerosols: Levitation experiments and numerical modeling. *Journal of Aerosol Science* **154**, 105760-105760, doi:10.1016/j.jaerosci.2021.105760 (2021).

Liu, L., Wei, J., Li, Y. & Ooi, A. Evaporation and dispersion of respiratory droplets from coughing. *Indoor Air* **27**, 179-190, doi:10.1111/ina.12297 (2017).

Chong, K. L. *et al.* Extended lifetime of respiratory droplets in a turbulent vapor puff and its implications on airborne disease transmission. *Physical Review Letters* **126**, doi:10.1103/PhysRevLett.126.034502 (2021).

Yang, X. *et al.* Transmission of pathogen-laden expiratory droplets in a coach bus. *Journal of Hazardous Materials* **397**, 122609-122609, doi:10.1016/j.jhazmat.2020.122609 (2020).

Feng, Y., Marchal, T., Sperry, T. & Yi, H. Influence of wind and relative humidity on the social distancing effectiveness to prevent COVID-19 airborne transmission: A numerical study. *Journal of Aerosol Science*, 105585-105585, doi:10.1016/j.jaerosci.2020.105585 (2020).

McDevitt, J., Rudnick, S., First, M. & Spengler, J. Role of absolute humidity in the inactivation of influenza viruses on stainless steel surfaces at elevated temperatures. *Applied and Environmental Microbiology* **76**, 3943-3947, doi:10.1128/aem.02674-09 (2010).

Metzger, S. *et al.* Aerosol water parameterisation: a single parameter framework. *Atmospheric Chemistry and Physics* **16**, 7213-7237, doi:10.5194/acp-16-7213-2016 (2016).

Metzger, S., Steil, B., Xu, L., Penner, J. E. & Lelieveld, J. New representation of water activity based on a single solute specific constant to parameterize the hygroscopic growth of aerosols in atmospheric models. *Atmospheric Chemistry and Physics* **12**, 5429-5446, doi:10.5194/acp-12-5429-2012 (2012).

Jarvis, M. C. Aerosol transmission of SARS-CoV-2: Physical principles and implications. *Frontiers in Public Health*, doi:10.3389/fpubh.2020.590041 (2020).

Morawska, L. *et al.* Size distribution and sites of origin of droplets expelled from the human respiratory tract during expiratory activities. *Journal of Aerosol Science* **40**, 256-269, doi:10.1016/j.jaerosci.2008.11.002 (2009).

Anfinrud, P., Stadnytskyi, V., Bax, C. E. & Bax, A. Visualizing speech-generated oral fluid droplets with laser light scattering. *New England Journal of Medicine* **382**, 2061-2063, doi:10.1056/NEJMc2007800 (2020).

Johnson, G. R. & Morawska, L. The mechanism of breath aerosol formation. *Journal of Aerosol Medicine and Pulmonary Drug Delivery* **22**, 229-237, doi:10.1089/jamp.2008.0720 (2009).

Fabian, P., Brain, J., Houseman, E. A., Gern, J. & Milton, D. K. Origin of exhaled breath particles from healthy and human rhinovirus-infected subjects. *Journal of Aerosol Medicine and Pulmonary Drug Delivery* **24**, 137-147, doi:10.1089/jamp.2010.0815 (2011).

Asadi, S. *et al.* Aerosol emission and superemission during human speech increase with voice loudness. *Scientific Reports* **9**, doi:10.1038/s41598-019-38808-z (2019).

Han, Z. Y., Weng, W. G. & Huang, Q. Y. Characterizations of particle size distribution of the droplets exhaled by sneeze. *Journal of the Royal Society Interface* **10**, doi:10.1098/rsif.2013.0560 (2013).

Hamed, R., Schenck, D. M. & Fiegel, J. Surface rheological properties alter aerosol formation from mucus mimetic surfaces. *Soft Matter* **16**, 7823-7834, doi:10.1039/d0sm01232g (2020).

Wei, J. J. & Li, Y. G. Airborne spread of infectious agents in the indoor environment. *American Journal of Infection Control* **44**, S102-S108, doi:10.1016/j.ajic.2016.06.003 (2016).

Mittal, R., Ni, R. & Seo, J.-H. The flow physics of COVID-19. *Journal of Fluid Mechanics* **894**, doi:10.1017/jfm.2020.330 (2020).

Gralton, J., Tovey, E., McLaws, M.-L. & Rawlinson, W. D. The role of particle size in aerosolised pathogen transmission: A review. *Journal of Infection* **62**, 1-13, doi:10.1016/j.jinf.2010.11.010 (2011).

Johnson, G. R. *et al.* Modality of human expired aerosol size distributions. *Journal of Aerosol Science* **42**, 839-851, doi:10.1016/j.jaerosci.2011.07.009 (2011).

Guo, Y. *et al.* Deposition of droplets from the trachea or bronchus in the respiratory tract during exhalation: A steady-state numerical investigation. *Aerosol Science and Technology* **54**, 869-879, doi:10.1080/02786826.2020.1772459 (2020).

Buonanno, G., Stabile, L. & Morawska, L. Estimation of airborne viral emission: Quanta emission rate of SARS-CoV-2 for infection risk assessment. *Environment International* **141**, 105794-105794, doi:10.1016/j.envint.2020.105794 (2020).

Hou, Y. X. J. *et al.* SARS-CoV-2 reverse genetics reveals a variable infection gradient in the respiratory tract. *Cell* **182**, 429-+, doi:10.1016/j.cell.2020.05.042 (2020).

Mutuku, J. K., Hou, W.-C. & Chen, W.-H. An overview of experiments and numerical simulations on airflow and aerosols deposition in human airways and the role of bioaerosol motion in COVID-19 transmission. *Aerosol and Air Quality Research* **20**, 1172-1196, doi:10.4209/aagr.2020.04.0185 (2020).

Anderson, W. H. *et al.* The relationship of mucus concentration (hydration) to mucus osmotic pressure and transport in chronic bronchitis. *American Journal of Respiratory and Critical Care Medicine* **192**, 182-190, doi:10.1164/rccm.201412-2230OC (2015).

Button, B. *et al.* A periciliary brush promotes the lung health byseparating the mucus layer from airway epithelia. *Science* **337**, 937-941, doi:10.1126/science.1223012 (2012).

Knowles, M. R. *et al.* Ion composition of airway surface liquid of patients with cystic fibrosis as compared with normal and disease-control subjects. *Journal of Clinical Investigation* **100**, 2588-2595, doi:10.1172/jci119802 (1997).

Jayaraman, S., Joo, N. S., Reitz, B., Wine, J. J. & Verkman, A. S. Submucosal gland secretions in airways from cystic fibrosis patients have normal Na$^+$ and pH but elevated viscosity. *Proceedings of the National Academy of Sciences of the United States of America* **98**, 8119-8123, doi:10.1073/pnas.131087598 (2001).

Markovetz, M. R. *et al.* Endotracheal tube mucus as a source of airway mucus for rheological study. *American Journal of Physiology-Lung Cellular and Molecular Physiology* **317**, L498-L509, doi:10.1152/ajplung.00238.2019 (2019).

Landry, J. S. & Eidelman, D. H. Airway surface liquid: End of the controversy? *Journal of General Physiology* **117**, 419-422, doi:10.1085/jgp.117.5.419 (2001).

Gal, J. Y., Fovet, Y. & Adib-Yadzi, M. About a synthetic saliva for in vitro studies. *Talanta* **53**, 1103-1115, doi:10.1016/s0039-9140(00)00618-4 (2001).

Kumar, B. *et al.* The composition, function and role of saliva in maintaining oral health: A review. *International Journal of Contemporary Dental and Medical Reviews*, 011217, doi:10.15713/ins.ijcdmr.121 (2017).

Li, X., Gupta, D., Eom, H.-J., Kim, H. & Ro, C.-U. Deliquescence and efflorescence behavior of individual NaCl and KCl mixture aerosol particles. *Atmospheric Environment* **82**, 36-43, doi:10.1016/j.atmosenv.2013.10.011 (2014).

Bzdek, B. R., Reid, J. P., Malila, J. & Prisle, N. L. The surface tension of surfactant-containing, finite volume droplets. *Proceedings of the National Academy of Sciences of the United States of America* **117**, 8335-8343, doi:10.1073/pnas.1915660117 (2020).

Larsson, P. *et al.* Surfactant protein A and albumin in particles in exhaled air. *Respiratory Medicine* **106**, 197-204, doi:10.1016/j.rmed.2011.10.008 (2012).

Schicht, M. *et al.* The distribution of human surfactant proteins within the oral cavity and their role during infectious diseases of the gingiva. *Annals of Anatomy-Anatomischer Anzeiger* **199**, 92-97, doi:10.1016/j.aanat.2014.05.040 (2015).

Larsson, P. *et al.* The effect of exhalation flow on endogenous particle emission and phospholipid composition. *Respiratory Physiology & Neurobiology* **243**, 39-46, doi:10.1016/j.resp.2017.05.003 (2017).

Komori, T. & Sakamoto, R. On Tanaka-Fillimore's kinetics swelling of gels. *Colloid and Polymer Science* **267**, 179-183, doi:10.1007/bf01410357 (1989).

Subbiah, B., Blank, U. K. M. & Morison, K. R. A review, analysis and extension of water activity data of sugars and model honey solutions. *Food Chemistry* **326**, doi:10.1016/j.foodchem.2020.126981 (2020).

Hernandez-Izquierdo, V. M. & Krochta, J. M. Thermoplastic processing of proteins for film formation - A review. *Journal of Food Science* **73**, R30-R39, doi:10.1111/j.1750-3841.2007.00636.x (2008).

Bhardwaj, R. & Agrawal, A. Likelihood of survival of coronavirus in a respiratory droplet deposited on a solid surface. *Physics of Fluids* **32**, doi:10.1063/5.0012009 (2020).