Golden-cheeked warbler habitat changes

RESEARCH ARTICLE

Spatiotemporal patterns in Golden-cheeked Warbler breeding habitat quality and quantity

Authors: Lindsay M. Dreiss*, Paul Sanchez-Navarro, Bryan Bird

Defenders of Wildlife, 1130 17th Street NW, Washington DC, USA

*Corresponding author: Lrosa@defenders.org

Acknowledgements: We thank J Malcom, T Niederman and L O'Donnell for their thoughtful
feedback in reviewing this work. We also thank the organizations that curate these publicly
available datasets and the community scientists that work to collect these data. This research was
conducted on secondary data sources. To our knowledge, no organisms were harmed in the
collection of the datasets used.

Data Availability: Analyses reported in this article can be reproduced using publicly available
data listed in Table 1. Final outputs are available on OSF at DOI 10.17605/OSF.IO/T4DJX

ABSTRACT

The Golden-cheeked Warbler, *Setophaga chrysoparia,* is a migratory songbird listed as

endangered under the federal Endangered Species Act that breeds exclusively in central Texas

and is heavily impacted by habitat conversion. The species relies on mixed Ashe-juniper and oak

woodlands for nest-building and shelter during spring and early summer months. Using land

cover data spanning the last 25 years, we conducted a geospatial analysis to quantify changes and

identify shifts in breeding habitat quantity and quality. Since 1985, 13% of all forests within the

warbler's breeding range were disturbed, with greater incidences near San Antonio (32%) and

Austin (24%) metropolitan areas. Additionally, there was a 45% decrease in high-quality habitat

(i.e., intact mixed or evergreen core forests) and a decrease in patch size. Habitats within

protected areas saw a less sharp decline in habitat quality and large increases in warbler

sightings, but these only represent 10% of all highest-quality habitat in the breeding range.

Drastic declines in habitat quality suggest that generalized metrics of conversion may

underestimate true habitat loss as degradation may impact the ecological viability of remaining

forests for warbler nesting. Further evidence suggests that the few protected areas within the

Texas range continue to play a significant role in warbler breeding. This information will assist

researchers and managers prioritizing conservation action and will inform upcoming species

status determinations.

Keywords: habitat quality, change detection, conservation, Setophaga chrysoparia

INTRODUCTION

Habitat loss and degradation through anthropogenic landscape modification are major drivers of declining global wildlife populations and serve as primary threats justifying species' listing under the United States Endangered Species Act (ESA; Hanski 2011, Bairlein 2016, Thompson et al. 2016, Horváth et al. 2019, Leu et al. 2019). Habitat disturbances are even more notable for Neotropical migrant songbirds that travel long and costly distances between breeding and non-breeding sites. In North America, the Breeding Bird Survey – a roadside census that has been conducted since 1966 – shows that half of migratory bird species are declining; declines in long-distance Neotropical migrants are more pronounced than those of birds migrating shorter distances (see North American Breeding Bird Survey). These species depend on multiple habitats at different points in space and time, and the reduction in the quality of one habitat can have far-reaching consequences for overall species persistence (Robbins et al. 1989, Zitske et al. 2011, Taylor and Stutchbury 2015, Jackson et al. 2019). Breeding season, though a small proportion of the annual cycles for many migratory birds, is significant due to the more direct association with recruitment and fitness of a species (La Sorte et al. 2017). Therefore, loss and fragmentation of breeding habitats in the United States and other northern locales may have particularly severe ecological implications for imperiled migratory bird populations, especially when breeding distribution is very restricted.

The Golden-cheeked Warbler (*Setophaga chrysoparia*, GCWA) is a migratory songbird listed as endangered under the federal ESA that is heavily impacted by forest conversion. The species is a classic habitat specialist, breeding exclusively in the Edwards Plateau of central Texas, commonly referred to as the Texas Hill Country, and preferring mature, mixed Ashe-juniper and oak woodlands (Pulich 1976, Ladd and Gass 1999, Long et al. 2016). These tree species provide

critical material for nest-building and shelter for main GCWA food sources. A wealth of

literature assessing the influence of habitat factors on measures of GCWA survival (i.e.,

presence, reproductive success, nest survival, and density) finds that forest composition, age, and

patch size are important to species success (Pulich 1976, Shaw and Atkinson 1990, U.S. FWS

1992, Jetté et al. 1998, Magness et al. 2006, Diamond 2007, Colón et al. 2019). Their specialized

preference for this already range-restricted forest type, as well as the high rates of habitat loss

from urban developments and transportation infrastructure, led to the listing of GCWA as

endangered by the U.S. Fish and Wildlife Service in 1990. Since listing, the threat of habitat loss

persists with an estimated 29% reduction in total GCWA breeding habitat between 2000-2010

(Duarte et al. 2013). As of 2020, a very small portion (1.5%) of Texas lands are protected or

managed in a way that is consistent with biodiversity conservation (GAP 1 or GAP 2, U.S.

Geological Survey 2019). As such, the threat of further habitat loss and degradation remains, and

continued monitoring and evaluation of landscapes and populations is necessary to understand

the progress of GCWA recovery (Eichenwald et al. 2020).

A mandate of the ESA (section 4(c)(2)), regular reviews of the best available science and

commercial information are conducted to revisit population trends, threats to recovery, and

accuracy of the listing. Science that demonstrates a range-wide understanding of available

breeding habitat conditions and distribution is critical to engaging federal action for proper

protections for species recovery (La Sorte et al. 2015). Though coarse, this information may be

used to estimate population size and viability and, in the case of habitat conversion, can also

serve to evaluate the status and trends of major threats to species recovery (McGowan et al.

2017). For GCWA, analyses are generally focused on habitat quantity, but additional nuances of

habitat quality may result in a more refined understanding of population dynamics and can help

managers prioritize habitats for conservation and restoration. Most importantly, regular updates

to species habitat quality, quantity, location, and use are necessary to understand longer-term

temporal trends for better informed conservation efforts and for consideration in regular federal

assessments revisiting species listing (and other) decisions. As such, up-to-date spatiotemporal

trends in both the quality and quantity of species habitat at larger ecological scales are important

for informing conservation management and supporting continued protections for threatened and

endangered species.

The last review for GCWA was in 2014 and since then, there have been questions regarding

species recovery and listing status. Currently available science on GCWA breeding habitat

quantity assesses temporal trends on short timeframes (no more than a decade) and is now a

decade out-of-date (Duarte et al. 2013). Additionally, available analyses of habitat quality are

restricted to fractions of the breeding range (Loomis Austin 2008, Heger and Hayes 2013). There

is a need for an amended breeding habitat assessment to reflect recent landscape changes and

data availability as well as to inform upcoming species status assessments and the next steps in

conservation planning.

The objective of this study was to conduct a more comprehensive spatiotemporal model of

GCWA breeding habitat distribution. As part of this study, we used geospatial data spanning the

last 25 years to 1) update range-wide dynamics in GCWA habitat quantity, 2) analyze

spatiotemporal patterns in habitat quality, and 3) compare findings with GCWA sightings and

with local protected areas for considerations of habitat use and conservation, respectively.

METHODS

This study focused on spatiotemporal changes to habitat loss and degradation for the entirety of

the Golden-cheeked Warbler (*Setophaga chrysoparia,* GCWA) breeding range. Breeding and

nesting activities are confined to central Texas, USA where ideal habitat varies in density and

cover. Generally, habitat is more common in the southern and eastern regions of the range.

Nesting habitat is generally defined by the tree species composition. Warblers nest in habitat

made up of mature Ashe-juniper and a combination of other species such as live oak (*Quercus*

fusiformis), Shallow-lobed oak (*Quercus breviloba),* Texas oak (*Q. buckleyi*), post oak (*Q.*

stellata), blackjack oak (*Q. marilandica*), Lacey oak (*Q. glaucoides*), shin oak (*Q. sinuata*),

sugarberry (*Celtis laevigata*), Texas ash (*Fraxinus texensis*), Nuttall's oak (*Quercus taxana),*

cedar elm (*Ulmus crassifolia*), escarpment cherry (*Prunus serotina var. eximia*), pecan (*Carya*

illinoinensis), and little walnut (*Juglans microcarpa*) (U.S. FWS 1992, Texas Parks and Wildlife

Dept). Quality habitat generally occurs in forest patches at least 100 hectares in size with

moderate to high density of older trees. Forests with greater variation in tree height, greater

average tree height, and greater density of deciduous oaks are also associated with higher

densities of GCWA (Wahl et al. 1990).

Data Acquisition

All spatial datasets used are publicly available and were analyzed with relation to GCWA range

as defined by the U.S. Geological Survey's Gap Analysis Program (Table 1). Data were acquired

in the summer of 2020 and analyzed using ArcPro v 2.3 (Esri, USA).

Spatial Analyses

Habitat Disturbance

Google Earth Engine implementation of the LandTrendr algorithm (Kennedy et al. 2018) was

used to identify loss of habitat within the GCWA breeding range between 1985 and 2018 from

Landsat imagery via breakpoints in temporal trends of NDVI (see Eichenwald et al. 2020).

Habitat loss was defined as the area where one habitat was degraded quickly over a short period

of time (including from natural or prescribed burns). For each year, we calculated the area of

disturbed and undisturbed habitat throughout the entire breeding range, in urban and non-urban

portions of the range, and in the metropolitan areas of Austin and San Antonio separately as

defined by U.S. Census urban area boundaries. We conducted identical calculations with forest

loss data from the National Land Cover Database to corroborate the analysis.

Habitat Quality

We applied a previously developed habitat assessment framework to determine location and

acreage of quality GCWA habitat for the entire breeding range in 1985 and the most current year

of the National Land Cover Dataset (2016, Heger and Hayes 2013). Coarser-resolution (250m)

models of historical land use and land cover for the contiguous U.S. were used for estimating

habitat in 1985 (Sohl et al. 2018). To account for differences in data resolution, the land cover

data from 1985 were resampled and masked by historic forest disturbance data (NLCD 2016)

assuming that all pixels labeled as either a) never experiencing a disturbance or b) experiencing a

disturbance after 1985 were forested in 1985. The framework for scoring habitat suitability is

based on a large body of literature citing forest composition, landscape fragmentation, and edge

effects as related to GCWA presence, survival and breeding success (e.g., Pulich 1976, U.S.

FWS 1992, Jetté et al. 1998, Ladd and Gass 1999, Magness et al. 2006, Peak 2007, Long et al.

2016, Reidy et al. 2017, Reidy et al. 2018, Colón et al. 2019). Additionally, Heger and Hayes

tested multiple models and confirmed that an evergreen and mixed forest-based model performed

better than models using only evergreen or mixed forest types. Habitats of highest quality are

intact mixed or evergreen forest cores. Factors and criteria used for scoring habitat quality

include:

Forest type: where mixed or evergreen forest types and deciduous forest within 100m of

mixed/evergreen forest received a 1. All other land cover types received a 0.

Landscape context: neighborhood statistics were used to determine the percent of forest land

cover in a 210m radius. Areas that were 80-100% forested received highest score (4) and areas 0-

20% forested the lowest (0).

Edge effect: scores were docked 1 if they were within 100m of the forest edge.

For each dataset, we calculated the amount of habitat by score throughout the entire breeding

range. Descriptive statistics were also generated to compare results in urban areas and

specifically for Austin and San Antonio metropolitan areas.

Hotspot Analysis

Occurrence data from open-source community science databases (iNaturalist and eBird) were

used to assess hotspots in GCWA sightings. Point locations were grouped by date: sightings

prior to 1995 (n = 647) were considered more closely linked to historic landscape patterns and

sightings after 2010 (n = 14,568) may give more insight on current spatial patterns. Sightings

during breeding season (April to August) were used. Additionally, only non-duplicate points

representing live observations were used if they were associated with an observation date and a

meaningful latitude and longitude (not the centroid of the state or county). Kernel density was

used to calculate density of GCWA sighting per square kilometer, with the top quartile of density

values representing 'hotspots' for GCWA nesting. Centrality and directional distribution of

sightings were also compared between the two time periods. We have high confidence in

drawing conclusions about breeding location based on sighting location because a warbler's

range is on average a 100-m radius around its nest, depending on the quality of habitat (Reidy et

al. 2018).

Protected Areas Overlay

Community science data and results from habitat quantity and quality analyses were used in

overlays to calculate descriptive statistics based on other landscape designations and coverages

from the protected areas database of the U.S. (PADUS v 2.0). We used U.S. Geological Survey's

Gap Analysis Program (GAP) codes, which are specific to the management intent to conserve

biodiversity. GAP 1 and 2 areas are managed in ways typically consistent with conservation and

are considered 'protected' in this context.

RESULTS

Before 1985, 25.64% of Golden-cheeked Warbler (GCWA) breeding range was covered by

forest lands (over 4.54 million acres). Between 1985 and 2016, 13% of all forests within the

warbler's breeding range were converted to other land uses (Figure 1). Forest conversion was

more extreme in parts of the range in metropolitan areas, with 24% forest loss in the Austin

metropolitan area and 32% loss in the San Antonio metropolitan area. Generally, for all regions,

there were greater rates of decline in more recent years. Habitat quality also declined during this

time period throughout the breeding range (Figure 2). In the 1980s, over one-tenth of the forested

habitat within GCWA breeding range was intact mixed or evergreen core forests. In 2016, high

quality habitat made up 5% of the breeding range, indicating a 45% decrease in the highest

quality breeding habitat (Figure 3). Remaining quality habitat is more fragmented, with

significantly smaller patch sizes than in 1985 (Figure 4; t = 1.96, p value <0.001). Generally,

quality habitat is more concentrated along the southeastern extent of the breeding range and

some forested areas to the northwest of Austin and in the northern parts of the breeding range

have improved since 1985 (Figure 2). Habitats within protected areas (i.e., GAP Code 1 & 2)

saw a less sharp decline in habitat quality from 27% to 20% of the breeding range (27%

decrease; Figure 3). However, protected habitats currently represent only 10% of all highest-

quality habitat in the breeding range.

GCWA sightings were generally spatially coincident with habitat quality. In the 1980s, 39% of

sightings were in high-quality habitat. As of 2020, sightings in high-quality habitat had dropped

to 28%, but this is still disproportionately high given that only 5% of the breeding range consists

of high-quality breeding habitat. The proportion of sightings in protected areas in the breeding

range has increased dramatically from 5% of sightings before 1995 to 59% after 2010. There

were small, localized shifts in the location of GCWA sighting hotspots between the 1980s and

2020, but breeding range-wide, the distribution of hotspots (centrality and dispersion) remains

the same (Figure 4). Sightings were once very concentrated to the southeastern portions of the

range, but are now less concentrated, with a few hotspots formed in western parts of the range.

Parks in and around San Antonio metropolitan area and Lost Maples Natural Area continue to be

hotspots for sightings. All but one of the hotspots were associated with a protected area (Figure

4).

DISCUSSION

We quantified the absolute change in forest cover within Golden-cheeked Warbler (GCWA)

breeding range and changes in breeding habitat quality over a 30-year period. Overall, we

estimated a 13% loss in breeding habitat with high spatial heterogeneity in landscape conversion

closely tied to human developments. This value is lower than other estimates and may be

reflective of changing forest dynamics that occur over the longer-term study period- this would

support our findings that greater decline has occurred in more recent years (Groce et al. 2010,

Duarte et al. 2013). Additionally, drastic declines in habitat quality suggest that 13% is an

underestimation of effective habitat loss as degradation may impact the ecological viability of

remaining forests for GCWA nesting. The amount of intact core forest habitat fell 45% in the 30-

year period leaving quality breeding sites concentrated along the southeastern extent of the

breeding range and in protected areas.

Human impact is on the rise in Texas landscapes and may compromise habitat quality. In recent

years, Texas has had the largest increases in population of any state in the U.S. (U.S. Census

Bureau 2020). In the 30-year period that was studied, the population grew 73.9% and growth is

projected to continue at a steady rate to 2050 (88.3% increase in the next three decades; Texas

Demographic Center 2019). Within GCWA breeding range, at least four counties are projected

to see population increases of over 100% by 2050, all of which coincide with areas of quality

habitat in the southeastern parts of the range: Williamson, Hayes, Comal, and Kendall counties.

Increased development pressures in the Texas Hill Country could continue to drive the trends of

GCWA habitat disturbance and degradation. Our data indicate that quality forests have

undergone fragmentation resulting in smaller habitat patches. Similar trends have been reported

more specifically for Ashe-juniper distributions across the state due to an increase in pastureland

and development (Diamond 1997). A reduction in canopy cover can lead to decreased nest

success for forest songbirds (Martin and Roper 1988, Trzcinski et al. 1999, Twedt et al. 2001).

Canopy cover is also essential to conceal GCWA nests located in the mid-story to upper

canopies of trees, thus reducing the probability of nest predation and parasitism (Reidy et al.

2008). Additionally, fragmentation of breeding habitat may represent barriers to dispersal of

birds and important genetic material (Lindsay et al. 2008). Hence, there is already evidence of

notable genetic differentiation among populations of GCWA, having important implications for

management of species like GCWA that are relatively vagile, but highly specialized in their

habitat preferences. Restoration and protection of connected patches may be the best option for

conserving or recovering such species (Young and Clarke 2000).

We found that only 10% of the highest quality forest habitat are in protected areas, creating both

challenges and opportunities. These lands, because they are managed in ways consistent with

biodiversity conservation, generally represent higher quality habitats with fewer human

disturbances (Rosa and Malcom 2020). Our findings indicate that protected areas within GCWA

breeding range also exhibited declines in quality, but degradation was buffered relative to the

overall range. As human populations grow and landscape conversion continues, protected areas

are expected to grow in importance. Nearly all (17 out of 18) of GCWA sighting hotspots from

our analysis were associated with a protected area. Additionally, the proportion of sightings that

occurred on protected areas saw a significant increase. It should be noted that public lands may

have a higher proportion of sighting simply due to their accessibility to observers. Nevertheless,

protected areas still had significantly more sightings than areas without protections

demonstrating that they are important to GCWA success. Public protected areas can play a

central role in habitat conservation efforts because they are more amenable to the application of

broad-scale management strategies that more closely align with species conservation. However,

the extent to which public protected areas can benefit migratory bird populations depends on

how well protected areas are represented within the breeding range (La Sorte et al. 2015).

Currently, areas managed for conservation (GAP status 1 and 2) represent 3.23% of the breeding

range. Lands with more intermediate mandates (GAP 3) provide a higher degree of flexibility for

the implementation of management recommendations more closely aligned with maintaining

biodiversity. However, these lands are also limited in the state of Texas (1.71%). Expansion of

protections to key habitats would require that resources be spent in agency land acquisition or in

private lands conservation.

Our findings demonstrate a need for strengthening current conservation measures and expanding

upon protections for GCWA habitat to ensure greater breeding success and, ultimately, species

recovery. Newer proposals to protect at least 30% of U.S. lands and waters by 2030 to address

the biodiversity and climate crises may provide additional opportunities for land designations

and conservation efforts for imperiled species like GCWA (Exec Order No 14008 2021, CA

Exec Order N-82-20 2020). While a majority of GCWA habitat conservation dollars have been

spent conserving GCWA breeding habitat on the outskirts of the cities of Austin and San

Antonio, our findings support previous work demonstrating higher rates of habitat conversion

near metropolitan areas (Duarte et al. 2013). Given the scarcity of public lands, the distribution

of intact forest habitat, and the relatively high amount of habitat loss and degradation occurring

in and around metropolitan areas, future GCWA habitat conservation efforts should be more

focused on supporting current protected areas and expanding protections to quality habitats in the

Balcones Canyonlands and Fort Hood areas and regions west of San Antonio and Fort Worth.

Additionally, projected species distribution models reflecting climate change impacts on tree

species indicate that the Texas Hill Country will continue to be a stronghold for Ashe-juniper

(with potential for population stabilization and maybe even increase/spread to the northeast;

McKenney et al. 2007). This suggests that efforts to conserve or restore quality GCWA habitat

will have long-term benefits.

We recognize the limitations of the analysis which equate all available mixed or evergreen

forests within the breeding range, and not strictly those with Ashe-juniper components, as

potentially suitable habitat for GCWA nesting. This is mainly due to current publicly available

data sources and lack of LiDAR or other advanced geospatial datasets that would clarify spectral

or structural differences in forest composition. Regardless, overall classification accuracies of the

habitat loss dataset followed methods that average a mean absolute error of less than 3%

(Kennedy et al. 2018). Additionally, datasets used for assessing habitat quality, though they

represent the most current version available, are already out of date. Collectively, this indicates

that our estimates for available habitat may be more liberal than in actuality. In the context of

federal species listing and review, landscape change analyses, habitat identification and

classification, and the characterization of trends over time must be considered. We selected

metrics that are easily applied to larger scales and interpretable to a wide audience of resource

managers and policy makers, regardless of biogeographic, ecological, or geospatial expertise and

training. Nonetheless, selection of metrics may in some cases influence how landscape change is

quantified and interpreted, particularly when the analysis is focused on characterizing or

understanding underlying landscape processes and species modeling.

Human landscape modification is likely to continue in the Texas Hill Country, but conservation

and land management actions can be taken to minimize further habitat loss and degradation in

GCWA breeding range. This information will assist researchers and managers in prioritizing

range-wide breeding habitat conservation efforts and highlights the significant role land

management for conservation biodiversity plays on the landscape. There remains a need to grow

the network of protected areas for GCWA restoration. Further, continued regular spatiotemporal

assessments of habitat quantity and quality are necessary to assess changes to species potential

for persistence and extrapolate population viability given these dynamics.

LITERATURE CITED

Bairlein, F. (2016). Migratory birds under threat. Science 354:547-548.

Colón, M.R., M.R. Hutchinson, A.M. Long, G. Voelker, and M.L. Morrison (2019).

Postbreeding Habitat Use by Golden-Cheeked Warblers (*Setophaga chrysoparia*). Western

North American Naturalist 79(3):337-346.

Diamond, D.D. (2007). Range-wide modeling of Golden-cheeked warbler habitat. Unpublished

report to TPWD. Columbia, Missouri: University of Missouri.

Diamond, D.D., G.A. Rowell, and D.P. Keddy-Hector (1995). Conservation of Ashe Juniper

(Juniperus ashei Buchholz) woodland of the central Texas hill country. Natural Areas Journal

15: 189-202.

Duarte, A., J.L.R. Jensen, J.S. Hatfield, and F.W. Weckerly (2013). Spatiotemporal variation in

range-wide golden-cheeked warbler breeding habitat. Ecosphere 4(12):152.

Eichenwald, A.J., M.J. Evans, and J.W. Malcom (2020). US imperiled species are most

vulnerable to habitat loss on private lands. Frontiers in Ecology and the Environment 18(8):

439-446.

Groce, J.E., H.A. Mathewson, M.L. Morrison, and N. Wilkins (2010) Scientific evaluation for

the 5-Year status review of the golden-cheeked warbler. Prepared for U.S. Fish and Wildlife

Service.

Hanski I. (2011). Habitat loss, the Dynamics of Biodiversity, and a Perspective on Conservation.

AMBIO 40:248-255.

Heger N., and T. Hayes (2013). Change in Golden-cheeked Warbler habitat through time.

Golden-cheeked Warbler Symposium, Austin, TX.

Jackson M.V., L.R. Carrasco, C.Y. Choi, J. Li, Z. Ma, D.S. Melville, T. Mu, H.B. Peng, B.K.

Woodworth, Z. Yang, L. Zhang, and R.A. Fuller (2019). Multiple habitat use by declining

migratory birds necessitates joined up conservation. Ecology and Evolution 9(5): 2505-

2515.

Jette L.A., T.J. Hayden, J.D. Cornelius (1998). Demographics of the golden-cheeked warbler

(*Dendroica Chrysoparia*) on Fort Hood, Texas. Construction Engineering Research

Laboratory, US Army Engineer Research and Development Center, Champaign, Illinois.

Technical Report 98/52.

Kennedy R.E., Z. Yang, N. Gorelick, J. Braaten, L. Cavalcante, W.B. Cohen, and S. Healy

(2018). Implementation of the LandTrendr Algorith on Google Earth Engine. Remote

Sensing 10(5): 691.

La Sorte, F.A., D. Fink, P.J. Blancher, A.D. Rodewald, V. Ruiz-Gutierrrez, K.V. Rosenberg,

W.M. Hochachka, P.H. Verburg, and S. Kelling (2017). Global change and the distributional

dynamics of migratory bird populations wintering in Central America. Global Change

Biology 23(12):5284-5296.

La Sorte, F.A., D. Fink, W.M. Hochachka, J.L. Aycrigg, KV. Rosenberg, A.D. Rodewald, N.E.

Bruns, A. Farnsworth, B.L. Sullivan, C. Wood, and S. Kelling (2015). Documenting

stewardship responsibilities across the annual cycle for birds on U.S. public lands. Ecological

Applications 25(1): 39-51.

Ladd, C.G. and L. Gass (1999). Golden-cheeked Warbler (*Dendroica chrysoparia*). Birds of

North America 420, Philadelphia.

Leu, M., A.M. Haines, C.E. Check, D.M. Costante, J.C. Evans, M.A. Hollingsworth, I.T.

Ritrovato, A.M. Rydberg, A.M. Sandercock, K.L. Thomas, and T.C. Treakle (2019).

Temporal analysis of threats causing species endangerment in the United States. Conservation

Science and Practice 1(8): e78.

Lindsay, D.L., K.R. Barr, R.F. Lance, S.A. Tweddale, T.J. Hayden, and P.L. Leberg (2008).

Habitat fragmentation and genetic diversity of an endangered, migratory songbird, the golden-

cheeked warbler (*Dendroica chrysoparia*). Molecular Ecology 17(9): 2122-2133.

Long, A.M., J.C. Newnam, M.R. Colon, K.N. Smith-Hicks, and M.L. Morrison (2016). Characteristics of Vegetation Used by Golden-Cheeked Warblers in Central Texas. Southeastern Naturalist 15(1):153-161.

Loomis Austin (2008). Mapping potential golden-cheeked warbler breeding habitat using remotely sensed forest canopy cover data. Report LAI Project No. 051001. Austin, TX: Loomis Austin.

Magness, D.R., R.N. Wilkins, and S. J Hejl (2006). Quantitative relationships among golden-cheeked warbler occurrence and landscape size, composition, and structure. Wildlife Society Bulletin 34:473-479.

Martin, T.E and J.J. Roper (1988). Nest predation and nest-site selection of a western population of the hermit thrush. The Condor 90: 51-57.

McGowan, C. P., N. Allan, J. Servoss, S. Hedwall, and B. Woldridge (2017). Incorporating population viability models into species status assessment and listing decisions under the U.S. Endangered Species Act. Global Ecology and Conservation 12: 119-130.

McKenney, D.W., J.H. Pedlar, K. Lawrence, K. Campbell, and M.F. Hutchinson (2007). Potential impacts of climate change on the distribution of North American trees. BioScience 57(11): 939-948.

Pulich, W.M. (1976). The Golden cheeked Warbler, A Bioecological Study. Austin, Texas: Texas Parks & Wildlife Dept. 172 pp.

Reidy, J.L., M.M. Stake, and F.R. Thompson (2008). Golden-cheeked warbler nest mortality and predators in urban and rural landscapes. The Condor 110: 458-466.

Reidy, J.L., F.R. Thompson III, L. O'Donnell (2017). Density and Nest Survival of Golden-

Cheeked Warblers: Spatial Scale Matters. The Journal of Wildlife Management 81(4): 678-

689.

Reidy, J.L., F.R. Thompson III, G.M. Connette, and L. O'Donnell (2018). Demographic rates of

Golden-cheeked Warblers in an urbanizing woodland preserve. The Condor 120: 249-264.

Robbins, C.S., J.R. Sauer, R.S. Greenberg, and S. Droege (1989). Population declines in North

American birds that migrate to the neotropics. Proceedings of the National Academy of

Science 86:7658-7662.

Rosa, L.D., and J.W. Malcom (2020). Getting to 30x30: guidelines for decision-makers.

https://dfnd.us/2Orbk7V

Shaw, D.M. and S.F. Atkinson (1990). An Introduction to the Use of Geographic Information

Systems for Ornithological Research. The Condor 92(3):564–570.

Sohl, T.L., R. Ryan, M. Bouchard, K. Sayler, J. Dornbierer, S. Wika, R. Quenzer, and A. Friesz

(2018). Modeled historical land use and land cover for the conterminous United States: 1938-

1992: U.S. Geological Survey data release.

Taylor C.M. and B.J.M. Stutchbury (2015). Effects of breeding versus winter habitat loss and

fragmentation on the population dynamics of a migratory songbird. Ecological Applications

26(2):424-437.

Texas Demographic Center (2019). Texas population projections 2010-2050:

https://demographics.texas.gov/Resources/publications/2019/20190128_PopProjectionsBrief.

pdf

Texas Parks and Wildlife Dept. Golden-Cheeked Warbler.

https://tpwd.texas.gov/publications/pwdpubs/media/pwd_bk_w7000_0013_golden_cheeked_warbler.pdf

Thompson, P.L.B. Rayfield, and A. Gonzalez (2016). Loss of habitat and connectivity erodes species diversity, ecosystem functioning, and stability in metacommunity networks. Ecogeography 40(1):98-108.

Trzcinski, M.K., L. Fahrig, and G. Merriam (1999). Independent effects of forest cover and fragmentation on the distribution of forest breeding birds. Ecological Applications 9(2): 586-593.

Twedt, D.J. and J. Henne-Kerr (2001). Artificial cavities enhance breeding bird densities in managed forests. Wildlife Society Bulletin 29: 680-687.

U.S. Census Bureau (2020). National Demographic Analysis: https://www.census.gov/programs-surveys/decennial-census/about/coverage-measurement/da.html

U.S. Fish and Wildlife Service (1992) Golden-Cheeked Warbler (*Dendroica Chrysoparia*) Recovery Plan. Albuquerque, New Mexico.U.S. Geological Survey (2019) Protected Areas Database of the US.

Wahl, R., D.D. Diamond, and D. Shaw (1990) The golden-cheeked warbler: a status review. Final report, submitted to the Office of Endangered Species, U. S. Fish and Wildlife Service, Albuquerque, New Mexico.

Young, A.C., and G.M. Clarke (2000). Genetics, demography and the viability of fragmented populations. Cambridge University Press, Cambridge.

447   Zitske, B.P., M.G. Betts, A.W. Diamond (2011). Negative Effects of Habitat Loss on Survival of

448   Migrant Warblers in a Forest Mosaic. Conservation Biology 25(5):993-1001.

454   **TABLES**

455   Table 1. Data acquired for spatiotemporal analyses on habitat disturbance, quality, and sightings.

Data	Source	Temporal Resolution	Spatial Resolution	Analysis
Landscape change	Eichenwald et al. 2020	1985-2018	30m	Habitat disturbance
Forest Change	MRLC National Land Cover Database	1985-2016	30m	Habitat disturbance
Modeled historic land use	Sohl et al. 2018	1985	250m	Habitat quality
NLCD	MRLC	2016	30m	Habitat quality
Warbler sightings	iNaturalist & eBird	1934-2020	point	Sighting Hotspots
Protected Areas	U.S. Geological Survey	2019	vector	All
Urban areas	U.S. Census	2019	vector	Habitat disturbance
Breeding range	U.S. Geological Survey GAP	2001	vector	Study area

Figure 1

  FIGURES

Figure 2

Figure 3

Figure 4

Figure 5

483
484

FIGURE CAPTIONS

Figure 1. Annual declines in forested land cover relative to a 1985 baseline for portions of the Golden-cheeked Warbler breeding range that fall outside of urban boundaries and portions that fall within the urban boundaries of San Antonio and Austin, respectively.

Figure 2. A map of overall change in Golden-cheeked Warbler breeding habitat quality between 1985 and 2016. Blues indicate areas of lower-quality habitat in 1985, white indicates high-quality habitat in 1985, and red indicates high-quality habitat areas that experienced a decline in habitat quality between 1985 and 2016.

Figure 3. Percent of the Golden-cheeked Warbler breeding range by habitat quality value (0 = low quality, 4 = high quality) for habitats throughout the entire range and for habitats that fall inside protected areas managed for biodiversity conservation (U.S. Geological Survey's Protected Areas Database of the U.S., GAP codes 1 and 2) for 1985 and 2016, respectively.

Figure 4. Histograms showing frequency of patch size for high quality habitat areas in 1985 (blue) and 2016 (red) indicate habitat fragmentation over the time period. Means of the two groups were significantly different at $\alpha = 0.05$ (p value <0.001).

Figure 5. A map comparing locations of high-quality habitat in 2016, protected areas managed for biodiversity conservation (U.S. Geological Survey's Protected Areas Database of the U.S.,

512   GAP codes 1 and 2), and hotspots for Golden-cheeked Warbler sightings between 1980 – 2000

513   (in blue) and 2010 – 2020 (in red).