Phylogenomic discordance suggests polytomies along the backbone of the large genus *Solanum*

Edeline Gagnon[1,2*], Rebecca Hilgenhof[1,2], Andrés Orejuela[1,2], Angela McDonnell[3], Gaurav Sablok[4,5], Xavier Aubriot[6], Leandro Giacomin[7], Yuri Gouvêa[8], Thamyris Bragionis[8], João Renato Stehmann[8], Lynn Bohs[9], Steven Dodsworth[10,11], Christopher Martine[12], Péter Poczai[4,13], Sandra Knapp[14], Tiina Särkinen[1]

[1]Royal Botanic Garden Edinburgh, 20A Inverleith Row, Edinburgh, EH3 5LR, United Kingdom

[2]School of Biological Sciences, University of Edinburgh, King's Buildings, Mayfield Road, Edinburgh, EH9 3JH, United Kingdom

[3]Negaunee Institute for Plant Conservation Science and Action, Chicago Botanic Garden, 1000 Lake Cook Rd, Glencoe, IL 60022, United States of America

[4]Finnish Museum of Natural History (Botany Unit), University of Helsinki, PO Box 7 FI-00014 Helsinki, Finland

[5]Organismal and Evolutionary Biology Research Programme (OEB), Viikki Plant Science Centre (ViPS), PO Box 65, FI-00014 University of Helsinki, Finland

[6]Université Paris-Saclay, CNRS, AgroParisTech, Écologie, Systématique et Évolution, 91405, Orsay, France

[7]Instituto de Ciências e Tecnologia das Águas & Herbário HSTM, Universidade Federal do Oeste do Pará, Rua Vera Paz, sn, Santarém, CEP 68040-255, PA, Brazil

[8]Departamento de Botânica, Instituto de Ciências Biológicas, Universidade Federal de Minas

Gerais – UFMG, Av. Antônio Carlos, 6627, Pampulha, Belo Horizonte, CEP 31270-901, MG,

Brazil

[9]Department of Biology, University of Utah, Salt Lake City, Utah, United States of America

[10]School of Life Sciences, University of Bedfordshire, University Square, Luton LU1 3JU,

United Kingdom

[11]Royal Botanic Gardens, Kew, Richmond TW9 3AE, Surrey, United Kingdom

[12]Department of Biology, Bucknell University, Lewisburg, PA 17837, United States of America

[13]Faculty of Environmental and Biological Sciences, University of Helsinki, FI-00014 Finland

[14]Department of Life Sciences, Natural History Museum, Cromwell Road, London SW7 5BD,

United Kingdom.

*Corresponding author, E-mail: edeline.gagnon@gmail.com

Short title: Phylogenomic discordance in *Solanum*

Abstract

Premise of the study: Evolutionary studies require solid phylogenetic frameworks, but increased volumes of phylogenomic data have revealed incongruent topologies among gene trees in many organisms both between and within genomes. Some of these incongruences indicate polytomies that may remain impossible to resolve. Here we investigate the degree of gene-tree discordance in *Solanum,* one of the largest flowering plant genera that includes the cultivated potato, tomato, and eggplant, as well as 24 minor crop plants.

Methods: A densely sampled species-level phylogeny of *Solanum* is built using unpublished and publicly available Sanger sequences comprising 60% of all accepted species (742 spp.) and nine regions (ITS, *waxy*, and seven plastid markers). The robustness of this topology is tested by examining a full plastome dataset with 140 species and a nuclear target-capture dataset with 39 species of *Solanum* (Angiosperms353 probe set).

Key results: While the taxonomic framework of *Solanum* remained stable, gene tree conflicts and discordance between phylogenetic trees generated from the target-capture and plastome datasets were observed. The latter correspond to regions with short internodal branches, and network analysis and polytomy tests suggest the backbone is composed of three polytomies found at different evolutionary depths. The strongest area of discordance, near the crown node of *Solanum,* could potentially represent a hard polytomy.

Conclusions: We argue that incomplete lineage sorting due to rapid diversification is the most likely cause for these polytomies, and that embracing the uncertainty that underlies them is crucial to understand the evolution of large and rapidly radiating lineages.

Recent advances in high-throughput sequencing have provided larger molecular datasets,

including entire genomes, for reconstructing evolutionary relationships (e.g. Ronco et al., 2021).

Considerable progress has been made since the publication of the first molecular-based

classification of orders and families of flowering plants (APG, 1998), with one of the most recent

examples including a phylogenetic tree of the entire Viridiplantae based on transcriptome data

from more than a thousand species (One Thousand Plant Transcriptomes Initiative, 2019). Whilst

large datasets have strengthened our understanding of evolutionary relationships and

classifications across the Tree of Life, several of them have demonstrated repeated cases of

persistent topological discordance across key nodes in birds (Suh et al., 2015; Suh, 2016),

mammals (Morgan et al., 2013; Romiguier et al., 2013; Simion et al., 2017), amphibians (Hime

et al., 2021), plants (Wickett et al., 2014; One Thousand Plant Transcriptomes Initiative, 2019),

and fungi (Kuramae et al., 2006). Whereas previous expectations were that these "soft

polytomies" would be improved with the addition of more data, their persistence after addition of

more taxonomic and molecular data have led some authors to suggest that they actually represent

"hard polytomies", i.e., extremely rapid divergence events of three or more lineages at the same

time or reticulate evolution due to species hybridisation and/or introgression. In an era where

obtaining genome-wide sampling of species for phylogenetic reconstruction has become

mainstream, the question about whether persistent topological discordance can be resolved with

more data or whether they reflect complex biological realities (Jeffroy et al., 2006; Philippe et

al., 2011) is becoming increasingly common.

Discordance in phylogenetic signal can be due to three general classes of effects (Wendel

and Doyle, 1998): (1) technical causes such as gene choice, sequencing error, model selection, or

poor taxonomic sampling (Philippe et al., 2011, 2017); (2) organism-level processes such as

rapid or convergent evolution, rapid diversification, incomplete lineage sorting (ILS), or

horizontal gene transfer (Degnan and Rosenberg, 2009), and (3) gene and genome-level

processes such as interlocus interactions and concerted evolution, intragenic recombination, use

of paralogous genes for analysis, and/or non-independence of sites used for analysis. Together,

these biological and non-biological processes can lead to conflicting phylogenetic signals

between different loci in the genome and hinder the recovery of the evolutionary history of a

group (Degnan and Rosenberg, 2009). Consequently, careful assessment of phylogenetic

discordance across mitochondrial, plastid, and nuclear datasets is critical for understanding

realistic evolutionary patterns in a group, as traditional statistical branch support measures fail to

reflect topological variation of the gene trees underlying a species tree (Liu et al., 2009; Kumar

et al., 2012).

Here we explore the presence of topological discordance in nuclear and plastome

datasets of the large and economically important angiosperm genus *Solanum* L. (Solanaceae)*,*

which includes 1,228 accepted species and several major crops and their wild relatives, including

potato, tomato and brinjal eggplant (aubergine), as well as at least 24 minor crop species

(Solanaceaesource.org, Nov. 2020). Building a robust species-level phylogeny for *Solanum* has

been challenging because of the sheer size of the genus, and because of persistent poorly

resolved nodes along the phylogenetic backbone. Bohs (2005) published the first plastid

phylogenetic analysis for *Solanum* and established a set of 12 highly supported clades based on

her strategic sampling of 112 species (9% of the total species number in the genus), spanning

morphological and geographic variation. As new studies have emerged with increased taxonomic

and genetic sampling (e.g., Levin et al., 2006; Weese and Bohs, 2007; Stern et al., 2011;

Särkinen et al., 2013; Tepe et al., 2016), the understanding of overall phylogenetic relationships

within *Solanum* has evolved to recognise three main clades: (1) the Thelopodium clade

containing three species sister to the rest of the genus, (2) Clade I containing c. 350 mostly

herbaceous and non-spiny species (including the Tomato, Petota, and Basarthrum clades that

contain the cultivated tomato, potato, and pepino, respectively), and (3) Clade II consisting of c.

900 predominantly spiny and shrubby species, including the cultivated brinjal eggplant (Table 1).

The two latter clades are further resolved into 10 major and 43 minor clades (Table 1).

Despite these advancements, phylogenetic relationships between many of the major

clades of *Solanum* have remained poorly resolved, mainly due to limitations in taxon and

molecular marker sampling. The most recent genus-wide phylogenetic study by Särkinen et al.

(2013), based on seven markers (two nuclear and 5 plastid) and fewer than half (34%) of the

species of *Solanum,* failed to resolve the relationships between major clades, especially within

Clade II and the large component Leptostemonum clade, which includes the Old World spiny

clade, comprising almost all spiny *Solanum* species that occur in the eastern hemisphere. To

reduce colonial connotations associated with this name, we hereafter refer to this clade as the

Eastern Hemisphere Spiny clade (EHS; Table 1).

To gain a better understanding of the evolutionary relationships of *Solanum,* we built a

new Sanger supermatrix that included 60% of the species of the genus and compared the

phylogenetic relationships obtained with the Sanger supermatrix with genus-wide plastid (PL)

and nuclear target-capture (TC) phylogenomic datasets. We ask: (1) Does a significant increase

in taxon sampling of the supermatrix dataset lead to significant changes in the circumscription of

major and minor clades in *Solanum*?; (2) Does increased gene sampling in both plastome and

nuclear data resolve previously identified polytomies between major clades?; (3) Is there

evidence of discordance within and between genomic datasets?; and (4) Are areas of high

discordance in the *Solanum* phylogeny better represented by polytomies rather than bifurcating

nodes? Comparison of the topologies from the different datasets, and results from discordance

analyses, a filtered supertree network, and polytomy tests lead us to suggest that some of the soft

polytomies of *Solanum* might be hard polytomies caused by rapid speciation and diversification

coupled with ILS. We discuss the consequences that such an interpretation has for investigating

the biogeography and morphological trait evolution across the economically important genus.

MATERIAL AND METHODS

Taxon sampling–

A Sanger sequence supermatrix was generated including all available sequences from

GenBank related to the genus *Solanum* for nine regions: the nuclear ribosomal internal

transcribed spacer (ITS), low-copy nuclear region *waxy* (i.e., GBSSI), two protein-coding plastid

genes *matK* and *ndhF*, and five non-coding plastid regions (*ndhF-rpl32*, *psbA-trnH*, *rpl32-trnL*,

trnS-G, and *trnT-L*). Only vouchered and verified samples were utilized. All sequences were

blasted against target regions in USEARCH v.11 (Edgar, 2010). Taxon names were checked

against SolanaceaeSource synonymy (solanaceaesource.org, Nov. 2020) and duplicate sequences

belonging to the same species were pruned out to retain a single individual per taxa. A total of

817 Sanger sequences were generated and added to the matrix, adding 129 previously unsampled

species and new data for 257 species (Appendix S1; see the Supplementary Data with this

article). Final species sampling across major and minor clades of *Solanum* varied from 13-100%,

with 742 species of *Solanum* (60% of the 1,228 currently accepted species, Nov 2020; Table 1).

Four species of *Jaltomata* Schltdl. were used as an outgroup (Appendix S1).

To assess phylogenetic discordance within *Solanum*, a set of species was selected for the

phylogenomic study to represent all 10 major and as many of the 43 minor clades of *Solanum* as

possible (Table 1), as well as the outgroup *Jaltomata*. The final sampling included 151 samples

for the plastome (**PL**) dataset (140 *Solanum* species; Table 1 and Appendix S2) and 40 samples

for the target-capture (**TC**) dataset (39 *Solanum* species; Table 1 and Appendix S3). For the PL

dataset, 86 samples were sequenced using low-coverage genome skimming, and the remaining

samples were downloaded from GenBank (Nov 2019). For the TC dataset, 12 samples were

sequenced as part of the Plant and Fungal Trees of Life project (Baker et al., 2021) using the

Angiosperms353 bait set (Johnson et al., 2019). In addition, 17 sequences were added from an

unpublished dataset provided by A. McDonnell and C. Martine. Sequences for the remaining 12

samples were extracted from the GenBank SRA archive using the SRA Toolkit 2.10.7

(https://github.com/ncbi/sra-tools; Appendix S3).

DNA extraction, library preparation & sequencing–

Supermatrix Sanger sequencings– DNA extractions for Sanger sequencing were done

using DNeasy plant mini extraction kits (Qiagen, Valencia, California, USA) or the FastDNA kit

(MP Biomedicals, Irvine, California, USA). Amplification of *waxy* followed Levin et al. (2005)

using two (waxyF with 1171R and 1058F with 2R) or four primer pairs (waxyF with Ex4R,

Ex4F with 1171R, 1058F with 3'N, and 3F with 2R). *trnT-L* was amplified with primers a-d and

c-f (Taberlet et al., 1991; Bohs and Olmstead, 2001; Bohs, 2004). *ndhF* amplification followed

Bohs and Olmstead (1997), *psbA-trnH* followed Sang et al. (1997), *matK* followed Rosario et al.

(2019), ITS and *trnS-G* followed Levin et al. (2006), and *rpl32-trnL* and *ndhF-rpl32* followed

Miller et al. (2009). Sequencing was carried out on ABI automated sequencers at the University

of Utah DNA sequencing facility (Salt Lake City, UT, USA), at the Natural History Museum

(London, United Kingdom), and at Myleus Biotecnologia (Belo Horizonte, Brazil). Contigs were

visually checked in Sequencher v.4.8 (GeneCodes, Ann Arbor, Michigan, USA) and Geneious

Prime 2020.1.1 (https://www.geneious.com). The combined matrix was 10,908 bp long

(Appendix S4). The two most densely sampled regions (*trnT-L* and ITS) included 84% and 82%

of the sampled species, respectively; *waxy* (54%) and ITS (67%) loci had the most parsimony

informative characters (Appendix S4).

PL and TC datasets– DNA for high-throughput sequencing was extracted using the low-

salt CTAB method (Arseneau et al., 2017) and quantified on a Qubit fluorometer (Thermo Fisher

Scientific, USA). Genome skimming was done at the Institute of Biotechnology, University of

Helsinki (Finland). A paired-end genomic library was constructed using the Nextera DNA

library preparation kit (Illumina, San Diego, CA, USA). Fragment analysis was conducted with

an Agilent Technologies 2100 Bioanalyzer using a DNA 1000 chip. Sequencing was performed

on an Illumina MiSeq platform from both ends with a read length of 150 bp. DNA extraction,

quantification, and sequencing for TC followed Johnson et al. (2019). All PL and TC reads have

been submitted to Genbank and the European Nucleotide Archive (Appendices S2-S3).

Phylogenetic analyses–

Overview of methodological strategy– Ten phylogenetic analyses with different

methodological strategies were compared across the supermatrix, PL and TC datasets, to test if

the phylogenetic results were robust despite these different choices (e.g., Philippe et al., 2011,

2017; Saarela et al., 2018; Duvall et al., 2020). The Sanger supermatrix analyses based on

Maximum Likelihood (ML) and Bayesian inference (BI) were used as a reference to compare

results from the PL and TC species trees because the Sanger supermatrix had the most complete

taxonomic sampling (Table 2). For the PL dataset, a total of four analysis were compared to test

the effect of missing data and sampling on the resulting phylogenies, as well as the effect of

different partitioning schemes in IQ-TREE 2 (Table 2; Minh, Schmidt, et al., 2020). For the TC

dataset, a total of four analyses were compared to test the effect of the phylogenetic method (ML

vs. coalescent methods), missing data, and taxonomic sampling on the resulting phylogenies

(Table 2). Full methods for all analyses are described below. All bioinformatic analyses were run

either on the Toby-G1 server at the Royal Botanical Gardens of Edinburgh, or the Crop Diversity

Server from the James Hutton Institute, in Dundee, Scotland, except for the supermatrix ML

analysis.

Supermatrix dataset– Sequences were aligned in MAFFT v.7 (Katoh et al., 2005),

manually checked, and optimised. Short multi-repeats and ambiguously aligned regions were

excluded manually or with trimAl (-gappyout method; Capella-Gutiérrez et al., 2009). Both ML

and BI analyses were run on individual loci, as well as on a combined plastid alignment (seven

loci in total) to check for topological incongruences, rogue taxa, and misidentified sequences.

Visual checks revealed a small number of clear mis-determinations and/or lab errors. A further

26 samples were removed based on high RogueNaRok scores (Aberer et al., 2013). Nuclear

sequence data (ITS and *waxy*) were identified for all known polyploid species (63 species,

Appendix S5), and subsequently examined to determine if there were any strong incongruences

with the results from the plastid loci. As none were found (Appendices S6-S7), sequences from

these species were kept in the final supermatrix analysis.

Maximum likelihood (ML) and Bayesian inference (BI) analyses were run on all nine

loci individually and on the combined plastid dataset (seven loci). ML analyses were run in

RaxML-HPC v.8.2.12 (Stamatakis, 2014) on XSEDE on CIPRES Science Gateway v.3.3 (Miller

et al., 2010), with 10 independent runs based on unique starting trees. The General Time

Reversible (GTR) model with CAT (Tavaré and Others, 1986; Stamatakis, 2006) was used for all

partitions. A total of 1,000 non-parametric bootstraps were run; bootstrap support (BS) ≥95%

was considered strong, 75-94% moderate, and 60-74% weak.

BI analyses were run using Beast v.2.6.3 (Bouckaert et al., 2019), with two parallel runs

sampling trees every 10,000 generations. ModelTest-NG (Darriba et al., 2020) was used to find

the most suitable nucleotide substitution model for the individual loci and combined plastid loci;

JC + G4 was specified for the *ITS* and *trnS-G* regions, GTR+G4 for the *psbA-trnH, trnL-T, rpL32*

and *matK* regions, and the GTR+I+G4 model for all other regions, as well as the combined plastid

dataset and the full supermatrix dataset. For all analyses, an uncorrelated log normal relaxed clock,

birth-death tree prior, and a normally distributed UCLD.mean prior was specified (mean 1,

SD=0.3). All runs were checked with Tracer v.1.7.1 (Rambaut et al., 2018) to ensure that adequate

effective sample sizes were reached (ESS >200). LogCombiner and TreeAnnotator were used to

generate the final maximum credibility tree with a 15% burn-in. Posterior probability (PP) values

≥0.95 were considered strong, and from 0.94 to 0.75 as moderate to weak.

The concatenated ML Sanger supermatrix analysis was run on a concatenated matrix, with

the same settings as described above in RaxML. The concatenated BI Sanger supermatrix was

analysed partitioning the dataset between ITS, *waxy* and the plastid genes. Modifications to the

analysis included a monophyletic constraint on *Solanum*, and four parallel runs that were run for

60 million generations with two chains, sampling trees every 10,000 generations. The ML best

tree was used as a starting topology to speed up convergence of the chains.

PL dataset– Paired reads from genome skimming were cleaned using BBDuk from the

BBTools suite (sourceforge.net/projects/bbmap/; ktrimright=t, k=27, hdist=1, edist=0, qtrim=rl,

trimq=20, minlength=36, trimbyoverlap=t, minoverlap=24, and qin=33). Sequence quality was

checked with FastQC (Andrews and Others, 2010) and MultiFastQC (Ewels et al., 2016). Plastome

assembly was done using de-novo assembly with Fast-Plast v.1.2.6

(https://github.com/mrmckain/Fast-Plast), and reference-guided assembly using GetOrganelle

v.1.6.2.e (Jin et al., 2020) with the high-coverage plastome sequence of *S. dulcamara* L. (GenBank

KY863443 (Amiryousefi et al., 2018). For GetOrganelle, the following settings were used: -w 0.6,

-R 20 -k 85, 95, 105, 127; for Fast-Plast, the Solanales Bow-tie index was used for the assembly.

Results from both methods were aligned in Geneious and visually checked to determine

consistency. Assembly quality was assessed using the reads identified from the Bow-tie step in the

Fast-Plast analysis, which were mapped against the final recovered plastome sequence using BWA

(Li and Durbin, 2010). Mean and standard deviation of coverage depth for each base pair was

determined by examining the same files in Geneious. Assemblies were annotated using both

Chlorobox GeSeq (Tillich et al., 2017) and the "Annotate from database" tool in Geneious using

the reference plastid genome of *S. dulcamara*. Results were compared to ensure that start and stop

codons for exon boundaries were congruent. Annotated plastomes were submitted to Genbank

(Appendix S2). A total of 55 full plastomes were assembled with a mean length of 155,498 bp

(max. 156,138 bp, min. 154,715 bp; Appendix S2), and a mean coverage of 158 (min. 22, max.

571; Appendix S2), and 28 partial plastomes (45,398-154,598 bp) with a mean coverage of 29

(min 4, max 96; Appendix S2). All plastomes had a highly conserved quadripartite structure, with

no loss, duplication, or expansion of gene families.

Plastomes from this study and those retrieved from Genbank were aligned in Geneious

using MAFFT (Katoh et al., 2005), visually checked, and corrected. A copy of the inverted repeat

(IRa) was removed prior to phylogenomic analyses, although 1,189 bp were kept at the beginning

of the region to be able to extract the gene that spans the boundary between the small single copy

(SSC) and IRa region. We then separated the plastome alignment into: (1) 79 protein-coding

regions, (2) 15 introns, (3) and 73 intergenic regions. For each dataset, the ambiguously aligned

regions and polyA repeats were removed, using visual checks for the exons and intron regions,

and the strict mode of trimAl (Capella-Gutiérrez et al., 2009) for the intergenic regions (Appendix

S8). Sequences shorter than 25% of the length of the aligned matrix for each region and columns

containing >75% of gaps were removed in trimAl (Capella-Gutiérrez et al., 2009) to avoid issues

with long branch attraction following Gardner et al. (2021). Two pseudogenes (*ycf*1 and *rps*19) at

the junction of IRa and Long Single Copy (LSC) (Amiryousefi et al., 2018), and four intergenic

regions with no parsimony informative characters were excluded from the final analysis. All

remaining loci alignments were concatenated together for the final PL phylogenetic analyses.

To test for the effect of missing data, two datasets were compared: a matrix with 151 taxa

containing all 140 species selected for this study with higher proportion of missing data (147,278

bp long with the second IR removed), and a matrix with 125 samples containing only complete

plastid sequences (Appendices S2 and S8).

ML searches were run on all PL datasets in IQ-TREE2 (Minh, Schmidt, et al., 2020) with

1,000 non-parametric bootstraps. Optimal substitution models were determined using –TEST in

IQ-TREE2 (Appendix S9). For both PL datasets, topologies from two different partitioning

schemes were also compared (unpartitioned vs. best-fit partition scheme based on PartitionFinder;

Lanfear et al., 2012) in IQ-TREE 2, to test if accounting for variation in substitution rate amongst

loci affected the phylogenetic results. BS values ≥95% were considered strong, 75-94% moderate,

and 60-74% weak.

TC dataset– Trimmomatic (Bolger et al., 2014) was used to trim reads (TruSeq3-PE-

simpleclip.fa:1:30:6, LEADING:30, TRAILING:30, SLIDINGWINDOW:4:30, MINLEN:36).

Read quality was checked with FastQC (Andrews and Others, 2010) and MultiFastQC (Ewels et

al., 2016). Over-represented repeat sequences were removed with CutAdapt (Martin, 2011).

HybPiper (Johnson et al., 2016) was used to produce reference-guided *de novo* assembles using

the reference provided by Johnson et al. (Johnson et al., 2019). Putative paralogs were identified

using the HybPiper script "paralog_retriever.py". Phylogenies were generated for all 45 loci for

which paralog warnings were found using MAFFT (Katoh et al., 2005) and FastTree (Price et al.,

2010). Five loci were deleted and several taxa whose paralogs caused paraphyly of clades were

excluded from 27 loci (one to seven taxa per loci). A single gene (g5299) presented a clear

duplication event and was divided into two separate matrices for downstream analyses.

Default HybPiper settings were used for all but three samples (*S. betaceum* Cav., *S.*

valdiviense Dunal, and *S. etuberosum* Lindl.), for which the coverage cutoff was reduced from

eight to four to maximise recovery of target genes. One sample (*S. terminale* Forssk.) was excluded

due to poor sequence quality. Only the exon dataset was analysed in downstream phylogenomic

analyses, because the transcriptome dataset showed large differences in the recovered flanking

regions of target loci between samples, likely due to post-transcriptional splicing and editing of

messenger RNA. The HybPiper script "fasta_merge.py" was used to concatenate all genes together

and produce a partition file. In summary, an average of 289 genes per sample were recovered for

the TC analysis (min 48, max 340) when the two samples with low numbers were excluded (*S.*

betaceum and *S. etuberosum*, Appendix S3). Furthermore, to reduce the effect of missing data and

long branch attraction, sequences shorter than 25% of the average length for the gene were

eliminated. The number of loci retained from the min04 and min20 datasets was 310 and 348

respectively, with the final aligned length varying between 242,272 bp and 261,975 bp (Appendix

S10).

The effect of missing data was tested by comparing two different sampling thresholds

based on the minimum number of taxa in each of the target genes alignments (min20 vs. min04,

i.e. a minimum of 20 taxa per gene and a minimum of four taxa per gene, respectively) using

HybPiper (Johnson et al., 2016) to retrieve and filter the genes.

ML analyses were run on both TC datasets in IQ-TREE2 (Minh et al., 2020) with

partitioning between loci. In addition, IQ-TREE2 was used to generate individual ML trees for

each loci, and the resulting phylogenetic trees were used for coalescent analyses with ASTRAL-

III v.5.7.3 (Appendix S9; Zhang et al., 2018), where tree nodes with <10% BS values were

collapsed using Newick Utilities v. 1.5.0 (Junier and Zdobnov, 2010). Trees with excessively long

branches were identified using phyx (Brown et al., 2017) by looking at tree lengths and root-to-tip

variation (command "pxlstr"); seven gene trees with excessively long branches were identified and

excluded for the min20 and ten for the min04 datasets, leading to a total of 303 and 338 gene trees

being used for the respective coalescent analyses. Branch support was assessed using local PP

support (Sayyari and Mirarab, 2016) calculated in ASTRAL, where PP values >0.95 were

considered strong, 0.75-0.94 weak to moderate, and ≤0.74 as unsupported.

Discordance analyses–

Comparison of resulting species trees– Topological congruence and discordances

between all 10 topologies generated were assessed visually by generating graphical

representations through custom R-scripts using the following packages: "ggtree" (Yu, 2020),

"stringr" (Wickham and Wickham, 2019), "ape"(Paradis and Schliep, 2019), "ggplot2"

(Villanueva and Chen, 2019) and "gridExtra"(Auguie and Antonov, 2017). To facilitate

comparisons, all trees were reduced to include the outgroup *Jaltomata* and 9 taxa representing

the following clades of *Solanum*, which were recovered in all analyses: Thelopodium,

Regmandra, Potato, Morelloid (as a representative of both the Dulcamaroid and Morelloid

clades), Archaesolanum , *S. anomalostemon* S.Knapp & M.Nee (species sister to Clade II),

Acanthophora (minor clade of the Leptostemonum) and two representatives of the EHS clade

(Table 1). The species sampled in the PL and TC datasets were identical for all except three

minor clades, in which different closely related species were sequenced (Acanthophora: *S.*

viarum Dunal/*S. capsicoides* All.; Morelloid: *S. opacum* A.Braun & C.D.Bouché/*S. americanum*

Mill.)

Concordance factors– Phylogenomic discordance was measured using gene concordance

factors (gCF) and site concordance factors (sCF) calculated in IQ-TREE 2 (Minh, Hahn, et al.,

2020). These metrics assess the proportion of gene trees that are concordant with different nodes

along the phylogenetic tree and the number of informative sites supporting alternative

topologies. Low gCF values can result from either limited information (i.e., short branches)

and/or genuine conflicting signal; low sCF values (~30%) indicate lack of phylogenetic

information in loci (Minh, Hahn, et al., 2020). The metrics were calculated using the TC

ASTRAL min20 topology (303 genes) and the PL IQ-Tree topology of 151 species

(unpartitioned) where sampling was reduced to 21 and 34 tips in TC and PL topologies,

respectively, retaining a single tip for each of the different minor and major clades. An additional

tip was retained for the Old Word Clade to visualize the gCF and sCF for the crown node of that

lineage.

Network analyses and polytomy tests– The presence of reticulate evolution and

conflicting signals in gene trees in the TC dataset was explored by generating a filtered supertree

network in SplitsTree 4 (Huson and Bryant, 2006) of the TC min20 dataset (303 genes)

collapsing branches with <75% local PP support with a minimum number of trees set to 50%

(151 trees). Polytomy tests were carried out in ASTRAL-III (Sayyari and Mirarab, 2018) using

ASTRAL topologies of the two datasets (min20 and min04). Gene trees were used to infer

quartet frequencies for all branches to determine the presence of polytomies while accounting for

ILS. The analysis was run twice to minimize gene tree error.

RESULTS

Phylogenetic analyses–

Congruent recovery of major clades– All three datasets, including the supermatrix and

the two phylogenomic datasets (PL and TC), recovered previously recognized major clades in

Solanum (Fig. 1-2a,c); a few minor clades, concentrated in Clade II, were found to be

polyphyletic in the supermatrix phylogeny, including the Mapiriense-Clandestinum,

Sisymbriifolium, Wendlandii-Allophyllum and Cyphomandropsis minor clades (Appendices

S11-S12); comparison with PL and TC phylogenies is not possible, as only one species of each

clade were sampled in these datasets. In Clade I, nearly all specimens of the Dulcamaroid clade

formed a monophyletic group. The only exception concerned *S. alphonsei* Dunal, sampled here

for the first time. In both the supermatrix and PL analyses, this species was sister to *S.*

valdviviense of the Valdiviense clade, with maximum branch support in the PL analyses (Fig 2.,

Appendix S13).

Despite these minor novelties, all analyses recovered the Thelopodium clade as sister to

the rest of *Solanum* (Fig. 1-2, Appendices S11-15). The Potato clade was strongly supported

across all analyses (Fig. 1-2, Appendices S11-S15), as was the Regmandra clade in supermatrix

and PL analyses (only one sample in TC phylogenies). Furthermore, all analyses recovered a

clade here referred to as DulMo that includes the Morelloid and Dulcamaroid clades (Figs. 1-2,

Appendices S11-S15). A new strongly supported clade, here referred to as VANAns clade and

comprising the Valdiviense (including *S. alphonsei*, see below), Archaesolanum, Normania, and

the African non-spiny clades, was found across all analyses (Figs. 1-2; Appendices S11-S15).

Clade II was supported as monophyletic across all topologies (Fig. 1-2 a,c), with

maximum branch support in all 10 species trees (Appendices S11-S15).While differences in

sampling prevent thorough comparisons of relationships between clades within Clade II, there

was no deep incongruences detected amongst topologies obtained with the supermatrix, PL, and

TC datasets (Fig. 1-2 a,c; Appendices S9-S15). Within Clade II, the large Leptostemonum clade

(the spiny solanums) was strongly supported in all cases (Fig. 1-2 a,c; Appendices S11-S15).

Incongruent relationships amongst clades and impact of different analyses– Overall, we

found that despite using different phylogenetic analyses and investigating the impact of missing

data and taxon sampling on the different datasets, these had little impact on the relationships

recovered amongst clades. The BI and ML supermatrix analyses were identical in terms of

composition and relationships of major clades (Fig. 3b), as were the four PL species trees (Fig. 3

d,e). There were some differences amongst the topologies of the TC datasets, but these

differences concerned branches which had little support (Fig. 3a,c,b). Between supermatrix, PL

and TC datasets, however, major incongruences between species trees were observed with

respect to the relationships among the main clades identified in the section above (Fig. 1-3).

While the BI and ML supermatrix phylogeny supported the monophyly of the previously

recognised Clade I that includes most non-spiny *Solanum* clades (Fig. 1, Appendices S11-12),

the PL and TC phylogenetic trees resolved clades associated with Clade I as a grade relative to

Clade II (Fig. 2a,c, Appendices S13-S15). This was in large part due to the unstable position of

the Regmandra clade that was subtended by a particularly short branch and resolved in different

positions along the backbone in all three datasets (Fig. 3). For example, the ML supermatrix

analysis recovered the Regmandra clade as sister to the Potato clade with strong to moderate

branch support (Fig. 3b), although the BI supermatrix analysis could not resolve whether the

Regmandra clade was sister DulMo+VANAns clade or the Potato clade (Fig. 3b, Appendix S12).

In contrast, the PL analyses resolved Regmandra as sister to the M clade + Clade II, with either

maximal or no branch support at all (Fig. 3). The TC species trees resolved Regmandra as sister

to the Potato clade, DulMo, and Clade II, with maximum support (Fig. 3). While one of the TC

ASTRAL analysis also recovered this topology with moderate support (local posterior

probability 0.82, Fig. 3), the other TC ASTRAL analysis resolved Regmandra as sister to the

VANAns clade, but without any branch support (local PP 0.4, Fig. 3).

The previously identified M Clade composed of the VANAns and DulMo clades were

not supported by all analyses (Fig. 3). While all PL ML analyses recovered the M clade with

maximum BS values (Fig. 3), none of the TC analyses recovered it. Instead, they resolved the

DulMo clade as sister to the Potato clade, with maximal BS or local PP support values (Fig. 3.)

Furthermore, the VANAns clade was recovered as sister to the rest of *Solanum* (excluding the

Thelopodium clade) with moderate support in the TC ML analyses. Placement of the VANAns

clade in the TC-ASTRAL-III analyses had low or no support value, being resolved as either

sister to DulMo, or sister to the rest of *Solanum,* excluding the Thelopodium clade (Fig. 3).

In addition, the position of the Potato clade within *Solanum* was incongruent between

datasets: whereas it was resolved as sister to Regmandra in the supermatrix analysis, it was

resolved as sister to the remaining *Solanum* in PL dataset, and sister to the DulMo clade in all TC

analyses (Fig. 3), all with strong branch support. The phylogenomic datasets also showed

incongruent positions for the Etuberosum clade within the larger Potato clade, where TC

analyses resolved it as sister to the Petota clade with maximum local PP support in the ASTRAL

analyses (Appendix S15); in the ML analyses, this position either had moderate BS values (76%)

or was found to be nested within the Petota clade with no branch support (Appendix S14). In

contrast, PL analyses placed Etuberosum clade as sister to the Tomato clade with maximum

branch support (Appendix S13).

Finally, the BI and ML supermatrix phylogenies resolved the morphologically unusual *S.*

anomalostemon as sister to the rest of Clade II (BS 95%, PP 1.0; Fig. 3, Appendices S11-S12).

This contrasts with results from previous analyses, which found it to be part of the Mapiriense

clade (Särkinen et al., 2015). PL analyses supported *S. anomalostemon* + Brevantherum clade as

sister to the rest of Clade II with high branch support (Appendix S13). *Solanum anomalostemon*

was also found to be sister to Clade II, although the Brevantherum clade was not included in the

TC analyses preventing a strict comparison (Fig. 3). Two other taxa were found to represent

single species lineage: *S. polygamum* as sister to the Leptostemonum clade and *S. euacanthum* as

sister to the EHS clade (Appendices S11-S12). Within the Leptostemonum clade, the EHS clade

was strongly supported in all analyses (Figs.1-3). There were however some minor differences in

species-level relationships for closely related species of the Eggplant clade and Anguivi Grade

(viz. *S. campylacanthum* Hochst. ex A.Rich., *S. melongena* L., *S. linnaeanum* Hepper & P.-

M.LJaeger, *S. dasyphyllum* Schum. & Thonn. and *S. aethiopicum* L.; Fig. 1-2a, c, Appendices

S11-S15).

Discordance analyses–

Concordance factors– Phylogenomic discordance was generally high across the PL and

TC topologies, with gCF values >50% in only three nodes in the PL phylogeny (*Solanum* as a

whole, *S. chilense* (Dunal) Reiche + *S. lycopersicum* L. or the Tomato clade, and *S. hieronymi*

Kuntze + *S. aridum* Morong in the Leptostemonum clade; Fig. 4). Elsewhere, along the

backbone of the PL phylogeny, gCF fell to 39% and below (8 nodes with gCF values 10% and

below), with the lowest values found near branch nodes that varied the most amongst the

different reconstructed species trees. This included the node subtending Regmandra (gCF 4%,

SCF 38%; Fig. 4), and that positioning Regmandra + DulMo + VANAns clade as sister to Clade

II (gCF 2%, SCF 31%). Similarly, low gCF and uninformative sCF values around 33% were

found across Clade II, including the node placing *S. hieronymi* + *S. aridum* as sister to the

Elaeagnifolium + Old World Minor clades (gCF 6 %, sCF 36 %; Fig. 4), as well as the

placement of the Erythrotrichum + Thomasiifolium clades within the large Leptostemonum clade

(gCF 5%, sCF 23%; Fig. 4).

Across the TC phylogeny, gCF and sCF values were slightly higher on average, with 3

nodes presenting values >50% for both metrics: one within the Petota clade (gCF 67%, SCF

69%; Fig. 4), one at the base of the Leptostemonum clade (gCF 64%, SCF 72%; Fig. 4), and

another at the base of the EHS clade within Leptostemonum (gCF 58%, SCF 75%; Fig. 4). Three

nodes had low gCF values of 10% or less, with again some of the lowest values located near the

base of the tree, including the relationship of Regmandra as sister to the VANAns clade (gCF

3%, sCF 39%; Fig. 4), or placement of Potato as sister to the DulMo clade (gCF 10%, sCF 41%;

Fig. 4), and the relationship of the Potato + DulMo clades as sister to Clade II (gCF 4%, sCF

41%; Fig. 4).

Network analyses and polytomy tests– High amount of reticulation/gene tree conflict was

recovered between major clades of *Solanum* previously assigned to Clade I (e.g., Thelopodium,

Regmandra, Potato, DulMo, VANAns), as well as with some lineage belonging to Clade II in the

filtered supertree network using the TC data with 303 genes (min20; Fig. 2b). The network

clearly supported the monophyly of the Leptostemonum and the EHS clade (Fig. 2b),

corresponding to the nodes with high gCF and sCF values in the TC ASTRAL phylogeny (N1

and N2, Fig. 4).

The polytomy tests carried out for the two TC ASTRAL datasets resulted in 10 nodes

each for which the null hypothesis of branch lengths equal to zero was accepted, suggesting they

should be collapsed into polytomies (Appendix S16); these nodes corresponded to the ones

subtending the Regmandra, Leptostemonum and EHS clades, but were also located within the

VANAns clade as well as within Clade II, the. Polytomies were also detected with the Petota

clade, including at the base of the Tomato clade (min4 dataset, Appendix S16), and at the base of

the Etuberosum + Petota +Tomato clade (min20 dataset, Appendix S16). Repeating the analysis

by collapsing nodes with <75% local PP support led to the collapse of 12 to 13 nodes across the

analyses, most of them affecting the same clades as in the previous runs, but also leading to the

collapse of the crown node of *Solanum*. The effective number of gene trees was too low when

nodes with <75% local PP support were collapsed to carry out the test for two nodes subtending

S. betaceum and *S. anomalostemon*, most likely related to the low number of genes recovered for

S. betaceum (Appendix S3).

DISCUSSION

The results of the ten phylogenetic analyses conducted here provide an updated evolutionary framework for the large and economically important genus *Solanum,* demonstrating that the major and minor clades within the group are stable (with a few noteworthy exceptions, see below). However, the strong levels of nuclear and nuclear-plastome discordance uncovered in the PL and TC analyses, in combination with the network analysis and polytomy tests, suggest that there are polytomies present along the backbone of the phylogeny. We first discuss the stability of the clades within *Solanum*, and the discovery of a few novel minor clades. We then examine the nuclear-plastome discordance and polytomies recovered and explore the possible causes underlying these, and their implications for the study of biogeography and trait evolution.

Updated evolutionary framework for *Solanum*

The supermatrix phylogeny, despite being based on only nine loci, nearly doubles the species sampling, confirming the monophyly of most major and minor clades established in previous analyses (Särkinen et al. 2013) and the polyphyly of three minor clades (Pachyphylla, Cyphomandropsis, and Allophyllum, the latter including species of Mapiriense-Clandestinum clade). It also reveals three new minor clades in *Solanum* comprising a single species each and confirms the placement of 129 previously unsampled species (e.g., *S. alphonsei* in the Valdiviense clade and *S. graveolens* in the Cyphomandra clade; Appendices S11-S12). Meanwhile, the phylogenomic analyses with increased gene sampling reveal a previously undetected major clade referred to as VANAns comprising of four minor clades (Valdiviense, Archaesolanum, Normania, and African non-spiny clades). Finally, our results did not support two previously resolved major clades due to nuclear-plastome discordance (Clade I and the M clade; Fig. 2). Detailed molecular systematic studies with increased taxon and genetic sampling

will be required to fully resolve the circumscription of all the major and minor clades recovered

with diagnostic features, including the new ones identified here (Hilgenhof et al. in prep).

Overall, our results establish that the taxonomic framework used in *Solanum* dividing the

large genus into major and minor clades is robust, based on both phylogenomic datasets

recovering the same major clades independent of methodological choices compared to the

Sanger sequence supermatrix (e.g., Thelopodium, Regmandra, Potato, DulMo, VANAns, Clade

II, Leptostemonum, and EHS clade). The major and minor clades currently used as informal

infrageneric groups in *Solanum* were first established by Bohs (2005) based on a single locus of

c. 2,000 bp in length (*ndhF*). Our results demonstrate that larger species and gene sampling

support the clades established nearly two decades ago (e.g. Weese and Bohs, 2007; Särkinen et

al., 2013). However, increased gene sampling provided by the two phylogenomic datasets does

not help to resolve any of the polytomies along the backbone of *Solanum* close to the crown node

and along the backbone of Clade II (Särkinen et al., 2013).

Nuclear & nuclear-plastome discordance

Our results reveal three regions of the *Solanum* phylogeny with gene discordance with

low gCF and sCF values in the PL and TC dataset (Fig. 4). These regions with nuclear

discordance include: (1) the backbone of *Solanum* near the crown node of the genus where major

clades previously identified as Clade I diverge (from here on referred to as Grade I); (2) the

backbone of the large Leptostemonum clade; and (3) the backbone of the EHS clade within the

Leptostemonum (Figs. 2b and 3). Many of the branches within these regions are extremely short

in both PL and TC phylogenomic datasets (Fig. 1-2, Appendices S11-S15), and network analyses

of the nuclear dataset reveals reticulation in one of them (Grade I, Fig. 2b). Polytomy tests

confirm that multiple nodes within all three regions should be collapsed in the TC dataset

(Appendix S16) and support the recognition of these regions as polytomies. Hence, we refer to

these three regions of the phylogeny as polytomies from hereon.

Further exploration of the polytomies reveal nuclear-plastome discordance within Grade

I, relating to the position and relationship between Regmandra, Potato, DulMo and VANAns

clades (Fig. 3-4). No signal of nuclear-plastome discordance was detected in the other

polytomies based on the species sampling presented here (Fig. 3-4), but increased species

sampling will be needed to confirm these results.

Altogether, our results indicate the presence of three polytomies which differ somewhat

in nature. The deepest of these polytomies along the backbone of *Solanum* near the crown node

shows high nuclear and nuclear-plastid discordance with reticulation evident even within the

nuclear phylogenomic dataset (Fig. 2b). This polytomy could be referred to as a hard polytomy,

because it will probably be difficult to resolve even with more genomic data, due to its deeper

position in the phylogeny in terms of evolutionary depth and time, the presence of clear nuclear-

plastome discordance, short branch lengths and evidence for reticulation within the nuclear

phylogenomic dataset. In contrast, the other two polytomies along the backbone of

Leptostemonum and the EHS clades are at shallower evolutionary depth and show nuclear

discordance only without clear/widespread reticulation in the nuclear dataset (Fig. 2b). These

polytomies represent simpler cases and may turn out to be possible to resolve with more genomic

data. In either case, to confirm whether the polytomies recovered here are truly "hard" or "soft",

denser taxon sampling and more genomic data will be required to carry out more rigorous tests

concerning the cause of the gene discordance observed here.

What is causing genomic discordance in our dataset?

Finding genomic discordance in our phylogenomic datasets is unsurprising, given that it

has also been found in many other phylogenomic studies in the Solanaceae, including *Nicotiana*

(Dodsworth et al., 2020), the Capsiceae (*Capsicum* and relatives; Spalink et al., 2018), subtribe

Iochrominae (Gates et al., 2018), *Jaltomata* (Wu et al., 2019)*,* and two studies of *Solanum*

involving the Tomato (Strickler et al., 2015; Pease et al., 2016) and Petota clades (Huang et al.,

2019). ILS was shown to be responsible for the widespread discordance found in phylogenomic

data in the diploid Tomato clade (Strickler et al., 2015; Pease et al., 2016), while hybridization

and introgression has been argued to be behind genomic discordance in Petota clade that includes

many polyploids (Huang et al., 2019).

Potential processes responsible for nuclear or nuclear-plastome discordance involve gene

introgression, ILS, hybridization, and polyploidisation; distinguishing between these remains

difficult even with increased genomic sampling involving custom bait sets (Larridon et al., 2020;

Koenen et al., 2021) or whole genome-sequences (Suh, 2016; Malinsky et al., 2018; Williams et

al., 2021). Comparison of the nuclear and plastome topologies in our study does not indicate any

obvious chloroplast capture events that could explain the observed nuclear-plastome discordance

along the backbone of *Solanum* near the crown node. Furthermore, cytogenetic and chromosome

studies show no evidence for genome duplication or polyploidy along the three polytomies

discovered here, despite the three-fold increase in genome size between the distantly related

potato (*S. tuberosum* L., Potato clade) and eggplant (*S. melongena,* Leptostemonum clade;

Barchi et al., 2019). Chromosome counts indicate that the ancestor of *Solanum* was diploid: a

large majority of *Solanum* species are reported to be diploid (>97% of the 506 species for which

chromosome counts are available), and mapping of ploidy level across the phylogeny indicates

that most of the lineages involved in the three polytomy regions identified here are diploid

(Chiarini et al., 2018). Polyploidy has arisen independently within the Archaesolanum, Petota,

Morelloid, Caroliniense, Elaeagnifolium and EHS minor clades within the larger

Leptostemonum clade (Chiarini et al., 2018), and hybridization/introgression has been argued to

be the case behind phylogenomic discordance found in the Petota clade (Huang et al., 2019).

Gene duplication could explain the signal recovered here for the EHS clade but is unlikely to

explain the discordance observed here. Save for one locus, our analyses did not detect the

presence of paralogs in our nuclear dataset.

Currently, the most likely explanation for the discordance along the backbone of *Solanum*

is due to ILS caused by rapid speciation. Two of the polytomies include the most species-rich

(Table 1) and rapidly diversifying lineages of *Solanum*, the Leptostemonum and the EHS clades

(Echeverría‑Londoño et al., 2020), whose crown ages have been estimated to be between 8-11

and 4-6 million years (Myr), respectively (Särkinen et al., 2013). The backbone of *Solanum* near

the crown node has been estimated to be almost twice as old as the Leptostemonum clade (13-17

Myr; Särkinen et al. 2013) yet shows a strong signal of nuclear-plastome discordance. While past

studies have not detected any increased rates of diversification near the crown node of *Solanum*,

detecting diversification rate shifts remains a challenge (Louca and Pennell, 2020), especially in

older nodes. Hence, we cannot fully exclude the option that ILS and rapid speciation has taken

place close to the crown node of the genus.

Presence of short internal branches is typical of ILS in lineages with large population

sizes and high mutation rates (Schrempf and Szöllősi, 2020). This fits with the biology of

Solanum in general, which is typically known to contain "weedy", disturbance-loving pioneer

species resilient to change. Many species are known to have large geographical ranges and

ecological amplitude, including globally distributed weeds from the Leptostemonum,

Brevantherun and Morelloid clades, such as *S. elaeagnifolium, S. caroliniense*, *S. torvum*, *S.*

erianthum, S. mauritianum, S. americanum and *S. nigrum* (Knapp et al., 2017, 2019; Cowie et

al., 2018; Särkinen et al., 2018). Some of the weedy characteristics found in these species include

the ability to improve fitness and defense traits in response to disturbance (Chavana et al., 2021),

as well as having allelopathic properties which allow them to establish themselves to the

detriment of native vegetation (Cowie et al., 2018). If such characteristics were present in

ancestral *Solanum*, they could have promoted rapid speciation across the globe, followed by

rapid morphological evolution and speciation within areas. The patterns observed here could

possibly be the result of three major rapid speciation "pulses" across the evolutionary history of

Solanum, involving lineages close to the crown node of *Solanum*, Leptostemonum, and the EHS

clade. The idea of an ecologically opportunistic ancestor is supported by the tendency of many of

the major clades near the crown node of *Solanum* to occupy periodically highly stressed and

disturbed habitats, including flooded varzea forests occupied by Thelopodium clade, hyper-arid

deserts occupied by Regmandra clade, and highly disturbed and dynamic open mid-elevation

Andean montane habitats occupied by DulMo clade, where landslides are among the most

common areas where many of the species are found (Knapp, 2013; Särkinen et al., 2018; Knapp

et al., 2019).

Future studies with larger datasets will be able to carry out additional tests, such as the

impact of using phylogenetic models that take into consideration the heterogeneity of molecular

sequence evolution (Williams et al., 2021), as well as different data types (Romiguier et al.,

2013; Reddy et al., 2017). Future studies will need to untangle how introgression and ILS are

potentially affecting the patterns of genomic discordance observed here at different phylogenetic

depths (Meleshko et al., 2021). Additional information about recombination, chromosome

structure, and genomic size and evolution of *Solanum* will also be useful to clearly define

coalescence genes in phylogenomic datasets, fundamental units in coalescent analyses which are

rarely examined (Springer and Gatesy, 2018). Currently, information about genome evolution in

Solanum is lacking, as only 62 species (5% of *Solanum*) are recorded in the plant DNA C-value

database (Pellicer and Leitch, 2020), and 86 species (7% of *Solanum)* have been studied with

chromosome banding and/or FISH techniques (Chiarini et al., 2018). Information about genome

size is missing for lineages such as the Thelopodium and Regmandra clades and for the majority

of species not directly related to major commercial crops.

***Implications for biogeographical and morphological studies in* Solanum**

The idea that well-supported and fully bifurcating phylogenies are a requisite for

evolutionary studies is built on the premise that such trees are the accurate way of representing

evolution. The shift in systematics from "tree"- to "bush"-like thinking, where polytomies and

reticulate patterns of evolution are considered as acceptable or real (Poczai, 2013; Mallet et al.,

2016; Edelman et al., 2019), comes from the accumulation of studies finding similar

unresolvable phylogenetic nodes, despite using different large-scale genomic sampling strategies

and various analytical methods (Suh, 2016). Given the difficulty of resolving short internal

branches in phylogenies and the rapid evolution of major clades in *Solanum,* it will be important

to adopt methods that incorporate polytomies and networks to conduct biogeographical and

morphological studies (Than et al., 2008; Solís-Lemus et al., 2017; Wen et al., 2018; Olave and

Meyer, 2020; Lutteropp et al., 2021).

In terms of biogeography, our inability to resolve relationships amongst the major

lineages in *Solanum*, especially along the backbone of *Solanum* near the crown node, has

implications for understanding the ancestral environment of *Solanum* and its major lineages.

Uncertainty amongst the relationships of major clades does not change the hypothesis that the

genus probably originated from South America and spread multiple times to Africa, Asia,

Australia, North America, and Europe (Olmstead and Palmer, 1997; Echeverría‐Londoño et al.,

2020). The polytomy near the crown node of *Solanum* does, however, cast uncertainty on the

specific region and habitat/biome that the major clades originated within the South American

continent. For example, the sister relationship of Regmandra and the Potato clade inferred by the

Sanger supermatrix analysis suggests that the wild ancestors of both potato and tomato evolved

from an ancestor adapted to survive in lomas deserts from coastal South America (Bennett, 2008;

Fig. 1). Yet, both nuclear and plastome phylogenomic datasets suggest that the Potato clade is

more closely related to the DulMo clade found to occur in tropical montane and subtropical

biomes (Fig. 3)

The hard polytomy along the backbone of *Solanum* also has important implications for

evolutionary biologists interested in trait evolution. Standard methods of trait evolution relying

on bifurcating trees may incorrectly infer how traits evolve (Hahn and Nakhleh, 2016). The

discordance between traits, gene trees, and species trees has been defined as hemiplasy (Avise

and Robinson, 2008), and studies have shown that depending on the level of ILS present in the

data, hemiplasy can lead to different interpretations of convergent evolution of traits across

phylogenetic trees (Mendes et al., 2016). While broad mapping of morphological traits on a

species-level phylogeny can help gain a rough understanding of phenotypic variation across

clades, careful study of gene tree topologies in relation to a trait of interest is essential to gain an

exact understanding of its evolutionary origin.

Our findings reflect results from recently published studies showing rapid morphological

innovation coinciding with areas of strong phylogenomic discordance in different plants and

animal groups (Parins-Fukuchi et al., 2021), where the signal of nuclear-plastome discordance

corresponds to strong ecological diversification and morphological innovation across major

clades in *Solanum* previously assigned to Clade I. The major clades involved in the nuclear-

plastome discordance along Grade I show large differences in their ecology as well as

morphology. Members of the Thelopodium, Regmandra, VANAns, Potato, and DulMo clades

occupy a wide range of tropical, montane and temperate habitats across South America, Africa

and Australia (Symon, 1994; Knapp, 2000; Bohs and Olmstead, 2001; Spooner et al., 2004,

2016, 2019; Bohs, 2005; Peralta et al., 2007; Bennett, 2008; Knapp, 2013; Knapp and

Vorontsova, 2016; Tepe et al., 2016; Särkinen et al., 2018; Knapp et al., 2019). Morphology

shows equally high polymorphism between these major clades across many traits, such as growth

form, which varies from single-stemmed wand-like shrubs (Thelopodium clade), annual herbs

(Regmandra, Potato, and Morelloid clade), woody climbers and shrubs (VANAns clade), and

herbaceous vines rooting along nodes (Potato clade). Similar patterns are observed in

inflorescence position and branching, corolla shape, stamen dimorphism, and anther shape

showing the presence of high polymorphism in these clades of which only some was retained in

Clade II (Hilgenhof et al. in prep). Testing the idea that this phenotypic diversity is linked to

ecological diversification will require the construction of detailed morphological and ecological

datasets to test if this pattern holds up in more formal and rigorous analyses.

CONCLUSION

We demonstrate the stability of the majority of the clades defined within *Solanum* and

uncover significant nuclear and nuclear-plastome discordance amongst relationships of major

clades in *Solanum* based on the first phylogenomic study of the genus with wide species

sampling. Three major polytomies are identified in *Solanum* based on the short branch lengths,

gene concordance factor results, and polytomy tests. Two of these polytomies correspond to the

biggest and most quickly diversifying lineages within *Solanum* (Leptostemonum and EHS

clades). The third polytomy along the backbone of *Solanum* near the crown node involves

reticulation and strong nuclear-plastome discordance and highlights great uncertainty in the

relationships between the Potato, DulMo, Regmandra, and VANAns clades. This region of

nuclear-plastome discordance corresponds with high ecological and morphological innovation

and we argue that it is most likely due to ILS and rapid speciation based on current knowledge of

genome evolution in *Solanum*. Future studies, even with full genome sequences and increased

taxon sampling, might not be able to resolve the polytomy near the crown node of *Solanum*

because the pattern of high reticulation combined with internodal short branches and it's older

age. Data on genome size and chromosome structure of the earliest branching lineages in

Solanum will be required to further explore the nature and causes of this hard polytomy. We

argue that acknowledging and embracing polytomies and reticulation is crucial if we are to

design research programs aimed at understanding the biology of large and rapidly radiating

lineages, such as the large and economically important *Solanum*.

Acknowledgements and funding

This work was supported by the Fonds de recherche du Québec en Nature et

Technologies postdoctoral fellowship and a grant from the Department of Biological Sciences of

the University of Moncton to EG, the Sibbald Trust fellowship to RH, the Ceiba Foundation to

AO, CNPq Conselho Nacional de Desenvolvimento Científico e Tecnológico awards

479921/2010-54 and 427198/2016-0 and Coordenação de Aperfeiçoamento de Pessoal de Nível

Superior CAPES/FAPESPA award 88881.159124/2017-01 to LLG, NSF through grant DEB-

0316614 "PBI Solanum: a worldwide treatment" to SK and LB, the Calleva Foundation &

Sackler Trust (Plant and Fungal Trees of Life Project at Kew), the LUOMUS Trigger and

Systematics Research Fund to PP, the OECD CRP and Eötvös Research Grant (MAEÖ-00074-

002/2021). Field sampling was supported by the Northern Territory Herbarium and the David

Burpee Endowment at Bucknell University (Australia), and National Geographic Society

Northern Europe Award GEFNE49-12 (Peru, TS). Peruvian specimens were collected and

sequenced under the permission of Ministerio de Agricultura, Dirección General Forestal y de

Fauna Silvestre (collection permits 084-2012-AG-DGFFSDGEFFS and 096-2017-

SERFOR/DGGSPFFS, and genetic resource permit 008-2014-MINAGRI-DGFFS/DGEFFS).

We thank Elliot Gardner for sharing scripts and advice on phylogenomic analyses with

HybPiper, Royce Steeves for providing advice on DNA extraction for genome skimming, Felix

Forest and Olivier Maurin for providing technical support and providing feedback on the

manuscript, and João R. Stehmann, Thais Almeida, Paul Gonzáles, and Maria Baden who greatly

contributed to fieldwork and sample acquisition. Finally, we would also like to thank the three

reviewers, including Stacey Smith and William J. Baker, who provided constructive reviews and

feedback that greatly improved the final version of this manuscript.

Author contributions

EG designed and performed the analyses of the paper, with guidance from PP, AO, SD

and TS; EG produced all figures, and wrote the article, with major contributions from TS, and

PP, SD, SK and XA. RH and TS helped in data gathering and analyses. All other authors

contributed data to the main analyses. All authors read and contributed to the final version of the

manuscript.

Data Availability Statement

Raw sequence data generated in this study are deposited in various archives, including Genbank

(https://www.ncbi.nlm.nih.gov/genbank/) and the European Nucleotide Archive

(https://www.ebi.ac.uk/ena/browser/home); full accession numbers are provided in Appendices

S1, S2 and S3. In addition, the 10 species trees generated for this study, as well as the alignments

used for the different phylogenetic analyses, including the concatenated Sanger supermatrix, the

plastome dataset and and the target capture datasets (min 04 and min20) are available via Data

Dryad, at the following link: (to be provided upon acceptance for publication).

Supporting Information

Additional supporting information may be found online in the Supporting Information

section at the end of the article:

Appendix S1. Supermatrix sample information, including voucher details and Genbank numbers

for sequences used.

Appendix S2. Plastome (PL) sample information, including voucher details and plastome

assemblies' results. Total length, as well as length for the long-single copy region (LSC), the

short-single copy region (SSC) and the two inverted repeat regions (IR1 and IR2) is shown;

statistics of mean coverage per base pair and standard deviation are also provided.

Appendix S3. Target-capture (TC) sample information, including voucher details and sequence

recovery statistics. The number of reads (NumReads), the number of reads mapped to the targets

(ReadsMapped), the percentage of reads on target (PctOnTarget), the number of genes with reads

(GenesMapped), the number of genes with contigs (GenesWithContigs), (GenesWithSeqs,

GenesAt25pct, GenesAt50pct, GenesAt75pct, GenesAt150pct, and the number of genes with

paralog warnings (ParalogWarnings) is shown.

Appendix S4. Supermatrix alignment details, with details about the nine regions selected for this

study. Number of species sampled per region, accumulative percentage of species sampled per

region, aligned length, proportions of parsimony informative characters (PI), and variable sites

(VS) per region in the dataset are indicated. Values are calculated with outgroups, and with

ambiguous regions and repeats excluded. bp=base pairs.

Appendix S5. List of polyploid taxa in *Solanum*.

Appendix S6. ML results for each of the nine individual loci and combined plastid loci. (a) ITS ;

(b) matK; (c) ndhF; (d) ndhF-rpL32; (e) psbA-trnH; (f) rpL32-trnL; (g) trnL-trnT; (h) trnS-trnG;

(i) waxy; (j) seven plastid loci. Nodes with bootstrap support equal and above 95% are in cyan,

and with branch support between 75% and 94% in red. Tips indicate species names, followed by

major and/or minor clade, as indicated in Table 1.

Appendix S7. BI results for each of the nine individual loci and combined plastid loci. (A) ITS ;

(B) matK; (C) ndhF, (D) ndhF-rpL32, (E) psbA-trnH (F) rpL32-trnL; (G) trnL-trnT; (H) trnS-

trnG; (I) waxy; (J) seven plastid loci. Nodes with posterior probability equal and above 0.95 are

in cyan, and nodes with posterior probabilities between 0.75 and 0.95 are in red. Tips indicate

species names, followed by major and/or minor clade, as indicated in Table 1.

Appendix S8. Plastome (PL) alignment statistics for plastome alignment. Data shows number of

sequences, trimming mode, the number of loci retained for coalescent analysis after checking for

excessive gene tree branch lengths, alignment length, number of informative and constant sites,

pairwise identity, average GC content, percentage of gaps, and average locus length for the exon,

intron and intergenic regions.

Appendix S9. Optimal substitution model used in ML analyses for the PL and TC datasets,

determined using ModelFinder in IQ-TREE2. For each locus, the number of taxa, sites,

informative sites, and invariable sites are indicated, as well as the model selected and the AICc

score. Worksheet titles correspond to the following: PLUnpartitioned = Models selected for PL

unpartitioned datasets, for 151 taxa and 125 taxa; PLBestPartScheme = Models selected for PL

datasets analysed according to the best-partition scheme; TCPartitioned_Min4= Models selected

for loci of the TC dataset, with minimum 4 taxa per loci; TCPartitioned_Min20= Models

selected for loci of the TC dataset, with minimum 20 taxa per loci.

Appendix S10. Target-capture (TC) alignment statistics. Loci excluded refer to the number of

excluded loci based on excessively long branch lengths, and loci retained is the final number of

loci retained for both ML and coalescent analyses. Empty sequences inserted refers to amount of

missing data. Min = minimum; Bp = base pairs. Appendix S11. Detailed RaxML of supermatrix

phylogenetic tree with 746 taxa. Nodes with bootstrap support equal and above 95% are in cyan,

and with branch support between 75% and 94% in red. Bootstrap support values for each node

indicated in italic. Tips indicate species names, followed by major and/or minor clade, as

indicated in Table 1.

Appendix S12. Detailed Bayesian inference (Beast) supermatrix phylogenetic tree with 746 taxa.

Nodes with posterior probability equal and above 0.95 are in cyan, and nodes with posterior

probabilities between 0.75 and 0.95 are in red. Posterior probability values for each indicated in

italic. Tips indicate species names, followed by major and/or minor clade, as indicated in Table

1.

Appendix S13. ML phylogenetic trees of plastome datasets. Nodes with bootstrap support equal

and above 95% are in cyan, and with branch support between 75% and 94% in red. Tips indicate

species names, followed by major and/or minor clade, as indicated in Table 1 a) 151 taxa, all

data, unpartitioned; b) 125 taxa, all data, unpartitioned; c) 151 taxa, all data, best partition

scheme; d) 125 taxa, all data, best partition scheme.

Appendix S14. ML phylogenetic trees of A353 target capture datasets (IQ-TREE2). Nodes with

bootstrap support equal and above 95% are in cyan, and with branch support between 75% and

94% in red. Tips indicate species names, followed by major and/or minor clade, as indicated in

Table 1. a) filtering threshold of a minimum of 4 taxa per loci; b) filtering threshold of a

minimum of 20 taxa per loci.

Appendix S15. Coalescent phylogenetic trees of A353 target-capture datasets (ASTRAL-III).

Nodes with multi-locus local posterior probability support equal and above 0.95 are in cyan, and

with support between 0.75 and 0.94 in red. Tips indicate species names, followed by major

and/or minor clade, as indicated in Table 1. a) filtering threshold of minimum of 4 taxa per loci;

b) filtering threshold of minimum of 20 taxa per loci.

Appendix S16. Polytomy test results with ASTRAL-III. a) Target Capture A353 species tree

ASTRAL-III, filtering threshold of minimum 4 taxa per loci, branches in gene trees with 10% or

less branch support collapsed; b) Target Capture A353, ASTRAL-III, filtering threshold of

minimum 4 taxa per loci, branches in gene trees with 75% or less branch support collapsed; c)

Target Capture A353, ASTRAL-III, filtering threshold of minimum 20 taxa per loci, branches in

gene trees with 10% or less branch support collapsed; d) Target Capture A353, ASTRAL-III,

filtering threshold of minimum 20 taxa per loci, branches in gene trees with 75% or less branch

support collapsed;

References

References

Aberer, A. J., D. Krompass, and A. Stamatakis. 2013. Pruning rogue taxa improves phylogenetic
accuracy: An efficient algorithm and webservice. *Systematic biology* 62: 162–166.
Amiryousefi, A., J. Hyvönen, and P. Poczai. 2018. The chloroplast genome sequence of
bittersweet (Solanum dulcamara): Plastid genome structure evolution in Solanaceae. *PloS*
one 13: e0196069.
Andrews, S., and Others. 2010. FastQC: a quality control tool for high throughput sequence data.
APG. 1998. An ordinal classification for the families of flowering plants. *Annals of the Missouri*
Botanical Garden. Missouri Botanical Garden 85: 531.
Arseneau, J.-R., R. Steeves, and M. Laflamme. 2017. Modified low-salt CTAB extraction of
high-quality DNA from contaminant-rich tissues. *Molecular ecology resources* 17: 686–
693.
Auguie, B., and A. Antonov. 2017. gridExtra: miscellaneous functions for "grid" graphics. *R*
package version 2.
Avise, J. C., and T. J. Robinson. 2008. Hemiplasy: a new term in the lexicon of phylogenetics.
Systematic biology 57: 503–507.
Baker, W. J., P. Bailey, V. Barber, A. Barker, S. Bellot, D. Bishop, L. R. Botigué, et al. 2021. A
Comprehensive Phylogenomic Platform for Exploring the Angiosperm Tree of Life.
Systematic biology.
Barchi, L., M. Pietrella, L. Venturini, A. Minio, L. Toppino, A. Acquadro, G. Andolfo, et al.
2019. A chromosome-anchored eggplant genome sequence reveals key events in
Solanaceae evolution. *Scientific reports* 9: 11769.
Bennett, J. R. 2008. Revision of *Solanum* section Regmandra (Solanaceae). *Edinburgh journal of*
botany 65: 69–112.
Bohs, L. 2004. A chloroplast DNA phylogeny of *Solanum* section Lasiocarpa. *Systematic botany*
29: 177–187.
Bohs, L. 2005. Major clades in *Solanum* based on ndhF sequence data. *Monographs in*
Systematic Botany 104: 27.
Bohs, L., and R. G. Olmstead. 2001. A reassessment of *Normania* and *Triguera* (Solanaceae).
Plant systematics and evolution 228: 33–48.
Bohs, L., and R. G. Olmstead. 1997. Phylogenetic relationships in *Solanum* (Solanaceae) based
on ndhF sequences. *Systematic botany* 22: 5.
Bolger, A. M., M. Lohse, and B. Usadel. 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics (Oxford, England)* 30: 2114–2120.
Bouckaert, R., T. G. Vaughan, J. Barido-Sottani, S. Duchêne, M. Fourment, A. Gavryushkina, J.
Heled, et al. 2019. BEAST 2.5: An advanced software platform for Bayesian
evolutionary analysis. *PLoS computational biology* 15: e1006650.
Brown, J. W., J. F. Walker, and S. A. Smith. 2017. Phyx: phylogenetic tools for unix.
Bioinformatics 33: 1886–1888.
Capella-Gutiérrez, S., J. M. Silla-Martínez, and T. Gabaldón. 2009. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics (Oxford,*
England) 25: 1972–1973.
Chavana, J., S. Singh, A. Vazquez, B. Christoffersen, A. Racelis, and R. R. Kariyat. 2021. Local
adaptation to continuous mowing makes the noxious weed Solanum elaeagnifolium a
superweed candidate by improving fitness and defense traits. *Scientific reports* 11: 6634.

Chiarini, F., F. Sazatornil, and G. Bernardello. 2018. Data reassessment in a phylogenetic
context gives insight into chromosome evolution in the giant genus Solanum
(Solanaceae). *Systematics and biodiversity* 16: 397–416.
Cowie, B. W., N. Venter, E. T. F. Witkowski, M. J. Byrne, and T. Olckers. 2018. A review of
Solanum mauritianum biocontrol: prospects, promise and problems: a way forward for
South Africa and globally. *Biocontrol* 63: 475–491.
Darriba, D., D. Posada, A. M. Kozlov, A. Stamatakis, B. Morel, and T. Flouri. 2020. ModelTest-
NG: A new and scalable tool for the selection of DNA and protein evolutionary models.
Molecular biology and evolution 37: 291–294.
Degnan, J. H., and N. A. Rosenberg. 2009. Gene tree discordance, phylogenetic inference and
the multispecies coalescent. *Trends in ecology & evolution* 24: 332–340.
Dodsworth, S., M. J. M. Christenhusz, J. G. Conran, M. S. Guignard, S. Knapp, M. Struebig, A.
R. Leitch, and M. W. Chase. 2020. Extensive plastid-nuclear discordance in a recent
radiation of *Nicotiana* section Suaveolentes (Solanaceae). *Botanical journal of the*
Linnean Society. Linnean Society of London 193: 546–559.
Duvall, M. R., S. V. Burke, and D. C. Clark. 2020. Plastome phylogenomics of Poaceae:
alternate topologies depend on alignment gaps. *Botanical journal of the Linnean Society.*
Linnean Society of London 192: 9–20.
Echeverría‑Londoño, S., T. Särkinen, I. S. Fenton, A. Purvis, and S. Knapp. 2020. Dynamism
and context‑dependency in diversification of the megadiverse plant genus *Solanum*
(Solanaceae). *Journal of systematics and evolution* 58: 767–782.
Edelman, N. B., P. B. Frandsen, M. Miyagi, B. Clavijo, J. Davey, R. B. Dikow, G. García-
Accinelli, et al. 2019. Genomic architecture and introgression shape a butterfly radiation.
Science 366: 594–599.
Edgar, R. C. 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*
26: 2460–2461.
Ewels, P., M. Magnusson, S. Lundin, and M. Käller. 2016. MultiQC: summarize analysis results
for multiple tools and samples in a single report. *Bioinformatics* 32: 3047–3048.
Gardner, E. M., M. Garner, R. Cowan, S. Dodsworth, N. Epitawalage, D. Arifiani, S. Sahromi, et
al. 2020. Repeated parallel losses of inflexed stamens in Moraceae: phylogenomics and
generic revision of the tribe Moreae and the reinstatement of the tribe Olmedieae
(Moraceae). *bioRxiv*.
Gates, D. J., D. Pilson, and S. D. Smith. 2018. Filtering of target sequence capture individuals
facilitates species tree construction in the plant subtribe Iochrominae (Solanaceae).
Molecular phylogenetics and evolution 123: 26–34.
Hahn, M. W., and L. Nakhleh. 2016. Irrational exuberance for resolved species trees. *Evolution;*
international journal of organic evolution 70: 7–17.
Hilgenhof, R., E. Gagnon, S. Knapp, X. Aubriot, E.J. Tepe, L. Bohs, L.L. Giacomin, Y.F.
Gouvea, J.R. Stehmann, A Orejuela, C.I. Orozco, I.E. Peralta, and T. Sarkinen. 2021.
Evolution of key morphological traits in *Solanum* (Solanaceae): identifying areas of
interest for further study. *American Journal of Botany*, in prep.
Hime, P. M., A. R. Lemmon, E. C. M. Lemmon, E. Prendini, J. M. Brown, R. C. Thomson, J. D.
Kratovil, et al. 2021. Phylogenomics reveals ancient gene tree discordance in the
amphibian tree of life. *Systematic biology* 70: 49–66.
Huang, B., H. Ruess, Q. Liang, C. Colleoni, and D. M. Spooner. 2019. Analyses of 202 plastid
genomes elucidate the phylogeny of *Solanum* section Petota. *Scientific reports* 9: 4454.

Huson, D. H., and D. Bryant. 2006. Application of phylogenetic networks in evolutionary
studies. *Molecular biology and evolution* 23: 254–267.
Jeffroy, O., H. Brinkmann, F. Delsuc, and H. Philippe. 2006. Phylogenomics: the beginning of
incongruence? *Trends in genetics: TIG* 22: 225–231.
Jin, J.-J., W.-B. Yu, J.-B. Yang, Y. Song, C. W. dePamphilis, T.-S. Yi, and D.-Z. Li. 2020.
GetOrganelle: a fast and versatile toolkit for accurate de novo assembly of organelle
genomes. *Genome biology* 21: 241.
Johnson, M. G., E. M. Gardner, Y. Liu, R. Medina, B. Goffinet, A. J. Shaw, N. J. C. Zerega, and
N. J. Wickett. 2016. HybPiper: Extracting coding sequence and introns for phylogenetics
from high-throughput sequencing reads using target enrichment. *Applications in plant*
sciences 4: 1600016.
Johnson, M. G., L. Pokorny, S. Dodsworth, L. R. Botigué, R. S. Cowan, A. Devault, W. L.
Eiserhardt, et al. 2019. A universal probe set for targeted sequencing of 353 nuclear
genes from any flowering plant designed using k-Medoids clustering. *Systematic biology*
68: 594–606.
Junier, T., and E. M. Zdobnov. 2010. The Newick utilities: high-throughput phylogenetic tree
processing in the UNIX shell. *Bioinformatics (Oxford, England)* 26: 1669–1670.
Katoh, K., K.-I. Kuma, H. Toh, and T. Miyata. 2005. MAFFT version 5: improvement in
accuracy of multiple sequence alignment. *Nucleic acids research* 33: 511–518.
Knapp, S. 2013. A revision of the Dulcamaroid clade of *Solanum* L. (Solanaceae). *PhytoKeys* 22:
1–432.
Knapp, S., G. E. Barboza, L. Bohs, and T. Särkinen. 2019. A revision of the Morelloid clade of
Solanum L. (Solanaceae) in North and Central America and the Caribbean. *PhytoKeys*
123: 1–144.
Knapp, S., and Others. 2000. A revision of *Solanum thelopodium* species group (section
Anthoresis sensu Seithe, pro parte): Solanaceae. *Bulletin of the Natural History Museum,*
Botany Series 30: 13–30.
Knapp, S., E. Sagona, A. K. Z. Carbonell, and F. Chiarini. 2017. A revision of the *Solanum*
elaeagnifolium clade (Elaeagnifolium clade; subgenus *Leptostemonum*, Solanaceae).
PhytoKeys: 1–104.
Knapp, S., and M. S. Vorontsova. 2016. A revision of the "African Non-Spiny" Clade of
Solanum L. (*Solanum* sections Afrosolanum Bitter, Benderianum Bitter, Lemurisolanum
Bitter, Lyciosolanum Bitter, Macronesiotes Bitter, and Quadrangulare Bitter:
Solanaceae). *PhytoKeys* 66: 1–142.
Koenen, E. J. M., D. I. Ojeda, and F. T. Bakker. 2021. The origin of the legumes is a complex
paleopolyploid phylogenomic tangle closely associated with the cretaceous–paleogene
(K–Pg) mass extinction event. *Systematic biology* 70(3): 508–526.
Kumar, S., A. J. Filipski, F. U. Battistuzzi, S. L. Kosakovsky Pond, and K. Tamura. 2012.
Statistics and truth in phylogenomics. *Molecular biology and evolution* 29: 457–472.
Kuramae, E. E., V. Robert, B. Snel, M. Weiss, and T. Boekhout. 2006. Phylogenomics reveal a
robust fungal tree of life. *FEMS yeast research* 6: 1213–1220.
Lanfear, R., B. Calcott, S. Y. W. Ho, and S. Guindon. 2012. Partitionfinder: combined selection
of partitioning schemes and substitution models for phylogenetic analyses. *Molecular*
biology and evolution 29: 1695–1701.

Larridon, I., T. Villaverde, A. R. Zuntini, L. Pokorny, G. E. Brewer, N. Epitawalage, I. Fairlie, et
al. 2020. Tackling Rapid Radiations With Targeted Sequencing. *Frontiers in plant*
science 10: 1655.
Levin, R. A., N. R. Myers, and L. Bohs. 2006. Phylogenetic relationships among the "spiny
solanums" (*Solanum* subgenus *Leptostemonum*, Solanaceae). *American journal of botany*
93: 157–169.
Levin, R. A., K. Watson, and L. Bohs. 2005. A four-gene study of evolutionary relationships in
Solanum section Acanthophora. *American journal of botany* 92: 603–612.
Li, H., and R. Durbin. 2010. Fast and accurate long-read alignment with Burrows–Wheeler
transform. *Bioinformatics* 26: 589–595.
Liu, L., L. Yu, L. Kubatko, D. K. Pearl, and S. V. Edwards. 2009. Coalescent methods for
estimating phylogenetic trees. *Molecular phylogenetics and evolution* 53: 320–328.
Louca, S., and M. W. Pennell. 2020. Extant timetrees are consistent with a myriad of
diversification histories. *Nature* 580: 502–505.
Lutteropp, S., C. Scornavacca, A. M. Kozlov, and B. Morel. 2021. NetRAX: Accurate and Fast
Maximum Likelihood Phylogenetic Network Inference. *bioRxiv*.
Malinsky, M., H. Svardal, A. M. Tyers, E. A. Miska, M. J. Genner, G. F. Turner, and R. Durbin.
2018. Whole-genome sequences of Malawi cichlids reveal multiple radiations
interconnected by gene flow. *Nature ecology & evolution* 2: 1940–1955.
Mallet, J., N. Besansky, and M. W. Hahn. 2016. How reticulated are species? *BioEssays: news*
and reviews in molecular, cellular and developmental biology.
Meleshko, O., M. D. Martin, T. S. Korneliussen, C. Schröck, P. Lamkowski, J. Schmutz, A.
Healey, et al. 2021. Extensive Genome-Wide Phylogenetic Discordance Is Due to
Incomplete Lineage Sorting and Not Ongoing Introgression in a Rapidly Radiated
Bryophyte Genus. *Molecular biology and evolution* 38: 2750–2766.
Mendes, F. K., Y. Hahn, and M. W. Hahn. 2016. Gene Tree Discordance Can Generate Patterns
of Diminishing Convergence over Time. *Molecular biology and evolution* 33: 3299–
3307.
Miller, J. S., A. Kamath, and R. A. Levin. 2009. Do multiple tortoises equal a hare? The utility of
nine noncoding plastid regions for species-level phylogenetics in tribe Lycieae
(Solanaceae). *Systematic botany* 34: 796–804.
Miller, M. A., W. Pfeiffer, and T. Schwartz. 2010. Creating the CIPRES Science Gateway for
inference of large phylogenetic trees. 2010 Gateway Computing Environments Workshop
(GCE), IEEE.
Minh, B. Q., M. W. Hahn, and R. Lanfear. 2020. New methods to calculate concordance factors
for phylogenomic datasets. *Molecular biology and evolution* 37: 2727–2733.
Minh, B. Q., H. A. Schmidt, O. Chernomor, D. Schrempf, M. D. Woodhams, A. von Haeseler,
and R. Lanfear. 2020. IQ-TREE 2: new models and efficient methods for phylogenetic
inference in the genomic era. *Molecular biology and evolution* 37: 1530–1534.
Morgan, C. C., P. G. Foster, A. E. Webb, D. Pisani, J. O. McInerney, and M. J. O'Connell. 2013.
Heterogeneous models place the root of the placental mammal phylogeny. *Molecular*
biology and evolution 30: 2145–2156.
Olave, M., and A. Meyer. 2020. Implementing large genomic single nucleotide polymorphism
data sets in phylogenetic network reconstructions: a case study of particularly rapid
radiations of cichlid fish. *Systematic biology* 69: 848–862.

One Thousand Plant Transcriptomes Initiative. 2019. One thousand plant transcriptomes and the
phylogenomics of green plants. *Nature* 574: 679–685.
Paradis, E., and K. Schliep. 2019. ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics* 35: 526–528.
Parins-Fukuchi, C., G. W. Stull, and S. A. Smith. 2021. Phylogenomic conflict coincides with
rapid morphological innovation. *Proceedings of the National Academy of Sciences of the*
United States of America 118.
Pease, J. B., D. C. Haak, M. W. Hahn, and L. C. Moyle. 2016. Phylogenomics reveal three
sources of adaptive variation during a rapid radiation. *PLoS biology* 14: e1002379.
Pellicer, J., and I. J. Leitch. 2020. The Plant DNA C-values database (release 7.1): an updated
online repository of plant genome size data for comparative studies. *The new phytologist*
226: 301–305.
Peralta, I., S. Knapp, and D. Spooner. 2007. The taxonomy of tomatoes: a revision of wild
tomatoes (*Solanum* L. section Lycopersicon (Mill.) Wettst.) and their outgroup relatives
(*Solanum* sections Juglandifolium (Rydb.) Child and Lycopersicoides (Child) Peralta).
Systematic botany monographs 84: 1–186.
Philippe, H., H. Brinkmann, D. V. Lavrov, D. T. J. Littlewood, M. Manuel, G. Wörheide, and D.
Baurain. 2011. Resolving difficult phylogenetic questions: why more sequences are not
enough. *PLoS biology* 9: e1000602.
Philippe, H., D. M. de Vienne, V. Ranwez, B. Roure, D. Baurain, and F. Delsuc. 2017. Pitfalls in
supermatrix phylogenomics. *European Journal of Taxonomy* 283. doi:
10.5852/ejt.2017.283
Poczai, P. 2013. To network or not to network, that is the question. *Journal of genetics* 92: 703–
705.
Price, M. N., P. S. Dehal, and A. P. Arkin. 2010. FastTree 2 – approximately maximum-
likelihood trees for large alignments. *PloS one* 5: e9490.
Rambaut, A., A. J. Drummond, D. Xie, G. Baele, and M. A. Suchard. 2018. Posterior
summarization in bayesian phylogenetics using Tracer 1.7. *Systematic biology* 67: 901–
904.
Reddy, S., R. T. Kimball, A. Pandey, P. A. Hosner, M. J. Braun, S. J. Hackett, K.-L. Han, et al.
2017. Why Do Phylogenomic data sets yield conflicting trees? Data type influences the
avian tree of life more than taxon sampling. *Systematic biology* 66: 857–879.
Romiguier, J., V. Ranwez, F. Delsuc, N. Galtier, and E. J. P. Douzery. 2013. Less is more in
mammalian phylogenomics: AT-rich genes minimize tree conflicts and unravel the root
of placental mammals. *Molecular biology and evolution* 30: 2134–2144.
Ronco, F., M. Matschiner, A. Böhne, A. Boila, H. H. Büscher, A. El Taher, A. Indermaur, et al.
2021. Drivers and dynamics of a massive adaptive radiation in cichlid fishes. *Nature* 589:
76–81.
Rosario, L. H., J. O. Rodríguez Padilla, D. R. Martínez, A. M. Grajales, J. A. Mercado Reyes, G.
J. Veintidós Feliu, B. Van Ee, and D. Siritunga. 2019. DNA barcoding of the Solanaceae
family in Puerto Rico including endangered and endemic species. *Journal of the*
American Society for Horticultural Science 144: 363–374.
Saarela, J. M., S. V. Burke, W. P. Wysocki, M. D. Barrett, L. G. Clark, J. M. Craine, P. M.
Peterson, et al. 2018. A 250 plastome phylogeny of the grass family (Poaceae):
topological support under different data partitions. *PeerJ* 6: e4299.

Sang, T., D. Crawford, and T. Stuessy. 1997. Chloroplast DNA phylogeny, reticulate evolution,
1060   and biogeography of Paeonia (Paeoniaceae). *American journal of botany* 84: 1120.
1061   Särkinen, T., G. E. Barboza, and S. Knapp. 2015. True Black nightshades: phylogeny and
1062   delimitation of the Morelloid clade of *Solanum*. *Taxon* 64: 945–958.
1063   Särkinen, T., L. Bohs, R. G. Olmstead, and S. Knapp. 2013. A phylogenetic framework for
1064   evolutionary study of the nightshades (Solanaceae): a dated 1000-tip tree. *BMC*
1065   *evolutionary biology* 13: 214.
1066   Särkinen, T., P. Poczai, G. E. Barboza, G. M. van der Weerden, M. Baden, and S. Knapp. 2018.
1067   A revision of the Old World Black Nightshades (Morelloid clade of *Solanum* L.,
1068   Solanaceae). *PhytoKeys*: 1–223.
1069   Sayyari, E., and S. Mirarab. 2016. Fast Coalescent-based computation of local branch support
1070   from quartet frequencies. *Molecular biology and evolution* 33: 1654–1668.
1071   Sayyari, E., and S. Mirarab. 2018. Testing for polytomies in phylogenetic species trees using
1072   quartet frequencies. *Genes* 9.
1073   Schrempf, D., and G. Szöllősi. 2020. The sources of phylogenetic conflicts. *Phylogenetics in the*
1074   *Genomic Era*: 3.1:1-3.1:23.
1075   Simion, P., H. Philippe, D. Baurain, M. Jager, D. J. Richter, A. Di Franco, B. Roure, et al. 2017.
1076   A large and consistent phylogenomic dataset supports sponges as the sister group to all
1077   other animals. *Current biology* 27: 958–967.
1078   Solís-Lemus, C., P. Bastide, and C. Ané. 2017. PhyloNetworks: a package for phylogenetic
1079   networks. *Molecular biology and evolution* 34: 3292–3298.
1080   Spalink, D., K. Stoffel, G. K. Walden, A. M. Hulse-Kemp, T. A. Hill, A. Van Deynze, and L.
1081   Bohs. 2018. Comparative transcriptomics and genomic patterns of discordance in
1082   Capsiceae (Solanaceae). *Molecular phylogenetics and evolution* 126: 293–302.
1083   Spooner, D. M., N. Alvarez, I. E. Peralta, and A. M. Clausen. 2016. Taxonomy of wild potatoes
1084   and their relatives in Southern South America (*Solanum* sect. Petota and Etuberosum).
1085   *Systematic botany monographs* 100: 1–240.
1086   Spooner, D. M., R. G. van den Berg, A. Rodrigues, J. B. Bamberg, R. J. Hijmans, and S. Lara-
1087   Cabrera. 2004. Wild potatoes (*Solanum* section Petota; Solanaceae) of North and Central
1088   America. *Systematic botany monographs* 68: 1–209.
1089   Spooner, D. M., S. Jansky, F. Rodríguez, R. Simon, M. Ames, D. Fajardo, and R. O. Castillo.
1090   2019. Taxonomy of wild potatoes in northern South America (*Solanum* section Petota).
1091   *Systematic botany monographs* 108: 1–305.
1092   Springer, M. S., and J. Gatesy. 2018. Delimiting Coalescence Genes (C-Genes) in Phylogenomic
1093   Data Sets. *Genes* 9.
1094   Stamatakis, A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
1095   large phylogenies. *Bioinformatics* 30: 1312–1313.
1096   Stamatakis, A. 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with
1097   thousands of taxa and mixed models. *Bioinformatics* 22: 2688–2690.
1098   Stern, S., M. de F. Agra, and L. Bohs. 2011. Molecular delimitation of clades within New World
1099   species of the "spiny solanums" (*Solanum* subg. Leptostemonum). *Taxon* 60: 1429–1441.
1100   Stern, S., and L. Bohs. 2012. An explosive innovation: phylogenetic relationships of *Solanum*
1101   section Gonatotrichum (Solanaceae). *PhytoKeys*: 89–98.
1102   Strickler, S. R., A. Bombarely, J. D. Munkvold, T. York, N. Menda, G. B. Martin, and L. A.
1103   Mueller. 2015. Comparative genomics and phylogenetic discordance of cultivated tomato
1104   and close wild relatives. *PeerJ* 3: e793.

Suh, A. 2016. The phylogenomic forest of bird trees contains a hard polytomy at the root of Neoaves. *Zoologica scripta* 45: 50–62.

Suh, A., L. Smeds, and H. Ellegren. 2015. The dynamics of incomplete lineage sorting across the ancient adaptive radiation of neoavian birds. *PLoS biology* 13: e1002224.

Symon, D. E. 1994. Kangaroo apples: *Solanum* sect. Archaesolanum. Adelaide, Australia: Published by the author.

Taberlet, P., L. Gielly, G. Pautou, and J. Bouvet. 1991. Universal primers for amplification of three non-coding regions of chloroplast DNA. *Plant molecular biology* 17: 1105–1109.

Tavaré, S. 1986. Some probabilistic and statistical problems in the analysis of DNA sequences. *Lectures on mathematics in the life sciences* 17: 57–86.

Tepe, E. J., G. J. Anderson, D. M. Spooner, and L. Bohs. 2016. Relationships among wild relatives of the tomato, potato, and pepino. *Taxon* 65: 262–276.

Than, C., D. Ruths, and L. Nakhleh. 2008. PhyloNet: a software package for analyzing and reconstructing reticulate evolutionary relationships. *BMC bioinformatics* 9: 322.

Tillich, M., P. Lehwark, T. Pellizzer, E. S. Ulbricht-Jones, A. Fischer, R. Bock, and S. Greiner. 2017. GeSeq-versatile and accurate annotation of organelle genomes. *Nucleic acids research* 45: W6–W11.

Villanueva, R. A. M., and Z. J. Chen. 2019. ggplot2: elegant graphics for data analysis (2nd ed.). *Measurement: interdisciplinary research and perspectives* 17: 160–167.

Weese, T. L., and L. Bohs. 2007. A three-gene phylogeny of the genus *Solanum* (Solanaceae). *Systematic botany* 32: 445–463.

Wen, D., Y. Yu, J. Zhu, and L. Nakhleh. 2018. Inferring phylogenetic networks using PhyloNet. *Systematic biology* 67: 735–740.

Wendel, J. F., and J. J. Doyle. 1998. Phylogenetic incongruence: window into genome history and molecular evolution. *In* D. E. Soltis, P. S. Soltis, and J. J. Doyle [eds.], Molecular Systematics of Plants II: DNA Sequencing, 265–296. Springer US, Boston, MA.

Wickett, N. J., S. Mirarab, N. Nguyen, T. Warnow, E. Carpenter, N. Matasci, S. Ayyampalayam, et al. 2014. Phylotranscriptomic analysis of the origin and early diversification of land plants. *Proceedings of the National Academy of Sciences of the United States of America* 111: E4859-68.

Wickham, H., and M. H. Wickham. 2019. Package 'stringr.'

Williams, T. A., D. Schrempf, G. J. Szöllősi, C. J. Cox, P. G. Foster, and T. M. Embley. 2021. Inferring the deep past from molecular data. *Genome biology and evolution* 13.

Wu, M., J. L. Kostyun, and L. C. Moyle. 2019. Genome sequence of *Jaltomata* addresses rapid reproductive trait evolution and enhances comparative genomics in the hyper-diverse Solanaceae. *Genome biology and evolution* 11: 335–349.

Yu, G. 2020. Using ggtree to visualize data on tree-like structures. *Current protocols in bioinformatics* 69: e96.

Zhang, C., M. Rabiee, E. Sayyari, and S. Mirarab. 2018. ASTRAL-III: polynomial time species tree reconstruction from partially resolved gene trees. *BMC bioinformatics* 19.

Tables

Table 1. Number of species and taxon sampling across major and minor clades of *Solanum.*

Clades are based on groups identified in previous molecular phylogenetic studies (Bohs, 2005;

Weese and Bohs, 2007; Stern et al., 2011; Stern and Bohs, 2012; Särkinen et al., 2013; Tepe et

al., 2016). Species number for each clade is based on current updated taxonomy in the

SolanaceaeSource database. The 19 clades sampled in the pruned trees for the principal

coordinate analysis in this study are in bold. New associated major clade names are given where

applicable. Rows shaded in gray represent major and minor clades belonging to Clade II. The

Eastern Hemisphere Spiny clade (EHS, formerly known as Old World spiny clade) comprises

almost all the spiny solanums occurring in the eastern hemisphere.

Minor clade	Associated major clade (Särkinen et al. 2013)	New associated major clade (This study)	Species	Sampled species (%)		
				Super-matrix	Plastome (PL)	Target Capture (TC)
Thelopodium	**Thelopodium**		**3**	**3 (100%)**	**1 (33%)**	**1 (33%)**
African non-spiny	M Clade	VANAns	14	5 (36%)	1 (7%)	-
Normania	**M Clade**	**VANAns**	**3**	**2 (67%)**	**1 (33%)**	**1 (33%)**
Archaesolanum	**M Clade**	**VANAns**	**8**	**8 (100%)**	**1 (13%)**	**1 (13%)**
Valdiviense	**M Clade**	**VANAns**	**1**	**1 (100%)**	**1 (100%)**	**1 (100%)**
Dulcamaroid	**M Clade**	**DulMo**	**45**	**25 (56%)**	**8 (18%)**	**1 (2%)**
Morelloid	**M Clade**	**DulMo**	**75**	**66 (88%)**	**15 (20%)**	**1 (1%)**
Regmandra	**Potato**	**Regmandra**	**12**	**6 (50%)**	**4 (33%)**	**1 (8%)**
Herpystichum	Potato		10	10 (100%)	-	-
Pteroidea	Potato		10	10 (100%)	1 (10%)	-
Oxycoccoides	Potato		1	1 (100%)	-	-
Articulatum	Potato		2	2 (100%)	-	-
Basarthrum	**Potato**		**16**	**10 (56%)**	**3 (19%)**	**3 (19%)**
Anarrhichomenum	Potato		12	8 (82%)		
Etuberosum	**Potato**		**3**	**2 (67%)**	**2 (67%)**	**1 (33%)**
Tomato	**Potato**		**7**	**14 (82%)**	**8 (47%)**	**3 (18%)**
Petota	**Potato**		**113**	**61 (54%)**	**38 (34%)**	**2 (2%)**
Clandestinum-Mapiriense	**Clandestinum-Mapiriense**		**3**	**3 (100%)**	**1 (33%)**	**1 (33%)**
Wendlandii-Allophyllum	**Wendlandii-Allophyllum**		**10**	**7 (70%)**	**1 (10%)**	**1 (10%)**

Nemorense	Nemorense	4	4 (100%)	1 (25%)	-
Pachyphylla	Cyphomandra	39	32 (82%)	1 (3%)	-
Cyphomandropsis	Cyphomandra	11	7 (64%)	1 (9%)	1 (9%)
Geminata	**Geminata**	**150**	**68 (45%)**	**5 (3%)**	**1 (1%)**
Reductum	Geminata	2	2 (100%)	1 (50%)	-
Brevantherum	Brevantherum	83	29 (35%)	3 (4%)	-
Gonatotrichum	Brevantherum	7	7 (100%)	1 (14%)	-
Inornatum	Brevantherum	5	2 (40%)	1 (20%)	-
Trachytrichium	Brevantherum	2	2 (100%)	-	-
Elaeagniifolium	**Leptostemonum**	**5**	**5 (100%)**	**1 (20%)**	**1 (20%)**
Micracantha	Leptostemonum	14	9 (64%)	1 (7%)	-
Torva	**Leptostemonum**	**54**	**34 (63%)**	**5 (9%)**	**1 (2%)**
Erythrotrichum	Leptostemonum	33	13 (39%)	1 (3%)	-
Thomasiifolium	Leptostemonum	9	4 (44%)	1 (11%)	-
Gardneri	Leptostemonum	10	8 (80%)	1 (10%)	-
Acanthophora	**Leptostemonum**	**22**	**13 (59%)**	**1 (5%)**	**-**
Lasiocarpa	Leptostemonum	12	12 (100%)	-	-
Sisymbriifolium	**Leptostemonum**	**4**	**4 (100%)**	**1 (25%)**	**1 (25%)**
Androceras	Leptostemonum	16	15 (94%)	-	-
Crinitum	Leptostemonum	23	10 (43%)	-	-
Bahamense	Leptostemonum	3	3 (100%)	-	-
Asterophorum	Leptostemonum	4	2 (50%)	-	-
Carolinense	Leptostemonum	11	8 (73%)	1 (9%)	-
Hieronymi	Leptostemonum	1	1 (100%)	1 (100%)	-
Eastern Hemisphere Spiny	**Leptostemonum**	**332**	**197 (59%)**	**24 (7%)**	**16 (5%)**
Campechiense	Leptostemonum	1	1 (100%)	-	-
Crotonoides	Leptostemonum	3	2 (67%)	1 (33%)	-
Multispinum	Leptostemonum	1	1 (100%)	1 (100%)	-
Unplaced	Leptostemonum	9	1 (13%)	-	-
	TOTAL:	1,228	746 (60%)	140 (11%)	39 (3%)

1156

1157 **Table 2. Overview of the 10 different analyses conducted across the Sanger supermatrix,**

1158 **plastome (PL) and target capture (TC) datasets.** Acronyms indicate how each analysis is

1159 referred to in the figures and text. ML = Maximum Likelihood; BI = Bayesian Inference, A353 =

1160 Angiosperms353 bait set. See Methods section for full details.

Dataset	Taxon & genomic sampling	Phylogenetic method	Partitioning scheme	Acronym
Supermatrix	746 taxa, 9 loci	ML: RaxML	--	Supermatrix ML
		BI : Beast2	--	Supermatrix BI
Plastome (PL)	151 taxa, full + partial plastomes	ML: IQ-TREE2	Unpartitioned	PL-151-UP
	151 taxa, full + partial plastomes	ML: IQ-TREE2	Best-Partition scheme	PL-151-BP
	125 taxa, full plastomes only	ML: IQ-TREE2	Unpartitioned	PL-125-UP
	125 taxa, full plastomes	ML: IQ-TREE2	Best-Partition scheme	PL-125-BP
Target Capture (TC) (A353)	40 taxa, 338 exons	ML: IQ-TREE2	--	TC-min04-ML
	40 taxa, 338 exons	Coalescent: ASTRAL-III	--	TC-min04-ASTRAL-III
	40 taxa, 303 exons	ML: IQ-TREE2	--	TC-min20-ML
	40 taxa, 303 exons	Coalescent: ASTRAL-III	--	TC-min20-ASTRAL-III

1161

Figures

Figure 1. Supermatrix phylogeny from Maximum Likelihood analysis (RaxML) of 742

Solanum **species based on two nuclear and seven plastid regions**. Bootstrap branch support

values are color coded: black = strong (0.95–1.0), white = moderate to weak support (0.75–0.94).

Dashed lines indicate in phylogeny indicate relationships that were not recovered in the TC and

PL analyses (see Figures 2-3). Clade names refer to major and minor clades discussed in the text

(see Table 1); dashed lines for clade labels indicate groups that were not recovered in the TC and

PL analyses.

1170

Figure 2. Comparison of *Solanum* clades recovered in plastome (PL) and target-capture

(TC) phylogenomic datasets. a) Plastome phylogeny from the unpartitioned maximum

likelihood analysis (PL-151-UP) based on 160 loci representing exons, introns and intergenic

regions; b) Filtered supertree network of the TC dataset (min20) based on 303 gene trees with a

50% minimum tree threshold. c) TC phylogeny with 40 species from coalescent analysis (TC-

min20-ASTRAL). Clades are shown in the same color in all three phylogenies to enable

comparison. Branch support values (BS values in (a) and local PP values in (c)) are color coded:

black = strong (0.95–1.0), white = moderate to weak (0.75–0.94). Scale bars = substitutions/site.

Collection or Genbank numbers are indicated in the PL phylogeny for duplicate species sampled

in the phylogenetic trees.

Figure 3. Comparison of *Solanum* clades recovered in the three different datasets. a) TC

Astral-III phylogeny of the min-20 dataset, with local posterior probabilities indicated at nodes;

b) ML and BI phylogenies of supermatrix dataset, with bootstrap support and posterior

probabilities indicated at nodes; c) TC ML phylogeny of the min 20 dataset, with local posterior

probabilities indicated at nodes; d) PL ML phylogenies of the unpartitioned and best partition-

scheme of the 151 taxa dataset, with bootstrap for each respective analysis is indicated at nodes;

e) TC ML phylogeny and Astral-III phylogeny of the min 04 dataset, with bootstrap support and

local posterior probabilities indicated at nodes; f) PL ML phylogenies of the unpartitioned and

best partition-scheme of the 125 taxa dataset, with bootstrap for each respective analysis

indicated at nodes.

Figure 4. **Discordance analyses within and between the plastome (PL) and target capture**

(TC) phylogenomic datasets across *Solanum*. Rooted TC ASTRAL phylogeny (left) and PL

IQ-TREE2 phylogeny (right) with gene concordance factor (gCF) and site concordance factor

(sCF) values shown as pie charts, above and below each node respectively; the PL topology is

the unpartitioned ML analysis of 151 taxa, whereas the TC topology is based on the analysis of

40 taxa and 303 genes recovered from the A353 bait set. Both trees have been pruned to retain a

single tip for each of the major and minor clades present within the PL and TC datasets. For gCF

pie charts, blue represents proportion of gene trees concordant with that branch (gCF), green is

proportion of gene trees concordant for 1^{st} alternative quartet topology (gDF1), yellow support

for 2^{nd} alternative quartet topology (gDF2), and red is the gene discordance support due to

polylphyly (gDFP). For the sCF pie charts: blue represents proportion of concordance across

sites (sCF), green support for 1^{st} alternative topology (quartet 1), and yellow support for 2^{nd}

alternative topology (quartet 2) as averaged over 100 sites. Percentages of gCF and sCF are

given above branches, in bold. Branch support (local posterior probability) values ≥0.95 are not

shown, and 0.94 and below are shown in italic grey, on the right; double-dash (--) indicates that

the branch support was unavailable due to rooting of the phylogenetic tree. The nodes

corresponding to the two clear clades identified in the filtered supertree network (see Fig. 2b) are

identified as "N1" and "N2".

1214