Mathematical Models of Retinitis Pigmentosa: The Trophic Factor Hypothesis

Paul A. Roberts[a,*]

[a] *School of Life Sciences, University of Sussex, John Maynard Smith Building, Brighton, BN1 9QG, UK*

Abstract

Retinitis pigmentosa (RP) is the term used to denote a group of inherited retinal-degenerative conditions that cause progressive sight loss. Individuals with this condition lose their light-sensitive photoreceptor cells, known as rods and cones, over a period of years to decades; degeneration starting in the retinal periphery, and spreading peripherally and centrally over time. RP is a rod-cone dystrophy, meaning that rod health and function are affected earlier and more severely than that of cones. Rods degenerate due to an underlying mutation, whereas the reasons for cone degeneration are unknown. A number of mechanisms have been proposed to explain secondary cone loss and the spatio-temporal patterns of retinal degeneration in RP. One of the most promising is the trophic factor hypothesis, which suggests that rods produce a factor necessary for cone survival, such that, when rods degenerate, cone degeneration follows. In this paper we formulate and analyse mathematical models of RP under the trophic factor hypothesis. These models are constructed as systems of reaction diffusion partial differential equations in one spatial dimension, and are solved and analysed using a combination of numerical and analytical methods. We predict the conditions under which cones will degenerate following the loss of a patch of rods from the retina, the critical trophic factor treatment rate required to prevent cone degeneration following rod loss and the spatio-temporal patterns of cone loss that would result if the trophic factor mechanism alone were responsible for retinal degeneration.

Keywords: Partial Differential Equations, Asymptotic Analysis, Retina, Photoreceptors, Rod-derived Cone Viability Factor

1. Introduction

The group of inherited retinal diseases, known collectively as retinitis pigmentosa (RP), have been the subject of many decades of research. Despite this attention, many of the mechanisms underpinning RP have yet to be conclusively determined, though a number have been hypothesised (Hamel, 2006; Hartong et al., 2006). In this paper, we formulate mathematical models to describe and predict retinal degeneration in RP for one of the leading hypotheses: the trophic factor hypothesis.

The retina is the tissue layer at the back of the eye responsible for light detection (Fig. 1(a)). Retinal light-detecting cells are known as photoreceptors, which come in two varieties: rods and cones. Rods confer peripheral, achromatic vision under low-light (scotopic) conditions, while cones confer high-acuity central, chromatic vision under well-lit (photopic) conditions (Oyster, 1999). Cones are mostly concentrated in the

---

*Corresponding author

Email address: `p.a.roberts@univ.oxon.org` (Paul A. Roberts)

centre of the retina, known as the fovea, whereas rods dominate the mid- and far-peripheral retina (Curcio et al., 1990) (Fig. 1(b)). Both rods and cones are composed of inner and outer segments, together with an axon, ending in a synaptic terminal. Outer segments (OSs) contain a series of discs in which light-sensitive photo-pigments are embedded (Oyster, 1999). Each day, OS tips are shed and phagocytosed by the underlying retinal pigment epithelium (RPE), while new discs are generated where the OS meets the inner segment (Oyster, 1999; Young, 1967, 1971, 1978; Young and Bok, 1969).

RP is a rod-cone dystrophy, meaning that rod function and health are affected earlier and more severely than cone function and health (though there do exist rarer cone-rod dystrophy forms, and forms in which rods and cones are affected on a similar time scale, Hamel, 2006; Hartong et al., 2006). This is often because it is the rods, rather than the cones, which express the disease-causing mutation (though mutations can alternatively be expressed in the cones or RPE, Hamel, 2006; Hartong et al., 2006). Retinal degeneration (and hence visual field loss) typically initiates in the mid- or far-periphery of the retina, starting with small patches of photoreceptor loss which spread and coalesce into larger regions, causing blind spots (scotomas). Degeneration spreads through the mid- and far-periphery until all peripheral vision is lost (tunnel vision), finally spreading to the central retina (macula), leading to the loss of central vision, and hence, total blindness (Grover et al., 1998; Hamel, 2006; Hartong et al., 2006).

Given that the disease-causing mutations are not typically expressed by cones, it is unclear what causes them to degenerate. Further, the reasons behind the spatio-temporal patterns of retinal degeneration in general, and for which any particular degenerate patch expands or remains stable, remain a mystery (Hamel, 2006; Hartong et al., 2006). A number of mechanisms have been proposed to explain these phenomena, most notably trophic factor depletion (Léveillard et al., 2004), oxygen toxicity (Stone et al., 1999), metabolic disregulation (Punzo et al., 2009), toxic substances (Ripps, 2002) and microglia (Gupta et al., 2003). In this paper, we consider the first of these, often denoted as the trophic factor hypothesis. This hypothesis suggests that rods produce a trophic factor necessary for cone survival, such that, when rods degenerate, the trophic factor is depleted, and cone degeneration follows. Such a factor, known as rod-derived cone viability factor (RdCVF), has been chemically identified by Léveillard et al. (2004), and shown to slow cone degeneration and preserve cone function in chick, rat and mouse models (Fintz et al., 2003; Léveillard et al., 2004; Mohand-Saïd et al., 1998, 2000, 1997; Yang et al., 2009).

In this paper, we construct mathematical models to describe and predict retinal degeneration and preservation under the trophic factor hypothesis. Our models are formulated as systems of reaction diffusion partial differential equations (PDEs) in one spatial dimension, and solved and analysed using a combination of numerical and analytical techniques. We use these models to address three key questions: (i) under what conditions will the loss of a patch of rods lead to cone degeneration?, (ii) what is the critical trophic factor treatment dose required to prevent cone degeneration following a patch of rod loss?, and (iii) what spatio-temporal patterns of cone degeneration can be explained via this mechanism alone?

We previously developed the first and only models to consider the oxygen toxicity hypothesis (Roberts et al., 2017, 2018). These models are formulated as systems of PDEs in one- and two-dimensions and were the first models of RP to take the key step of accounting for the photoreceptor distribution. Burns et al. (2002) developed the only (1D PDE) model to consider the toxic substance hypothesis. Lastly, Camacho *et al.* have produced a series of ordinary differential equation models of the trophic factor hypothesis, focussing on the biochemical details, rather than the spatial spread, of retinal degeneration (Colón Vélez et al., 2003; Camacho et al., 2010; Camacho and Wirkus, 2013; Camacho et al., 2014, 2016a,b,c). For further details on mathematical modelling of the retina, including RP, see our review: Roberts et al. (2016). The models developed and analysed in this paper are the first spatially-resolved models to explore the trophic factor hypothesis, or to account for photoreceptor distribution. As such, these are the first models capable of answering the three key questions posed above.

Figure 1: Diagrams of the human eye and retinal photoreceptor distribution (reproduced, with permission, from Roberts et al., 2017). (a) Diagram of the (right) human eye, viewed in the transverse plane, illustrating the model geometry. All models are posed on a domain spanning the region between the foveal centre, at $\theta = 0$, and the ora serrata, at $\theta = \Theta$, along the temporal horizontal meridian, where θ measures the eccentricity. Figure originally reproduced, with modifications, from `http://www.nei.nih.gov/health/coloboma/coloboma.asp`, courtesy: National Eye Institute, National Institutes of Health (NEI/NIH). (b) Measured and fitted photoreceptor profiles, along the temporal horizontal meridian, in the human retina. Cone profile: $\tilde{p}_c(\theta) = B_1 e^{-b_1\theta} + B_2 e^{-b_2\theta}$, and rod profile: $\tilde{p}_r(\theta) = B_3\theta e^{-b_3\theta}$ (see Table 1 for parameter values). The photoreceptor profile is the sum of the rod and cone profiles ($\tilde{p}_r(\theta) + \tilde{p}_c(\theta)$). Experimental data provided by Curcio and published in Curcio et al. (1990).

The remainder of this paper is structured as follows. In Section 2, we formulate and non-dimensionalise our mathematical models. In Section 3, we consider the steady-state model, performing asymptotic analyses (Section 3.1), and comparing numerical and analytical solutions (Section 3.2) in the untreated (Section 3.2.1) and treated (Section 3.2.2) cases. In Section 4, we consider numerical solutions to the full dynamic (time dependent) model, in both the untreated (Section 4.1) and treated (Section 4.2) cases. Lastly, in Section 5, we discuss our results and suggest directions for future research.

2. Model formulation

We begin by formulating a model to describe the photoreceptor and trophic factor dynamics within the human retina. Given that the retina lines the essentially spherical inner surface of the eye, we pose our model in spherical polar coordinates (r,θ,ϕ), with origin at the centre of the vitreous body and orientated such that the z-axis passes outwards through the centre of the fovea (see Fig. 1(a)). We make two geometrically simplifying assumptions. Firstly, we assume that the retina is symmetric about the z-axis, such that the azimuthal dimension, ϕ (rad), can be neglected. While this ignores the optic disc (from which photoreceptors are absent) and other inhomogeneities in rod and cone densities, this is a reasonable simplification to make in the first spatial model of this disease mechanism. Secondly, since the retina's width ($\sim$80–320 μm, Webvision) is approximately two orders of magnitude smaller than its radius of curvature ($\sim$1.2 cm, Oyster, 1999), we depth-average through the retina, assuming a fixed radius $r = R$ (m). Thus, we have reduced the geometry to (R, θ); posing our model on the one-dimensional domain $\theta \in [0, \Theta]$ (rad), where $\theta = 0$ (rad) is located at the foveal centre and $\theta = \Theta$ (rad) is located at the ora serrata.

We consider two scenarios, leading to alternate forms of the governing equations:

- Scenario 1: without (w/o) cone regeneration;

- Scenario 2: with cone regeneration.

In Scenario 1, we model the cone photoreceptor density, $p_c(\theta, t)$ (photoreceptors m^{-2}), with its associated initial value, $p_{c_{init}}(\theta)$ (photoreceptors m^{-2}). In this case, cone degeneration corresponds to the loss of whole cone photoreceptor cells. Thus, given that photoreceptors cannot be regrown once they are lost, recovery is impossible. In Scenario 2, we model the local cone OS length, $p_c^L(\theta, t)$ (m), with its associated initial value, $p_{c_{init}}^L(\theta)$ (m). In this case, cone degeneration corresponds to the shortening of cone OS, while leaving the rest of the photoreceptor intact, such that cone density remains fixed at $p_{c_{init}}(\theta)$ (photoreceptors m^{-2}) for all time. As such, cone regeneration is possible in the form of OS regrowth (see Guérin et al., 1993, and also Chrysostomou et al., 2008; Lee et al., 2012). Since prolonged exposure to adverse conditions will lead to the eventual loss of the whole photoreceptor, it is assumed in this scenario that adverse conditions are either mild or relatively brief.

We construct a system of partial differential equations (PDEs) in terms of the dependent variables: trophic factor concentration, $f(\theta, t)$ (M), rod photoreceptor density, $p_r(\theta, t)$ (photoreceptors m^{-2}), and cone photoreceptor density, $p_c(\theta, t)$ (photoreceptors m^{-2}), or local cone OS length, $p_c^L(\theta, t)$ (m); as functions of the independent variables: polar angle, θ (rad), and time, t (s), where $t > 0$ (s).

The trophic factor equation takes the following form,

$$\frac{\partial f}{\partial t} = \underbrace{\frac{D_f}{R^2 \sin(\theta)} \frac{\partial}{\partial \theta} \left(\sin(\theta) \frac{\partial f}{\partial \theta} \right)}_{\text{diffusion}} + \underbrace{\alpha p_r}_{\text{production}} + \underbrace{\xi T(\theta, t)}_{\text{treatment}} - \underbrace{\beta C(\theta, f, p_c)}_{\text{consumption}} - \underbrace{\eta f}_{\text{decay}}, \tag{1}$$

where R (m), the radial position of the retina, D_f (m^{2}s^{-1}), the trophic factor diffusivity, α (Mm^{2}photoreceptors^{-1}s^{-1}), the rate of trophic factor production by rods, ξ (M s^{-1}), the rate of supply of trophic factor through treatment, β (m^{2}photoreceptors^{-1}s^{-1}), the rate of trophic factor consumption by cones, and η (s^{-1}), the rate of trophic factor decay, are positive constants. The treatment function is defined as follows

$$T(\theta, t) = \begin{cases} H(t - t_{crit}) & \text{global treatment,} \\ (1 - [H(\theta_{treat_1} - \theta) + H(\theta - \theta_{treat_2})])H(t - t_{crit}) & \text{local treatment,} \end{cases}$$

where trophic factor (RdCVF) treatment is given from time $t = t_{crit} > 0$ (s), and can be applied either globally, across the whole domain $\theta \in [0, \Theta]$ (rad), or locally, within the region $\theta \in (\theta_{treat_1}, \theta_{treat_2})$ (rad) only. We define the Heaviside step function, $H(\cdot)$, such that

$$H(x) := \begin{cases} 0 & \text{if } x < 0, \\ 1 & \text{if } x \geq 0. \end{cases}$$

The trophic factor consumption function is defined as follows

$$C(\theta, f, p_c) = \begin{cases} f p_c & \text{w/o cone regeneration,} \\ f p_{c_{init}}(\theta) & \text{with cone regeneration,} \end{cases}$$

where $p_{c_{init}}(\theta)$ (photoreceptors m^{-2}), the initial cone density, is defined below. In Scenario 1, cone density may decrease, such that the rate of TF consumption changes both with decreasing $p_c(\theta, t)$ and changing $f(\theta, t)$. By contrast, in Scenario 2, the cone density remains fixed, such that the rate of TF consumption changes with $f(\theta, t)$ only.

The rod equation is as follows

$$\frac{\partial p_r}{\partial t} = \begin{cases} & 0 \quad \text{w/o mutation-induced rod degeneration,} \\ - & \underbrace{\phi_r p_r}_{\substack{\text{cell degeneration} \\ \text{(mutation-induced)}}} \quad \text{with mutation-induced rod degeneration,} \end{cases} \tag{2}$$

where ϕ_r (s^{-1}), the rate of mutation-induced rod degeneration, is a positive constant. We keep the rod density constant in the case where we seek to focus on the photoreceptor dynamics following the formation of a degenerate patch (from which rods and/or cones are absent), while an exponential decline in rods more faithfully represents the natural disease progression. Therefore, in the case without mutation-induced rod degeneration, the rod density remains at its initial value, $p_r(\theta, t) = p_{r_{init}}(\theta)$ (photoreceptors m^{-2}), while in the case with mutation-induced rod degeneration, $p_r(\theta, t) = p_{r_{init}}(\theta)e^{-\phi_r t}$ (photoreceptors m^{-2}), provided there is no delay in onset or interruption of degeneration.

In Scenario 1, the cone photoreceptor dynamics are given by

$$\frac{\partial p_c}{\partial t} = \underbrace{\delta p_c \lambda_2(f)}_{\substack{\text{cell degeneration} \\ \text{(TF starvation)}}}, \tag{3}$$

where δ (s^{-1}), the rate of trophic factor starvation-induced cone degeneration, is a positive constant.

In Scenario 2, the cone photoreceptor dynamics are given by

$$\frac{\partial p_c^L}{\partial t} = \left[\underbrace{\mu_1 \lambda_3(p_c^L)}_{\substack{\text{OS regrowth} \\ \text{(phase 1)}}} + \underbrace{\mu_2 \left(1 - \frac{p_c^L}{\tilde{p}_c^L}\right) \lambda_4(p_c^L)}_{\substack{\text{OS regrowth} \\ \text{(phase 2)}}} \right] \lambda_1(f) \lambda_5(p_{c_{init}}) - \underbrace{\delta_L \lambda_2(f) \lambda_6(p_c^L)}_{\substack{\text{OS degeneration} \\ \text{(TF starvation)}}}, \tag{4}$$

where $\tilde{p}_c^L$ (m), the healthy cone OS length, μ_1 (s^{-1}), the growth rate of cone OS in phase 1, μ_2 (s^{-1}), the growth rate of cone OS in phase 2, and δ_L (ms^{-1}), the rate of trophic factor starvation-induced cone OS degeneration, are positive constants. In Scenario 1, cone cells degenerate exponentially, while in Scenario 2, individual cone OS degenerate at a constant rate.

Model fitting to cone OS regrowth data from Guérin et al. (1993) reveals that cone OS regrowth is well-described by a two-phase model. Phase 1, constant growth, occurs for cone OS lengths between 0 and 0.33 as a proportion of their full length. In this early, rapid regrowth phase, new OS discs are regenerated, but OS shedding does not occur. Phase 2, hyperbolic growth, occurs for cone OS lengths between 0.33 and 1 as a proportion of their full length. In this later, slower regrowth phase, new OS discs continue to be regenerated, while OS shedding is re-established. The functions λ_{1-6} are given by

$$\lambda_1(f) = H(f - f_{crit}),$$
$$\lambda_2(f) = 1 - \lambda_1(f) = 1 - H(f - f_{crit}),$$
$$\lambda_3(p_c^L) = H(0.33\tilde{p}_c^L - p_c^L),$$
$$\lambda_4(p_c^L) = 1 - \lambda_3(p_c^L) = 1 - H(0.33\tilde{p}_c^L - p_c^L),$$
$$\lambda_5(p_{c_{init}}) = 1 - H(-p_{c_{init}}),$$
$$\lambda_6(p_c^L) = 1 - H(-p_c^L),$$

where f_{crit} (M), the trophic factor threshold concentration, is a positive constant.

The functions $\lambda_1(f)$ and $\lambda_2(f)$ determine when the cone regrowth and degeneration terms are active. Cone degeneration occurs when $f < f_{crit}$ (TF starvation) in both Scenarios 1 and 2, while cone (OS) regrowth occurs when $f \geq f_{crit}$ in Scenario 2, where TF levels are sufficient to maintain cones in health. The functions $\lambda_3(p_c^L)$ and $\lambda_4(p_c^L)$ determine whether phase 1 or phase 2 cone OS regrowth is active. Phase 1 is active when $p_c^L(\theta, t) \leq 0.33\tilde{p}_c^L$, while phase 2 is active when $p_c^L(\theta, t) > 0.33\tilde{p}_c^L$. The function $\lambda_5(p_{c_{init}})$ prevents cone OS regrowth in regions from which cones are absent initially. The function $\lambda_6(p_c^L)$ halts cone OS degeneration when their length reaches zero, preventing cone OS length becoming negative.

Lastly, we close the system by imposing boundary and initial conditions. We impose zero-flux boundary conditions at both the fovea ($\theta = 0$) and ora serrata ($\theta = \Theta$):

$$\frac{\partial f}{\partial \theta}(0, t) = 0 = \frac{\partial f}{\partial \theta}(\Theta, t). \tag{5}$$

A zero-flux condition is required by symmetry at $\theta = 0$, while the zero-flux condition at $\theta = \Theta$ is justified by the assumption that TF cannot escape from the retina at the ora serrata.

We impose the following initial conditions:

$$f(\theta, 0) = f_{init}(\theta), \qquad p_r(\theta, 0) = p_{r_{init}}(\theta) = F(\theta)\tilde{p}_r(\theta),$$
$$p_c(\theta, 0) = p_{c_{init}}(\theta) = G(\theta)\tilde{p}_c(\theta), \qquad p_c^L(\theta, 0) = p_{c_{init}}^L(\theta) = G(\theta)\tilde{p}_c^L, \tag{6}$$

where $f_{init}(\theta)$ (M) is the steady-state solution to Eqs. (1) and (5) with $p_r = p_{r_{init}}(\theta)$ (photoreceptors m^{-2}), $p_c = p_{c_{init}}(\theta)$ (photoreceptors m^{-2}) and $T(\theta, 0) = 0$ (no treatment), while $\tilde{p}_r(\theta)$ (photoreceptors m^{-2}), the healthy rod distribution, and $\tilde{p}_c(\theta)$ (photoreceptors m^{-2}), the healthy cone distribution are defined as

$$\tilde{p}_r(\theta) = B_3\theta e^{-b_3\theta},$$
$$\tilde{p}_c(\theta) = B_1 e^{-b_1\theta} + B_2 e^{-b_2\theta},$$

where B_1 (photoreceptors m^{-2}), B_2 (photoreceptors m^{-2}), B_3 (photoreceptors m^{-2} rad^{-1}), b_1 (rad^{-1}), b_2 (rad^{-1}) and b_3 (rad^{-1}) are positive constants, obtained by fitting to physiological data (see Fig. 1(b) and Curcio et al., 1990). We make the simplifying assumption that healthy cone OS length, $\tilde{p}_c^L$ (m), is constant across the retina. We consider two types of initial conditions for rods and cones: healthy and containing a degenerate patch. Which of these situations obtain is determined by the functions $F(\theta)$ and $G(\theta)$, which are given by

$$F(\theta) = \begin{cases} 1 & \text{if rods healthy initially,} \\ H(\theta_{r_1} - \theta) + H(\theta - \theta_{r_2}) & \text{if rods absent from } \theta \in (\theta_{r_1}, \theta_{r_2}), \end{cases}$$

$$G(\theta) = \begin{cases} 1 & \text{if cones healthy initially,} \\ H(\theta_{c_1} - \theta) + H(\theta - \theta_{c_2}) & \text{if cones absent from } \theta \in (\theta_{c_1}, \theta_{c_2}), \end{cases}$$

where θ_{r_1} (rad) and θ_{r_2} (rad) are the left- (central) and right-hand (peripheral) boundaries of a patch of rod degeneration, and θ_{c_1} (rad) and θ_{c_2} (rad) are the left- (central) and right-hand (peripheral) boundaries of a patch of cone degeneration.

Parameter values associated with the dimensional model can be found in Table 1. See the Supplementary Materials for justification of parameter values.

Table 1: Parameters employed in the dimensional model (Eqs. (1)–(6)). Parameters for which only the source is given are taken directly from the literature, those described as 'calculated' or 'fitted' are computed using data from the literature or from simulations, those described as 'estimated' are either estimated from relevant data in the literature or on the basis of biological reasoning, and those described as 'chosen' are free for us to choose. Values are given to three significant figures. See Supplementary Materials for justification of parameter values.

Parameter	Description	Value	Source
R	Retinal radial position	1.2×10^{-2} m	Oyster (1999)
Θ	Eccentricity of the ora serrata	1.33 rad	Curcio et al. (1990)
D_f	Trophic factor diffusivity	1.73×10^{-11} m^{2}s^{-1}	Estimated using data from Jürgens et al. (1994)
α	Rate of trophic factor production by rods	1.81×10^{-17} Mm^{2}photoreceptors^{-1}s^{-1}	Estimate using data from Eden et al. (2011)
ξ	Rate of trophic factor supply from treatment	0 to 1.72×10^{-6} Ms^{-1}	Chosen
β	Rate of trophic factor consumption by cones	4.62×10^{-12} m^{2}photoreceptors^{-1}s^{-1}	Estimate using data from Eden et al. (2011)
η	Rate of trophic factor decay	5.13×10^{-5} s^{-1}	Estimated using data from Dörrbaum et al. (2018) Eden et al. (2011)
ϕ_r	Rate of mutation-induced rod degeneration	2.10×10^{-8} s^{-1}	Estimated using data from Curcio et al. (1993)
δ	Rate of trophic factor starvation-induced cone degeneration	2.87×10^{-7} s^{-1}	Estimated using data from Kocaoglu et al. (2016)
μ_1	Growth rate of cone OS in phase 1	1.60×10^{-11} m s^{-1}	Calculated using data from Guérin et al. (1993)
μ_2	Growth rate of cone OS in phase 2	3.07×10^{-12} m s^{-1}	Calculated using data from Guérin et al. (1993)
δ_L	Rate of trophic factor starvation-induced cone OS degeneration	2.34×10^{-11} m s^{-1}	Estimated using data from Kocaoglu et al. (2016)
$\tilde{p}_c^L$	Healthy cone OS length	2.93×10^{-5} m	Kocaoglu et al. (2016)
f_{crit}	Trophic factor threshold concentration	3×10^{-9} to 3×10^{-5} M	Estimate
B_1	Cone profile parameter	1.73×10^{11} photoreceptors m^{-2}	Fitted using data from Curcio et al. (1990)
B_2	Cone profile parameter	1.76×10^{10} photoreceptors m^{-2}	Fitted using data from Curcio et al. (1990)
B_3	Rod profile parameter	8.84×10^{11} photoreceptors m^{-2}rad^{-1}	Fitted using data from Curcio et al. (1990)
b_1	Cone profile parameter	54.1 rad^{-1}	Fitted using data from Curcio et al. (1990)
b_2	Cone profile parameter	2.01 rad^{-1}	Fitted using data from Curcio et al. (1990)
b_3	Rod profile parameter	2.31 rad^{-1}	Fitted using data from Curcio et al. (1990)
$\tilde{f}_A$	Mean trophic factor concentration	1×10^{-4} M	Calculated from simulations
$\tilde{p}_A$	Mean photoreceptor density	1.11×10^{11} photoreceptors m^{-2}	Calculated using data from Curcio et al. (1990)
θ_{r_1}	Position of left-hand rod degenerate patch boundary	0 to Θ rad	Chosen
θ_{r_2}	Position of right-hand rod degenerate patch boundary	0 to Θ rad	Chosen
θ_{c_1}	Position of left-hand cone degenerate patch boundary	0 to Θ rad	Chosen
θ_{c_2}	Position of right-hand cone degenerate patch boundary	0 to Θ rad	Chosen
θ_{treat_1}	Position of left-hand limit of local treatment	0 to Θ rad	Chosen
θ_{treat_2}	Position of right-hand limit of local treatment	0 to Θ rad	Chosen
t_{crit}	Time at which treatment first applied	> 0 s	Chosen

2.1. Non-dimensionalisation

We recast the dimensional model in non-dimensional form to identify the dominant terms and simplify the equations. We scale the independent and dependent variables, and initial conditions as follows:

$$\theta = \Theta\theta^*, \quad t = \frac{t^*}{\delta}, \quad f = \tilde{f}_A f^*, \quad p_r = \tilde{p}_A p_r^*, \quad p_c = \tilde{p}_A p_c^*, \quad p_c^L = \tilde{p}_c^L p_c^{L*}, \quad f_{init} = \tilde{f}_A f_{init}^*.$$

Consequently, the scaled system is posed on the domain $\theta^* \in [0, 1]$. We rescale time by the rate of cone degeneration, δ, since this is the timescale of interest. The scaling factor $\tilde{f}_A$ is the mean trophic factor concentration under healthy conditions, found by taking the mean of the solution to Eqs. (1) and (5) at steady-state, with $p_r = \tilde{p}_r(\theta)$, $p_c = \tilde{p}_c(\theta)$ and $T(\theta) = 0$ (no treatment). The scaling factor $\tilde{p}_A$ is the mean total photoreceptor density (rods and cones) under healthy conditions.

We define the following dimensionless parameters:

$$D_f^* = \frac{D_f}{R^2\Theta^2\delta}, \quad \alpha^* = \frac{\tilde{p}_A}{\delta\tilde{f}_A}\alpha, \quad \beta^* = \frac{\tilde{p}_A}{\delta}\beta, \quad \eta^* = \frac{\eta}{\delta}, \quad \phi_r^* = \frac{\phi_r}{\delta}, \quad \mu_1^* = \frac{\mu_1}{\delta\tilde{p}_c^L},$$

$$\mu_2^* = \frac{\mu_2}{\delta\tilde{p}_c^L}, \quad f_{crit}^* = \frac{f_{crit}}{\tilde{f}_A}, \quad B_1^* = \frac{B_1}{\tilde{p}_A}, \quad B_2^* = \frac{B_2}{\tilde{p}_A}, \quad B_3^* = \frac{\Theta}{\tilde{p}_A}B_3, \quad b_1^* = \Theta b_1,$$

$$b_2^* = \Theta b_2, \quad b_3^* = \Theta b_3, \quad \theta_{r_1}^* = \frac{\theta_{r_1}}{\Theta}, \quad \theta_{r_2}^* = \frac{\theta_{r_2}}{\Theta}, \quad \theta_{c_1}^* = \frac{\theta_{c_1}}{\Theta}, \quad \theta_{c_2}^* = \frac{\theta_{c_2}}{\Theta},$$

$$\xi^* = \frac{\xi}{\delta\tilde{f}_A}, \quad \theta_{treat_1}^* = \frac{\theta_{treat_1}}{\Theta}, \quad \theta_{treat_2}^* = \frac{\theta_{treat_2}}{\Theta}, \quad t_{crit}^* = \delta t_{crit}, \quad \delta_L^* = \frac{\delta_L}{\delta\tilde{p}_c^L},$$

and note that $T(\theta, t) = T^*(\theta^*, t^*)$, where $T^*(\theta^*, t^*) = H(t^* - t_{crit}^*)$ or $(1 - [H(\theta_{treat_1}^* - \theta^*) + H(\theta^* - \theta_{treat_2}^*)])H(t^* - t_{crit}^*)$ (in the global and local treatment cases respectively); $C(\theta, f, p_c) = \tilde{p}_A\tilde{f}_A C^*(\theta^*, f^*, p_c^*)$, where $C^*(\theta^*, f^*, p_c^*) = p_c^* f^*$ or $p_{c_{init}}^*(\theta^*)f^*$ (in the without cone regeneration and with cone regeneration cases respectively); $\lambda_{1,2}(f) = \lambda_{1,2}^*(f^*)$, where $\lambda_1^*(f^*) = H(f^* - f_{crit}^*)$ and $\lambda_2^*(f^*) = 1 - H(f^* - f_{crit}^*)$; $\lambda_{3,4}(p_c^L) = \lambda_{3,4}^*(p_c^{L*})$, where $\lambda_3^*(p_c^*) = H(0.33 - p_c^{L*})$ and $\lambda_4^*(p_c^*) = 1 - H(0.33 - p_c^{L*})$; $\lambda_5(p_{c_{init}}) = \lambda_5^*(p_{c_{init}}^*)$, where $\lambda_5^*(p_{c_{init}}^*) = 1 - H(-p_{c_{init}}^*)$; $\lambda_6(p_c^L) = \lambda_6^*(p_c^{L*})$, where $\lambda_6^*(p_c^{L*}) = 1 - H(-p_c^{L*})$; $\tilde{p}_r(\theta) = \tilde{p}_A\tilde{p}_r^*(\theta^*)$, where $\tilde{p}_r^*(\theta^*) = B_3^*\theta^* e^{-b_3^*\theta^*}$; $\tilde{p}_c(\theta) = \tilde{p}_A\tilde{p}_c^*(\theta^*)$, where $\tilde{p}_c^*(\theta^*) = B_1^* e^{-b_1^*\theta^*} + B_2^* e^{-b_2^*\theta^*}$; $F(\theta) = F^*(\theta^*)$, where $F^*(\theta^*) = 1$ or $H(\theta_{r_1}^* - \theta^*) + H(\theta^* - \theta_{r_2}^*)$ (in the healthy and degenerate patch cases respectively); and $G(\theta) = G^*(\theta^*)$, where $G^*(\theta^*) = 1$ or $H(\theta_{c_1}^* - \theta^*) + H(\theta^* - \theta_{c_2}^*)$ (in the healthy and degenerate patch cases respectively).

Applying the above scalings to the dimensional governing equations, boundary and initial conditions (Eqs. (1)–(6)) and dropping the stars, we obtain the following dimensionless governing equations

$$\frac{\partial f}{\partial t} = \frac{D_f}{\sin(\Theta\theta)}\frac{\partial}{\partial\theta}\left(\sin(\Theta\theta)\frac{\partial f}{\partial\theta}\right) + \alpha p_r + \xi T(\theta, t) - \beta C(\theta, f, p_c) - \eta f, \tag{7}$$

$$\frac{\partial p_r}{\partial t} = \begin{cases} 0 & \text{w/o mutation-induced rod degeneration,} \\ -\phi_r p_r & \text{with mutation-induced rod degeneration,} \end{cases} \tag{8}$$

$$\frac{\partial p_c}{\partial t} = -p_c\lambda_2(f) \qquad\qquad\qquad \text{w/o cone regeneration,} \tag{9}$$

$$\frac{\partial p_c^L}{\partial t} = \left[\mu_1\lambda_3(p_c^L) + \mu_2\left(1 - p_c^L\right)\lambda_4(p_c^L)\right]\lambda_1(f)\lambda_5(p_{c_{init}}) - \delta_L\lambda_2(f)\lambda_6(p_c^L) \quad \text{with cone regeneration,} \tag{10}$$

boundary conditions,

$$\frac{\partial f}{\partial\theta}(0, t) = 0 = \frac{\partial f}{\partial\theta}(1, t), \tag{11}$$

and initial conditions,

$$f(\theta, 0) = f_{init}(\theta), \qquad p_r(\theta, 0) = p_{r_{init}}(\theta) = F(\theta)\tilde{p}_r(\theta),$$

$$p_c(\theta, 0) = p_{c_{init}}(\theta) = G(\theta)\tilde{p}_c(\theta), \qquad p_c^L(\theta, 0) = p_{c_{init}}^L(\theta) = G(\theta), \tag{12}$$

where $f_{init}(\theta)$ is the steady-state solution to Eqs. (7) and (11) with $p_r = p_{r_{init}}(\theta)$, $p_c = p_{c_{init}}(\theta)$ and $T(\theta) = 0$ (no treatment). See Table 2 for the dimensionless parameter values.

Table 2: Parameters employed in the non-dimensional model (Eqs. (7)–(12)). Values are given to three significant figures.

Parameter	Description	Value
D_f	Trophic factor diffusivity	0.237
α	Rate of trophic factor production by rods	7.01×10^4
ξ	Rate of trophic factor supply from treatment	0 to 6×10^4
β	Rate of trophic factor consumption by cones	1.79×10^6
η	Rate of trophic factor decay	1.79×10^2
ϕ_r	Rate of mutation-induced rod degeneration	7.33×10^{-2}
μ_1	Growth rate of cone OS in phase 1	1.90
μ_2	Growth rate of cone OS in phase 2	0.365
δ_L	Rate of trophic factor starvation-induced cone OS degeneration	2.79
f_{crit}	Trophic factor threshold concentration	3×10^{-5} or 0.3
B_1	Cone profile parameter	1.56
B_2	Cone profile parameter	0.158
B_3	Rod profile parameter	10.6
b_1	Cone profile parameter	71.8
b_2	Cone profile parameter	2.67
b_3	Rod profile parameter	3.06
θ_{r_1}	Position of left-hand rod degenerate patch boundary	0 to 1
θ_{r_2}	Position of right-hand rod degenerate patch boundary	0 to 1
θ_{c_1}	Position of left-hand cone degenerate patch boundary	0 to 1
θ_{c_2}	Position of right-hand cone degenerate patch boundary	0 to 1
θ_{treat_1}	Position of left-hand limit of local treatment	0 to 1
θ_{treat_2}	Position of right-hand limit of local treatment	0 to 1
t_{crit}	Time at which treatment first applied	> 0

3. Results: steady-state problem

In this section we consider the RdCVF Eqs. (7) and (11) at steady-state. Using a combination of asymptotic analysis and numerical simulations we seek to determine the spatial extent of a patch of cone loss, consequent upon the loss of a patch of rods, and the critical RdCVF treatment rate required to prevent cone degeneration following a patch of rod loss.

3.1. Asymptotic analyses

There are a variety of scenarios to be considered:

- Rod loss only / rod and cone loss: either rods alone are removed from a patch, or rods and cones are removed from the patch, providing upper and lower bounds, respectively, on the predicted spatial extent of cone loss and critical treatment rate (see Section 3.2.1 for further details);

- Wide/narrow patch of photoreceptor loss: wide patches must be treated in an asymptotically distinct way from narrow patches (see below), and represent late and early disease stages respectively;

- No treatment / treatment: in the untreated case we are interested in the extent of cone loss and in the treatment case we are interested in the critical RdCVF treatment rate required to prevent cone loss;

– Global/local treatment: RdCVF treatment may be applied globally, across the whole retina, or locally, within the degenerate rod patch only. We are interested to see to what extent the critical treatment rate varies with global vs. local application.

We consider 8 cases (of a possible 12):

1. Wide patch of rod loss without treatment;
2. Narrow patch of rod loss without treatment;
3. Wide patch of rod and cone loss without treatment;
4. Narrow patch of rod and cone loss without treatment;
5. Wide patch of rod loss with global treatment;
6. Wide patch of rod loss with local treatment;
7. Narrow patch of rod loss with global treatment;
8. Narrow patch of rod loss with local treatment.

We do not consider rod and cone loss with treatment (4 cases) since we are only interested in calculating the upper bound on the critical treatment rate.

In the next section (3.1.1) we will use the following variables and parameters, defined here for ease of reference:

- $\tilde{p}_{c_0}(\theta) = B_2 e^{-b_2\theta}$;

- $\tilde{p}_{r_0}(\theta) = B_3 \theta e^{-b_3\theta}$;

- $c_1 := \beta \tilde{p}_{c_0}(\theta_{r_1})/D_f = \beta B_2 e^{-b_2\theta_{r_1}}/D_f$;

- $\bar{c}_1 := \beta \tilde{p}_{c_0}(\theta_{r_2})/D_f = \beta B_2 e^{-b_2\theta_{r_2}}/D_f$;

- $c_2 := \alpha \tilde{p}_{r_0}(\theta_{r_1})/D_f = \alpha B_3 \theta_{r_1} e^{-b_3\theta_{r_1}}/D_f$;

- $\bar{c}_2 := \alpha \tilde{p}_{r_0}(\theta_{r_2})/D_f = \alpha B_3 \theta_{r_2} e^{-b_3\theta_{r_2}}/D_f$.

3.1.1. Wide patch of rod loss without treatment

In this section we perform an asymptotic analysis with the aim of deriving analytical expressions for the boundaries of cone loss following the formation of a degenerate rod patch. We assume that cone loss will occur in the region for which the trophic factor concentration $f < f_{crit}$ local to the degenerate rod patch. Thus, the boundaries of the degenerate cone patch will be given as the eccentricities at which $f = f_{crit}$. To determine where this occurs, we seek a leading order solution for $f(\theta)$ to Eqs. (7) and (11) at steady-state, for prescribed $p_r(\theta) = F(\theta)\tilde{p}_r(\theta)$ and $p_c(\theta) = G(\theta)\tilde{p}_c(\theta)$, where $F(\theta) = H(\theta_{r_1} - \theta) + H(\theta - \theta_{r_2})$ and $G(\theta) = 1$.

We choose $\epsilon = O(10^{-2})$, noting that we set $\epsilon = 10^{-2}$ for all plots, scaling parameters as follows: $\alpha = \epsilon^{-2}\alpha'$, $\beta = \epsilon^{-3}\beta'$, $\eta = \epsilon^{-1}\eta'$ and $b_1 = \epsilon^{-1}b_1'$. All remaining parameters are assumed to be $O(1)$. We also scale the dependent variable $p_c = \epsilon p_c'$, and assume $f = O(1)$ and $p_r = O(1)$. This scaling on the dependent variables is valid in the region $\theta \in (\epsilon, 1]$ here and in all rod loss only cases. For completeness, we note that there are two further regions with different scalings on the dependent variables in the interval $\theta \in [0, \epsilon]$. Analysis within these regions is beyond the scope of this study.

Applying the above scalings to the steady-state version of Eqn. (7) and dropping the dashes, we obtain

$$0 = \epsilon^2 D_f \frac{d^2 f}{d\theta^2} + \epsilon^2 D_f \Theta \cot(\Theta\theta) \frac{df}{d\theta} + \alpha p_r - \beta p_c f - \epsilon \eta f, \tag{13}$$

where,

$$\tilde{p}_r(\theta) = B_3 \theta e^{-b_3 \theta}, \tag{14}$$

$$\tilde{p}_c(\theta) = B_1 e^{-\epsilon^{-1} b_1 \theta} + B_2 e^{-b_2 \theta}. \tag{15}$$

We note that $\cot(\Theta\theta) \leq O(1)$ for $\theta \in (0.1, 1]$ and $\theta_{r_2} - \theta_{r_1} \leq O(1)$. Therefore, the following analysis is valid in the region $\theta \in (0.1, 1]$.

We close the system by imposing zero-flux boundary conditions as in Eqn. (11):

$$\frac{df}{d\theta}(0) = 0 = \frac{df}{d\theta}(1). \tag{16}$$

Next, we decompose the domain into left ($\theta < \theta_{r_1}$), centre ($\theta_{r_1} < \theta < \theta_{r_2}$) and right ($\theta > \theta_{r_2}$), outer (width $O(1)$) and inner (width $O(\epsilon)$) regions as depicted in Fig. 2(a). Discontinuities are introduced into Eqn. (13) at the edges of the degenerate rod patch (θ_{r_1} and θ_{r_2}), accounting for the left/centre/right division, while boundary layers are present either side of these discontinuities, where there is a sharp transition in $f(\theta)$, and also at the ends of the domain, to satisfy the zero-flux boundary conditions. In the following analysis we are not interested in the behaviour of the solution within the boundary layers at the ends of the domain. We label the remaining regions as follows from left to right: left-outer, left-inner, left-centre-inner, centre-outer, right-centre-inner, right-inner and right-outer (see Fig. 2(a)(ii)).

In what follows we seek leading order ($O(1)$) solutions to $f(\theta)$ in each of the outer and inner regions. This results in a number of unknown constants of integration, whose values are determined by a combination of matching and patching. Asymptotic matching ensures that the inner limit of the outer solutions match with the outer limit of the inner solutions where they meet, while patching ensures continuity of trophic factor concentration and flux across the discontinuities at θ_{r_1} and θ_{r_2} (Bender and Orszag, 1999).

We form the regular perturbation expansions:

$$f(\theta) = f_0(\theta) + \epsilon f_1(\theta) + O(\epsilon^2), \tag{17}$$

$$p_r(\theta) = p_{r_0}(\theta) + \epsilon p_{r_1}(\theta) + O(\epsilon^2), \tag{18}$$

$$p_c(\theta) = p_{c_0}(\theta) + \epsilon p_{c_1}(\theta) + O(\epsilon^2), \tag{19}$$

and seek $f_0(\theta)$, where $p_{r_0}(\theta) = F(\theta)\tilde{p}_{r_0}(\theta) = F(\theta)B_3\theta e^{-b_3\theta}$ and $p_{c_0}(\theta) = G(\theta)\tilde{p}_{c_0}(\theta) = G(\theta)B_2 e^{-b_2\theta}$.

Seeking leading order solutions to (13) in the left- and right-outer regions, we obtain,

$$f_0(\theta) = \frac{\alpha B_3 \theta e^{-b_3\theta}}{\beta B_2 e^{-b_2\theta}},$$

while, in the centre-outer region, we obtain,

$$f_0(\theta) = 0.$$

To find the leading order solution in the left-inner and left-centre-inner regions, we rescale the independent variable θ as $\hat{\theta} = \epsilon^{-1}(\theta - \theta_{r_1})$, such that the first diffusion term enters the dominant balance. Thus, Eqn. (13) becomes

$$0 = D_f \frac{d^2 f}{d\hat{\theta}^2} + \epsilon D_f \Theta \cot(\Theta(\theta_{r_1} + \epsilon\hat{\theta})) \frac{df}{d\hat{\theta}} + \alpha p_r - \beta p_c f - \epsilon \eta f. \tag{20}$$

Thus we obtain leading order solutions in the left-inner region,

$$f(\hat{\theta}) = A_1 e^{\sqrt{c_1}\hat{\theta}} + A_2 e^{-\sqrt{c_1}\hat{\theta}} + \frac{c_2}{c_1}, \tag{21}$$

Figure 2: Inner and outer region locations for the asymptotic analysis. (a) wide (width $O(1)$) patch of rod loss, (b) narrow (width $O(\epsilon)$) patch of rod loss, (c) wide (width $O(1)$) patch of rod and cone loss, (d) narrow (width $O(\epsilon^{1/2})$) patch of rod and cone loss. The full domain is shown in the left-hand column, while the right-hand column magnifies the region from which photoreceptors have been removed. The green dashed vertical lines mark the boundaries of rod (and cone) loss, while the solid magenta vertical lines demarcate the approximate limits of the boundary layers. To obtain $f(\theta)$, Eqs. (7) and (11) were solved using the finite difference method, with 4001 mesh points, where $p_r(\theta) = F(\theta)\tilde{p}_r(\theta)$, $p_c(\theta) = G(\theta)\tilde{p}_c(\theta)$, $F(\theta) = H(\theta_{r_1} - \theta) + H(\theta - \theta_{r_2})$, $G(\theta) = 1$ in (a) and (b), and $G(\theta) = H(\theta_{c_1} - \theta) + H(\theta - \theta_{c_2})$ in (c) and (d), and without treatment, cone regeneration or mutation-induced rod degeneration. Parameter values: $\xi = 0$; (a) $(\theta_{r_1}, \theta_{r_2}) = (0.4, 0.6)$; (b) $(\theta_{r_1}, \theta_{r_2}) = (0.49, 0.51)$; (c) $(\theta_{r_1}, \theta_{r_2}) = (\theta_{c_1}, \theta_{c_2}) = (0.2, 0.8)$; and (d) $(\theta_{r_1}, \theta_{r_2}) = (\theta_{c_1}, \theta_{c_2}) = (0.4, 0.6)$. Remaining parameter values as in Table 2.

and in the left-centre-inner region,

$$f(\hat{\theta}) = A_3 e^{\sqrt{c_1}\hat{\theta}} + A_4 e^{-\sqrt{c_1}\hat{\theta}}, \tag{22}$$

where $c_1 = \beta\tilde{p}_{c_0}(\theta_{r_1})/D_f = \beta B_2 e^{-b_2\theta_{r_1}}/D_f$, $c_2 = \alpha\tilde{p}_{r_0}(\theta_{r_1})/D_f = \alpha B_3 \theta_{r_1} e^{-b_3\theta_{r_1}}/D_f$, and A_1–A_4 are constants of integration to be determined.

To find the leading order solution in the right-centre-inner and right-inner regions, we rescale the independent variable θ as $\bar{\theta} = \epsilon^{-1}(\theta - \theta_{r_2})$, obtaining the equivalent equation to Eqn. (20), but with $\hat{\theta} \mapsto \bar{\theta}$ and $\theta_{r_1} \mapsto \theta_{r_2}$.

This gives us leading order solutions in the right-centre-inner region,

$$f(\bar{\theta}) = A_5 e^{\sqrt{\bar{c}_1}\bar{\theta}} + A_6 e^{-\sqrt{\bar{c}_1}\bar{\theta}}, \tag{23}$$

and in the right-inner region,

$$f(\bar{\theta}) = A_7 e^{\sqrt{\bar{c}_1}\bar{\theta}} + A_8 e^{-\sqrt{\bar{c}_1}\bar{\theta}} + \frac{\bar{c}_2}{\bar{c}_1}, \tag{24}$$

where $\bar{c}_1 = \beta\tilde{p}_{c_0}(\theta_{r_2})/D_f = \beta B_2 e^{-b_2\theta_{r_2}}/D_f$, $\bar{c}_2 = \alpha\tilde{p}_{r_0}(\theta_{r_2})/D_f = \alpha B_3 \theta_{r_2} e^{-b_3\theta_{r_2}}/D_f$, and A_5–A_8 are constants of integration to be determined.

To obtain values for A_1–A_8 we first match and then patch. We match using Van Dyke's matching rule, which states that the m term inner expansion of the n term outer solution, should equal the n term outer expansion of the m term inner solution. In this case, since we are dealing with leading order solutions only, $m = n = 1$. Matching the left-outer and left-inner solutions, we form the one term inner expansion of the one term outer solution,

$$f(\hat{\theta}) \sim \frac{\alpha B_3 \theta_{r_1} e^{-b_3\theta_{r_1}}}{\beta B_2 e^{-b_2\theta_{r_1}}} = \frac{c_2}{c_1},$$

and the one term outer expansion of one term inner solution,

$$f(\theta) \sim A_1 e^{\sqrt{c_1}\epsilon^{-1}(\theta-\theta_{r_1})} + A_2 e^{-\sqrt{c_1}\epsilon^{-1}(\theta-\theta_{r_1})} + \frac{c_2}{c_1}.$$

Since, $\theta - \theta_{r_1} > 0$ in the left-hand regions, the first term is exponentially small and so can be neglected, while the second term is exponentially large and so we must have $A_2 = 0$. Matching the left-centre-inner and centre-outer solutions, the centre-outer and right-centre-inner solutions, and the right-inner and right-outer solutions in a similar fashion, we obtain $A_3 = 0$, $A_6 = 0$, and $A_7 = 0$ respectively.

Patching across $\theta = \theta_{r_1}$ and $\theta = \theta_{r_2}$, we obtain expression for A_1, A_4, A_5 and A_8 in terms of known parameters. Substituting for A_1–A_8 into Eqs. (21)–(24), we obtain the full set of leading order outer and inner solutions as follows —

Left- and Right-Outer:

$$f_0(\theta) = \frac{\alpha B_3 \theta e^{-b_3\theta}}{\beta B_2 e^{-b_2\theta}}, \tag{25}$$

Left-Inner:

$$f_0(\theta) = \frac{c_2}{c_1}\left(1 - \frac{1}{2}e^{\epsilon^{-1}\sqrt{c_1}(\theta-\theta_{r_1})}\right), \tag{26}$$

Left-Centre-Inner:

$$f_0(\theta) = \frac{c_2}{2c_1} e^{-\epsilon^{-1} \sqrt{c_1}(\theta-\theta_{r_1})}, \tag{27}$$

Centre-Outer:

$$f_0(\theta) = 0, \tag{28}$$

Right-Centre-Inner:

$$f_0(\theta) = \frac{\bar{c}_2}{2\bar{c}_1} e^{\epsilon^{-1} \sqrt{\bar{c}_1}(\theta-\theta_{r_2})}, \tag{29}$$

Right-Inner:

$$f_0(\theta) = \frac{\bar{c}_2}{\bar{c}_1} \left(1 - \frac{1}{2} e^{-\epsilon^{-1} \sqrt{\bar{c}_1}(\theta-\theta_{r_2})} \right). \tag{30}$$

Next we calculate leading order composite solutions, providing uniformly valid expansions across the left-outer and left-inner regions (left-composite); across the left-centre-inner, centre-outer and right-centre-inner regions (centre-composite), and across the right-inner and right-outer regions (right-composite). Composite solutions are calculated by adding together the relevant outer and inner solutions and subtracting one copy of the common terms to avoid double counting (see Bender and Orszag, 1999, for further details). We obtain the following composite solutions —

Left-Composite:

$$f_{0_{left-comp}}(\theta) = \frac{\alpha B_3 \theta e^{-b_3 \theta}}{\beta B_2 e^{-b_2 \theta}} - \frac{c_2}{2c_1} e^{\epsilon^{-1} \sqrt{c_1}(\theta-\theta_{r_1})}, \tag{31}$$

Centre-Composite:

$$f_{0_{centre-comp}}(\theta) = \frac{c_2}{2c_1} e^{-\epsilon^{-1} \sqrt{c_1}(\theta-\theta_{r_1})} + \frac{\bar{c}_2}{2\bar{c}_1} e^{\epsilon^{-1} \sqrt{\bar{c}_1}(\theta-\theta_{r_2})}, \tag{32}$$

Right-Composite:

$$f_{0_{right-comp}}(\theta) = \frac{\alpha B_3 \theta e^{-b_3 \theta}}{\beta B_2 e^{-b_2 \theta}} - \frac{\bar{c}_2}{2\bar{c}_1} e^{-\epsilon^{-1} \sqrt{\bar{c}_1}(\theta-\theta_{r_2})}, \tag{33}$$

Lastly, we derive expressions for the cone degenerate patch boundaries. As discussed above, we assume that the boundaries of a cone degenerate patch, θ_{crit_1} (left-hand boundary) and θ_{crit_2} (right-hand boundary), occur where $f(\theta) = f_{crit}$. This cannot occur in the outer regions, since we have chosen f_{crit} such that $f(\theta) > f_{crit}$ in the healthy retina (left- and right-outer regions), while $f_0(\theta) = 0 < f_{crit}$ in the centre-outer region. Therefore, we must have $f(\theta) = f_{crit}$ somewhere in the left-inner or left-centre-inner region (θ_{crit_1}), and somewhere in the right-centre-inner or right-inner region (θ_{crit_2}). Setting $f(\theta) = f_{crit}$ in Eqs. (26), (27), (29) and (30), and rearranging, we obtain —

Minimal Cone Degenerate Patch Left Boundary:

$$\theta_{crit_1} = \theta_{r_1} + \text{sgn}\left(f_{crit} - \frac{c_2}{2c_1} \right) \frac{\epsilon}{\sqrt{c_1}} \log\left(1 - \left| \frac{2c_1 f_{crit}}{c_2} - 1 \right| \right), \tag{34}$$

Maximal Cone Degenerate Patch Right Boundary:

$$\theta_{crit_2} = \theta_{r_2} - \text{sgn}\left(f_{crit} - \frac{\bar{c}_2}{2\bar{c}_1}\right) \frac{\epsilon}{\sqrt{\bar{c}_1}} \log\left(1 - \left|\frac{2\bar{c}_1 f_{crit}}{\bar{c}_2} - 1\right|\right), \tag{35}$$

where sgn($\cdot$) is the signum (or sign) function, defined such that,

$$\text{sgn}(x) := \begin{cases} -1 & \text{if } x < 0, \\ 0 & \text{if } x = 0, \\ 1 & \text{if } x > 0. \end{cases}$$

We have combined the expressions for θ_{crit_1} resulting from Eqs. (26) and (27), to give Eqn. (34), and similarly for θ_{crit_2} with Eqs. (29) and (30) to give Eqn. (35). The position of the cone degenerate patch boundaries relative to the rod degenerate patch boundaries, and hence the question of whether the cone degenerate patch lies within or exceeds the rod degenerate patch, depends upon the value of f_{crit}, and is given by the following inequalities —

- when $f_{crit} \geq c_2/(2c_1)$, $\theta_{crit_1} \leq \theta_{r_1}$;

- when $f_{crit} < c_2/(2c_1)$, $\theta_{crit_1} > \theta_{r_1}$;

- when $f_{crit} \geq \bar{c}_2/(2\bar{c}_1)$, $\theta_{crit_2} \geq \theta_{r_2}$;

- when $f_{crit} < \bar{c}_2/(2\bar{c}_1)$, $\theta_{crit_2} < \theta_{r_2}$.

Thus, as would be expected intuitively, cone degenerate patches are wider for larger values of f_{crit}.

For asymptotic analysis Cases 2–8, see Appendix A.

3.2. Numerical and analytical solutions

In this section we compare numerical solutions and analytical approximations to the steady-state problem in which rods (or rods and cones) have been lost from the interval $\theta \in (\theta_{r_1}, \theta_{r_2})$, with and without treatment. Analytical solutions provide deeper insight into the dominant mechanisms determining the behaviour of our models and are cheap to compute, while numerical solutions serve to confirm their accuracy and are more computationally expensive. To obtain numerical solutions we solve Eqs. (7) and (11) for $f(\theta)$ at steady-state using the Matlab routine fsolve, which employs a Trust–Region–Dogleg algorithm. Analytical approximations are derived in Section 3.1.

We consider the 8 cases described in Section 3.1, dividing each of these cases into the two subcases:

1. Low (homogeneous) RdCVF threshold concentration $f_{crit} = 3 \times 10^{-5}$: chosen such that $f_{crit} < f(\theta)$ for all $\theta \in [0, 1]$ in the healthy retina (where $p_r(\theta) = \tilde{p}_r(\theta)$ and $p_c(\theta) = \tilde{p}_c(\theta)$);
2. High (heterogeneous) RdCVF threshold concentration $f_{crit} = f_{crit}(\theta) = f_{crit_1} H(0.13 - \theta) + f_{crit_2}(1 - H(0.13 - \theta))$: where $f_{crit_1} = 3 \times 10^{-5}$, $f_{crit_2} = 0.3$ and the switch point, $\theta = 0.13$, are chosen such that $f_{crit}(\theta) < f(\theta)$ for all $\theta \in [0, 1]$ in the healthy retina.

The value of f_{crit} has not yet been measured; therefore, we consider the two most likely scenarios. In the first (low f_{crit}) scenario, the threshold is spatially uniform and chosen to be low enough such that cones will not degenerate in the healthy retina. In the second (high f_{crit}) scenario, we assume that the cone-rich fovea is afforded special protection against low RdCVF levels, such that f_{crit} is lower in that region ($\theta \leq 0.13$) than in the rest of the retina, where rods, and hence RdCVF, are more plentiful and cone density, and hence the rate of RdCVF uptake, is low. For notational simplicity, we shall refer to the high f_{crit} scenario simply as $f_{crit} = 0.3$ in what follows. Note that throughout this section, we consider only region $\theta \in (0.13, 1]$ in the high f_{crit} scenario.

3.2.1. No treatment

We begin by considering the untreated case, in which $\xi = 0$. Here we are interested in predicting the interval over which cones will degenerate following the removal of a patch of rods from the interval $\theta \in (\theta_{r_1}, \theta_{r_2})$, and the conditions under which cone degeneration will expand beyond the rod degenerate patch. The boundaries of the resultant patch of cone loss are calculated as the positions at which $f(\theta) = f_{crit}$ (between which $f(\theta) < f_{crit}$) local to the patch of rod loss. In the full dynamic problem (see Section 4.1), $f(\theta)$ will rise as cones degenerate, shrinking the interval over which $f(\theta) < f_{crit}$. Here, we derive upper and lower bounds on the cone degenerate patch width. The upper bound is found in the case where only rods are removed, such that cones continue to consume RdCVF within the degenerate patch, expanding the interval over which $f(\theta) < f_{crit}$. The lower bound is found in the case where both rods and cones are removed from $\theta \in (\theta_{r_1}, \theta_{r_2})$ (where $\theta_{c_1} = \theta_{r_1}$ and $\theta_{c_2} = \theta_{r_2}$), such that cones do not consume RdCVF within the degenerate patch, reducing the interval over which $f(\theta) < f_{crit}$. We note that, in calculating the lower bound this way, we assume that cones do not degenerate beyond the rod degenerate patch. Where cone loss does exceed the rod patch, this method acts as a confirmation that this will occur, rather than as a lower bound.

Fig. 3 shows analytical solutions for the distance between the left-hand rod and cone degenerate patch boundaries, $\theta_{r_1} - \theta_{crit_1}$, as a function of θ_{r_1}, and similarly for the distance between the right-hand rod and cone degenerate patch boundaries, $\theta_{crit_2} - \theta_{r_2}$, as a function of θ_{r_2}. We consider the rod loss only ((a) and (b)), and rod and cone loss cases ((c) and (d)), corresponding to the upper and lower bounds on cone degenerate patch width respectively. For each of these cases, we consider the subcases $f_{crit} = 0.3$ ((a) and (c)) and $f_{crit} = 3 \times 10^{-5}$ ((b) and (d)).

The equations plotted in Fig. 3 ($\theta_{r_1} - \theta_{crit_1}$ and $\theta_{crit_2} - \theta_{r_2}$, where θ_{crit_1} and θ_{crit_2} are as given in Eqs. (34) and (35) in (a) and (b), and Eqs. (A.21)–(A.24) in (c) and (d)) are all monotone decreasing functions of θ_{r_1} and θ_{r_2}. However, θ_{crit_1} and θ_{crit_2} cannot in practice go outside of the retina and hence cannot exceed the interval $\theta \in [0, 1]$. Therefore, we set $\theta_{crit_1} = 1$ where $\theta_{crit_1} > 1$ and $\theta_{crit_2} = 0$ where $\theta_{crit_2} < 1$. This results in upward jumps on the right-hand side of the domain in $\theta_{r_1} - \theta_{crit_1}$ in (a)–(d) and an upward jump on the left-hand side of the domain in $\theta_{crit_2} - \theta_{r_2}$ in (d). Apart from these jumps, the curves for $\theta_{r_1} - \theta_{crit_1}$ and $\theta_{crit_2} - \theta_{r_2}$ are identical. Note that, for the wide patch case, the predicted position of one cone degenerate patch boundary does not depend upon the other cone degenerate patch boundary (unlike in the narrow patch case), since the boundary layers surrounding the rod degenerate patch boundaries are separated by a central outer region, such that they do not 'interact'. The maximum spatial extent of cone loss remains within the boundaries of rod loss in (b) and (d), but may exceed it close to the fovea (centred at $\theta = 0$) in (a) and (c).

Fig. 4 shows numerical and analytical bounds on cone degenerate patch size for the wide rod degenerate patch case. The cone degenerate patch width, $\theta_{crit_2} - \theta_{crit_1}$, increases monotonically with increasing rod degenerate patch width, $\theta_{r_2} - \theta_{r_1}$, and with decreasing rod degenerate patch centre position, $(\theta_{r_1} + \theta_{r_2})/2$. For narrower rod loss patches in (c), and especially (d), the predicted lower bound on cone degenerate patch width is zero. The difference between rod and cone degenerate patch widths, $(\theta_{crit_2} - \theta_{crit_1}) - (\theta_{r_2} - \theta_{r_1})$, is a monotone (not strictly) decreasing function of rod degenerate patch centre position. The cone degenerate patch width is less than the rod degenerate patch width except in (a) (rod loss only with $f_{crit} = 0.3$), where it may exceed rod degenerate patch width close to the fovea, doing so for a wider range of patch positions for narrower rod loss patches. The predicted cone degenerate patch width is lower for the rod and cone loss case ((a) and (b), lower bound) than for the rod loss only case ((c) and (d), upper bound) as would be expected. Numerical and analytical results match closely, except for the width 0.1 rod and cone degenerate patch case in (c), where the wide patch assumption breaks down, since $0.1 \sim O(\epsilon^{1/2})$.

Fig. 5 shows numerical and analytical bounds on cone degenerate patch size for the narrow rod degenerate patch case. We plot only the $f_{crit} = 0.3$ subcase here since, for the $f_{crit} = 3 \times 10^{-5}$ subcase, the predicted cone loss patch width is always zero. As in Fig. 4, cone degenerate patch width increases monotonically with

Figure 3: Distance between rod and minimum/maximum cone degenerate patch boundaries — wide rod degenerate patch case. Each panel shows the variation in the distances between the left-hand rod and minimum cone degenerate patch boundaries, $\theta_{r_1} - \theta_{crit_1}$, and between the right-hand rod and maximum cone degenerate patch boundaries, $\theta_{crit_2} - \theta_{r_2}$, with θ_{r_1} and θ_{r_2} respectively. (a) rod loss only with $f_{crit} = 0.3$, (b) rod loss only with $f_{crit} = 3 \times 10^{-5}$, (c) rod and cone loss with $f_{crit} = 0.3$, (d) rod and cone loss with $f_{crit} = 3 \times 10^{-5}$. Note the different scales on the y-axes. The maximum spatial extent of cone loss remains within the boundaries of rod loss in (b) and (d), but may exceed it close to the fovea (centred at $\theta = 0$) in (a) and (c). Curves are plotted using Eqs. (34) and (35) in (a) and (b), and using Eqs. (A.21)–(A.24) in (c) and (d). The parameter $\epsilon = 10^{-2}$. Remaining parameter values as in Table 2.

Figure 4: Numerical and analytical bounds on cone degenerate patch size using the steady-state model — wide rod degenerate patch case. (a)–(d) first row: maximum/minimum cone degenerate patch width, $\theta_{crit_2} - \theta_{crit_1}$; second row: difference between rod and maximum/minimum cone degenerate patch widths, $(\theta_{crit_2} - \theta_{crit_1}) - (\theta_{r_2} - \theta_{r_1})$; columns 1 and 2: variation of patch widths and width differences in $(\theta_{r_1}, \theta_{r_2})$ parameter space; columns 3–5: variation of patch widths and width differences with rod degenerate patch centre position, $(\theta_{r_1} + \theta_{r_2})/2$, each curve representing a constant rod degenerate patch width, $\theta_{r_2} - \theta_{r_1}$; columns 1 and 3: numerical solutions; columns 2 and 5: analytical approximations; column 4: numerical (solid curves) and analytical (dashed curves) solutions compared. (a) rod loss only with $f_{crit} = 0.3$, (b) rod loss only with $f_{crit} = 3 \times 10^{-5}$, (c) rod and cone loss with $f_{crit} = 0.3$, (d) rod and cone loss with $f_{crit} = 3 \times 10^{-5}$. (a) and (b) upper bounds on cone degenerate patch widths; (c) and (d) lower bounds on cone degenerate patch widths. Cone degenerate patch width increases monotonically with increasing rod degenerate patch width and with decreasing rod degenerate patch centre position. Cone degenerate patch width is less than rod degenerate patch width except in (a), where it may exceed rod degenerate patch width close to the fovea (centred at $\theta = 0$). Numerical solutions were obtained by calculating the eccentricities at which $f(\theta) = f_{crit}$, solving Eqs. (7) and (11) for $f(\theta)$ at steady-state using the finite difference method, with 4001 mesh points, where $p_r(\theta) = F(\theta)\tilde{p}_r(\theta)$, $p_c(\theta) = G(\theta)\tilde{p}_c(\theta)$, $F(\theta) = H(\theta_{r_1} - \theta) + H(\theta - \theta_{r_2})$, $G(\theta) = 1$ in (a) and (b), and $G(\theta) = H(\theta_{c_1} - \theta) + H(\theta - \theta_{c_2})$, with $\theta_{c_1} = \theta_{r_1}$ and $\theta_{c_2} = \theta_{r_2}$, in (c) and (d), and without treatment or cone regeneration. Analytical solutions are plotted using Eqs. (34) and (35) in (a) and (b), and using Eqs. (A.21)–(A.24) in (c) and (d). Parameter values: $\xi = 0$ and $\epsilon = 10^{-2}$. Remaining parameter values as in Table 2.

increasing rod degenerate patch width and with decreasing rod degenerate patch centre position, while the difference between rod and cone degenerate patch widths is a monotone (not strictly) decreasing function of rod degenerate patch centre position. The cone degenerate patch width may exceed the rod degenerate patch width close to the fovea in (a). Numerical and analytical solutions agree well in (a) and on the right-hand side in (b); however, the scaling assumptions break down close to the fovea, such that the analytical solution predicts cone loss on left-hand side in (b), where the numerical solution remains zero.

3.2.2. Treatment

In this section we compare numerical and analytical estimates for the critical treatment rate, ξ_{crit}. This is the minimum RdCVF treatment rate required to prevent cone degeneration in a given scenario. We consider just the rod loss only case, since this provides an upper bound on ξ_{crit}, which is of greater clinical relevance, given that this would theoretically guarantee the avoidance of cone degeneration under the trophic factor hypothesis. We also consider analytical estimates for the eccentricity of the minimum trophic factor concentration local to the rod degenerate patch, θ_{crit}, when $\xi = \xi_{crit}$. This is the point at which cone loss will initiate if ξ drops below ξ_{crit}. Numerical solutions are obtained using the Matlab routine fminsearch (which uses a simplex search method) to find the value of ξ at which the minimum value of $f(\theta)$ local to the rod degenerate patch is equal to f_{crit}. This involves solving Eqs. (7) and (11) for $f(\theta)$ at steady-state using fsolve at each iteration. Analytical solutions are obtained by employing the Matlab routine fsolve to solve the implicit equations (A.46) and (A.47) for a wide patch of rod loss with global treatment, (A.59) and (A.60) for a wide patch of rod loss with local treatment, (A.71) and (A.72) for a narrow patch of rod loss with global treatment, and (A.83) and (A.84) for a narrow patch of rod loss with local treatment.

Fig. 6 shows numerical and analytical estimates of the critical treatment rate for the wide rod degenerate patch case. The critical treatment rate increases monotonically with decreasing rod degenerate patch centre position and left boundary position in all cases. The critical treatment rate depends almost exclusively upon the rod degenerate patch left boundary position, rather than the rod degenerate patch width or right boundary position. Numerical and analytical solutions agree well.

Fig. 7 shows numerical and analytical estimates of the critical treatment rate for the narrow rod degenerate patch case. We consider only the $f_{crit} = 0.3$ subcase here, since there is no cone loss for $f_{crit} = 3 \times 10^{-5}$ and hence no treatment is required in this subcase. As with the wide rod degenerate patch case, the critical treatment rate increases monotonically with decreasing rod degenerate patch centre position and left boundary position in all cases. Unlike in the wide rod degenerate patch case, the critical treatment rate depends upon both the rod degenerate patch left and right boundary positions, and hence upon the patch width (compare Fig. 6), increasing monotonically with increasing rod degenerate patch width. Numerical and analytical solutions match closely.

Fig. 8 compares analytical estimates of the critical treatment rate for the global and local treatment cases. The critical treatment rate is almost identical for global and local treatment in the wide patch case, for both $f_{crit} = 0.3$ (a) and $f_{crit} = 3 \times 10^{-5}$ (b), while the critical treatment rate is higher for local than for global treatment for the narrowest patches close to the fovea (centred at $\theta = 0$) in the narrow patch case (with $f_{crit} = 0.3$) (c).

Fig. 9 shows analytical estimates of the normalised eccentricity of the minimum trophic factor concentration, $\hat{\theta}_{crit}$, for both the wide and narrow rod degenerate patch cases. The variable $\hat{\theta}_{crit}$ increases monotonically with decreasing rod degenerate patch width, and with increasing rod degenerate patch centre position and left boundary position in all cases. Further, $\hat{\theta}_{crit}$ remains close to 0 for patches of width $\theta_{r_2} - \theta_{r_1} \geq 0.2$ ((a)–(d)) and close to 0.5 for narrow patches ((e) and (f)).

Figure 5: Numerical and analytical bounds on cone degenerate patch size using the steady-state model — narrow rod degenerate patch case. Graphs show variation of maximum/minimum cone degenerate patch widths, and differences between rod and maximum/minimum cone degenerate patch widths with rod degenerate patch centre position, $(\theta_{r_1} + \theta_{r_2})/2$, each curve representing a constant rod degenerate patch width, $\theta_{r_2} - \theta_{r_1}$. (a) and (b) first row: maximum/minimum cone degenerate patch width, $\theta_{crit_2} - \theta_{crit_1}$; second row: difference between rod and maximum/minimum cone degenerate patch widths, $(\theta_{crit_2} - \theta_{crit_1}) - (\theta_{r_2} - \theta_{r_1})$; column 1: numerical solutions; column 3: analytical approximations; column 2: numerical (solid curves) and analytical (dashed curves) solutions compared. (a) rod loss only: upper bounds on cone degenerate patch widths; (b) rod and cone loss: lower bounds on cone degenerate patch widths. Cone degenerate patch width increases monotonically with increasing rod degenerate patch width and with decreasing rod degenerate patch centre position. Cone degenerate patch width may exceed rod degenerate patch width close to the fovea (centred at $\theta = 0$). Numerical solutions were obtained by calculating the eccentricities at which $f(\theta) = f_{crit}$, solving Eqs. (7) and (11) for $f(\theta)$ at steady-state using the finite difference method, with 4001 mesh points, where $p_r(\theta) = F(\theta)\tilde{p}_r(\theta)$, $p_c(\theta) = G(\theta)\tilde{p}_c(\theta)$, $F(\theta) = H(\theta_{r_1} - \theta) + H(\theta - \theta_{r_2})$, $G(\theta) = 1$ in (a), and $G(\theta) = H(\theta_{c_1} - \theta) + H(\theta - \theta_{c_2})$, with $\theta_{c_1} = \theta_{r_1}$ and $\theta_{c_2} = \theta_{r_2}$, in (b), and without treatment or cone regeneration. Analytical solutions are plotted using Eqs. (A.7)–(A.10) in (a), and using Eqs. (A.31)–(A.34) in (b). Parameter values: $f_{crit} = 0.3$, $\xi = 0$ and $\epsilon = 10^{-2}$. Remaining parameter values as in Table 2.

Figure 6: Numerical and analytical estimates of the critical treatment rate using the steady-state model — wide rod degenerate patch case. Rod loss only. All plots show the critical treatment rate, ξ_{crit}. (a) and (c) global treatment, (b) and (d) local treatment; (a) and (b) $f_{crit} = 0.3$; (c) and (d) $f_{crit} = 3\times10^{-5}$. (a)–(d) column 1: numerical (top) and analytical (bottom) estimates of the variation in the critical treatment rate over $(\theta_{r_1}, \theta_{r_2})$ parameter space; columns 2–4: variation in the critical treatment rate with rod degenerate patch centre position, $(\theta_{r_1} + \theta_{r_2})/2$ (top), and with rod degenerate patch left boundary position, θ_{r_1} (bottom), each curve representing a constant rod degenerate patch width, $\theta_{r_2} - \theta_{r_1}$; column 2: numerical solutions; column 4: analytical approximations; column 3: numerical (solid curves) and analytical (dashed curves) solutions compared. The critical treatment rate increases monotonically with decreasing rod degenerate patch centre/left boundary position in all cases. The critical treatment rate depends almost entirely upon the rod degenerate patch left boundary position, rather than the rod degenerate patch width/right boundary position. Numerical solutions were obtained by using the Matlab routine fminsearch to calculate the critical treatment rate at which $\min(f(\theta)) = f_{crit}$ within $\theta \in [\theta_{r_1}, \theta_{r_2}]$, solving Eqs. (7) and (11) for $f(\theta)$ at steady-state at each iteration using the finite difference method, with 401 mesh points, where $p_r(\theta) = F(\theta)\tilde{p}_r(\theta)$, $p_c(\theta) = G(\theta)\tilde{p}_c(\theta)$, $F(\theta) = H(\theta_{r_1} - \theta) + H(\theta - \theta_{r_2})$, $G(\theta) = 1$, $T(\theta) = 1$ for (a) and (c), and $T(\theta) = 1 - F(\theta)$ for (b) and (d). Analytical solutions are obtained by implicitly solving Eqs. (A.46) and (A.47) in (a) and (c), and Eqs. (A.59) and (A.60) in (b) and (d). The parameter $\epsilon = 10^{-2}$. Remaining parameter values as in Table 2.

Figure 7: Numerical and analytical estimates of the critical treatment rate using the steady-state model — narrow rod degenerate patch case. Rod loss only. All plots show the critical treatment rate, ξ_{crit}. (a) global treatment, (b) local treatment. (a) and (b) variation in the critical treatment rate with rod degenerate patch centre position, $(\theta_{r_1} + \theta_{r_2})/2$ (top), and with rod degenerate patch left boundary position, θ_{r_1} (bottom), each curve representing a constant rod degenerate patch width, $\theta_{r_2} - \theta_{r_1}$; column 1: numerical solutions; column 3: analytical approximations; column 2: numerical (solid curves) and analytical (dashed curves) solutions compared. The critical treatment rate depends upon both the rod degenerate patch left and right boundary positions, and hence the patch width (compare Fig. 6). The critical treatment rate increases monotonically with increasing rod degenerate patch width and with decreasing patch centre/left boundary position in all cases. Numerical solutions were obtained by using the Matlab routine fminsearch to calculate the critical treatment rate at which $\min(f(\theta)) = f_{crit}$ within $\theta \in [\theta_{r_1}, \theta_{r_2}]$, solving Eqs. (7) and (11) for $f(\theta)$ at steady-state at each iteration using the finite difference method, with 1001 mesh points, where $p_r(\theta) = F(\theta)\tilde{p}_r(\theta)$, $p_c(\theta) = G(\theta)\tilde{p}_c(\theta)$, $F(\theta) = H(\theta_{r_1} - \theta) + H(\theta - \theta_{r_2})$, $G(\theta) = 1$, $T(\theta) = 1$ for (a) and $T(\theta) = 1 - F(\theta)$ for (b). Analytical solutions are obtained by implicitly solving Eqs. (A.71) and (A.72) in (a) and Eqs. (A.83) and (A.84) in (b). Parameter values: $f_{crit} = 0.3$ and $\epsilon = 10^{-2}$. Remaining parameter values as in Table 2.

Figure 8: Comparison of critical treatment rates under global and local treatment — wide and narrow rod degenerate patch cases. All plots show analytical estimates of the variation in critical treatment rate, ξ_{crit}, with rod degenerate patch left boundary position, θ_{r_1}, using the steady-state model. Each curve represents a constant rod degenerate patch width, $\theta_{r_2} - \theta_{r_1}$. Solid curves: global treatment; dashed curves: local treatment. (a) wide rod degenerate patch with $f_{crit} = 0.3$, (b) wide patch with $f_{crit} = 3 \times 10^{-5}$, (c) narrow patch with $f_{crit} = 0.3$. The critical treatment rate is almost identical for global and local treatment in the wide patch case, while the critical treatment rate is higher for local than for global treatment for the narrowest patches close to the fovea (centred at $\theta = 0$) in the narrow patch case. Analytical solutions and parameter values are identical to those for the corresponding cases in Figs. 6 and 7.

4. Results: dynamic problem

Having explored the steady-state problem, we now consider the full dynamic problem, consisting of Eqs. (7)–(9) and (11)–(12) in the no cone regeneration/no treatment case, and Eqs. (7), (8) and (10)–(12) in the cone regeneration/treatment case. In both cases, equations are solved using the method of lines. This involves discretising in space and then integrating in time using the Matlab routine ode15s, a variable-step, variable-order solver. The initial trophic factor profile, $f(\theta, 0) = f_{init}(\theta)$, is calculated by solving Eqs. (7) and (11) at steady-state using the Matlab routine fsolve with $p_r = p_{r_{init}}(\theta)$ and $p_c = p_{c_{init}}(\theta)$.

There are a variety of scenarios to be considered:

- No rod loss patch / rod loss patch: either the retina is healthy initially or there is a single patch of rod loss;

  - Wide/narrow patch: representing late and early disease stages respectively;
  - Patch eccentricity: we will consider patches on the left-hand side (parafovea/perifovea), centre (mid-periphery) and right-hand side (far-periphery) of the domain;

- No mutation-induced rod loss / mutation-induced rod loss: in the absence of mutation-induced rod loss, photoreceptor (cone) degeneration is due solely to RdCVF starvation, whereas, in the presence of mutation-induced rod loss, rods also degenerate exponentially across the domain over time;

- No treatment / treatment: we consider both the natural evolution of retinal degeneration in the absence of treatment, and the effect of RdCVF treatment upon cone degeneration and cone OS regeneration;

  - Global/local treatment: as in Section 3, RdCVF treatment may be applied globally, across the whole retina, or locally, within a degenerate rod patch only.

Unlike in Section 3, we do not consider the scenario in which both rods and cones have been lost from a patch initially, since, as demonstrated in that section, cone loss seldom proceeds to exceed rod loss.

We consider the following 5 cases (/combinations of scenarios):

Figure 9: Analytical estimates of the eccentricity of the minimum trophic factor concentration using the steady-state model — wide and narrow rod degenerate patch cases. Rod loss only. All plots show the normalised eccentricity of the minimum trophic factor concentration, $\hat{\theta}_{crit} = (\theta_{crit} - \theta_{r_1})/(\theta_{r_2} - \theta_{r_1})$. (a), (c) and (e) global treatment; (b), (d) and (f) local treatment; (a)–(d) wide patch; (e) and (f) narrow patch; (a), (b), (e) and (f) $f_{crit} = 0.3$; (c) and (d) $f_{crit} = 3 \times 10^{-5}$. (a)–(d) column 1: variation in $\hat{\theta}_{crit}$ over $(\theta_{r_1}, \theta_{r_2})$ parameter space. (a)–(f) column 2: variation in $\hat{\theta}_{crit}$ with rod degenerate patch centre position, $(\theta_{r_1} + \theta_{r_2})/2$; column 3: variation in $\hat{\theta}_{crit}$ with rod degenerate patch left boundary position, θ_{r_1}; each curve represents a constant rod degenerate patch width, $\theta_{r_2} - \theta_{r_1}$. $\hat{\theta}_{crit}$ increases monotonically with decreasing rod degenerate patch width and with increasing rod degenerate patch centre/left boundary position in all cases. $\hat{\theta}_{crit}$ remains close to 0 for patches of width $\theta_{r_2} - \theta_{r_1} \geq 0.2$ ((a)–(d)) and close to 0.5 for narrow patches ((e) and (f)). Analytical solutions are obtained by implicitly solving Eqs. (A.46) and (A.47) in (a) and (c), Eqs. (A.59) and (A.60) in (b) and (d), Eqs. (A.71) and (A.72) in (e), and Eqs. (A.83) and (A.84) in (f). The parameter $\epsilon = 10^{-2}$. Remaining parameter values as in Table 2.

1. Wide and narrow patches of rod loss, at left/central/right positions, without mutation-induced rod loss or treatment;
2. Mutation-induced rod loss without patch loss or treatment;
3. Wide and narrow patches of rod loss, at left/central/right positions, with mutation-induced rod loss and without treatment;
4. Wide and narrow patches of rod loss, at left-hand positions, with global and local treatment, and without mutation-induced rod loss;
5. Mutation-induced rod loss with global treatment and without patch loss.

As in Section 3, we also consider the subcases $f_{crit} = 3 \times 10^{-5}$ and $f_{crit} = 0.3$. Unlike in Section 3, we now solve and plot across the whole domain for the $f_{crit} = 0.3$ subcase.

4.1. No treatment

Fig. 10 shows RdCVF and photoreceptor dynamics following the complete removal of rods from the interval $(\theta_{r_1}, \theta_{r_2})$. Cones degenerate within all wide patches of rod loss in (b) ($f_{crit} = 3 \times 10^{-5}$) and (c) ($f_{crit} = 0.3$), and within all narrow patches of rod loss in (c), but only in the left-hand narrow patch of rod loss in (b). The boundaries of cone loss remain within those of rod loss for all patches in (b) and for central and right patches in (c); however, cone loss exceeds rod loss on the left-hand side of the left wide patch and on both the left- and right-hand sides of the left narrow patch in (c). This agrees nicely with our predictions in Figs. 4 and 5.

Fig. 11 shows RdCVF and photoreceptor dynamics with mutation-induced rod degeneration. Cone degeneration initiates at the fovea ($\theta = 0$) in (a) ($f_{crit} = 3 \times 10^{-5}$) and at $\theta = 0.13$ in (b) ($f_{crit} = 0.3$), spreading peripherally (rightwards) in both cases. The retina is mostly preserved for $\theta \le 0.13$ in (b) due to the lower value of $f_{crit} = f_{crit_1} = 3 \times 10^{-5}$ in that region (see discussion in Section 3.2 for more details); however, given enough time, photoreceptors will also degenerate in this region, as in (a). The rate at which cone degeneration spreads across the domain accelerates over time as it propagates peripherally.

Fig. 12 shows RdCVF and photoreceptor dynamics with mutation-induced rod degeneration, following the complete removal of rods from the interval $(\theta_{r_1}, \theta_{r_2})$. Cone degeneration initiates at the fovea ($\theta = 0$) in (a) ($f_{crit} = 3 \times 10^{-5}$) and at $\theta = 0.13$ in (b) ($f_{crit} = 0.3$), spreading peripherally (rightwards) in both cases as in Fig. 12. Mutation-induced rod loss drives cone degenerate patch expansion and precipitates the formation of cone degenerate patches where they did not previously form ((b) centre and right narrow patches).

4.2. Treatment

Fig. 13 shows the effects of treatment upon RdCVF and cone OS dynamics following the complete removal of rods from the interval $(\theta_{r_1}, \theta_{r_2})$. The treatment rate, ξ, is chosen to lie above the critical value, ξ_{crit}, in each case. Treatment results in the complete recovery of cone OSs in all cases, in agreement with our steady-state analytical and numerical predictions (see Figs. 6 and 7).

Fig. 14 shows the effects of treatment upon cone OS dynamics with mutation-induced rod degeneration. Strong treatment ($\xi = 6 \times 10^4$) results in the complete recovery of cone OSs in both (a) ($f_{crit} = 3 \times 10^{-5}$) and (b) ($f_{crit} = 0.3$), while weak treatment ($\xi = 4$ in (a) and $\xi = 4 \times 10^4$ in (b)) delays cone OS degeneration in (a), and provides permanent cone OS recovery for $\theta \le 0.13$ and temporary cone OS recovery for $\theta > 0.13$ in (b).

Figure 10: Simulations of RdCVF and photoreceptor dynamics following the complete removal of rods from the interval $(\theta_{r_1}, \theta_{r_2})$. (a) left-to-right: trophic factor concentration, $f(\theta, t)$, rod density, $p_r(\theta, t)$, and the natural logarithm of the cone density, $\log(p_c(\theta, t))$ (colour scale lower threshold: $\log(\min(p_{c_{init}}(\theta))))$, for the case $f_{crit} = 3 \times 10^{-5}$ and $(\theta_{r_1}, \theta_{r_2}) = (0.4, 0.6)$ (the same as for the top-middle panel in (b)). (b) proportional cone loss, $p_c(\theta, t)/\tilde{p}_c(\theta)$, for the case $f_{crit} = 3 \times 10^{-5}$. (c) proportional cone loss for the case $f_{crit} = 0.3$. Rods are removed from wide (width $O(1)$) and narrow (width $O(\epsilon)$) intervals, on the left-hand side (parafovea/perifovea), centre (mid-periphery) and right-hand side (far-periphery) of the domain. All simulations span the period of one year in dimensional variables. Cones degenerate within all wide patches of rod loss ((b) and (c)), and within all narrow patches of rod loss in (c), but only in the left-hand narrow patch of rod loss in (b). Eqs. (7)–(9) and (11)–(12) were solved using the method of lines, with 4001 mesh points, $F(\theta) = H(\theta_{r_1} - \theta) + H(\theta - \theta_{r_2})$ and $G(\theta) = 1$, and without treatment, cone regeneration or mutation-induced rod degeneration. The parameter $\xi = 0$. Remaining parameter values as in Table 2.

Figure 11: Simulations of RdCVF and photoreceptor dynamics with mutation-induced rod degeneration. (a) and (b) top, left-to-right: trophic factor concentration, $f(\theta, t)$, rod density, $p_r(\theta, t)$, and the natural logarithm of the cone density, $\log(p_c(\theta, t))$ (colour scale lower threshold: $\log(\min(p_{c_{init}}(\theta)))$); bottom: proportional cone loss, $p_c(\theta, t)/\tilde{p}_c(\theta)$. (a) $f_{crit} = 3 \times 10^{-5}$, simulation spans ~22.1 years in dimensional variables, (b) $f_{crit} = 0.3$, simulation spans ~5.5 years in dimensional variables. Cone degeneration initiates at the fovea ($\theta = 0$) in (a) and at $\theta = 0.13$ in (b), spreading peripherally (rightwards) in both cases. Eqs. (7)–(9) and (11)–(12) were solved using the method of lines, with $F(\theta) = 1$ and $G(\theta) = 1$, and without treatment or cone regeneration. (a) 4001 mesh points, (b) 401 mesh points. The parameter $\xi = 0$. Remaining parameter values as in Table 2.

Figure 12: Simulations of RdCVF and photoreceptor dynamics with mutation-induced rod degeneration, following the complete removal of rods from the interval $(\theta_{r_1}, \theta_{r_2})$. (a) left-to-right: trophic factor concentration, $f(\theta, t)$, rod density, $p_r(\theta, t)$, and the natural logarithm of the cone density, $\log(p_c(\theta, t))$ (colour scale lower threshold: $\log(\min(p_{c_{init}}(\theta)))$), for the case $f_{crit} = 3 \times 10^{-5}$ and $(\theta_{r_1}, \theta_{r_2}) = (0.4, 0.6)$ (the same as for the top-middle panel in (b)). (b) proportional cone loss, $p_c(\theta, t)/\tilde{p}_c(\theta)$, for the case $f_{crit} = 3 \times 10^{-5}$. (c) proportional cone loss for the case $f_{crit} = 0.3$. Rods are removed from wide (width $O(1)$) and narrow (width $O(\epsilon)$) intervals, on the left-hand side (parafovea/perifovea), centre (mid-periphery) and right-hand side (far-periphery) of the domain. (a) and (b) simulation spans ~22.1 years in dimensional variables, (c) simulation spans ~5.5 years in dimensional variables. Cone degeneration initiates at the fovea ($\theta = 0$) in (a) and (b), and at $\theta = 0.13$ in (c), spreading peripherally (rightwards) in both cases. Patches of cone loss form and expand in all cases, initiating within patches of rod loss (compare with Fig. 10). Eqs. (7)–(9) and (11)–(12) were solved using the method of lines, with $F(\theta) = H(\theta_{r_1} - \theta) + H(\theta - \theta_{r_2})$ and $G(\theta) = 1$, and without treatment or cone regeneration. (a) and (b) 4001 mesh points, (c) 401 mesh points. The parameter $\xi = 0$. Remaining parameter values as in Table 2.

Figure 13: Simulations of the effects of treatment upon RdCVF and cone OS dynamics following the complete removal of rods from the interval $(\theta_{r_1}, \theta_{r_2})$. Panels show trophic factor concentration, $f(\theta, t)$, and cone OS length, $p_c^L(\theta, t)$, for global treatment (columns 1 and 2), $T(\theta, t) =$, and local treatment (columns 3 and 4), $T(\theta, t) =$, following the removal of rods from wide (width $O(1)$) patches (rows 1 and 3), (a) $(\theta_{r_1}, \theta_{r_2}) = (0.1, 0.3)$ and (b) $(\theta_{r_1}, \theta_{r_2}) = (0.2, 0.4)$, and narrow (width $O(\epsilon)$) patches (rows 2 and 4), (a) $(\theta_{r_1}, \theta_{r_2}) = (0.1, 0.12)$ and (b) $(\theta_{r_1}, \theta_{r_2}) = (0.2, 0.22)$. Black and white horizontal lines mark the time point, t_{crit}, at which treatment is introduced. (a) $f_{crit} = 3 \times 10^{-5}$, (b) $f_{crit} = 0.3$. All simulations span the period of ~2.2 years in dimensional variables. Treatment results in the complete recovery of cone OSs in all cases. Eqs. (7), (8) and (10)–(12) were solved using the method of lines, with 401 mesh points, $F(\theta) = H(\theta_{r_1} - \theta) + H(\theta - \theta_{r_2})$ and $G(\theta) = 1$, and without mutation-induced rod degeneration. Parameter values: $\xi = 6 \times 10^4$ and $t_{crit} = 4.53$ (=1 year in dimensional variables). Remaining parameter values as in Table 2.

Figure 14: Simulations of the effects of treatment upon cone OS dynamics with mutation-induced rod degeneration. Panels show cone OS length, $p_c^L(\theta, t)$, with no treatment (column 1), weak treatment (column 2) and strong treatment (column 3). Black horizontal lines mark the time point, t_{crit}, at which treatment is introduced. (a) $f_{crit} = 3 \times 10^{-5}$ and $t_{crit} = 81.5$, while the simulation spans a period of 24 years in dimensional variables; (b) $f_{crit} = 0.3$ and $t_{crit} = 9.05$, while the simulation spans a period of 12 years in dimensional variables. Strong treatment results in the complete recovery of cone OSs in both (a) and (b), while weak treatment delays cone OS degeneration in (a), and provides permanent cone OS recovery for $\theta \leq 0.13$ and temporary cone OS recovery for $\theta > 0.13$ in (b). Eqs. (7), (8) and (10)–(12) were solved using the method of lines, with 401 mesh points, $F(\theta) = 1$, $G(\theta) = 1$ and $T(\theta, t) =$. Parameter values: (a) and (b) column 1, $\xi = 0$; (a) column 2, $\xi = 4$; (b) column 2, $\xi = 4 \times 10^4$; (a) and (b) column 3, $\xi = 6 \times 10^4$. Remaining parameter values as in Table 2.

5. Discussion

Despite decades of research, the mechanisms underlying secondary cone loss and the spread of photoreceptor degeneration in retinitis pigmentosa (RP) have yet to be conclusively determined; though a number of candidate mechanisms have been hypothesised. One such hypothesis — the trophic factor hypothesis — suggests that rod photoreceptors produce a chemical which cone photoreceptors require to survive, such that when rods are lost, the chemical is depleted, and cone degeneration follows. Such a trophic factor, rod-derived cone viability factor (RdCVF), has been chemically identified by Léveillard et al. (2004), and was found to slow the degeneration and preserve the function of cones in rat, mouse and chick models (Fintz et al., 2003; Léveillard et al., 2004; Mohand-Saïd et al., 1998, 2000, 1997; Yang et al., 2009).

In this paper, we have formulated and analysed reaction-diffusion partial differential equation (PDE) models of RP, in which we considered RdCVF starvation as a potential mechanism of cone loss. As such, this paper complements experimental and clinical studies by isolating a biochemical mechanism in a way that would be almost impossible to achieve experimentally, allowing us to predict the *in vivo* dynamics of photoreceptor loss that would result if this mechanism (RdCVF starvation) alone were responsible for retinal degeneration (subsequent to the direct effects of the underlying mutation). We used our models to answer two key questions: (i) what are the dynamics of cone loss following rod loss?, and (ii) what is the critical RdCVF treatment rate required to prevent cone loss following rod loss?

We began by considering the steady-state version of the model (Section 3), first without (Section 3.2.1) and then with (Section 3.2.2) RdCVF treatment, using a combination of numerical solutions and asymptotic analysis, achieving excellent agreement between the two.

In the untreated scenario we considered four cases: (i) wide patch of rod loss, (ii) narrow patch of rod loss, (iii) wide patch of rod and cone loss, and (iv) narrow patch of rod and cone loss; and for each of these cases the two sub-cases: (a) low trophic factor threshold concentration, $f_{crit} = 3 \times 10^{-5}$, and (b) high trophic factor threshold concentration, $f_{crit} = 0.3$ (Figs. 3–5). In all cases it was found that cone degenerate patch width increases monotonically with increasing rod degenerate patch width and with decreasing rod degenerate patch centre eccentricity, while the difference between rod and cone degenerate patch widths decreases monotonically with rod degenerate patch centre eccentricity. Cone degenerate patch width was predicted to be less than rod degenerate patch width, except for wide and narrow rod loss patches with $f_{crit} = 0.3$, where the cone degenerate patch width may exceed rod degenerate patch width close to the fovea. These results make sense in the light of the fact that the ratio of rods to cones increases monotonically with increasing eccentricity, such that trohpic factor is more scarce toward the centre of the retina, leading to the formation of larger cone degenerate patches.

In the treated scenario we considered two cases: (i) wide patch of rod loss, and (ii) narrow patch of rod loss; and for each of these cases the two sub-cases: (a) low trophic factor threshold concentration, $f_{crit} = 3 \times 10^{-5}$, and (b) high trophic factor threshold concentration, $f_{crit} = 0.3$ (Figs. 6–9). In all cases the critical treatment rate, ξ_{crit}, increases monotonically with decreasing rod degenerate patch centre and left boundary eccentricity. As above, this is to be expected from consideration of the ratio of rods to cones, as endogenously produced trophic factor is more scarce towards the retinal centre. In the wide patch case, the critical treatment rate depends almost entirely upon the rod degenerate patch left boundary position, while being independent of rod degenerate patch width (Fig. 6). By contrast, in the narrow patch case, the critical treatment rate depends upon both the rod degenerate patch left and right boundary positions, and hence upon the rod degenerate patch width, the critical treatment rate increasing monotonically with increasing rod degenerate patch width (Fig. 7). The reason for this difference becomes clear from considering the normalised eccentricity of the minimum trophic factor concentration, $\hat{\theta}_{crit}$, in Fig. 9. For wide patches, $\hat{\theta}_{crit}$ is close to the left-hand rod degenerate patch boundary ($\hat{\theta}_{crit} \in (0, 0.2)$ is in the left-centre-inner region, where only the left-hand boundary exerts an influence); whereas, for narrow patches, $\hat{\theta}_{crit}$ is close to the

centre of the rod degenerate patch ($\hat{\theta}_{crit} \sim 0.5$ is in centre-inner region, upon which both boundaries exert an influence). $\hat{\theta}_{crit}$ is close to the left- rather than the right-hand rod degenerate patch boundary in the wide patch case because the rod to cone ratio is lower there. $\hat{\theta}_{crit}$ increases monotonically with decreasing rod degenerate patch width and with increasing rod degenerate patch centre and left boundary position in all cases. Knowing the location of $\hat{\theta}_{crit}$ is important as it tells us which region of the retina is most at risk of degeneration, suggesting where to spatially target treatment. Comparison of critical treatment rates in the global and local treatment cases shows that ξ_{crit} is almost identical for global and local treatment in the wide patch case, while the critical treatment rate is higher for local than for global treatment for the narrowest patches close to the fovea in the narrow patch case (Fig. 8). Therefore, a targeted treatment, which uses less trophic factor, could be just as effective as a global treatment in most cases.

Next we considered numerical solutions to the dynamic, time-dependent, version of the model (Section 4), first without (Section 4.1) and then with (Section 4.2) RdCVF treatment.

In the untreated scenario we considered three cases: (i) wide and narrow patches of rod loss without mutation-induced rod degeneration, (ii) mutation-induced rod degeneration without patch loss, and (iii) wide and narrow patches of rod loss with mutation-induced rod degeneration; and for each of these cases the two sub-cases: (a) low trophic factor threshold concentration, $f_{crit} = 3 \times 10^{-5}$, and (b) high trophic factor threshold concentration, $f_{crit} = 0.3$ (Figs. 10–12). In Case (i) it was found that cone degeneration always occurs within wide rod degenerate patches, whereas cone degeneration may not occur in narrow rod degenerate patches away from the retinal centre for $f_{crit} = 3 \times 10^{-5}$ (Fig. 10). Further, for $f_{crit} = 0.3$, the spatial extent of cone loss may exceed that of rod loss in near-central rod degenerate patches. This is in good agreement with our predictions from the steady-state problem in Section 3. In Case (ii) cone degeneration initiates at the retinal centre and spreads peripherally, accelerating as it does so (Fig. 11). This does not replicate any of the spatio-temporal patterns of visual field loss (and hence retinal degeneration) observed by Grover et al. (1998) (see also Roberts et al., 2018), suggesting that other mechanisms are necessary, either in place of or together with trophic factor starvation-induced degeneration, to explain *in vivo* disease progression. In Case (iii) mutation-induced rod degeneration was shown to precipitation the formation of cone degenerate patches at the locations of narrow rod degenerate patches, where they would not have otherwise formed, and to drive the expansion of cone degenerate patches (Fig. 12). This occurs because the loss of rods reduces the rate of production of trophic factor, pushing RdCVF concentrations below the critical threshold in regions which were previously well-supplied.

As a hypothesis to explain the spread of photoreceptor degeneration, the trophic factor mechanism is self-limiting. This is because, following the loss of rods and consequent drop in RdCVF concentrations, the loss of cones decreases trophic factor demand until RdCVF levels are healthy once more. This is in contrast to the oxygen toxicity mechanism, considered by the author in previous publications (see Roberts et al., 2017, 2018), which is self-reinforcing; toxically high oxygen levels causing photoreceptor degeneration, leading to a further rise in oxygen levels, and so on.

In the treated scenario we considered two cases: (i) wide and narrow patches of rod loss, without mutation-induced rod degeneration, with global and local treatment, and (ii) mutation-induced rod degeneration, without patch loss, with weak and strong global treatment; and for each of these cases the two sub-cases: (a) low trophic factor threshold concentration, $f_{crit} = 3 \times 10^{-5}$, and (b) high trophic factor threshold concentration, $f_{crit} = 0.3$ (Figs. 13–14). In Case (i), the treatment rate, ξ, was chosen to lie above the critical treatment rate, ξ_{crit}, predicted from the steady-state model, resulting in the complete recovery of cone outer segments (OSs) in all cases as expected (Fig. 13). In Case (ii), strong treatment ($\xi = 6 \times 10^4$) resulted in the complete recovery of cone OSs, while weak treatment ($\xi = 4$ or $\xi = 4 \times 10^4$) delayed or temporarily reversed cone OS degeneration (Fig. 14). While the possible rates of RdCVF treatment have yet to be measured, our models provide useful predictions of the critical treatment rate required to prevent or

reverse damage to cones and the possible effects of different treatment rates.

In conclusion, we have constructed novel mathematical models of RP to describe and predict retinal degeneration, and the effects of treatment, under the trophic factor hypothesis. We predicted the spatial extent of cone degeneration, following the loss of a patch of rods, and the critical RdCVF treatment rate required to prevent cone loss in this case. Lastly, we predicted the spatio-temporal patterns of cone degeneration that would result from this mechanism, demonstrating that additional mechanisms are required to explain the patterns seen *in vivo*.

Appendix A. Asymptotic analyses (continued)

Here we consider Cases 2-8 from Section 3.1. In each case the same approach is taken as in Section 3.1.1. Where the analysis differs from that in Section 3.1.1 we explain the difference, otherwise the results are simply stated.

In the sections that follow (Appendix A.1–Appendix A.7) we will use the following parameters (in addition to those defined in Section 3.1), defined here for ease of reference:

- $K := e^{\epsilon^{-1}\sqrt{c_1}(\theta_{r_2}-\theta_{r_1})}$;

- $B := 1 + \epsilon^{\frac{1}{2}}\sqrt{\dfrac{\eta}{D_f c_1}}$;

- $\bar{B} := 1 + \epsilon^{\frac{1}{2}}\sqrt{\dfrac{\eta}{D_f \bar{c}_1}}$;

- $Q := \epsilon^{\frac{1}{2}}\sqrt{\dfrac{\eta}{D_f c_1}} - 1 = B - 2$;

- $\bar{Q} := \epsilon^{\frac{1}{2}}\sqrt{\dfrac{\eta}{D_f \bar{c}_1}} - 1 = \bar{B} - 2$;

- $M := e^{\epsilon^{-1/2}\sqrt{\frac{\eta}{D_f}}(\theta_{r_2}-\theta_{r_1})}$.

Appendix A.1. Narrow patch of rod loss without treatment

In this case there is no centre-outer region and the left-centre-inner and right-centre-inner regions coalesce into a single centre-inner region (see Fig. 2(b)).

Left- and Right-Outer:

$$f_0(\theta) = \frac{\alpha B_3 \theta e^{-b_3 \theta}}{\beta B_2 e^{-b_2 \theta}}, \tag{A.1}$$

Left-Inner:

$$f_0(\theta) = A_1 e^{\epsilon^{-1}\sqrt{c_1}(\theta-\theta_{r_1})} + \frac{c_2}{c_1}, \tag{A.2}$$

Centre-Inner:

$$f_0(\theta) = A_3 e^{-\epsilon^{-1}\sqrt{c_1}(\theta-\theta_{r_1})} + A_4 e^{\epsilon^{-1}\sqrt{c_1}(\theta-\theta_{r_2})}, \tag{A.3}$$

Right-Inner:

$$f_0(\theta) = A_8 e^{-\epsilon^{-1}\sqrt{\bar{c}_1}(\theta-\theta_{r_2})} + \frac{\bar{c}_2}{\bar{c}_1}, \tag{A.4}$$

where:

$$A_1 = \left(1 + \sqrt{\frac{c_1}{\bar{c}_1}}\right)^{-1}\left[\frac{c_2}{2c_1}K^{-2}\left(\sqrt{\frac{c_1}{\bar{c}_1}} - 1\right) + \frac{\bar{c}_2}{\bar{c}_1}K^{-1}\right] - \frac{c_2}{2c_1},$$

$$A_3 = \frac{c_2}{2c_1},$$

$$A_4 = \left(1 + \sqrt{\frac{c_1}{\bar{c}_1}}\right)^{-1}\left[\frac{c_2}{2c_1}K^{-1}\left(\sqrt{\frac{c_1}{\bar{c}_1}} - 1\right) + \frac{\bar{c}_2}{\bar{c}_1}\right],$$

$$A_8 = -\left(1 + \sqrt{\frac{c_1}{\bar{c}_1}}\right)^{-1}\left[\frac{c_2}{2c_1}K^{-1}\left(\sqrt{\frac{c_1}{\bar{c}_1}} - 1\right) + \frac{\bar{c}_2}{\bar{c}_1}\right]\sqrt{\frac{c_1}{\bar{c}_1}} + \frac{c_2}{2c_1}\sqrt{\frac{c_1}{\bar{c}_1}}K^{-1},$$

Left-Composite:

$$f_{0_{left-comp}}(\theta) = \frac{\alpha B_3 \theta e^{-b_3\theta}}{\beta B_2 e^{-b_2\theta}} + A_1 e^{\epsilon^{-1}\sqrt{c_1}(\theta-\theta_{r_1})}, \tag{A.5}$$

Centre-Composite is identical with the centre-inner,
Right-Composite:

$$f_{0_{right-comp}}(\theta) = \frac{\alpha B_3 \theta e^{-b_3\theta}}{\beta B_2 e^{-b_2\theta}} + A_8 e^{-\epsilon^{-1}\sqrt{\bar{c}_1}(\theta-\theta_{r_2})}, \tag{A.6}$$

Minimal Cone Degenerate Patch Left Boundary:

$$\theta_{crit_1} = \theta_{r_1} + \frac{\epsilon}{\sqrt{c_1}}\log\left(A_1^{-1}\left(f_{crit} - \frac{c_2}{c_1}\right)\right), \text{ for } f_{crit} \geq A_1 + \frac{c_2}{c_1}, \tag{A.7}$$

$$f_{crit} = A_3 e^{-\epsilon^{-1}\sqrt{c_1}(\theta_{crit_1}-\theta_{r_1})} + A_4 e^{\epsilon^{-1}\sqrt{c_1}(\theta_{crit_1}-\theta_{r_2})}, \text{ for } f_{crit} \leq A_1 + \frac{c_2}{c_1}, \tag{A.8}$$

Maximal Cone Degenerate Patch Right Boundary:

$$f_{crit} = A_3 e^{-\epsilon^{-1}\sqrt{c_1}(\theta_{crit_2}-\theta_{r_1})} + A_4 e^{\epsilon^{-1}\sqrt{c_1}(\theta_{crit_2}-\theta_{r_2})}, \text{ for } f_{crit} \leq A_8 + \frac{\bar{c}_2}{\bar{c}_1}, \tag{A.9}$$

$$\theta_{crit_2} = \theta_{r_2} - \frac{\epsilon}{\sqrt{\bar{c}_1}}\log\left(A_8^{-1}\left(f_{crit} - \frac{\bar{c}_2}{\bar{c}_1}\right)\right), \text{ for } f_{crit} \geq A_8 + \frac{\bar{c}_2}{\bar{c}_1}, \tag{A.10}$$

where Eqs. (A.8) and (A.9) must be solved implicitly for θ_{crit_1} and θ_{crit_2} respectively,

- when $f_{crit} \geq A_1 + \frac{c_2}{c_1}$, $\theta_{crit_1} \leq \theta_{r_1}$;

- when $f_{crit} < A_1 + \frac{c_2}{c_1}$, $\theta_{crit_1} > \theta_{r_1}$;

- when $f_{crit} \geq A_8 + \frac{\bar{c}_2}{\bar{c}_1}$, $\theta_{crit_2} \geq \theta_{r_2}$;

- when $f_{crit} < A_8 + \frac{\bar{c}_2}{\bar{c}_1}$, $\theta_{crit_2} < \theta_{r_2}$.

Appendix A.2. Wide patch of rod and cone loss without treatment

In this case, $G(\theta) = H(\theta_{c_1} - \theta) + H(\theta - \theta_{c_2})$ (and $F(\theta) = H(\theta_{r_1} - \theta) + H(\theta - \theta_{r_2})$ as before), where $\theta_{c_1} = \theta_{r_1}$ and $\theta_{c_2} = \theta_{r_2}$. The asymptotics for the rod and cone loss case, both here and in Appendix A.3, is valid in the region $\theta \in (\sim 0.16, 1]$.

We decompose into the same regions as in Section 3.1.1 (see Fig. 2(c)); however, the left- and right-centre-inner regions are $O(\epsilon^{1/2})$ width in this case. This is because, in the absence of cones in the central region, we must seek a dominant balance between the diffusion and decay terms. Thus, the new scaling on θ in the left-centre-inner region is: $\hat{\theta}' = \epsilon^{-1/2}(\theta - \theta_{r_1})$; while the new scaling on θ in right-centre-inner region is: $\bar{\theta}' = \epsilon^{-1/2}(\theta - \theta_{r_2})$; while the regular perturbation expansion for f in the central region is

$$f(\theta) = f_0(\theta) + \epsilon^{1/2} f_1(\theta) + O(\epsilon). \tag{A.11}$$

Perturbation expansions are not required for rods and cones in this region since they are absent here. The scalings on θ and the regular perturbation expansions in the left-inner and right-inner regions are the same as in 3.1.1.

Left- and Right-Outer:

$$f_0(\theta) = \frac{\alpha B_3 \theta e^{-b_3 \theta}}{\beta B_2 e^{-b_2 \theta}}, \tag{A.12}$$

Left-Inner:

$$f_0(\theta) = \frac{c_2}{c_1} \left(1 + \left[\left(1 + \epsilon^{1/2} \sqrt{\frac{\eta}{D_f c_1}} \right)^{-1} - 1 \right] e^{\epsilon^{-1} \sqrt{c_1}(\theta - \theta_{r_1})} \right), \tag{A.13}$$

Left-Centre-Inner:

$$f_0(\theta) = \frac{c_2}{c_1} \left(1 + \epsilon^{1/2} \sqrt{\frac{\eta}{D_f c_1}} \right)^{-1} e^{-\epsilon^{-1/2} \sqrt{\frac{\eta}{D_f}}(\theta - \theta_{r_1})}, \tag{A.14}$$

Centre-Outer:

$$f_0(\theta) = 0, \tag{A.15}$$

Right-Centre-Inner:

$$f_0(\theta) = \frac{\bar{c}_2}{\bar{c}_1} \left(1 + \epsilon^{1/2} \sqrt{\frac{\eta}{D_f \bar{c}_1}} \right)^{-1} e^{\epsilon^{-1/2} \sqrt{\frac{\eta}{D_f}}(\theta - \theta_{r_2})}, \tag{A.16}$$

Right-Inner:

$$f_0(\theta) = \frac{\bar{c}_2}{\bar{c}_1} \left(1 + \left[\left(1 + \epsilon^{1/2} \sqrt{\frac{\eta}{D_f \bar{c}_1}} \right)^{-1} - 1 \right] e^{-\epsilon^{-1} \sqrt{\bar{c}_1}(\theta - \theta_{r_2})} \right), \tag{A.17}$$

Left-Composite:

$$f_{0_{left-comp}}(\theta) = \frac{\alpha B_3 \theta e^{-b_3 \theta}}{\beta B_2 e^{-b_2 \theta}} + \frac{c_2}{c_1} \left[\left(1 + \epsilon^{1/2} \sqrt{\frac{\eta}{D_f c_1}} \right)^{-1} - 1 \right] e^{\epsilon^{-1} \sqrt{c_1}(\theta - \theta_{r_1})}, \tag{A.18}$$

Centre-Composite:

$$f_{0_{centre-comp}}(\theta) = \frac{c_2}{c_1}\left(1 + \epsilon^{1/2}\sqrt{\frac{\eta}{D_f c_1}}\right)^{-1} e^{-\epsilon^{-1/2}\sqrt{\frac{\eta}{D_f}}(\theta-\theta_{r_1})} + \frac{\bar{c}_2}{\bar{c}_1}\left(1 + \epsilon^{1/2}\sqrt{\frac{\eta}{D_f \bar{c}_1}}\right)^{-1} e^{\epsilon^{-1/2}\sqrt{\frac{\eta}{D_f}}(\theta-\theta_{r_2})}, \tag{A.19}$$

Right-Composite:

$$f_{0_{right-comp}}(\theta) = \frac{\alpha B_3 \theta e^{-b_3\theta}}{\beta B_2 e^{-b_2\theta}} + \frac{\bar{c}_2}{\bar{c}_1}\left[\left(1 + \epsilon^{1/2}\sqrt{\frac{\eta}{D_f \bar{c}_1}}\right)^{-1} - 1\right] e^{-\epsilon^{-1}\sqrt{\bar{c}_1}(\theta-\theta_{r_2})}, \tag{A.20}$$

Maximal Cone Degenerate Patch Left Boundary:

$$\theta_{crit_1} = \theta_{r_1} + \frac{\epsilon}{\sqrt{c_1}}\log\left(\left[\left(1 + \epsilon^{1/2}\sqrt{\frac{\eta}{D_f c_1}}\right)^{-1} - 1\right]^{-1}\left(\frac{c_1 f_{crit}}{c_2} - 1\right)\right), \text{ for } f_{crit} \geq \frac{c_2}{c_1}\left(1 + \epsilon^{1/2}\sqrt{\frac{\eta}{D_f c_1}}\right)^{-1}, \tag{A.21}$$

$$\theta_{crit_1} = \theta_{r_1} - \epsilon^{1/2}\sqrt{\frac{D_f}{\eta}}\log\left(\left(1 + \epsilon^{1/2}\sqrt{\frac{\eta}{D_f c_1}}\right)\frac{c_1 f_{crit}}{c_2}\right), \text{ for } f_{crit} \leq \frac{c_2}{c_1}\left(1 + \epsilon^{1/2}\sqrt{\frac{\eta}{D_f c_1}}\right)^{-1}, \tag{A.22}$$

Minimal Cone Degenerate Patch Right Boundary:

$$\theta_{crit_2} = \theta_{r_2} + \epsilon^{1/2}\sqrt{\frac{D_f}{\eta}}\log\left(\left(1 + \epsilon^{1/2}\sqrt{\frac{\eta}{D_f \bar{c}_1}}\right)\frac{\bar{c}_1 f_{crit}}{\bar{c}_2}\right), \text{ for } f_{crit} \leq \frac{\bar{c}_2}{\bar{c}_1}\left(1 + \epsilon^{1/2}\sqrt{\frac{\eta}{D_f \bar{c}_1}}\right)^{-1}, \tag{A.23}$$

$$\theta_{crit_2} = \theta_{r_2} - \frac{\epsilon}{\sqrt{\bar{c}_1}}\log\left(\left[\left(1 + \epsilon^{1/2}\sqrt{\frac{\eta}{D_f \bar{c}_1}}\right)^{-1} - 1\right]^{-1}\left(\frac{\bar{c}_1 f_{crit}}{\bar{c}_2} - 1\right)\right), \text{ for } f_{crit} \geq \frac{\bar{c}_2}{\bar{c}_1}\left(1 + \epsilon^{1/2}\sqrt{\frac{\eta}{D_f \bar{c}_1}}\right)^{-1}, \tag{A.24}$$

where the maximal and minimal labels above are the other way around to those in the rod loss only cases (which estimate maximal cone degenerate patch width), since here, for the rod and cone loss case, we are estimating the minimal cone degenerate patch width,

- when $f_{crit} \geq \frac{c_2}{c_1}\left(1 + \epsilon^{1/2}\sqrt{\frac{\eta}{D_f c_1}}\right)^{-1}$, $\theta_{crit_1} \leq \theta_{r_1}$;

- when $f_{crit} < \frac{c_2}{c_1}\left(1 + \epsilon^{1/2}\sqrt{\frac{\eta}{D_f c_1}}\right)^{-1}$, $\theta_{crit_1} > \theta_{r_1}$;

- when $f_{crit} \geq \frac{\bar{c}_2}{\bar{c}_1}\left(1 + \epsilon^{1/2}\sqrt{\frac{\eta}{D_f \bar{c}_1}}\right)^{-1}$, $\theta_{crit_2} \geq \theta_{r_2}$;

- when $f_{crit} < \frac{\bar{c}_2}{\bar{c}_1}\left(1 + \epsilon^{1/2}\sqrt{\frac{\eta}{D_f \bar{c}_1}}\right)^{-1}$, $\theta_{crit_2} < \theta_{r_2}$.

Appendix A.3. Narrow patch of rod and cone loss without treatment

As in Appendix A.1 there is no centre-outer region and the left-centre-inner and right-centre-inner regions coalesce into a single centre-inner region (see Fig. 2(d)).

Left- and Right-Outer:

$$f_0(\theta) = \frac{\alpha B_3 \theta e^{-b_3\theta}}{\beta B_2 e^{-b_2\theta}},$$ (A.25)

Left-Inner:

$$f_0(\theta) = A_1 e^{\epsilon^{-1}\sqrt{c_1}(\theta-\theta_{r_1})} + \frac{c_2}{c_1},$$ (A.26)

Centre-Inner:

$$f_0(\theta) = A_3 e^{-\epsilon^{-1/2}\sqrt{\frac{\eta}{D_f}}(\theta-\theta_{r_1})} + A_4 e^{\epsilon^{-1/2}\sqrt{\frac{\eta}{D_f}}(\theta-\theta_{r_2})},$$ (A.27)

Right-Inner:

$$f_0(\theta) = A_8 e^{-\epsilon^{-1}\sqrt{\bar{c}_1}(\theta-\theta_{r_2})} + \frac{\bar{c}_2}{\bar{c}_1},$$ (A.28)

where:

$$A_1 = (1 + B^{-1}Q)\frac{B^{-1}\bar{Q}M^{-1}\frac{c_2}{c_1} + \frac{\bar{c}_2}{\bar{c}_1}}{\bar{B}M - B^{-1}Q\bar{Q}M^{-1}} + (B^{-1} - 1)\frac{c_2}{c_1},$$

$$A_3 = B^{-1}\left[\frac{c_2}{c_1} + Q\frac{B^{-1}\bar{Q}M^{-1}\frac{c_2}{c_1} + \frac{\bar{c}_2}{\bar{c}_1}}{\bar{B}M - B^{-1}Q\bar{Q}M^{-1}}\right],$$

$$A_4 = \frac{B^{-1}\bar{Q}\frac{c_2}{c_1} + M\frac{\bar{c}_2}{\bar{c}_1}}{\bar{B}M - B^{-1}Q\bar{Q}M^{-1}},$$

$$A_8 = (M + B^{-1}QM^{-1})\frac{B^{-1}\bar{Q}M^{-1}\frac{c_2}{c_1} + \frac{\bar{c}_2}{\bar{c}_1}}{\bar{B}M - B^{-1}Q\bar{Q}M^{-1}} + B^{-1}M^{-1}\frac{c_2}{c_1} - \frac{\bar{c}_2}{\bar{c}_1},$$

Left-Composite:

$$f_{0_{left-comp}}(\theta) = \frac{\alpha B_3 \theta e^{-b_3\theta}}{\beta B_2 e^{-b_2\theta}} + A_1 e^{\epsilon^{-1}\sqrt{c_1}(\theta-\theta_{r_1})},$$ (A.29)

Centre-Composite is identical with the centre-inner.
Right-Composite:

$$f_{0_{right-comp}}(\theta) = \frac{\alpha B_3 \theta e^{-b_3\theta}}{\beta B_2 e^{-b_2\theta}} + A_8 e^{-\epsilon^{-1}\sqrt{\bar{c}_1}(\theta-\theta_{r_2})},$$ (A.30)

Maximal Cone Degenerate Patch Left Boundary:

$$\theta_{crit_1} = \theta_{r_1} + \frac{\epsilon}{\sqrt{c_1}}\log\left(A_1^{-1}\left(f_{crit} - \frac{c_2}{c_1}\right)\right), \text{ for } f_{crit} \geq A_1 + \frac{c_2}{c_1},$$ (A.31)

$$f_{crit} = A_3 e^{-\epsilon^{-1}\sqrt{c_1}(\theta_{crit_1}-\theta_{r_1})} + A_4 e^{\epsilon^{-1}\sqrt{c_1}(\theta_{crit_1}-\theta_{r_2})}, \text{ for } f_{crit} \leq A_1 + \frac{c_2}{c_1},$$ (A.32)

Minimal Cone Degenerate Patch Right Boundary:

$$f_{crit} = A_3 e^{-\epsilon^{-1} \sqrt{\bar{c}_1}(\theta_{crit_2} - \theta_{r_1})} + A_4 e^{\epsilon^{-1} \sqrt{\bar{c}_1}(\theta_{crit_2} - \theta_{r_2})}, \text{ for } f_{crit} \leq A_8 + \frac{\bar{c}_2}{\bar{c}_1}, \tag{A.33}$$

$$\theta_{crit_2} = \theta_{r_2} - \frac{\epsilon}{\sqrt{\bar{c}_1}} \log\left(A_8^{-1}\left(f_{crit} - \frac{\bar{c}_2}{\bar{c}_1}\right)\right), \text{ for } f_{crit} \geq A_8 + \frac{\bar{c}_2}{\bar{c}_1}, \tag{A.34}$$

where Eqs. (A.32) and (A.33) must be solved implicitly for θ_{crit_1} and θ_{crit_2} respectively. The maximal and minimal labels above are the other way around to those in the rod loss only cases (which estimate maximal cone degenerate patch width), since here, for the rod and cone loss case, we are estimating the minimal cone degenerate patch width,

- when $f_{crit} \geq A_1 + \frac{c_2}{c_1}$, $\theta_{crit_1} \leq \theta_{r_1}$;

- when $f_{crit} < A_1 + \frac{c_2}{c_1}$, $\theta_{crit_1} > \theta_{r_1}$;

- when $f_{crit} \geq A_8 + \frac{\bar{c}_2}{\bar{c}_1}$, $\theta_{crit_2} \geq \theta_{r_2}$;

- when $f_{crit} < A_8 + \frac{\bar{c}_2}{\bar{c}_1}$, $\theta_{crit_2} < \theta_{r_2}$.

Appendix A.4. Wide patch of rod loss with global treatment

This case is the same as that in Section 3.1.1, except that treatment is applied across the whole domain ($T(\theta) = 1$) at rate ξ. Thus, we decompose the domain into the same regions as in the untreated case (see Fig. 2(a)). We rescale $\xi = \epsilon^{-2}\xi'$, where $\xi' = O(1)$, here and in Appendix A.5–Appendix A.7, dropping the dash.

Left- and Right-Outer:

$$f_0(\theta) = \frac{\alpha B_3 \theta e^{-b_3 \theta} + \xi}{\beta B_2 e^{-b_2 \theta}}, \tag{A.35}$$

Left-Inner:

$$f_0(\theta) = \frac{c_2}{c_1}\left(1 - \frac{1}{2}e^{\epsilon^{-1}\sqrt{c_1}(\theta - \theta_{r_1})}\right) + \frac{\xi}{D_f c_1}, \tag{A.36}$$

Left-Centre-Inner:

$$f_0(\theta) = \frac{c_2}{2c_1}e^{-\epsilon^{-1}\sqrt{c_1}(\theta - \theta_{r_1})} + \frac{\xi}{D_f c_1}, \tag{A.37}$$

Centre-Outer:

$$f_0(\theta) = \frac{\xi}{\beta B_2 e^{-b_2 \theta}}, \tag{A.38}$$

Right-Centre-Inner:

$$f_0(\theta) = \frac{\bar{c}_2}{2\bar{c}_1}e^{\epsilon^{-1}\sqrt{\bar{c}_1}(\theta - \theta_{r_2})} + \frac{\xi}{D_f \bar{c}_1}, \tag{A.39}$$

Right-Inner:

$$f_0(\theta) = \frac{\bar{c}_2}{\bar{c}_1}\left(1 - \frac{1}{2}e^{-\epsilon^{-1}\sqrt{\bar{c}_1}(\theta - \theta_{r_2})}\right) + \frac{\xi}{D_f \bar{c}_1}, \tag{A.40}$$

Left-Composite:

$$f_{0_{left-comp}}(\theta) = \frac{\alpha B_3 \theta e^{-b_3 \theta} + \xi}{\beta B_2 e^{-b_2 \theta}} - \frac{c_2}{2c_1} e^{\epsilon^{-1} \sqrt{c_1}(\theta - \theta_{r_1})}, \tag{A.41}$$

Centre-Composite:

$$f_{0_{centre-comp}}(\theta) = \frac{c_2}{2c_1} e^{-\epsilon^{-1} \sqrt{c_1}(\theta - \theta_{r_1})} + \frac{\xi}{\beta B_2 e^{-b_2 \theta}} + \frac{\bar{c}_2}{2\bar{c}_1} e^{\epsilon^{-1} \sqrt{\bar{c}_1}(\theta - \theta_{r_2})}, \tag{A.42}$$

Right-Composite:

$$f_{0_{right-comp}}(\theta) = \frac{\alpha B_3 \theta e^{-b_3 \theta} + \xi}{\beta B_2 e^{-b_2 \theta}} - \frac{\bar{c}_2}{2\bar{c}_1} e^{-\epsilon^{-1} \sqrt{\bar{c}_1}(\theta - \theta_{r_2})}, \tag{A.43}$$

Minimal Cone Degenerate Patch Left Boundary:

$$\theta_{crit_1} = \theta_{r_1} + \text{sgn}\left(f_{crit} - \frac{c_2}{2c_1} - \frac{\xi}{D_f c_1}\right) \frac{\epsilon}{\sqrt{c_1}} \log\left(1 - \left|\frac{2c_1 f_{crit}}{c_2} - \frac{2\xi}{D_f c_2} - 1\right|\right), \tag{A.44}$$

Maximal Cone Degenerate Patch Right Boundary:

$$\theta_{crit_2} = \theta_{r_2} - \text{sgn}\left(f_{crit} - \frac{\bar{c}_2}{2\bar{c}_1} - \frac{\xi}{D_f \bar{c}_1}\right) \frac{\epsilon}{\sqrt{\bar{c}_1}} \log\left(1 - \left|\frac{2\bar{c}_1 f_{crit}}{\bar{c}_2} - \frac{2\xi}{D_f \bar{c}_2} - 1\right|\right), \tag{A.45}$$

- when $f_{crit} \geq c_2/(2c_1) + \xi/(D_f c_1)$, $\theta_{crit_1} \leq \theta_{r_1}$;

- when $f_{crit} < c_2/(2c_1) + \xi/(D_f c_1)$, $\theta_{crit_1} > \theta_{r_1}$;

- when $f_{crit} \geq \bar{c}_2/(2\bar{c}_1) + \xi/(D_f \bar{c}_1)$, $\theta_{crit_2} \geq \theta_{r_2}$;

- when $f_{crit} < \bar{c}_2/(2\bar{c}_1) + \xi/(D_f \bar{c}_1)$, $\theta_{crit_2} < \theta_{r_2}$.

We calculate the critical treatment rate, ξ_{crit}, as the minimal treatment rate required to prevent cone loss, that is, to keep $f \geq f_{crit}$ local to the patch of rod loss. The minimum trophic factor concentration is achieved at some θ_{crit} within the interval $\theta_{crit} \in [\theta_{r_1}, \theta_{r_2}]$. Thus, ξ_{crit} and θ_{crit} can be found by setting $\xi = \xi_{crit}$ and $\theta = \theta_{crit}$ in both Eqn. (A.42) and in the equation obtained by setting $f'_{0_{centre-comp}}(\theta) = 0$ from Eqn. (A.42), to provide,

Critical treatment rate:

$$\xi_{crit} = \beta B_2 e^{-b_2 \theta_{crit}} \left(f_{crit} - \frac{c_2}{2c_1} e^{-\epsilon^{-1} \sqrt{c_1}(\theta_{crit} - \theta_{r_1})} - \frac{\bar{c}_2}{2\bar{c}_1} e^{\epsilon^{-1} \sqrt{\bar{c}_1}(\theta_{crit} - \theta_{r_2})}\right), \tag{A.46}$$

Eccentricity of the minimum trophic factor concentration:

$$0 = -\frac{\epsilon^{-1} c_2}{2\sqrt{c_1}} e^{-\epsilon^{-1} \sqrt{c_1}(\theta_{crit} - \theta_{r_1})} + \frac{\xi_{crit} b_2}{\beta B_2 e^{-b_2 \theta_{crit}}} + \frac{\epsilon^{-1} \bar{c}_2}{2\sqrt{\bar{c}_1}} e^{\epsilon^{-1} \sqrt{\bar{c}_1}(\theta_{crit} - \theta_{r_2})}, \tag{A.47}$$

solving Eqs. (A.46) and (A.47) simultaneously.

Appendix A.5. Wide patch of rod loss with local treatment

This case is the same as that in Appendix A.4, except that treatment is only applied locally, within the degenerate rod patch ($T(\theta) = 1 - F(\theta)$), rather than globally, across the whole domain. We decompose the domain into the same regions as in the untreated and global treatment cases (see Fig. 2(a)).

Left- and Right-Outer:

$$f_0(\theta) = \frac{\alpha B_3 \theta e^{-b_3\theta}}{\beta B_2 e^{-b_2\theta}}, \tag{A.48}$$

Left-Inner:

$$f_0(\theta) = \frac{c_2}{c_1} + \left(\frac{\xi}{2D_f c_1} - \frac{c_2}{2c_1}\right) e^{\epsilon^{-1}\sqrt{c_1}(\theta - \theta_{r_1})}, \tag{A.49}$$

Left-Centre-Inner:

$$f_0(\theta) = \left(\frac{c_2}{2c_1} - \frac{\xi}{2D_f c_1}\right) e^{-\epsilon^{-1}\sqrt{c_1}(\theta - \theta_{r_1})} + \frac{\xi}{D_f c_1}, \tag{A.50}$$

Centre-Outer:

$$f_0(\theta) = \frac{\xi}{\beta B_2 e^{-b_2\theta}}, \tag{A.51}$$

Right-Centre-Inner:

$$f_0(\theta) = \left(\frac{\bar{c}_2}{2\bar{c}_1} - \frac{\xi}{2D_f \bar{c}_1}\right) e^{\epsilon^{-1}\sqrt{\bar{c}_1}(\theta - \theta_{r_2})} + \frac{\xi}{D_f \bar{c}_1}, \tag{A.52}$$

Right-Inner:

$$f_0(\theta) = \frac{\bar{c}_2}{\bar{c}_1} + \left(\frac{\xi}{2D_f \bar{c}_1} - \frac{\bar{c}_2}{2\bar{c}_1}\right) e^{-\epsilon^{-1}\sqrt{\bar{c}_1}(\theta - \theta_{r_2})}, \tag{A.53}$$

Left-Composite:

$$f_{0_{left-comp}}(\theta) = \frac{\alpha B_3 \theta e^{-b_3\theta}}{\beta B_2 e^{-b_2\theta}} + \left(\frac{\xi}{2D_f c_1} - \frac{c_2}{2c_1}\right) e^{\epsilon^{-1}\sqrt{c_1}(\theta - \theta_{r_1})}, \tag{A.54}$$

Centre-Composite:

$$f_{0_{centre-comp}}(\theta) = \left(\frac{c_2}{2c_1} - \frac{\xi}{2D_f c_1}\right) e^{-\epsilon^{-1}\sqrt{c_1}(\theta - \theta_{r_1})} + \frac{\xi}{\beta B_2 e^{-b_2\theta}} + \left(\frac{\bar{c}_2}{2\bar{c}_1} - \frac{\xi}{2D_f \bar{c}_1}\right) e^{\epsilon^{-1}\sqrt{\bar{c}_1}(\theta - \theta_{r_2})}, \tag{A.55}$$

Right-Composite:

$$f_{0_{right-comp}}(\theta) = \frac{\alpha B_3 \theta e^{-b_3\theta}}{\beta B_2 e^{-b_2\theta}} + \left(\frac{\xi}{2D_f \bar{c}_1} - \frac{\bar{c}_2}{2\bar{c}_1}\right) e^{-\epsilon^{-1}\sqrt{\bar{c}_1}(\theta - \theta_{r_2})}, \tag{A.56}$$

Minimal Cone Degenerate Patch Left Boundary:

$$\theta_{crit_1} = \theta_{r_1} + \text{sgn}\left(f_{crit} - \frac{c_2}{2c_1} - \frac{\xi}{2D_f c_1}\right) \frac{\epsilon}{\sqrt{c_1}} \log\left(\left(1 - \frac{\xi}{D_f c_2}\right)^{-1}\left(1 - \frac{\xi}{D_f c_2} - \left|\frac{2c_1 f_{crit}}{c_2} - 1 - \frac{\xi}{D_f c_2}\right|\right)\right), \tag{A.57}$$

Maximal Cone Degenerate Patch Right Boundary:

$$\theta_{crit_2} = \theta_{r_2} - \text{sgn}\left(f_{crit} - \frac{\bar{c}_2}{2\bar{c}_1} - \frac{\xi}{2D_f \bar{c}_1}\right) \frac{\epsilon}{\sqrt{\bar{c}_1}} \log\left(\left(1 - \frac{\xi}{D_f \bar{c}_2}\right)^{-1}\left(1 - \frac{\xi}{D_f \bar{c}_2} - \left|\frac{2\bar{c}_1 f_{crit}}{\bar{c}_2} - 1 - \frac{\xi}{D_f \bar{c}_2}\right|\right)\right), \tag{A.58}$$

- when $f_{crit} \geq c_2/(2c_1) + \xi/(2D_f c_1)$, $\theta_{crit_1} \leq \theta_{r_1}$;

- when $f_{crit} < c_2/(2c_1) + \xi/(2D_f c_1)$, $\theta_{crit_1} > \theta_{r_1}$;

- when $f_{crit} \geq \bar{c}_2/(2\bar{c}_1) + \xi/(2D_f \bar{c}_1)$, $\theta_{crit_2} \geq \theta_{r_2}$;

- when $f_{crit} < \bar{c}_2/(2\bar{c}_1) + \xi/(2D_f \bar{c}_1)$, $\theta_{crit_2} < \theta_{r_2}$.

Critical treatment rate:

$$\xi_{crit} = \left(f_{crit} - \frac{c_2}{2c_1} e^{-\epsilon^{-1} \sqrt{c_1}(\theta_{crit}-\theta_{r_1})} - \frac{\bar{c}_2}{2\bar{c}_1} e^{\epsilon^{-1} \sqrt{\bar{c}_1}(\theta_{crit}-\theta_{r_2})} \right) \times$$

$$\left(\frac{1}{\beta B_2 e^{-b_2 \theta_{crit}}} - \frac{1}{2D_f c_1} e^{-\epsilon^{-1} \sqrt{c_1}(\theta_{crit}-\theta_{r_1})} - \frac{1}{2D_f \bar{c}_1} e^{\epsilon^{-1} \sqrt{\bar{c}_1}(\theta_{crit}-\theta_{r_2})} \right)^{-1}, \tag{A.59}$$

Eccentricity of the minimum trophic factor concentration:

$$0 = -\epsilon^{-1} \sqrt{c_1} \left(\frac{c_2}{2c_1} - \frac{\xi_{crit}}{2D_f c_1} \right) e^{-\epsilon^{-1} \sqrt{c_1}(\theta_{crit}-\theta_{r_1})} + \frac{\xi_{crit} b_2}{\beta B_2 e^{-b_2 \theta_{crit}}} + \epsilon^{-1} \sqrt{\bar{c}_1} \left(\frac{\bar{c}_2}{2\bar{c}_1} - \frac{\xi_{crit}}{2D_f \bar{c}_1} \right) e^{\epsilon^{-1} \sqrt{\bar{c}_1}(\theta_{crit}-\theta_{r_2})}, \tag{A.60}$$

solving Eqs. (A.59) and (A.60) simultaneously for ξ_{crit} and θ_{crit}.

Appendix A.6. Narrow patch of rod loss with global treatment

This case is the same as that in Appendix A.1, except that treatment is applied across the whole domain ($T(\theta) = 1$) at rate ξ. Thus, we decompose the domain into the same regions as in the untreated case (see Fig. 2(b)).

Left- and Right-Outer:

$$f_0(\theta) = \frac{\alpha B_3 \theta e^{-b_3 \theta} + \xi}{\beta B_2 e^{-b_2 \theta}}, \tag{A.61}$$

Left-Inner:

$$f_0(\theta) = A_1 e^{\epsilon^{-1} \sqrt{c_1}(\theta-\theta_{r_1})} + \frac{c_2}{c_1} + \frac{\xi}{D_f c_1}, \tag{A.62}$$

Centre-Inner:

$$f_0(\theta) = A_3 e^{-\epsilon^{-1} \sqrt{c_1}(\theta-\theta_{r_1})} + A_4 e^{\epsilon^{-1} \sqrt{c_1}(\theta-\theta_{r_2})} + \frac{\xi}{D_f c_1}, \tag{A.63}$$

Right-Inner:

$$f_0(\theta) = A_8 e^{-\epsilon^{-1} \sqrt{\bar{c}_1}(\theta-\theta_{r_2})} + \frac{\bar{c}_2}{\bar{c}_1} + \frac{\xi}{D_f \bar{c}_1}, \tag{A.64}$$

where:

$$A_1 = \left(1 + \sqrt{\frac{c_1}{\bar{c}_1}}\right)^{-1} \left[\frac{c_2}{2c_1}K^{-2}\left(\sqrt{\frac{c_1}{\bar{c}_1}} - 1\right) + K^{-1}\left(\frac{\bar{c}_2}{\bar{c}_1} + \frac{\xi}{D_f\bar{c}_1} - \frac{\xi}{D_f c_1}\right)\right] - \frac{c_2}{2c_1},$$

$$A_3 = \frac{c_2}{2c_1},$$

$$A_4 = \left(1 + \sqrt{\frac{c_1}{\bar{c}_1}}\right)^{-1} \left[\frac{c_2}{2c_1}K^{-1}\left(\sqrt{\frac{c_1}{\bar{c}_1}} - 1\right) + \left(\frac{\bar{c}_2}{\bar{c}_1} + \frac{\xi}{D_f\bar{c}_1} - \frac{\xi}{D_f c_1}\right)\right],$$

$$A_8 = -\left(1 + \sqrt{\frac{c_1}{\bar{c}_1}}\right)^{-1} \left[\frac{c_2}{2c_1}K^{-1}\left(\sqrt{\frac{c_1}{\bar{c}_1}} - 1\right) + \left(\frac{\bar{c}_2}{\bar{c}_1} + \frac{\xi}{D_f\bar{c}_1} - \frac{\xi}{D_f c_1}\right)\right]\sqrt{\frac{c_1}{\bar{c}_1}} + \frac{c_2}{2c_1}\sqrt{\frac{c_1}{\bar{c}_1}}K^{-1},$$

Left-Composite:

$$f_{0_{left-comp}}(\theta) = \frac{\alpha B_3 \theta e^{-b_3\theta} + \xi}{\beta B_2 e^{-b_2\theta}} + A_1 e^{\epsilon^{-1}\sqrt{c_1}(\theta-\theta_{r_1})}, \tag{A.65}$$

Centre-Composite is identical with the centre-inner.
Right-Composite:

$$f_{0_{right-comp}}(\theta) = \frac{\alpha B_3 \theta e^{-b_3\theta} + \xi}{\beta B_2 e^{-b_2\theta}} + A_8 e^{-\epsilon^{-1}\sqrt{\bar{c}_1}(\theta-\theta_{r_2})}, \tag{A.66}$$

Minimal Cone Degenerate Patch Left Boundary:

$$\theta_{crit_1} = \theta_{r_1} + \frac{\epsilon}{\sqrt{c_1}}\log\left(A_1^{-1}\left(f_{crit} - \frac{c_2}{c_1} - \frac{\xi}{D_f c_1}\right)\right), \text{ for } f_{crit} \geq A_1 + \frac{c_2}{c_1} + \frac{\xi}{D_f c_1}, \tag{A.67}$$

$$f_{crit} = A_3 e^{-\epsilon^{-1}\sqrt{c_1}(\theta_{crit_1}-\theta_{r_1})} + A_4 e^{\epsilon^{-1}\sqrt{c_1}(\theta_{crit_1}-\theta_{r_2})} + \frac{\xi}{D_f c_1}, \text{ for } f_{crit} \leq A_1 + \frac{c_2}{c_1} + \frac{\xi}{D_f c_1}, \tag{A.68}$$

Maximal Cone Degenerate Patch Right Boundary:

$$f_{crit} = A_3 e^{-\epsilon^{-1}\sqrt{c_1}(\theta_{crit_2}-\theta_{r_1})} + A_4 e^{\epsilon^{-1}\sqrt{c_1}(\theta_{crit_2}-\theta_{r_2})} + \frac{\xi}{D_f c_1}, \text{ for } f_{crit} \leq A_8 + \frac{\bar{c}_2}{\bar{c}_1} + \frac{\xi}{D_f \bar{c}_1}, \tag{A.69}$$

$$\theta_{crit_2} = \theta_{r_2} - \frac{\epsilon}{\sqrt{\bar{c}_1}}\log\left(A_8^{-1}\left(f_{crit} - \frac{\bar{c}_2}{\bar{c}_1} - \frac{\xi}{D_f \bar{c}_1}\right)\right), \text{ for } f_{crit} \geq A_8 + \frac{\bar{c}_2}{\bar{c}_1} + \frac{\xi}{D_f \bar{c}_1}, \tag{A.70}$$

where Eqs. (A.68) and (A.69) must be solved implicitly for θ_{crit_1} and θ_{crit_2} respectively,

- when $f_{crit} \geq A_1 + \frac{c_2}{c_1} + \frac{\xi}{D_f c_1}$, $\theta_{crit_1} \leq \theta_{r_1}$;

- when $f_{crit} < A_1 + \frac{c_2}{c_1} + \frac{\xi}{D_f c_1}$, $\theta_{crit_1} > \theta_{r_1}$;

- when $f_{crit} \geq A_8 + \frac{\bar{c}_2}{\bar{c}_1} + \frac{\xi}{D_f \bar{c}_1}$, $\theta_{crit_2} \geq \theta_{r_2}$;

- when $f_{crit} < A_8 + \frac{\bar{c}_2}{\bar{c}_1} + \frac{\xi}{D_f \bar{c}_1}$, $\theta_{crit_2} < \theta_{r_2}$.

Critical treatment rate:

$$\xi_{crit} = \left(f_{crit} - \left(1 + \sqrt{\frac{c_1}{\bar{c}_1}}\right)^{-1} \left[\frac{c_2}{2c_1} K^{-2} \left(\sqrt{\frac{c_1}{\bar{c}_1}} - 1\right) + K^{-1}\frac{\bar{c}_2}{\bar{c}_1} \right] e^{\epsilon^{-1}\sqrt{c_1}(\theta_{crit}-\theta_{r_1})} - \frac{c_2}{2c_1} e^{-\epsilon^{-1}\sqrt{c_1}(\theta_{crit}-\theta_{r_1})} \right) \times$$

$$\left(\frac{1}{D_f c_1} + \left(1 + \sqrt{\frac{c_1}{\bar{c}_1}}\right)^{-1} K^{-1} \left(\frac{1}{D_f \bar{c}_1} - \frac{1}{D_f c_1}\right) e^{\epsilon^{-1}\sqrt{c_1}(\theta_{crit}-\theta_{r_1})} \right)^{-1}, \tag{A.71}$$

Eccentricity of the minimum trophic factor concentration:

$$\theta_{crit} = \theta_{r_1} - \frac{\epsilon}{2\sqrt{c_1}} \log\left(\frac{2c_1}{c_2}\left(1 + \sqrt{\frac{c_1}{\bar{c}_1}}\right)^{-1} \left[\frac{c_2}{2c_1} K^{-2}\left(\sqrt{\frac{c_1}{\bar{c}_1}} - 1\right) + K^{-1}\left(\frac{\bar{c}_2}{\bar{c}_1} + \frac{\xi_{crit}}{D_f \bar{c}_1} - \frac{\xi_{crit}}{D_f c_1}\right) \right] \right), \tag{A.72}$$

solving Eqs. (A.71) and (A.72) simultaneously for ξ_{crit} and θ_{crit}.

Appendix A.7. Narrow patch of rod loss with local treatment

This case is the same as that in Appendix A.6, except that treatment is only applied locally, within the degenerate rod patch ($T(\theta) = 1 - F(\theta)$), rather than globally, across the whole domain. We decompose the domain into the same regions as in the untreated and global treatment cases (see Fig. 2(b)).

Left- and Right-Outer:

$$f_0(\theta) = \frac{\alpha B_3 \theta e^{-b_3\theta}}{\beta B_2 e^{-b_2\theta}}, \tag{A.73}$$

Left-Inner:

$$f_0(\theta) = A_1 e^{\epsilon^{-1}\sqrt{c_1}(\theta-\theta_{r_1})} + \frac{c_2}{c_1}, \tag{A.74}$$

Centre-Inner:

$$f_0(\theta) = A_3 e^{-\epsilon^{-1}\sqrt{c_1}(\theta-\theta_{r_1})} + A_4 e^{\epsilon^{-1}\sqrt{c_1}(\theta-\theta_{r_2})} + \frac{\xi}{D_f c_1}, \tag{A.75}$$

Right-Inner:

$$f_0(\theta) = A_8 e^{-\epsilon^{-1}\sqrt{\bar{c}_1}(\theta-\theta_{r_2})} + \frac{\bar{c}_2}{\bar{c}_1}, \tag{A.76}$$

where:

$$A_1 = \left(1 + \sqrt{\frac{c_1}{\bar{c}_1}}\right)^{-1} \left[\frac{K^{-2}}{2}\left(\frac{c_2}{c_1} - \frac{\xi}{D_f c_1}\right)\left(\sqrt{\frac{c_1}{\bar{c}_1}} - 1\right) + K^{-1}\left(\frac{\bar{c}_2}{\bar{c}_1} - \frac{\xi}{D_f c_1}\right) \right] - \frac{1}{2}\left(\frac{c_2}{c_1} - \frac{\xi}{D_f c_1}\right),$$

$$A_3 = \frac{1}{2}\left(\frac{c_2}{c_1} - \frac{\xi}{D_f c_1}\right),$$

$$A_4 = \left(1 + \sqrt{\frac{c_1}{\bar{c}_1}}\right)^{-1} \left[\frac{K^{-1}}{2}\left(\frac{c_2}{c_1} - \frac{\xi}{D_f c_1}\right)\left(\sqrt{\frac{c_1}{\bar{c}_1}} - 1\right) + \left(\frac{\bar{c}_2}{\bar{c}_1} - \frac{\xi}{D_f c_1}\right) \right],$$

$$A_8 = -\left(1 + \sqrt{\frac{c_1}{\bar{c}_1}}\right)^{-1} \left[\frac{K^{-1}}{2}\left(\frac{c_2}{c_1} - \frac{\xi}{D_f c_1}\right)\left(\sqrt{\frac{c_1}{\bar{c}_1}} - 1\right) + \left(\frac{\bar{c}_2}{\bar{c}_1} - \frac{\xi}{D_f c_1}\right) \right]\sqrt{\frac{c_1}{\bar{c}_1}} + \sqrt{\frac{c_1}{\bar{c}_1}}\frac{K^{-1}}{2}\left(\frac{c_2}{c_1} - \frac{\xi}{D_f c_1}\right),$$

Left-Composite:

$$f_{0_{left-comp}}(\theta) = \frac{\alpha B_3 \theta e^{-b_3\theta}}{\beta B_2 e^{-b_2\theta}} + A_1 e^{\epsilon^{-1}\sqrt{c_1}(\theta-\theta_{r_1})}, \tag{A.77}$$

Centre-Composite is identical with the centre-inner.
Right-Composite:

$$f_{0_{right-comp}}(\theta) = \frac{\alpha B_3 \theta e^{-b_3\theta}}{\beta B_2 e^{-b_2\theta}} + A_8 e^{-\epsilon^{-1}\sqrt{c_1}(\theta-\theta_{r_2})}, \tag{A.78}$$

The Left- and Right-Composite solutions are the same here as for the narrow patch without treatment, except that the expressions for A_1 and A_8 are different.
Minimal Cone Degenerate Patch Left Boundary:

$$\theta_{crit_1} = \theta_{r_1} + \frac{\epsilon}{\sqrt{c_1}} \log\left(A_1^{-1}\left(f_{crit} - \frac{c_2}{c_1}\right)\right), \text{ for } f_{crit} \geq A_1 + \frac{c_2}{c_1}, \tag{A.79}$$

$$f_{crit} = A_3 e^{-\epsilon^{-1}\sqrt{c_1}(\theta_{crit_1}-\theta_{r_1})} + A_4 e^{\epsilon^{-1}\sqrt{c_1}(\theta_{crit_1}-\theta_{r_2})}, \text{ for } f_{crit} \leq A_1 + \frac{c_2}{c_1}, \tag{A.80}$$

Maximal Cone Degenerate Patch Right Boundary:

$$f_{crit} = A_3 e^{-\epsilon^{-1}\sqrt{c_1}(\theta_{crit_2}-\theta_{r_1})} + A_4 e^{\epsilon^{-1}\sqrt{c_1}(\theta_{crit_2}-\theta_{r_2})}, \text{ for } f_{crit} \leq A_8 + \frac{\bar{c}_2}{\bar{c}_1}, \tag{A.81}$$

$$\theta_{crit_2} = \theta_{r_2} - \frac{\epsilon}{\sqrt{\bar{c}_1}} \log\left(A_8^{-1}\left(f_{crit} - \frac{\bar{c}_2}{\bar{c}_1}\right)\right), \text{ for } f_{crit} \geq A_8 + \frac{\bar{c}_2}{\bar{c}_1}, \tag{A.82}$$

where Eqs. (A.80) and (A.81) must be solved implicitly for θ_{crit_1} and θ_{crit_2} respectively,

- when $f_{crit} \geq A_1 + \frac{c_2}{c_1}$, $\theta_{crit_1} \leq \theta_{r_1}$;

- when $f_{crit} < A_1 + \frac{c_2}{c_1}$, $\theta_{crit_1} > \theta_{r_1}$;

- when $f_{crit} \geq A_8 + \frac{\bar{c}_2}{\bar{c}_1}$, $\theta_{crit_2} \geq \theta_{r_2}$;

- when $f_{crit} < A_8 + \frac{\bar{c}_2}{\bar{c}_1}$, $\theta_{crit_2} < \theta_{r_2}$.

Critical treatment rate:

$$\xi_{crit} = \left(f_{crit} - \left(1 + \sqrt{\frac{c_1}{\bar{c}_1}}\right)^{-1}\left[\frac{c_2}{2c_1}K^{-2}\left(\sqrt{\frac{c_1}{\bar{c}_1}} - 1\right) + K^{-1}\frac{\bar{c}_2}{\bar{c}_1}\right]e^{\epsilon^{-1}\sqrt{c_1}(\theta_{crit}-\theta_{r_1})} - \frac{c_2}{2c_1}e^{-\epsilon^{-1}\sqrt{c_1}(\theta_{crit}-\theta_{r_1})}\right) \times$$

$$\left(\left(1 + \sqrt{\frac{c_1}{\bar{c}_1}}\right)^{-1}\left[-\left(\sqrt{\frac{c_1}{\bar{c}_1}} - 1\right)\frac{K^{-2}}{2D_f c_1} - \frac{K^{-1}}{D_f c_1}\right]e^{\epsilon^{-1}\sqrt{c_1}(\theta_{crit}-\theta_{r_1})} - \frac{1}{2D_f c_1}e^{-\epsilon^{-1}\sqrt{c_1}(\theta_{crit}-\theta_{r_1})} + \frac{1}{D_f c_1}\right)^{-1}, \tag{A.83}$$

Eccentricity of the minimum trophic factor concentration:

$$\theta_{crit} = \theta_{r_1} - \frac{\epsilon}{2\sqrt{c_1}} \log\left(2\left(\frac{c_2}{c_1} - \frac{\xi_{crit}}{D_f c_1}\right)^{-1}\left(1 + \sqrt{\frac{c_1}{\bar{c}_1}}\right)^{-1}\left[\left(\frac{c_2}{c_1} - \frac{\xi_{crit}}{D_f c_1}\right)\frac{K^{-2}}{2}\left(\sqrt{\frac{c_1}{\bar{c}_1}} - 1\right) + K^{-1}\left(\frac{\bar{c}_2}{\bar{c}_1} - \frac{\xi_{crit}}{D_f c_1}\right)\right]\right), \tag{A.84}$$

solving Eqs. (A.83) and (A.84) simultaneously for ξ_{crit} and θ_{crit}.

References

Bender, C.M., Orszag, S.A., 1999. Advanced Mathematical Methods for Scientists and Engineers I: Asymptotic Methods and Perturbation Theory. Springer.

Burns, J., Clarke, G., Lumsden, C.J., 2002. Photoreceptor death: Spatiotemporal patterns arising from one-hit death kinetics and a diffusible cell death factor. Bull. Math. Biol. 64, 1117–1145.

Camacho, E.T., Colón Vélez, M.A., Hernández, D.J., Bernier, U.R., van Laarhoven, J., Wirkus, S., 2010. A mathematical model for photoreceptor interactions. J. Theor. Biol. 267, 638–646.

Camacho, E.T., Léveillard, T., Sahel, J.A., Wirkus, S., 2016a. Mathematical model of the role of RdCVF in the coexistence of rods and cones in a healthy eye. Bull. Math. Biol. 78, 1394–1409.

Camacho, E.T., Melara, L.A., Villalobos, M.C., Wirkus, S., 2014. Optimal control in the treatment of retinitis pigmentosa. Bull. Math. Biol. 76, 292–313.

Camacho, E.T., Punzo, C., Wirkus, S.A., 2016b. Quantifying the metabolic contribution to photoreceptor death in retinitis pigmentosa via a mathematical model. J. Theor. Biol. 408, 75–87.

Camacho, E.T., Radulescu, A., Wirkus, S., 2016c. Bifurcation analysis of a photoreceptor interaction model for retinitis pigmentosa. Commun. Nonlinear Sci. Numer. Simulat. 38, 267–276.

Camacho, E.T., Wirkus, S., 2013. Tracing the progression of retinitis pigmentosa via photoreceptor interactions. J. Theor. Biol. 317, 105–118.

Chrysostomou, V., Stone, J., Stowe, S., Barnett, N.L., Valter, K., 2008. The status of cones in the rhodopsin mutant P23H-3 retina: Light-regulated damage and repair in parallel with rods. Invest. Ophthalmol. Vis. Sci. 49, 1116–1125. doi:`https://doi.org/10.1167/iovs.07-1158`.

Colón Vélez, M.A., Hernández, D.J., Bernier, U.R., van Laarhoven, J., Camacho, E.T., 2003. Mathematical Models for Photoreceptor Interactions. Technical Report. Cornell University, Department of Biological Statistics and Computational Biology.

Curcio, C.A., Millican, C.L., Allen, K.A., Kalina, R.E., 1993. Aging of the human photoreceptor mosaic: evidence for selective vulnerability of rods in central retina. Invest. Ophthalmol. Vis. Sci. 34, 3278–3296.

Curcio, C.A., Sloan, K.R., Kalina, R.E., Hendrickson, A.E., 1990. Human photoreceptor topography. J. Comp. Neurol. 292, 497–523. doi:`10.1002/cne.902920402`.

Dörrbaum, A.R., Kochen, L., Langer, J.D., Schuman, E.M., 2018. Local and global influences on protein turnover in neurons and glia. eLife 7, e34202. doi:`10.7554/eLife.34202`.

Eden, E., Geva-Zatorsky, N., Issaeva, I., Cohen, A., Dekel, E., Danon, T., Cohen, L., Mayo, A., Alon, U., 2011. Proteome half-life dynamics in living human cells. Science 331, 764–768. doi:`10.1126/science.1199784`.

Fintz, A.C., Audo, I., Hicks, D., Mohand-Saïd, S., Léveillard, T., Sahel, J., 2003. Partial characterization of retina-derived cone neuroprotection in two culture models of photoreceptor degeneration. Invest. Ophthalmol. Vis. Sci. 44, 818–825.

Grover, S., Fishman, G.A., Brown Jr, J., 1998. Patterns of visual field progression in patients with retinitis pigmentosa. Ophthalmology 105, 1069–1075.

Guérin, C.J., Lewis, G.P., Fisher, S.K., Anderson, D.H., 1993. Recovery of photoreceptor outer segment length and analysis of membrane assembly rates in regenerating primate photoreceptor outer segments. Invest. Ophthalmol. Vis. Sci. 34, 175–183.

Gupta, N., Brown, K.E., Milam, A.H., 2003. Activated microglia in human retinitis pigmentosa, late-onset retinal degeneration, and age-related macular degeneration. Exp. Eye Res. 76, 463–471.

Hamel, C., 2006. Retinitis pigmentosa. Orphanet. J. Rare Dis. 1, 40.

Hartong, D.T., Berson, E.L., Dryja, T.P., 2006. Retinitis pigmentosa. Lancet 368, 1795–1809.

Jürgens, K.D., Peters, T., Gros, G., 1994. Diffusivity of myoglobin in intact skeletal muscle cells. Proc. Natl. Acad. Sci. 91, 3829–3833. doi:`10.1073/pnas.91.9.3829`.

Kocaoglu, O.P., Liu, Z., Zhang, F., Kurokawa, K., Jonnal, R.S., Miller, D.T., 2016. Photoreceptor disc shedding in the living human eye. Biomed. Opt. Express 7, 4554–4568. doi:`10.1364/BOE.7.004554`.

Lee, D.C., Vazquez-Chona, F.R., Ferrell, W.D., Tam, B.M., Jones, B.W., Marc, R.E., Moritz, O.L., 2012. Dysmorphic photoreceptors in a P23H mutant rhodopsin model of retinitis pigmentosa are metabolically active and capable of regenerating to reverse retinal degeneration. J. Neurosci. 32, 2121–2128. doi:`https://doi.org/10.1523/JNEUROSCI.4752-11.2012`.

Léveillard, T., Mohand-Saïd, S., Lorentz, O., Hicks, D., Fintz, A.C., Clérin, E., Simonutti, M., Forster, V., Cavusoglu, N., Chalmel, F., Dollé, P., Poch, O., Lambrou, G., Sahel, J.A., 2004. Identification and characterization of rod-derived cone viability factor. Nat. Genet. 36, 755–759. doi:`https://doi.org/10.1038/ng1386`.

Mohand-Saïd, S., Deudon-Combe, A., Hicks, D., Simonutti, M., Forster, V., Fintz, A.C., Léveillard, T., Dreyfus, H., Sahel, J.A., 1998. Normal retina releases a diffusible factor stimulating cone survival in the retinal degeneration mouse. Proc. Natl. Acad. Sci. 95, 8357–8362.

Mohand-Saïd, S., Hicks, D., Dreyfus, H., Sahel, J.A., 2000. Selective transplantation of rods delays cone loss in a retinitis pigmentosa model. Arch. Ophthalmol. 118, 807–811.

Mohand-Saïd, S., Hicks, D., Simonutti, M., Tran-Minh, D., Deudon-Combe, A., Dreyfus, H., Silverman, M.S., Ogilvie, J.M., Tenkova, T., Sahel, J., 1997. Photoreceptor transplants increase host cone survival in the retinal degeneration (rd) mouse. Ophthalmic Res. 29, 290–297.

Oyster, C.W., 1999. The Human Eye: Structure and Function. Sinauer Associates Inc.

Punzo, C., Kornacker, K., Cepko, C.L., 2009. Stimulation of the insulin/mTOR pathway delays cone death in a mouse model of retinitis pigmentosa. Nat. Neurosci. 12, 44–52.

Ripps, H., 2002. Cell death in retinitis pigmentosa: Gap junctions and the 'bystander' effect. Exp. Eye Res. 74, 327–336.

Roberts, P.A., Gaffney, E.A., Luthert, P.J., Foss, A.J.E., Byrne, H.M., 2016. Mathematical and computational models of the retina in health, development and disease. Prog. Retin. Eye. Res. 53, 48–69. doi:`https://doi.org/10.1016/j.preteyeres.2016.04.001`.

Roberts, P.A., Gaffney, E.A., Luthert, P.J., Foss, A.J.E., Byrne, H.M., 2017. Mathematical models of retinitis pigmentosa: The oxygen toxicity hypothesis. J. Theor. Biol. 425, 53–71. doi:`https://doi.org/10.1016/j.jtbi.2017.05.006`.

Roberts, P.A., Gaffney, E.A., Whiteley, J.P., Luthert, P.J., Foss, A.J.E., Byrne, H.M., 2018. Predictive mathematical models for the spread and treatment of hyperoxia-induced photoreceptor degeneration in retinitis pigmentosa. Invest. Ophthalmol. Vis. Sci. 59, 1238–1249. doi:`10.1167/iovs.17-23177`.

Stone, J., Maslim, J., Valter-Kocsi, K., Mervin, K., Bowers, F., Chu, Y., Barnett, N., Provis, J., Lewis, G., Fisher, S.K., Bistid, S., Gargini, C., Cervetto, L., Merin, S., Pe'er, J., 1999. Mechanisms of photoreceptor death and survival in mammalian retina. Prog. Retin. Eye Res. 18(6), 689–735.

Yang, Y., Mohand-Said, S., Danan, A., Simonutti, M., Fontaine, V., Clerin, E., Picaud, S., Leveillard, T., Sahel, J.A., 2009. Functional cone rescue by RdCVF protein in a dominant model of retinitis pigmentosa. Mol. Ther. 17, 787–795.

Young, R.W., 1967. The renewal of photoreceptor cell outer segments. J. Cell Biol. 33, 61–72.

Young, R.W., 1971. The renewal of rod and cone outer segments in the rhesus monkey. J. Cell Biol. 49, 303–318.

Young, R.W., 1978. The daily rhythm of shedding and degradation of rod and cone outer segment membranes in the chick retina. Invest. Ophthalmol. Vis. Sci. 17, 105–16.

Young, R.W., Bok, D., 1969. Participation of the retinal pigment epithelium in the rod outer segment renewal process. J. Cell Biol. 42, 392–403.