Lipid membranes modulate the activity of RNA through sequence-specific interactions

Authors: Tomasz Czerniak[a] and James P Saenz[a]#

[a] Technische Universität Dresden, B CUBE Center for Molecular Bioengineering, 01307 Dresden, Germany

#corresponding author and lead contact: james.saenz@tu-dresden.de

Abstract

RNA is a ubiquitous biomolecule that can serve as both catalyst and information carrier. Understanding how RNA activity is controlled and how it in turn regulates bioactivity is crucial for elucidating its physiological roles and potential applications in synthetic biology. Here we show that lipid membranes can act as RNA organization platforms, introducing a novel mechanism for ribo-regulation. The activity of R3C ribozyme can be modified by the presence of lipid membranes, with direct RNA-lipid interactions dependent on RNA sequence, structure and length. In particular, the presence of guanine in short RNAs is crucial for RNA-lipid interactions, while double-stranded RNAs further increase lipid-binding affinity. Lastly, by artificially modifying the R3C-substrate sequence to enhance membrane binding we unexpectedly generated a lipid-sensitive riboswitch. These findings introduce RNA-lipid interactions as a tool for developing riboswitches and novel RNA-based lipid biosensors, and bear significant implications for RNA World scenarios for the origin of life.

Introduction

RNA performs diverse functions ranging from information storage, to regulation of other biomolecules, to directly catalysing biochemical reactions. The functional versatility of RNA has implications for understanding plausible scenarios for the origin of life[1–3], and for developing tools in synthetic biology[4–6]. Research aimed at understanding how an RNA World could have emerged has motivated development of ribozymes with functions including RNA ligation[7–10], replication[11,12] and other activities[13–16]. The experimental development of functional RNAs raises the possibility of RNA-based synthetic systems. For both synthetic life and understanding the origin of self-replicating organisms, RNA has intrinsic appeal: it can serve functions of both DNA (information storage) and proteins (enzymes), obviating the need for translation machineries and protein chaperones. Furthermore, RNA does not undergo irreversible denaturation like proteins, lending robustness against a broad range of physical and chemical conditions. In order to design a system based on RNA it is essential to be able to coordinate RNA activity in space and time.

Key to harnessing the functional versatility of RNA is in understanding how to spatially and temporally modulate its properties and to selectively modulate the activity of different RNAs within one system. The physicochemical environment surrounding an RNA molecule is a central determinant of its structure, stability and activity. Spontaneous RNA hydrolysis and ligation, as well as the catalytic RNA activity are sensitive to pH[17], ionic strength[18,19] and RNA concentration changes[11], among other parameters. Similarly, RNA activity can be modulated by interactions with molecules such as ions, proteins, and other nucleic acids[20]. Thus, one approach to determining RNA activity could be via

45   tunable interactions with binding partners that affect RNA's structure, concentration or chemical
46   microenvironment.
47
One mechanism for modulating RNA activity could be through direct RNA-lipid interactions[21–23].
Because of their amphiphilic nature, lipids spontaneously self-assemble into membranous structures
that can encapsulate RNA into protected and selective microcompartments[24–26]. Alternatively, direct
RNA-lipid interactions could localize RNAs to the membrane surfaces, increasing its local concentration
and reducing dimensionality for intermolecular interactions[21]. Lastly, localization to a lipid surface
brings RNA into a physicochemically unique microenvironment with sharp gradients of hydrophobicity,
electrical permittivity, and water activity. Through these effects, RNA-lipid interactions could provide
a powerful mechanism for modulating RNA activity.
56
The first functional RNA-lipid interaction was described more than 40 years ago[27], with subsequent
research revealing various factors that facilitate nucleic acid-lipid binding[28–38]. More recently, specific
RNA sequences have been generated through SELEX with affinity for fluid membranes comprised of
phospholipids and cholesterol[39–41]. Interestingly, mixtures of RNAs were also shown to bind to
membranes that are in a solid crystalline (gel) phase[42,43]. These studies revealed that while most
randomized mixtures of RNA sequences can bind to gel membranes, there is a relatively small chemical
space of oligomers that have affinity for fluid membranes. Thus, conceptually, gel membranes could
provide a platform for modulating the activity of a diverse range of RNAs. However, effects of gel
membranes on RNA activity and the sequence selectivity of such interactions are relatively unexplored.
66
This study reports the effect of lipid membranes on RNA catalytic activity. We show that RNA-lipid
binding depends on the RNA primary sequence, secondary structure, and length. Using the trans-acting
R3C ligase ribozyme we observed that R3C-lipid binding changes ribozyme activity in a concentration-
dependent manner. Lipid binding assays show that the interaction of short RNA sequences with gel
membranes is sequence-specific and depends on guanine content and presence of double-stranded
structures. Lastly, modification of R3C's substrate sequence increased tunability of the R3C based
reactions, through a lipid-dependent riboswitch. Our findings prove that membranes can serve as
platforms for RNA-activity modulation, which can contribute to the development of RNA-based
biosensors and riboswitches. This approach introduces new tools for molecular and synthetic biology
and raises the prospect of previously unrecognized roles for RNA-lipid interactions in the origin and
evolution of life.
78

Results

80
RNA-lipid binding changes probability of RNA-RNA interactions
Building on previous observations that RNA binds to lipid membranes[27,39–42], we hypothesized that
RNA-membrane interactions could influence ribozyme activity by changes of RNA local concentrations
on the membrane surface and/or influencing RNA conformations. Indeed, RNA oligonucleotides can
bind to lipid membranes in the gel phase in the presence of the divalent cations, leading to aggregation
into large RNA-lipid assemblies (**Fig. 1a**) [28,36,42,43]. Depletion of divalent ions or increasing temperature
above the lipid melting temperature (i.e. producing fluid instead of gel membranes) reduces RNA-
membrane binding and aggregation. This reversible aggregation could provide a tunable mechanism
to concentrate RNAs. To test this, we asked whether RNAs are in fact concentrated in the presence of
gel membranes, and how that influences trans-acting-ribozyme activity.

To determine changes in local RNA concentration upon interaction with lipid membranes, we relied
on UV-mediated crosslinking. All membranes in this study were comprised of
dipalmitoylphosphatidylcholine (DPPC) which are in a solid crystalline (gel) phase at the chosen
experimental temperature of 24 °C. Nucleic acid bases absorb UV light, producing chemical changes
that yield base-base covalent bonds in a distance dependent manner, yielding insights into the
structure and interactions of nucleic acids[44–46]. We first observed the effect of gel-membranes on the
crosslinking of a single defined RNA sequence, the R3C ligase, which predominately forms one
structure (R3C ligase, **Fig. 1d**). When subjected to UV, the R3C ligase shows a slight increase of
crosslinking in the presence of low lipid concentrations compared to the reaction in the absence of
lipids. A further increase of lipid concentration led to a decrease of crosslinking efficiency, possibly
through dilution of the RNA species on the surface of the lipid membranes. This effect was inversely
correlated with RNA-lipid binding efficiency (**Fig. 1b**). On the other hand, a mixture of RNA oligomers
with randomized sequences which is able to create a more diverse range of intermolecular structures
(**Fig. 1d**) showed continuously increasing and overall higher UV-crosslinking efficiency in the presence
of lipids (**Fig. 1c**). The contrasting effect of crosslinking for R3C and randomized oligomers might be
explained by the influence of lipid membranes on RNA interactions: membrane binding could
potentially stabilize base-pairing[47] and thereby enhance crosslinking efficiency, whereas the decrease
of crosslinking might be caused by a dilution effect at higher membrane surface areas or lipid-based
steric hindrance. These observations demonstrate that RNA-membrane interactions can influence
RNA-RNA interactions in a manner that is dependent on lipid concentration and RNA sequence
diversity. This further suggests that membrane binding could have an effect on trans-acting ribozyme
activity derived from base pairing.

RNA-lipid interactions influence RNA catalytic activity
To investigate the functional consequence of RNA-lipid binding, we tested the effect of lipids on the
trans-acting R3C ligase ribozyme activity. R3C ligase is a relatively short ribozyme that catalyzes ligation
of substrate strands to the ribozyme[10] (**Fig. 2a**). Since R3C is also part of an RNA self-replication system
[11,12], the effects of lipids on R3C is also interesting with regard to the emergence, evolution and artificial
synthesis of autonomous self-replicating systems. The R3C reaction rate in the absence of membranes
was 11 pM/min, with reaction constant rate of k = 6.4×10^{-4} min^{-1}**.** Addition of lipid vesicles led to an
increase of the reaction rate (14 pM/min, +29%), which could plausibly be due to increased ligase-
substrate interaction on the membrane either through increased concentrations at the membrane
surface, or through stabilization of base pairing[47]. At the highest lipid concentration the reaction rate
dropped to 8 pM/min (-27%) (**Fig. 2b**). A decrease in reaction rate was correlated with further R3C
ligase binding to the lipid membranes (>100 lipid/R3C, **Fig. 1b**) and we speculate that this may be
caused by steric hindrance from the lipid vesicles. These results show that gel-membranes can
influence the reaction rate of a trans-acting ribozyme in a lipid concentration-dependent manner.

R3C ligase and its substrate bind differently to the membrane
Previous findings determined effect of lipid-R3C ligase interaction on its activity. To further understand
how R3C and substrate membrane binding might contribute to changes in ligation rates, lipid-binding
affinities of the ligase and its substrate were determined. To quantify binding to the membranes we
calculated lipid – buffer partition coefficients (K). The short 12 nt substrate shows a more than 3 orders
of magnitude lower partition coefficient compared with the 84 nt R3C ligase (**Fig. 2c**). Thus, in this
system substrate-membrane binding is essentially negligible compared with R3C-membrane binding,
suggesting that the increased activity observed in **Figure 2b** was due to an effect of the membrane on

the R3C ligase-substrate complex stability, rather than through enhanced substrate-lipid interactions
on the membrane surface.
139
The large difference in membrane partitioning between the R3C ligase and its substrate raised the
possibility that RNA features such as length, base composition or structure could influence membrane
binding (**Fig. 2c**). Understanding the features that influence membrane interactions could allow us to
engineer RNAs with higher membrane affinity, and thereby potentially enhance activity by
concentrating RNAs at the membrane surface. Consequently, we next investigated what features of
RNA influence its interaction with lipid membranes.
146
147 **RNA sequence influences binding to gel membranes**
148 Interactions of RNA with gel phase membranes have been previously shown to exhibit low sequence-
149 specificity, based on binding of oligos with randomized sequences[42,43]. However, the large difference
150 in binding for R3C and its substrate suggested to us that RNA-gel membrane interactions might in fact
151 be dependent on RNA composition or structure.
152
We first evaluated how the base content influences membrane binding of short RNA species. 19nt
oligomers with only one type of nucleotide were tested together with a random sequence as a control.
To quantify relative changes of RNA-membrane binding, we calculated the RNA binding efficiency as
the percentage of total RNA bound to membranes at a fixed oligo to lipid ratio. Remarkably, while the
19xG oligomer binds very efficiently to the gel phase membrane (84% binding efficiency), oligomers of
A, C or U showed practically no binding. A mixture of 19-mer RNA with randomized sequences showed
moderate binding (33% binding efficiency), as did one with AG-repeats (**Fig. 3a**), most probably due to
the introduction of guanine. Guanine could conceivably enhance binding either through direct
interactions with the membrane, or alternatively by promoting G≡C base pairing, thereby influencing
structure. The enhanced binding of an oligomer with only AG-repeats suggests that membrane binding
can be influenced by direct guanine-membrane interactions.
164
We next examined the effect of deleting a specific base from mixtures of 40-mer oligos with
randomized sequences. Depletion of A or U led to slight decrease of membrane binding, whereas
removal of G or C showed a significant binding decrease with highest effect for guanine-depleted
oligomers. The decrease in binding affinity in all of the base-depleted oligos indicated that base-pairing
may influence membrane binding (**Fig. 3b**). However, the larger effect of G and C depletion indicates
that base pairing is not the only factor responsible for membrane binding consistent with our previous
observation that G-content alone can influence binding. Finally, by comparing 19-mer and 40-mer
randomized oligos, we observe that increasing RNA length also led to enhanced binding, possibly due
to increased length and G content (**Fig. 3 a,b**).
174
To further determine how guanine content affects membrane binding, lipid-buffer partition
coefficients were determined for short oligomers with varying G-content. RNA membrane-buffer
partition coefficients indicate how well a particular molecule binds to the membrane by comparing the
relative amount of RNA in buffer and on the lipid membrane. The RNA oligos measured in this study
had partition coefficients around 10^5, which is in the range of partition coefficients determined for
hydrophobic peptides[48]. Remarkably, addition of 2 G's per oligo led to a 5-fold increase of the partition
coefficient relative to guanine-depleted RNA. Oligos with 4 guanine residues showed the highest
partition coefficient, plateauing with additional guanosines and a small decrease of binding for 7

guanine/oligo. This result shows that the presence of guanine affects RNA-lipid binding and even low
guanine content significantly increases membrane binding efficiency (**Fig. 3c**).

RNA-membrane binding is dependent on RNA structure
Our observations of the effect of base depletion on oligomer binding suggested that base-pairing may
be a key factor in membrane binding. To investigate the effect of base-pairing-derived secondary
structures on membrane interactions, we estimated binding efficiency for ssRNA and dsRNA
(composed of 100% complementary sequences) species. We observed that not only was the
membrane binding efficiency higher for dsRNA compared with ssRNA species, but also that only dsRNA
promoted vesicle aggregation (**Fig. 4a**). It is interesting to note that the difference in binding efficiency
of the two ssRNAs, which have identical base composition but different sequence, is consistent with
the importance of sequence as key determinate of membrane binding. We further observed that
dsRNA interacts with fewer lipids than ssRNA, implying different models of binding (**Fig. 4b**). Thus,
membrane binding efficiency can be dependent on the propensity of an RNA molecule to form intra-
or inter-molecular structures through base pairing.

To gain further insight into the effect of base pairing-dependent intra- and inter-molecular
interactions, we estimated the lipid to nucleotide binding ratios of ssRNA and dsRNA (**Fig. 4c**). By
calculating the molecular ratio of lipid/nucleotide binding we introduce a normalization factor that
allows us to directly compare membrane binding for oligos of varying length. Furthermore, varying
lipid to nucleotide stoichiometry reflects changes in the average lipid/nucleotide interaction strength.
With this estimate we compared lipid/nucleotide binding for ssRNA and dsRNA alongside oligos with
varying guanine content from our previous analysis (**Fig. 3c**). The differences between ssRNA and
dsRNA, all containing 9 guanines per oligo reiterate our previous observation that sequence variation
(with constant base content) and base pairing can both influence binding. However, the reason for a
change in lipid/nucleotide ratio for oligos with varying guanine content was not readily apparent.
Analysis of oligos with varying guanine content by native non-denaturing gel revealed that the
presence of guanine promotes a larger spread in migration, indicating that guanines promote
intermolecular interactions. These observations show that guanine can enhance binding by promoting
intermolecular interactions, and further confirm the importance of base pairing in RNA-lipid
interactions.

Modifying R3C substrate sequence enhances membrane binding and modifies reaction rates
In previous experiments we observed that the binding affinity of the short substrate was >3 orders of
magnitude lower than for the R3C ligase and that reaction rates correlated strongly with R3C-
membrane binding (**Fig. 2c, d**). We therefore reasoned that by modifying the R3C ligase substrate to
enhance membrane binding, we might be able to further modulate ligase activity through increased
RNA density at the membrane surface. Having determined that guanine content is a key factor
influencing RNA-lipid binding (**Fig. 3**), we modified the R3C substrate sequence to increase lipid
membrane binding efficiency by addition of a 5'-overhang with varying amounts of AG repeats (**Fig.**
5a).

We first assessed the effect of modifying the R3C ligase substrate on ligation rate in the absence of
membranes (**Fig. 5a**). We observed that increasing the guanine content decreased ligase activity,
suggesting that 5' modification of the substrate was inhibiting the reaction either through steric
hindrance, or promotion of inhibitory inter- or intra-molecular interactions. To determine if AG-rich

substrates are forming intra-molecular structures[49] and/or exhibiting reduced binding with the R3C
ligase, electrophoresis in native conditions with and without ligase was conducted (**Fig. 5b**). Firstly, the
non-modified short substrate migrated as one band and was fully bound to the R3C ligase. Secondly,
4x(AG) 20 nt substrate variant migrates as two bands, which suggests that part of the substrate
molecules have some folding, which might bias binding to the R3C ligase – in the presence of ligase
30% of the substrate was non-bound, which correlated with a 27% reaction rate drop. For the longest
substrate variant (31 nt) electrophoretic mobility suggested higher folding, since only one band is
present and migrates half-way between the non-folded 12 nt and 20 nt substrates. 70% of the 9x(AG)
substrate is not bound to the ligase, which correlates with a R3C reaction rate decrease. Lastly, the AG-
rich oligomer does not co-migrate with the R3C ligase, thus the inhibition effect is not based on non-
specific substrate-ligase interactions (not shown). These observations show that increasing guanine
content reduced R3C ligase-substrate binding, likely accounting for decreased activity.
241
We next determined whether varying guanine content through 5' modification of the substrate
influenced membrane binding and, consequently, ligation activity in the presence of membranes. As
expected, the longer guanine-containing R3C substrates showed significantly higher membrane
binding compared with the shorter variant (**Fig. 5c**). To determine if binding of the substrates to the
membrane enhanced ligase activity, R3C reaction rates were determined in the presence of the gel
lipid membranes (**Fig. 5d**). For modified substrates, the reaction rates increased significantly to the
level of the unmodified substrate. Interestingly, the low-G substrate (2G) shows no significant reaction
increase with the introduction of membranes, indicating that the rescue of activity of the other
guanine variants is not due to increased substrate-ligase interactions through enhanced binding to the
membrane, but rather due to interaction of the membrane with 5' overhangs. Thus, substrate-
membrane interaction increases activity by rescuing the inhibitory effect of the guanine-rich 5'
overhangs (**Fig. 5e**).
254
255 In summary, by modifying the R3C ribozyme substrate with an additional short sequence containing
256 guanine residues, we increased membrane affinity. The addition of guanines, however, also introduced
257 an inhibitory effect on ligation activity that was reversed through the introduction of gel-membranes
258 to the reaction. Thus, we unexpectedly produced a lipid-sensitive riboswitch-like system that could be
259 tuned in a lipid concentration-dependent manner.
260
261 **Discussion**
262
We demonstrated that RNA-gel membrane interactions are dependent on the primary sequence and
secondary structure of ribonucleic acids. Varying guanosine content of RNAs changed the membrane-
buffer partitioning by several orders of magnitude. Additionally, double stranded RNAs exhibited
higher membrane partitioning than single stranded RNAs, indicating that secondary structure is a
critical factor in RNA-membrane interactions. Although gel-membranes are not widely observed in
living cells, RNA-gel membrane interactions can be employed for ribo-regulation in synthetic biological
systems. Gel membranes also plausibly accumulated in ancient prebiotic scenarios, potentially serving
as an organizational scaffold, as has been proposed for "ribofilms" on mineral surfaces[50]. Furthermore,
understanding the factors involved in RNA-gel membrane interactions can provide insight for designing
RNA sequences with affinity for physiologically-relevant fluid membranes. Indeed, our finding that
guanosine is a key factor in promoting RNA-gel membrane partitioning is consistent with previous

speculation that guanosine residues might play a role in the binding of RNA aptamers for fluid
membranes[39]. Taken together, the sequence and structural dependence of RNA-gel membrane
interactions shown here provides the necessary framework to design synthetic RNAs with varying
membrane affinities.
278
Our observation of riboswitch-like behavior suggests a novel mechanism for regulation of RNA through
membrane interactions. Riboswitches are RNAs that have regulatory effects that are derived from
sensitivity to physicochemical conditions or through binding with an interaction partner. Riboswitch
regulation through temperature changes, interaction with nucleic acids and small molecules have been
described previously[51–55]. However, lipid sensitive riboswitches have not been documented. It is now
plausible that spontaneously self-assembling lipid membranes in ancient environments could have
served as an organizational scaffold regulating the activity of RNAs in a sequence and structure
dependent manner. Furthermore, the importance of guanosine in membrane binding suggests a
possible mechanism for its selection as one of the canonical nucleic acids. In extant organisms,
synthetic or naturally occurring RNAs exhibiting sensitivity to lipids could potentially act as biosensors
or riboswitches. By demonstrating that RNA-gel membrane interactions can induce riboswitch-like
behaviors, we raise the possibility that such functionalities could be an ancient mechanism of ribo-
regulation and may be present among the myriad of uncharacterized small RNAs in modern cells.
292
RNA-membrane interactions provide a basis for selection of RNAs based on sequence and structure.
The induction of RNA-membrane interactions allows spatial segregation or co-localization of RNAs
based on differential membrane affinities, affording an additional degree of control over localization
of RNAs in cell-free synthetic systems. RNA-membrane localization in a gel membrane system could be
controlled by shifting the temperature above and below the membrane gel-liquid transition
temperature, thereby turning on and off RNA-membrane binding. Membrane localization could also
be regulated dynamically through enzymatic modification[56–58] of RNA. Lipid-dependent sorting and
aggregation of RNAs could have played a role in primordial scenarios, where one of the biggest
challenges was to increase concentration of molecules. The sequence and structural selectivity of RNAs
for membranes could also be employed in the design of novel RNA membrane biosensors. Selective
RNA-membrane interactions would also influence the sequence space explored by evolving ribozymes
and may give rise to novel functions. Looking forward, RNA-membrane interactions can provide tools
for selective ribo-regulation in synthetic systems, development of membrane biosensors, and might
facilitate the emergence of novel functions through artificial and natural evolution of ribozymes in the
presence of membranes.
308
In conclusion, we reveal that lipids, which are present in every modern cell and were plausibly part of
a prebiotic world[59–61], can interact with RNA and change its activity. These findings have significant
application in fields such as synthetic biology where merging the selective affinities of aptamers with
ribozyme activity (aptazymes) is currently a developing field[62–64]. Our results provide a foundation for
exploring the basis for membrane-RNA interactions with physiologically relevant fluid membranes
exhibited by synthetic aptamers[16,65,66] thereby providing new regulatory mechanisms in synthetic
biology through the specific and anchor-free localization of RNA molecules within lipid compartments.
Nonetheless, insights from the present study can already be implemented in bioengineering
applications such as in improving mRNA drug delivery mechanisms and introducing lipid sensitive ribo-
regulation to synthetic ribozyme networks. Furthermore, the interaction between RNA and lipid
membranes could have provided an evolutionary pressure in ancient times through selection for base

composition or secondary structural features. Our observations show that RNA-lipid interactions have applications in designing ribozyme-based metabolic circuits, lipid-based riboswitches, and provide new mechanisms for evolution of function in an RNA World.

Acknowledgements:

We would like to thank Mario Mörl, Ilya Leventel, Robert Ernst, Andre Nadler, Dora Tang, Grzegorz Chwastek and Mike Thompson for helpful discussions and feedback. This work was supported by the B CUBE of the TU Dresden, a Simons Foundation Fellowship (to J.S.), a German Federal Ministry of Education and Research BMBF grant (to J.S., project 03Z22EN12), and a VW Foundation ''Life'' grant (to J.S., project 93090).

The authors declare no conflict of interests.

References

1. Pace, N. R. & Marsh, T. L. RNA catalysis and the origin of life. *Origins Life Evol B* **16**, 97–116 (1985).

2. Gilbert, W. Origin of life: The RNA world. *Nature* **319**, 618–618 (1986).

3. Rich, A. On the problems of evolution and biochemical information transfer. *Horizons in biochemistry* 103–126 (1962).

4. Breaker, R. R. Engineered allosteric ribozymes as biosensor components. *Curr Opin Biotech* **13**, 31–39 (2002).

5. Wittmann, A. & Suess, B. Engineered riboswitches: Expanding researchers' toolbox with synthetic RNA regulators. *Febs Lett* **586**, 2076–2083 (2012).

6. Isaacs, F. J., Dwyer, D. J. & Collins, J. J. RNA synthetic biology. *Nat Biotechnol* **24**, 545–554 (2006).

7. Bartel, D. & Szostak, J. Isolation of new ribozymes from a large pool of random sequences. *Science* **261**, 1411–1418 (1993).

8. Ekland, E., Szostak, J. & Bartel, D. Structurally complex and highly active RNA ligases derived from random RNA sequences. *Science* **269**, 364–370 (1995).

9. Robertson, M. P., Hesselberth, J. R. & Ellington, A. D. Optimization and optimality of a short ribozyme ligase that joins non-Watson–Crick base pairings. *Rna* **7**, 513–523 (2001).

10. Rogers, J. & Joyce, G. F. The effect of cytidine on the structure and function of an RNA ligase ribozyme. *Rna* **7**, 395–404 (2001).

11. Paul, N. & Joyce, G. F. A self-replicating ligase ribozyme. *Proc National Acad Sci* **99**,
12733–12740 (2002).

12. Lincoln, T. A. & Joyce, G. F. Self-Sustained Replication of an RNA Enzyme. *Science* **323**,
1229–1232 (2009).

13. Saragliadis, A., Krajewski, S. S., Rehm, C., Narberhaus, F. & Hartig, J. S. Thermozymes:
synthetic RNA thermometers based on ribozyme activity. *Rna Biol* **10**, 1009–1016 (2013).

14. Ferretti, A. C. & Joyce, G. F. Kinetic Properties of an RNA Enzyme That Undergoes Self-
Sustained Exponential Amplification. *Biochemistry-us* **52**, 1227–1235 (2013).

15. Lam, B. J. & Joyce, G. F. Autocatalytic aptazymes enable ligand-dependent exponential
amplification of RNA. *Nat Biotechnol* **27**, 288–292 (2009).

16. Janas, T., Janas, T. & Yarus, M. A membrane transporter for tryptophan composed of
RNA. *Rna* **10**, 1541–1549 (2004).

17. Pino, S., Ciciriello, F., Costanzo, G. & Mauro, E. D. Nonenzymatic RNA Ligation in Water*.
J Biol Chem **283**, 36494–36503 (2008).

18. Guerrier-Takada, C., Gardiner, K., Marsh, T., Pace, N. & Altman, S. The RNA moiety of
ribonuclease P is the catalytic subunit of the enzyme. *Cell* **35**, 849–857 (1983).

19. Anella, F. & Danelon, C. Prebiotic Factors Influencing the Activity of a Ligase Ribozyme.
Life **7**, 17 (2017).

20. Dai, X., Zhang, S. & Zaleta-Rivera, K. RNA: interactions drive functionalities. *Mol Biol Rep*
47, 1413–1434 (2020).

21. Müller, U. F. & Bartel, D. P. Improved polymerase ribozyme efficiency on hydrophobic
assemblies. *Rna* **14**, 552–562 (2008).

22. Anella, F. & Danelon, C. Prebiotic Factors Influencing the Activity of a Ligase Ribozyme.
Life **7**, 17 (2017).

23. Anella, F. & Danelon, C. Reconciling Ligase Ribozyme Activity with Fatty Acid Vesicle
Stability. *Life* **4**, 929–943 (2014).

24. Chen, I. A., Salehi-Ashtiani, K. & Szostak, J. W. RNA Catalysis in Model Protocell Vesicles. *J*
Am Chem Soc **127**, 13213–13219 (2005).

25. Mansy, S. S. *et al.* Template-directed synthesis of a genetic polymer in a model protocell.
Nature **454**, 122–125 (2008).

26. Bianconi, G., Zhao, K., Chen, I. A. & Nowak, M. A. Selection for Replicases in Protocells.
Plos Comput Biol **9**, e1003051 (2013).

27. Budker, V. G., Kazatchkov, Yu. A. & Naumova, L. P. Polynucleotides adsorb on
mitochondrial and model lipid membranes in the presence of bivalent cations. *Febs Lett* **95**,
143–146 (1978).

28. McManus, J. J., Rädler, J. O. & Dawson, K. A. Does Calcium Turn a Zwitterionic Lipid
Cationic? *J Phys Chem B* **107**, 9869–9875 (2003).

29. Kharakoz, D. P., Khusainova, R. S., Gorelov, A. V. & Dawson, K. A. Stoichiometry of
dipalmitoylphosphatidylcholine-DNA interaction in the presence of Ca2+: a temperature-
scanning ultrasonic study. *Febs Lett* **446**, 27–29 (1999).

30. Gromelski, S. & Brezesinski, G. Adsorption of DNA to zwitterionic DMPE monolayers
mediated by magnesium ions. *Phys Chem Chem Phys* **6**, 5551–5556 (2004).

31. Uhríková, D. *et al.* The structure of DNA–DOPC aggregates formed in presence of calcium
and magnesium ions: A small-angle synchrotron X-ray diffraction study. *Biochimica Et
Biophysica Acta Bba - Biomembr* **1713**, 15–28 (2005).

32. Gromelski, S. & Brezesinski, G. DNA Condensation and Interaction with Zwitterionic
Phospholipids Mediated by Divalent Cations. *Langmuir* **22**, 6293–6301 (2006).

33. McLoughlin, D. *et al.* Surface Complexation of DNA with Insoluble Monolayers. Influence
of Divalent Counterions. *Langmuir* **21**, 1900–1907 (2005).

34. Michanek, A. *et al.* RNA and DNA Association to Zwitterionic and Charged Monolayers at
the Air–Liquid Interface. *Langmuir* **28**, 9621–9633 (2012).

35. Michanek, A., Kristen, N., Höök, F., Nylander, T. & Sparr, E. RNA and DNA interactions
with zwitterionic and charged lipid membranes — A DSC and QCM-D study. *Biochimica Et
Biophysica Acta Bba - Biomembr* **1798**, 829–838 (2010).

36. Marty, R., N'soukpoé-Kossi, C. N., Charbonneau, D. M., Kreplak, L. & Tajmir-Riahi, H.-A.
Structural characterization of cationic lipid–tRNA complexes. *Nucleic Acids Res* **37**, 5197–
5207 (2009).

37. Suga, K., Tanabe, T., Tomita, H., Shimanouchi, T. & Umakoshi, H. Conformational change
of single-stranded RNAs induced by liposome binding. *Nucleic Acids Res* **39**, 8891–8900
(2011).

38. Lu, D. & Rhodes, D. G. Binding of phosphorothioate oligonucleotides to zwitterionic
liposomes. *Biochimica Et Biophysica Acta Bba - Biomembr* **1563**, 45–52 (2002).

39. Khvorova, A., Kwak, Y.-G., Tamkun, M., Majerfeld, I. & Yarus, M. RNAs that bind and
change the permeability of phospholipid membranes. *Proc National Acad Sci* **96**, 10649–
10654 (1999).

40. Vlassov, A. & Yarus, M. Interaction of RNA with Phospholipid Membranes. *Mol Biol+* **36**,
389–393 (2002).

41. Vlassov, A., Khvorova, A. & Yarus, M. Binding and disruption of phospholipid bilayers by
supramolecular RNA complexes. *Proc National Acad Sci* **98**, 7706–7711 (2001).

42. Janas, T., Janas, T. & Yarus, M. Specific RNA binding to ordered phospholipid bilayers.
Nucleic Acids Res **34**, 2128–2136 (2006).

43. Pannwitt, S., Slama, K., Depoix, F., Helm, M. & Schneider, D. Against Expectations:
Unassisted RNA Adsorption onto Negatively Charged Lipid Bilayers. *Langmuir* **35**, 14704–
14711 (2019).

44. Quarless, S. A. & Cantor, C. R. Analysis of RNA structure by ultraviolet crosslinking and
denaturation gel electrophoresis. *Anal Biochem* **147**, 296–300 (1985).

45. Behlen, L. S., Sampson, J. R. & Uhlenbeck, O. C. An ultraviolet light-induced crosslink in
yeast tRNA phe. *Nucleic Acids Res* **20**, 4055–4059 (1992).

46. Nejedlý, K., Kittner, R., Pospíšilová, Š. & Kypr, J. Crosslinking of the complementary
strands of DNA by UV light: dependence on the oligonucleotide composition of the UV
irradiated DNA. *Biochimica Et Biophysica Acta Bba - Gene Struct Expr* **1517**, 365–375 (2001).

47. Lengyel, A., Uhríková, D., Klacsová, M. & Balgavý, P. DNA condensation and its thermal
stability influenced by phospholipid bilayer and divalent cations. *Colloids Surfaces B
Biointerfaces* **86**, 212–217 (2011).

48. White, S. H., Wimley, W. C., Ladokhin, A. S. & Hristova, K. [4] Protein folding in
membranes: Determining energetics of peptide-bilayer interactions. *Methods Enzymol* **295**,
62–87 (1998).

49. Lee, J. S., Evans, D. H. & Morgan, A. R. Polypurine DNAs and RNAs form secondary
structures which may be tetra-stranded. *Nucleic Acids Res* **8**, 4305–4320 (1980).

50. Baross, J. A. & Martin, W. F. The Ribofilm as a Concept for Life's Origins. *Cell* **162**, 13–15
(2015).

51. Johansson, J. *et al.* An RNA Thermosensor Controls Expression of Virulence Genes in
Listeria monocytogenes. *Cell* **110**, 551–561 (2002).

52. Winkler, W., Nahvi, A. & Breaker, R. R. Thiamine derivatives bind messenger RNAs
directly to regulate bacterial gene expression. *Nature* **419**, 952–956 (2002).

53. Morita, M. T. *et al.* Translational induction of heat shock transcription factor ς32:
evidence for a built-in RNA thermosensor. *Gene Dev* **13**, 655–665 (1999).

54. Schilling, O., Langbein, I., Müller, M., Schmalisch, M. H. & Stülke, J. A protein-dependent
riboswitch controlling ptsGHI operon expression in Bacillus subtilis: RNA structure rather
than sequence provides interaction specificity. *Nucleic Acids Res* **32**, 2853–2864 (2004).

55. Nahvi, A. *et al.* Genetic Control by a Metabolite Binding mRNA. *Chem Biol* **9**, 1043–1049
(2002).

457 56. Boo, S. H. & Kim, Y. K. The emerging role of RNA modifications in the regulation of mRNA
458 stability. *Exp Mol Medicine* **52**, 400–408 (2020).

459 57. Edwards, A. M., Addo, M. A. & Santos, P. C. D. Extracurricular Functions of tRNA
460 Modifications in Microorganisms. *Genes-basel* **11**, 907 (2020).

461 58. Zhou, W., Karcher, D. & Bock, R. Importance of adenosine-to-inosine editing adjacent to
462 the anticodon in an Arabidopsis alanine tRNA under environmental stress. *Nucleic Acids Res*
463 **41**, 3362–3372 (2013).

464 59. Patel, B. H., Percivalle, C., Ritson, D. J., Duffy, C. D. & Sutherland, J. D. Common origins of
465 RNA, protein and lipid precursors in a cyanosulfidic protometabolism. *Nat Chem* **7**, 301–307
466 (2015).

467 60. Rushdi, A. I. & Simoneit, B. R. T. Lipid Formation by Aqueous Fischer-Tropsch-Type
468 Synthesis over a Temperature Range of 100 to 400 °C. *Origins Life Evol B* **31**, 103–118 (2001).

469 61. Brady, R. C. & Pettit, R. Mechanism of the Fischer-Tropsch reaction. The chain
470 propagation step. *J Am Chem Soc* **103**, 1287–1289 (1981).

471 62. Robertson, M. P. & Ellington, A. D. Design and optimization of effector-activated
472 ribozyme ligases. *Nucleic Acids Res* **28**, 1751–1759 (2000).

473 63. Debiais, M., Lelievre, A., Smietana, M. & Müller, S. Splitting aptamers and nucleic acid
474 enzymes for the development of advanced biosensors. *Nucleic Acids Res* **48**, 3400–3422
475 (2020).

476 64. Stifel, J., Spöring, M. & Hartig, J. S. Expanding the toolbox of synthetic riboswitches with
477 guanine-dependent aptazymes. *Synthetic Biology* **4**, ysy022- (2019).

478 65. Vlassov, A., Khvorova, A. & Yarus, M. Binding and disruption of phospholipid bilayers by
479 supramolecular RNA complexes. *Proc National Acad Sci* **98**, 7706–7711 (2001).

480 66. Janas, T. & Yarus, M. Visualization of membrane RNAs. *Rna* **9**, 1353–1361 (2003).

Materials and methods

Materials

Dipalmitoylphosphatidylcholine (DPPC) was purchased from Avanti Polar Lipids and was used without further purification. HEPES, $MgCl_2$ and $CaCl_2$ were purchased from CarlRoth and were at least >99% pure. All solutions were prepared in MiliQ water, Millipore.

R3C ligase ribozyme was expressed using pRZ plasmid and T7 RNA polymerase and purified using denaturing PAGE and electroelution. 5' 6-FAM labelled R3C substrates as well as short oligomers used in binding assays were obtained from Integrated DNA Technologies and used without further purification.

Sequence name	Sequence (5' – 3')
R3C	GAGACCGUAAUGAGUAGUACUUAUUAUGCUCGAUUGUUCGUAAG AACAGUUUGAAUGGGUUGAAUAUAGUGAGGAUCCGCUAGC
Short substrate	6_FAM_CGACUCACUAUA
4xAG substrate	6_FAM_A**G**A**G**A**G**A**G**CGACUCACUAUA
9xAG substrate	6_FAM_A**G**A**G**A**G**A**G**A**G**A**G**A**G**A**G**ACGACUCACUAUA
2G substrate	6_FAM_HHHHHHHH**G**HHH**G**HHHHHHCGACUCACUAUA
0G	HHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHH
2G	HHHHHHHHHHHHHHH**G**HHH**G**HHHHHHHHHHHHHHHH
4G	HHHHHHHHHHH**G**HHH**G**HHH**G**HHH**G**HHHHHHHHHHH
5G	HHHHHHHHH**G**HHH**G**HHH**G**HHH**G**HHH**G**HHHHHHHHHH
7G	HHHHHH**G**HHH**G**HHH**G**HHH**G**HHH**G**HHH**G**HHHHHH
9G	HH**G**HHH**G**HHH**G**HHH**G**HHH**G**HHH**G**HHH**G**HHH**G**HH
40xN	NNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNN
40xN - G	HHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHH
40xN - A	BBBBBBBBBBBBBBBBBBBBBBBBBBBBBBBBBBBBBBBB
40xN - C	DDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDD
40xN - U	VVVVVVVVVVVVVVVVVVVVVVVVVVVVVVVVVVVVVVVV
19xG	**GGGGGGGGGGGGGGGGGGG**
19xA	AAAAAAAAAAAAAAAAAAA
19xC	CCCCCCCCCCCCCCCCCCC
19xU	UUUUUUUUUUUUUUUUUUU
19xN	NNNNNNNNNNNNNNNNNNN
AG repeats	6_FAM_AGAGAGAGAGAGAGAGAGA
ssRNA (ACUG)	ACUGACUGACUGACUGACUGACUGACUGACUG
ssRNA (CAGU)	CAGUCAGUCAGUCAGUCAGUCAGUCAGUCAGU

B – C, G or U

D – A, G or U

H – A, C or U

V – A, C or G

Methods

All of the RNA incubations were prepared in Sarstedt DNA low bind tubes at the constant temperature of 24 °C in buffer comprised of 10 mM HEPES pH 7, 5 mM $CaCl_2$, and 5 mM $MgCl_2$. RNA concentration was measured by measuring absorbance at 260nm (A_{260}, SPARK 20M, TECAN). Before every incubation RNA was preheated in SafeLock Eppendorf tubes (90 °C, 5 minutes) to ensure unfolded structures. Samples after UV crosslinking and R3C activity assays were ethanol precipitated.

Lipid preparation

DPPC chloroform stock was pipetted into a glass vial and briefly evaporated under a steady flow of nitrogen gas. To remove organic solvent residues, a lipid film was dried under vacuum overnight. Lipid film hydration (in reaction buffer) method was used to obtain multi layer DPPC vesicles (MLV DPPC). Buffer-lipid mixtures were shaken at 60 °C, well above the melting temperature of DPPC (T_m = 42 °C), for one hour. A cloudy liposome suspension was freeze and thawed 10x and extruded 17x through 100 nm polycarbonate filters (Millipore) to reduce multi-lamelarity and achieve a consistent size distribution of vesicles. Lipid vesicle stocks were kept in 4 °C. Stock lipid concentrations were confirmed using a phosphate assay.

Dynamic light scattering

The size distribution of DPPC vesicles was estimated using a ZetaSizer Nano in 173° backscatter with polydisperse solution calculation multi-mode. Final concentration of lipids was 10 uM and RNA amount was kept at a ratio of 10 lipids / nucleotide. Results were plotted as the size distribution in number of detected species function (NumberPSD).

UV-crosslinking

RNA was incubated for 10 minutes in the reaction buffer with or without lipids followed by 10 minutes incubation under UV-B light (300 mW LEUVA77N50KKU00 LED 305 nm) from a distance of 1.5 – 2 cm. Samples were analysed on denaturing PAGE with SybrGold poststaining.

R3C activity assays

5 pmol of R3C ligase was mixed with 0.5 pmol of the substrate to ensure saturation of the system (pseudo 1^{st} order reaction) and decrease any batch-to-batch differences coming from R3C purification. Incubation was conducted at various concentrations of DPPC vesicles and samples were ethanol precipitated after 30, 60, 90 and 120 minutes. Reactions were quantified on denaturing PAGE using FIJI software. Product and substrate intensities (I_P and I_S, respectively) were quantified and the concentration of product at time t was calculated:

$$[P]_t = [S]_0 * \left(\frac{I_P}{I_P + I_S} \right)$$

Concentration of product was plotted as a function of time and then fitted with a linear function (fixed [0;0] point) to determine reaction rates.

RNA-lipid binding assays

Fig. 3a,b binding assays were conducted using liposome-coated magnetic beads. Fig. 2b,c, Fig. 4 binding assays were conducted using an ultracentrifugation method.

Magnetic beads binding assay

Magnetic beads (DynaBeads streptavidin T1) were coupled with DPPC liposomes doped with 0.5 mol%

biotynoyl cap PE (Avanti Polar Lipids). Lipid concentration on non-diluted beads was estimated using a

phosphate assay and a theoretical calculation and came out to be 800 uM. 25 pmol of 19nt or 5 pmol

of 40nt RNA species were incubated with 2x diluted magnetic beads for 30'. After incubation the

supernatant was separated from beads using magnetic field. The amount of RNA was estimated either

using absorbance or Qubit miRNA quantification kit (ThermoScientific).

Ultracentrifugation binding assay

RNA was incubated with various concentrations of DPPC vesicles for 30'. After incubation

Ultracentrifugation was conducted to separate vesicles from solution (125 000 x G, 40', 24 °C). 9 uL of

supernatant (out of 30 uL reaction mix) was pulled and incubated with 1.5 uL of 0.5 M EDTA (90 °C, 5').

The amount of RNA was measured by Qubit miRNA quantification kit using a plate reader (SPARK 20M,

TECAN).

Measured fluorescence intensity values were plotted as a function of lipid/oligomer or lipid/nucleotide

ratios (we assumed that RNA interacts only with the outer membrane leaflet) and fitted using Hill's fit

(OriginLab software):

575
$$f(x) = Start + (End - Start)\frac{x^n}{x^n + k^n}$$

Where Start and End are function plateau values, k is inflection point of curve and n is cooperativity

index. Inflection point of the curve (k) is a parameter which we used for further data analysis.

RNA-lipid binding partition coefficient determination

RNA was incubated with high excess of DPPC vesicles (5 mM, 2.5 mM outer leaflet) for 30 minutes.

After incubation ultracentrifugation was conducted to separate vesicles from solution (125 000 x G,

40', 24 °C). 10 uL out of 30 uL was taken as a supernatant sample, and the rest of the supernatant was

discarded to leave only the pellet. RNA from supernatant and pellet samples was ethanol precipitated

and analyzed using denaturing PAGE. The partition coefficient K was calculated according to following

formula:

588
$$K = \frac{\frac{[R_L]}{[R_L] + [L]}}{\frac{[R_w]}{[R_w] + [W]}}$$

590 **Where:**

591 **$[R_L]$ – RNA bound to the membranes**

592 **$[R_w]$ – non-bound RNA**

593 **$[L]$ – outer lipid concentration**

594 **$[W]$ – water concentration**

Fig. 1   RNA-gel membrane binding has different phenotypes. Introduction of a randomized pool of RNA oligos to gel lipid membranes promotes vesicle aggregation a divalent cation-dependent process **(a)**. R3C ligase (84 nt) shows a small increase in UV-crosslinking at low lipid concentration, whereas crosslinking efficiency decreases at higher lipid concentration. Binding and UV crosslinking are inversely correlated and show similar inflection points with respect to lipid concentration **(b)**. Random RNA oligos (75-85 nt) show increasing crosslinking efficiency in the presence of lipids with a slight decrease at the highest lipid concentration; crosslinking and lipid binding are closely correlated **(c)**. The R3C ligase forms primarily one structure, whereas randomized RNA forms intermolecular complexes **(d)**.

a) R3C ligase activity determination

Slope = reaction rate

b) R3C ligase reaction rate in function of lipid content

c) Lipid binding and characteristics of R3C reactants

Oligo	log(K)	Nucleotide %				Length
		G	A	C	U	
R3C	6.74 ± 0.13	27	28	13	31	84 nt
Substrate	3.39 ± 0.15	8	33	33	25	12 nt

Fig. 2   Lipid membranes influence R3C ligase activity through lipid-ligase interactions. R3C ligase reaction rates were quantified by measuring the amount of product within 2 hours incubation time **(a)**. R3C reaction rate is dependent on the presence of gel membranes - increased activity is observed for lower lipid concentration followed by drop in activity at higher lipid concentration **(b)**. The 12 nt R3C substrate shows several orders of magnitude lower membrane-buffer partition coefficient compared with the R3C ligase, which could derived from different compositional or structural characteristics of both RNA species **(c)**.

Fig. 3   RNA-gel membrane binding is RNA sequence-dependent. Short 19 nucleotide oligomers show sequence-dependent binding to the gel membranes, with highest binding observed for pure G oligomer followed by oligos with randomized sequence or AG repeats. Oligomers containing only A, C or U bases show negligible binding to the lipid membranes **(a)**. Increase in oligonucleotide length leads to a further increase in binding efficiency (compare **(a)** 19xN and **(b)** 40xN). Depletion of one type of nucleotide within oligomers caused a significant decrease in binding efficiency. Lack of guanine and cytidine decreased binding by 43 and 31 percentage points, respectively, whereas lack of adenine or uracil showed a less pronounced effect **(b)**. Oligonucleotides with increasing G content showed higher lipid – buffer partition coefficients compared with G-deficient oligo with the highest value observed for the 4xG oligomer (0.9 log unit) **(c)**.

Fig. 4   RNA-gel membrane binding depends on RNA structure. The membrane binding of RNA's which can't create any inter- and intramolecular structures (36 nucleotide ACUG or CAGU repeats) was examined together with 100% complementary double stranded structures (mixed ACUG and CAGU oligos). Single stranded RNAs do not cause lipid vesicle aggregation despite moderate membrane binding, whereas double stranded RNA species show both high binding and lipid aggregation **(a)**. RNA:lipid binding ratios are different for single stranded RNAs compared to double stranded species **(b)**. Oligomers with different nucleotide composition show different lipid/nucleotide binding stoichiometry. Analysis of oligomers by native PAGE gel indicates that the presence of guanosine promotes inter-molecular interactions, suggesting, that base-pairing enhances the average strength of nucleotide-lipid interactions **(c)**.

Fig. 5   Modification of R3C substrate sequences enables tunability of ligation rate. Modified R3C reaction substrates were synthesized in order to increase binding to the membrane. The non-catalytic 5′ end of the molecule was amended with tails with varying G content **(a)**. Reactions for all AG-modified substrates, but not 2xG, showed lower reaction rate constants compared with the unmodified substrate **(a, table)**. The mechanism for inhibition of activity was evaluated by analysis of R3C reactants using native PAGE **(b)**. The unmodified short substrate was entirely bound to the ligase, whereas AG-modified substrates show mobility abnormalities, which suggest substrate folding **(b, magenta points)**. Modification of the substrate sequence through addition of a 5′ overhang increased binding to the membrane **(c)**. Addition of lipid membranes to reactions with the G-rich substrates increased reaction rates to the level of the unmodified substrate, except for the 2xG substrate which showed no change **(d)**. Decrease of the R3C reaction rates in the presence of G-rich substrates might be based on the inhibition of the catalytic part of the substrate by the 5′ G-rich overhang **(e)**.