Lipid membranes modulate the activity of RNA through sequence-specific interactions

Authors: Tomasz Czerniak[a] and James P Saenz[a]#

[a] Technische Universität Dresden, B CUBE Center for Molecular Bioengineering, 01307 Dresden, Germany

#corresponding author and lead contact: james.saenz@tu-dresden.de

Abstract

RNA is a ubiquitous biomolecule that can serve as both catalyst and information carrier. Understanding how RNA activity is controlled and how it in turn regulates bioactivity is crucial for elucidating its physiological roles and potential applications in synthetic biology. Here we show that lipid membranes can act as RNA organization platforms, introducing a novel mechanism for ribo-regulation. The activity of R3C ribozyme can be modified by the presence of lipid membranes, with direct RNA-lipid interactions dependent on RNA sequence, structure and length. In particular, the presence of guanine in short RNAs is crucial for RNA-lipid interactions, while double-stranded RNAs further increase lipid-binding affinity. Lastly, by artificially modifying the R3C-substrate sequence to enhance membrane binding we unexpectedly generated a lipid-sensitive riboswitch. These findings introduce RNA-lipid interactions as a tool for developing riboswitches and novel RNA-based lipid biosensors, and bear significant implications for RNA World scenarios for the origin of life.

Introduction

RNA performs diverse functions ranging from information storage to regulation of other biomolecules and direct catalysis of biochemical reactions. The functional versatility of RNA has implications for understanding plausible scenarios for the origin of life[1–3], and for developing tools in synthetic biology[4–6]. Research aimed at understanding how an RNA World could have emerged has motivated development of ribozymes with functions including RNA ligation[7–10], replication[11,12] and other activities[13–16]. The experimental development of functional RNAs raises the possibility of RNA-based synthetic systems. For both synthetic life and understanding the origin of self-replicating organisms, RNA has intrinsic appeal: it can serve functions of both DNA (information storage) and proteins (enzymes), obviating the need for translation machineries and protein chaperones. Furthermore, RNA does not undergo irreversible denaturation like proteins, lending robustness against a broad range of physical and chemical conditions. In order to design a system based on RNA, however, it is essential to be able to coordinate RNA activity in space and time.

Key to harnessing the functional versatility of RNA is understanding how to spatially and temporally modulate its properties and to selectively modulate the activity of different RNAs within one system. The physicochemical environment surrounding an RNA molecule is a central determinant of its structure, stability and activity. Spontaneous RNA hydrolysis and ligation, as well as catalytic RNA activity are sensitive to pH[17], ionic strength[18,19] and RNA concentration changes[11], among other parameters. Similarly, RNA activity can be modulated by interactions with molecules such as ions, proteins, and other nucleic acids[20]. Thus, one approach to regulating RNA activity could be via tunable

45   interactions with binding partners that affect RNA structure, concentration or chemical
46   microenvironment.
47
One mechanism for modulating RNA activity could be through direct RNA-lipid interactions[19,21,22].
Because of their amphiphilic nature, lipids spontaneously self-assemble into membranous structures
that can encapsulate RNA into protected and selective microcompartments[23–25]. Alternatively, direct
RNA-lipid interactions could localize RNAs to membrane surfaces, increasing its local concentration
and reducing dimensionality for intermolecular interactions[21]. Lastly, localization to a lipid surface
brings RNA into a physicochemically unique microenvironment with sharp gradients of hydrophobicity,
electrical permittivity, and water activity. Through these effects, RNA-lipid interactions could provide
a powerful mechanism for modulating RNA activity.
56
The first functional RNA-lipid interaction was described more than 40 years ago[26], with subsequent
research revealing various factors that facilitate nucleic acid-lipid binding[27–37]. More recently, specific
RNA sequences have been generated through SELEX with affinity for fluid membranes comprised of
phospholipids and cholesterol[38–40]. Interestingly, mixtures of RNAs have also been shown to bind to
membranes that are in a solid crystalline (gel) phase[41,42]. These studies revealed that, while most
randomized mixtures of RNA sequences can bind to gel membranes, there is a relatively small chemical
space of oligomers that have affinity for fluid membranes. Thus, conceptually, gel membranes could
provide a platform for modulating the activity of a diverse range of RNAs. However, the effects of gel
membranes on RNA activity and the sequence selectivity of such interactions are relatively unexplored.
66
This study reports the effect of lipid membranes on RNA catalytic activity. We show that RNA-lipid
binding depends on the primary sequence, secondary structure, and length of RNA. Using the trans-
acting R3C ligase ribozyme, we observed that R3C-lipid binding changes ribozyme activity in a
concentration-dependent manner. Lipid binding assays show that the interaction of short RNA
sequences with gel membranes is sequence specific and depends on guanine content and the presence
of double-stranded structures. Lastly, modification of R3C's substrate sequence increased the
tunability of R3C-based reactions through a lipid-dependent riboswitch. Our findings demonstrate that
membranes can serve as platforms for RNA-activity modulation, which could contribute to the
development of RNA-based biosensors and riboswitches. This approach introduces new tools for
molecular and synthetic biology and raises the prospect of previously unrecognized roles for RNA-lipid
interactions in the origin and evolution of life.
78

Results

The discovery that RNA can catalyse reactions in addition to encoding information[18,43], opened new
directions for engineering life and the possibility of protocells emerging from an RNA world[2]. But, a
key missing ingredient for RNA-based systems is a mechanism to organize RNAs and regulate their
activity. We hypothesized that RNA-membrane interactions could influence ribozyme activity by
changing local RNA concentrations at the membrane surface or influencing RNA conformations.
Indeed, RNA oligonucleotides can bind to lipid membranes in the gel phase in the presence of divalent
cations, leading to aggregation into large visible RNA-lipid assemblies (**Fig. 1a**) [27,35,41,42]. Depletion of
divalent cations or increasing temperature above the lipid melting temperature (i.e. producing fluid
instead of gel membranes) reduced RNA-membrane binding and aggregation (**Fig. 1a**). This reversible
aggregation could provide a tunable mechanism to concentrate and regulate RNAs in simple bottom
up synthetic systems, or in a prebiotic environment.

RNA-lipid binding changes the probability of RNA-RNA interactions

To determine whether local RNA concentration is influenced by interaction with lipid membranes, we relied on UV-mediated crosslinking. All membranes in this study were composed of dipalmitoylphosphatidylcholine (DPPC), which is in a solid crystalline (gel) phase at the experimental temperature of 24°C. Nucleic acid bases absorb UV light, producing chemical changes that yield base-base covalent bonds in a distance-dependent manner, yielding insights into the structure and interactions of nucleic acids[44–46]. We first observed the effect of gel membranes on the crosslinking of a single defined RNA sequence, the R3C ligase, which predominately forms one structure (R3C ligase, **Fig. 1d**). When subjected to UV, the R3C ligase shows slightly increased crosslinking in the presence of low lipid concentrations. A further increase in lipid concentration led to decreased crosslinking efficiency, possibly through dilution of the RNA species on the surface of the lipid membranes. This effect was inversely correlated with RNA-lipid binding efficiency (**Fig. 1b**). In contrast, a mixture of RNA oligomers with randomized sequences which is able to create a more diverse range of intermolecular structures (**Fig. 1d**) showed continuously increasing and overall higher UV-crosslinking efficiency in the presence of lipids (**Fig. 1c**). The contrasting effect of crosslinking for R3C and randomized oligomers might be explained by the influence of lipid membranes on RNA interactions: membrane binding could potentially stabilize base-pairing[47] and thereby enhance crosslinking efficiency, whereas the decrease of crosslinking might be caused by a dilution effect at higher membrane surface areas or by lipid-based steric hindrance. These observations demonstrate that RNA-membrane interactions can influence RNA-RNA interactions in a manner that is dependent on lipid concentration and RNA sequence diversity. This further suggests that membrane binding could have an effect on trans-acting ribozyme activity derived from base pairing.

RNA-lipid interactions influence RNA catalytic activity

To investigate the functional consequence of RNA-lipid binding, we tested the effect of lipids on the activity of the trans-acting R3C ligase ribozyme. R3C ligase is a relatively short ribozyme that catalyzes ligation of substrate strands to the ribozyme[10](**Fig. 2a**). Since R3C is also part of an RNA self-replication system [11,12], the effects of lipids on R3C are also interesting with regard to the emergence, evolution and artificial synthesis of autonomous self-replicating systems. The R3C reaction rate in the absence of membranes was 11 pM/min, with a reaction constant rate of k = 6.4×10^{-4} min^{-1}. Addition of lipid vesicles led to an increased reaction rate (14 pM/min, +29%), which could plausibly be due to increased ligase-substrate interaction on the membrane either through increased concentrations at the membrane surface or through stabilization of base pairing[47]. At the highest lipid concentration, the reaction rate dropped to 8 pM/min (-27%) (**Fig. 2b**). A decrease in reaction rate was correlated with further R3C ligase binding to the lipid membranes (>100 lipid/R3C, **Fig. 1b**) and we speculate that this may be caused by steric hindrance from the lipid vesicles. These results show that gel membranes can influence the reaction rate of a trans-acting ribozyme in a lipid concentration-dependent manner.

R3C ligase and its substrate bind differently to the membrane

To further understand how R3C and substrate membrane binding might contribute to changes in ligation rates, we analyzed lipid-binding affinities of the ligase and its substrate. To quantify binding to the membranes, we calculated lipid–buffer partition coefficients (K). The partition coefficient of the short 12 nt substrate was more than 3 orders of magnitude lower than that of the 84 nt R3C ligase (**Fig. 2c**). Thus, in this system, substrate-membrane binding is essentially negligible compared with R3C-membrane binding. Combined with our observation that RNA-RNA interactions are stabilized by

137 membrane binding (Fig. 1c), this indicates that the increased activity observed in **Figure 2b** was due to
138 an effect of the membrane on the R3C ligase-substrate complex stability, rather than through
139 substrate-lipid interactions on the membrane surface. Thus, membranes can enhance ribozyme
140 activity by promoting the stability of reaction intermediates.
141
RNA sequence influences binding to gel membranes
Understanding the features that influence RNA-lipid interactions could allow us to engineer RNAs with
higher membrane affinity, and thereby potentially enhance activity by concentrating RNAs at the
membrane surface. Interactions of RNA with gel-phase membranes have been previously shown to
exhibit low sequence-specificity, based on binding of oligos with randomized sequences[41,42]. However,
the large difference in binding for R3C and its substrate suggests that RNA-gel membrane interactions
might in fact be dependent on RNA composition or structure.
149
To investigate which features of RNA influence its interaction with lipid membranes we first evaluated
how base content influences membrane binding of short RNA species. 19nt oligomers with only one
type of nucleotide were tested together with a random sequence as a control. To quantify relative
changes of RNA-membrane binding, we calculated the RNA binding efficiency as a percentage of total
RNA bound to membranes at a fixed oligo-to-lipid ratio. Remarkably, whereas the 19xG oligomer
bound very efficiently to the gel-phase membrane (84% binding efficiency), oligomers of A, C or U
showed practically no binding. A mixture of 19-mer RNA with randomized sequences showed
moderate binding (33% binding efficiency), as did one with AG-repeats (**Fig. 3a**), most probably due to
the introduction of guanine. Guanine could conceivably enhance binding either through direct
interactions with the membrane, or alternatively by promoting G≡C base pairing or presence of G-
quadruplexes, thereby influencing structure. The enhanced binding of an oligomer with only AG-
repeats suggests that membrane binding can be influenced by direct guanine-membrane
interactions[48].
163
We next examined the effect of deleting a specific base from mixtures of 40-mer oligos with
randomized sequences. Depletion of A, C or U led to a slight decrease in membrane binding, whereas
removal of G showed a significant binding decrease. The decrease in binding affinity in all of the base-
depleted oligos (**Fig. 3b**) indicates that base-pairing may influence membrane binding. However, the
larger effect of G depletion indicates that either depletion of G causes highest bias on RNA base-pairing
or base pairing is not the only factor responsible for membrane binding, consistent with our previous
observation that G-content alone can influence binding (Fig. 3a). Finally, by comparing 19-mer and 40-
mer randomized oligos, we observed that increasing RNA length also led to enhanced binding, possibly
due to increased length and G content (**Fig. 3 a,b**).
173
To further determine how guanine content affects membrane binding, lipid-buffer partition
coefficients were determined for short oligomers with varying G-content. RNA membrane-buffer
partition coefficients indicate how well a particular molecule binds to the membrane by comparing the
relative amount of RNA in buffer and on the lipid membrane. The RNA oligos measured in this study
had partition coefficients around 10^5, which is in the range of partition coefficients determined for
hydrophobic peptides[49]. Remarkably, addition of 2 G's per oligo led to a 2.7-fold increase of the
partition coefficient relative to guanine-depleted RNA. Oligos with 4 and more guanine residues
showed a further increase of partition coefficient values with highest value for 9 G's. This result shows

that the presence of guanine affects RNA-lipid binding and even low guanine content significantly
increases membrane binding efficiency (**Fig. 3c**).
184

RNA-membrane binding is dependent on RNA structure
Our observations of the effect of base depletion on oligomer binding suggested that base-pairing may
be a key factor in membrane binding. To investigate the effect of base-pairing-derived secondary
structures on membrane interactions, we estimated binding efficiency for ssRNA and dsRNA
(composed of 100% complementary sequences) species. We observed that not only was the
membrane binding efficiency higher for dsRNA compared with ssRNA species, but also that only dsRNA
promoted vesicle aggregation (**Fig. 4a**). It is interesting to note that the difference in binding efficiency
of the two ssRNAs, which have identical base composition but different sequence, is consistent with
the importance of sequence as a key determinant of membrane binding. We further observed that
dsRNA interacts with fewer lipids than ssRNA, implying different models of binding (**Fig. 4b**). Thus,
membrane binding efficiency can be dependent on the propensity of an RNA molecule to form intra-
or inter-molecular structures through base pairing.

To gain further insight into the effect of base pairing-dependent intra- and inter-molecular
interactions, we estimated the lipid-to-nucleotide binding ratios of ssRNA and dsRNA (**Fig. 4c**). By
calculating the molecular ratio of lipid/nucleotide binding, we introduce a normalization factor that
allows us to directly compare membrane binding for oligos of varying length. Furthermore, varying
lipid to nucleotide stoichiometry reflects changes in the average lipid/nucleotide interaction strength.
With this estimate, we compared lipid/nucleotide binding for ssRNA and dsRNA alongside oligos with
varying guanine content from our previous analysis (**Fig. 3c**). The differences between ssRNA and
dsRNA, all containing 9 guanines per oligo, confirmed our previous observation that sequence variation
(with constant base content) and base pairing can both influence binding. However, the reason for a
change in lipid/nucleotide ratio for oligos with varying guanine content was not readily apparent.
Analysis of oligos with varying guanine content by native non-denaturing gel revealed that the
presence of guanine promotes a larger spread in migration (**Fig. 4c, inset**), indicating that guanines
promote intermolecular interactions. These observations show that guanine can enhance binding by
promoting intermolecular interactions, and further confirm the importance of base pairing in RNA-
lipid interactions.

Modifying R3C substrate sequence enhances membrane binding and modifies reaction rates
In previous experiments, we observed that the binding affinity of the short substrate was >3 orders of
magnitude lower than for the R3C ligase and that reaction rates correlated strongly with R3C-
membrane binding (**Fig. 2c, d**). We therefore reasoned that by modifying the R3C ligase substrate to
enhance membrane binding, we might be able to further modulate ligase activity through increased
RNA density at the membrane surface. Having determined that guanine content is a key factor
influencing RNA-lipid binding (**Fig. 3**), we modified the R3C substrate sequence to increase lipid
membrane binding efficiency by addition of a 5'-overhang with varying amounts of AG repeats (**Fig.
5a**).

We first assessed the effect of modifying the R3C ligase substrate on ligation rate in the absence of
membranes (**Fig. 5a**). We observed that increasing the guanine content decreased ligase activity,
suggesting that 5' modification of the substrate was inhibiting the reaction either through steric
hindrance or promotion of inhibitory inter- or intra-molecular interactions. To determine whether AG-

rich substrates form intra-molecular structures[50] or exhibit reduced binding with the R3C ligase, we
performed electrophoresis in native conditions with and without ligase (**Fig. 5b**). Firstly, the non-
modified short substrate migrated as one band and was fully bound to the R3C ligase. Secondly, a
4x(AG) 20 nt substrate variant migrated as two bands, which suggests that part of the substrate
molecules have some folding that might bias binding to the R3C ligase. In the presence of ligase, 30%
of the substrate was non-bound, which correlated with a 22% reduction in reaction rate. For the
longest substrate variant (31 nt), electrophoretic mobility suggested higher folding, since only one
band was present and migrated half-way between the non-folded 12 nt and 20 nt substrates. 70% of
the 9x(AG) substrate was not bound to the ligase, which correlates with a reduced R3C reaction rate.
Lastly, the AG-rich oligomer did not co-migrate with the R3C ligase, indicating that the inhibition effect
is not based on non-specific substrate-ligase interactions (not shown). Therefore, increasing guanine
content reduces R3C ligase-substrate binding, likely accounting for decreased activity.
240
We next investigated whether varying guanine content through 5' modification of the substrate
influences membrane binding and, consequently, ligation activity in the presence of membranes. As
expected, the longer guanine-containing R3C substrates showed significantly higher membrane
binding compared with the shorter variant (**Fig. 5c**). To determine whether binding of the substrates
to the membrane enhances ligase activity, we determined R3C reaction rates in the presence of the
gel lipid membranes (**Fig. 5d**). For modified substrates, the reaction rates increased significantly to the
level of the unmodified substrate. Interestingly, none of the modified substrates exhibited higher
activity than the unmodified substrate, indicating that the rescue of activity of the other guanine
variants is not due to increased substrate-ligase interactions through enhanced binding to the
membrane, but rather due to interaction of the membrane with 5' overhangs. Thus, substrate-
membrane interaction increases activity by rescuing the inhibitory effect of the guanine-rich 5'
overhangs (**Fig. 5e**).
253
254   In summary, by modifying the R3C ribozyme substrate with an additional short sequence containing
255   guanine residues, we increased membrane affinity. The addition of guanines, however, also introduced
256   an inhibitory effect on ligation activity that was reversed through the introduction of gel-membranes
257   to the reaction. Thus, we unexpectedly produced a lipid-sensitive riboswitch-like system that could be
258   tuned in a lipid concentration-dependent manner.
259
260   **Discussion**
261
We have demonstrated that RNA-gel membrane interactions are dependent on the primary sequence
and secondary structure of ribonucleic acids. Varying the guanosine content of RNAs changed
membrane-buffer partitioning by several orders of magnitude. Additionally, double-stranded RNAs
exhibited higher membrane partitioning than single-stranded RNAs, indicating that secondary
structure is a critical factor in RNA-membrane interactions. Although gel-membranes are not widely
observed in living cells, RNA-gel membrane interactions could be employed for ribo-regulation in
synthetic biological systems. Gel membranes also plausibly accumulated in ancient prebiotic scenarios,
potentially serving as an organizational scaffold, as has been proposed for "ribofilms" on mineral
surfaces[51]. Furthermore, understanding the factors involved in RNA-gel membrane interactions should
provide insights that can be used to design RNA sequences with affinity for physiologically relevant
fluid membranes. Indeed, our finding that guanosine is a key factor in promoting RNA-gel membrane

partitioning is consistent with previous speculation that guanosine residues might play a role in the
binding of RNA aptamers to fluid membranes[38]. Taken together, the sequence and structural
dependence of RNA-gel membrane interactions shown here provides the necessary framework to
design synthetic RNAs with varying membrane affinities.
277
Our observation of riboswitch-like behavior suggests a novel mechanism for regulation of RNA through
membrane interactions. Riboswitches are RNAs with regulatory effects that are derived from
sensitivity to physicochemical conditions or through binding with an interaction partner. Riboswitch
regulation through temperature changes, interaction with nucleic acids and small molecules have been
described previously[52–56]. However, lipid sensitive riboswitches have not been documented. It is now
plausible that spontaneously self-assembling lipid membranes in ancient environments could have
served as an organizational scaffold regulating the activity of RNAs in a sequence- and structure-
dependent manner. Furthermore, the importance of guanosine in membrane binding suggests a
possible mechanism for its selection as one of the canonical nucleic acids. In extant organisms,
synthetic or naturally occurring RNAs exhibiting sensitivity to lipids could potentially act as biosensors
or riboswitches. By demonstrating that RNA-gel membrane interactions can induce riboswitch-like
behaviors, we raise the possibility that such functionalities could be an ancient mechanism of ribo-
regulation and may be present among the myriad uncharacterized small RNAs in modern cells.
291
RNA-membrane interactions provide a means to select RNAs by sequence and structure. For example,
diverse RNA species can be segregated or co-localized based on their differential membrane affinities.
Furthermore, selective RNA-membrane localization can be controlled by shifting the temperature
above and below the membrane gel-liquid transition temperature, thereby turning on and off RNA-
membrane binding. Such lipid-dependent sorting of RNAs could provide a mechanism to selectively
control the concentration of reactants in primordial scenarios and in synthetic RNA-based systems.
Selective RNA-membrane interactions would also influence the sequence space explored by evolving
ribozymes. Ribozymes generated through natural or artificial selection in the presence of lipid
membranes may, therefore, evolve enhanced or novel functions. The sequence and structural
selectivity of RNAs for membranes can also be implemented in the design of novel RNA membrane
biosensors. Looking forward, RNA-membrane interactions can provide tools for selective ribo-
regulation in synthetic systems, development of membrane biosensors, and might facilitate the
emergence of novel functions through artificial and natural evolution of ribozymes in the presence of
membranes.
306
In conclusion, our findings reveal that lipids, which are present in every modern cell and were plausibly
part of a prebiotic world[57–59], can interact with RNA and change its activity. These findings have
significant applications in fields such as synthetic biology, where merging the selective affinities of
aptamers with ribozyme activity (aptazymes) is currently a developing field[60–62]. Our results provide a
foundation for exploring the basis of membrane-RNA interactions with physiologically relevant fluid
membranes exhibited by synthetic aptamers[16,40,63] thereby providing new regulatory mechanisms in
synthetic biology through the specific and anchor-free localization of RNA molecules within lipid
compartments. Furthermore, insights from the present study can already be implemented in
bioengineering applications such as in improving mRNA drug delivery mechanisms and introducing
lipid sensitive ribo-regulation to synthetic ribozyme networks. Finally, the interaction between RNA
and lipid membranes could have provided an evolutionary pressure in ancient times through selection
for base composition or secondary structural features.

319
320
Acknowledgements:
322 We would like to thank Mario Mörl, Ilya Levental, Robert Ernst, Andre Nadler, Dora Tang, Grzegorz
323 Chwastek, Michał Grzybek and Mike Thompson for helpful discussions and feedback. This work was
324 supported by the B CUBE of the TU Dresden, a Simons Foundation Fellowship (to J.S.), a German
325 Federal Ministry of Education and Research BMBF grant (to J.S., project 03Z22EN12), and a VW
326 Foundation ''Life'' grant (to J.S., project 93090).
327
328 The authors declare no conflict of interests.
329

References

1. Pace, N. R. & Marsh, T. L. RNA catalysis and the origin of life. *Origins Life Evol B* **16,** 97–116 (1985).

2. Gilbert, W. Origin of life: The RNA world. *Nature* **319,** 618–618 (1986).

3. Rich, A. On the problems of evolution and biochemical information transfer. *Horizons in biochemistry* 103–126 (1962).

4. Breaker, R. R. Engineered allosteric ribozymes as biosensor components. *Curr Opin Biotech* **13,** 31–39 (2002).

5. Wittmann, A. & Suess, B. Engineered riboswitches: Expanding researchers' toolbox with synthetic RNA regulators. *Febs Lett* **586,** 2076–2083 (2012).

6. Isaacs, F. J., Dwyer, D. J. & Collins, J. J. RNA synthetic biology. *Nat Biotechnol* **24,** 545–554 (2006).

7. Bartel, D. & Szostak, J. Isolation of new ribozymes from a large pool of random sequences. *Science* **261,** 1411–1418 (1993).

8. Ekland, E., Szostak, J. & Bartel, D. Structurally complex and highly active RNA ligases derived from random RNA sequences. *Science* **269,** 364–370 (1995).

9. Robertson, M. P., Hesselberth, J. R. & Ellington, A. D. Optimization and optimality of a short ribozyme ligase that joins non-Watson–Crick base pairings. *Rna* **7,** 513–523 (2001).

10. Rogers, J. & Joyce, G. F. The effect of cytidine on the structure and function of an RNA ligase ribozyme. *Rna* **7,** 395–404 (2001).

11. Paul, N. & Joyce, G. F. A self-replicating ligase ribozyme. *Proc National Acad Sci* **99,** 12733–12740 (2002).

12. Lincoln, T. A. & Joyce, G. F. Self-Sustained Replication of an RNA Enzyme. *Science* **323,** 1229–1232 (2009).

13. Saragliadis, A., Krajewski, S. S., Rehm, C., Narberhaus, F. & Hartig, J. S. Thermozymes:
synthetic RNA thermometers based on ribozyme activity. *Rna Biol* **10,** 1009–1016 (2013).

14. Ferretti, A. C. & Joyce, G. F. Kinetic Properties of an RNA Enzyme That Undergoes Self-
Sustained Exponential Amplification. *Biochemistry-us* **52,** 1227–1235 (2013).

15. Lam, B. J. & Joyce, G. F. Autocatalytic aptazymes enable ligand-dependent exponential
amplification of RNA. *Nat Biotechnol* **27,** 288–292 (2009).

16. Janas, T., Janas, T. & Yarus, M. A membrane transporter for tryptophan composed of
RNA. *Rna* **10,** 1541–1549 (2004).

17. Pino, S., Ciciriello, F., Costanzo, G. & Mauro, E. D. Nonenzymatic RNA Ligation in Water*.
J Biol Chem **283,** 36494–36503 (2008).

18. Guerrier-Takada, C., Gardiner, K., Marsh, T., Pace, N. & Altman, S. The RNA moiety of
ribonuclease P is the catalytic subunit of the enzyme. *Cell* **35,** 849–857 (1983).

19. Anella, F. & Danelon, C. Prebiotic Factors Influencing the Activity of a Ligase Ribozyme.
Life **7,** 17 (2017).

20. Dai, X., Zhang, S. & Zaleta-Rivera, K. RNA: interactions drive functionalities. *Mol Biol Rep*
47, 1413–1434 (2020).

21. Müller, U. F. & Bartel, D. P. Improved polymerase ribozyme efficiency on hydrophobic
assemblies. *Rna* **14,** 552–562 (2008).

22. Anella, F. & Danelon, C. Reconciling Ligase Ribozyme Activity with Fatty Acid Vesicle
Stability. *Life* **4,** 929–943 (2014).

23. Chen, I. A., Salehi-Ashtiani, K. & Szostak, J. W. RNA Catalysis in Model Protocell Vesicles. *J
Am Chem Soc* **127,** 13213–13219 (2005).

24. Mansy, S. S., Schrum, J. P., Krishnamurthy, M., Tobé, S., Treco, D. A. & Szostak, J. W.
Template-directed synthesis of a genetic polymer in a model protocell. *Nature* **454,** 122–125
(2008).

25. Bianconi, G., Zhao, K., Chen, I. A. & Nowak, M. A. Selection for Replicases in Protocells.
Plos Comput Biol **9,** e1003051 (2013).

26. Budker, V. G., Kazatchkov, Yu. A. & Naumova, L. P. Polynucleotides adsorb on
mitochondrial and model lipid membranes in the presence of bivalent cations. *Febs Lett* **95,**
143–146 (1978).

27. McManus, J. J., Rädler, J. O. & Dawson, K. A. Does Calcium Turn a Zwitterionic Lipid
Cationic? *J Phys Chem B* **107,** 9869–9875 (2003).

28. Kharakoz, D. P., Khusainova, R. S., Gorelov, A. V. & Dawson, K. A. Stoichiometry of
dipalmitoylphosphatidylcholine-DNA interaction in the presence of Ca2+: a temperature-
scanning ultrasonic study. *Febs Lett* **446,** 27–29 (1999).

29. Gromelski, S. & Brezesinski, G. Adsorption of DNA to zwitterionic DMPE monolayers
mediated by magnesium ions. *Phys Chem Chem Phys* **6,** 5551–5556 (2004).

30. Uhríková, D., Hanulová, M., Funari, S. S., Khusainova, R. S., Šeršeň, F. & Balgavý, P. The
structure of DNA–DOPC aggregates formed in presence of calcium and magnesium ions: A
small-angle synchrotron X-ray diffraction study. *Biochimica Et Biophysica Acta Bba -*
Biomembr **1713,** 15–28 (2005).

31. Gromelski, S. & Brezesinski, G. DNA Condensation and Interaction with Zwitterionic
Phospholipids Mediated by Divalent Cations. *Langmuir* **22,** 6293–6301 (2006).

32. McLoughlin, D., Dias, R., Lindman, B., Cardenas, M., Nylander, T., Dawson, K., Miguel, M.
& Langevin, D. Surface Complexation of DNA with Insoluble Monolayers. Influence of
Divalent Counterions. *Langmuir* **21,** 1900–1907 (2005).

33. Michanek, A., Yanez, M., Wacklin, H., Hughes, A., Nylander, T. & Sparr, E. RNA and DNA
Association to Zwitterionic and Charged Monolayers at the Air–Liquid Interface. *Langmuir*
28, 9621–9633 (2012).

34. Michanek, A., Kristen, N., Höök, F., Nylander, T. & Sparr, E. RNA and DNA interactions
with zwitterionic and charged lipid membranes — A DSC and QCM-D study. *Biochimica Et*
Biophysica Acta Bba - Biomembr **1798,** 829–838 (2010).

35. Marty, R., N'soukpoé-Kossi, C. N., Charbonneau, D. M., Kreplak, L. & Tajmir-Riahi, H.-A.
Structural characterization of cationic lipid–tRNA complexes. *Nucleic Acids Res* **37,** 5197–
5207 (2009).

36. Suga, K., Tanabe, T., Tomita, H., Shimanouchi, T. & Umakoshi, H. Conformational change
of single-stranded RNAs induced by liposome binding. *Nucleic Acids Res* **39,** 8891–8900
(2011).

37. Lu, D. & Rhodes, D. G. Binding of phosphorothioate oligonucleotides to zwitterionic
liposomes. *Biochimica Et Biophysica Acta Bba - Biomembr* **1563,** 45–52 (2002).

38. Khvorova, A., Kwak, Y.-G., Tamkun, M., Majerfeld, I. & Yarus, M. RNAs that bind and
change the permeability of phospholipid membranes. *Proc National Acad Sci* **96,** 10649–
10654 (1999).

39. Vlassov, A. & Yarus, M. Interaction of RNA with Phospholipid Membranes. *Mol Biol+* **36,**
389–393 (2002).

40. Vlassov, A., Khvorova, A. & Yarus, M. Binding and disruption of phospholipid bilayers by
supramolecular RNA complexes. *Proc National Acad Sci* **98,** 7706–7711 (2001).

41. Janas, T., Janas, T. & Yarus, M. Specific RNA binding to ordered phospholipid bilayers.
Nucleic Acids Res **34,** 2128–2136 (2006).

42. Pannwitt, S., Slama, K., Depoix, F., Helm, M. & Schneider, D. Against Expectations:
Unassisted RNA Adsorption onto Negatively Charged Lipid Bilayers. *Langmuir* **35,** 14704–
14711 (2019).

43. Kruger, K., Grabowski, P. J., Zaug, A. J., Sands, J., Gottschling, D. E. & Cech, T. R. Self-
splicing RNA: Autoexcision and autocyclization of the ribosomal RNA intervening sequence
of tetrahymena. *Cell* **31,** 147–157 (1982).

44. Quarless, S. A. & Cantor, C. R. Analysis of RNA structure by ultraviolet crosslinking and
denaturation gel electrophoresis. *Anal Biochem* **147,** 296–300 (1985).

45. Behlen, L. S., Sampson, J. R. & Uhlenbeck, O. C. An ultraviolet light-induced crosslink in
yeast tRNA phe. *Nucleic Acids Res* **20,** 4055–4059 (1992).

46. Nejedlý, K., Kittner, R., Pospíšilová, Š. & Kypr, J. Crosslinking of the complementary
strands of DNA by UV light: dependence on the oligonucleotide composition of the UV
irradiated DNA. *Biochimica Et Biophysica Acta Bba - Gene Struct Expr* **1517,** 365–375 (2001).

47. Lengyel, A., Uhríková, D., Klacsová, M. & Balgavý, P. DNA condensation and its thermal
stability influenced by phospholipid bilayer and divalent cations. *Colloids Surfaces B
Biointerfaces* **86,** 212–217 (2011).

48. Black, R. A., Blosser, M. C., Stottrup, B. L., Tavakley, R., Deamer, D. W. & Keller, S. L.
Nucleobases bind to and stabilize aggregates of a prebiotic amphiphile, providing a viable
mechanism for the emergence of protocells. *Proc National Acad Sci* **110,** 13272–13276
(2013).

49. White, S. H., Wimley, W. C., Ladokhin, A. S. & Hristova, K. [4] Protein folding in
membranes: Determining energetics of peptide-bilayer interactions. *Methods Enzymol* **295,**
62–87 (1998).

50. Lee, J. S., Evans, D. H. & Morgan, A. R. Polypurine DNAs and RNAs form secondary
structures which may be tetra-stranded. *Nucleic Acids Res* **8,** 4305–4320 (1980).

51. Baross, J. A. & Martin, W. F. The Ribofilm as a Concept for Life's Origins. *Cell* **162,** 13–15
(2015).

52. Johansson, J., Mandin, P., Renzoni, A., Chiaruttini, C., Springer, M. & Cossart, P. An RNA
Thermosensor Controls Expression of Virulence Genes in Listeria monocytogenes. *Cell* **110,**
551–561 (2002).

53. Winkler, W., Nahvi, A. & Breaker, R. R. Thiamine derivatives bind messenger RNAs
directly to regulate bacterial gene expression. *Nature* **419,** 952–956 (2002).

54. Morita, M. T., Tanaka, Y., Kodama, T. S., Kyogoku, Y., Yanagi, H. & Yura, T. Translational
induction of heat shock transcription factor ç32: evidence for a built-in RNA thermosensor.
Gene Dev **13,** 655–665 (1999).

55. Schilling, O., Langbein, I., Müller, M., Schmalisch, M. H. & Stülke, J. A protein-dependent
riboswitch controlling ptsGHI operon expression in Bacillus subtilis: RNA structure rather
than sequence provides interaction specificity. *Nucleic Acids Res* **32,** 2853–2864 (2004).

56. Nahvi, A., Sudarsan, N., Ebert, M. S., Zou, X., Brown, K. L. & Breaker, R. R. Genetic Control
by a Metabolite Binding mRNA. *Chem Biol* **9,** 1043–1049 (2002).

57. Patel, B. H., Percivalle, C., Ritson, D. J., Duffy, C. D. & Sutherland, J. D. Common origins of
RNA, protein and lipid precursors in a cyanosulfidic protometabolism. *Nat Chem* **7,** 301–307
(2015).

58. Rushdi, A. I. & Simoneit, B. R. T. Lipid Formation by Aqueous Fischer-Tropsch-Type
Synthesis over a Temperature Range of 100 to 400 °C. *Origins Life Evol B* **31,** 103–118 (2001).

59. Brady, R. C. & Pettit, R. Mechanism of the Fischer-Tropsch reaction. The chain
propagation step. *J Am Chem Soc* **103,** 1287–1289 (1981).

60. Robertson, M. P. & Ellington, A. D. Design and optimization of effector-activated
ribozyme ligases. *Nucleic Acids Res* **28,** 1751–1759 (2000).

61. Debiais, M., Lelievre, A., Smietana, M. & Müller, S. Splitting aptamers and nucleic acid
enzymes for the development of advanced biosensors. *Nucleic Acids Res* **48,** 3400–3422
(2020).

62. Stifel, J., Spöring, M. & Hartig, J. S. Expanding the toolbox of synthetic riboswitches with
guanine-dependent aptazymes. *Synthetic Biology* **4,** ysy022- (2019).

63. Janas, T. & Yarus, M. Visualization of membrane RNAs. *Rna* **9,** 1353–1361 (2003).

64. Walker, S. C., Avis, J. M. & Conn, G. L. General plasmids for producing RNA in vitro
transcripts with homogeneous ends. *Nucleic Acids Res* **31,** e82–e82 (2003).

65. Chen, P. S., Toribara, T. Y. & Warner, H. Microdetermination of Phosphorus. *Anal Chem*
28, 1756–1758 (1956).

Materials and methods

Materials

Dipalmitoylphosphatidylcholine (DPPC) and biotinylated phosphatidylethanolamine (18:1 biotinyl cap PE) were purchased from Avanti Polar Lipids (USA) and were used without further purification. HEPES, $MgCl_2$ and $CaCl_2$ were purchased from CarlRoth (Germany) and were at least >99% pure. All solutions were prepared in MilliQ water, Merck Millipore (USA).

The R3C ligase construct[10] was cloned into the pRZ plasmid[64] using the InFusion cloning system (Takara Bio, Japan). To ensure correct length of transcript, pRZ-R3C plasmid was treated with EcoRI-HF (NEB, USA). RNA wasexpressed using T7 RNA polymerase (homemade, MPI-CBG, Germany) at 37 °C overnight, followed by 10 cycles of 5' 60 °C -> 5' 24 °C (HDV cleavage of construct to release pure R3C) and DNAse I treatment (Thermo Scientific, USA). 75-85 randomised oligomer was expressed using artificial DNA template (Eurofins Genomics, Germany) – both R3C and randomer were purified using phenol-chloroform-isoamyl alcohol extraction, denaturing PAGE and electroelution. 5' 6-FAM labelled R3C substrates as well as other short oligomers used in binding assays were obtained from Integrated DNA Technologies (USA) and used without further purification.

Sequence name	Sequence (5' – 3')
R3C	GAGACCGUAAUGAGUAGUACUUAUUAUGCUCGAUUGUUCGUAAG AACAGUUUGAAUGGGUUGAAUAUAGUGAGGAUCCGCUAGC
Short substrate	6_FAM_CGACUCACUAUA
4xAG substrate	6_FAM_A**G**A**G**A**G**A**G**CGACUCACUAUA
9xAG substrate	6_FAM_A**G**A**G**A**G**A**G**A**G**A**G**A**G**A**G**A**G**ACGACUCACUAUA
2G substrate	6_FAM_HHHHHHHH**G**HHH**G**HHHHHHCGACUCACUAUA
0G	HHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHH
2G	HHHHHHHHHHHHHHH**G**HHH**G**HHHHHHHHHHHHHHH
4G	HHHHHHHHHHH**G**HHH**G**HHH**G**HHH**G**HHHHHHHHHHH
5G	HHHHHHHHH**G**HHH**G**HHH**G**HHH**G**HHH**G**HHHHHHHHHH
7G	HHHHHH**G**HHH**G**HHH**G**HHH**G**HHH**G**HHH**G**HHHHHH
9G	HH**G**HHH**G**HHH**G**HHH**G**HHH**G**HHH**G**HHH**G**HHH**G**HH
40xN	NNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNN
40xN - G	HHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHH
40xN - A	BBBBBBBBBBBBBBBBBBBBBBBBBBBBBBBBBBBBBBBBB
40xN - C	DDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDDD
40xN - U	VVVVVVVVVVVVVVVVVVVVVVVVVVVVVVVVVVVVVVVVV
19xG	**GGGGGGGGGGGGGGGGGGG**
19xA	AAAAAAAAAAAAAAAAAAA
19xC	CCCCCCCCCCCCCCCCCCC
19xU	UUUUUUUUUUUUUUUUUUU
19xN	NNNNNNNNNNNNNNNNNNN
AG repeats	6_FAM_AGAGAGAGAGAGAGAGAGA
ssRNA (ACUG)	ACUGACUGACUGACUGACUGACUGACUGACUG
ssRNA (CAGU)	CAGUCAGUCAGUCAGUCAGUCAGUCAGUCAGU

B – C, G or U

D – A, G or U

H – A, C or U

V – A, C or G

Methods

All of the RNA incubations were prepared in DNA low bind tubes (Sarstedt, Germany) at the constant

temperature of 24 °C in buffer comprised of 10 mM HEPES pH 7, 5 mM $CaCl_2$, and 5 mM $MgCl_2$. RNA

concentration was determined by measuring absorbance at 260 nm (SPARK 20M, TECAN). Before every

incubation RNA was preheated in SafeLock tubes (90 °C, 5 minutes; Eppendorf, Germany) to ensure

unfolded structures. Denaturing PAGE analysis was performed using 10% 19:1

acrylamide:bisacrylamide gel composition, whereas native gels were composed of 8-10% 20:1

acrylamide:bisacrylamide mix. Before application on the denaturing PAGE all of the samples were

ethanol precipitated.

Structure of R3C ligase from Fig. 1d was generated using RNA structure Fold tool from Mathews lab

(https://rna.urmc.rochester.edu/RNAstructureWeb/).

Lipid preparation

DPPC chloroform stock was pipetted into a glass vial and briefly evaporated under a steady flow of

nitrogen gas. To remove organic solvent residues, a lipid film was dried under vacuum overnight. To

obtain multi-layer DPPC vesicles (MLV DPPC), lipid films were hydrated in reaction buffer, with a final

lipid concentration of 10 mM in 1 mL. Buffer-lipid mixtures were shaken at 60 °C, above the melting

temperature of DPPC (T_m = 42 °C), for one hour. A cloudy liposome suspension was freeze-and-thawed

10x and extruded 17x through 100 nm polycarbonate filters (Merck Millipore, USA) to reduce multi-

lamellarity and achieve a consistent size distribution of vesicles. Lipid vesicle stocks were kept at 4 °C.

Stock lipid concentrations were confirmed using a phosphate assay[65].

Dynamic light scattering

The size distribution of DPPC vesicles was estimated using a ZetaSizer Nano in 173° backscatter with

multiple narrow mode (high resolution) analysis. Final concentration of lipids was 10 μM and the

amount of RNA was fixed at a ratio of 10 lipids / nucleotide. Results were plotted as the size distribution

in the number of detected species (NumberPSD).

UV-crosslinking

RNA was incubated for 10 minutes in the reaction buffer with or without lipids followed by 10 minutes

incubation under UV-B light (300 mW LEUVA77N50KKU00 LED 305 nm) from a distance of 1.5 – 2 cm.

Samples were analysed on denaturing PAGE with SybrGold poststaining.

R3C activity assays

5 pmol of R3C ligase was mixed with 0.5 pmol of the substrate to ensure saturation of the system

(pseudo 1st order reaction) and decrease any batch-to-batch differences coming from R3C purification.

Final reaction volume was adjusted to 30 μl. Incubation was conducted at various concentrations of

DPPC vesicles and samples were ethanol precipitated after 30, 60, 90 and 120 minutes. Reactions were

quantified on denaturing PAGE using FIJI software. Product and substrate intensities (I_P and I_S,
respectively) were quantified and the concentration of product at time t was calculated:

$$[P]_t = [S]_0 * \left(\frac{I_P}{I_P + I_S} \right)$$

To determine reaction rates (M/min), concentration of the product was plotted as a function of time
and then fitted with a linear function (fixed [0;0] point).

RNA-lipid binding assays
Binding assays for the results in Fig. 2c (R3C ligase), Fig. 3a, b, c and Fig. 4a were conducted using
liposome-coated magnetic beads. Binding assays for the results in Fig. 1b, c and 4b, c were conducted
using an ultracentrifugation method.

Magnetic beads binding assay
Magnetic beads (DynaBeads streptavidin T1, Invitrogen) were coupled with DPPC liposomes doped
with 0.5 mol% biotynoyl cap PE. Lipid concentration on non-diluted beads was estimated using a
phosphate assay[65]. Liposome conjugated beads final working concentration was of 800 μM lipid.

25 pmol of 19 nt or 10 pmol of 40/36 nt RNAs were incubated with 2 – 1.5x diluted magnetic beads for
30'. After incubation the supernatant was separated from the beads using a magnet. The amount of
RNA left in solution was estimated either using absorbance or Qubit miRNA quantification kit
(ThermoScientific, USA). The reported binding efficiencies (Fig. 3a, b) correspond to 15 μL of liposome-
coated beads in a 30 μL reaction volume. The partition coefficient K (Fig. 3c, Fig. 4a) was calculated
according to following formula:

577
578

$$K = \frac{\dfrac{[R_L]}{[R_L] + [L]}}{\dfrac{[R_w]}{[R_w] + [W]}} \qquad \text{Eq. 1}$$

579
580

582   **Assuming high excess of [L] and [W] we simplified the equation to:**

584
585

$$K = \frac{\dfrac{[R_L]}{[L]}}{\dfrac{[R_w]}{[W]}} \qquad \text{Eq. 2}$$

586
587

Where:
[R_L] – RNA bound to the membranes
[R_W] – non-bound RNA
[L] – outer lipid concentration
[W] – water concentration

Ultracentrifugation binding assay

RNA was incubated with various concentrations of DPPC vesicles for 30'. After incubation
ultracentrifugation was conducted to separate vesicles from the solution (125 000 x G, 40', 24 °C). 9
μL of supernatant (out of 30 μL reaction mix) was pulled and incubated with 1.5 μL of 0.5 M EDTA (90
°C, 5'). After incubation 150 μl of 1xQubit dye was added to the solution and the amount of RNA was
measured using a plate reader (SPARK 20M, TECAN).
601
602   Measured fluorescence intensity values were plotted as a function of lipid/oligomer or lipid/nucleotide
603   ratios (we assumed that RNA interacts only with the outer membrane leaflet, so final outer lipid
604   concentration is equal half of the total lipid amount) and fitted using Hill's fit (OriginLab software):
605
606
$$f(x) = Start + (End - Start)\frac{x^n}{x^n + k^n} \quad \text{Eq. 3}$$

607
608   Where Start and End are function plateau values, k is inflection point of curve and n is cooperativity
609   index. Inflection point of the curve (k) is a parameter which we used for further data analysis.
610
RNA-lipid binding partition coefficient determination for R3C substrates
5 pmol of RNA was incubated with an excess of DPPC vesicles (5 mM, 2.5 mM outer leaflet) for 30
minutes. After incubation ultracentrifugation was conducted to separate vesicles from solution (125
000 x G, 40', 24 °C). 10 μL out of 30 μL was taken as a supernatant sample, and the rest of the
supernatant was discarded to leave only the pellet. RNA from supernatant and pellet samples was
ethanol precipitated and analyzed using denaturing PAGE. Partition coefficient values (Fig. 2c
"substrate", Fig. 5c) were calculated using Eq. 2.
618
619
620

Fig. 1 **RNA-lipid interactions induce reversible aggregation of vesicles and enhance intermolecular RNA interactions. (a)** RNA-induced vesicular aggregation can be observed through changes in vesicle size distribution measured by dynamic light scattering (left), or visually as depicted in the cuvette images (right). The introduction of a randomized pool of RNA oligos to gel lipid membranes (gel phase = **G**) induces vesicle aggregation, which can be reversed either through chelation of divalent cations with EDTA or by increasing the temperature to achieve membranes in a liquid phase (liquid phase = **L**). **(b)** R3C ligase (84 nt) shows a small increase in UV-crosslinking at low lipid concentration, whereas crosslinking efficiency decreases at higher lipid concentration. Binding and UV crosslinking are inversely correlated and show similar inflection points with respect to lipid concentration. **(c)** Randomized RNA oligos (75-85 nt) show increasing crosslinking efficiency in the presence of lipids with a slight decrease at the highest lipid concentration; crosslinking and lipid binding are closely correlated. **(d)** Analysis of RNAs by native PAGE gel demonstrates that the R3C ligase forms primarily one structure, whereas randomized RNA forms diverse intermolecular complexes.

Fig. 2 **Lipid membranes influence R3C ligase activity through lipid-ligase interactions. (a)** R3C ligase reaction rates were quantified by measuring the amount of product within a 2 hour incubation time. **(b)** R3C reaction rate is dependent on the presence of gel membranes - increased activity is observed for lower lipid concentration followed by drop in activity at higher lipid concentration. **(c)** The 12 nt R3C substrate shows several orders of magnitude lower membrane-buffer partition coefficient log(K) compared with the R3C ligase, which could be derived from different compositional or structural characteristics of both RNA species.

Fig. 3 **RNA-lipid binding is RNA sequence-dependent. (a)** Short 19 nucleotide oligomers show sequence-dependent binding to the gel membranes, with highest binding observed for the pure G oligomer followed by oligos with randomized sequence or AG repeats. Oligomers containing only A, C or U bases show negligible binding to the lipid membranes. **(b)** Oligonucleotide length also influences binding efficiency as seen in the comparison of **(a)** 19xN and **(b)** 40xN. Depletion of one type of nucleotide within oligomers lowers binding efficiency. Lack of guanine decreased binding by 57%, whereas lack of adenine, uracil or cytidine showed a less pronounced effect. **(c)** Oligonucleotides with increasing G content showed higher lipid – buffer partition coefficients compared to a G-deficient oligo with the highest value observed for the 9xG oligomer.

Fig. 4 **RNA-gel membrane binding depends on RNA structure. (a)** The effect of RNA base-pairing structure on lipid vesicle aggregation (left) and membrane binding (right) was evaluated by comparing RNA's with limited ability to create inter- and intramolecular structures (ssRNA: 36 nucleotide ACUG or CAGU repeats) with 100% complementary double stranded RNAs (dsRNA: mixed ACUG and CAGU oligos). Single stranded RNAs do not cause lipid vesicle aggregation despite moderate membrane binding, whereas double stranded RNA species show both high binding and vesicle aggregation. **(b)** RNA:lipid binding ratios (shown in table) estimated from binding curves of ssRNA and dsRNA (plotted) are affected by base pairing and sequence. **(c)** Oligomers with different nucleotide composition show different lipid/nucleotide binding stoichiometry. Analysis of oligomers by native PAGE gel **(c, inset)** indicates that the presence of guanine promotes inter-molecular interactions, suggesting, that base-pairing enhances the average strength of nucleotide-lipid interactions.

Fig. 5   Modification of the R3C substrate sequences enables lipid-dependent tunability of ligation rate. (a) Modified R3C reaction substrates were synthesized in order to increase binding to the membrane. The catalytic 3' end of the molecule was enriched with 5' tails with varying G content. Reactions for modified substrates showed lower reaction rate constants compared with the unmodified substrate. **(b)** The mechanism for inhibition of activity was evaluated by analysis of R3C reactants using native PAGE. The unmodified short substrate is entirely bound to the ligase, whereas AG-modified substrates show mobility abnormalities, which suggest substrate folding (magenta points). **(c)** Modification of the substrate sequence through addition of a 5' overhang increases binding to the membrane. **(d)** Addition of lipid membranes to reactions with the G-rich substrates increases reaction rates to the level of the unmodified substrate. **(e)** We propose that the decrease of R3C reaction rates in the presence of G-rich substrates is based on the inhibition of the catalytic part of the substrate by the 5' G-rich overhang. The presence of lipid membranes not only increases R3C-substrate stability, but also screens the 5'rich substrate part enabling the R3C reaction to proceed.