Gene Conversion Facilitates the Adaptive Evolution of Self-Resistance in Highly Toxic Newts

Kerry L. Gendreau[1]*, Angela D. Hornsby[1,2], Michael T. J. Hague[3], and Joel W. McGlothlin[1]

[1]Department of Biological Sciences, Virginia Tech, Blacksburg, United States
[2]Philip L. Wright Zoological Museum, Division of Biological Sciences, University of Montana, Missoula, United States
[3]Division of Biological Sciences, University of Montana, Missoula, MT, United States

*Corresponding author: kerryg@vt.edu

Abstract

Reconstructing the histories of complex adaptations and identifying the evolutionary mechanisms underlying their origins are two of the primary goals of evolutionary biology. *Taricha* newts, which contain high concentrations of the deadly toxin tetrodotoxin (TTX) as an antipredator defense, have evolved resistance to self-intoxication, which is a complex adaptation requiring changes in six paralogs of the voltage-gated sodium channel (Na_v) gene family, the physiological target of TTX. Here, we reconstruct the origins of TTX self-resistance by sequencing the entire Na_v gene family in newts and related salamanders. We show that moderate TTX resistance evolved early in the salamander lineage in three of the six Na_v paralogs, preceding the proposed appearance of tetrodotoxic newts by ~100 million years. TTX-bearing newts possess additional unique substitutions across the entire Na_v gene family that provide physiological TTX resistance. These substitutions coincide with signatures of positive selection and relaxed purifying selection, as well as gene conversion events, that together likely facilitated their evolution. We also identify a novel exon duplication within Na_v1.4 encoding an expressed TTX-binding site. Two resistance-conferring changes within newts appear to have spread via nonallelic gene conversion: in one case, one codon was copied between paralogs, and in the second, multiple substitutions were homogenized between the duplicate exons of Na_v1.4. Our results demonstrate that gene conversion can accelerate the coordinated evolution of gene families in response to a common selection pressure.

Key words: gene conversion, adaptation, coevolutionary arms races, molecular evolution, newts, toxin resistance

Introduction

Fitting evolutionary models to molecular sequences in a phylogenetic context can help piece together the key steps in adaptive evolution and uncover the relative contributions of selection and other evolutionary mechanisms to adaptive phenotypic evolution (Smith et al. 2020). Comparative studies of convergence, or the repeated evolution of characters within different lineages undergoing the same environmental challenges, provide powerful evidence of both adaptation and connections between genetic and

phenotypic change (Losos 2011). Investigations into the molecular basis of convergence have revealed

multiple occurrences of parallelism, where different lineages have evolved changes within the same

proteins, and occasionally at the same amino acid sites, in response to shared selective pressures, such as

insects that have evolved the ability to feed on toxic plants (Zhen et al. 2012) and populations of ducks

and humans living at high elevations (Graham and McCracken 2019). Such patterns support important

roles for both positive selection and constraint in the origin of complex adaptations (reviewed by Storz

2016).

48   Resistance to tetrodotoxin (TTX), a potent neurotoxin, has evolved convergently in several

distantly related organisms, including pufferfish, snakes, and newts (reviewed by Soong and Venkatesh

2006; Toledo et al. 2016), and therefore offers an ideal system to investigate the molecular basis of

adaptive evolution (Arbuckle et al. 2017). The genetic basis of TTX resistance, which is well established

in tetrodotoxic puffer fish (Jost et al. 2008) and in snakes that consume TTX-bearing prey (Geffeney et al.

2002; Feldman et al. 2012; McGlothlin et al. 2014; McGlothlin et al. 2016), involves amino acid

substitutions in the toxin's target, voltage-gated sodium channels. While voltage-gated sodium channels

are commonly abbreviated as *SCNA* genes and Na_v proteins, hereafter we simplify this nomenclature by

abbreviating both as Na_v. Na_v channels are responsible for the initiation and propagation of action

potentials in excitable cells and are composed of four domains (DI–DIV), each comprising six

transmembrane helices and a pore-loop region (the P-loop; Fux et al. 2018). The four P-loops, one in each

domain, form a pore within the membranes of excitable cells to selectively allow sodium ions to cross

when the channel is open. TTX exerts its effects by binding to the P-loops of sensitive channels and

preventing sodium entry into cells, thus blocking action potentials. Gene duplication events have resulted

in six Na_v paralogs, each with tissue-specific expression, that are shared across all tetrapods (table 1),

with additional lineage-specific duplications occurring in amniotes (Widmark et al. 2011; Zakon 2012).

Because the structure of these paralogs is highly conserved, each has the potential to be blocked by TTX

if it lacks resistance-conferring substitutions.

66   Species that possess or consume TTX must either have a full complement of resistant paralogs or

otherwise shield sodium channels from contact with the toxin. Indeed, resistant substitutions are present

in all eight of the Na_v genes within the genomes of multiple species of TTX-bearing pufferfish (from the

family Tetraodontidae; Jost et al. 2008) and in six of the nine Na_v genes in *Thamnophis sirtalis* snakes

that consume TTX-bearing *Taricha* newts (McGlothlin et al. 2014; Perry et al. 2018). The three brain

channels $Na_v1.1$, $Na_v1.2$, and $Na_v1.3$ of snakes remain TTX sensitive, but presumably are protected from

TTX by the blood-brain barrier (McGlothlin et al. 2014). In snakes, the evolution of extreme TTX

resistance appears to follow a predictable, stepwise substitution pattern across TTX-exposed members of

the Na_v gene family, with substitutions in heart and peripheral nerve channels preceding those in the

Table 1. Nomenclature for Voltage-Gated Sodium Channel Genes

Gene name	Protein name	Tissue expression[a]
SCN1A	$Na_V1.1$	Brain
SCN2A	$Na_V1.2$	Brain
SCN3A	$Na_V1.3$	Brain
SCN4A	$Na_V1.4$	Muscle
SCN5A	$Na_V1.5$	Heart
SCN8A	$Na_V1.6$	Brain / peripheral nervous system

[a] Patterns of tissue expression are inferred from studies of gene orthologs in mammals (reviewed in Yu and Catterall 2003).

muscle channel gene, $Na_v1.4$, which evolves resistance only in snakes locked in coevolutionary arms

races with highly tetrodotoxic amphibian prey (Feldman et al. 2012; McGlothlin et al. 2016; Perry et al.

2018).

Less is known about the evolutionary history of TTX resistance in *Taricha* newts, the highly toxic

coevolutionary partner of *Thamnophis* (Brodie and Brodie 1990; Brodie et al. 2002; Williams et al. 2010;

Hague et al. 2020). The extreme toxicity of *Taricha*, which has been elaborated by the ongoing

coevolutionary arms race with garter snakes, builds upon lower levels of toxicity that evolved ~30 million

years ago (mya) within "modern" newts (tribe Molgini; Hanifin and Gilly 2015; divergence date

estimated by Hime et al. 2021). The evolution of toxicity necessitates the evolution of toxin

autoresistance so that a prey species is not incapacitated by its own antipredator defense (Jost et al. 2008;

Toledo et al. 2016; Tarvin et al. 2017; Márquez et al. 2019). Understanding the timing and details of this

autoresistance can shed light on the genetic processes underlying the predator-prey arms race. Hanifin and

Gilly (2015) compared the sequences of one sodium channel gene, the muscle paralog $Na_v1.4$, across

several salamander species and identified substitutions in the P-loops of DIII and DIV that provide

extreme TTX resistance to the muscles of TTX-bearing newts. Importantly, the sister group of these toxic

newts had substitutions in the same gene providing more moderate resistance, indicating that the

evolution of autoresistance in a common ancestor paved the way for the evolution of extreme toxicity.

More recently, Vaelli et al. (2020) used transcriptome sequencing to characterize the genetic basis of

physiological resistance to TTX in *Taricha granulosa* and identified substitutions within TTX-binding

regions in the other five Na_v paralogs, many of which occur in within the P-loop of DI. However, because

it is unknown whether other salamander species possess TTX resistance in these paralogs, the order and

timing of the evolutionary events leading to autoresistance in toxic newts is still unknown. Furthermore,

no studies to date have applied evolutionary models to test for the relative importance of mechanisms

such as positive selection, relaxed constraint, and interlocus gene conversion in the evolution of newt

TTX resistance.

Here, we trace the evolutionary history of the entire Na_v gene family across the salamander

phylogeny to show the order in which resistant substitutions appeared. Using published genome

sequences and newly generated sequence data, we characterize the genomic structure of Na_v genes in

newts and their relatives, inferring the timing of resistant substitutions leading to the extreme TTX

resistance observed across all Na_v paralogs in *Taricha* newts (Vaelli et al. 2020). We estimate rates of

synonymous and nonsynonymous substitution to identify positive selection. In addition, we assess the

potential of nonallelic gene conversion, a process by which sequence is copied from one paralog to

another (Chen et al. 2007), to act as a source of adaptive variation. Combining these data provides insight

into the evolutionary mechanisms underlying the origin of a uniquely potent chemical defense.

Results

Genomic Structure of Voltage-Gated Sodium Channels

We used targeted sequence capture to characterize Na_v sequences from the genomes of five salamander

species (order Urodela), including three TTX-bearing newts (family Salamandridae, subfamily

Pleurodelinae, tribe Molgini), *Notophthalmus viridescens*, *Taricha torosa*, and *Taricha granulosa* (*n* = 3

diploid individuals of each species), and two salamanders that do not possess tetrodotoxin,

Cryptobranchus alleganiensis (Crypotobranchidae) and *Plethodon cinereus* (Plethodontidae, *n* = 2 each).

We also identified Na_v sequences within two publicly available salamander genome sequences:

Ambystoma mexicanum (Ambystomatidae; Smith et al. 2019; AmexG.v6 assembly) and *Pleurodeles waltl*

(Salamandridae; Elewa et al. 2017) and a full-body transcriptome from the fire salamander *Salamandra*

salamandra (Salamandridae; Goedbloed et al. 2017; BioProject accession PRJNA607429), all three of

which appear to lack tetrodotoxin (Hanifin 2010). The split between *Cryptobranchus* (suborder

Cryptobranchoidea) and all the other salamanders in our study (members of suborder Salamandroidea)

represents the most ancient division in the phylogeny of extant salamanders (~160 mya; Hime et al.

2021).

We identified six Na_v genes in the genomes of all salamander species, which is consistent with

observations in other amphibians (Zakon et al. 2011). Hereafter, we use the exon delineation introduced

in Widmark et al. (2011), where domain I (DI) is encoded by exons 2-9, domain II (DII) is encoded by

exons 13-16, domain III (DIII) is encoded by exons 18-23, and domain IV (DIV) is encoded by exons 25-

26. We obtained near full-length assemblies for all paralogs (table S1); however, a few exons containing

TTX-binding sites, including exon 15 (encoding the DII P-loop) of Na_v1.2 from *N. viridescens* and exon

22 (encoding part of the DIII P-loop) of Na_v1.2 for several newt species, were missing from our

assemblies. Polymorphism was rare in our assemblies and we observed few nonsynonymous mutations

within the newt genomes, but we found slightly elevated polymorphism in *N. viridescens* relative to other

Table 2. Locations of Na$_v$ Genes in the *Ambystoma mexicanum* AmexG.v6 Genome Assembly.

Gene	Chromosome	Start	End	Strand	Length (bp)
Na$_v$1.1	9q	503,107,904	503,797,933	+	690,029
Na$_v$1.2	9q	507,685,503	508,688,108	-	1,002,605
Na$_v$1.3	9q	509,830,827	510,576,927	+	746,100
Na$_v$1.4	13q	113,343,821	115,707,581	+	2,363,760
Na$_v$1.4 exon 26b	13q	115,891,071	115,892,251	+	1,180
Na$_v$1.5	2p	562,159,042	563,030,267	-	871,225
Na$_v$1.6	3q	465,212,114	466,692,047	-	1,479,933

species (table S2). No heterozygosity or nonsynonymous polymorphisms were observed within any of the
known TTX-binding P-loop regions within any of the species sequenced for this study.
Synteny of Na$_v$ genes in *A. mexicanum* is conserved relative to other tetrapods (fig. S1), which
allowed us to use the *A. mexicanum* sequences as a baseline to confidently identify Na$_v$ paralogs in all
species. Three of the paralogs, Na$_v$1.1, Na$_v$1.2, and Na$_v$1.3 are arrayed in tandem on *A. mexicanum*
chromosome 9, with Na$_v$1.2 inverted relative to its neighboring paralogs (table 2, fig. S1). The additional
three paralogs, Na$_v$1.4, Na$_v$1.5, and Na$_v$1.6, are each located on separate chromosomes (table 2). In the
gene family tree built from amino acid sequences, all salamander Na$_v$ proteins formed a monophyletic
clade (bootstrap support > 98%) with the corresponding orthologs from the genomes of the frogs *Xenopus*
tropicalis and *Nanorana parkeri*, which we included as outgroups (fig. 1). The gene family tree
constructed from the nucleotide coding sequences of these genes yielded a similar topology, with each
salamander Na$_v$ ortholog forming a monophyletic clade. However, in the nucleotide gene family tree, the
three *X. tropicalis* nerve channels Na$_v$1.1, Na$_v$1.2, and Na$_v$1.3 formed a monophyletic clade that is distinct
from the salamander sequences (bootstrap support 86%; fig. S2). The same tree topology was resolved
when partitioning for the third codon position (see Supplemental Data on Dryad).
149
Partial Duplication of Na$_v$1.4 and Evolution of TTX Resistance in Duplicated Domains
Our search of the *A. mexicanum* genome revealed a partial tandem duplication of the 3' end of the Na$_v$1.4
gene, including the full coding region of exon 26, located ~180,000 base pairs downstream of the full-
length Na$_v$1.4 gene on the same DNA strand (table 2). Both exon 26 copies are similar in length to each
other and to exon 26 of other paralogs, encoding open reading frames of approximately 390 base pairs
without introduced stop codons. Exon 26 is the 3'-terminal exon of the Na$_v$1.4 gene and encodes the TTX-
binding P-loop region of DIV. Hereafter, we refer to the duplicate exons as 26a (more proximal to exon
25) and 26b (more distal to exon 25) and duplicate P-loop regions as DIVa (more proximal to exon 25)

and DIVb (more distal to exon 25). We also found this duplicated exon in Na$_v$1.4 orthologs within the

genomes of salamanders *P. cinereus*, *P. waltl*, *N. viridescens*, *T. torosa*, and *T. granulosa* and in

published transcriptomes of *Tylototriton wenxianensis* and *Bolitoglossa vallecula*, but not in the

transcriptome of *Hynobius retardatus* or in the genomes of *C. alleganiensis* or the frogs *X. tropicalis* or

N. parkeri. This pattern suggests that the duplication event likely took place after the split of

Cryptobranchoidea and Salamandroidea (fig. 2). Within the *S. salamandra* transcriptome, we found four

Fig. 1. Evolutionary relationships among salamander voltage-gated sodium channels. Midpoint rooted maximum likelihood tree constructed from alignment of 2188 amino acids from full coding sequence translations of salamander sodium channel genes, including sequences from two frogs (*Nanorana parkeri* and *Xenopus tropicalis*) and one fish (*Danio rerio*) as outgroups. Black circles indicate nodes with >90% bootstrap support.

Fig. 2. Ancestral duplication and convergent evolution of TTX resistance in Na$_v$1.4 terminal exon 26. Maximum likelihood tree constructed from 1050 bp nucleotide alignment of Na$_v$1.4 exon 26 identified in salamander genomes and transcriptomes. "Exon 26a" and "exon 26b" in tip labels refer to the exon copy more proximal and more distal to exon 25, respectively. Black circles indicate node bootstrap support >95%. "R" at tips indicates the presence of substitutions conferring extreme (orange) and moderate (blue) TTX resistance. The same tree topology was resolved using an alignment of amino acids translations of these sequences.

unique RNA sequences transcribed from the Na$_v$1.4 locus, with alternative splicing of exon 17 and

alternative encoding of either exon 26a or 26b. Genome-mapped reads of multi-tissue transcriptomes of

A. mexicanum (Bryant et al. 2017; Caballero-Pérez et al. 2018; Nowoshilow et al. 2018) indicate that

these alternative transcripts have similar expression profiles across various tissues. Taken together, these

observations provide evidence that the duplication of exon 26 led to the creation of functional splice

variants in these salamanders.

While numerous nonsynonymous substitutions differentiated the duplicated $Na_v1.4$ exon 26

sequences relative to the original sequences within each genome, we found that identical substitutions

conferring extreme TTX-resistance to toxic newts (Hanifin and Gilly 2015) were present in both exons

from the genomes of all three TTX-bearing newts but not in other, less toxic salamander species (fig. 2).

Also consistent with the results of Hanifin and Gilly (2015), we found resistant substitutions conferring

moderate TTX resistance in exon 26a of *P. waltl*, *T. wenxianensis*, and *S. salamandra*; however, we

observed no resistant substitutions in exon 26b outside of the toxic newt clade.

Evolution of TTX Resistance in Salamanders

We characterized levels of TTX resistance in each Na_v paralog as extreme, moderate, and TTX-sensitive

based on previous site-directed mutagenesis experiments in which substitutions were introduced to TTX-

sensitive Na_v channels and cross-membrane Na^+ current was measured *in vitro* in the presence and

absence of TTX (table 3). Our results confirm that *T. granulosa* has six paralogs with extreme TTX

resistance (table 3, fig. 3). Our findings are consistent with those reported by Vaelli et al. (2020) with one

exception: we associate DIV (encoded by exon 26) substitutions A1529G, G1533V, and G1533A with

$Na_v1.1$ and Q1524E, G1533R, and G1533Q with $Na_v1.2$ based on synteny mapping (fig. S1), gene trees

(figs. 1, S2), and a phylogeny created from a coding sequence alignment of exon 26 (fig. S3), whereas the

previous study reversed these assignments. We also show that substitutions with extreme TTX resistance

are present in all six Na_v paralogs in two other species of highly toxic newt, *T. torosa* and *N. viridescens*,

indicating that the common ancestor of these three species possessed extreme TTX resistance. Many of

the substitutions in toxic newts parallel those found in TTX-bearing fish and in snakes that consume

tetrodotoxic amphibians (table 3).

No salamander species outside the clade of highly toxic newts possessed a full complement of

TTX-resistant Na_v paralogs, indicating that the evolution of full physiological resistance coincided with

the origin of extreme toxicity. However, we found at least three paralogs with moderate or extreme TTX

resistance in all salamander species we examined, indicating that the evolution of TTX resistance in newts

built upon more ancient changes that first appeared in their non-newt relatives. Substitutions conferring

moderate or extreme resistance were observed within the heart channel $Na_v1.5$ and brain/nerve channels

$Na_v1.1$ and $Na_v1.6$ of all salamander species, with additional resistance-conferring substitutions evolving

within TTX-bearing newts. As first shown by Hanifin and Gilly (2015), moderate resistance was present

in the skeletal muscle channel $Na_v1.4$ of *S. salamandra* and *P. waltl*, but not in the three other salamander

species we examined. Although our outgroup, the frog *X. tropicalis*, also contained a highly resistant

substitution in Na$_v$1.2, we found no evidence for resistance in this paralog in any salamanders outside of

tetrodotoxic newts.

Based on our ancestral sequence reconstructions, the most recent common ancestor of all

salamanders had three TTX resistant sodium channels: Na$_v$1.1 (brain, moderately resistant), Na$_v$1.5 (heart,

highly resistant), and Na$_v$1.6 (brain/peripheral nerves, moderately resistant; figs. 3, S4-S9). Moderate

resistance in the muscle channel Na$_v$1.4 appeared between 75–130 mya, after the divergence of

Ambystomatidae and Salamandridae (the family consisting of true salamanders and newts; Hanifin and

Gilly 2015). This gain in muscle resistance coincided with the appearance of two highly resistant

substitutions in DI of Na$_v$1.5, which are present in all Salamandridae. Extreme TTX resistance across all

Na$_v$ paralogs evolved more recently, occurring approximately 30 mya, after the split between primitive

newts, which include *Pleurodeles*, and modern newts (Hime et al. 2021), which include all TTX-bearing

species. Over this time period, TTX resistance evolved in Na$_v$1.2 and Na$_v$1.3, and multiple additional

resistant mutations appeared and became fixed in Na$_v$1.1, Na$_v$1.4, and Na$_v$1.6.

Table 3. List of TTX Resistance-Conferring Substitutions Observed in Salamanders

Substitution[a]	Tetrodotoxic newts	Non-tetrodotoxic salamanders	Pufferfish and snakes[b]	Fold-change in TTX sensitivity	Resistance category	Citation
Y401C (DI)	Na$_v$1.1 Na$_v$1.2	Na$_v$1.5	Pufferfish: Na$_v$1.4a Na$_v$1.5La Na$_v$1.5Lb	~2500×	Extreme	Venkatesh et al. (2005); Jost et al. (2008)
Y401A (DI)	Na$_v$1.1 Na$_v$1.3 Na$_v$1.6	Na$_v$1.1	Pufferfish: Na$_v$1.1La Na$_v$1.6b	>600×	Extreme	Jost et al. (2008); Vaelli et al. (2020)
Y401S (DI)	Na$_v$1.5	Na$_v$1.5	-	~7000×	Extreme	Leffler et al. (2005)
D1532S (DIV) G1533D (DIV)	Na$_v$1.4		Snake: Na$_v$1.4	-	Extreme[c]	Feldman et al. (2012); Hanifin and Gilly (2015)
E758D (DII)	-	Na$_v$1.5	Pufferfish: Na$_v$1.4b	~3000×	Extreme	Bricelj et al. (2005)
M1240T (DIII)	Na$_v$1.1 Na$_v$1.4	Na$_v$1.1 Na$_v$1.5	Pufferfish: Na$_v$1.1La Na$_v$1.1Lb Na$_v$1.4a Na$_v$1.4b	~15×	Moderate	Jost et al. (2008)
V1233I (DIII)	Na$_v$1.6	Na$_v$1.6	Snake: Na$_v$1.6	~2×	Moderate	McGlothlin et al. (2014); Vaelli et al. (2020)
I1525V (DIV)	Na$_v$1.6	Na$_v$1.5 Na$_v$1.6	Snake: Na$_v$1.6	~2×	Moderate	Geffeney et al. (2005); McGlothlin et al. (2014); Vaelli et al. (2020)
I1525S (DIV)	Na$_v$1.4	Na$_v$1.4		-	Moderate[c]	Hanifin and Gilly (2015)
I1525T (DIV)	-	Na$_v$1.4 Na$_v$1.5		~7.7×	Moderate	Du et al. (2009)

[a] Amino acid sites are in reference to rat Na$_v$1.4 (accession number AAA41682)
[b] Parallel substitutions in Na$_v$ channels of TTX-bearing pufferfish and in snakes that consume TTX-bearing prey
[c] Resistance category determined from action potentials recorded from salamander muscle fibers

Fig. 3. Distribution of TTX-resistance conferring substitutions in salamander voltage-gated sodium channels. Toxic newt species are indicated with orange branches ("High TTX exposure"). Amino acids associated with Na$^+$ selectivity (400D, 755E, 1237K, 1529A) are shown in bold. Dots indicate identity with consensus sequences. Blue boxes indicate paralogs inferred to have moderate (~2-15-fold) resistance, orange boxes indicate paralogs with extreme (>300-fold) resistance, and grey boxes indicate paralogs without resistance or with insufficient sequence data. Substitutions known to confer TTX resistance are highlighted with respective colors. Extreme resistance in a paralog can result from the presence of one highly resistant substitution or the combination of multiple moderately resistant substitutions. Exon duplication has led to an additional TTX-binding domain (DIVb) in Na$_v$1.4 of all salamanders except *C. alleganiensis*. We did not identify the sequence encoding DIII of Na$_v$1.2 for any of the newts, however Vaelli et al. (2020) report that, in *T. granulosa*, this domain is identical in amino acid sequence to other salamanders. Amino acid sites are in reference to rat Na$_v$1.4 (accession number AAA41682).

Selective Regimes and Evolutionary Rates

In order to characterize the selective regimes acting on Na$_v$ genes, we used the codeml program in PAML

(Yang 2007) to fit models of selection to Na$_v$ codon alignments and compared nested models using

likelihood ratio tests (LRTs). We tested for site-specific positive selection within all amphibians by

comparing two sets of nested models: one set using a discrete distribution of ω values either with or

without positive selection (M1a vs. M2a; Yang et al. 2005); and another set fitting a continuous

distribution of ω values under purifying selection only, or adding categories of unconstrained evolution

and positive selection (M7 vs. M8 and M8a vs. M8; Yang et al. 2000). Parameter estimates and LRT

results for site models are summarized in table S3. To test for selection within toxic newts, we fit branch

(Yang 1998) and branch-site models (Zhang et al. 2005) to our datasets, which allowed ω to vary both

among codon sites and between toxic newts and other amphibians (summarized in table S4).

All models estimated relatively low ratios of nonsynonymous to synonymous substitution (d$_N$/d$_S$,

or ω ratios) for all Na$_v$ paralogs (average ω ratios from branch models ranged from 0.05 to 0.25),

indicating pervasive purifying selection. Based on LRTs comparing one ω-value models (which allow

only for one ω value across the entire phylogeny) to branch models (allowing for a different ω ratio in the

toxic newt clade relative to other amphibians), we found that ω ratios were significantly (P < 0.01) higher

in toxic newts for Na$_v$1.1, Na$_v$1.3, and Na$_v$1.4 and non-significant (P > 0.01) for Na$_v$1.2, Na$_v$1.5, and

Na$_v$1.6 (fig. 4A, table S4). The largest difference in ω ratios in the branch test was observed for the

muscle channel Na$_v$1.4 (newt ω = 0.23; all salamanders ω = 0.10), which appears to be due to both an

increase in the proportion of unconstrained sites as well as a larger number of estimated sites undergoing

positive selection (fig. 4B). However, the posterior probability support for positive selection at many of

these sites was low, and LRTs from branch-site models indicated significant evidence for a shift in

positive selection only in paralog Na$_v$1.3 (tables S3, S4).

Our site and branch-site models identified a number of TTX-binding sites with elevated ω ratios

(tables 4, S5, S6). Because of the small number of species in our study, we had low power to detect

statistically significant positive selection (Yang et al. 2000; Anisimova et al. 2001; Kosakovsky Pond and

Frost 2005), and the posterior probabilities provided low-to-moderate support for positive selection at

most sites. We also note that the false positive rates of site selection models can be inflated due to gene

conversion events (Casola and Hahn 2009), among-site variation in d$_S$ (Kosakovsky Pond and Frost

2005), and multinucleotide mutations (Venkat et al. 2018). However, results were consistent between the

M2a and M8 models, and the codons that were identified suggest that positive selection may have been

important for observed substitutions in TTX-binding regions.

Within the brain channels Na$_v$1.1, Na$_v$1.3, and Na$_v$1.6, putative positive selection was detected by

site models at site 401 within the DI P-loop, indicating selection acting across all salamanders rather than

specifically within toxic newts (table 4). Replacement of the aromatic amino acid at site 401 with a non-

aromatic amino acid can substantially impact TTX binding capacity (Leffler et al. 2005; Venkatesh et al.

2005; Vaelli et al. 2020), and we observed non-aromatic substitutions at this site within all brain channels

of highly toxic newts (fig. 3, table 3). The codon sequence for site 401 was variable across many

salamanders lacking TTX; however, almost all of the nonsynonymous changes observed outside of TTX-

bearing newts were biochemically conservative (both phenylalanine and tyrosine are aromatic and do not

affect TTX binding; Sunami et al. 2000), with the exception of the 401A observed in *P. cinereus* $Na_v1.1$.

This conservative variation likely contributes to the signal of diversifying selection acting on this codon

in less toxic salamander lineages. Site models also suggested positive selection acting on site 1533 in DIV

of $Na_v1.1$ and $Na_v1.2$ (table 4). Although the substitutions present at site 1533 in these newt paralogs have

not been tested experimentally for their effects on TTX binding, Maruta et al. (2008) showed that a

G1533T substitution at this site led to a moderate (~2-3-fold) decrease in TTX binding affinity, and

substitutions at this site are common in TTX-resistant channels (Geffeney et al. 2005; Jost et al. 2008;

Feldman et al. 2012; McGlothlin et al. 2016), suggesting that these non-synonymous changes likely also

reduce TTX binding affinity in newts.

266   Within the toxic newt lineage, we identified putative positive selection acting on three known

TTX-binding sites: 1240 and 1532 in DIII and DIV of muscle channel $Na_v1.4$ and 1529 in DIV of brain

channel $Na_v1.1$ (table 4). In $Na_v1.4$, sites 1240 and 1532 contain resistance-conferring substitutions

exclusively within toxic newts and these substitutions have been associated with extreme TTX resistance

in *Taricha* muscle fibers (Hanifin and Gilly 2015). In $Na_v1.1$, site 1529 encodes part of the Na^+ selectivity

filter (comprised of interacting amino acids DEKA – sites 400, 755, 1237, and 1529), which is highly

conserved across Na_v paralogs. An A1529G substitution (resulting in a DEKG filter) is present in $Na_v1.1$

of *A. mexicanum* and all members of the toxic newt clade. While this substitution does not appear to

affect Na^+ selectivity or to be sufficient in preventing TTX from binding (Jost et al. 2008), it may alter

channel firing properties, as this A1529G substitution resulted in substantially higher Na^+ currents relative

to the wild-type when introduced into a mammalian $Na_v1.4$ channel (Jost et al. 2008). The same alanine

to glycine substitution has been observed in Na_v channels of TTX-bearing flatworms (Jeziorski et al.

1997) and pufferfish (Jost et al. 2008), which suggests that it may play a role in TTX resistance in these

organisms.

280   Outside of TTX-binding regions, we found evidence for putative positive selection in similar

281   regions across multiple Na_v paralogs (tables S5, S6). For $Na_v1.4$, the majority of sites under positive

282   selection reside in terminal exon 26a that encodes the DIVa P-loop (exon 26b was excluded from this

283   analysis due to its absence in some species). For most other paralogs, the largest clusters of sites with

284   elevated d_N/d_S ratios were within the DI L5 turret (the extracellular loop upstream of the DI P-loop,

285   encoded by exons 6 and 7) and the DIII L5 turret upstream of the DIII P-loop encoded by exon 21.

Fig. 4. Tests for shifts in selective pressure on salamander voltage-gated sodium channels. (A) PAML branch models comparing estimates of gene-wide ω (d_N/d_S) within TTX-bearing newts (black circles) and other salamanders (grey circles). (B) PAML site and branch-site model estimates of proportions of sites under purifying selection ($0 < \omega < 0.05$ in both lineages), unconstrained evolution ($\omega = 1$ in both lineages), and positive selection or relaxed purifying selection ($0 < \omega < 0.05$ in salamanders and $\omega \geq 1$ in TTX-bearing newts). Significant differences based on likelihood ratio tests are indicated with * ($p < 0.05$) and ** ($p < 0.01$). Although likelihood-ratio tests were non-significant for Na$_V$1.1 and Na$_V$1.4 (tables S3, S4), PAML identified a large proportion of amino acid coding sites within these paralogs with elevated d_N/d_S ratios in toxic newts.

Table 4. P-loop Sites with Elevated ω Values in All Salamanders and in Toxic Newts.

	Na$_V$1.1	Na$_V$1.2	Na$_V$1.3	Na$_V$1.4	Na$_V$1.5	Na$_V$1.6
Tissue expression	Brain	Brain	Brain	Muscle	Heart	Brain / PNS
P-loop sites under positive selection in all salamanders[b,d]	Y401A/C (DI)[a] G1533V (DIV)[a]	G1533R (DIV)[a]	Y401A (DI)[a]	-	739 (DII)	Y401A (DI)[a]
P-loop sites under positive selection in toxic newts[c,d]	1224 (DIII) A1529G (DIV)[a]	-	T759S (DII)	W756Y (DII) M1240T (DIII)[a] 1517 (DIV) 1519 (DIV) D1532S (DIV)*,[a]	-	-

* Empirical Bayes posterior probability > 0.95
[a] Non-conservative amino acid substitutions associated with TTX resistance in toxic newts
[b] Sites identified with site models using empirical Bayes method (posterior probability > 0.50)
[c] Sites identified with branch-site models using empirical Bayes method (posterior probability > 0.50)
[d] Because we had low power to detect positive selection at specific sites due to the small number of species in our study, these results should be interpreted with caution

These sites may facilitate interaction with other proteins, or alternatively, some of these sites identified by

the branch-site model may be selected to compensate for biochemical changes produced by TTX-resistant

mutations.

Gene Conversion Events

We also tested for evidence of nonallelic gene conversion as a mechanism of sequence evolution

contributing to adaptive evolution in Na$_v$ genes using the program GENECONV. Nonallelic or ectopic

gene conversion results from an interlocus exchange of DNA that can occur between closely related

sequences during double-stranded break repair (Hansen et al. 2000). GENECONV uses the information in

a multiple sequence alignment to identify regions of similarity shared between two sequences that is

higher than expected by chance based on comparisons to permuted alignments (Sawyer 1999). We

selected this program to detect gene conversion because of its low false positive rates and robustness to

shifts in selective pressure (Posada and Crandall 2001; Bay and Bielawski 2011). Because nonallelic gene

conversion is more likely to occur between paralogs residing on the same chromosome (Semple and

Fig. 5. Gene conversion within TTX binding regions. (A) Gene conversion was detected between DIVa and DIVb P-loops within duplicate exons 26a and 26b of Na$_v$1.4 in *T. torosa*. Substitutions conferring TTX resistance are highlighted in yellow and blue. While one moderately resistant substitution is present in DIVa of less toxic salamanders (1525T/S), three identical substitutions are present in both DIVa and DIVb of highly toxic newts. (B) Gene conversion was also detected between the DI P-loops of paralogs Na$_v$1.1 and Na$_v$1.3 in both *T. torosa* and *T. granulosa*. In *Notophthalmus*, both paralogs have different resistant substitutions (401C in Na$_v$1.1 – highlighted in green; 401A in Na$_v$1.3 – highlighted in orange). This putative gene conversion may have homogenized TTX-resistant substitutions among these paralogs within *Taricha* newts. Site numbers are in reference to amino acids of rat Na$_v$1.4 (accession number AAA41682). Stars at branches indicate the inferred transfer of substitutions associated with moderate resistance (blue text) or extreme resistance (orange text) via non-allelic gene conversion.

Wolfe 1999; Drouin 2002; Benovoy and Drouin 2009), we limited our search to events between the

tandem duplicates Na$_v$1.1, Na$_v$1.2, and Na$_v$1.3 and between exons 26a and 26b of Na$_v$1.4 within each

salamander genome. While few gene conversion events were detected within each species by

GENECONV, all of the regions that were detected within *Taricha* newts contain TTX-binding sites with

substitutions associated with TTX resistance, including the DIVa and DIVb P-loops of Na$_v$1.4 (fig. 5,

table 5). We observed three TTX resistant amino acids (I1525S, D1532S, G1533D) within the DIVa and

DIVb P-loops of Na$_v$1.4 in toxic newt genomes. In contrast, *P. waltl*, a closely related but non-

tetrodotoxic newt, contains one moderately TTX-resistant amino acid (I1525S) in the DIVa P-loop and no

resistant amino acids in the DIVb P-loop (fig. 5A). These differences involve four identical nucleotide

changes at homologous sites in both exon duplicates. Our short reads from the genomes of *N. viridescens*

and *Taricha* newts mapped onto each of these exon assemblies across putative recombination break points

with high (> 50-fold) coverage, lending support for sequence convergence rather than an assembly error.

We did not detect gene conversion between these exons within the genomes of *T. granulosa* or *N.*

viridescens; however, this may be due to the low power of GENECONV to detect conversion (Bay and

Bielawski 2011), particularly in the presence of low sequence diversity and when the conversion tract is

Table 5. Nonallelic Gene Conversion Events

Na$_v$1.1 – Na$_v$1.2 – Na$_v$1.3 Coding Sequences

Species	Gene 1 – Gene 2	BC p-value[a]	Protein begin[b]	Protein end[b]	P-loop	Exon	Length (bp)	Total poly (bp)[c]	Num diffs (bp)[d]	Total diffs (bp)[e]
A. mexicanum	Na$_v$1.1 - Na$_v$1.2	0.01	12	42	-	1	93	48	5	794
	Na$_v$1.1 - Na$_v$1.3	0.01	45	84	-	1	128	54	5	722
	Na$_v$1.2 - Na$_v$1.3	0.00	203	405	DI	5; 6; 7; 8 19; 20;	>626	74	2	902
	Na$_v$1.1 - Na$_v$1.2	0.00	1059	1193	-	21	>407	134	25	794
	Na$_v$1.1 - Na$_v$1.3	0.02	1104	1144	-	20	127	37	1	722
	Na$_v$1.1 - Na$_v$1.2[f]	**0.02**	**1208**	**1232**	**DIII**	**21**	**84**	**35**	**2**	**794**
C. alleganiensis	Na$_v$1.1 - Na$_v$1.2	0.00	1111	1232	DIII	20;21	>376	153	23	621
	Na$_v$1.2 - Na$_v$1.3[f]	**0.02**	**1478**	**1611**	**DIV**	**26**	**406**	**164**	**32**	**693**
	Na$_v$1.1 - Na$_v$1.2	0.00	1531	1652	DIV	26	371	177	27	621
N. viridescens	Na$_v$1.1 - Na$_v$1.3	0.00	500	512	-	10	89	41	1	733
	Na$_v$1.1 - Na$_v$1.2	0.01	1104	1199	-	16	292	109	20	796
P. cinereus	Na$_v$1.1 - Na$_v$1.2	0.00	12	32	-	1	64	31	0	838
	Na$_v$1.1 - Na$_v$1.3	0.00	362	499	DI	8; 9; 10	>576	88	13	786
	Na$_v$1.2 - Na$_v$1.3	0.02	573	599	-	13	83	29	2	897
	Na$_v$1.1 - Na$_v$1.3	0.00	610	667	-	14	177	68	8	786
	Na$_v$1.1 - Na$_v$1.2	0.00	1135	1305	DIII	20; 21; 22; 23; 24	>524	138	9	838
	Na$_v$1.1 - Na$_v$1.3	0.03	1430	1473	-	26	134	46	5	786
P. waltl	Na$_v$1.1 - Na$_v$1.3	0.03	1369	1386	-	25	56	30	0	738
T. granulosa	**Na$_v$1.1 - Na$_v$1.3[g]**	**0.01**	**374**	**408**	**DI**	**8**	**105**	**41**	**2**	**725**
T. torosa	**Na$_v$1.1 - Na$_v$1.3[g]**	**0.01**	**374**	**408**	**DI**	**8**	**105**	**41**	**2**	**725**

Na$_v$1.4 Exon 26

Species	Gene	BC p-value[a]	Protein begin[b]	Protein end[b]	P-loop	Exon	Length (bp)	Total poly (bp)[c]	Num diffs (bp)[d]	Total diffs (bp)[e]
P. cinereus	Na$_v$1.4	0.01	1509	1534	DIV	26a-26b	77	26	0	154
T. torosa	**Na$_v$1.4[g]**	**0.02**	**1518**	**1536**	**DIV**	**26a-26b**	**56**	**22**	**1**	**178**

[a] Global Bonferroni-corrected (BC) p-values from 10,000 simulations
[b] Protein begin and end sites are in reference to the rat Na$_v$1.4 channel (accession number AAA41682)
[c] Polymorphism within the converted region across all species
[d] Pairwise differences between the two paralogs within the converted region
[e] Total pairwise differences across the entire length of the two paralogs
[f] Putative gene conversion events associated with the loss of TTX resistance
[g] Putative gene conversion events associated with the gain of TTX resistance

shorter than ~100 bp (Posada and Crandall 2001; McGrath et al. 2009). Together, these results suggest

that the three resistant amino acids accumulated together in one exon copy followed by conversion of the

other exon ~30 mya in a toxic newt ancestor.

We also detected gene conversion between the DI P-loops of paralogs $Na_v1.1$ and $Na_v1.3$ within

the genomes of both *Taricha* species (fig. 5B, table 5). TTX-resistant substitutions are identical in the DI

P-loops of $Na_v1.1$ and $Na_v1.3$ in both *Taricha* species (401A, encoded by a GCT codon), while $Na_v1.1$ of

N. viridescens contains a different codon at this position (TGC, encoding 401C), consistent with gene

conversion occurring at this locus in *Taricha.* Both $Na_v1.1$ and $Na_v1.3$ of non-tetrodotoxic *P. waltl* newts

encode a TTX-sensitive tyrosine at this locus. It is unclear whether putative gene conversion event(s)

occurred before or after resistance evolved in both paralogs. Gene conversion may have converted a non-

resistant channel to a resistant channel in an ancestral *Taricha*, while *N. viridescens* independently

acquired a Y401A substitution, or it may have homogenized substitutions within two channels that had

previously evolved resistance in an ancestor of all toxic newts. In *Taricha* newts, the transition from

either tyrosine or cysteine to alanine required multiple nucleotide substitutions in both paralogs, making

gene conversion a likely explanation for the observed substitution patterns.

We detected additional gene conversion events that may have involved a non-resistant paralog

acting as a donor to a resistant paralog, leading to the loss of TTX resistance in *A. mexicanum* and *C.*

alleganiensis paralogs. A resistant substitution is present in the DIII P-loop of $Na_v1.1$ in most

salamanders but is absent in $Na_v1.1$ of *A. mexicanum* (fig. 3), and we detected gene conversion in an

adjacent region between $Na_v1.1$ and the non-resistant $Na_v1.2$ paralog within the *A. mexicanum* genome

(table 5, fig. S10A). Similarly, *C. alleganiensis* $Na_v1.2$ contains a non-resistant 1533G within DIV of all

six paralogs, while $Na_v1.2$ of *X. tropicalis* frogs encodes a putatively TTX-resistant 1533L, and gene

conversion between paralogs $Na_v1.2$ and $Na_v1.3$ may have facilitated the loss of this substitution in

salamanders (fig. S10B)

Discussion

Here we show that TTX-bearing newts have evolved resistance to their own toxicity through

multiple parallel changes in their Na_v genes and that, similar to snakes that consume TTX-containing prey

(McGlothlin et al. 2016; Perry et al. 2018), some of the resistance in this taxon is ancient, first appearing

in an early salamander. However, while substitutions conferring moderate TTX resistance are present in

some non-toxic salamander genomes, only the TTX-bearing newts have substitutions conferring high

resistance across all six of their Na_v paralogs, and several of these channels harbor multiple resistant

substitutions in more than one domain. Many of the substitutions conferring resistance to toxic newts are

also present in Na_v paralogs of TTX-resistant pufferfish (Jost et al. 2008) and snakes (McGlothlin et al.

2016). Similar to pufferfish, newts appear to require resistance within all of their brain/nerve channels in

addition to their hearts and muscles. This feature apparently distinguishes toxic prey from their predators,

whose brain channels lack resistant substitutions (McGlothlin et al. 2014). This molecular parallelism

emphasizes the strong structural and functional constraints on this gene family, which appear to limit the

evolution of TTX resistance to a small number of predictable pathways, leading to convergent and

parallel changes across multiple taxa (Feldman et al. 2012). We show that the evolution of extreme TTX

resistance is accompanied by a shift in signatures of selection in four out of six paralogs, with suggestive

evidence of positive selection acting directly on TTX-binding sites. Finally, the evolution of physiological

resistance appears to have been facilitated by at least two instances of non-allelic gene conversion, which

acted to introduce TTX-resistance substitutions from one paralog (or exon duplicate) to another,

providing a rare example of the facilitation of adaptive evolution via non-allelic gene conversion.

Our reconstruction of the history of TTX resistance in salamanders reveals the ancient origins of

moderate resistance in nerve channels $Na_v1.1$ and $Na_v1.6$ and high resistance in the heart channel $Na_v1.5$,

which arose ~160 mya. Resistant substitutions in the muscle channel $Na_v1.4$ evolved later, becoming

fixed in the clade including all newts and *S. salamandra* between 75–130 mya. This was followed by the

accumulation of additional substitutions in DIII, DIVa, and DIVb of $Na_v1.4$ within members of the highly

toxic newt clade, providing their muscles with resistance to much higher concentrations of TTX relative

to salamanders lacking these mutations (Hanifin and Gilly 2015). Substitutions in DI conferring extreme

TTX resistance to the brain/nerve channels $Na_v1.1$, $Na_v1.2$, $Na_v1.3$, and $Na_v1.6$ also evolved more

recently (~30 mya) and they are limited to the toxic newt clade with only one exception, $Na_v1.1$ of *P.*

cinereus, which may have arisen independently in this lineage. Toxic newts also have unique

substitutions in DIV of $Na_v1.1$ and $Na_v1.2$, which may provide additional resistance to their brain

channels or may compensate for structural or functional changes resulting from resistant substitutions in

DI or in other regions of the protein. The widespread presence of TTX in modern newts suggests that

possession of TTX evolved in the common ancestor of this clade (Hanifin 2010). This is supported by our

observation of shared TTX-resistance substitutions across all Na_v paralogs in highly toxic newts.

However, the highly toxic newts included in this study are limited to North American species and do not

include Asian or European newts (e.g. *Cynops* and *Triturus* spp.) known to have TTX (Hanifin 2010).

Sequencing the Na_v genes of these species will reveal if extreme resistance in the nerve channels is

specific to North American newts or arose earlier in an ancestral newt species.

Whether the TTX-resistance substitutions observed in newt relatives are adaptive and what

selective pressures act to retain them remains to be determined. In ancestral salamandrids, resistance in

three or four of the six Na_v proteins may have provided tolerance to low levels of TTX, facilitating the

evolution of extreme toxicity in the modern newts, although it is unclear whether these substitutions arose

in response to TTX exposure or as a side effect of selection for another aspect of channel function.

Environmental exposure to TTX could potentially occur from TTX-bearing prey, such as terrestrial

Bipalium flatworms (Stokes et al. 2014) or from TTX-producing bacteria (Vaelli et al. 2020). In a feeding

study on *B. adventitium* flatworms, Ducey et al. (1999) demonstrated that while all salamanders rejected

the flatworms when they were first presented, some habituated *Ambystoma* and *Plethodon* individuals

were able to consume them with only minor symptoms of TTX poisoning, including apparent mucus

production and numbing of the mouth. Although the amount of TTX present in the worms was not

measured, this observation supports the conjecture that these substitutions play a protective role against

consuming toxic prey.

Our power to detect sites undergoing positive selection was low due to the small number of

species available for our analyses and the high sequence conservation among orthologs (Yang et al. 2000;

Anisimova et al. 2001). However, our analyses provided evidence consistent with positive selection

acting on TTX-binding sites, including sites implicated in the extreme TTX resistance of salamander

muscle fibers. Our branch models indicate that several Na_v paralogs have higher ω ratios within the

tetrodotoxic newt clade relative to other salamanders. This increase in ω ratio appears to be mainly the

result of relaxed purifying selection rather than positive selection, as few sites were estimated to have $\omega >$

1 by branch-site models. Muscle channel Na_v1.4 had the highest ω ratio in toxic newts, coinciding with a

relatively high proportion (~0.06) of sites detected as undergoing positive selection in toxic newts and

purifying selection in other salamanders, although the likelihood ratio test was non-significant (table S4).

This pattern may derive from ongoing positive selection on Na_v1.4 resulting from the coevolutionary

arms race between newts and snakes. Increased ω ratios in toxic newts coincide with the appearance of

highly TTX-resistant substitutions within the DI P-loops of Na_v1.1, Na_v1.3, and Na_v1.6, DIV of Na_v1.1,

and the DIII, DIVa, and DIVb P-loops in Na_v1.4. While the pattern of these substitutions suggests that

they are adaptive changes that occurred specifically within the tetrodotoxic newt clade, our branch-site

models detected positive selection on TTX-binding sites only within the DIV P-loop of Na_v1.1 and the

DIII and DIVa P-loops of Na_v1.4, while our site models detected positive selection on site 401 within

paralogs Na_v1.1, Na_v1.3, and Na_v1.6 of all salamanders. This may be due to the statistically conservative

nature of the branch-site model and the tendency of PAML to detect ongoing diversifying selection rather

than rare positive selection events (Yang and dos Reis 2010). Furthermore, the site and branch-site

models do not distinguish between biochemically conservative and non-conservative amino acid changes.

Nevertheless, the detection of relatively high ω ratios within the toxic newt clade along with site-specific

positive selection at known TTX-binding sites provides strong evidence that these substitutions are

adaptive.

In addition to the six Na$_v$ paralogs described in amphibians (Zakon 2012), our sequence data

revealed the presence of a partial duplication of the Na$_v$1.4 gene in salamanders that includes the entirety

of exon 26, encoding the DIV P-loop, which likely occurred in an ancestor of Salamandroidea. The

maintained open reading frame and shared expression patterns of transcripts encoding exons 26a and 26b

in *A. mexicanum* tissues suggests that this duplicate region is functional in salamander muscles. In some

insects that feed on toxic plants, the appearance of resistance-conferring substitutions is accompanied by

one or more duplications of the genes that the toxin targets (Zhen et al. 2012; Petschenka et al. 2017).

Petschenka et al. (2017) show that in at least one species, such gene duplication precedes resistance, and

they suggest that gene duplication may help to alleviate the potential decrease in fitness incurred by the

insects due to the negative pleiotropic effects of toxin-resistant mutations. Similarly, resistance in DIV of

salamander Na$_v$1.4 appears after the duplication of this domain, and only the TTX-bearing newts have

resistant substitutions in both exon copies, lending support to this hypothesis and raising the possibility

that the evolution of physiological resistance in salamanders may have been mediated in part by this

genomic novelty.

We observed a rare case of the generation of adaptive variants via non-allelic gene conversion,

which occurred both between the duplicated exons of Na$_v$1.4 and between paralogs Na$_v$1.1 and Na$_v$1.3.

Gene conversion is often thought to play a role in constraint, preserving the core functions of gene

families (Chapman et al. 2006) and reducing deleterious mutation loads (Ohta 1989; Khakhlova and Bock

2006), and has also been associated with the diversification of major histocompatibility complex genes in

mammals (Kuhner et al. 1991; Go et al. 2003) and the introduction of deleterious nonsynonymous

mutations into different parts of the genome (Galtier et al. 2009; Casola et al. 2012). However, the

potential for gene conversion to facilitate adaptation is less widely appreciated. Theory suggests that gene

duplication and subsequent gene conversion may allow for movement between adaptive peaks via the

accumulation of beneficial mutations in one gene copy that can be transferred to a favorable genetic

background (Hansen et al. 2000). This process has been implicated in the adaptive evolution of hypoxia

tolerance in high-altitude Tibetan wolves (Signore et al. 2019) and in heavy metal tolerance in

Arabidopsis (Hanikenne et al. 2013). We observed a rare case of the generation of adaptive variants via

non-allelic gene conversion, which occurred both between the duplicated exons of Na$_v$1.4 and between

paralogs Na$_v$1.1 and Na$_v$1.3. In the brain channel genes Na$_v$1.1 and Na$_v$1.3, this likely occurred between

two genes that had previously evolved resistance, homogenizing the substitutions between the two

paralogs in *Taricha* newts. In the muscle channel gene Na$_v$1.4, three identical amino acids were present

within both duplicated DIV TTX-binding domains in toxic newts, while only one resistant amino acid

was present in a single DIV domain in their non-tetrodotoxic salamander relatives, suggesting that

resistant substitutions accumulated in one exon copy in a toxic newt ancestor and subsequently spread to

the other exon copy via gene conversion. Both Hanifin and Gilly (2015) and Du et al. (2009) have shown

that the single resistant amino acid observed in less toxic salamanders confers only low levels of TTX

resistance, while the combination of the three amino acids found in highly toxic newts confers extreme

resistance. As splice variants encoding the alternative exons appear to share similar expression patterns in

salamanders, both exon copies should require extreme resistance in species exposed to high TTX

concentrations. The concerted evolution of TTX resistance among homologous P-loop domains of $Na_v1.4$

may have expedited the evolution of extreme resistance in newt muscles, requiring new resistant

mutations to appear in only one of these domains before being transferred to the other copy.

The degree of parallel molecular evolution among members of the Na_v gene family and across

lineages provides insight into the constraints on Na_v nucleotide sequence as well as the evolvability of the

TTX-resistance phenotype. Our results reveal that similar to their coevolutionary partners, *Thamnophis*

garter snakes, *Taricha* newts evolved extreme TTX resistance through a stepwise process that built upon

ancient changes that were in place millions of years before the arms race began. However, the pattern of

TTX resistance evolution in newts also displays important differences from that of their predators. First,

perhaps because of the constitutive presence of TTX, newts display extreme levels of resistance even in

channels that are expressed in the central nervous system, which are protected by the blood-brain barrier

in species that encounter TTX in their diet. Second, our analysis indicates that many substitutions may

have become fixed relatively close to one another in evolutionary time within the clade of modern newts.

This is in contrast to snakes, where key changes were separated by millions of years. Due to our lack of

sampling of newt species outside of North America, however, further work is necessary to understand the

timing of these changes on a finer scale. We also show that while positive selection appears to be a strong

driving force of the evolution of TTX auto-resistance in newts, gene conversion may have sped up the

process of adaptive evolution in some Na_v paralogs, and constraints have limited the possible locations

and types of resistant substitutions to a small subset of realized genetic changes. Taken together, our

results emphasize the interplay among selection, constraint, and historical contingency in the evolution of

complex adaptations.

Methods

Sequencing and Annotation of Voltage-Gated Sodium Channel Paralogs

We identified Na_v genes in the two publicly available salamander genome assemblies, *A. mexicanum*

(Smith et al. 2019; AmexG.v6 assembly) and *P. waltl* (Elewa et al. 2017), and in one full-body

transcriptome from the fire salamander *S. salamandra* (Goedbloed et al. 2017; BioProject accession

PRJNA607429) using the reciprocal best BLAST hit method (Moreno-Hagelsieb and Latimer 2007) with

queries of Na_v sequences from *X. tropicalis* (Hellsten et al. 2010) and salamanders (Hanifin and Gilly

2015). We confirmed assignments of each amphibian Na_v paralog based on nucleotide alignments of the

coding sequences with *Xenopus* sequences, as well as synteny of the chromosomal segments containing

Na_v genes (fig. S1). These Na_v annotations were then used to design targeted sequencing probes and to

subsequently assign paralog identity to our de novo salamander assemblies. We used Geneious v10.2.3

for sequence visualization and to create DNA and protein alignments (Kearse et al. 2012). We also

performed BLASTn searches of published transcriptome assemblies of *Tylototriton wenxianensis*

(PRJNA323392), *Bolitoglossa vallecula* (PRJNA419601), and *Hynobius retardatus* (Matsunami et al.

2015; PRJDB2409). However, we were unable to identify the full set of Na_v paralogs within these three

assemblies, possibly due to their tissue specificity. Therefore, we used the sequences for probe design and

to identify exon duplications in $Na_v1.4$ but excluded them from PAML analyses.

In order to design targeted sequencing probes, we compiled partial and complete amphibian Na_v

sequences obtained from NCBI databases and additional published sources (table S7) into a single

FASTA file. Each individual FASTA entry included a single exon and, when available, up to 200 bp of

intron sequence upstream and downstream of each exon to aid in paralog assignment. Using

RepeatMasker (Smit et al. 2013-2015), we replaced transposable elements and other sequence repeats

with Ns, and subsequently filtered out sequences < 120 bp in length, as well as those with more than 25%

missing or ambiguous characters. We submitted this masked and filtered file, which was 465 kb in total

length, to Agilent Technologies (Santa Clara, CA, USA) for custom probe design using the SureSelect

tool, resulting in 7518 unique 120-mer probes. Probe sequences are available in our Dryad repository.

We obtained DNA samples from adult individuals of three TTX-bearing species (n=3 for each

species): *T. torosa* from Hopland, CA, *T. granulosa* from Benton, OR, and *N. viridescens* from Mountain

Lake, VA, and from two additional salamander species presumed to lack TTX ($n = 2$ for each species): *P.*

cinereus collected in Mountain Lake, VA and *C. alleganiensis* collected in southwestern VA. We

extracted genomic DNA using the DNeasy Blood & Tissue kit (Qiagen Inc., Valencia, CA) and prepared

sequencing libraries using the SureSelect[XT] Target Enrichment system for Illumina paired-end

multiplexed sequencing from Agilent Technologies (Santa Clara, CA, USA), following the protocol for

low input (200 ng) DNA samples. We used a Covaris M220 Focused-ultrasonicator to shear 200 ng of

purified whole genomic DNA from each sample into ~250 bp fragments using the following settings:

duty factor 10%, peak incident power 75 w, 200 cycles per burst, and treatment time of 160 seconds. We

followed the Agilent SureSelect[XT] Target Enrichment kit protocol for end repair, adaptor ligation,

amplification, hybridization and bead capture (using the custom SureSelect probes described above),

indexing, and purification. We quantified the resulting enriched, indexed libraries with qPCR and

combined them in equimolar concentrations into one final library pool, which was submitted to the

Genomics Sequencing Center at Virginia Tech for sequencing on an Illumina MiSeq 300-cycle v2 with

150 bp paired-end reads. Prior to alignment, we trimmed Illumina reads of TruSeq3 adapter sequences,

removed bases with a phred64 quality score less than 3, and filtered out subsequent reads shorter than 100

bp using Trimmomatic version 0.33 (Bolger et al. 2014). We used SPAdes 3.6.0 (Bankevich et al. 2012)

to create de novo assemblies of the trimmed and filtered reads. Each Na_v paralog had > 10-fold sequence

coverage, with an average of 32-fold coverage across all species and paralogs (table S1). After filtering,

all reads had Phred Q-scores >30 for >99% of sites.

Because we designed the sequencing probes to capture only small portions of the Na_v intronic

regions flanking exons (regions more likely to be conserved across species), each exon was assembled

into a separate scaffold. For each individual from our sequencing trial, we created BLAST databases from

the assembled de novo scaffolds and performed BLAST searches using single exons from the *Ambystoma*

Na_v genes as queries. We created 26 separate nucleotide alignments, one for each individual Na_v exon,

including sequences from *Ambystoma*, *Pleurodeles*, and our de novo assemblies using MAFFT v 7.450

(Katoh and Standley 2013) and created consensus neighbor-joining trees with a Tamura-Nei genetic

distance model using the Geneious Tree Builder (Kearse et al. 2012) with 1000 bootstrap replicates and

an 80% support threshold. The resulting tree topologies were used to assign paralog identity to each of the

exons. When necessary, we included exon and intron sequences from additional species in the alignments

to resolve the topology of the trees.

We concatenated all exons from each paralog into full coding sequences. Based on alignment

with full-length *Xenopus* sequences, all salamander Na_v coding sequences collected for this study were

>90% complete with the exception of two sequences from the *Salamandra salamandra* transcriptome

(paralogs $Na_v1.1$ and $Na_v1.2$, which were 68.8% and 70.7% complete respectively; table S1). For each

species, we used Na_v paralogs sequenced from the genome of a single individual for our downstream

analyses, based on completeness of the assembly.

Determination of TTX Resistance Levels

TTX sensitivity is commonly measured *in vitro* by using site-directed mutagenesis to introduce mutations

of interest into a TTX-sensitive Na_v channel, followed by expression in *Xenopus* oocyte or HEK 293 cells

and the application of patch-clamp whole-cell recordings to measure channel current in the presence of

TTX. The fold-change in TTX sensitivity is then calculated by taking the ratio of the IC_{50} values, or the

TTX concentration at which 50% of the Na_v channels are blocked, of mutated and wild-type channels (see

table 3 for references).

Another line of evidence for resistance in salamander muscle channels ($Na_v1.4$) comes from

Hanifin and Gilly (2015), who estimated TTX sensitivity by recording action potentials generated from

salamander muscle fibers and estimating the amount of TTX required to diminish the rise of the action

potential, associating these relative changes with the presence and absence of substitutions in TTX-

binding sites. They associated moderate TTX resistance (reduced sensitivity to 0.010 μM TTX) with the

presence of DIII substitution M1240T and extreme resistance (low sensitivity to 300 μM TTX) with the

presence of DIII and DIV substitutions M1240T, D1532S, and G1533D. We categorize levels of TTX

resistance conferred by Na_v substitutions as extreme or moderate based on the results of Hanifin and Gilly

(2015) as well as the data summarized in table 3.

Phylogenetic Analyses and Identification of Site-Specific Evolutionary Rates

We constructed phylogenetic trees for the entire Na_v gene family using our *de novo* assembled sequences

as well as sequences from the genomes of *A. mexicanum*, *P. waltl*, the whole-body transcriptome of *S.*

salamandra, two frog genomes: *X. tropicalis* (Hellsten et al. 2010) and *Nanorana parkeri* (Sun et al.

2015), and one fish genome: *Danio rerio* (Howe et al. 2013). We created an amino acid alignment of

translated coding sequences using MAFFT v 7.450 (Katoh and Standley 2013) and constructed maximum

likelihood trees from these alignments with RAxML v8.2.11 (Stamatakis 2014) in Geneious using a

GAMMA BLOSSUM62 protein model and estimated clade support with 100 bootstrap replicates. In

order to improve the accuracy of the nucleotide alignment, we used this amino acid alignments to guide

codon alignments with PAL2NAL v14 (Suyama et al. 2006). We identified the best fitting substitution

models for the nucleotide alignment using jModelTest 2.1.10 v20160303 (Darriba et al. 2012) and

constructed maximum likelihood trees of the coding sequences with RAxML v8.2.11. The two models

with the lowest AIC scores were (1) the transition model with unequal base frequencies, a gamma shape

parameter, and some proportion of invariable sites (TIM2+I+G) and (2) a general time-reversible model

with a gamma shape parameter and a proportion of invariable sites (GTR+I+G), which is nearly identical

to TIM2+I+G, but includes two additional rate parameters (Posada 2008). Because the two models are

nearly equivalent and both fit our data equally well, we chose the GTR+I+G model for its ease of

implementation across different programs. We repeated these methods for each individual Na_v paralog for

downstream analyses in PAML, excluding the sequences from the outgroup species *N. parkeri* and *D.*

rerio. We estimated chronograms in Figs. 3 and 5 using the chronos( ) function of the ape v5.4-1 package

in R using a phylogenetic tree from Pyron (2014) and divergence dates from Hime et al. (2021; see Dryad

repository for R code).

579   We estimated synonymous (d_S) and nonsynonymous (d_N) substitution rates, as well as the d_N/d_S

ratios (ω) for each paralog using codeml in PAML v4.8 (Yang 2007). In order to test for changes in

selective regimes in the Na_v genes between salamanders and highly toxic newts (*Notophthalmus* and

Taricha species), we fit the following models to our Na_v alignments: (1) one-ratio models (allowing for a

single ω ratio among all sites and all branches of the phylogeny), (2) branch models (allowing for separate

ω ratios for the foreground [toxic newts] and background [other salamanders]), (3) branch-site neutral

models (allowing ω to vary both between toxic newts and other salamanders and among sites, with two

possible categories: $0 < \omega < 1$ and $\omega = 1$), and (4) branch-site models (allowing ω to vary both between

toxic newts and other salamanders and among sites, with three possible categories: $0 < \omega < 1$, $\omega = 1$, and

$\omega > 1$, the latter being allowed only within toxic newts). To test for site-specific positive selection among

all salamanders, we fit two sets of nested models: (1) discrete ω ratio models M1a vs. M2a, which

allowing for ω to vary among sites, with either two possible categories: $0 < \omega < 1$ and $\omega = 1$ or three

possible categories: $0 < \omega < 1$, $\omega = 1$, and $\omega > 1$; and (2) continuous ω ratio models M7, M8a, and M8,

which fit ω ratios of sites into a beta distribution, with ncatG (the number of ω values in the beta

distribution) set to 5. We used F3x4 codon models, which estimate individual nucleotide frequencies for

each of the three codon positions, and allowed codeml to estimate ω and transition-transversion rates (κ).

The outputs of these models were used to estimate and compare gene-wide ω between toxic newts and

salamanders (one-ratio and branch models), to identify sites under positive selection in all salamanders

(site and neutral site models), and to identify sites under positive selection or with elevated ω in toxic

newts relative to other salamanders (branch-site neutral and branch-site models). We also created

ancestral sequence reconstructions by specifying RateAncestor = 1 within codeml configuration files. We

report posterior probability estimates for ancestral sequence reconstructions using models with the highest

likelihood scores (Table S3), which were the neutral site models (M8a) for all Na$_v$ paralogs except

Na$_v$1.3, for which the site selection model M8 had the most support (Fig. S4-S9). We performed all of the

above analyses both including and excluding the two *S. salamandra* paralogs with a large number of gaps

(Na$_v$1.1 and Na$_v$1.2). Maximum likelihood parameter estimates were largely congruent using these

different datasets. Therefore, we present results from the gene trees including *S. salamandra* here.

Detection of Gene Conversion

We used the program GENECONV to detect potential nonallelic gene conversion events between Na$_v$

paralogs. We used a full codon alignment of all six Na$_v$ paralogs from 7 of the 8 salamanders included in

the study (excluding *S. salamandra* due to missing data), but targeted our search to include only genes

Na$_v$1.1, Na$_v$1.2, and Na$_v$1.3 within each species under the assumption that gene conversion is more likely

to occur between closely related sequences that reside on the same chromosome. This increased the power

of detecting gene conversion from multiple pairwise comparisons. We also performed a separate search

for gene conversion events between duplicate exons 26a and 26b of Na$_v$1.4. For this analysis, we used a

codon alignment of Na$_v$1.4 exons 26a and 26b from 10 salamander species. We assigned a mismatch

penalty using gscale=1 and used corrected *p*-values from 10,000 permutations to determine significance.

Acknowledgments

We thank Miguel Vences for helpful feedback on the manuscript and for providing an early draft of the *Salamandra*

salamandra transcriptome. We also thank Vincent Farallo, Edmund Brodie III, William Hopkins, and Brian Case for

providing DNA samples for this study, Kaitlyn Malewicz, Mercedes Collins, and Emily Orr for assistance with lab
work, John Abramyan, Edmund Brodie III, Charles Hanifin, and Andrew Kern for helpful conversations and
Matthew Hahn for advice on analyses and comments on the manuscript. Hellbender (*Cryptobranchus*) samples were
collected by Brian Case and William Hopkins in cooperation with the Virginia Department of Wildlife Resources
(DWR) and the U.S. Forest Service, and with assistance from J.D. Kleopfer. Hellbenders were handled under
scientific collecting permit #060465 and approved Virginia Tech IACUC procedures (protocol #16-162). We thank
the Departments of Fish and Wildlife in Oregon and California and the Virginia DWR for scientific collecting
permits to MTJH (063-18, SC-11937 and 048104 respectively). This work was supported by awards from the
National Science Foundation to JWM (DEB-1457463 and IOS-1755055), and to MTJH (DEB-1601296), by an R.C.
Lewontin award from the Society for the Study of Evolution to KLG, and by a graduate fellowship award from the
Interfaces of Global Change program at Virginia Tech to KLG.

Data Availability

633   Newly generated sequences have been submitted to the SRA database in NCBI under BioProject accession
634   PRJNA732671. Sequence alignments and configuration files for GENECONV analyses have been deposited at
635   https://github.com/kerrygendreau/Newt-TTX-Resistance.git and archived in a Dryad repository at
636   http://doi.org/10.5061/dryad.kprr4xh4v.

References

Anisimova M, Bielawski JP, Yang Z. 2001. Accuracy and power of the likelihood ratio test in detecting
adaptive molecular evolution. *Mol Biol Evol* 18(8):1585-1592.

Arbuckle K, Rodríguez de la Vega RC, Casewell NR. 2017. Coevolution takes the sting out of it:
Evolutionary biology and mechanisms of toxin resistance in animals. *Toxicon* 140:118-131.

Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI,
Pham S, Prjbelski AD, et al. 2012. SPAdes: A new genome assembly algorithm and its applications to
single-cell sequencing. *J Comput Biol* 19(5):455-477.

Bay RA, Bielawski JP. 2011. Recombination detection under evolutionary scenarios relevant to
functional divergence. *J Mol Evol* 73(5-6):273-286.

Benovoy D, Drouin G. 2009. Ectopic gene conversions in the human genome. *Genomics* 93(1):27-32.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics 30(15):2114-2120.

Bricelj VM, Connell L, Konoki K, MacQuarrie SP, Scheuer T, Catterall WA, Trainer VL. 2005. Sodium
channel mutation leading to saxitoxin resistance in clams increases risk of PSP. *Nature* 434(7034):763-
767.

Brodie ED, III, Brodie ED, Jr. 1990. Tetrodotoxin resistance in garter snakes: an evolutionary response of
predators to dangerous prey. *Evolution* 44(3):651-659.

Brodie ED, Jr., Ridenhour BJ, Brodie ED, III. 2002. The evolutionary response of predators to dangerous
prey: hotspots and coldspots in the geographic mosaic of coevolution between garter snakes and newts.
Evolution 56(10):2067-2082.

Bryant DM, Johnson K, DiTommaso T, Tickle T, Couger MB, Payzin-Dogru D, Lee TJ, Leigh ND, Kuo
T-H, Davis FG, et al. 2017. A tissue-mapped axolotl de novo transcriptome enables identification of limb
regeneration factors. *Cell Rep* 18(3):762-776.

Caballero-Pérez J, Espinal-Centeno A, Falcon F, García-Ortega LF, Curiel-Quesada E, Cruz-Hernández
A, Bako L, Chen X, Martínez O, Alberto Arteaga-Vázquez M, et al. 2018. Transcriptional landscapes of
Axolotl (*Ambystoma mexicanum*). *Dev Biol* 433(2):227-239.

Casola C, Hahn MW. 2009. Gene conversion among paralogs results in moderate false detection of
positive selection using likelihood methods. *J Mol Evol* 68(6):679-687.

Casola C, Zekonyte U, Phillips AD, Cooper DN, Hahn MW. 2012. Interlocus gene conversion events
introduce deleterious mutations into at least 1% of human genes associated with inherited disease.
Genome Research 22(3):429-435.

Chapman BA, Bowers JE, Feltus FA, Paterson AH. 2006. Buffering of crucial functions by paleologous
duplicated genes may contribute cyclicality to angiosperm genome duplication. *P Natl Acad Sci USA*
103(8):2730.

Chen J-M, Cooper DN, Chuzhanova N, Férec C, Patrinos GP. 2007. Gene conversion: mechanisms,
evolution and human disease. 8(10):762-775.

Darriba D, Taboada GL, Doallo R, Posada D. 2012. jModelTest 2: more models, new heuristics and
parallel computing. *Nat Methods* 9(8):772-772.

Drouin G. 2002. Characterization of the gene conversions between the multigene family members of the
yeast genome. *J Mol Evol* 55(1):14-23.

Du Y, Nomura Y, Liu Z, Huang ZY, Dong K. 2009. Functional expression of an arachnid sodium channel
reveals residues responsible for tetrodotoxin resistance in invertebrate sodium channels. *Journal of
Biological Chemistry* 284(49):33869-33875.

Ducey PK, Messere M, Lapoint K, Noce S. 1999. Lumbricid prey and potential herpetofaunal predators
of the invading terrestrial flatworm Bipalium adventitium (Turbellaria: Tricladida: Terricola). *Am Midl
Nat* 141(2):305-314.

Elewa A, Wang H, Talavera-López C, Joven A, Brito G, Kumar A, Hameed LS, Penrad-Mobayed M,
Yao Z, Zamani N, et al. 2017. Reading and editing the Pleurodeles waltl genome reveals novel features of
tetrapod regeneration. *Nat Commun* 8(1):2286.

Feldman CR, Brodie ED, Jr., Brodie ED, III, Pfrender ME. 2012. Constraint shapes convergence in
tetrodotoxin-resistant sodium channels of snakes. *Proc Natl Acad Sci USA* 109(12):4556-4561.

Fux JE, Mehta A, Moffat J, Spafford JD. 2018. Eukaryotic voltage-gated sodium channels: on their
origins, asymmetries, losses, diversification and adaptations. *Front Physiol* 9:1406.

Galtier N, Duret L, Glémin S, Ranwez V. 2009. GC-biased gene conversion promotes the fixation of
deleterious amino acid changes in primates. *Trends in Genetics* 25(1):1-5.

Geffeney SL, Brodie ED, Jr., Ruben PC, Brodie ED, III. 2002. Mechanisms of adaptation in a predator-
prey arms race: TTX-resistant sodium channels. *Science* 297(5585):1336-1339.

Geffeney SL, Fujimoto E, Brodie ED, III, Brodie ED, Jr., Ruben PC. 2005. Evolutionary diversification
of TTX-resistant sodium channels in a predator-prey interaction. *Nature* 434(7034):759-763.

Go Y, Satta Y, Kawamoto Y, Rakotoarisoa G, Randrianjafy A, Koyama N, Hirai H. 2003. Frequent
segmental sequence exchanges and rapid gene duplication characterize the MHC class I genes in lemurs.
Immunogenetics 55(7):450-461.

Goedbloed DJ, Czypionka T, Altmüller J, Rodriguez A, Küpfer E, Segev O, Blaustein L, Templeton AR,
Nolte AW, Steinfartz S. 2017. Parallel habitat acclimatization is realized by the expression of different
genes in two closely related salamander species (genus *Salamandra*). *Heredity* 119(6):429-437.

Graham AM, McCracken KG. 2019. Convergent evolution on the hypoxia-inducible factor (HIF)
pathway genes EGLN1 and EPAS1 in high-altitude ducks. *Heredity* 122(6):819-832.

Hague MTJ, Stokes AN, Feldman CR, Brodie Jr. ED, Brodie III ED. 2020. The geographic mosaic of
arms race coevolution is closely matched to prey population structure. *Evolution Letters* 4(4):317-332.

Hanifin CT. 2010. The chemical and evolutionary ecology of tetrodotoxin (TTX) toxicity in terrestrial
vertebrates. *Marine Drugs* 8(3):577-593.

Hanifin CT, Gilly WF. 2015. Evolutionary history of a complex adaptation: tetrodotoxin resistance in
salamanders. *Evolution* 69(1):232-244.

Hanikenne M, Kroymann J, Trampczynska A, Bernal M, Motte P, Clemens S, Krämer U. 2013. Hard
selective sweep and ectopic gene conversion in a gene cluster affording environmental adaptation. *PLOS
Genet* 9(8):e1003707.

Hansen TF, Carter AJR, Chiu C-H. 2000. Gene conversion may aid adaptive peak shifts. *J Theor Biol*
207(4):495-511.

Hellsten U, Harland RM, Gilchrist MJ, Hendrix D, Jurka J, Kapitonov V, Ovcharenko I, Putnam NH, Shu
S, Taher L, et al. 2010. The genome of the western clawed frog *Xenopus tropicalis*. *Science*
328(5978):633-636.

Hime PM, Lemmon AR, Lemmon ECM, Prendini E, Brown JM, Thomson RC, Kratovil JD, Noonan BP,
Pyron RA, Peloso PLV, et al. 2021. Phylogenomics reveals ancient gene tree discordance in the
amphibian tree of life. *Syst Biol* 70(1):49-66.

Howe K, Clark MD, Torroja CF, Torrance J, Berthelot C, Muffato M, Collins JE, Humphray S, McLaren
K, Matthews L, et al. 2013. The zebrafish reference genome sequence and its relationship to the human
genome. *Nature* 496(7446):498-503.

Jeziorski MC, Greenberg RM, Anderson PAV. 1997. Cloning of a putative voltage-gated sodium channel
from the turbellarian flatworm *Bdelloura candida*. *Parasitology* 115(3):289-296.

Jost MC, Hillis DM, Lu Y, Kyle JW, Fozzard HA, Zakon HH. 2008. Toxin-resistant sodium channels:
parallel adaptive evolution across a complete gene family. *Mol Biol Evol* 25(6):1016-1024.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in
performance and usability. *Mol Biol Evol* 30(4):772-780.

Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz
S, Duran C, et al. 2012. Geneious Basic: An integrated and extendable desktop software platform for the
organization and analysis of sequence data. *Bioinformatics* 28(12):1647-1649.

Khakhlova O, Bock R. 2006. Elimination of deleterious mutations in plastid genomes by gene
conversion. *Plant J* 46(1):85-94.

Kosakovsky Pond SL, Frost SDW. 2005. Not so different after all: a comparison of methods for detecting
amino acid sites under selection. *Mol Biol Evol* 22(5):1208-1222.

Kuhner MK, Lawlor DA, Ennis PD, Parham P. 1991. Gene conversion in the evolution of the human and
chimpanzee MHC class I loci. *Tissue Antigens* 38(4):152-164.

Leffler A, Herzog RI, Dib-Hajj SD, Waxman SG, Cummins TR. 2005. Pharmacological properties of
neuronal TTX-resistant sodium channels and the role of a critical serine pore residue. *Pflug Arch Eur J*
Phy 451(3):454-463.

Losos JB. 2011. Convergence, adaptation, and constraint. *Evolution* 65(7):1827-1840.

Márquez R, Ramírez-Castañeda V, Amézquita A. 2019. Does batrachotoxin autoresistance coevolve with
toxicity in *Phyllobates* poison-dart frogs? *Evolution* 73(2):390-400.

Maruta S, Yamaoka K, Yotsu-Yamashita M. 2008. Two critical residues in P-loop regions of pufferfish
Na$^+$ channels on TTX sensitivity. *Toxicon* 51(3):381-387.

Matsunami M, Kitano J, Kishida O, Michimae H, Miura T, Nishimura K. 2015. Transcriptome analysis of
predator- and prey-induced phenotypic plasticity in the Hokkaido salamander (*Hynobius retardatus*).
Molecular Ecology 24(12):3064-3076.

McGlothlin JW, Chuckalovcak JP, Janes DE, Edwards SV, Feldman CR, Brodie ED, Jr., Pfrender ME,
Brodie ED, III. 2014. Parallel evolution of tetrodotoxin resistance in three voltage-gated sodium channel
genes in *Thamnophis sirtalis*. *Mol Biol Evol* 31(11):2386-2846.

McGlothlin JW, Kobiela ME, Feldman CR, Castoe TA, Vonk FJ, Richardson MK, Brodie ED, Jr.,
Pfrender ME, Brodie ED, III. 2016. Historical contingency in a multigene family facilitates adaptive
evolution of toxin resistance. *Curr Biol* 26:1616-1621.

McGrath CL, Casola C, Hahn MW. 2009. Minimal effect of ectopic gene conversion among recent
duplicates in four mammalian genomes. *Genetics* 182(2):615-622.

Moreno-Hagelsieb G, Latimer K. 2007. Choosing BLAST options for better detection of orthologs as
reciprocal best hits. *Bioinformatics* 24(3):319-324.

Nowoshilow S, Schloissnig S, Fei J-F, Dahl A, Pang AWC, Pippel M, Winkler S, Hastie AR, Young G,
Roscito JG, et al. 2018. The axolotl genome and the evolution of key tissue formation regulators. *Nature*
554(7690):50-55.

Ohta T. 1989. The mutational load of a multigene family with uniform members. *Genet Res* 53(2):141-
145.

Perry BW, Card DC, McGlothlin JW, Pasquesi GIM, Adams RH, Schield DR, Hales NR, Corbin AB,
Demuth JP, Hoffmann FG, et al. 2018. Molecular adaptations for sensing and securing prey and insight
into amniote genome diversity from the garter snake genome. *Genome Biol Evol* 10(8):2110-2129.

Petschenka G, Wagschal V, von Tschirnhaus M, Donath A, Dobler S. 2017. Convergently evolved toxic
secondary metabolites in plants drive the parallel molecular evolution of insect resistance. *Am Nat*
190(S1):S29-S43.

Posada D. 2008. jModelTest: Phylogenetic model averaging. *Mol Biol Evol* 25(7):1253-1256.

Posada D, Crandall KA. 2001. Evaluation of methods for detecting recombination from DNA sequences:
computer simulations. *P Natl Acad Sci USA* 98(24):13757-13762.

Pyron RA. 2014. Biogeographic analysis reveals ancient continental vicariance and recent oceanic
dispersal in amphibians. *Syst Biol* 63(5):779-797.

Sawyer SA. 1999. GENECONV: a computer package for the statistical detection of gene conversion.
http://www.math.wustl.edu/~sawyer.

Semple C, Wolfe KH. 1999. Gene duplication and gene conversion in the Caenorhabditis elegans
genome. *J Mol Evol* 48(5):555-564.

Signore AV, Yang Y-Z, Yang Q-Y, Qin G, Moriyama H, Ge R-L, Storz JF. 2019. Adaptive changes in
hemoglobin function in high-altitude Tibetan canids were derived via gene conversion and introgression.
Mol Biol Evol 36(10):2227-2237.

Smit A, Hubley R, Green P. 2013-2015. RepeatMasker Open-4.0.

Smith JJ, Timoshevskaya N, Timoshevskiy VA, Keinath MC, Hardy D, Voss SR. 2019. A chromosome-
scale assembly of the axolotl genome. *Genome Research* 29(2):317-324.

Smith SD, Pennell MW, Dunn CW, Edwards SV. 2020. Phylogenetics is the new genetics (for most of
biodiversity). *Trends Ecol Evol* 35(5):415-425.

Soong TW, Venkatesh B. 2006. Adaptive evolution of tetrodotoxin resistance in animals. *Trends in
Genetics* 22(11):621-626.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. *Bioinformatics* 30(9):1312-1313.

Stokes AN, Ducey PK, Neuman-Lee L, Hanifin CT, French SS, Pfrender ME, Brodie ED, III, Brodie ED,
Jr. 2014. Confirmation and distribution of tetrodotoxin for the first time in terrestrial Invertebrates: two
terrestrial flatworm species (*Bipalium adventitium* and *Bipalium kewense*). *PLOS One* 9(6):e100718.

Storz JF. 2016. Causes of molecular convergence and parallelism in protein evolution. *Nature Reviews
Genetics* 17(4):239-250.

Sun YB, Xiong ZJ, Xiang XY, Liu SP, Zhou WW, Tu XL, Zhong L, Wang L, Wu DD, Zhang BL, et al.
2015. Whole-genome sequence of the Tibetan frog *Nanorana parkeri* and the comparative evolution of
tetrapod genomes. *P Natl Acad Sci USA* 112(11):E1257.

Sunami A, Glaaser IW, Fozzard HA. 2000. A critical residue for isoform difference in tetrodotoxin
affinity is a molecular determinant of the external access path for local anesthetics in the cardiac sodium
channel. *P Natl Acad Sci USA* 97(5):2326-2331.

Suyama M, Torrents D, Bork P. 2006. PAL2NAL: robust conversion of protein sequence alignments into
the corresponding codon alignments. *Nucleic Acids Research* 34(Web Server issue):W609-W612.

Tarvin RD, Borghese CM, Sachs W, Santos JC, Lu Y, O'Connell LA, Cannatella DC, Harris RA, Zakon
HH. 2017. Interacting amino acid replacements allow poison frogs to evolve epibatidine resistance.
Science 357(6357):1261-1266.

Toledo G, Hanifin C, Geffeney S, Brodie ED, III. 2016. Convergent evolution of tetrodotoxin-resistant
sodium channels in predators and prey. *Curr Top Membr* 78:87-113.

Vaelli PM, Theis KR, Williams JE, O'Connell LA, Foster JA, Eisthen HL. 2020. The skin microbiome
facilitates adaptive tetrodotoxin production in poisonous newts. *eLife* 9:e53898.

Venkat A, Hahn MW, Thornton JW. 2018. Multinucleotide mutations cause false inferences of lineage-
specific positive selection. *Nat Ecol Evol* 2(8):1280-1288.

Venkatesh B, Lu SQ, Dandona N, See SL, Brenner S, Soong TW. 2005. Genetic basis of tetrodotoxin
resistance in pufferfishes. *Curr Biol* 15(22):2069-2072.

Widmark J, Sundström G, Ocampo Daza D, Larhammar D. 2011. Differential evolution of voltage-gated
sodium channels in tetrapods and teleost fishes. *Mol Biol Evol* 28(1):859-871.

Williams BL, Hanifin CT, Brodie ED, Jr., Brodie ED, III. 2010. Tetrodotoxin affects survival probability
of rough-skinned newts (*Taricha granulosa*) faced with TTX-resistant garter snake predators
(*Thamnophis sirtalis*). *Chemoecology* 20(4):285-290.

Yang Z. 1998. Likelihood ratio tests for detecting positive selection and application to primate lysozyme
evolution. *Mol Biol Evol* 15(5):568-573.

Yang Z, dos Reis M. 2010. Statistical properties of the branch-site test of positive selection. *Mol Biol
Evol* 28(3):1217-1228.

Yang Z, Nielsen R, Goldman N, Pedersen A-MK. 2000. Codon-substitution models for heterogeneous
selection pressure at amino acid sites. *Genetics* 155(1):431-449.

Yang Z, Wong WSW, Nielsen R. 2005. Bayes empirical Bayes inference of amino acid sites under
positive selection. *Mol Biol Evol* 22(4):1107-1118.

Yang ZH. 2007. PAML 4: Phylogenetic analysis by maximum likelihood. *Mol Biol Evol* 24(8):1586-
1591.

Yu FH, Catterall WA. 2003. Overview of the voltage-gated sodium channel family. *Genome Biol*
4(3):207.

Zakon HH. 2012. Adaptive evolution of voltage-gated sodium channels: The first 800 million years. *P
Natl Acad Sci USA* 109(Supplement 1):10619-10625.

Zakon HH, Jost MC, Lu Y. 2011. Expansion of voltage-dependent Na$^+$ channel gene family in early
tetrapods coincided with the emergence of terrestriality and increased brain complexity. *Mol Biol Evol*
28(4):1415-1424.

Zhang J, Nielsen R, Yang Z. 2005. Evaluation of an improved branch-site likelihood method for detecting
positive selection at the molecular level. *Mol Biol Evol* 22(12):2472-2479.

Zhen Y, Aardema ML, Medina EM, Schumer M, Andolfatto P. 2012. Parallel molecular evolution in an
herbivore community. *Science* 337(6102):1634-1637.
845