A covariation analysis reveals elements of selectivity in quorum sensing systems

Wellington Miranda, S[1], Cong, Q[2, 3, 4, 5], Schaefer, AL[1], MacLeod, EK[1], Zimenko, A[1], Baker, D[2, 3, 6], & Greenberg, EP[1, *]

1. Department of Microbiology, University of Washington, Seattle, WA 98195, USA
2. Department of Biochemistry, University of Washington, Seattle, WA 98105, USA
3. Institute for Protein Design, University of Washington, Seattle, WA 98105, USA
4. Eugene McDermott Center for Human Growth and Development, University of Texas Southwestern Medical Center, Dallas, Texas 75390, USA
5. Department of Biophysics, University of Texas Southwestern Medical Center, Dallas, TX 75390, USA
6. Howard Hughes Medical Institute, University of Washington, Seattle, WA 98105, USA

*Corresponding author, epgreen@uw.edu

Abstract

Many bacteria communicate with kin and coordinate group behaviors through a form of cell-cell

signaling called acyl-homoserine lactone (AHL) quorum sensing (QS). In these systems, a signal

synthase produces an AHL to which its paired receptor selectively responds. Selectivity is fundamental

to cell signaling. Despite its importance, it has been challenging to determine how this selectivity is

achieved and how AHL QS systems evolve and diversify. We hypothesized that we could use

covariation within the protein sequences of AHL synthases and receptors to identify selectivity residues.

We began by identifying about 6,000 unique synthase-receptor pairs. We then used the protein

sequences of these pairs to identify covariation patterns and mapped the patterns onto the LasI/R system

from *Pseudomonas aeruginosa* PAO1. The covarying residues in both proteins cluster around the ligand

binding sites. We demonstrate that these residues are involved in system selectivity toward the cognate

signal and go on to engineer the Las system to both produce and respond to an alternate AHL signal. We

have thus demonstrated a new application for covariation methods and have deepened our understanding

of how communication systems evolve and diversify.

Quorum sensing (QS) is a widespread form of cell-cell signaling that bacteria use to coordinate

the production of public goods including toxins, antibiotics, bioluminescence, and secreted enzymes[1,2].

Many Proteobacteria[3] and Nitrospirae[4] employ a form of QS based on acyl-homoserine lactone (AHL)

signals. AHL QS systems consist of two proteins: a LuxI-type signal synthase and a LuxR-type receptor

(Figure 1a). The signal synthase produces an AHL from S-adenosylmethionine (SAM) and an acyl-acyl

carrier protein (ACP) for some LuxI-type synthases or an acyl-coenzyme A (CoA) substrate for others[5]

(Fig. 1b). AHL signals can freely diffuse through cell membranes[6,7] and at low cell density the QS

system is "off". At high cell density, the signal accumulates and binds the LuxR-type receptor which is a

cytosolic transcription factor that regulates gene expression in response to signal binding.

AHL signals share a conserved lactone core, but vary in the acyl moiety which can be a fatty

acid ranging from 4 to 20 carbons long, with potential oxidation on the C3 carbon and varying degrees

of unsaturation, or can have an aromatic or branched structure[8]. This variability in the acyl portion of the

signal confers selectivity to the system. Typically, a LuxI-type synthase produces a primary AHL to

which its paired LuxR-type receptor selectively responds[9]. Selectivity is critical to cell signaling in order

to avoid undesired cross-talk or spurious outputs[10]. In the case of QS, selectivity ensures bacteria

cooperate only with kin cells.

Despite its importance, we know little about how QS systems achieve selectivity or how they

evolve and diversify to use new signals. Although the conserved amino acids essential for synthase and

receptor activity are well described[11,12], residues that dictate selectivity are often different from those

that are required for activity[13]. Due to the low amino acid sequence identity between LuxI/R

homologues, it has been difficult to determine how QS systems discriminate between various AHL

signals[14].

We hypothesized that we could use covariation patterns to identify QS selectivity residues. Such

methods have been used to identify amino acid residues that interact with each other within proteins and

between proteins that physically bind each other[15-17]. Here, we endeavored to expand these methods to

assess the interaction between AHL synthases and receptors. While AHL synthases and receptors do not

physically interact, they interact indirectly via binding to a shared cognate signal and are believed to

coevolve to maintain this shared signal recognition[9]. Phylogenetic analyses also support coevolution of

synthases and receptors[18,19]. We therefore hypothesized that we could identify amino acid residues that

covary between QS synthases and receptors, and further, that the covarying residues would be those

responsible for signal selectivity.

We used a statistical method, GREMLIN[20], to measure covariation within the sequences of AHL

synthase-receptor pairs and mapped the covarying residues onto the LasI/R QS system of *Pseudomonas*

aeruginosa PAO1. By mutating the top-scoring residues identified by GREMLIN, we demonstrate that

they are indeed important for signal selectivity and, further, that these residues can be used to rationally

engineer LasI/R to produce and respond to a non-native signal. We thus demonstrate a new application

for powerful covariation methods and at the same time identify determinants of QS selectivity.

Results

Covariation patterns in QS systems. To begin our analysis, we gathered select protein sequences for

known synthase-receptor pairs (Supplementary Table 1) and used these sequences to search the

European Nucleotide Archive (ENA) database[21] from the European Bioinformatics Institute and the

Integrated Microbial Genomes and Microbiomes (IMG/M) database[22] from the Joint Genome Institute

(JGI) for additional synthase-receptor pairs. The genes for synthase-receptor pairs are frequently co-

located on the genome, and organisms can harbor more than one complete QS system[14]. To increase the

likelihood of identifying true pairs, we required that the two genes be separated by no more than two

coding sequences. A total of 6,360 non-identical pairs were identified. We further discarded pairs that

were more than 90% identical to another pair, resulting in 3,489 representative AHL synthase-receptor

pairs.

We aligned these sequences to the LasI/R QS system from *P. aeruginosa* PAO1. Not only is *P.

aeruginosa* a clinically important pathogen, the Las system is well-studied and crystal structures have

been solved for both LasI[23] and LasR[24], making this a particularly useful model system for our studies.

We connected the sequences of synthase and receptor from each pair and used GREMLIN to analyze

covariation patterns in these sequences (Supplementary Fig. 1). We applied Average Product Correction

(APC) to the GREMLIN covariance coefficients, a common technique shown to improve the accuracy

of coevolution analyses[25]. We performed the same analysis by aligning the synthase-receptor pairs to the

LuxI/R system from *Vibrio fischeri* MJ11. The top-ranking coevolving residue pairs overlap

significantly between the LasI/R and LuxI/R systems (62.5% in common among the top 0.05% residue

pairs) (Supplementary Fig. 1). We integrated the analyses for the LasI/R and LuxI/R systems by using

the higher score for each residue pair and the top 10 residue pairs are shown in Fig. 2a. As a control, we

randomly paired the synthases and receptors from different species and reanalyzed them using

GREMLIN. While top-scoring covarying residues had a minimal GREMLIN score (with APC) of 0.09,

the highest score from the randomized control was 0.08 (Fig. 2b). This control provides a guideline for

our analysis; residues with a GREMLIN score (with APC) above or near the maximal score for the

randomized control are likely to meaningful.

Top-scoring residues cluster near ligand-binding pockets. For both LasI and LasR, the top-scoring

covarying residues cluster around the ligand-binding pocket. For LasR, the top-scoring residues map

exclusively to the ligand-binding domain with an average distance of 5.0 Å from the co-crystalized

native ligand *N*-3-oxo-dodecanoyl-L-homoserine lactone (3OC12-HSL) (Fig. 2c). In contrast, the

residues identified in the randomized control are scattered throughout LasR, including three residues in

the DNA-binding domain, and are an average distance of 17.8 Å from 3OC12-HSL (Fig. 2d).

In LasI, the top-scoring covarying residues cluster around the hydrophobic pocket thought to bind

the fatty acyl substrate (Fig. 2e) and are an average distance of 3.7 Å from an acyl substrate modeled

into the LasI structure[23]. As with LasR, the residues identified in the randomized control are scattered

throughout LasI, with many of the residues exposed to solvent (Fig. 2f). The randomized control

residues in LasI are over three times further from the fatty acyl substrate, mean distance = 11.7Å,

compared to the covarying residues.

Due to their location near the ligand-binding pockets, several of the covarying residues have been

previously studied in various LasI/R homologues. Encouragingly, many of these residues have been

reported to be important for protein activity and, in some cases, for selectivity. We have summarized

several of these studies in Supplementary Tables 2 and 3.

LasR mutations alter selectivity. To determine whether residues identified by GREMLIN are involved

in LasR selectivity, we mutated a selection of the top-scoring amino acids, G38, R61, A127, S129, and

L130, to the most common natural variants at each position (Supplementary Table 4). By expressing

LasR in *Escherichia coli* and measuring its activity against a previously reported panel of 19 AHL

signals[26], we were able to quickly prioritize mutants for further study. The majority of our LasR mutants

retained the ability to respond to AHLs and all mutants had an altered selectivity profile when compared

to wild-type (Supplementary Fig. 2).

We, and others, have previously demonstrated that compared to native activity, QS receptor

sensitivity and selectivity can be altered when in *E. coli*[26,27]. We therefore engineered several mutations

into *lasR* on the *P. aeruginosa* PAO-SC4 chromosome to confirm our findings. *P. aeruginosa* PAO-SC4

is an AHL synthase-null mutant which we use here to measure LasR activity in response to exogenously

provided AHL signals. The *lasR* mutations largely had the same effect on activity and selectivity in *P.*

aeruginosa as they did when *lasR* was expressed *E. coli* (Supplementary Fig. 3). Of note, LasR^{A127L} had

an increased sensitivity to numerous signals (Fig. 3a-d), potentially through increased hydrophobic

interactions with the fatty acyl chain of the AHLs. Consistent with its role in a water-mediated hydrogen

bond with the C3 oxygen of 3OC12-HSL, and with previous studies[28,29], LasR R61 mutants were less

responsive to oxo-substituted AHLs, but maintained wild-type or better levels of activation by

unsubstituted AHLs (Fig. 3a-d and Supplementary Fig. 3).

The mutations also affected the sensitivity of LasR to 3OC12-HSL (Supplementary Fig. 3).

Interestingly, two of our mutants were more sensitive to 3OC12-HSL than wild-type LasR. LasR^{A127L}

was roughly 3-fold more sensitive and LasR^{L130F} was 2-fold more sensitive (Fig. 3e). This increased

sensitivity came at the cost of decreased selectivity for both of these mutations. In fact, many of our

single amino acid mutants displayed reduced selectivity compared to wild-type LasR (Supplementary

Fig. 2 and Supplementary Fig. 3).

LasI mutations alter activity and selectivity. Similar to LasR, we focused our LasI mutations on the

top-scoring positions: L102, T142, T145, and L157 (Supplementary Table 5). We expressed wild-type

or mutated *lasI* on a low copy number plasmid in the AHL synthase-null *P. aeruginosa* PAO-SC4 and

extracted AHLs produced by these bacteria from culture fluid. While bioassays are commonly used for

the detection of AHLs[30], they suffer from multiple drawbacks. In particular, bioassays are not equally

sensitive to all AHLs and typically cannot be used to determine which AHLs are produced and in what

ratio. To screen our LasI mutants for altered activity and selectivity, we developed a thin layer

chromatography (TLC) method based on our existing high performance liquid chromatography (HPLC)

radiotracer assay[31]. In this method, the C1 position in the homoserine lactone ring is labeled with ^{14}C.

The label is incorporated into AHLs at a ratio of one ^{14}C per AHL molecule. This results in unbiased

detection of all AHLs produced. While the established method uses HPLC to separate and detect AHLs

one sample at a time, we can run nine samples per TLC, resulting in a more high-throughput assay.

Using our TLC method, we confirmed that *lasI* directs the synthesis of the same primary product

whether it is expressed on a plasmid or from the chromosome (Supplementary Fig. 4). HPLC analysis of

matched extracts confirmed that the major LasI product observed by TLC is 3OC12-HSL. As expected,

an empty vector control did not produce detectable AHLs, nor did we detect radioactivity in a media-

only control. We then screened the activity each *lasI* mutant by TLC (Supplementary Fig. 4). Several

mutants produced little or no detectable AHLs, while some appeared to produce more 3OC12-HSL than

wild-type LasI. We analyzed select extracts by both TLC and HPLC and found that the results were

consistent between the two methods, further validating the TLC method.

Based on our TLC results we selected one mutant, LasI^{L157W}, for further study by HPLC. We

found that LasI^{L157W} produces equal amounts of two ^{14}C-AHLs that elute in the fractions of *N*-3-oxo-

decanoyl-L-homoserine lactone (3OC10-HSL) and *N*-3-oxo-octanoyl-L-homoserine lactone (3OC8-

HSL) along with a lesser amount of 3OC12-HSL (Fig. 4). These findings demonstrate that the covarying

residues influence LasI activity and selectivity, and that a single mutation is sufficient to significantly

alter LasI selectivity.

Multiple mutations can "rewire" LasI/R selectivity. In general, multiple mutations are required to

generate a protein with orthogonal selectivity[15,17,28]. In non-QS proteins, altered selectivity has been

engineered by swapping the covarying residues in one homolog to the identities in another[15,17]. Here, we

seek to "rewire" LasI/R to use an orthogonal signal. We targeted the MupI/R system from *Pseudomonas*

fluorescens NCIMB 10586, which uses the signal 3OC10-HSL[32]. MupI and MupR share 52% and 39%

identity with LasI and LasR, respectively.

LasR and MupR differ at eight covariation sites in the ligand-binding domain with a GREMLIN

score (with APC) > 0.08 (Supplementary Fig. 5). LasR modified to contain all eight mutations was

inactive. However, there were several intermediate mutants that displayed an increased response to

3OC10-HSL. We identified three mutations that are sufficient for this increased sensitivity: L125F,

A127M, and L130F (Fig. 5a). LasR$^{L125F, A127M, L130F}$ is over 20-fold more sensitive to 3OC10-HSL than

wild-type LasR. The L125F mutation appears to be the primary driver of this altered selectivity (Fig.

5b,c and Supplementary Fig. 5). All "MupR-like" LasR mutants responded to 3OC12-HSL with similar

sensitivity to wild-type LasR (Supplementary Fig. 5).

LasI differs from MupI at five high-scoring covariation residues: LasI M125, T145, M152, V159,

and N181 (Supplementary Fig. 6), the first 3 of which line the LasI acyl-binding pocket (Fig. 5d).

Swapping these three residues for their MupI identities resulted in a synthase that has substantially

altered selectivity. LasI$^{M125I, T145S, M152L}$ produces ~2-fold more 3OC10-HSL than 3OC12-HSL (Fig. 5e).

The M125I mutation alone was sufficient to relax LasI's selectivity, resulting in a synthase that produces

roughly equal amounts of 3OC10-HSL and 3OC12-HSL. As a comparison, we measured the activity of

mupI expressed in *P. aeruginosa*, and found it produces 9:1 3OC10-HSL:3OC12-HSL (Fig. 5e and

Supplementary Fig. 6). All single and double "MupI-like" LasI mutants retained AHL synthase activity,

but only those mutants that contain the M125I mutation displayed increased 3OC10-HSL production

relative to 3OC12-HSL (Supplementary Fig. 6).

Discussion

Despite decades of study, it has been challenging to determine how AHL QS systems distinguish

between signals. We hypothesized that we could identify covariation patterns in AHL QS systems and

that these patterns would illuminate residues important for signal selectivity. By analyzing the sequences

of 6,360 unique QS systems, we identified amino acids that strongly covary between AHL synthases and

receptors. The top-scoring residues in our analysis cluster near the ligand-binding pockets for both

proteins and are more than three times closer to the signal molecule compared to top-scoring residues in

a randomized control. We focused our study on *P. aeruginosa* LasI/R. Through targeted mutations in the

top-scoring covarying residues we demonstrate that these amino acids are indeed determinants of signal

selectivity. We have thus validated a new application of covariation analysis for proteins that interact

indirectly and not through direct binding to one-another. Additionally, these strong covariation results

further support the view that AHL synthases and receptors coevolve.

For both the synthase, LasI, and the receptor, LasR, a single amino acid substitution is sufficient

to significantly alter selectivity. Interestingly, our mutations also revealed that LasR is not optimized to

be as sensitive as possible to its native 3OC12-HSL signal. The increase in sensitivity of specific

mutants came at the cost of decreased selectivity, which suggests that QS systems may evolve to balance

these two properties. Furthermore, increased sensitivity to the native signal may lead to premature

activation of the QS regulon, which would likely decrease fitness[33]. The mutants generated in our study

provide us with the tools to directly address these questions and assess the impact of sensitivity and

selectivity on QS function.

We also demonstrated that we can use covarying residues to rationally engineer a QS system to

produce and respond to a signal of our choosing. By mutating the covarying residues in LasI/R, we

improved the sensitivity of LasR to 3OC10-HSL over 20-fold and increased the production of 3OC10-

HSL by LasI roughly 15-fold. For both the synthase and receptor, a single amino acid substitution was

the primary driver of the altered selectivity. This was surprising given the low sequence identity between

LasI/R and MupI/R. These findings suggest new QS systems might evolve with relative ease. Further,

the ability to engineer QS selectivity could be beneficial to synthetic biology where AHL signaling is a

powerful tool to build biological circuits[34].

Though we were able to substantially increase the 3OC10-HSL activity of LasI/R, our mutants retained their native 3OC12-HSL activity. We have thus generated a promiscuous system with broadened selectivity. Similarly, a directed evolution study of the AHL receptor LuxR found that it evolves through promiscuous intermediates[13]. This has also been observed in other systems, such as toxin-antitoxin systems[17]. Proteins tend to evolve through broadly active intermediates before gaining new specificity. In this way, the system maintains functionality *en route* to altered selectivity. Quorum sensing systems appear to follow these same trends.

One limitation we faced is a lack of close LasI/R homologs with known signals. It has been demonstrated that "supporting" residues, i.e. residues within a protein that covary with the selectivity residues, may indirectly impact selectivity by influencing the orientation of selectivity residues[17]. Thus, given the large differences in sequence identity between the Las and Mup systems, there are likely other residues that must be mutated to fully swap selectivity. The identification of a more closely related system to LasI/R may provide a better starting point for engineering altered selectivity. Alternatively, our mutants could be further evolved though saturating mutagenesis and/or *in vitro* evolution.

Collectively, our results provide insight into AHL QS selectivity and will help us predict signal selectivity in newly identified QS systems, in metagenomes, and in naturally occurring QS variants such as those found in clinical isolates. More broadly, we have gained insight into how AHL QS systems evolve and diversify.

Methods

Identification of quorum sensing systems. Starting from 24 pairs of manually curated QS synthases and receptors (Supplementary Table 1), we searched for homologs in complete bacterial genomes using BLAST (e-value <0.01)[35]. We filtered the BLAST hits by sequence identity (>30%) to the query and the alignment coverage (query coverage >0.75 and hit coverage >0.75), and the filtered hits were aligned by

Clustal Omega[36]. We selected the LasI/R system from *Pseudomonas aeruginosa* PAO1 as the target and

mapped the Multiple Sequence Alignments (MSA) to the target system. We built sequence profiles from

the MSA with HMMER[37] and hmmbuild for the QS synthases and receptors, respectively. The sequence

profiles were then used to search against the European Nucleotide Archive database[21] and the Integrated

Microbial Genomes and Microbiomes database[22] from Joint Genome Institute using HMMER

hmmsearch. A total of 149,837 and 5,046,620 homologs were found in these databases for the QS

synthase and receptor, receptively. Because the synthases and receptors of the known QS systems

frequently locate near each other in the genome, we kept synthase-receptor gene pairs that are separated

by no more than two other open reading frames (ORFs) in the genome or contig. A total of 14,980

synthase-receptor gene pairs were identified and they represent 6,360 non-identical QS systems. In

another attempt, we carried out the same procedure using the LuxI/R system from *Vibrio fischeri* MJ11

as the target system. A similar number of QS systems were identified.

Identification of covarying residues. We connected the synthase and receptor protein sequences for

each QS system we found in the databases and derived the alignments between these QS systems to the

target QS system (LasI/R) from the hmmsearch result. We filtered the MSA for synthase-receptor pairs

by sequence identity (maximal identify for remaining sequences <=90%) and gap ratio in each sequence

(maximal gap ratio <=25%). We applied GREMLIN to analyze the covariation in the MSA[20], and the

GREMLIN coefficients were normalized using Average Product Correction (APC)[25] as we described

previously[16]. The GREMLIN coefficients after APC were used as measures for covariation signals

between synthase and receptor amino acid residues. As a control, we connected each synthase sequence

with a randomly selected receptor sequence and performed the covariation analysis in the same way.

We mapped the top-scoring covarying residues in the LasI/R system onto the crystal structures

for each protein. Reported distances between residues and ligands are the shortest distance between any

non-hydrogen atoms. For LasR, distances were calculated using PDB 6V7X. For LasI, distances were

calculated using a LasI structure with 3-oxo-C12-acyl-phosphopantetheine modeled into the acyl-

binding pocket[23]. Reported distances for LasI are between residues and the acyl portion of the modeled

substrate.

Bacterial strains, plasmids, and culture conditions. Bacterial strains and plasmids are listed in

Supplementary Table 6. Unless otherwise specified, *Pseudomonas aeruginosa* and *Escherichia coli*

were grown in lysogeny broth (LB) (10 g tryptone, 5 g yeast extract, 5 g NaCl per liter) buffered with 50

mM 3-(*N*-morpholino) propanesulfonic acid (MOPS) (pH 7) (LB/MOPS) or on LB agar (LB plus 1.5%

Bacto agar)[26]. Liquid cultures were grown at 37°C with shaking. For radiotracer thin layer

chromatography (TLC) experiments, *P. aeruginosa* was grown in Jensen's medium with 0.3%

glycerol[31].

For plasmid selection and maintenance, antibiotics were used at the following concentrations: *P.*

aeruginosa, 30 µg per mL gentamicin (Gm) and 150 µg per mL carbenicillin (Cb); *E. coli* 10 µg per mL

Gm and 100 µg per mL ampicillin (Ap). BD Difco Pseudomonas Isolation Agar (PIA) was prepared

according to manufacturer directions and supplemented with 100 µg per mL Gm as needed. Where

needed for gene expression L-arabinose (0.4% w/v) was added.

All chemicals and reagents were obtained from commercial sources. AHLs were dissolved either

in dimethyl sulfoxide (DMSO) or in ethyl acetate (EtAc) acidified with glacial acetic acid (0.01% v/v).

AHLs in DMSO were used at <=1% of the final culture volume and AHLs dissolved in EtAc were dried

on the bottom of the culture vessel prior to addition of the bacterial culture. DMSO or acidified EtAc

was used as a vehicle control where appropriate.

Plasmid and strain construction. pJN-lasI and pJN-RBSmupI were constructed using *E. coli*-mediated

DNA assembly[38]. Briefly, for pJN-lasI, *lasI* was amplified from *P. aeruginosa* PAO1 genomic DNA

(gDNA) using primers lasI-pJN-F and lasI-pJN-R (Supplementary Table 6). pJN105 was amplified

using the reverse complement of these primers. The resulting PCR products were treated with the

restriction enzyme DpnI to remove the parent template. Both PCR products were then used to transform

E. coli (NEB 5alpha). The resulting constructs were confirmed by Sanger sequencing. For pJN-

RBSmupI, we began by amplifying *mupI* from *Pseudomonas fluorescens* Migula (ATCC 49323) gDNA

using primers mupI-F and mupI-R. We then used primers mupI-pJN-F and mupI-pJN-R to amplify the

mupI PCR product and used the reverse complement of these two primers to amplify pJN-RBSlasI. The

resulting PCR products were treated the same as for pJN-lasI. We constructed pJN-RBSlasI using

restriction digestion. The *lasI* gene, including its upstream ribosomal binding site (RBS), was amplified

from *P. aeruginosa* PAO1 gDNA using primers RBS-lasI-F and lasI-pJN-R. pJN-lasI and the RBS-*lasI*

PCR product were digested using NheI and SacI, gel or column purified respectively, ligated by T4

DNA ligase, and transformed into NEB 5alpha. The resulting constructs were confirmed by Sanger

sequencing. Plasmids were introduced into *E. coli* by using heat shock and were introduced into *P.*

aeruginosa by electroporation.

Point-mutations were introduced to *lasI* and *lasR* on pJN-lasI and JNL or pEXG2-lasR,

respectively, using site directed mutagenesis by PCR. Primers were designed to amplify each plasmid

while introducing the desired mutation(s). The resulting PCR products were treated with DpnI and were

then used to transform NEB 5alpha. Plasmids from the resulting colonies were screened for the desired

mutations by Sanger sequencing. To mutate *lasR* on the *P. aeruginosa* PAO-SC4 chromosome, *E. coli*

S17-1 was used to deliver pEXG2-lasR containing various *lasR* mutations to PAO-SC4 via conjugation

and potential mutants were isolated as previously described[39]. All mutations were confirmed by PCR

amplification of *lasR* from the genome followed by Sanger sequencing.

LasR activity measurements. LasR activity was measured in *E. coli* containing pJNL and pPROBE-

P$_{rsaL}$ or in *P. aeruginosa* PAO-SC4 containing pPROBE-P$_{rsaL}$ using previously reported methods[26].

Briefly, overnight-grown cultures were diluted 1:100 and grown back to log-phase. For *E. coli*, cultures

were grown to an optical density at 600 nm (OD) of 0.3, treated with L-arabinose (0.4%), and incubated

with AHLs for 4 h. For *P. aeruginosa*, cultures were grown to an OD between 0.05 and 0.3, were diluted

to an OD of 0.01 and then incubated with AHLs for 16 to 18 h. LasR activity was measured as GFP

fluorescence (excitation 490 nm, emission 520 nm, gain 50) using a Synergy H1 microplate reader

(Biotek Instruments). Activity measurements were normalized by dividing by OD and subtracting

background values (fluorescence per OD for cultures incubated with vehicle control). Concentrations of

half maximal activation, EC$_{50}$, were calculated using GraphPad Prism.

TLC screening for AHLs. Cultures of *P. aeruginosa* PAO1Δ*rhlI* or of *P. aeruginosa* PAO-SC4 with

pJN-empty or with wild-type or mutated pJN-lasI were grown overnight in Jensen's medium with 0.3%

glycerol. Overnight cultures were used to inoculate fresh medium (1% v/v). When the OD reached 0.5,

lasI expression was induced with arabinose (0.4%) and 1.1 mL cultures were incubated with 1.1 μCi/mL

L-[1-^{14}C]-methionine (^{14}C-methionine) for 90 min[31]. Cells were pelleted by centrifugation and 1 mL of

supernatant fluid was extracted twice with 2 mL acidified EtAc. The extracts were dried under N$_2$ and

resuspended in 15 μL acidified EtAc. Five μL of each extract was spotted on an aluminum backed C18-

W-silica TLC plate (Sorbtech). AHLs were separated using 70% methanol in water, then the TLC plate

was dried and exposed to a phosphor screen for at least 16 h. Phosphor screens were imaged with a

Sapphire Biomolecular Imager (Azure Biosystems). To confirm TLC findings, select extracts were

dried, suspended in methanol and analyzed by C18-reversed-phase high performance liquid

chromatography (HPLC) using a previously reported method[31].

HPLC radiotracer assays for LasI activity. For better detection of AHLs, we slightly modified the

radiolabeling protocol detailed above, modeling it after a previously published method[40]. Cultures of *P.*

aeruginosa PAO-SC4 with wild-type or mutated pJN-RBSlasI were grown overnight in LB/MOPS.

Overnight cultures were used to inoculate 5 mL LB/MOPS (1% v/v). After 2 h, *lasI* expression was

induced with arabinose (0.4%) and cultures were grown to OD 0.7. Cells were centrifuged at 5,000 rpm

for 10 min, and pellets were suspended in 1.1 mL phosphate buffered saline (PBS) with 10 mM glucose.

After shaking incubation at 37°C for 10 min, 1.1 μCi ^{14}C-methionine was added to the cell suspension.

Cell suspensions were incubated with radiolabel for 2 h, after which cells were pelleted by

centrifugation and 1 mL supernatant fluid was extracted twice with 2 mL acidified EtAc. Radiolabeled

AHLs were dried under N_2 and suspended in methanol. One-third of each extract was analyzed by

reversed-phase HPLC using a gradient of 10 to 100% methanol-in-water[31].

References

1. Whiteley, M., Diggle, S.P. & Greenberg, E.P. Progress in and promise of bacterial quorum sensing research. *Nature* **551**, 313-320 (2017).

2. Waters, C.M. & Bassler, B.L. Quroum sensing: Cell-to-cell communication in bacteria. *Annu. Rev. Cell Dev. Biol.* **21**, 319-346 (2005).

3. Case, R.J., Labbate, M. & Kjelleberg, S. AHL-driven quorum-sensing circuits: Their frequency and function among the Proteobacteria. *ISME J.* **2**, 345-349 (2008).

4. Mellbye, B.L. *et al.* Acyl-homoserine lactone production in nitrifying bacteria of the genera *Nitrosospira*, *Nitrobacter*, and *Nitrospira* identified via a survey of putative quorum-sensing genes. *Appl. Environ. Microbiol.* **83**, e01540-01517 (2017).

5. Schaefer, A.L. *et al.* A new class of homoserine lactone quorum-sensing signals. *Nature* **454**, 595-599 (2008).

6. Kaplan, H.B. & Greenberg, E.P. Diffusion of autoinducer is involved in regulation of the *Vibrio fischeri* luminescence system. *J. Bacteriol.* **163**, 1210 (1985).

7. Pearson, J.P., Van Delden, C. & Iglewski, B.H. Active efflux and diffusion are involved in transport of *Pseudomonas aeruginosa* cell-to-cell signals. *J. Bacteriol.* **181**, 1203 (1999).

8. Rajput, A., Kaur, K. & Kumar, M. SigMol: Repertoire of quorum sensing signaling molecules in prokaryotes. *Nucleic Acids Res.* **44**, D634-639 (2016).

9. Aframian, N. & Eldar, A. A bacterial tower of babel: Quorum-sensing signaling diversity and its evolution. *Annu. Rev. Microbiol.* **74**, 587-606 (2020).

10. Laub, M.T. Keeping signals straight: How cells process information and make decisions. *PLoS Biol.* **14**, e1002519 (2016).

11. Parsek, M.R., Schaefer, A.L. & Greenberg, E.P. Analysis of random and site-directed mutations in *rhlI*, a *Pseudomonas aeruginosa* gene encoding an acylhomoserine lactone synthase. *Mol. Microbiol.* **26**, 301-310 (1997).

12. Zhang, R.-G. *et al.* Structure of a bacterial quorum-sensing transcription factor complexed with pheromone and DNA. *Nature* **417**, 971-974 (2002).

13. Collins, C.H., Arnold, F.H. & Leadbetter, J.R. Directed evolution of *Vibrio fischeri* LuxR for increased sensitivity to a broad spectrum of acyl-homoserine lactones. *Mol. Microbiol.* **55**, 712-723 (2005).

14. Fuqua, C., Winans, S.C. & Greenberg, E.P. Census and consensus in bacterial ecosystems: The LuxR-LuxI family of quorum-sensing transcriptional regulators. *Annu. Rev. Microbiol.* **50**, 727-751 (1996).

15. Skerker, J.M. *et al.* Rewiring the specificity of two-component signal transduction systems. *Cell*
133, 1043-1054 (2008).

16. Ovchinnikov, S., Kamisetty, H. & Baker, D. Robust and accurate prediction of residue–residue
interactions across protein interfaces using evolutionary information. *eLife* **3**, e02030 (2014).

17. Aakre, C.D. *et al.* Evolving new protein-protein interaction specificity through promiscuous
intermediates. *Cell* **163**, 594-606 (2015).

18. Gray, K.M. & Garey, J.R. The evolution of bacterial LuxI and LuxR quorum sensing regulators.
Microbiology **147**, 2379-2387 (2001).

19. Lerat, E. & Moran, N.A. The evolutionary history of quorum-sensing systems in bacteria. *Mol. Biol.*
Evol. **21**, 903-913 (2004).

20. Kamisetty, H., Ovchinnikov, S. & Baker, D. Assessing the utility of coevolution-based residue–
residue contact predictions in a sequence- and structure-rich era. *PNAS* **110**, 15674 (2013).

21. Amid, C. *et al.* The European Nucleotide Archive in 2019. *Nucleic Acids Res.* **48**, D70-D76 (2019).

22. Chen, I.-M.A. *et al.* The IMG/M data management and analysis system v.6.0: New tools and
advanced capabilities. *Nucleic Acids Res.* **49**, D751-D763 (2020).

23. Gould, T.A., Schweizer, H.P. & Churchill, M.E.A. Structure of the *Pseudomonas aeruginosa* acyl-
homoserinelactone synthase LasI. *Mol. Microbiol.* **53**, 1135-1146 (2004).

24. Zou, Y. & Nair, S.K. Molecular basis for the recognition of structurally distinct autoinducer mimics
by the *Pseudomonas aeruginosa* LasR quorum-sensing signaling receptor. *Chem. Biol.* **16**, 961-
970 (2009).

25. Buslje, C.M. *et al.* Correction for phylogeny, small number of observations and data redundancy
improves the identification of coevolving amino acid pairs using mutual information.
Bioinformatics **25**, 1125-1131 (2009).

26. Wellington, S. & Greenberg, E.P. Quorum sensing signal selectivity and the potential for
interspecies cross talk. *mBio* **10**, e00146-00119 (2019).

27. Moore, J.D. *et al.* A comparative analysis of synthetic quorum sensing modulators in *Pseudomonas*
aeruginosa: New insights into mechanism, active efflux susceptibility, phenotypic response, and
next-generation ligand design. *J. Am. Chem. Soc.* **137**, 14626-14639 (2015).

28. Collins, C.H., Leadbetter, J.R. & Arnold, F.H. Dual selection enhances the signaling specificity of a
variant of the quorum-sensing transcriptional activator LuxR. *Nat. Biotechnol.* **24**, 708-712
(2006).

29. Gerdt, J.P. *et al.* Unraveling the contributions of hydrogen-bonding interactions to the activity of
native and non-native ligands in the quorum-sensing receptor LasR. *Org. Biomol. Chem.* **13**,
1453-1462 (2015).

30. Chu, W. *et al.* Bioassays of quorum sensing compounds using *Agrobacterium tumefaciens* and
Chromobacterium violaceum in *Quorum Sensing: Methods and Protocols* Vol. 692 Ch. 1
(Humana Press, Totowa, NJ, 2011).

31. Schaefer, A.L., Harwood, C.S. & Greenberg, E.P. "Hot Stuff": The many uses of a radiolabel assay
in detecting acyl-homoserine lactone quorum-sensing signals in *Quorum Sensing: Methods and*
Protocols Vol. 1673 Ch. 3 (Springer New York, New York, NY, 2018).

32. Hothersall, J. *et al.* Manipulation of quorum sensing regulation in *Pseudomonas fluorescens*
NCIMB 10586 to increase mupirocin production. *Appl. Microbiol. Biotechnol.* **90**, 1017-1026
(2011).

33. Darch, S.E. *et al.* Density-dependent fitness benefits in quorum-sensing bacterial populations. *PNAS*
109, 8259-8263 (2012).

34. Davis, R.M., Muller, R.Y. & Haynes, K.A. Can the natural diversity of quorum-sensing advance
synthetic biology? *Front. Bioeng. Biotechnol.* **3**, e1-e30 (2015).

35. Altschul, S.F. *et al.* Basic local alignment search tool. *J. Mol. Biol.* **215**, 403-410 (1990).

36. Sievers, F. & Higgins, D.G. The clustal omega multiple alignment package. *Methods Mol. Biol.*
2231, 3-16 (2021).

37. Eddy, S.R. A new generation of homology search tools based on probabilistic inference. *Genome*
Inform. **23**, 205-211 (2009).

38. Kostylev, M. *et al.* Cloning should be simple: *Escherichia coli* DH5α-mediated assembly of
multiple DNA fragments with short end homologies. *PLoS One* **10**, e0137466 (2015).

39. Kostylev, M. *et al.* Evolution of the *Pseudomonas aeruginosa* quorum-sensing hierarchy. *PNAS*
116, 7027 (2019).

40. Leadbetter, J.R. & Greenberg, E.P. Metabolism of acyl-homoserine lactone quorum-sensing signals
by *Variovorax paradoxus*. *J. Bacteriol.* **182**, 6921-6926 (2000).

Acknowledgements

We thank Mair Churchill for sharing her lab's LasI structure modeled with an acyl substrate. This work was supported by grants NIH R35GM136218 to EPG, Yeast Program Grant 5 P41 GM103533-24, Washington Research Foundation fellowship to QC, and Helen Hay Whitney Foundation fellowship to SWM.

Author contributions

SWM, QC, DB, and EPG conceived of and designed the investigation. QC carried out bioinformatic analyses. SWM, EM, and AZ constructed mutants and measured their activity. SWM and ALS developed and conducted radioassays. SWM together with QC, ALS, EPG, and DB interpreted data and wrote the manuscript.

Competing interests

The authors declare no competing interests.

Fig. 1 | Schematic of acyl-homoserine lactone (AHL) quorum sensing (QS). a) AHL QS circuits

consist of a signal synthase, a LuxI homolog (LasI in this cartoon), that produces an AHL signal. At low

cell densities, the system exists in an "off" state. At high cell densities, the AHL concentration increases

and the signal binds its cytosolic receptor, a LuxR homolog (LasR in this cartoon), which functions as a

transcription factor. **b)** LuxI-type synthases produce AHL signals from two substrates: S-adenosyl

methionine (SAM) and an activated organic acid in the form of an acyl-acyl carrier protein (ACP) or

acyl-coenzyme A (CoA). SAM provides the lactone core, which is conserved across all AHL signals,

while the acyl-ACP (shown here) provides an acyl moiety which varies between signaling systems. In

this example, the synthase LasI produces *N*-3-oxo-dodecanoyl-L-homoserine lactone (3OC12-HSL).

Covarying residues		
Synthase amino acid position	Receptor amino acid position	Top GREMLIN score
142	127	0.109
145	36	0.103
102	38	0.100
145	61	0.096
157	129	0.096
102	61	0.092
140	130	0.091
103	91	0.091
152	36	0.090
144	76	0.090

Randomized control		
Synthase amino acid position	Receptor amino acid position	GREMLIN score
137	203	0.080
123	175	0.079
183	28	0.070
62	58	0.068
38	102	0.067
93	114	0.065
110	151	0.066
180	138	0.065
175	193	0.065
183	59	0.065

Fig. 2 | Covarying residues identified in LasI/R. a) Top-scoring covarying residues in LasI (synthase)

and LasR (receptor) along with the top GREMLIN score (with APC) for each residue pair based an

integrated analysis of the Las and Lux systems. **b)** Top-scoring residues in a randomized control,

mapped onto LasI/R, along with the GREMLIN score (with APC) for each pair. **c)** Top-scoring

covarying residues mapped onto LasR (covarying residues in orange, 3OC12-HSL in silver; PDB 3IX3)

and **e)** LasI (covarying residues in blue; PDB 1RO5). Top-scoring residues in the randomized control are

mapped onto **d)** LasR and **f)** LasI as in panels **c** and **e**.

Fig. 3 | Activity of LasR mutants. Activity of chromosomal *lasR* mutants in *P. aeruginosa* PAO-SC4 pPROBE-P$_{rsaL}$ in response to **a)** 3OC12-HSL, **b)** *N*-dodecanoyl-L-homoserine lactone (C12-HSL), **c)** *N*-3-oxo-tetradecanoyl-L-homoserine lactone (3OC14-HSL) or **d)** *N*-3-oxo-hexadecanoyl-L-homoserine lactone (3OC16-HSL). Indicated mutations are amino acid substitutions. Wild-type (WT) is shown in black, LasR^{A127L} in purple, and LasR^{R61L} in blue. The horizontal axis indicates AHL concentration. LasR activity is reported on the vertical axis as relative fluorescence units normalized by optical density at 600 nm (RFU/OD x 1,000). Data are the mean and standard deviation of three biological replicates and are representative of three independent experiments. **e)** Concentration for half-maximal activation (EC$_{50}$) of 3OC12-HSL for *P. aeruginosa* PAO-SC4 LasR mutants calculated from data in Supplementary Fig. 3. Data are the mean and SEM of three (mutants) or four (wild-type) independent experiments. Sensitivity of the mutants compared to LasR^{WT} is calculated by dividing WT EC$_{50}$ by mutant EC$_{50}$.

Fig. 4 | Activity of LasI mutants. HPLC analysis of radiolabeled AHLs extracted from *P. aeruginosa*

PAO-SC4 harboring **a)** pJN-RBSlasI[WT] or **b)** pJN-RBSlasI[L157W]. The horizontal axis denotes the HPLC

fraction number (fractions 1-14 are not shown). The methanol gradient is indicated as a dashed line

plotted on the right vertical axis. The left vertical axis indicates the amount of radioactivity (counts per

minute (cpm)) in each fraction. Data are representative of two (L157W) or three (WT) independent

experiments. Arrow 1 indicates the fraction in which 3OC8-HSL elutes, arrow 2 indicates the fraction in

which 3OC10-HSL elutes, and arrow 3 indicates the fraction in which 3OC12-HSL elutes.

Fig. 5 | Reprogramming LasI/R selectivity. a) Residues mutated in LasR shown as orange sticks in the LasR structure (PDB 3IX3). 3OC12-HSL is shown in grey. **b)** LasR activity in response to 3OC10-HSL measured in *E. coli* harboring pJNL (wild-type, WT, or with indicated mutations) and pPROBE-P$_{rsaL}$. Data are the mean and standard deviation of three biological replicates and are representative of three independent experiments. **c)** Concentration of half maximal activity (EC$_{50}$) of 3OC10-HSL for LasR, calculated from data shown in panel **b**. Data are mean and SEM. Sensitivity of mutants compared to LasR^{WT} is calculated by dividing the WT EC$_{50}$ by mutant EC$_{50}$. **d)** Residues mutated in LasI shown as blue sticks in the LasI structure (PDB 1ROH). **e)** Relative amount of AHLs produced by *P. aeruginosa* PAO-SC4 harboring pJN-RBSlasI (WT or with the indicated mutations) or pJN-RBSmupI. Ratios were calculated from HPLC data shown in Supplementary Fig. 6. Bars show mean and standard deviation. The dashed line indicates equal production of 3OC10-HSL and 3OC12-HSL.