Interactions between Diameter, Myelination and the Na/K pump Affect Axonal Resilience to High Frequency Spiking

Yunliang Zang & Eve Marder[*]

Volen Center and Department of Biology, Brandeis University, Waltham, MA 02454, USA

[*]marder@brandeis.edu

Keywords: computational model; action potential; node of Ranvier; electrogenic pumps; sodium dynamics

Running title: Axonal resilience to high frequency firing

Abstract

Axons reliably conduct action potentials between neurons and/or other targets. Axons have widely variable diameters and can be myelinated or unmyelinated. Although the effect of these factors on propagation speed is well studied, how they constrain axonal resilience to high frequency spiking is incompletely understood. Maximal firing frequencies range from ~ 1 Hz to > 300 Hz across

neurons, but the process by which Na/K pumps counteract Na^+ influx is slow, and it is unclear the

extent to which slow Na^+ removal is compatible with high frequency spiking. Modeling the process

of Na^+ removal in unmyelinated and myelinated axons shows that both increasing diameter and

myelination slow down $[Na^+]$ accumulation and increase axonal resilience to high frequency

spiking. Increasing pump density alleviates $[Na^+]$ accumulation, but can paradoxically reduce the

resilience. We speculate that $[Na^+]$ accumulation may contribute to fatigue after continuous high

frequency firing.

Significance

The reliability of spike propagation in axons is determined by complex interactions between ionic

currents, ion pumps and morphological properties. We use compartment-based modeling to

uncover that interactions of diameter, myelination and the Na/K pump determine axonal resilience

to high frequency spiking. The Na/K pump can play a double-edged sword role in affecting axonal

excitability. Our findings suggest that slow sodium removal influences axonal resilience to high

frequency spiking, and may also contribute to fatigue.

Introduction

Axons usually reliably conduct information (spikes) to other neurons, muscles and glands. The largest diameter axons are found in some invertebrates, where the squid giant axon and arthropod giant fibers are parts of rapid escape systems. In the mammalian nervous system, axon diameters can differ by a factor of > 100. Some are covered with myelin sheaths but others are not [1]. Both myelination and large axon diameter increase spike propagation speed [2, 3], and the latter factor can also increase axonal resilience to noise perturbation [4].

Neuronal firing frequency is not static, but variations in firing rates are used to encode a wide range of signals such as inputs from sensory organs and command signals from motor cortex. The maximal firing frequency, critical for neuronal functional capacity, ranges from ~ 1 Hz to > 300 Hz [5-8]. However, it remains unclear how axonal physical properties constrain axonal resilience to high frequency firing. During action potentials, Na^+ flows into axons and then ATP is used by Na/K pumps to pump out the excess Na^+. Given the limited energy supply in the brain, evolutionary strategies including shortening spike propagation distance, reducing Na^+- and K^+-current overlap, and sparse coding, etc. have helped minimize energy expenditure [9-13]. However, the observation that high frequency spiking tends to occur in larger diameter axons is confusing because it exerts a heavy burden on the energy supply in the brain [5, 14, 15]. Additionally, the process by which Na/K pumps remove Na^+ is slow [16-22]. In neurons, it takes seconds or even minutes to remove the excess Na^+ after Na^+ elevation [16-21]. It remains unknown how these directly correlated processes, fast spiking and slow Na^+ removal, interact in the control of reliable spike generation and propagation. Consequently, by modeling the process of Na^+ removal in

54 unmyelinated and myelinated axons, we systematically explored the effects of diameter,

55 myelination and Na/K pump density on spiking reliability.

Results

Fig. 1. Spike-triggered [Na$^+$] rise and decay in unmyelinated and myelinated axon models. (A)
Schematic of the unmyelinated (left) and the myelinated (right) axon models. Black traces show
spike waveforms recorded at the center of each model. (B) Spike-triggered [Na$^+$] rise and decay
in the unmyelinated (left) and the myelinated (right) axon models. In the unmyelinated axon model,
[Na$^+$] was recorded at the center of the axon. In the myelinated axon model, [Na$^+$] at the central
node of Ranvier and the myelinated compartment (10.5 μm far) are shown. Initial parts of [Na$^+$]
changes are enlarged in the inset. The Na/K pump density is 0.5 pmol/cm^2, referred to as baseline
condition hereafter. (C) The effects of diameter and pump density on [Na$^+$] rise and decay in the
unmyelinated (left) and the myelinated (right) axon models. Note the very different time scales for
results in the two models.

Different [Na$^+$] Decay Dynamics in Unmyelinated and Myelinated Axons. We implemented

multicompartment model of axons in NEURON (23). Both unmyelinated and myelinated axon

models include I_{Na}, I_{KD}, I_{KA}, I_{leak} and the Na/K pump. All channels and pumps were evenly spread

in the unmyelinated axon model and at the nodes of Ranvier in the myelinated axon model. For

the myelinated compartments, there is a significantly lower density of I_{leak} and C_m to simulate the

effect of insulation (Methods). The model structures are shown in Fig. 1A.

Spikes propagate rapidly in both models. In the unmyelinated axon model, the propagation

increases from 0.16 m/s to 0.39 m/s when the axonal diameter increases from 0.2 μm to 1 μm. In

the myelinated axon model, the speed increases from 2.5 m/s to 10 m/s accordingly. Upon the

arrival of spikes, Na^+ channels open and extracellular Na^+ flows into axons to raise the intracellular

$[Na^+]$ immediately (Fig. 1B). Then K^+ currents are activated to repolarize the spikes. Compared

with the other ionic currents, the magnitude of the Na/K pump current is extremely small during a

single spike, but it decays very slowly. Eventually, the Na/K pump gradually removes Na^+ from

the axon, leading to a slow, many second, $[Na^+]$ decay in the unmyelinated axon model (Fig. 1B,

left). In contrast, in the myelinated axon model, the spike-triggered $[Na^+]$ elevation at the node of

Ranvier decays on the time scale of hundreds of ms (Fig. 1B, right). The Na/K pump densities in

two models are the same and thus the quick decay is not a result of super-efficient pumps. In the

myelinated axon model, Na^+ channels are placed only in the nodes of Ranvier and there is no Na^+

influx in myelinated compartments. But, the slightly elevated $[Na^+]$ in the myelinated compartment

and the rapid decay of nodal $[Na^+]$ are due to the axial diffusion of Na^+ into neighboring myelinated

compartments, rather than immediate pump action.

Because of the large diffusion coefficient (24) and the lack of buffering (16), $[Na^+]$ in the radial

direction becomes rapidly homogeneous. Thus, spike-triggered $[Na^+]$ elevation can be well

predicted by the surface-to-volume ratio (SVR ∝ 1/diameter). In both models, spike-triggered $[Na^+]$

elevation decreases with diameter (Fig. 1C). Increasing the pump density speeds up the $[Na^+]$

decay in the unmyelinated axon model, but has no obvious effect on the nodal [Na$^+$] decay in the

myelinated axon model, further confirming that the quick decay is not a result of the Na/K pump.

[Na$^+$] Accumulation in the Unmyelinated Axon at High Firing Frequencies. Compared with

the basal level before spiking ($\sim$ 10 mM), the [Na$^+$] elevation caused by a single spike is small and

insufficient to affect axonal excitability (Fig. 1B). However, neurons can fire at high frequencies

when animals receive stimuli or undergo behavior changes (7, 25-27). With continuous high

frequency spiking, [Na$^+$] can accumulate if subsequent spikes occur before the complete recovery

of a preceding spike elicited [Na$^+$] rise. In the unmyelinated axon model, [Na$^+$] significantly

accumulates within just 100 spikes, and the accumulation speed increases with firing frequency

because of a faster Na$^+$ influx rate (Fig. 2A, above). Larger SVRs also make [Na$^+$] accumulate

faster in thin axons. In parallel with [Na$^+$] accumulation, the Na$^+$ reversal potential (driving force

of Na$^+$ current) becomes more negative and the spike amplitude decreases (Fig. 2A, middle and

bottom).

Fig. 2. The effects of diameter and pump density on [Na$^+$] accumulation and resilience to high frequency spiking in the unmyelinated axon model. (A) The course of [Na$^+$] accumulation (above), Na$^+$ reversal potential decrease (middle) and membrane potential peak reduction (bottom) in models with different diameters (0.2, 0.6 and 1.0 μm) and firing frequencies (1, 40 and 90 Hz). Spike threshold is -30 mV. Na/K pump density is 0.5 pmol/cm^2 (baseline). (B) Steady state [Na$^+$] accumulation (above) and Na$^+$ reversal potential reduction (bottom) at different firing frequencies. Symbol codes diameter and color codes pump density. (C) The effect of pump density on the resilience to high frequency spiking, measured by the number of spikes required to cause spike failure (nspike_fail), when firing at 70 and 80 Hz. 'shunt' corresponds to neutral pumps, meaning pumps remove Na$^+$, but carry 0 net current. The absence of data points means spikes don't fail. (D) The course of Na$^+$ reversal potential decrease (left) and outward pump current increase (right) with different pump densities. Diameter is 0.6 μm and firing frequency is 80 Hz.

When the model fires over a sufficiently long period, Na$^+$ influx rate decreases and efflux rate

increases because of elevated [Na$^+$]. Eventually, Na$^+$ influx rate and efflux rate balance each other

and [Na$^+$] reaches a steady state. The steady state [Na$^+$] increases with firing frequency until the

occurrence of spike failure at high frequencies (Fig. 2B). Within the low frequency range, the

steady state $[Na^+]$ is independent of diameter despite different accumulation speeds. The rate of

Na^+ influx per surface area is only determined by the firing frequency because of identical Na^+

channel density. The Na/K pump-caused net Na^+ efflux rate is determined by

$-3k_1[pump][Na^+]^3 + 3k_2[pumpNa]$ (Methods). Because pump density is also identical, the

model needs identical $[Na^+]$ to match influx rate and reach a steady state irrespective of diameters.

Na^+ influx rate becomes diameter dependent when spiking fails at high frequencies. Accordingly,

different $[Na^+]$ are required to reach a steady state. Increasing pump density can reduce $[Na^+]$

accumulation and alleviate the reversal potential decrease, because the Na^+ efflux rate becomes

larger (Fig. 2B).

As a result of incomplete recovery from inactivation of Na^+ channels and the accumulated $[Na^+]$,

spikes fail intermittently at high frequencies. This gradual occurrence of failure is alleviated in the

absence of $[Na^+]$ accumulation (data not shown). An example of spike failure is shown in Fig. 2A

when the models fire at 90 Hz. $[Na^+]$ accumulates slowly in thick axons and in thick axons it

always requires more spikes to trigger spike failure than in thin axons, demonstrating that thick

axons are more resilient to high frequency firing (Fig. 2A, C). At higher frequencies, $[Na^+]$

accumulates faster and Na^+ channels recover less from inactivation. Consequently, fewer spikes

are required to trigger spike failure and accordingly the resilience (equivalent to excitability)

decreases (Fig. 2C).

Surprisingly, although increasing pump density alleviates $[Na^+]$ accumulation, it doesn't

consistently increase the resilience (Fig. 2C). We tested the pump density in the range of 1–10

times of the baseline level (0.5 pmol/cm^2). Increasing pump density doesn't obviously affect the

resilience of the 0.2-μm-thick axon model. Although the resilience of 0.6- and 1.0-μm-thick axons

is increased by pump density at 70 Hz, their resilience decreases with pump density at 80 Hz.

When removing excess Na^+, the pumps carry a negative current. More pumps alleviate $[Na^+]$

accumulation and the resultant reversal potential decrease, but they also provide a larger outward

current to decrease the axonal excitability directly (Fig. 2D). To confirm that the larger outward

current accounts for the paradoxical effect of pump density on the resilience, we 'shunt' the pump

current (net current becomes neutral instead of outward, but still pumps out Na^+). After shunting

this current, increasing pump density consistently increases the resilience (an example at 80 Hz is

given in Fig. 2C). Thus, the effect of pump density on the resilience depends on the balance

between $[Na^+]$ accumulation and increased outward current.

Myelination Slows down $[Na^+]$ accumulation. In contrast to the slow decay of $[Na^+]$ in the

unmyelinated axon, $[Na^+]$ at the node of Ranvier in the myelinated axon shows a quick decay (Fig.

1B). Can nodal $[Na^+]$ accumulate at high firing frequencies? It does accumulate to reduce the Na^+

reversal potential and spike amplitude peak in parallel (Fig. 3A). As explained previously, the

quick decay of nodal $[Na^+]$ is a result of dilution, rather than being removed: nodal Na^+ influx

quickly diffuses into neighboring myelinated compartments. The diluted $[Na^+]$ is still higher than

the pre-spiking level (Fig. 3A at 1 Hz, inset). Nodal Na^+ influx not only needs to fill the nodes of

Ranvier, but also the neighboring myelinated compartments, which is equivalent to decreasing the

SVR. Consequently, nodal $[Na^+]$ still accumulates, but at a much slower speed. Compared with

the unmyelinated axon model, the $[Na^+]$ elevation and the Na^+ reversal potential decrease are

negligible for 100 spikes in the myelinated axon model (Fig. 3B). Instead, it requires thousands of

spikes to significantly change nodal $[Na^+]$. Therefore, by acting like an 'anti-flood water reservoir',

myelin slows down the nodal $[Na^+]$ accumulation and the excitability reduction.

Similar to the unmyelinated axon model, nodal [Na⁺] accumulates faster at higher firing

frequencies (Fig, 3A). Although the [Na⁺] accumulation speed decreases with diameter, the steady

Fig. 3. Myelination slows down $[Na^+]$ accumulation to increase the axonal resilience to high
frequency spiking. (A) The course of $[Na^+]$ accumulation (above), Na^+ reversal potential decrease
(middle) and membrane potential peak reduction (bottom) at the node of Ranvier in the myelinated
axon model with different diameters (0.2, 0.6 and 1.0 μm) and firing frequencies (1, 40 and 90 Hz).
Fig. 2A shows 100 spike caused changes, but here shows 5000 spike caused changes. At 1 Hz, the
$[Na^+]$ in myelinated compartment (10.5 μm far) is also enlarged and shown in the inset when the
diameter is 0.2 μm. Spike threshold is -30 mV. (B) 100-spike caused $[Na^+]$ accumulation (left) and
Na^+ reversal potential decrease (right) in the unmyelinated axon and the myelinated axon (node of
Ranvier), with diameter of 0.2 μm. In A and B, the Na/K pump density is 0.5 pmol/cm² (baseline).
(C) Steady state $[Na^+]$ elevation (above) and Na^+ reversal potential decrease (bottom) at different
firing frequencies. Symbol codes diameter and color codes pump density. (D) The effect of pump
density on the axonal resilience to high frequency spiking, measured by the number of spikes
required to cause spike failure (nspike_fail), when firing at 90 and 100 Hz. 'shunt' corresponds to
neutral pumps. The absence of data points means spikes don't fail.

190 state [Na$^+$] is diameter independent until the occurrence of intermittent spike failure at high

191 frequencies (Fig. 3C). The [Na$^+$] accumulation and the Na$^+$ reversal potential decrease can be

192 alleviated by increasing pump density. Thick axons are consistently more resilient to high

193 frequency spiking (Fig. 3D). Still, increasing pump density doesn't consistently increase resilience,

194 but a consistent increase can be achieved by shunting the pump current.

Discussion

Neuronal firing frequency often varies over a broad range to encode a wide range of signals. For

a given neuron, its maximal firing frequency is usually much higher than the speed at which

intracellular Na$^+$ is pumped out of the cell. Therefore, while short periods of high frequency firing

should present no problem for most neurons, continuous high frequency spiking could result in

compromised function, as intracellular [Na$^+$] accumulates. Hence, we explore the dynamic

relationships between fast spiking and slow Na$^+$ removal for reliable spike generation and

propagation.

The importance of the Na/K ATPase in regulating neuronal activities has been called into

attention in recent years (28). It can set neuronal excitability and contribute to maintaining

synchronous epileptiform bursting in developing CA3 neurons (29). By interacting with h-current,

it regulates bursting activity in the leech heartbeat central pattern generator (30). There have also

been theoretical efforts to study the pump's role in controlling the transition between different

firing patterns (31-33) and the transition between normal and pathological brain states (34-36).

However, to the best of our knowledge, how the pump interacts with axon diameter and myelin to

remove [Na$^+$] influx and affect axonal excitability hasn't been explored previously.

Axonal morphology can exert a critical constraint on a neuron's maximal firing frequency.

Thick and myelinated axons are more resilient to high frequency spiking, because of their slow

[Na$^+$] accumulation and the resultant modest excitability reduction. In thin unmyelinated axons,

spiking may still be reliable at low frequencies owing to limited [Na$^+$] accumulation and

excitability reduction, but the reliability may be severely reduced when the axons fire at high

frequencies. This may explain why high frequency spiking tends to occur in thick axons, despite a

higher energetical cost of information transmission (5, 14).

Increasing the Na/K pump density to remove Na$^+$ more efficiently is expected to facilitate axonal

resilience to high frequency spiking. Existing data argue that Na$^+$ removal after spiking is slow.

For example, in axon initial segments of L5 pyramidal neurons, as a result of Na$^+$ diffusing into

the soma and myelin wrapped compartments, elevated [Na$^+$] requires several seconds to roughly

recover (16). In mitral cell tuft dendrites, after a short period of high frequency stimulation, it takes

minutes for the Na/K ATPase to pump out the Na$^+$ influx (17). This slow [Na$^+$] removal may be

functionally important because it enables integration and binding of multiple chemosensory stimuli

over a prolonged time scale (37). We used pump densities that should fall into a reasonable

biological range, and the [Na$^+$] accumulation at high firing frequencies shown here is consistent

with recent Na$^+$ imaging data (17, 20). Importantly, according to these simulation results,

increasing pumping efficiency is a two-edged sword. On the one hand, it does alleviate [Na$^+$]

accumulation; on the other hand, it reduces axonal excitability by carrying a larger outward current.

This work illustrates the limitations of using conventional frequency-current curves generated

with a short-term current injection to estimate the reliability of spike propagation (38, 39). Neither

of these measures reflects the accommodation of neuronal excitability after continuous firing

caused changes such as $[Na^+]$ (explored in this work) and $[Ca^{2+}]$ accumulation (showing a biphasic

decay (40, 41)).

In the models explored here, $[Na^+]$ accumulation caused spike failure starting at 60 Hz in

unmyelinated axons. Specifically, when the axon diameter was 0.2 μm with the pump density of

0.5 pmol/cm^2, 1.1 sec continuous firing (66 spikes) elicited spike failure because of $[Na^+]$

accumulation. If only several spikes are required to encode a signal at this frequency, then $[Na^+]$

accumulation is unlikely to be an issue. However, with less recovery from inactivation of Na^+

channels at higher frequencies, spike failure reduces to 0.5 sec at 70 Hz (35 spikes), 0.19 sec at 80

Hz (15 spikes), 0.06 sec at 90 Hz (5 spikes) and 0.03 sec at 100 Hz (3 spikes).

In these modeled myelinated axons, spike failure was seen starting at 90 Hz. Specifically, when

the axon diameter was 0.2 μm with the pump density of 0.5 pmol/cm^2, failure only occurred after

continuous firing for 54 sec at 90 Hz (4821 spikes) and 8.7 sec at 100 Hz (873 spikes). Given that

myelinated axons are usually much thicker (1, 42), spike propagation in myelinated axons should

be extremely robust to fast firing caused $[Na^+]$ accumulation under usual physiological conditions.

However, under conditions requiring neurons to fire for extended time at high frequencies, $[Na^+]$

accumulation may still compromise the function of axons. Of course, the frequency threshold and

the number of spikes required to cause spike failure will depend on the specifics of the expression

of neuronal conductances, and will likely show variability as a consequence of variable channel

densities (43). Additionally, intrinsic channel noise can cause spike failure (4), and the normal

function of the Na/K ATPases requires ATP, the supply of which has a limit (35). Both a shortage

of ATP after continuous firing and channel noise can further reduce axonal excitability.

Interestingly, a recent study observed intermittent spike propagation failure *in vivo* in dorsal root

ganglia of mice during sustained tactile stimuli (27). [Na$^+$] accumulation may be another important

factor causing the failure apart from the channel noise.

Axon diameters can be up to 20 μm in the human peripheral nervous system and 1.5 mm in

squid giant axons (1). For larger diameter axons, the conclusions still hold, because increasing

diameter just further slows down [Na$^+$] accumulation. The localization of Na/K pumps in

myelinated axons is still controversial (1, 44, 45). The paradoxical effect of pump density on the

axonal resilience to high frequency spiking is preserved even if pumps are distributed between

nodes of Ranvier (data not shown), as additive pumps increase the leakiness of the internodal

compartments to reduce nodal excitability.

Fatigue is a deeply mysterious property that we all experience, and undoubtedly involves

numerous cellular mechanisms (46). Significant [Na$^+$] accumulation has been observed in muscle

cells and contributes to muscle fatigue after intense exercise (47, 48). Likewise, it is interesting to

speculate that [Na$^+$] accumulation that occurs as a consequence of concerted and sustained

neuronal activity could contribute to fatigue as we experience it.

ACKNOWLEDGMENTS. The authors would like to thank the constructive suggestions from
the members of Marder lab. This work was supported by the grant R35NS097343.

References

1.   S. G. Waxman, J. D. Kocsis, P. K. Stys, *The Axon: Structure, Function, and Pathophysiology*
(Oxford University Press, 1995).
2.   D. Debanne, E. Campanac, A. Bialowas, E. Carlier, G. Alcaraz, Axon physiology. Physiol Rev
91, 555-602 (2011).
3.   H. Schmidt, T. R. Knosche, Action potential propagation and synchronisation in
myelinated axons. PLoS Comput Biol **15**, e1007004 (2019).
4.   A. A. Faisal, J. A. White, S. B. Laughlin, Ion-channel noise places limits on the
miniaturization of the brain's wiring. Curr Biol **15**, 1143-1149 (2005).
5.   J. A. Perge, J. E. Niven, E. Mugnaini, V. Balasubramanian, P. Sterling, Why do axons differ
in caliber? J Neurosci **32**, 626-638 (2012).

6.   B. Wang *et al.*, Firing Frequency Maxima of Fast-Spiking Neurons in Human, Monkey, and
Mouse Neocortex. Front Cell Neurosci **10**, 239 (2016).
7.   S. Chen, G. J. Augustine, P. Chadderton, Serial processing of kinematic signals by
cerebellar circuitry during voluntary whisking. Nat Commun **8**, 232 (2017).
8.   A. Hasenstaub, S. Otte, E. Callaway, T. J. Sejnowski, Metabolic cost as a unifying principle
governing neuronal biophysics. Proc Natl Acad Sci USA **107**, 12329-12334 (2010).
9.   S. B. Laughlin, Energy as a constraint on the coding and processing of sensory information.
Curr Opin Neurobiol **11**, 475-480 (2001).
10.   W. B. Levy, R. A. Baxter, Energy efficient neural codes. Neural Comput **8**, 531-543 (1996).
11.   H. Alle, A. Roth, J. R. Geiger, Energy-efficient action potentials in hippocampal mossy
fibers. Science **325**, 1405-1408 (2009).
12.   F. J. Heras, J. Anderson, S. B. Laughlin, J. E. Niven, Voltage-dependent K^+ channels improve
the energy efficiency of signalling in blowfly photoreceptors. J R Soc Interface **14** (2017).
13.   H. Hu, F. C. Roth, D. Vandael, P. Jonas, Complementary Tuning of Na^+ and K^+ Channel
Gating Underlies Fast and Energy-Efficient Action Potentials in GABAergic Interneuron
Axons. Neuron **98**, 156-165 e156 (2018).
14.   C. Howarth, P. Gleeson, D. Attwell, Updated energy budgets for neural computation in
the neocortex and cerebellum. J Cereb Blood Flow Metab **32**, 1222-1232 (2012).
15.   R. Nortley, D. Attwell, Control of brain energy supply by astrocytes. Curr Opin Neurobiol
47, 80-85 (2017).
16.   I. A. Fleidervish, N. Lasser-Ross, M. J. Gutnick, W. N. Ross, Na^+ imaging reveals little
difference in action potential-evoked Na^+ influx between axon and soma. Nat Neurosci
13, 852-860 (2010).
17.   A. Zylbertal, A. Kahan, Y. Ben-Shaul, Y. Yarom, S. Wagner, Prolonged Intracellular Na^+
Dynamics Govern Electrical Activity in Accessory Olfactory Bulb Mitral Cells. PLoS Biol **13**,
e1002319 (2015).
18.   A. Zylbertal, Y. Yarom, S. Wagner, The Slow Dynamics of Intracellular Sodium
Concentration Increase the Time Window of Neuronal Integration: A Simulation Study.
Front Comput Neurosci **11**, 85 (2017).
19.   D. Ziemens, F. Oschmann, N. J. Gerkau, C. R. Rose, Heterogeneity of Activity-Induced
Sodium Transients between Astrocytes of the Mouse Hippocampus and Neocortex:
Mechanisms and Consequences. J Neurosci **39**, 2620-2634 (2019).
20.   T. Ona-Jodar, N. J. Gerkau, S. Sara Aghvami, C. R. Rose, V. Egger, Two-Photon Na^+ Imaging
Reports Somatically Evoked Action Potentials in Rat Olfactory Bulb Mitral and Granule Cell
Neurites. Front Cell Neurosci **11**, 50 (2017).
21.   B. Moshrefi-Ravasdjani, D. Ziemens, N. Pape, M. Färfers, C. R. Rose, Action Potential Firing
Induces Sodium Transients in Macroglial Cells of the Mouse Corpus Callosum. Neuroglia
1, 106-125 (2018).
22.   Y. Zang *et al.*, Theoretical investigation of the mechanism of heart failure using a canine
ventricular cell model: especially the role of up-regulated CaMKII and SR Ca^{2+} leak. J Mol
Cell Cardiol **56**, 34-43 (2013).
23.   M. L. Hines, N. T. Carnevale, The NEURON simulation environment. Neural Comput **9**,
1179-1209 (1997).

24. M. J. Kushmerick, R. J. Podolsky, Ionic mobility in muscle cells. Science **166**, 1297-1298
(1969).
25. L. S. Tang *et al.*, Precise temperature compensation of phase in a rhythmic motor pattern.
PLoS Biol **8**, e1000469 (2010).
26. J. C. Athilingam, R. Ben-Shalom, C. M. Keeshen, V. S. Sohal, K. J. Bender, Serotonin
enhances excitability and gamma frequency temporal integration in mouse prefrontal
fast-spiking interneurons. Elife **6**, e31991 (2017).
27. D. Al-Basha, S. A. Prescott, Intermittent Failure of Spike Propagation in Primary Afferent
Neurons during Tactile Stimulation. J Neurosci **39**, 9927-9939 (2019).
28. M. D. Forrest, The sodium-potassium pump is an information processing element in brain
computation. Frontiers in Physiology **5** (2014).
29. L. R. Shao, R. Janicot, C. E. Stafstrom, Na^+-K^+-ATPase functions in the developing
hippocampus: regional differences in CA1 and CA3 neuronal excitability and role in
epileptiform network bursting. J Neurophysiol **125**, 1-11 (2021).
30. D. Kueh, W. H. Barnett, G. S. Cymbalyuk, R. L. Calabrese, Na^+/K^+ pump interacts with the
h-current to control bursting activity in central pattern generator neurons of leeches. Elife
5, e19322 (2016).
31. M. D. Forrest, M. J. Wall, D. A. Press, J. Feng, The sodium-potassium pump controls the
intrinsic firing of the cerebellar Purkinje neuron. PLoS One **7**, e51169 (2012).
32. A. O. Komendantov, C. C. Canavier, Electrical coupling between model midbrain
dopamine neurons: effects on firing pattern and synchrony. J Neurophysiol **87**, 1526-1541
(2002).
33. C. C. Canavier, Sodium dynamics underlying burst firing and putative mechanisms for the
regulation of the firing pattern in midbrain dopamine neurons: a computational approach.
J Comput Neurosci **6**, 49-69 (1999).
34. G. P. Krishnan, G. Filatov, A. Shilnikov, M. Bazhenov, Electrogenic properties of the Na^+/K^+
ATPase control transitions between normal and pathological brain states. J Neurophysiol
113, 3356-3374 (2015).
35. G. Le Masson, S. Przedborski, L. F. Abbott, A computational model of motor neuron
degeneration. Neuron **83**, 975-988 (2014).
36. J. S. Coggan, G. K. Ocker, T. J. Sejnowski, S. A. Prescott, Explaining pathological changes in
axonal excitability through dynamical analysis of conductance-based models. J Neural Eng
8, 065002 (2011).
37. A. Zylbertal, Y. Yarom, S. Wagner, Synchronous Infra-Slow Bursting in the Mouse
Accessory Olfactory Bulb Emerge from Interplay between Intrinsic Neuronal Dynamics
and Network Connectivity. J Neurosci **37**, 2656-2672 (2017).
38. N. W. Gouwens *et al.*, Classification of electrophysiological and morphological neuron
types in the mouse visual cortex. Nat Neurosci **22**, 1182-1195 (2019).
39. A. Foust, M. Popovic, D. Zecevic, D. A. McCormick, Action Potentials Initiate in the Axon
Initial Segment and Propagate through Axon Collaterals Reliably in Cerebellar Purkinje
Neurons. The Journal of Neuroscience **30**, 6891-6902 (2010).
40. H. Schmidt, K. M. Stiefel, P. Racay, B. Schwaller, J. Eilers, Mutational analysis of dendritic
Ca^{2+} kinetics in rodent Purkinje cells: role of parvalbumin and calbindin D28k. J Physiol
551, 13-32 (2003).

41. Y. Zang, S. Dieudonne, E. De Schutter, Voltage- and Branch-Specific Climbing Fiber
Responses in Purkinje Cells. Cell Rep **24**, 1536-1549 (2018).
42. S. Lee *et al.*, A culture system to study oligodendrocyte myelination processes using
engineered nanofibers. Nat Methods **9**, 917-922 (2012).
43. D. J. Schulz, J. M. Goaillard, E. Marder, Variable channel expression in identified single and
electrically coupled neurons in different animals. Nat Neurosci **9**, 356-362 (2006).
44. S. G. Waxman, J. M. Ritchie, Molecular dissection of the myelinated axon. Ann Neurol **33**,
121-136 (1993).
45. E. A. Young *et al.*, Imaging correlates of decreased axonal Na$^+$/K$^+$ ATPase in chronic
multiple sclerosis lesions. Ann Neurol **63**, 428-435 (2008).
46. J. J. Wan, Z. Qin, P. Y. Wang, Y. Sun, X. Liu, Muscle fatigue: general understanding and
treatment. Exp Mol Med **49**, e384 (2017).
47. S. P. Cairns, M. I. Lindinger, Do multiple ionic interactions contribute to skeletal muscle
fatigue? J Physiol **586**, 4039-4054 (2008).
48. T. Clausen, Quantification of Na$^+$,K$^+$ pumps and their transport rate in skeletal muscle:
functional significance. J Gen Physiol **142**, 327-345 (2013).
49. Y. Zhang, D. Bucher, F. Nadim, Ionic mechanisms underlying history-dependence of
conduction delay in an unmyelinated axon. Elife **6**, e25382 (2017).
387

Methods

Axonal Morphology. We modeled a segment of axon with the length of 303 μm. The model was

implemented in NEURON (23). The model code will be available from ModelDB. The simulations

used an integration step of 0.01 ms. The unmyelinated model was discretized into 15 segments in

the longitudinal direction. Vm and [Na$^+$] were recorded at the central segment. For the myelinated

axon model, there are 3 nodes of Ranvier flanked with 4 myelinated compartments (Figure 1). The

lengths of the nodes of Ranvier are 1 μm (nseg = 1). The middle two myelinated compartments

are 100 μm long each (nseg = 5 each) and the other two myelinated compartments are 50 μm each

(nseg = 3 each). Vm and [Na$^+$] were recorded at the central node and its neighboring myelinated

compartment. In the radial direction, there are 4 shells to model Na$^+$ diffusion (23). The Na$^+$

diffusion coefficient is 0.6 μm^2/ms (24) and there is no buffering (16), leading to a homogeneous

[Na$^+$] in the radial direction. In both peripheral and central nervous systems, unmyelinated axons

are usually thinner than myelinated axons. In the central nervous system, axons > 0.2 μm are

usually myelinated (1). To delineate the effect of both diameter and myelination on [Na$^+$]

accumulation, we chose the same range of diameters between 0.2–1.0 μm.

Ionic Channels and the Na/K pump. The model has a minimal set of ionic channels: fast Na

current (I_{Na}), delayed rectifier K current (I_{KD}), A-type K current (I_{KA}) and leak current (I_{leak}). They

have standard kinetic formulations in the form of $I = g_{max}m^p h^q (V - E)$; $\tau_m \dot{m} = m_\infty(V) -$

m; $\tau_h \dot{h} = h_\infty(V) - h$; g_{max} represents maximal conductance; m and h represent activation and

inactivation gates; p and q represent corresponding gate exponents; E represents reversal

potential; τ_m and τ_h represent corresponding time constants; All the channel kinetics were

modified from (49). For I_{Na}, $g_{max} = 0.015\ S/cm^2$; $p = 3$; $q = 1$; $m_\infty =$

$1/(1 + exp(-(V + 38)/8.5))$; $\tau_m = 0.132/(\cosh((V + 27)/7.5) + 0.003/(1 +$

$\exp(-(V + 27)/5)))$; $h_\infty = 1/(1 + exp(V + 47/6))$; $\tau_h = 10/cosh((V + 42)/15$; $E_{Na} =$

$RT/F * \ln([Na^+]_o/[Na^+]_i)$; $[Na^+]_o$ is fixed to be 140 mM; $[Na^+]_i$ (also written as [Na$^+$] in the

main text) was dynamically computed by I_{Na} and the Na/K pump; For I_{KD}, $g_{max} = 0.216\ S/cm^2$;

$p = 4$; $q = 0$; $m_\infty = (-0.01 * (V + 45.7)/(exp(-(V + 45.7)/10) - 1))/((-0.01 * (V +$

$45.7)/(exp(-(V + 45.7)/10) - 1)) + 0.125 * exp(-(V + 55.7)/80))$; $\tau_m = 1/((-0.01 *$

$(V + 45.7)/(exp(-(V + 45.7)/10) - 1)) + 0.125 * exp(-(V + 55.7)/80))$; $E_K =$ -70 mV;

For I_{KA}, $g_{max} = 0.02\ S/cm^2$; $p = 3$; $q = 1$; $m_\infty = (0.0761 * exp((V + 94.22)/31.84)/(1 +$

$exp((V + 1.17)/28.93)))^{1/3}$; $\tau_m = 0.3632 + 1.158/(1 + exp((V + 55.96)/20.12))$; $h_\infty =$

$1/(1 + exp(V + 53.3/14.54))^4$; $\tau_h = 1.24 + 2.678/(1 + exp((V + 50)/16.027))$; For

I_{leak}, $g_{max} = 1.25 \times 10^{-4}\ S/cm^2$; $p = 0$; $q = 0$; $E_{leak} =$ -65 mV; the Na/K pump model is

similar with that used in (17, 37), described by the following kinetic schemes: $[pump] + 3[Na^+]_i$

$\xleftrightarrow{k_1,k_2} [pumpNa]$; $[pumpNa] \xleftrightarrow{k_3,k_4} [pump] + 3[Na^+]_o$; $pump_{total} = [pump] + [pumpNa]$;

$k_1 = 2\ mM^{-3}msec^{-1}$; $k_2 = 0.001\ msec^{-1}$; $k_3 = 0.6\ msec^{-1}$; $k_4 = 0.437\ mM^{-3}msec^{-1}$;

By these parameters, the steady state [Na⁺] is ~ 10 mM when the axon models are silent.

For the unmyelinated axon model, all defined currents and the Na/K pump were evenly

distributed with C_m = 1 µF/cm² and R_a = 120 Ωcm. For the myelinated axon model, the ionic

currents and the Na/K pump were placed in the nodes of Ranvier (1, 44). For simplicity, we did

not model juxtaparanodal regions. At the nodes of Ranvier, C_m = 1 µF/cm²; On myelinated

compartments, there is only a smaller I_{leak} with g_{max} = 1.25×10⁻⁶ S/cm² and effective C_m = 0.01

µF/cm²; R_a is 120 Ωcm. In both the unmyelinated and the myelinated axon models, the current

densities are identical so that we could explore the role of diameter and myelination on Na⁺

accumulation. The $pump_{total}$ is in the range of 0.5–5 pmol/cm². To show the effect of outward

pump current on axonal excitability, in Fig. 2 and 3, we 'shunt' the pump current by making it

neural, which means it carries 0 net current. In the unmyelinated axon model, spikes were triggered

by stimulating the left end to -30 mV when Na current was blocked; In the myelinated axon model,

spikes were triggered by stimulating the first node of Ranvier to -30 mV when Na⁺ current was

blocked.