1

Main Manuscript for

Interactions among Diameter, Myelination and the Na/K pump Affect
Axonal Resilience to High Frequency Spiking

Yunliang Zang & Eve Marder*

Volen Center and Department of Biology, Brandeis University, Waltham, MA 02454, USA

*Eve Marder

Email: marder@brandeis.edu

Author Contributions: YZ and EM designed this research; YZ implemented model building, model
simulation, data analysis, figure generation; YZ and EM wrote and revised the manuscript.
12
13 **Competing Interest Statement:** The authors declare no competing financial interests
14
Classification: Major - Biological Sciences; minor - Neuroscience

Keywords: computational model; action potential; node of Ranvier; electrogenic pumps; sodium

dynamics

This PDF file includes:
Main Text
Figures 1 to 4
21
22

Abstract

Axons reliably conduct action potentials between neurons and/or other targets. Axons have widely variable diameters and can be myelinated or unmyelinated. Although the effect of these factors on propagation speed is well studied, how they constrain axonal resilience to high frequency spiking is incompletely understood. Maximal firing frequencies range from ~ 1 Hz to > 300 Hz across neurons, but the process by which Na/K pumps counteract Na^+ influx is slow, and it is unclear the extent to which slow Na^+ removal is compatible with high frequency spiking. Modeling the process of Na^+ removal shows that large diameter axons are more resilient to high frequency spikes than small diameter axons, because of their slow Na^+ accumulation. In myelinated axons, the myelinated compartments between nodes of Ranvier act as a 'reservoir' to slow Na^+ accumulation and increase the reliability of axonal propagation. We now find that slowing the activation of K^+ current can increase the Na^+ influx rate, and the effect of minimizing the overlap between Na^+- and K^+-currents on spike propagation resilience depends on complex interactions among diameter, myelination and the Na/K pump density. Our results suggest that, in neurons with different channel gating kinetic parameters, different strategies may be required to improve the reliability of axonal propagation.

Significance Statement

The reliability of spike propagation in axons is determined by complex interactions among ionic currents, ion pumps and morphological properties. We use compartment-based modeling to reveal that interactions of diameter, myelination and the Na/K pump determine the reliability of high frequency spike propagation. By acting as a 'reservoir' of nodal Na^+ influx, myelinated compartments efficiently increase propagation reliability. Although spike broadening was thought to oppose fast spiking, its effect on spike propagation is complicated, depending on the balance of Na^+ channel inactivation gate recovery, Na^+ influx and axial charge. Our findings suggest that slow Na^+ removal influences axonal resilience to high frequency spike propagation, and that different strategies may be required to overcome this constraint in different neurons.

Main Text

Introduction

Axons usually reliably conduct information (spikes) to other neurons, muscles and glands. The largest diameter axons are found in some invertebrates, where the squid giant axon and arthropod giant fibers are parts of rapid escape systems. In the mammalian nervous system, axon diameters can differ by a factor of > 100. Some are covered with myelin sheaths but others are not [1]. Both myelination and large axon diameter increase spike propagation speed [2, 3], and the latter factor can also increase axonal resilience to noise perturbation [4].

Neuronal firing frequency is not static, but variations in firing rates are used to encode a wide range of signals such as inputs from sensory organs and command signals from motor cortex. The maximal firing frequency, critical for neuronal functional capacity, ranges from ~ 1 Hz to > 300 Hz [5-8]. However, it remains unclear how axonal physical properties constrain axonal resilience to high frequency firing. During action potentials, Na^+ flows into axons and then ATP is used by Na/K pumps to pump out the excess Na^+. Given the limited energy supply in the brain, evolutionary strategies including shortening spike propagation distance, sparse coding and reducing Na^+- and K^+-current overlap etc. have helped minimize energy expenditure [9-13]. However, the observation that high frequency spiking tends to occur in large diameter axons [5] is confusing because fast spiking in large diameter axons causes more Na^+ influx and exerts a heavy burden on the energy supply in the brain [14, 15]. Additionally, the process by which Na/K pumps remove Na^+ is slow [16-22]. In neurons, it takes seconds or even minutes to remove the excess Na^+ after $[Na^+]$ elevation [16-21]. It remains unknown how these directly correlated processes, fast spiking and slow Na^+ removal, interact in the control of reliable spike generation

72 and propagation. Furthermore, it is unclear whether reducing Na^+- and K^+-current overlap can

73 consistently decrease the rate of Na^+ influx and accordingly enhance axonal reliability to propagate

74 high frequency spikes. Consequently, by modeling the process of Na^+ removal in unmyelinated

75 and myelinated axons, we systematically explored the effects of diameter, myelination and Na/K

76 pump density on spike propagation reliability.

Results

Different [Na^+] Decay Dynamics in Unmyelinated and Myelinated Axons. We implemented

multicompartment models of axons in NEURON (23). Both the unmyelinated and myelinated axon

models include I_{Na}, I_{KD}, I_{KA}, I_{leak} and the Na/K pump. All channels and pumps were evenly spread

in the unmyelinated axon model and at the nodes of Ranvier in the myelinated axon model. For

the myelinated compartments, there is a significantly lower density of I_{leak} and C_m to simulate the

effect of insulation (Materials and Methods). The model structures are shown in Fig. 1A.

Spikes propagate rapidly in both models. In the unmyelinated axon model, the propagation

increases from 0.1 m/s to 0.3 m/s when the axonal diameter increases from 0.2 μm to 1 μm. In the

myelinated axon model, the speed increases from 0.87 m/s to 2.1 m/s accordingly. Upon the arrival

of spikes, Na^+ channels open and extracellular Na^+ flows into axons to raise the intracellular [Na^+]

immediately (Fig. 1B). Then K^+ currents are activated to repolarize the spikes. Compared with the

other ionic currents, the magnitude of the Na/K pump current is extremely small during a single

spike, but it decays very slowly. Eventually, the Na/K pump gradually removes Na^+ from the axon,

leading to a slow, many second, [Na^+] decay in the unmyelinated axon model (Fig. 1B, left). In

contrast, in the myelinated axon model, the spike-triggered [Na^+] elevation at the node of Ranvier

decays on the time scale of hundreds of ms (Fig. 1B, right). The Na/K pump densities in the two

models are the same and thus the quick decay is not a result of super-efficient pumps. In the

myelinated axon model, Na^+ channels are placed only in the nodes of Ranvier and there is no

transmembrane Na^+ influx in the myelinated compartments. But, the slightly elevated $[Na^+]$ in the

myelinated compartment and the rapid decay of nodal $[Na^+]$ are due to the axial diffusion of Na^+

into neighboring myelinated compartments, rather than immediate pump action.

Because of the large diffusion coefficient (24) and the lack of buffering (16), $[Na^+]$ in the radial

direction becomes rapidly homogeneous. Thus, spike-triggered $[Na^+]$ elevation can be well

predicted by the surface-to-volume ratio (SVR ∝ 1/diameter). In both models, spike-triggered $[Na^+]$

elevation decreases with diameter (Fig. 1C). Increasing the pump density speeds up the $[Na^+]$

decay in the unmyelinated axon model, but has no obvious effect on the nodal $[Na^+]$ decay in the

myelinated axon model, further confirming that the quick decay is not a result of the Na/K pump.

Na^+ Accumulation in the Unmyelinated Axon at High Firing Frequencies. Compared with the

basal level before spiking (~ 10 mM), the $[Na^+]$ elevation caused by a single spike is small and

insufficient to affect axonal excitability (Fig. 1B). However, neurons can fire at high frequencies

in certain behavioral contexts (7, 25-27). With continuous high frequency spiking, Na^+ can

accumulate if subsequent spikes occur before the complete recovery of a preceding spike elicited

$[Na^+]$ rise. In the unmyelinated axon model, Na^+ significantly accumulates within just 100 spikes,

and the accumulation speed increases with firing frequency because of a faster Na^+ influx rate (Fig.

2A). Larger SVRs can also make Na^+ accumulate faster in thin axons. Note that the seemingly

slower Na^+ accumulation in the 0.2-μm-diameter axon at 60 Hz is due to earlier propagation failure.

In parallel with Na^+ accumulation, the Na^+ equilibrium potential (driving force of Na^+ current)

becomes more negative and the action potential peak decreases (Fig. 2B, C). With increased firing

frequency, propagation starts to fail. The failure starts from 30 Hz in the 0.2-μm-diameter axon

(Fig. 2C), but from 60 Hz in larger diameter axons. A repertoire of propagation failure patterns

were observed. In Fig. 2D1, every other spike propagated through the axon, but a more

complicated propagation failure pattern is shown in Fig. 2D2.

When the model fires over a sufficiently long period, the Na^+ influx rate and efflux rate

gradually balance each other to make $[Na^+]$ reach a steady state. Despite different accumulation

speeds, the steady state $[Na^+]$ increases with firing frequency and is independent of diameter until

propagation starts to fail (Fig. 2E). The rate of Na^+ influx per surface area is only determined by

the firing frequency because of identical Na^+ channel density. The Na/K pump-caused net Na^+

efflux rate is determined by $-3k_1[pump][Na^+]^3 + 3k_2[pumpNa]$. Because the pump density is

also identical, the model needs identical $[Na^+]$ to match influx rate and reach a steady state

irrespective of diameters. If propagation fails, the steady state $[Na^+]$ depends on the actual pattern

of spikes that propagate through the axon. For example, at 70 Hz with baseline pump density, the

steady state $[Na^+]$ is the same in the 0.6- and 1.0-μm-diameter axons but lower in the 0.2-μm-

diameter axon, because of their specific propagation failure patterns (Fig. 2E). When pumps

remove excess Na^+, they also carry a negative current, which gradually repolarizes the membrane

potential during spikes along with Na^+ accumulation. The factors causing the gradual occurrence

of failure includes the Na^+ channel's incomplete recovery from inactivation, decreased driving

force due to Na^+ accumulation, and increased outward pump current during continuous spiking.

This propagation failure can be eliminated or alleviated when Na^+ accumulation is not considered

in the model. Increasing pump density consistently reduces Na^+ accumulation (Fig. 2E) to alleviate

the equilibrium potential decrease, but also enhances the outward pump current to directly decrease

the axonal excitability (Fig. 2F).

Na$^+$ accumulates slowly in large diameter axons and in these axons it always requires more

spikes to trigger propagation failure than in small diameter axons, making these axons more

resilient to high frequency firing (Fig. 2G). At higher frequencies, Na$^+$ accumulates faster and Na$^+$

channels recover less from inactivation. Consequently, fewer spikes are required to trigger

propagation failure and accordingly the resilience decreases (Fig. 2G). With higher pump densities

in the model, the axonal resilience to high frequency spiking is increased (Fig. 2 G), suggesting

that the alleviated decrease in Na$^+$ equilibrium potential prevails against the enhanced outward

pump current (Fig. 2F).

Myelinated Compartments Slow Na$^+$ Accumulation. In contrast to the slow decay of [Na$^+$] in

the unmyelinated axon, the [Na$^+$] at the node of Ranvier in the myelinated axon shows a quick

decay (Fig. 1B). Can nodal [Na$^+$] elevate at high firing frequencies? It does elevate to reduce the

Na$^+$ equilibrium potential and action potential peak in parallel (Fig. 3A-C). As explained

previously, the quick decay of nodal [Na$^+$] is a result of nodal Na$^+$ quickly diffusing into

neighboring myelinated compartments rather than being removed. After diffusion, the nodal [Na$^+$]

is still higher than the pre-spiking level (Fig. 1B). In other words, nodal Na$^+$ influx not only needs

to fill the nodes of Ranvier, but also the neighboring myelinated compartments, which is equivalent

to decreasing the SVR. Consequently, the nodal Na$^+$ still accumulates, but at a much slower rate

(Fig. 3A, F). It requires thousands of spikes to significantly change nodal [Na$^+$] in myelinated

axons. Na$^+$ in the myelinated compartments flows back during the removal of Na$^+$ by nodal Na/K

pumps. Therefore, by acting like an 'anti-flood water reservoir', myelinated compartments slow

down the nodal Na$^+$ accumulation and the excitability reduction.

Similar to the case in unmyelinated axon models, nodal Na$^+$ accumulates faster at higher firing

frequencies (Fig, 3A). Propagation failure starts from 50 Hz in the 0.2-μm-diameter axon, but from

60 Hz in larger diameter axons. In Fig. 3D1, every other spike can propagate through the axon.

However, only every fourth spike can successfully propagate in Fig. 3D2. The steady state $[Na^+]$

is diameter independent until propagation fails at high frequencies (Fig. 3E). When firing at 80 Hz

with baseline pump density, the steady state $[Na^+]$ is still the same in axons with different diameters,

because they eventually have the same propagation patterns.

Because Na^+ accumulates more slowly in myelinated axons compared with unmyelinated axons

(Fig. 3F), the propagation is more reliable in myelinated axons (Fig. 3G). Large diameter axons

are consistently more resilient and the resilience decreases with firing frequency. Similar to

unmyelinated axon models, more pumps increase the propagation resilience.

The Effects of Na^+-and K^+-Current Overlap on Propagation Resilience. Previous work has

shown that the temporal overlap between the Na^+- and K^+-currents strongly affects Na^+ influx

during spiking (8). Can minimizing their temporal overlap consistently enhance axonal resilience

by reducing the rates of Na^+ influx and Na^+ accumulation?

Slowing the activation of I_{KD} was reported to increase spike duration and decrease Na^+ influx

per spike (8). However, slowing I_{KD} activation increases spike triggered Na^+ influx instead of

decreasing it in our model (Fig. 4A, c→d). By systematically varying I_{KD} activation time constant

in the range of 0.2-1.4 times of the baseline value, we observed a U-shaped relationship between

Na^+ influx rate and I_{KD} activation rate. Slowing the activation rate initially decreases the Na^+ influx

rate, but further slowing it increases the Na^+ influx rate (Fig. 4A). This occurs because slowing I_{KD}

activation increases spike duration to widen the Na^+ current profile; on the other hand, slower

repolarization reduces the Na^+ channel driving force to decrease the current amplitude during the

repolarization phase. Their balance determines the net change of Na^+ influx rate caused by slowing

the I_{KD} activation.

Apart from the U-shaped Na^+ influx rate changes, slowing I_{KD} activation consistently increases

spike duration, which results in changes that both favor and oppose spike propagation. Broader

spikes provide more axial charge to neighboring segments per spike to favor spike propagation,

but they also cause larger Na^+ channel inactivation and less recovery to oppose spiking. These

factors interact to make the effect of slowing I_{KD} activation on propagation reliability complex

(Fig. 4B, C). In both unmyelinated and myelinated axon models, slowing I_{KD} activation from a to

b consistently increases the propagation resilience. However, when slowing I_{KD} activation from b

to c, the resilience increases at low firing rates but decreases at high firing rates in both models.

Slowing I_{KD} activation from c to d consistently reduces the resilience in both models. These

complex effects reflect the interactions among Na^+ influx rate, spike duration and Na^+ channel

inactivation gate state changes.

We also explored the effect of speeding Na^+ current inactivation on propagation reliability.

Faster Na^+ current inactivation consistently reduces spike duration and Na^+ influx, but it also has

complex effects on the resilience because of the interactions among the above-mentioned opposing

factors (Fig. S1). The effects of diameter and myelination on the speed of Na^+ accumulation and

propagation resilience hold in all the simulations we have done.

Discussion

Neuronal firing frequency often varies over a broad range to encode a wide range of signals. For

a given neuron, the rate of Na^+ entry at its high firing frequency is usually much higher than the

rate at which excess Na^+ is pumped out of the cell. Therefore, while short periods of high frequency

firing should present no problem for most neurons, continuous high frequency spiking could result

in compromised function, as intracellular $[Na^+]$ increases. Hence, we explore the dynamic

relationships between fast spiking and slow Na^+ removal for reliable spike generation and

propagation.

The importance of the Na/K ATPase in regulating neuronal activities has attracted attention in

recent years (28). It can set neuronal excitability and contribute to maintaining synchronous

epileptiform bursting in developing CA3 neurons (29). By interacting with h-current, it regulates

bursting activity in the leech heartbeat central pattern generator (30). There have also been

theoretical efforts to study the pump's role in controlling the transition between different firing

patterns (31-33) and the transition between normal and pathological brain states (34-37). However,

to the best of our knowledge, how the pump interacts with axon diameter and myelin to remove

Na^+ influx and affect axonal excitability has not been previously explored.

Axonal morphology can exert a critical constraint on a neuron's maximal firing frequency.

Large diameter and myelinated axons are intrinsically advantageous to propagate spikes because

of their larger space constants. In this work, we highlight another advantage in propagating

continuous high frequency spikes. Lower SVRs and slower Na^+ accumulation make them more

resilient to high frequency spiking. This may explain why fast-spiking parvalbumin interneurons

are extensively myelinated (38) and why high frequency spiking tends to occur in large diameter

axons, despite a higher energy cost of information transmission (5, 14, 15). Note that rapid

redistribution of ions such as Na^+ can occur at other sites and in other processes. For example, the

soma, acting as a Na^+ sink, can enhance reliable spike initiation in axon initial segments (16).

Redistribution of chloride in dendrites is also shown to be important for chloride equilibrium

potential and synaptic inhibition (39).

Increasing the Na/K pump density to remove Na$^+$ more efficiently is expected to facilitate axonal

resilience to high frequency spiking. Existing data argue that Na$^+$ removal after spiking is slow. In

axon initial segments of L5 pyramidal neurons, as a result of Na$^+$ diffusing into the soma and

myelin wrapped compartments, elevated [Na$^+$] requires several seconds to approximately recover

(16). In mitral cell tuft dendrites, after a short period of high frequency stimulation, it takes minutes

for the Na/K ATPase to pump out the accumulated Na$^+$ (17). This slow Na$^+$ removal may be

functionally important because it enables integration and binding of multiple chemosensory stimuli

over a prolonged time scale (40). We used pump densities that should fall into a reasonable

biological range, and the Na$^+$ accumulation at high firing frequencies shown here is consistent with

recent Na$^+$ imaging data (17, 20).

This work illustrates the limitations of using conventional frequency-current curves generated

with a short-term current injection to estimate the reliability of spike propagation (41, 42). Neither

of these measures reflects the accommodation of neuronal excitability after continuous firing

caused changes such as Na$^+$ (explored in this work) and Ca^{2+} accumulation (showing a biphasic

decay (43, 44)).

In the models explored here, myelinated axons are resilient to high frequency spiking elicited

Na$^+$ accumulation. Given that myelinated axons are usually much thicker (1, 45), spike propagation

in myelinated axons should be extremely robust to fast spiking caused Na$^+$ accumulation under

usual physiological conditions. However, under conditions requiring neurons to fire for extended

time at high frequencies, Na$^+$ accumulation may still compromise the function of axons. Of course,

the frequency threshold and the number of spikes required to cause spike failure will depend on

the specifics of the expression of neuronal conductances, and will likely show variability as a

consequence of variable channel densities (46). Additionally, intrinsic channel noise can cause

spike failure (4, 27), and the normal function of the Na/K ATPases requires ATP, the supply of

which has a limit (35). Both a shortage of ATP after continuous firing and channel noise can further

reduce axonal reliability of propagating spikes.

In recent years, a trade-off has been proposed between a neuron's capability to spike rapidly

and the high energetic cost of spiking (8, 47). To achieve high spike rates (narrow spikes), neurons

need to consume more energy to remove the larger Na^+ influx, coproduct of fast spiking. However,

in our simulations, we find a more complicated relationship between Na^+ influx and spike duration

(Fig. 4). Furthermore, although narrow spikes can facilitate high frequency spike generation, they

are more vulnerable to Na^+ accumulation or noise disturbance during propagation. This work

implies that different strategies may be required in different types of neurons to guarantee reliable

spike propagation for their functional requirements.

Materials and Methods

Axonal Morphology. We modeled a segment of axon with the length of 3131 μm. The model was

implemented in NEURON (23). The model code is available from ModelDB (accession number:

267009). The simulations used an integration step of 0.05 ms. The unmyelinated model was

discretized into 51 segments in the longitudinal direction. For the myelinated axon model, there

are 31 nodes of Ranvier flanked with 32 myelinated compartments (Figure 1). The lengths of the

nodes of Ranvier are 1 μm (nseg = 1). Each myelinated compartment is 100 μm long (nseg = 3),

except the outer two myelinated compartments are 50 μm each (nseg = 3). In the radial direction,

there are 4 shells to model Na^+ diffusion (23). The Na^+ diffusion coefficient is 0.6 $μm^2$/ms (24)

and there is no buffering (16), leading to a homogeneous $[Na^+]$ in the radial direction. In both

peripheral and central nervous systems, unmyelinated axons are usually thinner than myelinated

axons. In the central nervous system, axons > 0.2 μm are usually myelinated (1). To delineate the

effect of both diameter and myelination on Na⁺ accumulation, we chose the same range of

diameters between 0.2–1.0 μm.

Ionic Channels and the Na/K pump. The model has a minimal set of ionic channels: fast Na

current (I$_{Na}$), delayed rectifier K current (I$_{KD}$), A-type K current (I$_{KA}$) and leak current (I$_{leak}$). They

have standard kinetic formulations in the form of $I = g_{max}m^p h^q(V - E)$; $\tau_m \dot{m} = m_\infty(V) -$

m; $\tau_h \dot{h} = h_\infty(V) - h$; g_{max} represents maximal conductance; m and h represent activation and

inactivation gates; p and q represent corresponding gate exponents; E represents equilibrium

potential; τ_m and τ_h represent corresponding time constants; All the channel kinetics were

modified from (48). For I$_{Na}$, $g_{max} = 0.015\,S/cm^2$; $p = 3$; $q = 1$; $m_\infty =$

$1/(1 + exp(-(V + 38)/8.5))$; $\tau_m = 0.132/(\cosh((V + 27)/7.5) + 0.003/(1 +$

$\exp(-(V + 27)/5)))$; $h_\infty = 1/(1 + exp(V + 47/6))$; $\tau_h = 10/\cosh((V + 42)/15$; $E_{Na} =$

$RT/F * \ln([Na^+]_o/[Na^+]_i)$; $[Na^+]_o$ is fixed to be 140 mM; $[Na^+]_i$ (also written as $[Na^+]$ in the

main text) was dynamically computed by I$_{Na}$ and the Na/K pump; For I$_{KD}$, $g_{max} = 0.216\,S/cm^2$;

$p = 4$; $q = 0$; $m_\infty = (-0.01 * (V + 45.7)/(exp(-(V + 45.7)/10) - 1))/((-0.01 * (V +$

$45.7)/(exp(-(V + 45.7)/10) - 1)) + 0.125 * exp(-(V + 55.7)/80))$; $\tau_m = 1/((-0.01 *$

$(V + 45.7)/(exp(-(V + 45.7)/10) - 1)) + 0.125 * exp(-(V + 55.7)/80))$; $E_K =$ -70 mV;

For I$_{KA}$, $g_{max} = 0.02\,S/cm^2$; $p = 3$; $q = 1$; $m_\infty = (0.0761 * exp((V + 94.22)/31.84)/(1 +$

$exp((V + 1.17)/28.93)))^{1/3}$; $\tau_m = 0.3632 + 1.158/(1 + exp((V + 55.96)/20.12))$; $h_\infty =$

$1/(1 + exp(V + 53.3/14.54))^4$; $\tau_h = 1.24 + 2.678/(1 + exp((V + 50)/16.027))$; For

I$_{leak}$, $g_{max} = 1.25 \times 10^{-4}\,S/cm^2$; $p = 0$; $q = 0$; $E_{leak} =$ -65 mV; the Na/K pump model is

similar with that used in (17, 40), described by the following kinetic schemes: $[pump] + 3[Na^+]_i$

$\xleftrightarrow{k_1,k_2} [pumpNa]$; $[pumpNa] \xleftrightarrow{k_3,k_4} [pump] + 3[Na^+]_o$; $pump_{total} = [pump] + [pumpNa]$;

$k_1 = 2\ mM^{-3}msec^{-1}$; $k_2 = 0.001\ msec^{-1}$; $k_3 = 0.6\ msec^{-1}$; $k_4 = 0.437\ mM^{-3}msec^{-1}$;

By these parameters, the steady state [Na$^+$] is ~ 10 mM when the axon models are silent. For Fig.

4, a scaling factor in the range of 0.2 -1.4 was multiplied with τ_m at all voltages for I$_{KD}$.

For the unmyelinated axon model, all defined currents and the Na/K pump were evenly

distributed with C$_m$ = 1 μF/cm^2 and R$_a$ = 120 Ωcm. For the myelinated axon model, the ionic

currents and the Na/K pump were placed in the nodes of Ranvier (1, 49). For simplicity, we did

not model juxtaparanodal regions. At the nodes of Ranvier, C$_m$ = 1 μF/cm^2; On myelinated

compartments, there is only a smaller I$_{leak}$ with g_{max} = 1.25×10^{-6} S/cm^2 and effective C$_m$ = 0.01

μF/cm^2; R$_a$ is 120 Ωcm. In both the unmyelinated and the myelinated axon models, the current

densities are identical so that we could explore the role of diameter and myelination on Na$^+$

accumulation. The $pump_{total}$ is in the range of 0.5–5 pmol/cm^2. We did not explicitly simulate

the axon initial segment in our model. Instead, we applied strong enough current at the starting

end of the unmyelinated model and the first node of Ranvier in the myelinated model to reliably

trigger spikes.

Acknowledgments

313 The authors would like to thank the constructive suggestions from the reviewers and from the
314 members of Marder lab. This work was supported by the grant R35NS097343.

References

1.   S. G. Waxman, J. D. Kocsis, P. K. Stys, *The Axon: Structure, Function, and Pathophysiology*
(Oxford University Press, 1995).
2.   D. Debanne, E. Campanac, A. Bialowas, E. Carlier, G. Alcaraz, Axon physiology. Physiol Rev
91, 555-602 (2011).
3.   H. Schmidt, T. R. Knosche, Action potential propagation and synchronisation in
myelinated axons. PLoS Comput Biol **15**, e1007004 (2019).

4. A. A. Faisal, J. A. White, S. B. Laughlin, Ion-channel noise places limits on the miniaturization of the brain's wiring. Curr Biol **15**, 1143-1149 (2005).

5. J. A. Perge, J. E. Niven, E. Mugnaini, V. Balasubramanian, P. Sterling, Why do axons differ in caliber? J Neurosci **32**, 626-638 (2012).

6. B. Wang *et al.*, Firing Frequency Maxima of Fast-Spiking Neurons in Human, Monkey, and Mouse Neocortex. Front Cell Neurosci **10**, 239 (2016).

7. S. Chen, G. J. Augustine, P. Chadderton, Serial processing of kinematic signals by cerebellar circuitry during voluntary whisking. Nat Commun **8**, 232 (2017).

8. A. Hasenstaub, S. Otte, E. Callaway, T. J. Sejnowski, Metabolic cost as a unifying principle governing neuronal biophysics. Proc Natl Acad Sci USA **107**, 12329-12334 (2010).

9. S. B. Laughlin, Energy as a constraint on the coding and processing of sensory information. Curr Opin Neurobiol **11**, 475-480 (2001).

10. W. B. Levy, R. A. Baxter, Energy efficient neural codes. Neural Comput **8**, 531-543 (1996).

11. H. Alle, A. Roth, J. R. Geiger, Energy-efficient action potentials in hippocampal mossy fibers. Science **325**, 1405-1408 (2009).

12. F. J. Heras, J. Anderson, S. B. Laughlin, J. E. Niven, Voltage-dependent K^+ channels improve the energy efficiency of signalling in blowfly photoreceptors. J R Soc Interface **14** (2017).

13. H. Hu, F. C. Roth, D. Vandael, P. Jonas, Complementary Tuning of Na^+ and K^+ Channel Gating Underlies Fast and Energy-Efficient Action Potentials in GABAergic Interneuron Axons. Neuron **98**, 156-165 e156 (2018).

14. C. Howarth, P. Gleeson, D. Attwell, Updated energy budgets for neural computation in the neocortex and cerebellum. J Cereb Blood Flow Metab **32**, 1222-1232 (2012).

15. R. Nortley, D. Attwell, Control of brain energy supply by astrocytes. Curr Opin Neurobiol **47**, 80-85 (2017).

16. I. A. Fleidervish, N. Lasser-Ross, M. J. Gutnick, W. N. Ross, Na^+ imaging reveals little difference in action potential-evoked Na^+ influx between axon and soma. Nat Neurosci **13**, 852-860 (2010).

17. A. Zylbertal, A. Kahan, Y. Ben-Shaul, Y. Yarom, S. Wagner, Prolonged Intracellular Na^+ Dynamics Govern Electrical Activity in Accessory Olfactory Bulb Mitral Cells. PLoS Biol **13**, e1002319 (2015).

18. A. Zylbertal, Y. Yarom, S. Wagner, The Slow Dynamics of Intracellular Sodium Concentration Increase the Time Window of Neuronal Integration: A Simulation Study. Front Comput Neurosci **11**, 85 (2017).

19. D. Ziemens, F. Oschmann, N. J. Gerkau, C. R. Rose, Heterogeneity of Activity-Induced Sodium Transients between Astrocytes of the Mouse Hippocampus and Neocortex: Mechanisms and Consequences. J Neurosci **39**, 2620-2634 (2019).

20. T. Ona-Jodar, N. J. Gerkau, S. Sara Aghvami, C. R. Rose, V. Egger, Two-Photon Na^+ Imaging Reports Somatically Evoked Action Potentials in Rat Olfactory Bulb Mitral and Granule Cell Neurites. Front Cell Neurosci **11**, 50 (2017).

21. B. Moshrefi-Ravasdjani, D. Ziemens, N. Pape, M. Färfers, C. R. Rose, Action Potential Firing Induces Sodium Transients in Macroglial Cells of the Mouse Corpus Callosum. Neuroglia **1**, 106-125 (2018).

22. Y. Zang *et al.*, Theoretical investigation of the mechanism of heart failure using a canine
ventricular cell model: especially the role of up-regulated CaMKII and SR Ca^{2+} leak. J Mol
Cell Cardiol **56**, 34-43 (2013).
23. M. L. Hines, N. T. Carnevale, The NEURON simulation environment. Neural Comput **9**,
1179-1209 (1997).
24. M. J. Kushmerick, R. J. Podolsky, Ionic mobility in muscle cells. Science **166**, 1297-1298
(1969).
25. L. S. Tang *et al.*, Precise temperature compensation of phase in a rhythmic motor pattern.
PLoS Biol **8**, e1000469 (2010).
26. J. C. Athilingam, R. Ben-Shalom, C. M. Keeshen, V. S. Sohal, K. J. Bender, Serotonin
enhances excitability and gamma frequency temporal integration in mouse prefrontal
fast-spiking interneurons. Elife **6**, e31991 (2017).
27. D. Al-Basha, S. A. Prescott, Intermittent Failure of Spike Propagation in Primary Afferent
Neurons during Tactile Stimulation. J Neurosci **39**, 9927-9939 (2019).
28. M. D. Forrest, The sodium-potassium pump is an information processing element in brain
computation. Frontiers in Physiology **5** (2014).
29. L. R. Shao, R. Janicot, C. E. Stafstrom, Na^+-K^+-ATPase functions in the developing
hippocampus: regional differences in CA1 and CA3 neuronal excitability and role in
epileptiform network bursting. J Neurophysiol **125**, 1-11 (2021).
30. D. Kueh, W. H. Barnett, G. S. Cymbalyuk, R. L. Calabrese, Na^+/K^+ pump interacts with the
h-current to control bursting activity in central pattern generator neurons of leeches. Elife
5, e19322 (2016).
31. M. D. Forrest, M. J. Wall, D. A. Press, J. Feng, The sodium-potassium pump controls the
intrinsic firing of the cerebellar Purkinje neuron. PLoS One **7**, e51169 (2012).
32. A. O. Komendantov, C. C. Canavier, Electrical coupling between model midbrain
dopamine neurons: effects on firing pattern and synchrony. J Neurophysiol **87**, 1526-1541
(2002).
33. C. C. Canavier, Sodium dynamics underlying burst firing and putative mechanisms for the
regulation of the firing pattern in midbrain dopamine neurons: a computational approach.
J Comput Neurosci **6**, 49-69 (1999).
34. G. P. Krishnan, G. Filatov, A. Shilnikov, M. Bazhenov, Electrogenic properties of the Na^+/K^+
ATPase control transitions between normal and pathological brain states. J Neurophysiol
113, 3356-3374 (2015).
35. G. Le Masson, S. Przedborski, L. F. Abbott, A computational model of motor neuron
degeneration. Neuron **83**, 975-988 (2014).
36. J. S. Coggan, G. K. Ocker, T. J. Sejnowski, S. A. Prescott, Explaining pathological changes in
axonal excitability through dynamical analysis of conductance-based models. J Neural Eng
8, 065002 (2011).
37. N. Yu, C. E. Morris, B. Joos, A. Longtin, Spontaneous excitation patterns computed for
axons with injury-like impairments of sodium channels and Na/K pumps. PLoS Comput
Biol **8**, e1002664 (2012).
38. J. Stedehouder *et al.*, Fast-spiking Parvalbumin Interneurons are Frequently Myelinated
in the Cerebral Cortex of Mice and Humans. Cereb Cortex **27**, 5001-5013 (2017).

39.   N. Doyon *et al.*, Efficacy of synaptic inhibition depends on multiple, dynamically
interacting mechanisms implicated in chloride homeostasis. PLoS Comput Biol **7**,
e1002149 (2011).

40.   A. Zylbertal, Y. Yarom, S. Wagner, Synchronous Infra-Slow Bursting in the Mouse
Accessory Olfactory Bulb Emerge from Interplay between Intrinsic Neuronal Dynamics
and Network Connectivity. J Neurosci **37**, 2656-2672 (2017).

41.   N. W. Gouwens *et al.*, Classification of electrophysiological and morphological neuron
types in the mouse visual cortex. Nat Neurosci **22**, 1182-1195 (2019).

42.   A. Foust, M. Popovic, D. Zecevic, D. A. McCormick, Action Potentials Initiate in the Axon
Initial Segment and Propagate through Axon Collaterals Reliably in Cerebellar Purkinje
Neurons. The Journal of Neuroscience **30**, 6891-6902 (2010).

43.   H. Schmidt, K. M. Stiefel, P. Racay, B. Schwaller, J. Eilers, Mutational analysis of dendritic
Ca^{2+} kinetics in rodent Purkinje cells: role of parvalbumin and calbindin D28k. J Physiol
551, 13-32 (2003).

44.   Y. Zang, S. Dieudonne, E. De Schutter, Voltage- and Branch-Specific Climbing Fiber
Responses in Purkinje Cells. Cell Rep **24**, 1536-1549 (2018).

45.   S. Lee *et al.*, A culture system to study oligodendrocyte myelination processes using
engineered nanofibers. Nat Methods **9**, 917-922 (2012).

46.   D. J. Schulz, J. M. Goaillard, E. Marder, Variable channel expression in identified single and
electrically coupled neurons in different animals. Nat Neurosci **9**, 356-362 (2006).

47.   B. C. Carter, B. P. Bean, Sodium entry during action potentials of mammalian neurons:
incomplete inactivation and reduced metabolic efficiency in fast-spiking neurons. Neuron
64, 898-909 (2009).

48.   Y. Zhang, D. Bucher, F. Nadim, Ionic mechanisms underlying history-dependence of
conduction delay in an unmyelinated axon. Elife **6**, e25382 (2017).

49.   S. G. Waxman, J. M. Ritchie, Molecular dissection of the myelinated axon. Ann Neurol **33**,
121-136 (1993).

Figures and Tables

Figure 1. Spike-triggered [Na$^+$] rise and decay in unmyelinated and myelinated axon models. (A)
Schematics of the unmyelinated (left) and the myelinated (right) axon models. Black traces show
spike waveforms in two models (50 μm from the distal axon end; In myelinated model, it is also a
node of Ranvier). (B) Spike-triggered [Na$^+$] rise and decay in the unmyelinated (left) and the
myelinated (right) axon models. Black traces were measured at corresponding voltage recording
sites. Green trace shows [Na$^+$] in the myelinated compartment (16.7 μm from the node of Ranvier)
and the initial part is enlarged in the inset. The Na/K pump density is 0.5 pmol/cm^2, referred to as
baseline condition hereafter. (C) The effects of diameter and pump density on [Na$^+$] rise and decay
in the unmyelinated (left) and the myelinated (right, at the node of Ranvier) axon models. Note the
very different time scales in the two models.

Figure 2. The effects of diameter and pump density on Na$^+$ accumulation and resilience to high frequency spiking in the unmyelinated axon model. A-C show [Na$^+$] increase, Na$^+$ equilibrium potential decrease and membrane potential peak reduction (bottom) in models with different diameters and firing frequencies. (D) Voltage traces and interspike intervals (ISIs) at distal axon for 40 Hz (D1) and 50 Hz (D2) spike signals in the 0.2-μm-diameter axon. (E) Steady state [Na$^+$] elevation at different stimulation frequencies (left). Right panel shows ISIs at distal axon for 70 Hz spike signals with baseline pump density. (F) Na$^+$ equilibrium potential decrease (black) and outward pump current increase (blue) with different pump densities. 80 Hz spike signals in the 0.6-μm-diameter axon. In E & F, dot and star represent 1 and 10 times of baseline pump density, respectively. (G) The effect of pump density on axonal resilience to high frequency spiking, measured by the number of spikes required to cause propagation failure (nspike_fail), when firing at 70 and 80 Hz. The absence of data points means propagation doesn't fail under corresponding conditions.

Figure 3. Myelination slows Na$^+$ accumulation to increase axonal resilience to high frequency spiking. A-C show [Na$^+$] increase, Na$^+$ equilibrium potential decrease and membrane potential peak reduction at the node of Ranvier in the myelinated axon model with different diameters and firing frequencies. (D) Voltage traces and ISIs at distal axon for 60 Hz (D1) and 90 Hz (D2) spike signals in the 0.2-μm-diameter axon. (E) Steady state [Na$^+$] elevation at different stimulation frequencies (left). Dot and star represent 1 and 10 times of baseline pump density, respectively. Right panel shows ISIs at distal axon for 80 Hz spike signals with baseline pump density. (F) 100-spike caused Na$^+$ accumulation in the unmyelinated axon and the myelinated axon (node of Ranvier). The diameter is 0.2 μm, with baseline pump density. (G) The effect of pump density on axonal resilience to high frequency spiking, when firing at 70 Hz and 80 Hz. The absence of data points means propagation doesn't fail under corresponding conditions.

Figure 4. Complex effects of slowing I_{KD} activation on Na^+ influx rate and propagation resilience. (A) U-shaped dependence between Na^+ influx rate and I_{KD} activation rate. y axis (left panel) measures single spike triggered Na^+ influx per μm^2. c represents the baseline model (magenta); a, b and d represent 0.2, 0.6 and 1.4 times of the baseline I_{KD} activation time constant, respectively. Changes of spike shape, current profile and Na^+ channel inactivation gate state with slowed I_{KD} activation. The effects of delayed activation of I_{KD} on propagation resilience in unmyelinated and myelinated models are shown in B and C, respectively. a - d correspond to conditions defined in A. Resilience changes are measured by comparing symbols of the same color (coding diameter). For example, from a to b at 40 Hz in the unmyelinated model, black stars are higher than black dots, suggesting increased resilience in the 0.2-μm-diameter axon. The resilience also increases in larger diameter axons, because slowed I_{KD} activation eliminates propagation failure. The absence of data points means propagation doesn't fail under corresponding conditions.

490   Supporting Information

Figure S1. The effects of increasing the rate of Na⁺ current inactivation on propagation resilience. Star represents the baseline model and dot represents 40% decreased inactivation time constant of Na⁺ current. In unmyelinated axon models, at 30 Hz, faster inactivation decreases the resilience in the 0.2-μm-diameter axon model (dots are lower than stars). At 80 Hz, faster inactivation increases the resilience (dots are higher than stars). The same trend is observed in myelinated axon models. The absence of data points means propagation doesn't fail.