Quantifying isolation-by-resistance and connectivity in dendritic ecological networks

Tyler K. Chafin[1,*], Steven M. Mussmann[2,†], Marlis R. Douglas[2], and Michael E. Douglas[2]

[1]Biomathematics and Statistics Scotland, Edinburgh, EH49 3FD, United Kingdom

[2]Department of Biological Sciences, University of Arkansas, Fayetteville, AR, 72701, USA

[†]Current Address: Abernathy Fish Technology Center, U.S. Fish & Wildlife Service, Longview, WA, 98632, USA

Keywords: graph-based algorithms; stream hierarchy; riverscape genetics; environmental resistance

2 **ABSTRACT**

1.   A central theme in landscape ecology is the translation of individual movements within a

population by deconstructing/interpreting the components of its topographical environment.

Most such endeavors rely heavily on the concept of 'landscape resistance' – a composite of an

6 arbitrary number of features/covariates that, when identified/compiled, yield a 'surface' inversely

related to net movement. However, the statistical methodologies underlying this compilation

8 have limited applicability when applied to dendritic ecological networks (DENs), including

riverscapes.

10 2.   Herein we provide an analytical framework (ResistNet) that more appropriately annotates

DEN segments by first aligning individual genetic distances with environmental covariates

12 within a graph structure, then employing a genetic algorithm to optimise a composite model.

3.   We evaluated the efficacy of our method by first testing it *in silico* across an array of

14 sampling designs, spatial trajectories, and levels of complexity, then applying it in an empirical

case study involving 13,218 ddRAD loci from N=762 Speckled Dace (Leuciscidae: *Rhinichthys*

16 *osculus*), sampled across N=78 Colorado River localities. By doing so, we underscored the

utility of ResistNet within a large-scale conservation study, as well as identified prerequisites for

18 its appropriate application.

4.   Our contemporary framework not only allows an interpretation of meta-population/meta-

20 community structure across DENs, but also highlights several innovative applications. These are:

(a) Expanding an ongoing study design, and thus its hypotheses, into yet unsampled temporal

22 and/or spatial arenas, and; (b) Promoting multi-species management through comparative

analyses that extend across species and/or drainages.

1. INTRODUCTION

26 The physical structure of watersheds not only defines form and function in freshwater fishes

(Douglas & Matthews, 1992; Jackson et al., 2001), but also their intraspecific evolutionary

28 trajectories and population dynamics (Hopken et al., 2013). As a result, patterns of

diversification differ markedly from those found in terrestrial land- and seascapes (Manel et al.,

30 2020; Martinez et al., 2018). The unique constraints of riverine networks serve not only to define

dispersal pathways but also to dictate genetic relatedness within/ among aquatic biota (Meffe &

32 Vrijenhoek, 1988). Network 'connectivity' (=potential for movement), and 'complexity'

(=branching rates/ nonlinear topology) control numerous population processes, such as: Genetic

34 divergence (Hopken et al., 2013), colonization probabilities (Falke et al., 2012; Labonne et al.,

2008), and metapopulation components (Campbell Grant, 2011; Fagan, 2002).

36   In general, complex dendritic networks may promote intraspecific genetic diversities

(Thomaz et al., 2016), and divergences among distal populations (Chiu, Li, et al., 2020). Yet, the

38 unidirectional flows in riverscapes differ from other such dendritic ecological networks (DENs),

and thus biases dispersal (i.e., up *versus* downstream). This in turn promotes asymmetric gene

40 flow and source-sink metapopulation structure (Campbell Grant et al., 2007). Allelic diversity

increases in a downstream direction (Blanchet et al., 2020; Paz-Vinas et al., 2015; Paz-Vinas &

42 Blanchet, 2015), whereas unique genetic variation is most often retained within peripheral

headwater populations (Chiu, Nukazawa, et al., 2020; Morrissey & De Kerckhove, 2009).

44   Conversely, effective population size (Altermatt & Fronhofer, 2018) and genetic

variability (Thomaz et al., 2016) seemingly decrease within unimpeded linear systems which, in

46 turn, impacts both resilience and stability of lineages (Mari et al., 2014; Terui et al., 2019), as

well as proliferation capacity (Harvey et al., 2019). However, numerous considerations can alter

48   these expectations, such as: Dispersal bias (Chiu, Li, et al., 2020), nonequilibrium processes

       (Pilger et al., 2017), geomorphic/ climatic regimes (Douglas et al., 1999; Osborne et al., 2014),

50   and abiotic species-level interactions (Paz-Vinas et al., 2015; Rodríguez-González et al., 2019) .

           It is thus not surprising that evaluation of stream distances alone rarely captures

52   intraspecific genetic patterns. In a comparison of 12 salmonid species, isolation-by-distance

       (IBD) varied from r^2=0.06–0.87 (Bradbury & Bentzen, 2007), and from 0.0001–0.44 among

54   three darter species (Camak & Piller, 2018). A larger-scale study involving 31 co-distributed

       stream fishes across 7 families identified a significant relationship between among-site F_{st} and

56   log-transformed river network distances (Mantel r=0.11–0.88), with stream hierarchy explaining

       the greatest proportion of genetic variance (Zbinden et al., 2022a). Although such deviations are

58   partly driven by stochastic or nonequilibrium processes (Hutchison & Templeton, 1999), their

       residuals should also reflect spatial features that impede individual movements (Keis et al., 2013;

60   Tang et al., 2020). Thus, divergence can be considered as a general function of the movement-

       cost along stream segments, i.e., 'isolation-by-resistance'(IBR)(B. H. McRae, 2006; B. H.

62   McRae & Beier, 2007).

           The IBR model applied within dendritic networks emphasizes specific attributes of

64   individual edges or stream segments, with resistance amplified by an array of localized factors

       (Fullerton et al., 2010; Hughes et al., 2009): Rate of elevational change (Lowe et al., 2006), flow

66   regime (Rolls et al., 2016), channel width/ catchment area (Ma et al., 2020), water composition

       (Fourtune et al., 2016), and seasonal hydroclimate (Brauer et al., 2016, 2018). In addition,

68   organisms that never (or rarely) occupy upstream or headwater conditions can still be directly

       affected by those habitats (Frissell et al., 1986; Fullerton et al., 2010; Nadeau & Rains, 2007).

70   Most linear streams are inherently predisposed to fragmentation (Brauer & Beheregaray,

2020; Fagan et al., 2002; Fagan, 2002), a situation particularly exacerbated, given that relatively

72   few remain free-flowing (Grill et al., 2019). Water storage also impacts habitat suitability

(Chafin et al., 2019; Fraser et al., 2019), with populations shifting trajectories accordingly

74   (Dibble et al., 2021). Dams and diversions extirpate populations and promote homogenization by

disrupting homing behaviors (Baggio et al., 2018), with species of limited mobility particularly

76   impacted (Coleman et al., 2018). Again, vulnerability and the specific nature of impacts vary by

context and species (Prunier et al., 2018).

1.1 | *Landscape genetic methods applied to riverscapes*

80   The mechanics of landscape genetics/genomics often translate poorly to dendritic networks

(Davis et al., 2018; Grummer et al., 2019). At issue is the riverscape structure upon which

82   genetic relationships can be mapped. Frequent *ad hoc* approaches often lack an explicit spatial

framework, or rely upon spatial methodology is landscape-based (Cooke et al., 2014; Galbraith

84   et al., 2015; Kanno et al., 2011). Often, these methods do so by contrasting a genetic distance

matrix against pairwise differences in local conditions, often using the Mantel test and

86   derivatives (Douglas & Endler, 1982; Smouse et al., 1986), thus neglecting the space through

which individuals must travel between points, restricting their riverscape-applicability (Davis et

88   al., 2018).

On the other hand, path-based methods (as herein) consider the cumulative costs of

90   movement, with resistance being represented as a composite of contributing environmental

factors (Zeller et al., 2012) that allow for habitat-related metrics to be derived, such as: Least-

92   cost paths, net flux between points, and movement(B. McRae et al., 2016). Operationally, it

involves summarizing spatial variables as a raster image of n x m cells (analogous to pixels),

94   with assigned values representing 'resistance' (Davis et al., 2018).

We propose a graph structure to translate this concept to riverscapes, with edges

96   annotated with environmental attributes connecting the nodes (i.e., populations or junctions)

(Eros et al., 2011; Peterson et al., 2013). Few approaches have considered resistance within a

98   spatially explicit, graph-theoretic framework, and even fewer have incorporated multiple

covariates within an hypothesis-testing framework [although see (White et al., 2020)]. Here, an

100   heuristic approach that explores complex parameter space is required, particularly given the

sensitivity of resistance to parameterization, and the difficulties with the *a priori* weighting of

102   specific variables within a larger pool of candidates (Spear et al., 2010). These approaches have

previously been developed for landscapes, using algorithms for model exploration and selection

104   (Peterman, 2018), and proven successful in both empirical and *in silico* scenarios (Kimmig et al.,

2020; Peterman et al., 2019; Winiarski et al., 2020).

106   Here, we employ a similar resistance perspective to quantify riverscape genetics/

genomics and do so by exploiting large-scale, contemporary efforts that quantify/ classify stream

108   variables (Grill et al., 2019; Linke et al., 2019; McManamay et al., 2018; Troia & McManamay,

2020). We utilised our package (ResistNet) to evaluate a case study involving Speckled Dace

110   (Leuciscidae, *Rhinichthys osculus*), the most widely distributed freshwater fish endemic within

the most endangered and regulated river system of North America, the Colorado River (Grill et

112   al., 2019; Minckley & Deacon, 1968; Nilsson et al., 2005). To accomplish this, we analyzed a

large ddRAD dataset to test the following hypotheses: 1) Populations are spatially structured

114   within the topology of the dendritic network; 2) Genetic differentiation is modulated by

environmental attributes, representing edges within the river network; and 3) Spatio-genetic

116   patterns cannot be explained by distance alone, but rather by a model of IBR.

118   **2. METHODS**

ResistNet facilitates high-throughput analyses by automating the association between

120   environmental correlates and genetic differentiation in dendritic networks (Figure 1). Source

code and documentation is available at: github.com/tkchafin/resistnet.

2.1 | *Composite resistance models optimized within networks*

124   ResistNet combines a genetic algorithm with multiple environmental variables as a mechanism

to parameterize resistance (Figure 1), similar to that found in ResistanceGA (Peterman, 2018),

126   but with several important differences. Genetic algorithms are useful for model optimisation and

selection of features in that they are less impacted by local optima when compared to other hill-

128   climbing techniques (Prügel-Bennett, 2004). They are employed to simultaneously evaluate a

population of models by using log-likelihood or AIC to gauge *fitness*. Here, *individuals* (as

130   randomly-sampled model configurations) consist of *chromosomes*, with *genes* (as parameter

choices) represented by 1) Variable inclusion/exclusion; 2) Variable transformations (using

132   functions available in ResistanceGA; Supplementary Figure S1); 3) Shape parameters for

transformation; or 4) Variable weights (with user-defined ranges).

134   For an optimisation problem involving n-environmental parameters, an individual

chromosome has length $4n$. Fitness values (e.g., AIC) then determine the survival of individual

136   models and their contribution to successive generations, with stochasticity represented by

mutation (random changes to parameter values) and *recombination* (parameter choices by

138   multiple individuals combined as offspring). Both of these occur following user-defined

probabilities. Optimisation continues until a threshold number of generations have passed

140   without an increase in population fitness.

The workflow used by ResistNet (Figure 1) is summarized as follows:

142   1.   Network structure and environmental data are imported as shapefiles. Environmental

data for contiguous stream reaches (=edges) are summarised and rescaled from 0−10, (per

144   Peterman, 2018).

2.   The initial population has a user-defined size and randomly seeded, which initialises the

146   genetic algorithm (as implemented using DEAP; De Rainville et al., 2012). Individuals

(as chromosomes) contain transformation, shape, and weight parameters to ensure an

148   optimal scale per a potential non-linear relationship between gene flow and each variable.

This process also allows the relationship in a composite resistance model to vary

150   according to relative influence and directionality (e.g., positive/ negative).

3.   Cumulative effective resistance is then computed pairwise between nodes as a weighted

152   sum of (transformed/ scaled) individual covariates (McRae et al., 2008; Thiele et al.,

2018). By default, each covariate is individually aggregated pairwise between samples as

154   the sum of values along the Dijkstra least-distance path, although custom behaviours can

be user-specified. This results in a single composite pairwise resistance matrix for each

156   model.

4.   Individual fitness is calculated via a mixed effects maximum likelihood population

158   effects (MLPE) model (Clarke et al., 2002; Van Strien et al., 2012). This allows the

pairwise genetic and resistance matrices to be compared and fitted using the MLPE

160   function from ResistanceGA. Pairwise distances are inherently non-independent, a

condition MLPE models were specifically formulated to accommodate, and are thus

162   effective for landscape genetic model selection (Peterman & Pope, 2021; Shirk et al.,

2018). Fitness for the genetic algorithm is assessed as log-likelihood, AIC, or marginal

164   R^2, but may also be computed using ΔAIC_{null} (i.e., the change in AIC values as compared

to a random effects null model).

166   5. Individuals are selected for successive generations via 'tournament selection,' a

deterministic method whereby random sets of individuals (user-specified size n) are

168   compared (=tournaments), with 'winners' being retained.

6. Mutation (=changing parameter values) and recombination (=exchanging parameter

170   combinations among models) occur at generational intervals, with the respective

probabilities of each being user-specified.

172   7. Steps 2-6 are repeated for a user-specified number of generations, or until user-specified

generations fail to increase the maximum fitness beyond a given threshold.

174   8. After stopping criteria are met, Akaike weights are computed for the highest fitness

models in the population, with these being retained to meet a cumulative user-specified

176   Akaike weight threshold (default value=0.95). Model-averaged resistance and multi-

model relative variable importance values are subsequently computed using the chosen

178   model set (below).

180   Multi-model inference has several benefits: It avoids the inherent subjectivity of model

selection, in that estimates from every candidate model are integrated proportional to their

182   approximation of the observed genetic distances [using the Akaike weight (w_i for i models)],

while also more concisely incorporating both uncertainty and information from closely ranked

184   models. Additionally, it also allows the quantification of relative variable importance (RVI)

across models composed of varying numbers of explanatory variables, and does so by deriving

186   the sum of Akaike weights (SW) as a robust measure of RVI in a multi-model inference

framework (Giam & Olden, 2016).

2.2 | *Simulation-based validation*

190   To validate the capacity of ResistNet to accurately recover a known model, we employed a

simulation scheme involving three river networks of varying complexity (near-linear, moderately

192   branched, and highly branched; Figure 1) and four scenarios involving different environmental

covariates (Table 1), to include distance-only models, and those varying in weights. Three

194   sampling levels (10, 30, and 50) were evaluated, with 20 replicates per level. Simulations were

conducted using *simResistNet.py*, an auxiliary software that simplifies validation with a specific

196   riverscape and its combination of environmental variables. Input consists of shapefiles for the

three template networks, and simulation parameters (as tab-delimited text) specifying covariates

198   for inclusion, their weights, and any transformations that must be applied. These are employed to

compute pairwise resistance distances between sampled points, which are then normalised

200   between 01.

        ResistNet was run at 600 generations maximum on each replicate, with an ending

202   condition of 20 generations of continuous failure to increase maximum observed fitness, a fixed

population size of $N=600$, a variable weight floor of $N=0.2$, a maximum transformation shape of

204    $n=0,5$, and mutation/ recombination probabilities of 0.8. To facilitate replication, the evaluation

procedure is implemented as a Snakemake workflow (Mölder et al., 2021), enabling users to

206   apply it to be applied by users to their own input networks and environmental models.

We examined model-averaged weights (MAW) and multi-model relative variable

208   importance values (RVI) to gauge performance metrics against input conditions. These were

assessed in three ways: (1) Summarizing among-replicate variation using median RVI and mean

210   MAW; (2) Re-computing model averages as an Akaike-weighted ensemble model, including all

sampled models among replicates; and (3) Re-computing model averages using only the best

212   AIC model from each replicate run. For ensemble models, only models below a cumulative

Akaike threshold of 0.95 were retained.

2.3 / Rhinichthys osculus *as a case study*

216   We focused on the small-bodied *Rhinichthys osculus*, broadly endemic to western North

America within all major drainage basins west of the Rocky Mountains (Hubbs & Miller, 1948;

218   Minckley et al., 1986; Oakey et al., 2004; Smith & Dowling, 2008). As such, it is remarkably

adaptive, with a range extending from high-altitude/ cold-water Pacific Northwest streams

220   (Pfrender et al., 2004; Wiesenfeld et al., 2018), to endorheic basins vulnerable to stochastic and

anthropogenic influences in the Mojave/ Sonoran deserts (Mussmann, Douglas, Oakey, et al.,

222   2020). Click or tap here to enter text.Its extinction vulnerability is underscored by the

contemporaneous elimination of two lineages within the 20[th] century, and four additional

224   subspecies listed under the United States Endangered Species Act of 1973, as amended (16

U.S.C. 1531-1544, 87 Stat. 884). Our case study explores the utility of ResistNet in categorizing

226   extant conservation and management challenged for this species, with a focus on distributions

localised within the Colorado River Basin.

228   A matrix of unlinked single nucleotide polymorphisms (SNPs) was derived from existing

datasets (Mussmann, 2018). Briefly, genomic DNA was extracted from fin tissue non-lethally

230   sampled from *N*=762 individuals distributed throughout 78 Colorado River Basin localities

        (mean=9.8/site) (Figure 2; Supplementary Table S1; Mussmann, 2018). Library preparation

232   followed the ddRAD protocol (Peterson et al. 2012), using *Pst*I and *Msp*I enzymes (New

        England Biolabs) with 144 samples pooled per 1x100bp lane for sequencing on the Illumina

234   HiSeq 2000 (University of Wisconsin Biotechnology Center), or HiSeq 4000 (University of

        Oregon Genomics & Cell Characterization Core Facility).

236   Reads were demultiplexed and assembled via *de novo* clustering in Stacks v.2.41

        (Catchen et al., 2013). Samples were grouped hierarchically into *N*=5 subregions: Upper

238   Colorado River Basin (*N*=221); Virgin River (*N*=223); Grand Canyon and tributaries (*N*=149);

        Lower Colorado River Basin (*N*=105); and Little Colorado River (*N*=64). Trial runs using

240   STACKS yielded optimal parameters (Paris et al., 2017) for minimum number of raw reads needed

        per stack (m=4), as well as number of mismatches between stacks for identification of putative

242   loci (M=2), and catalog formation (n=2). Loci appearing in <50% of individuals per subregion

        were removed, as were loci with an observed heterozygosity >0.6, and a minor allele frequency

244   (MAF) <0.01. These measures effectively removed putative paralogs and minimally informative

        loci (Eaton, 2014; McKinney et al., 2017).

        2.3.1 | *Population structure*

248   To provide context for riverscape resistance and genetic differentiation, we first gauged

        population structure using an aspatial clustering method (ADMIXTURE v1.3.0; Alexander et al.,

250   2009). Data were filtered to one SNP/ locus, with samples partitioned by region. In doing so, we

        first evaluated models containing *K*=1-20 possible sub-populations across 20 replicates, executed

252   in parallel (Mussmann, Douglas, Chafin, et al., 2020). Replicates were clustered and evaluated

(Clumpak; Kopelman et al., 2015) to identify potentially discordant modes within a single *K*

254   value. Major clusters were identified using a threshold of 90%. Optimal *K* was chosen by cross-

validation (Alexander et al., 2009).

2.3.1 | *Riverscape genetics using ResistNet*

258   Inputs for ResistNet were derived from the SNP dataset (formatted as VCF) and an input

shapefile using autoStreamTree (Chafin et al., 2023). Samples were first grouped by locality and

260   mapped to a stream layer representing all of North America, provided by (RiverATLAS; Linke

et al., 2019). Pairwise genetic distances were computed as Weir and Cockerham's F_{ST} (Weir &

262   Cockerham, 1984), linearised (=F_{ST}/1- F_{ST}; Rousset; 1997). We subsequently selected a subset

of environmental attributes (from N=271; RiverATLAS) as potential covariates for ResistNet.

264   We first excluded those redundant, invariant, or at irrelevant spatial scales for this study, then

additionally removed those exhibiting significant collinearity (defined as $r^2 > 0.7$). This resulted

266   in N=20 environmental and anthropogenic covariates (Table 1).

ResistNet was implemented both globally and individually for each subregion, using 20

268   independent replicates for each. We used a population size of N=1000, with other settings

retained from the simulation-based validation.

3. RESULTS

272   3.1 | *Simulation-based validation*

Ensemble models were accurate or contained the correct model in all instances, with exception

274   of sim2 and sim3 for the least topologically complex network (net2; Figure 2), where the correct

model was only chosen by 20-60% of replicates (Supplementary Figure S1; Supplementary

276   Table S2). In these cases, ensemble models erroneously excluded SGR (stream gradient; Figure

         2). In all instances, convergence occurred in <20 iterations, with a minor sample size effect

278   observed in the delayed convergence for cases more sparsely sampled (i.e., N=10 samples;

         Supplementary Figure S2). As a composite across simulated scenarios, the correct model was

280   recovered in >90% of replicates. It occurred in100% of ensemble models for net1 when N=30 or

         50, with lower accuracy at N=10 (65% replicates correct and in 75% of ensemble models;

282   Supplementary Table S3).

               Accuracies were lowest for net2 (50% of ensemble models correct and 50-75%

284   containing the correct model). For net3, which had the greater spatial extent and topological

         complexity, 50-75% of ensemble models were correct, but 100% contained the correct model

286   (Supplementary Table S3). Of note, when covariates were erroneously included, they were

         seemingly selected at a lower weight (Supplementary Figure S3), with much less impact on the

288   model-averaged composite resistances. As above, these cases could generally be identified (and

         subsequently adjudicated) by examining the consensus, or lack thereof, among replicates and

290   ensemble approaches (Figure 2).

               Regarding the methods for summarising replicates, ensemble approaches (ensemble_all

292   and ensemble_best), which employ model-averaging across replicates), offered a means to

         consolidate discordant replicates, particularly in a few instances where only a minority of

294   replicates recovered the true model (Supplementary Figure S1). As such, they generally

         performed better when considered across scenarios (Supplementary Table S3).

296   The most sparsely sampled datasets (N=10 samples) had the highest rate of false

         covariate inclusion (Figure 2), slowest convergence (Supplementary Figure S3), and greatest

298   error in recovering the simulated variable weights (Supplementary Figure S4). They also

displayed reduced variation among replicates in model fitness (measured as AIC; Supplementary

300　Figure S5), and often displayed a stronger correlation of the recovered resistance matrix with the

input genetic distances (assessed as Mantel's r; Supplementary Figure S6). This suggests a

302　reduced accuracy at smaller sample sizes is not necessarily due to a loss of model optimisation

due to insufficient sample, but rather an inability to discriminate amongst multiple model optima

304　which fit equally (or better) than the explicitly simulated conditions. We also note the impact of

sample size on covariate selection varied among the three simulated networks, suggesting

306　potential impacts due to spatial dimension, topology, or local conditions relative to the selected

covariates.

308　　　　The most simple of our simulated models [distance-only (sim1) and a two-covariate

model with equal weights (sim2)] were most consistently recovered (Figure 2). In contrast, the

310　performance of the models across different network complexities was more nuanced. Here, the

least complex topological network (net2) exhibited the lowest accuracy, both in terms of inferred

312　variable importance (Figure 2) and model-averaged weights (Supplementary Figure S3).

314　3.2 | *Empirical case study in* Rhinichthys osculus

3.2.1 | *Assembly and population structure*

316　The global assembly had 13,218 loci with a 25.94x mean depth and 27.6% missing data after

filtering. ADMIXTURE identified $K=30$ populations segregating within the five subregions, and

318　broadly corresponding to major tributaries in all cases (Figure 3; Supplementary Figure S6). The

Upper Colorado River Basin was divided into $K=6$ groups, while the Virgin River Basin was

320　partitioned into $K=7$. The remaining subregions were divided into $K=5$ linearly distributed

through the Grand Canyon, $K=8$ in the Lower Colorado, and $K=4$ in the Little Colorado (Figure

322   3; Supplementary Figure S6).

324   3.2.2 | *Fitted distances and effective resistance networks*

The fitted, linearized F_{ST} values accurately predicted observed pairwise F_{ST} in all subregions

326   (Mantel $r=0.94$-0.99; significant per 1,000 permutations and $p<0.001$), with exception of the

Grand Canyon, where correlation was weaker ($r=0.34$) although still significant ($p<0.001$; Figure

328   3). This indicates that genetic distances mirror network topology, a necessary condition for the

appropriate application of ResistNet. As such, ADMIXTURE results generally align well with

330   network topology such that patterns were concordant with fitted, edge-wise F_{ST} (Figure 3).

        Model-averaged resistances projected onto the stream network (Figure 4) likewise

332   showed a strong concordance with ADMIXTURE probabilities and fitted F_{ST} (Figure 3). As

such, pairwise resistance significantly predicted F_{ST} in all cases ($r=0.34$-0.89) with exception of

334   the Grand Canyon ($r=0.19$). However, resistance models were only marginally better than

distance-only models in most subregions, the exception being the Little Colorado

336   (Supplementary Figure S7).

        Little variation was seen in relative importance (RVI) of covariates when median RVI

338   were compared among replicates using ensemble approaches (Supplementary Figure S8).

However, the implicated covariates (using an RVI threshold of 0.8) varied considerably among

340   subregion (Figure 4), with RUN and TMP commonly selected (each chosen in 4/5 subregions).

        Specific covariates selected using this threshold for each subregion varied between 3-9:

342   Virgin River ($N=4$; RUN, TMP, PRE, CRP), Little Colorado ($N=3$; RDD, RUN, TMP), Grand

Canyon ($N=4$; RUN, TMP, AET, PST), Upper Colorado ($N=3$; TMP, REV, PST), and Lower

344   Colorado (*N*=9; RUN, REV, AET, PRE, RDD, DOR, LKA, POP, SGR). Interestingly,

hydrologic distance had minimal contribution across subregions, with a small RVI only present

346   in the Little Colorado and Upper Colorado subregions (Figure 4), but in a minority of replicates

therein (Supplementary Figure S8).

348   The global model, which combined all subregions, selected 14 covariates representing an

amalgam of those in the individual subregion models (Supplementary Figure S9). However,

350   effective resistances as model-averages did not show a clear spatial correlation with fitted F_{ST}

and offered a comparatively weak explanation for observed population assignment probabilities

352   (Figure 3). Given this, we suggest interpretation should be restricted to subregion analyses.

354   **4. DISCUSSION**

Characterising the spatial distribution of genetic variability within riverscapes serves as a

356   blueprint for the interaction of biodiversity with the environment. Deconstructing this

relationship provides a mechanism that can quantify those features which dictate individual

358   movements, thereby informing conservation efforts (Cayuela et al., 2018; Van Strien et al.,

2014). To promote this, we introduce ResistNet, a novel method tailored to parse these

360   relationships within spatial networks such as riverscapes. Our simulation-based authentication

demonstrates that ResistNet does indeed recover true models of environmental resistance from

362   with high fidelity from pairwise genetic/genomic distances. However, and importantly, our

simulated scenarios are idealised, and inherently meet the analytical assumptions underlying the

364   model. Likewise, our case study applied to *Rhinichthys* in the Colorado River Basin underscores

both its utility and limitations, which we discuss herein.

As with any method, the applicability of ResistNet must be careful considered within the context of the system, with biological characteristics, the landscape/ riverscape, and the scale and

quality of available data being accounted for. Optimization of a resistance model will be poor when 'true' correlations among genetic and resistance distances are minimal (Winiarski et al.,

2020), or when they are poorly distributed according to network hierarchy (Hopken et al., 2013). This underscores the expectation that topology serves as a reasonable constraint for

metapopulation dynamics, a condition unfulfilled for freshwater species with terrestrial or aerial dispersal (Alp et al., 2012; Bohonak & Jenkins, 2003; Phillipsen et al., 2015). Perhaps less

obvious are obligate aquatic species that display highly variable dispersal traits and/or behaviours (Comte & Olden, 2018; Tonkin et al., 2018), with emergent patterns of population

genetic structure displayed as a cascading effect (Zbinden et al., 2022b). Here, intraspecific genetic differentiation is depicted as a spectrum, even among co-distributed species (Zbinden et

al., 2022a), such that it is governed on one hand by differential gene flow within a connected metapopulation, and on the other by isolated patches subject to vicariant processes (Meffe &

Vrijenhoek, 1988).

       A fundamental assumption of ResistNet, and indeed all methods leveraging the concept

382   of 'landscape resistance,' is that individuals act as discrete particles modelled as a 'random-walk,' where flux is dictated by resistance values along movement paths that are, by definition,

384   additive, isotropic, and time-invariant (Kumar et al., 2022; B. H. McRae et al., 2008). Emergent patterns of genetic divergence are thus assumed to follow a stepping-stone model (Kimura &

386   Weiss, 1964),that increases as a monotone function of cumulative resistance. However, genetic distances (serving as model input) display a dynamic evolutionary process over non-

388   contemporary timescales. Historic drainage evolution may thus drive cryptic deviations in the

model, in that intermittent connections, divergences, and/ or tributary captures may yield genetic

390 patterns at odds with modern hydrography (Douglas et al., 1999). Individual movements via

human-mediated translocations will also bias results by altering spatio-genetic associations

392 (Chafin, Zbinden, et al., 2021). It is thus critical to consider both timescale and degree to which

the riverscape serves as an evolutionary template (Epps & Keyghobadi, 2015).

394 Another necessary consideration is the timescale represented by the chosen genetic

distance, which must likewise be appropriate for sampling strategy, genetic marker, and the

396 biological context of the study (Landguth et al., 2010; Shirk et al., 2017). Although

autoStreamTree supports several alternate distance metrics, such as chord distance (Cavalli-

398 Sforza & Edwards, 1967) or Jost's D (Jost, 2008), we did not evaluate them in a comparative

sense as input to ResistNet. It is also important to note that ResistNet implicitly assumes that the

400 contribution of a given environmental feature to effective resistance is spatially homogenous,

and that it varies at a scale relevant to the spatial extent and genomic resolution of the study. This

402 may account for the comparatively lower performance of ResistNet in simulations involving our

smallest and least topologically complex network (i.e., 'Net 2'; Figure 2). Smaller sample sizes

404 also carry a greater risk of spurious correlations (Cushman & Landguth, 2010), a trend

observable in our simulations as well (Figure 2, Supplementary Table S2, S3). Hence, it is

406 crucial that the sampling design is aligned with both genetic and environmental processes under

study (Dauphin et al., 2023). This may translate, for example, to separating samples, thereby

408 more appropriately targeting scale-dependent effects, albeit at a potential loss of statistical power

(Keller et al., 2013; Razgour et al., 2014). We explore this topic in the context of our empirical

410 case study below.

Finally, we emphasise the need for environmental data to be properly curated. The scale

412   and accessibility of such data has recently expanded for riverscape (HydroATLAS; Linke et al.,

2019)., We recommend this database (and the shapefiles within) as a starting point for rapid

414   exploratory analysis using ResistNet models. We note additional datasets collate probability of

intermittence (Messager et al., 2021), and instream barriers (Belletti et al., 2020; Grill et al.,

416   2019) as well. The latter also include several quantitative indices as proxies for the influence of

dams (e.g., a factor driving flow regulation). However, the genetic algorithm used by ResistNet

418   can be computationally expensive with a large number of covariates, despite being mitigated

through parallel execution. Care should thus be taken to include only explanatory covariates.

420   Another pervasive confounding factor is multicollinearity (Graham, 2003), which (as

above) can vary as a function of sampling configuration (Graves et al., 2013; Prunier et al.,

422   2015). We took a commonly-used pairwise, correlative approach to reduce the number of input

features for Rhinichthys (Dormann et al., 2013). Other exclusion techniques taken in this context

424   could involve variation inflation factors (Blair et al., 2013), random forests (Zbinden et al.,

2023), or forward-selection based on redundancy analysis (Chafin et al., 2023); all of which

426   could minimise computational complexity by eliminating redundancy in model space, yet with

the potential for important data to be reduced (Graham, 2003). Other alternative may seek to

428   either engineer synthetic predictors as combinations of collinear variables, or partition unique

versus common effects (Dormann et al., 2013; Prunier et al., 2015). Implementation would

430   require pre-processing. Some engineering objectives are integrated within ResistNet as

alternative aggregation functions. Although the default is to compute variable-specific resistance

432   as a sum of edge-wise values along the pairwise least-cost path, alternatives such as coefficients

of variation or arithmetic/ harmonic means may be selected, though care must be taken that these

434   are appropriate. In summary, effective use of ResistNet requires careful consideration of genetic

       distance metrics, data treatment, as well as judicious management of trade-offs between variable

436   inclusion and computational expense.

438   *Empirical case study in* Rhinichthys

       Our case study not only provides insights specific to our study species, but also general

440   guidelines for applying ResistNet to empirical data such as ours. With regard to drainage-wide

       population structure, all subregions had reaches with high values of fitted-F_{ST} that corresponded

442   with similarly elevated environmental resistance (Figure 3, 4), to include those representing

       relatively well-connected riverscapes. IBR was generally a better predictor than (IBD (Figure 4,

444   Supplementary Figure S7). Hydrologic distances were likewise non-explanatory as a component

       of resistance models, implicating environmentally-mediated connectivity as more important for

446   intraspecific divergences. This, in turn, complements analyses of population structure (Figure 3,

       Supplementary Figure S6), and juxtaposes well with prior observations designating unique and

448   relictual lineages (Mussmann, Douglas, Oakey, et al., 2020; Oakey et al., 2004).

              Numerous covariates strongly implicated for *Rhinichthys* were linked to connectivity

450   though flow, drought tolerance, and seasonal persistence of streams (e.g., RUN, TMP, AET,

       PRE), suggests a role for flow regime/ ephemerality (Figure 4). Unsurprisingly, hydrology is a

452   driver of morphological diversification in numerous freshwater systems (Bruckerhoff &

       Magoulick, 2017; Langerhans, 2008; Meyers & Belk, 2014), and its importance in sustaining

454   native fish assemblages has been documented across southwestern streams (Kominoski et al.,

       2018; Propst et al., 2008; Propst & Gido, 2004). The large weights attributed to variables that

456   define stream persistence clearly reflect the impacts of aridity and persistent drought, both of

which are omnipresent across contemporary desert riverscapes (Comte et al., 2022).

458   Covariates relating to contemporary (e.g., anthropogenic) effects such as impoundment

and flow regulation were also chosen (Figure 4). Here, factors relating to anthropogenic land use

460   [e.g., road density (RDD); pasture extent (PST); population density (POP)]; lake/ reservoir extent

[limnicity (LKA); upstream reservoir volume (REV)], and indices of hydrologic manipulation

462   [degree of regulation (DOR)], most strongly contributed to composite resistance models. Several

are functionally related, such as: Impoundment as a mechanism for flow regulation/ flood

464   dampening (DOR), and determining reservoir volume (REV), channel width, and morphology

(Grams et al., 2020; Walker et al., 2020). Anthropogenic factors promoting habitat instability in

466   the Colorado River have previously been implicated as modulating species boundaries in other

endemic leuciscids (Chafin et al., 2019). These altered conditions may also change the

468   distribution and suitability of low flow refugia, serving in turn to extirpate peripheral populations

(Bezzerides & Bestgen, 2002; Magoulick & Kobza, 2003; Schlosser, 1995). Altered conditions

470   also promote the establishment of non-native species (Dibble et al., 2021; Gido & Propst, 1999;

Whitney et al., 2017), which may in turn elevate predation rates and promote competition (Marsh

472   & Douglas, 1997; Seegert et al., 2014). Here, predation pressure has been shown to curtail

invasion success of *Rhinichthys osculus* post-introduction (Harvey et al., 2004), thus suggesting

474   a potential role for non-natives in biotic resistance.

An observation of broader relevance to the applicability of ResistNet for *Rhinichthys* is

476   the relative fit of optimised resistance networks and, by extension, the applicability of IBR varies

substantially as an explanatory model by scale and subregion. The basin-wide analysis, in

478   particular, was a poor fit (Supplementary Figure S9), which we hypothesized as reflecting the

transection of historical or vicariant processes not captured by extant connectivity. This is

480   evidenced by the apparent 'plateau' in the global relationship of F_{ST} with hydrologic distance

(Supplementary Figure S9b), suggesting the relative importance of genetic drift and gene flow as

482   being scale dependent (Hutchison & Templeton, 1999; Keller et al., 2013). Here, inflated genetic

divergences may also emerge as vestiges of historic barriers, such as Pleistocene lava flows

484   within the Grand Canyon region (Duffield et al., 2006).

        Broad-scale hydrologic evolution, as driven by tectonism and vulcanism, are a hallmark

486   of the region (Minckley et al., 1986), with genetic vestiges also well-characterised in other co-

occuring species (Chafin, Douglas, et al., 2021). More recent population dynamics could also

488   lead to a situation where genetic divergences fail to juxtapose with extant conditions, a possibly

explanation of high divergences displayed by Vermillion Creek, which may have inadvertently

490   escaped large-scale, indiscriminate rotenone poisoning as a mechanism to exterminate non-

native species in the Green River in the 1950s-60s (Holden, 1991). Finally, it is also possible that

492   environmental and/ or anthropogenic components of riverscape resistance may vary at smaller

spatial scales, especially in species widely distributed.

494   Another confounding factor could be admixture involving unsampled or distant sites

(Jaquiéry et al., 2011). For example, intergradation between morphological forms of *Rhinichthys*

496   *osculus* in the Little Colorado and Lower Colorado subregions has been hypothesised as being

driven by historic stream capture along the Mogollon Rim, with historically-transient

498   connections possible as well between some Upper Colorado River Basin tributaries and the Bear

or Snake rivers (Loxterman & Keeley, 2012; Smith & Dowling, 2008). Historic reticulation

500   violates an assumption of our resistance model, in that connectivity does not involve edges

represented by the network. If such effects are indeed unrecognized, then F_{ST}-values fitted along

502 that stretch would be incorrectly inflated. Our strategy herein was to define units for analysis

from apparent population structure (Mussmann, 2018), phylogeny (Oakey et al., 2004), and a

504 biogeographic understanding of the region (Minckley et al., 1986). However, quantitative

approaches to construct input matrices could employ an iterative resampling procedure

506 (Goldberg & Waits, 2010) to identify appropriate spatial thresholds or topological constraints

(Savary et al., 2021).

Conclusion

510 IBR may not be appropriate for all systems – and indeed, our *Rhinichthys* case study

demonstrates the potential for intraspecific variation – and explanatory models may thus be

512 conflated or wholly redundant (Zbinden et al., 2022a). However, ResistNet can achieve very

high accuracies (>95%), when scale is appropriately considered along both spatial and

514 topological dimensions. These limitations serve as a necessary context from which to develop

additional applications, or serve as avenues for future research, as do emerging developments in

516 riverscape genetics (White et al., 2020). One extension for locus-wise analysis (e.g., Chafin et

al., 2023), is to identify functional genomic associations that can promote isolation-by-adaptation

518 rather than environmental resistance or limitation via dispersal (Orsini et al., 2013). Adaptive

genetic variation is a relatively underexplored theme in riverscape genetics (Davis et al., 2018),

520 due largely to a dearth of methods for detecting 'outlier' loci against the backdrop of a unique

autocorrelative structure of networks.

522   Another advantage of ResistNet is the potential to extrapolate and predict genetic

responses in unsampled space. Here, optimized resistance models may be applied across any

524 dendritic ecological network (or riverscapes, as herein), to predict effective resistance in regions

within which genetic data has yet to be obtained. This could promote *a priori* hypotheses of

526   population and phylogeographic structure that can supplement transecting regions with elevated

divergences or inform analyses requiring assignment priors (Hubisz et al., 2009). It could also

528   extrapolate connectivity as an invasive (Lovell et al., 2021), or project temporal changes due to

predicted climate change or other disturbance (Nakajima et al., 2023). Importantly, the approach

530   can be applied to any type of pairwise distance matrix. For example, autoStreamTree could be

applied to examine edge-wise distances or topological importance in generating differences

532   within community data (Eros et al., 2011), with ResistNet then applied to examine how

environmental resistance defines metacommunity structure. Again, any form of spatially

534   structured ecological network can be generalized for such accommodation.

The potential of ResistNet to accelerate automation in riverscape genetics can also

536   encourage large-scale comparative riverscape studies involve variation among species in spatial

or temporal trends, the results of which could vary substantially as a function of trait synergism

538   or life histories (Comte et al., 2014; McManamay & Frimpong, 2015; Troia et al., 2019). While

rare, such comparative studies are critical for testing ecological and evolutionary questions

540   (Blanchet et al., 2020; Fourtune et al., 2016; Paz-Vinas et al., 2018; Pilger et al., 2017; Zbinden

et al., 2022b, 2023). Further, a comparative framework also allows those commonalities driving

542   community-level structure or shared biogeographic breaks to clarify, which may supplement

conservation planning at contemporary spatial or taxonomic scales (Linke et al., 2011; Paz-Vinas

544   et al., 2018), and effectively promote multi-species management objectives (Nielsen et al., 2017).

546   **ACKNOWLEDGEMENTS**

Computational resources were provided by the Arkansas High Performance Computing Center

(AHPCC). MED and MRD are supported by the University of Arkansas via endowments (MED: 21st Chair Century Global Change Biology; MRD: Bruker Life Sciences Professorship). TKC is

supported by the Scottish Government's Rural and Environment Science and Analytical Services Division (RESAS). We also thank the members of numerous state, federal, and tribal entities

throughout the Colorado River Basin who provided tissue samples which enabled this research. The use of trade, product, industry, or firm names is for informative purposes only and does not

constitute an endorsement by the U.S. Government or the U.S. Fish and Wildlife Service. Links to non-Service websites do not imply any official U.S. Fish and Wildlife Service endorsement of

the opinions or ideas expressed therein or guarantee the validity of the information provided. The findings, conclusions, and opinions expressed in this article represent those of the authors, and

do not necessarily represent the views of the U.S. Fish & Wildlife Service.

AUTHOR CONTRIBUTIONS

All authors contributed to study design; SMM, MRD, and MED performed lab work and

562 collected data; SMM performed bioinformatics and assembly; TKC wrote software and designed

methodology; SMM and TKC analysed the data; TKC drafted the initial manuscript. All authors

564 contributed critically to the drafts, interpretation of results, and approved of the manuscript in its

final form.

568 **DATA AVAILABILITY**

All source code for ResistNet is available on GitHub: https://github.com/tkchafin/resistnet.

570   Simulated and empirical datasets used to validate ResistNet, including Snakemake workflows to

replicate analyses, are available via the Open Science Framework: https://osf.io/4uxfj . Raw

572   sequence reads are deposited in the NCBI SRA under BioProject PRJA656098.

REFERENCES

Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome Research*, *19*(9), 1655–1664.

Alp, M., Keller, I., Westram, A. M., & Robinson, C. T. (2012). How river structure and biological traits influence gene flow: A population genetic study of two stream invertebrates with differing dispersal abilities. *Freshwater Biology*, *57*(5), 969–981. https://doi.org/10.1111/j.1365-2427.2012.02758.x

Altermatt, F., & Fronhofer, E. A. (2018). Dispersal in dendritic networks: Ecological consequences on the spatial distribution of population densities. *Freshwater Biology*, *63*(1), 22–32. https://doi.org/10.1111/fwb.12951

Baggio, R. A., Araujo, S. B. L., Ayllón, D., & Boeger, W. A. (2018). Dams cause genetic homogenization in populations of fish that present homing behavior: Evidence from a demogenetic individual-based model. *Ecological Modelling*, *384*, 209–220. https://doi.org/10.1016/j.ecolmodel.2018.06.019

Belletti, B., Garcia de Leaniz, C., Jones, J., Bizzi, S., Börger, L., Segura, G., Castelletti, A., van de Bund, W., Aarestrup, K., Barry, J., Belka, K., Berkhuysen, A., Birnie-Gauvin, K., Bussettini, M., Carolli, M., Consuegra, S., Dopico, E., Feierfeil, T., Fernández, S., … Zalewski, M. (2020). More than one million barriers fragment Europe's rivers. *Nature*, *588*(7838), 436–441. https://doi.org/10.1038/s41586-020-3005-2

Bezzerides, N., & Bestgen, K. (2002). Status review of Roundtail chub (*Gila robusta*), Flannelmouth sucker (*Catostomus latipinnis*), and Bluehead sucker (*Catostomus discobolus*) in the Colorado River Basin. *Final Report to the U.S. Bureau of Reclamation*.

Blair, C., Jiménez Arcos, V. H., Mendez de la Cruz, F. R., & Murphy, R. W. (2013). Landscape genetics of Leaf-Toed Geckos in the tropical dry forest of Northern Mexico. *PLoS ONE*, *8*(2). https://doi.org/10.1371/journal.pone.0057433

Blanchet, S., Prunier, J. G., Paz-Vinas, I., Saint-Pé, K., Rey, O., Raffard, A., Mathieu-Bégné, E., Loot, G., Fourtune, L., & Dubut, V. (2020). A river runs through it: The causes, consequences, and management of intraspecific diversity in river networks. *Evolutionary Applications*, *13*(6), 1195–1213. https://doi.org/10.1111/eva.12941

Bohonak, A. J., & Jenkins, D. G. (2003). Ecological and evolutionary significance of dispersal by freshwater invertebrates. *Ecology Letters*, *6*(8), 783–796. https://doi.org/10.1046/j.1461-0248.2003.00486.x

Bradbury, I. R., & Bentzen, P. (2007). Non-linear genetic isolation by distance: Implications for dispersal estimation in anadromous and marine fish populations. *Marine Ecology Progress Series*, *340*, 245–257. https://doi.org/10.3354/meps340245

Brauer, C. J., & Beheregaray, L. B. (2020). Recent and rapid anthropogenic habitat fragmentation increases extinction risk for freshwater biodiversity. *Evolutionary Applications*, *13*(10), 2857–2869. https://doi.org/10.1111/eva.13128

Brauer, C. J., Hammer, M. P., & Beheregaray, L. B. (2016). Riverscape genomics of a threatened fish across a hydroclimatically heterogeneous river basin. *Molecular Ecology*, *61*, 5093–5113. https://doi.org/10.1111/mec.13830

Brauer, C. J., Unmack, P. J., Smith, S., Bernatchez, L., & Beheregaray, L. B. (2018). On the roles of landscape heterogeneity and environmental variation in determining population genomic structure in a dendritic system. *Molecular Ecology*, *27*(17), 3484–3497. https://doi.org/10.1111/mec.14808

618   Bruckerhoff, L. A., & Magoulick, D. D. (2017). Hydrologic regimes as potential drivers of morphologic divergence in fish. *Evolutionary Ecology*, *31*(4), 517–531.

620   https://doi.org/10.1007/s10682-017-9897-0

Camak, D. T., & Piller, K. R. (2018). Going with the flow: Testing the role of habitat isolation

622   among three ecologically divergent darter species. *Copeia*, *106*(2), 375–387. https://doi.org/10.1643/CG-17-623

624   Campbell Grant, E. H. (2011). Structural complexity, movement bias, and metapopulation extinction risk in dendritic ecological networks. *Journal of the North American*

626   *Benthological Society*, *30*(1), 252–258. https://doi.org/10.1899/09-120.1

Campbell Grant, E. H., Lowe, W. H., & Fagan, W. F. (2007). Living in the branches: Population

628   dynamics and ecological processes in dendritic networks. *Ecology Letters*, *10*(2), 165–175. https://doi.org/10.1111/j.1461-0248.2006.01007.x

630   Cavalli-Sforza, L. L., & Edwards, A. W. (1967). Phylogenetic analysis: Models and estimation procedures. *American Journal of Human Genetics*, *19*(3), 550–570.

632   https://doi.org/10.2307/2406616

Cayuela, H., Rougemont, Q., Prunier, J. G., Moore, J. S., Clobert, J., Besnard, A., & Bernatchez,

634   L. (2018). Demographic and genetic approaches to study dispersal in wild animal populations: A methodological review. *Molecular Ecology*, *27*(20), 3976–4010.

636   https://doi.org/10.1111/mec.14848

Chafin, T. K., Douglas, M. R., Bangs, M. R., Martin, B. T., Mussmann, S. M., & Douglas, M. E.

638   (2021). Taxonomic uncertainty and the anomaly zone: Phylogenomics disentangle a rapid radiation to resolve contentious species (*Gila robusta* complex) in the Colorado River.

640   *Genome Biology and Evolution*, *13*(9), evab200. https://doi.org/10.1101/692509

Chafin, T. K., Douglas, M. R., Martin, B. T., & Douglas, M. E. (2019). Hybridization drives

642   genetic erosion in sympatric desert fishes of western North America. *Heredity*, *123*, 759–773.

644   Chafin, T. K., Mussmann, S. M., Douglas, M. R., & Douglas, M. E. (2023). autoStreamTree: genomic variant data fitted to geospatial networks . *BioRxiv*, *542562*.

646   Chafin, T. K., Zbinden, Z. D., Douglas, M. R., Martin, B. T., Middaugh, C. R., Gray, M. C., Ballard, J. R., & Douglas, M. E. (2021). Spatial population genetics in heavily managed

648   species: Separating patterns of historical translocation from contemporary gene flow in white-tailed deer. *Evolutionary Applications*, *14*(6), 1673–1689.

650   https://doi.org/10.1111/eva.13233

Chiu, M. C., Li, B., Nukazawa, K., Resh, V. H., Carvajal, T., & Watanabe, K. (2020). Branching

networks can have opposing influences on genetic variation in riverine metapopulations. *Diversity and Distributions*, *26*(12), 1813–1824. https://doi.org/10.1111/ddi.13160

Chiu, M. C., Nukazawa, K., Carvajal, T., Resh, V. H., Li, B., & Watanabe, K. (2020). Simulation modeling reveals the evolutionary role of landscape shape and species dispersal on genetic

variation within a metapopulation. *Ecography*, *43*(12), 1891–1901. https://doi.org/10.1111/ecog.05118

Coleman, R. A., Gauffre, B., Pavlova, A., Beheregaray, L. B., Kearns, J., Lyon, J., Sasaki, M., Leblois, R., Sgro, C., & Sunnucks, P. (2018). Artificial barriers prevent genetic recovery of

small isolated populations of a low-mobility freshwater fish. *Heredity*, *120*(6), 515–532. https://doi.org/10.1038/s41437-017-0008-3

Comte, L., Murienne, J., & Grenouillet, G. (2014). Species traits and phylogenetic conservatism of climate-induced range shifts in stream fishes. *Nature Communications*, *5*, 2023.
https://doi.org/10.1038/ncomms6053

Comte, L., & Olden, J. D. (2018). Evidence for dispersal syndromes in freshwater fishes.
666 *Proceedings of the Royal Society B: Biological Sciences*, *285*, 20172214. https://doi.org/10.1098/rspb.2017.2214

668 Comte, L., Olden, J. D., Lischka, S., & Dickson, B. G. (2022). Multi-scale threat assessment of riverine ecosystems in the Colorado River Basin. *Ecological Indicators*, *138*.
670 https://doi.org/10.1016/j.ecolind.2022.108840

Cooke, G. M., Landguth, E. L., & Beheregaray, L. B. (2014). Riverscape genetics identifies
672 replicated ecological divergence across an Amazonian ecotone. *Evolution*, *68*(7), 1947–1960.

674 Cushman, S. A., & Landguth, E. L. (2010). Spurious correlations and inference in landscape genetics. *Molecular Ecology*, *19*(17), 3592–3602.

676 Dauphin, B., Rellstab, C., Wüest, R. O., Karger, D. N., Holderegger, R., Gugerli, F., & Manel, S. (2023). Re-thinking the environment in landscape genomics. *Trends in Ecology and*
678 *Evolution*, *38*(3), 261–274. https://doi.org/10.1016/j.tree.2022.10.010

Davis, C. D., Epps, C. W., Flitcroft, R. L., & Banks, M. A. (2018). Refining and defining
680 riverscape genetics: How rivers influence population genetic structure. *WIREs Water*, *5*(2), e1269.

682 De Rainville, F.-M., Fortin, F.-A., Gardner, M.-A., Parizeau, M., & Gagné, C. (2012). Deap: A python framework for evolutionary algorithms. *Proceedings of the 14th Annual Conference*
684 *Companion on Genetic and Evolutionary Computation*, 85–92.

Dibble, K. L., Yackulic, C. B., Schmidt, J. C., Kennedy, T. A., & Bestgen, K. R. (2021). Water
686 storage decisions will determine the distribution and persistence of imperiled river fishes. *Ecological Applications*, *31*(2), e02279. https://doi.org/10.1002/eap.2279

688 Dormann, C. F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J. R. G., Gruber, B., Lafourcade, B., Leitão, P. J., Münkemüller, T., Mcclean, C., Osborne, P. E.,
690 Reineking, B., Schröder, B., Skidmore, A. K., Zurell, D., & Lautenbach, S. (2013). Collinearity: A review of methods to deal with it and a simulation study evaluating their
692 performance. *Ecography*, *36*(1), 27–46. https://doi.org/10.1111/j.1600-0587.2012.07348.x

Douglas, M. E., & Endler, J. A. (1982). Quantitative matrix comparisons in ecological and
694 evolutionary investigations. *Journal of Theoretical Biology*, *99*, 777–795. https://doi.org/doi: 10.1016/0022-5193(82)90197-7

696 Douglas, M. E., & Matthews, W. J. (1992). Does morphology predict ecology? Hypothesis testing within a freshwater stream fish assemblage. *Oikos*, *65*(2), 213.
698 https://doi.org/10.2307/3545012

Douglas, M. E., Minckley, W. L., & DeMarais, B. D. (1999). Did vicariance mold phenotypes of
western North American fishes? Evidence from Gila River cyprinids. *Evolution*, *53*(1), 238–246.

Duffield, W., Riggs, N., Kaufman, D., Champion, D., Fenton, C., Forman, S., McIntosh, W., Hereford, R., Plescia, J., & Ort, M. (2006). Multiple constraints on the age of a Pleistocene
lava dam across the Little Colorado River at Grand Falls, Arizona. *Bulletin of the Geological Society of America*, *118*(3–4), 421–429. https://doi.org/10.1130/B25814.1

Eaton, D. A. R. (2014). PyRAD: assembly of de novo RADseq loci for phylogenetic analyses. *Bioinformatics*, *30*(13), 1844–1849.

Epps, C. W., & Keyghobadi, N. (2015). Landscape genetics in a changing world: Disentangling historical and contemporary influences and inferring change. *Molecular Ecology*, *24*(24),
6021–6040.

    Eros, T., Schmera, D., & Schick, R. S. (2011). Network thinking in riverscape conservation - A
712   graph-based approach. *Biological Conservation*, *144*(1), 184–192. https://doi.org/10.1016/j.biocon.2010.08.013

714   Fagan, W. F. (2002). Connectivity, fragmentation, and extinction risk in dendritic metapopulations. *Ecology*, *83*(12), 3243–3249. https://doi.org/10.1890/0012-
716   9658(2002)083[3243:CFAERI]2.0.CO;2

    Fagan, W., Unmack, P., Burgess, C., & Minckley, W. (2002). Rarity, fragmentation, and
718   extinction risk in desert fishes. *Ecology*, *82*(12), 3250–3256. https://doi.org/10.1890/0012-9658(2002)083[3250:RFAERI]2.0.CO;2

720   Falke, J. A., Bailey, L. L., Fausch, K. D., & Bestgen, K. R. (2012). Colonization and extinction in dynamic habitats: An occupancy approach for a Great Plains stream fish assemblage.
722   *Ecology*, *93*(4), 858–867. https://doi.org/10.1890/11-1515.1

    Fourtune, L., Paz-Vinas, I., Loot, G., Prunier, J. G., & Blanchet, S. (2016). Lessons from the
724   fish: a multi-species analysis reveals common processes underlying similar species-genetic diversity correlations. *Freshwater Biology*, *61*(11), 1830–1845.
726   https://doi.org/10.1111/fwb.12826

    Fraser, G. S., Bestgen, K. R., Winkelman, D. L., & Thompson, K. G. (2019). Temperature—not
728   flow—predicts native fish reproduction with implications for climate change. *Transactions of the American Fisheries Society*, *148*(3), 509–527. https://doi.org/10.1002/tafs.10151

730   Frissell, C. A., Liss, W. J., Warren, C. E., & Hurley, M. D. (1986). A hierarchical framework for stream habitat classification: Viewing streams in a watershed context. *Environmental*
732   *Management*, *10*, 199–214. https://doi.org/10.1007/BF01867358

    Fullerton, A. H., Burnett, K. M., Steel, E. A., Flitcroft, R. L., Pess, G. R., Feist, B. E., Torgersen,
734   C. E., Miller, D. J., & Sanderson, B. L. (2010). Hydrological connectivity for riverine fish: Measurement challenges and research opportunities. *Freshwater Biology*, *55*(11), 2215–
736   2237. https://doi.org/10.1111/j.1365-2427.2010.02448.x

    Galbraith, H. S., Zanatta, D. T., & Wilson, C. C. (2015). Comparative analysis of riverscape
738   genetic structure in rare, threatened and common freshwater mussels. *Conservation Genetics*, *16*(4), 845–857. https://doi.org/10.1007/s10592-015-0705-5

740   Giam, X., & Olden, J. D. (2016). Quantifying variable importance in a multimodel inference framework. *Methods in Ecology and Evolution*, *7*(4), 388–397.
742   https://doi.org/10.1111/2041-210X.12492

    Gido, K. B., & Propst, D. L. (1999). Habitat use and association of native and nonnative fishes in
744   the San Juan River, New Mexico and Utah. *Copeia*, *2*, 321–332. https://doi.org/10.2307/1447478

746   Goldberg, C. S., & Waits, L. P. (2010). Comparative landscape genetics of two pond-breeding amphibian species in a highly modified agricultural landscape. *Molecular Ecology*, *19*(17),
748   3650–3663. https://doi.org/10.1111/j.1365-294X.2010.04673.x

    Graham, M. H. (2003). Confronting multicollinearity in ecological multiple regression. *Ecology*,
750   *84*(11), 2809–2815. https://doi.org/10.1890/02-3114

    Grams, P. E., Dean, D. J., Walker, A. E., Kasprak, A., & Schmidt, J. C. (2020). The roles of
752   flood magnitude and duration in controlling channel width and complexity on the Green

River in Canyonlands, Utah, USA. *Geomorphology*, *371*, 107438.
754   https://doi.org/10.1016/j.geomorph.2020.107438

Graves, T. A., Beier, P., & Royle, J. A. (2013). Current approaches using genetic distances
756   produce poor estimates of landscape resistance to interindividual dispersal. *Molecular
         Ecology*, *22*(15), 3888–3903. https://doi.org/10.1111/mec.12348

758 Grill, G., Lehner, B., Thieme, M., Geenen, B., Tickner, D., Antonelli, F., Babu, S., Borrelli, P.,
         Cheng, L., Crochetiere, H., Ehalt Macedo, H., Filgueiras, R., Goichot, M., Higgins, J.,
760   Hogan, Z., Lip, B., McClain, M. E., Meng, J., Mulligan, M., … Zarfl, C. (2019). Mapping
         the world's free-flowing rivers. *Nature*, *569*(7755), 215–221.
762   https://doi.org/10.1038/s41586-019-1111-9

Grummer, J. A., Beheregaray, L. B., Bernatchez, L., Hand, B. K., Luikart, G., Narum, S. R., &
764   Taylor, E. B. (2019). Aquatic landscape genomics and environmental effects on genetic
         variation. *Trends in Ecology & Evolution*, *34*(7), 641–654.
766   https://doi.org/10.1016/j.tree.2019.02.013

Harvey, B. C., White, J. L., & Nakamoto, R. J. (2004). An emergent multiple predator effect may
768   enhance biotic resistance in a stream fish assemblage. *Ecology*, *85*(1), 127–133.
         https://doi.org/10.1890/03-3018

770 Harvey, E., Harvey, M. G., Singhal, S., & Rabosky, D. L. (2019). Beyond reproductive isolation:
         Demographic controls on the speciation process. *Annual Review of Ecology, Evolution, and
772   Systematics*, *50*, 75–95.

Holden, P. B. (1991). Ghosts of the Green River: impacts of Green River poisoning on
774   management of native fishes. In W. L. Minckley, J. E. Deacon, & S. L. Udall (Eds.), *Battle
         Against Extinction: Native Fish Management in the American West* (pp. 43–54). University
776   of Arizona Press. https://doi.org/10.1016/0006-3207(93)90090-n

Hopken, M. W., Douglas, M. R., & Douglas, M. E. (2013). Stream hierarchy defines riverscape
778   genetics of a North American desert fish. *Molecular Ecology*, *22*, 956–971.
         https://doi.org/10.1111/mec.12156

780 Hubbs, C. L., & Miller, R. R. (1948). The zoological evidence: correlation between fish
         distribution and hydrogeographic history in the desert basin of western North America.
782   *Bulletin of the University of Utah*, *10*(7), 17–166.

Hubisz, M. J., Falush, D., Stephens, M., & Pritchard, J. K. (2009). Inferring weak population
784   structure with the assistance of sample group information. *Molecular Ecology Resources*,
         9(5), 1322–1332. https://doi.org/10.1111/j.1755-0998.2009.02591.x

786 Hughes, J. M., Schmidt, D. J., & Finn, D. S. (2009). Genes in streams: Using DNA to understand
         the movement of freshwater fauna and their riverine habitat. *BioScience*, *59*(7), 573–583.
788   https://doi.org/10.1525/bio.2009.59.7.8

Hutchison, D. W., & Templeton, A. R. (1999). Correlation of pairwise genetic and geographic
790   distance measures: Inferring the relative influences of gene flow and drift on the distribution
         of genetic variability. *Evolution*, *53*(6), 1898–1914. https://doi.org/10.1111/j.1558-
792   5646.1999.tb04571.x

Jackson, D. A., Peres-Neto, P. R., & Olden, J. D. (2001). What controls who is where in
794   freshwater fish communities - The roles of biotic, abiotic, and spatial factors. *Canadian
         Journal of Fisheries and Aquatic Sciences*, *58*(1), 157–170. https://doi.org/10.1139/cjfas-
796   58-1-157

Jaquiéry, J., Broquet, T., Hirzel, A. H., Yearsley, J., & Perrin, N. (2011). Inferring landscape
effects on dispersal from genetic distances: How far can we go? *Molecular Ecology*, *20*(4),
692–705. https://doi.org/10.1111/j.1365-294X.2010.04966.x

Jost, L. (2008). Gst and its relatives do not measure differentiation. *Molecular Ecology*, *17*(18),
4015–4026. https://doi.org/10.1111/j.1365-294X.2008.03887.x

Kanno, Y., Vokoun, J. C., & Letcher, B. H. (2011). Fine-scale population structure and
riverscape genetics of brook trout (*Salvelinus fontinalis*) distributed continuously along
headwater channel networks. *Molecular Ecology*, *20*(18), 3711–3729.
https://doi.org/10.1111/j.1365-294X.2011.05210.x

Keis, M., Remm, J., Ho, S. Y. W., Davison, J., Tammeleht, E., Tumanov, I. L., Saveljev, A. P.,
Männil, P., Kojola, I., Abramov, A. V., Margus, T., & Saarma, U. (2013). Complete
mitochondrial genomes and a novel spatial genetic method reveal cryptic phylogeographical
structure and migration patterns among brown bears in north-western Eurasia. *Journal of
Biogeography*, *40*(5), 915–927. https://doi.org/10.1111/jbi.12043

Keller, D., Holderegger, R., & Van Strien, M. J. (2013). Spatial scale affects landscape genetic
analysis of a wetland grasshopper. *Molecular Ecology*, *22*(9), 2467–2482.
https://doi.org/10.1111/mec.12265

Kimmig, S. E., Beninde, J., Brandt, M., Schleimer, A., Kramer-Schadt, S., Hofer, H., Börner, K.,
Schulze, C., Wittstatt, U., Heddergott, M., Halczok, T., Staubach, C., & Frantz, A. C.
(2020). Beyond the landscape: Resistance modelling infers physical and behavioural gene
flow barriers to a mobile carnivore across a metropolitan area. *Molecular Ecology*, *29*(3),
466–484.

Kimura, M., & Weiss, G. H. (1964). The stepping stone model of population structure and the
decrease of genetic correlation with distance. *Genetics*, *49*, 561–576.

Kominoski, J. S., Ruhí, A., Hagler, M. M., Petersen, K., Sabo, J. L., Sinha, T.,
Sankarasubramanian, A., & Olden, J. D. (2018). Patterns and drivers of fish extirpations in
rivers of the American Southwest and Southeast. *Global Change Biology*, *24*(3), 1175–
1185. https://doi.org/10.1111/gcb.13940

Kopelman, N. M., Mayzel, J., Jakobsson, M., Rosenberg, N. A., & Mayrose, I. (2015). Clumpak:
a program for identifying clustering modes and packaging population structure inferences
across K. *Molecular Ecology Resources*, *15*(5), 1179–1191.

Kumar, S. U., Turnbull, J., Davies, O. H., Hodgetts, T., & Cushman, S. A. (2022). Moving
beyond landscape resistance: considerations for the future of connectivity modelling and
conservation science. *Landscape Ecology*, *37*(10), 2465–2480.
https://doi.org/10.1007/s10980-022-01504-x

Labonne, J., Ravigné, V., Parisi, B., & Gaucherel, C. (2008). Linking dendritic network
structures to population demogenetics: The downside of connectivity. *Oikos*, *117*(10),
1479–1490. https://doi.org/10.1111/j.0030-1299.2008.16976.x

Landguth, E. L., Cushman, S. A., Schwartz, M. K., McKelvey, K. S., Murphy, M., & Luikart, G.
(2010). Quantifying the lag time to detect barriers in landscape genetics. *Molecular
Ecology*, *19*(19), 4179–4191.

Langerhans, R. B. (2008). Predictability of phenotypic differentiation across flow regimes in
fishes. *Integrative and Comparative Biology*, *48*(6), 750–768.
https://doi.org/10.1093/icb/icn092

Linke, S., Lehner, B., Ouellet Dallaire, C., Ariwi, J., Grill, G., Anand, M., Beames, P., Burchard-
Levine, V., Maxwell, S., Moidu, H., Tan, F., & Thieme, M. (2019). Global hydro-

environmental sub-basin and river reach characteristics at high spatial resolution. *Scientific Data*, *6*(1), 283. https://doi.org/10.1038/s41597-019-0300-6

Linke, S., Turak, E., & Nel, J. (2011). Freshwater conservation planning: The case for systematic approaches. *Freshwater Biology*, *56*(1), 6–20. https://doi.org/10.1111/j.1365-2427.2010.02456.x

Lovell, R. S. L., Blackburn, T. M., Dyer, E. E., & Pigot, A. L. (2021). Environmental resistance predicts the spread of alien species. *Nature Ecology and Evolution*, *5*, 322–329. https://doi.org/10.1038/s41559-020-01376-x

Lowe, W. H., Likens, G. E., McPeek, M. A., & Buso, D. C. (2006). Linking direct and indirect data on dispersal: Isolation by slope in a headwater stream salamander. *Ecology*, *87*(2), 334–339. https://doi.org/10.1890/05-0232

Loxterman, J. L., & Keeley, E. R. (2012). Watershed boundaries and geographic isolation: Patterns of diversification in cutthroat trout from western North America. *BMC Evolutionary Biology*, *12*(1), 38. https://doi.org/10.1186/1471-2148-12-38

Ma, C., Shen, Y., Bearup, D., Fagan, W. F., & Liao, J. (2020). Spatial variation in branch size promotes metapopulation persistence in dendritic river networks. *Freshwater Biology*, *65*(3), 426–434. https://doi.org/10.1111/fwb.13435

Magoulick, D. D., & Kobza, R. M. (2003). The role of refugia for fishes during drought: A review and synthesis. *Freshwater Biology*, *48*(7), 1186–1198. https://doi.org/10.1046/j.1365-2427.2003.01089.x

Manel, S., Guerin, P. E., Mouillot, D., Blanchet, S., Velez, L., Albouy, C., & Pellissier, L. (2020). Global determinants of freshwater and marine fish genetic diversity. *Nature Communications*, *11*(1), 1–9. https://doi.org/10.1038/s41467-020-14409-7

Mari, L., Casagrandi, R., Bertuzzo, E., Rinaldo, A., & Gatto, M. (2014). Metapopulation persistence and species spread in river networks. *Ecology Letters*, *17*(4), 426–434. https://doi.org/10.1111/ele.12242

Marsh, P. C., & Douglas, M. E. (1997). Predation by introduced fishes on endangered humpback chub and other native species in the Little Colorado River, Arizona. *Transactions of the American Fisheries Society*, *126*(April 2016), 343–346. https://doi.org/10.1577/1548-8659(1997)126<0343:PBIFOE>2.3.CO;2

Martinez, A. S., Willoughby, J. R., & Christie, M. R. (2018). Genetic diversity in fishes is influenced by habitat type and life-history variation. *Ecology and Evolution*, *8*(23), 12022–12031. https://doi.org/10.1002/ece3.4661

McKinney, G. J., Larson, W. A., Seeb, L. W., & Seeb, J. E. (2017). RADseq provides unprecedented insights into molecular ecology and evolutionary genetics: comment on Breaking RAD by Lowry et al (2016). *Molecular Ecology*, *17*(3), 356–361. https://doi.org/10.1111/ijlh.12426

McManamay, R. A., & Frimpong, E. A. (2015). Hydrologic filtering of fish life history strategies across the United States: Implications for stream flow alteration. *Ecological Applications*, *25*(1), 243–263. https://doi.org/10.1890/14-0247.1

McManamay, R. A., Troia, M. J., DeRolph, C. R., Sheldon, A. O., Barnett, A. R., Kao, S. C., & Anderson, M. G. (2018). A stream classification system to explore the physical habitat diversity and anthropogenic impacts in riverscapes of the eastern United States. *PLoS ONE*, *13*(6), e0198439. https://doi.org/10.1371/journal.pone.0198439

McRae, B. H. (2006). Isolation By Resistance. *Evolution*, *60*(8), 1551. https://doi.org/10.1554/05-321.1

890 McRae, B. H., & Beier, P. (2007). Circuit theory predicts gene flow in plant and animal populations. *Proceedings of the National Academy of Sciences of the United States of America*, *104*(50), 19885–19890. https://doi.org/10.1073/pnas.0706568104

892 McRae, B. H., Dickson, B. G., Keitt, T. H., & Shah, V. B. (2008). Using circuit theory to model connectivity in ecology, evolution, and conservation. *Ecology*, *89*(10), 2712–2724.

894 https://doi.org/10.1890/07-1861.1

McRae, B., Shah, V., & Edelman, A. (2016). Circuitscape: modeling landscape connectivity to

896 promote conservation and human health. *The Nature Conservancy*.

Meffe, G. K., & Vrijenhoek, R. C. (1988). Conservation genetics in the management of desert

898 fishes. *Conservation Biology*, *2*(2), 157–169.

Messager, M. L., Lehner, B., Cockburn, C., Lamouroux, N., Pella, H., Snelder, T., Tockner, K.,

Trautmann, T., Watt, C., & Datry, T. (2021). Global prevalence of non-perennial rivers and streams. *Nature*, *594*(7863), 391–397. https://doi.org/10.1038/s41586-021-03565-5

Meyers, P. J., & Belk, M. C. (2014). Shape variation in a benthic stream fish across flow regimes. *Hydrobiologia*, *738*(1), 147–154. https://doi.org/10.1007/s10750-014-1926-1

Minckley, W. L., Hendrickson, D. A., & Bond, C. E. (1986). Geography of western North American freshwater fishes: description and relationships to intracontinental tectonism. In

C. H. Hocutt & E. O. Wiley (Eds.), *The Zoogeography of North American Freshwater Fishes* (pp. 519–613). John Wiley and Sons.

Minckley, W. L., & Deacon, J. E. (1968). Southwestern fishes and the enigma of 'endangered species'. *Science*, *159*(3822), 1424–1432.

Mölder, F., Jablonski, K. P., Letcher, B., Hall, M. B., Tomkins-Tinch, C. H., Sochat, V., Forster, J., Lee, S., Twardziok, S. O., Kanitz, A., Wilm, A., Holtgrewe, M., Rahmann, S., Nahnsen,

S., & Köster, J. (2021). Sustainable data analysis with Snakemake. *F1000Research*, *10*, 33. https://doi.org/10.12688/f1000research.29032.1

Morrissey, M. B., & De Kerckhove, D. T. (2009). The maintenance of genetic variation due to asymmetric gene flow in dendritic metapopulations. *American Naturalist*, *174*(6), 875–889.

https://doi.org/10.1086/648311

Mussmann, S. M. (2018). Diversification across a dynamic landscape: Phylogeography and

918 riverscape genetics of Speckled Dace (*Rhinichthys osculus*) in Western North America. *ProQuest Dissertations and Theses*.

920 Mussmann, S. M., Douglas, M. R., Chafin, T. K., & Douglas, M. E. (2020). AdmixPipe: Population analyses in ADMIXTURE for non-model organisms. *BMC Bioinformatics*, *21*,

922 337.

Mussmann, S. M., Douglas, M. R., Oakey, D. D., & Douglas, M. E. (2020). Defining relictual

924 biodiversity: Conservation units in speckled dace (Leuciscidae: *Rhinichthys osculus*) of the Greater Death Valley ecosystem. *Ecology and Evolution*, *10*(19), 10798–10817.

926 https://doi.org/10.1002/ece3.6736

Nadeau, T. L., & Rains, M. C. (2007). Hydrological connectivity of headwaters to downstream

928 waters: Introduction to the featured collection. *Journal of the American Water Resources Association*, *43*(1), 1–4. https://doi.org/10.1111/j.1752-1688.2007.00001.x

930 Nakajima, S., Suzuki, H., Nakatsugawa, M., Matsuo, A., Hirota, S. K., Suyama, Y., & Nakamura, F. (2023). Inferring future changes in gene flow under climate change in

932 riverscapes: a pilot case study in fluvial sculpin. *Landscape Ecology*, *38*(5), 1351–1362. https://doi.org/10.1007/s10980-023-01633-x

934 Nielsen, E. S., Beger, M., Henriques, R., Selkoe, K. A., & von der Heyden, S. (2017). Multispecies genetic objectives in spatial conservation planning. *Conservation Biology*,
936 *31*(4), 872–882. https://doi.org/10.1111/cobi.12875

Nilsson, C., Reidy, C. A., Dynesius, M., & Revenga, C. (2005). Fragmentation and flow
938 regulation of the world's large river systems. *Science*, *308*(5720), 405–408. https://doi.org/10.1126/science.1107887

940 Oakey, D. D., Douglas, M. E., & Douglas, M. R. (2004). Small fish in a large landscape: Diversification of *Rhinichthys osculus* (Cyprinidae) in western North America. *Copeia*,
942 *2004*(2), 207–221. https://doi.org/10.1643/CG-02-264R1

Orsini, L., Vanoverbeke, J., Swillen, I., Mergeay, J., & De Meester, L. (2013). Drivers of
944 population genetic differentiation in the wild: Isolation by dispersal limitation, isolation by adaptation and isolation by colonization. *Molecular Ecology*, *22*(24), 5983–5999.
946 https://doi.org/10.1111/mec.12561

Osborne, M. J., Perkin, J. S., Gido, K. B., & Turner, T. F. (2014). Comparative riverscape
948 genetics reveals reservoirs of genetic diversity for conservation and restoration of Great Plains fishes. *Molecular Ecology*, *23*(23), 5663–5679. https://doi.org/10.1111/mec.12970

950 Paris, J. R., Stevens, J. R., & Catchen, J. M. (2017). Lost in parameter space: A road map for stacks. *Methods in Ecology and Evolution*, *8*(10), 1360–1373. https://doi.org/10.1111/2041-
952 210X.12775

Paz-Vinas, I., & Blanchet, S. (2015). Dendritic connectivity shapes spatial patterns of genetic
954 diversity: A simulation-based study. *Journal of Evolutionary Biology*, *28*(4), 986–994. https://doi.org/10.1111/jeb.12626

956 Paz-Vinas, I., Loot, G., Hermoso, V., Veyssiere, C., Poulet, N., Grenouillet, G., & Blanchet, S. (2018). Systematic conservation planning for intraspecific genetic diversity. *Proceedings of
958 the Royal Society B: Biological Sciences*, *285*, 20172746. https://doi.org/10.1101/105544

Paz-Vinas, I., Loot, G., Stevens, V. M., & Blanchet, S. (2015). Evolutionary processes driving
960 spatial patterns of intraspecific genetic diversity in river ecosystems. *Molecular Ecology*, *24*(18), 4586–4604. https://doi.org/10.1111/mec.13345

962 Peterman, W. E. (2018). ResistanceGA: An R package for the optimization of resistance surfaces using genetic algorithms. *Methods in Ecology and Evolution*, *9*(6), 1638–1647.
964 https://doi.org/10.1111/2041-210X.12984

Peterman, W. E., & Pope, N. S. (2021). The use and misuse of regression models in landscape
966 genetic analyses. *Molecular Ecology*, *30*(1), 37–47. https://doi.org/10.1111/mec.15716

Peterman, W. E., Winiarski, K. J., Moore, C. E., Carvalho, C. da S., Gilbert, A. L., & Spear, S. F.
968 (2019). A comparison of popular approaches to optimize landscape resistance surfaces. *Landscape Ecology*, *34*(9), 2197–2208. https://doi.org/10.1007/s10980-019-00870-3

970 Peterson, E. E., ver Hoef, J. M., Isaak, D. J., Falke, J. A., Fortin, M. J., Jordan, C. E., McNyset, K., Monestiez, P., Ruesch, A. S., Sengupta, A., Som, N., Steel, E. A., Theobald, D. M.,
972 Torgersen, C. E., & Wenger, S. J. (2013). Modelling dendritic ecological networks in space: An integrated network perspective. *Ecology Letters*, *16*(5), 707–719.
974 https://doi.org/10.1111/ele.12084

Pfrender, M. E., Hicks, J., & Lynch, M. (2004). Biogeographic patterns and current distribution
of molecular-genetic variation among populations of speckled dace, *Rhinichthys osculus* (Girard). *Molecular Phylogenetics and Evolution*, *30*(3), 490–502.
https://doi.org/10.1016/S1055-7903(03)00242-2

Phillipsen, I. C., Kirk, E. H., Bogan, M. T., Mims, M. C., Olden, J. D., & Lytle, D. A. (2015). Dispersal ability and habitat requirements determine landscape-level genetic patterns in desert aquatic insects. *Molecular Ecology*, *24*(1), 54–69. https://doi.org/10.1111/mec.13003

Pilger, T. J., Gido, K. B., Propst, D. L., Whitney, J. E., & Turner, T. F. (2017). River network architecture, genetic effective size and distributional patterns predict differences in genetic
structure across species in a dryland stream fish community. *Molecular Ecology*, *26*(10), 2687–2697. https://doi.org/10.1111/mec.14079

Propst, D. L., & Gido, K. B. (2004). Responses of native and nonnative fishes to natural flow regime mimicry in the San Juan River. *Transactions of the American Fisheries Society*,
133(4), 922–931. https://doi.org/10.1577/t03-057.1

Propst, D. L., Gido, K. B., & Stefferud, J. A. (2008). Natural Flow Regimes , Nonnative Fishes ,
and Native Fish Persistence in Arid-Land River Systems. *Ecological Applications*, *18*(5), 1236–1252.

Prügel-Bennett, A. (2004). When a genetic algorithm outperforms hill-climbing. *Theoretical Computer Science*, *320*(1), 135–153. https://doi.org/10.1016/j.tcs.2004.03.038

Prunier, J. G., Colyn, M., Legendre, X., Nimon, K. F., & Flamand, M. C. (2015). Multicollinearity in spatial genetics: Separating the wheat from the chaff using
commonality analyses. *Molecular Ecology*, *24*(2), 263–283. https://doi.org/10.1111/mec.13029

Prunier, J. G., Dubut, V., Loot, G., Tudesque, L., & Blanchet, S. (2018). The relative contribution of river network structure and anthropogenic stressors to spatial patterns of
genetic diversity in two freshwater fishes: A multiple-stressors approach. *Freshwater Biology*, *63*(1), 6–21. https://doi.org/10.1111/fwb.13034

Razgour, O., Rebelo, H., Puechmaille, S. J., Juste, J., Ibáñez, C., Kiefer, A., Burke, T., Dawson, D. A., & Jones, G. (2014). Scale-dependent effects of landscape variables on gene flow and
population structure in bats. *Diversity and Distributions*, *20*(10), 1173–1185. https://doi.org/10.1111/ddi.12200

Rodríguez-González, P. M., García, C., Albuquerque, A., Monteiro-Henriques, T., Faria, C., Guimarães, J. B., Mendonça, D., Simões, F., Ferreira, M. T., Mendes, A., Matos, J., &
Almeida, M. H. (2019). A spatial stream-network approach assists in managing the remnant genetic diversity of riparian forests. *Scientific Reports*, *9*(1), 1–10.
https://doi.org/10.1038/s41598-019-43132-7

Rolls, R. J., Heino, J., & Chessman, B. C. (2016). Unravelling the joint effects of flow regime,
1012 climatic variability and dispersal mode on beta diversity of riverine communities. *Freshwater Biology*, *61*(8), 1350–1364. https://doi.org/10.1111/fwb.12793

1014 Rousset, F. (1997). Genetic differentiation and estimation of gene flow from F-statistics under isolation by distance. *Genetics*, *145*, 1219–1228.

1016 Savary, P., Foltête, J. C., Moal, H., Vuidel, G., & Garnier, S. (2021). Analysing landscape effects on dispersal networks and gene flow with genetic graphs. *Molecular Ecology Resources*,
1018 *21*(4), 1167–1185. https://doi.org/10.1111/1755-0998.13333

Schlosser, I. J. (1995). Critical landscape attributes that influence fish population dynamics in
1020 headwater streams. *Hydrobiologia*, *303*(1–3), 71–81. https://doi.org/10.1007/BF00034045

Seegert, S. E. Z., Rosi-Marshall, E. J., Baxter, C. V., Kennedy, T. A., Hall, R. O., & Cross, W. F.
1022 (2014). High diet overlap between native small-bodied fishes and nonnative fathead minnow in the Colorado River, Grand Canyon, Arizona. *Transactions of the American*
1024 *Fisheries Society*, *143*(4), 1072–1083. https://doi.org/10.1080/00028487.2014.901250

Seymour, M., Fronhofer, E. A., & Altermatt, F. (2015). Dendritic network structure and dispersal affect temporal dynamics of diversity and species persistence. *Oikos*, *124*(7), 908–916. https://doi.org/10.1111/oik.02354

Shirk, A. J., Landguth, E. L., & Cushman, S. A. (2017). A comparison of individual-based genetic distance metrics for landscape genetics. *Molecular Ecology Resources*, *17*(6), 1308–1317. https://doi.org/10.1111/1755-0998.12684

Shirk, A. J., Landguth, E. L., & Cushman, S. A. (2018). A comparison of regression methods for model selection in individual-based landscape genetic analysis. *Molecular Ecology Resources*, *18*(1), 55–67. https://doi.org/10.1111/1755-0998.12709

Smith, G. R., & Dowling, T. E. (2008). Correlating hydrographic events and divergence times of speckled dace (Rhinichthys: Teleostei: Cyprinidae) in the Colorado River drainage. *Geological Society of America Special Papers*, *439*(303), 301–317. https://doi.org/10.1130/2008.2439(13).

Smouse, P. E., Long, J. C., & Sokal, R. R. (1986). Multiple regression and correlation extensions of the Mantel test of matrix correspondence. *Systematic Zoology*, *35*(4), 627. https://doi.org/10.2307/2413122

Spear, S. F., Balkenhol, N., Fortin, M. J., McRae, B. H., & Scribner, K. (2010). Use of resistance surfaces for landscape genetic studies: Considerations for parameterization and analysis. *Molecular Ecology*, *19*(17), 3576–3591. https://doi.org/10.1111/j.1365-294X.2010.04657.x

Tang, Q., Fung, T., & Rheindt, F. E. (2020). ResDisMapper: An r package for fine-scale mapping of resistance to dispersal. *Molecular Ecology Resources*, *20*(3), 819–831. https://doi.org/10.1111/1755-0998.13127

Terui, A., Ishiyama, N., & Urabe, H. (2019). Metapopulation stability in branching river network. *Proceedings of the National Academy of Sciences of the United States of America*, *115*(26), E5963–E5969. https://doi.org/10.1073/pnas.1821107116

Thiele, J., Buchholz, S., & Schirmel, J. (2018). Using resistance distance from circuit theory to model dispersal through habitat corridors. *Journal of Plant Ecology*, *11*(3), 385–393. https://doi.org/10.1093/jpe/rtx004

Thomaz, A. T., Christie, M. R., & Knowles, L. L. (2016). The architecture of river networks can drive the evolutionary dynamics of aquatic populations. *Evolution*, *70*(3), 731–739. https://doi.org/10.1111/evo.12883

Tonkin, J. D., Altermatt, F., Finn, D. S., Heino, J., Olden, J. D., Pauls, S. U., & Lytle, D. A. (2018). The role of dispersal in river network metacommunities: Patterns, processes, and pathways. *Freshwater Biology*, *63*(1), 141–163. https://doi.org/10.1111/fwb.13037

Troia, M. J., Kaz, A. L., Niemeyer, J. C., & Giam, X. (2019). Species traits and reduced habitat suitability limit efficacy of climate change refugia in streams. *Nature Ecology and Evolution*, *3*(9), 1321–1330. https://doi.org/10.1038/s41559-019-0970-7

Troia, M. J., & McManamay, R. A. (2020). Biogeographic classification of streams using fish community– and trait–environment relationships. *Diversity and Distributions*, *26*(1), 108–125. https://doi.org/10.1111/ddi.13001

Van Strien, M. J., Keller, D., Holderegger, R., Ghazoul, J., Kienast, F., & Bolliger, J. (2014). Landscape genetics as a tool for conservation planning: Predicting the effects of landscape change on gene flow. *Ecological Applications*, *24*(2), 327–339. https://doi.org/10.1890/13-0442.1

1070 Walker, A. E., Moore, J. N., Grams, P. E., Dean, D. J., & Schmidt, J. C. (2020). Channel narrowing by inset floodplain formation of the lower Green River in the Canyonlands region, Utah. *GSA Bulletin*, *132*(11–12), 2333–2352. https://doi.org/10.1130/b35233.1

1072 Weir, B., & Cockerham, C. C. (1984). Estimating F-Statistics for the analysis of population structure. *Evolution*, *38*(6), 1358–1370. https://doi.org/10.2307/2410030

1074 White, S. L., Hanks, E. M., & Wagner, T. (2020). A novel quantitative framework for riverscape genetics. *Ecological Applications*, *30*(7), 1–17. https://doi.org/10.1002/eap.2147

1076 Whitney, J. E., Whittier, J. B., & Paukert, C. P. (2017). Environmental niche models for riverine desert fishes and their similarity according to phylogeny and functionality. *Ecosphere*, *8*(1), 1078 e01658. https://doi.org/10.1002/ecs2.1658

Wiesenfeld, J. C., Goodman, D. H., & Kinziger, A. P. (2018). Riverscape genetics identifies 1080 speckled dace (Rhinichthys osculus) cryptic diversity in the Klamath–Trinity Basin. *Conservation Genetics*, *19*(1), 111–127. https://doi.org/10.1007/s10592-017-1027-6

1082 Winiarski, K. J., Peterman, W. E., & McGarigal, K. (2020). Evaluation of the R package 'resistancega': A promising approach towards the accurate optimization of landscape 1084 resistance surfaces. *Molecular Ecology Resources*, *20*(6), 1583–1596. https://doi.org/10.1111/1755-0998.13217

1086 Zbinden, Z. D., Douglas, M. R., Chafin, T. K., & Douglas, M. E. (2022a). Riverscape community genomics: A comparative analytical approach to identify common drivers of 1088 spatial structure. *Molecular Ecology*, *In press*. https://doi.org/10.1111/mec.16806

Zbinden, Z. D., Douglas, M. R., Chafin, T. K., & Douglas, M. E. (2022b). Trait-based predictors 1090 of genomic diversity within freshwater fish communities of the Ozarks. *BioRxiv*. https://doi.org/10.1101/2022.10.26.513843

1092 Zbinden, Z. D., Douglas, M. R., Chafin, T. K., & Douglas, M. E. (2023). A community genomics approach to natural hybridization. *Proceedings of the Royal Society B: Biological Sciences*, 1094 *290*(1999), 20230768. https://doi.org/10.1098/rspb.2023.0768

Figures and Tables

Table 1: RiverATLAS variables used as covariates for modeling effective resistance networks in ResistNet. Inclusion as potential covariates simulation-based validation and empirical case study in *Rhinichthys osculus* is indicated as "*". Weights (*w*) are provided for each of the simulated scenarios (Sim1-4).

Var.	Description	Empirical	Simulation	*w*Sim1	*w*Sim2	*w*Sim3	*w*Sim4
AET	Actual evapotranspiration; equivalent to 'soil water stress coefficient'	*	-	-	-	-	-
ARI	Aridity index	*	-	-	-	-	-
CRP	Cropland extent as percentage of reach catchment	*	-	-	-	-	-
DOR	Degree/ index of anthropogenic flow regulation	*	*	-	-	0.5	0.5
FOR	Forest landcover percentage in reach catchment	*	-	-	-	-	-
HFT	Standardized human footprint index	*	-	-	-	-	-
KAR	Percentage of catchment exposed karst terrain	*	-	-	-	-	-
LEN	Reach length in kilometers	*	-	1.0	-	-	-
LKA	Limnicity (percent lake area in reach catchment)	*	-	-	-	-	-
PAC	Percentage of reach catchment in protected areas	*	-	-	-	-	-
POP	Scaled human population size in 2010	*	-	-	-	-	-
PRE	Precipitation, as annual average in reach catchment	*	*	-	-	-	-
PST	Pasture extent as percentage of reach catchment	*	-	-	-	-	-
RDD	Road crossing density	*	*	-	-	-	0.5
RIV	River volume along reach segment	*	-	-	-	-	-
REV	Reservoir volume in watershed upstream of reach	*	-	-	-	-	-
RUN	Annual average land surface discharge and runoff	*	-	-	-	-	-
SGR	Stream gradient; ratio of elevation differential in reach and length	*	-	-	1.0	1.0	1.0
SLP	Average slope degree in catchment	*	-	-	-	-	-
SOC	Soil organic carbon content	*	-	-	-	-	-
TMP	Mean annual temperature	*	*	-	1.0	1.0	1.0
URB	Urban extent	-	*	-	-	-	-

1102   **Figure 1**: Overview of ResistNet and its validation environments. (a) Inputs consist of a spatial network with *n* covariates encoded as
reach-level attributes (as shapefile, *.shp*), and pairwise genetic distances for georeferences samples. These are used to construct a
1104   minimal graph representation as the primary data structure of ResistNet. A population of models randomly initialises the genetic
algorithm, with fitness computed using a Maximum-Likelihood Population Effects (MLPE) model applied to the composite,
1106   transformed 'effective' resistance values. Optimisation proceeds iteratively following random sampling (mutation) of parameter
values, swapping between models (recombination), and 'selection' to generate offspring. (b) Variable transformation occurs given
1108   eight functions, or a subset thereof (per user configuration). (c) Three input networks of varying topological complexity and spatial
extent served as templates from which to simulate distance matrices to test ResistNet (Net1-3).

1110

Figure 2: Summary of recovered relative variable importance (RVI) from simulated scenarios. RVIs were derived from four pre-determined 'known' scenarios (Sim1-4), tested within three distinct template networks (Net1-3). Results are reported as: (1) Ensemble (all): a model-averaged RVI computed from all models chosen across 20 independent replicates; (2) Ensemble (best): A model-averaged RVI calculated exclusively from the 'optimum' models in each replicate, determined by the Akaike Information Criterion (AIC); and (3) Median: The median RVI amongst all replicates. Each scenario was modeled using three different sample sizes: 10, 30, and 50, represented by blue, red, and green points, respectively.

.

Figure 3: Sampling localities and riverscape genetic structure of *N*=78 sampled populations of Speckled Dace (*Rhinichthys osculus*) in the Colorado River Basin, as divided into five subregions: (A) Virgin River; (B) Little Colorado River; (C) Grand Canyon; (D) Upper Colorado River Basin; (E) Lower Colorado River Basin (Gila and Bill Williams Rivers). Subregion networks are displayed with edges colored by fitted F_{ST} (linearised), and sampling locations as mean among-individual assignment probabilities to ancestry clusters derived by ADMIXTURE. Pairwise observed F_{ST} (Obs.) is plotted against pairwise fitted F_{ST} (Pre.), with significance of Mantel's *r* indicated at $p<0.05$ (*), $p<0.01$ (**), and $p<0.001$ (***).

Figure 4: Environmental resistance models and model-averaged resistance networks for Speckled Dace (*Rhinichthys osculus*) from five subregions of the Colorado River. Shown for each subregion (A-E) are model-averaged relative variable importance (RVI) as the sum of Akaike weights for an ensemble model of the best (per AIC) model sampled in each of 20 independent replicates. Here, an RVI threshold of 0.8 is shown in gray. Model-averaged resistance values are plotted on the spatial networks (green=low; red=high), and plotted pairwise against F_{ST} [i.e., isolation-by-resistance (IBR)], with significance of Mantel's *r* indicated at $p<0.05$ (*), $p<0.01$ (**), and $p<0.001$ (***).

 Supplementary Material

Table S1: Locality information (=Locality) and sample sizes (=*N*) for *N*=78 sampling localities
 of Speckled Dace (*Rhinichthys osculus*) in subregions of the Colorado River Basin.

Site	Locality	Subregion	*N*
182	Grand Canyon RM182	Grand Canyon	10
ANR	Animas River	Upper CRB	6
ARU	Aravaipa Creek; San Pedro tributary	Lower CRB	12
BAC	Grand Canyon RM87-117	Grand Canyon	10
BBC	Big Beaver Creek – White River	Upper CRB	9
BBL	Meadow Valley Wash	Virgin	9
BDM	Beaver Dam Wash	Virgin	6
BDS	Beaver Dam Wash	Virgin	10
BDW	Mainstem Virgin River	Virgin	10
BFT	Blacks Fork	Upper CRB	10
BIG	Big Sandy River	Upper CRB	10
BOU	Grand Canyon RM87-117	Grand Canyon	10
CAM	Camp Spring – White River	Virgin	18
CCN	Beaver Dam Wash	Virgin	4
CHE	Chevelon Creek; Little Colorado trib.	Little Colorado	12
CHU	Grand Canyon RM225-249	Grand Canyon	9
CLC	Meadow Valley Wash	Virgin	10
CYC	Chinle Wash; Colorado R. tributary	Upper CRB	10
DES	Desolation Canyon; Green R. mainstem	Upper CRB	7
DIA	Grand Canyon RM225-249	Grand Canyon	10
DOL	Dolores River	Upper CRB	12
EAG	Eagle Creek	Lower CRB	12
ECC	East Clear Creek; Little Colorado trib	Little Colorado	12
EFL	East Fork, Little Colorado River	Little Colorado	10
ELK	Elk River (Yampa River Tributary)	Upper CRB	10
ELV	Grand Canyon RM87-117	Grand Canyon	10
EVR	Meadow Valley Wash	Virgin	7
FER	Ferron Creek – San Rafael River	Upper CRB	8
FOS	Fossil Creek; Verde River tributary	Lower CRB	10
FRA	Francis Creek; Bill Williams R. tributary	Lower CRB	12
FRE	Fremont River; Green River tributary	Upper CRB	10
GLN	San Francisco River at Glenwood	Lower CRB	9
HAM	Ham's Fork; Green River	Upper CRB	10
HAV	Havasu Creek, Grand Canyon	Grand Canyon	10
HFK	Ham's Fork; Green River	Upper CRB	10
ISP	Indian Springs; White River; Virgin-Muddy tributary	Virgin	8
KAN	Kanab Creek; Grand Canyon	Grand Canyon	10
KWA	Grand Canyon RM225-249	Grand Canyon	10
LAV	Mainstem Virgin River	Virgin	5

LCR	Little Colorado River confluence	Grand Canyon	10
LIT	Mainstem Virgin River	Virgin	7
LYC	Little Yampa Canyon; Yampa River	Upper CRB	10
MAT	Grand Canyon RM143-189	Grand Canyon	10
MAY	Maynard Spring – Pahranagat Valley	Virgin	7
MBD	Mainstem Virgin River	Virgin	8
MCE	McElmo Creek – San Juan River	Upper CRB	10
MOA	Moapa River; Virgin-Muddy tributary	Virgin	22
MOD	Mainstem Virgin River	Virgin	8
MVW	Meadow Valley Wash	Virgin	10
MUC	Muddy Creek – Blacks Fork River	Upper CRB	10
NCC	San Simon	Lower CRB	10
NFV	Upper Virgin River	Virgin	10
NTH	Upper Virgin River	Virgin	10
NVC	Vermillion Creek	Upper CRB	10
PAH	Pahranagat River; Virgin-Muddy basin	Virgin	10
PAR	Paria River	Upper CRB	12
PIP	Grand Canyon RM87-117	Grand Canyon	10
RYC	Ryckman Creek upstream of Hill Creek Confluence	Upper CRB	10
SCG	Mainstem Virgin River	Virgin	9
SIC	Silver Creek; Little Colorado tributary	Little Colorado	10
SHN	Shinumo Creek; Colorado R. tributary	Grand Canyon	10
SMF	Smith's Fork; Green River	Upper CRB	7
SPG	Spring Creek; Verde River tributary	Lower CRB	10
SUN	Sunnyside Creek; White River; Virgin-Muddy	Virgin	10
SUR	Grand Canyon RM225-249	Grand Canyon	10
SYC	Sycamore Creek; Agua Fria tributary	Lower CRB	10
TSA	Chinle Wash; Colorado R. tributary	Upper CRB	10
TUL	Tularosa Creek – San Francisco River	Lower CRB	10
VEL	Burn's Ranch – Pahranagat Valley	Virgin	7
VSC	Mainstem Virgin River	Virgin	10
WEN	East Fork, Little Colorado R.	Little Colorado	10
WFA	Mainstem Virgin River	Virgin	8
WFG	West Fork of the Gila River at cliff dwellings	Lower CRB	10
WIL	East Clear Creek; Little Colorado trib	Little Colorado	10
WIT	Grand Canyon RM143-189	Grand Canyon	10
WPC	Island Park below Whirlpool Canyon; Yampa River	Upper CRB	10
WHY	Chinle Wash; Colorado R. tributary	Upper CRB	10
YAM	Yampa River (Green R. tributary)	Upper CRB	10

1144 **Table S2**: Accuracy of ResistNet validations runs performed using four simulated scenarios (s1-s4) on three template networks (net1-3), and three sample sizes (10, 30, 50). Shown for each are
1146 the proportion of replicates for which the model-averaged output accurately captured the exact simulated scenario ("Exact Match"), or encapsulated the simulated scenario (e.g., having an
1148 additional covariate implicated; "Correct Model Included"). The same metrics are also shown for the ensemble_all and ensemble_best approaches, but as true (1) or false (0).

1150

| Network | Samples | Simulation | Correct Model Included | | | Exact Match | | |
			Ensemble-all	Ensemble-best	Replicates	Ensemble-all	Ensemble-best	Replicates
net1	10	s1	1	1	1	0	0	0.7
		s2	1	1	0.95	1	1	0.85
		s3	1	1	0.9	1	1	0.7
		s4	1	1	0.8	1	1	0.35
	30	s1	1	1	1	1	1	1
		s2	1	1	1	1	1	1
		s3	1	1	1	1	1	0.85
		s4	1	1	0.95	1	1	0.85
	50	s1	1	1	1	1	1	0.9
		s2	1	1	1	1	1	1
		s3	1	1	0.95	1	1	0.8
		s4	1	1	0.9	1	1	0.9
net2	10	s1	1	1	1	1	1	0.8
		s2	1	1	0.3	1	1	0.25
		s3	0	0	0.6	0	0	0.45
		s4	1	1	0.65	0	0	0.3
	30	s1	1	1	1	1	1	1
		s2	0	0	0.2	0	0	0.2
		s3	0	0	0.35	0	0	0.3
		s4	1	1	0.75	1	1	0.6
	50	s1	1	1	1	1	1	1
		s2	0	0	0.2	0	0	0.2
		s3	0	0	0.3	0	0	0.3
		s4	1	1	0.8	1	1	0.6
net3	10	s1	1	1	1	1	1	0.7
		s2	1	1	0.8	0	0	0.45
		s3	1	1	0.7	1	1	0.3
		s4	1	1	0.45	1	1	0.3
	30	s1	1	1	1	1	1	0.8
		s2	1	1	0.9	1	1	0.65
		s3	1	1	0.8	0	0	0.4

	s4	1	1	0.55	1	1	0.45
50	s1	1	1	1	1	1	0.95
	s2	1	1	1	1	1	0.95
	s3	1	1	0.6	0	0	0.35
	s4	1	1	0.55	0	0	0.25

Table S3: Accuracy of ResistNet validations runs performed using four simulated scenarios (s1-s4) on three template networks (net1-3), and three sample sizes (10, 30, 50). Shows are results summarised across all simulated scenarios, with the proportion of replicates or ensemble models (ensemble_best and ensemble_all) for which the model-averaged output accurately captured the exact simulated scenario ("Exact Match"), or encapsulated the simulated scenario (e.g., having an additional covariate implicated; "Correct Model Included").

| Network | Samples | Correct Model Included | | | Exact Match | | |
		Ensemble-all	Ensemble-best	Replicates	Ensemble-all	Ensemble-best	Replicates
net1	10	1	1	0.9125	0.75	0.75	0.65
	30	1	1	0.9875	1	1	0.925
	50	1	1	0.9625	1	1	0.9
net2	10	0.75	0.75	0.6375	0.5	0.5	0.45
	30	0.5	0.5	0.575	0.5	0.5	0.525
	50	0.5	0.5	0.575	0.5	0.5	0.525
net3	10	1	1	0.7375	0.75	0.75	0.4375
	30	1	1	0.8125	0.75	0.75	0.575
	50	1	1	0.7875	0.5	0.5	0.625

1158 **Figure S1**: Validation of ResistNet using three networks and four simulated scenarios. The proportion of replicates (green) is shown, illustrating the model-averaged result that accurately recovered the known conditions (light green) in contrast to those containing the
1160 known condition along with additional spurious covariates (dark green). Results are also presented for across-replicate ensemble models (blue) and ensemble models constructed exclusively from the best AIC models selected for each replicate (referred to as
1162 "ensembleBest"; red). Note that the results are further divided by sample size (10, 30, or 50).

1164

Figure S2: Model optimization results for simulation-based validation of ResistNet conducted on three networks and four simulated scenarios. Results are presented for simulations with 10 (blue), 30 (red), and 50 (green) samples, each performed in 20 independent replicates. The fitness values (Y-axis) represent AIC, shown as absolute values transformed onto a consistent scale across replicates using min-max normalisation.

Figure S3: Summary of model-averaged weights (MAW) from simulated scenarios. MAWs were derived from four pre-determined 'known' scenarios (Sim1-4), tested within three distinct template networks (Net1-3). Results are reported as: (1) Ensemble (all): a MAW computed from all models chosen across 20 independent replicates; (2) Ensemble (best): MAW calculated exclusively from the 'optimum' models in each replicate, determined by the Akaike Information Criterion (AIC); and (3) Median: The mean MAW amongst all replicates. Each scenario was modeled using three different sample sizes: 10, 30, and 50, represented by blue, red, and green points, respectively.

Figure S4: AIC values across replicates for simulation-based validation of ResistNet, performed on three networks and four simulated scenarios. Results are partitioned by simulation scenario (rows) and sample size (columns; 10, 30, or 50 samples), with results shown by network (x-axis).

1184 **Figure S5:** Correlation of output pairwise resistance matrices among replicate ResistNet runs (*N*=20) with simulated input distance matrices across four simulated scenarios (s1-4) run within three template networks (tc1-3), at varying sample sizes (10, 30, 50),
1186 reported as Mantel's *r*.

1188

Figure S6: Population structure among Colorado River Basin (CRB) Speckled Dace populations. Upper Colorado/ Green River aggregates (*K*=6) represent: Paria River (PAR), Chinle Wash (CYC-WHY), San Juan+San Rafael+Fremont rivers (DOL-FER), Lower-Mid Green River (DES-BIG), Smiths Fork (SMF), and Vermillion Creek (NVC). The Lower CRB, represented by the Gila+Bill Williams River drainages (*K*=8) include: the Bill Williams River (FRA), Verde River (FOS-SPG), Agua Fria (SYC), San Pedro (ARU), San Simon (NCC), Eagle Creek (EAG), San Francisco River (GLN-TUL), and Upper Gila River (WFG). The Little Colorado River (*K*=4) encompasses East Clear Creek (WIL-ECC), Chevelon Creek (CHE), Silver Creek (SIC), and the East Fork (WEN-EFL). Grand Canyon (*K*=5): distributed linearly along the length of the canyon, with the exception of Havasu Creek (HAV). Virgin River (*K*=7): upper Virgin River (NTH-NFV), Mainstem Virgin River (LAV-WFA; BDW-VSC), Beaver Dam Wash (BDM-CCN), Meadow Valley Wash (BBL-MVW), Moapa River (MOA), Pahranagat (PAH-VEL), and White River (ISP-CAM).

1200   **Figure S7**: Isolation-by-distance (IBD) for Speckled Dace (*Rhinichthys osculus*) populations
       sampled from five subregions of the Colorado River Basin, plotted as the pairwise geographic/
1202   hydrologic distance (in km) and linearised F_{ST}. Correlation is reported as Mantel's *r*, with
       significance indicated at $p<0.05$ (*), $p<0.01$ (**), and $p<0.001$ (***).

1204

1206   **Figure S8**: Model-averaged relative variable importance (RVI) and weights (MAW) for environmental resistance models derived
using ResistNet applied to Speckled Dace (*Rhinichthys osculus*) from five subregions of the Colorado River. Shown for each
1208   subregion (A-E) are RVIs and MAWs reported from an ensemble of all sampled models (ensemble_all; blue), an ensemble of the best
model sampled within each replicate (ensemble_best; red), and as the mean (MAW) or median (RVI) of replicates (green).

1210

1212 **Figure S9**: Environmental resistance models and model-averaged resistance network for
Speckled Dace (*Rhinichthys osculus*) evaluated globally across *N*=78 sampling localities in the
1214 Colorado River. Shown are model-averaged relative variable importances (RVIs) as the sum of
Akaike weights for an ensemble model of the best (per AIC) model sampled in each of 20
1216 independent replicates. Here, an RVI threshold of 0.8 is shown in gray. Model-averaged
resistance values are plotted on the spatial networks (green=low; red=high), and plotted pairwise
1218 against F_{ST} [i.e., isolation-by-resistance (IBR)].

1220