Symbiodiniaceae conduct under natural bleaching stress during advanced gametogenesis

stages of the mesophotic coral *Alveopora allingi*

Gal Eyal[1,2,*], Lee Eyal-Shaham[3,4], Yossi Loya[3]

[1] ARC Centre of Excellence for Coral Reef Studies, School of Biological Sciences, The

University of Queensland, St Lucia 4072, QLD, Australia. Correspond to:

GAL4596@GMAIL.COM

[2] The Mina & Everard Goodman Faculty of Life Sciences, Bar-Ilan University, Ramat Gan

5290002, Israel

[3] School of Zoology, Tel-Aviv University, Ramat Aviv 6997801, Israel

[4] The Interuniversity Institute for Marine Sciences of Eilat, Eilat 88103, Israel

Keywords: Mesophotic Coral Ecosystems (MCEs), Coral Bleaching, Coral Reproduction,

Host-Symbiont Interaction, Coral-Algae Symbiosis, Red Sea

Abstract

The mesophotic coral *Alveopora allingi* from the northern Gulf of Eilat/Aqaba, Red

Sea, is affected by year-round partial coral-bleaching events. During these events, the

migration of Symbiodiniaceae takes place from the coral-host mesoglea to the

developed oocytes in bleached parts of colonies of *A. allingi* but not in the non-bleached

parts. Additionally, these oocytes are abnormal, missing part of the structural material

of the peripheral areas and are also significantly larger in the bleached areas of the

colonies. Hence, we suggest a parasitic behavior of the symbionts or a commensalism

relationship which enhance symbionts' needs during bleaching periods and may boost

the gametogenesis development in these corals. We propose that evolutionarily, this

behavior may greatly contribute to the symbiont community survival throughout the

bleaching period, and it can also be beneficial for the host's persistence and adaptation
to bleaching through the acquisition of a specific symbiont community following the
bleaching event.

Introduction

Coral reefs have experienced increasing stress over the last four decades due to local
anthropogenic perturbations and global climate change, resulting in severe damage to
coral populations and their reproduction (Loya 2004,2007; Carpenter et al. 2008; Baird
et al. 2009; Harrison 2011; Hughes et al. 2017). The symbiotic relationship between
corals and their algal endosymbionts (Symbiodiniaceae) is a key factor in the
evolutionary success of hermatypic corals (Wooldridge 2010). This close association
between primary producer and consumer enables the tight nutrient recycling that is
thought to explain the high productivity of coral reefs (Hoegh-Guldberg 1999;
Wooldridge 2013; Muller-Parker et al. 2015). Consequently, environmental and
physiological conditions that result in changes in the relationship between animal host
and Symbiodiniaceae may have profound ecological and physiological effects (Szmant
and Gassman 1990). The loss of Symbiodiniaceae - a phenomenon described as coral
bleaching - can cause acute damage to the colony (Glynn 1993; Brown 1997; Loya et
al. 2001; Suggett and Smith 2020). Coral bleaching has been observed in response to a
diverse range of stressors and is associated with both anthropogenic and natural
disturbances (Fitt and Warner 1995; Glynn 1996; Kushmaro et al. 1996; Hoegh-
Guldberg 1999; Rowan 2004; Hughes et al. 2017; Hughes et al. 2018). Severe and
prolonged bleaching can cause partial to total colony death, resulting in diminished reef
growth, transformation of reef-building communities to degraded alternate states,
increased bioerosion and, ultimately, the disappearance of reef structures (Glynn 1996).
The future state of shallow reefs is glooming and the framework building of coral

communities is expected to shift over to low-relief and less complex entities (Knowlton

2001; Pandolfi et al. 2003; Hughes et al. 2007).

While shallow reefs have recently been intensively affected by mass coral-bleaching

(Hughes et al. 2018), very few mesophotic coral ecosystems (MCEs; 30-150 m depth)

were reported to have undergone bleaching during these events (Baker et al. 2016;

Frade et al. 2018). One explanation could be that of the 'deep reef refuge' hypothesis

(DRRH), which suggests that deep ecosystems are more protected from the

disturbances that affect shallow reefs, and consequently could provide a viable

reproductive source for shallow-reef areas following disturbance (Bongaerts et al.

2010). MCEs too, however, are not immune to bleaching (Frade et al. 2018) or to other

disturbances (Rocha et al. 2018; Pinheiro et al. 2019), which in many cases have been

overlooked due to technical difficulties. One of the most important questions facing

scientists, policy-makers, and the general public is that of why there has been an

apparent increase in the incidence of coral bleaching in the last four decades (Goreau

and Hayes 1994; Hoegh-Guldberg 1999; Hughes et al. 2017) and what we can do about

it?

Nevertheless, although highly impacted by anthropogenic stressors, the shallow corals

of the Gulf of Eilat/Aqaba (GoE/A) (northern Red Sea) are considered heat-tolerant and

resilient to thermal stress (Fine et al. 2013; Krueger et al. 2017; Osman et al. 2018).

They have been described as comprising 'super-corals' that do not experience natural

bleaching (Grottoli et al. 2017). Nonetheless, *Stylophora pistillata,* the most abundant

depth-generalist coral in the coral reefs of Eilat, suffers from periodic non-fatal

bleaching in the summer months in the upper MCEs (Nir et al. 2014), and recent

observations have revealed multiple-species bleaching-events on these deeper

ecosystems (Eyal et al. 2019).

Symbiotic relationship between the coral host and its Symbiodiniaceae community are

complex and essential for the development of reef corals (Trench 1971; LaJeunesse

2020). Nevertheless, these relationships are not necessarily always beneficial for both

the host and the symbionts. Parasitism in cnidarian symbiosis under stress conditions

was suggested in the past but supported only by indirect evidences (Baker et al. 2018;

Peng et al. 2020). Here we use observations from coral reproductive study to present

another potential hypothesis of parasitism of the symbionts or/and commensalism

relationship between the host and its symbionts.

Sexual reproduction is the most critical part of any living taxa (Kondrashov 1988).

Although energetically costly, the benefit of increasing fitness and genetic diversity is

priceless. Evolution processes through sexual reproduction and gene-shuffling is the

primary adaptive capability to environmental changes and enabling extension to and

occupation of new ecological niches (Rundle et al. 2006). In shallow corals, diverse

strategies of sexuality were discovered during recent years (Baird et al. 2009; Harrison

2011; Eyal-Shaham et al. 2019; Eyal-Shaham et al. 2020) and references within) but

our knowledge of the reproduction of MCE corals remains limited (Shlesinger and Loya

2019).

The purpose of this research was to detect possible natural bleaching effects on the

development of gametogenesis in mesophotic corals. Hence, we examine the

reproductive ecology of the mesophotic coral, *A. allingi*, under bleaching stress and

describe a novel host-symbiont behavior interaction under bleaching conditions.

Methods

Study area, sampling procedures, and histological processes

Coral pairs from the same genets of non-bleached (pigmented) and visually bleached

A. allingi were sampled in front of the Interuniversity Institute for Marine Sciences in

Eilat, Israel (IUI) (29°30′N, 034°55′E), at a depth of 60 m, during the end of their 2011

reproductive season (October 2011). This is a well-developed reef with a wide variety

of coral species, and *A. allingi* is one of the most abundant species in the area (Eyal-

Shaham et al. 2016). As part of a larger project described in Eyal-Shaham et al. (2016),

once a month, during the full moon, nubbins (4–5 cm in length) were randomly sampled

from 4–6 healthy colonies of *A. allingi* using SCUBA technical diving and Closed-

Circuit Rebreathers (CCR). Only in October 2021 we found partially bleached colonies

(two different genet pairs of pigmented and visually bleached nubbins). The sampled

colonies were >20 m distant from one another. The collected nubbins were placed in a

sealed nylon bag filled with seawater and immediately after collection were fixed in a

4% formaldehyde solution in seawater for 24–48 h. The nubbins were then rinsed in tap

water for 15 minutes and transferred to 70% ethyl alcohol. The decalcification process

was carried out following the protocol of Eyal-Shaham et al. (2016). A piece of tissue

was taken from each decalcified nubbin and 6 μm thick latitudinal histological serial

sections were prepared and dyed with Mayer's hematoxylin and Putt's eosin (H&E

stain) to highlight the reproduction structures.

<u>Histological and statistical analyses</u>

The histological analysis was performed using a Nikon Eclipse 90i microscope and NIS

Elements D 3.2 software (Nikon Instruments Inc.). A detailed study of gonad structure,

size, and development was conducted by analyzing the histological cross-sections [see

details in Eyal-Shaham et al. (2016)]. Detailed identification of the oocytes (Oc),

spermaries (Sp), nucleus and nucleoli (N), and Symbiodiniaceae (Z) were conducted.

The examination included size measurements of the oocytes (longest axes), performed

only when the nucleoli were present, and identification of the developmental stages of

gametocytes within the polyps. Mean oocyte size of each developmental stage was

calculated from all oocytes presented in the samples from the bleached and the healthy

groups.

Differences in oocyte sizes within partly bleached colonies were calculated in R

language and environment (R Development Core Team 2014) using 1,000 bootstrap

samples of the values to indicate the difference between the median oocyte size of the

bleached part and that of the control part.

Results and Discussion

During coral bleaching events that occur at 50-60 m depth, the depth-specialist *A.*

allingi experiences severe visual bleaching in parts of the branches of otherwise healthy

colonies (Fig. 1a, b). Unlike the periodic summer bleaching of *S. pistillata*, also

described from the same reef (Nir et al. 2014), and in which the colonies recover after

the summer ends, *A. allingi* presents a gradual degradation of the bleached tissue, with

mortality of the entire bleached part of the colonies occurring within a few months.

Our analysis of the histological sections of partly-bleached colonies revealed a

phenomenon that to our knowledge has never previously been described:

Symbiodiniaceae migration to the oocyte cytoplasm [i.e., in the visually bleached part

of the colonies, Symbiodiniaceae were observed in the oocytes' cytoplasm, which

presented an irregular shape and imperfect cytoplasm with missing parts of the

structural material of the peripheral areas, while the healthy (non-bleached) portions of

the same colonies presented normally-developed oocytes (Fig. 1c-f)]. Additionally, the

bleached coral's oocytes were found to be significantly larger than those of the non-

bleached parts (Fig. 2; Mann-Whitney Test, T=544, P<0.001), which suggests a faster

reproduction development in the impacted parts of these colonies. Although

Symbiodiniaceae can be directly transmitted from the parent colony (i.e. vertical

transmission) in some species (Lesser et al. 2013; Muller-Parker et al. 2015), the

observed change in shape (i.e., larger sizes and imperfect cytoplasm with missing parts

of the structural material of the peripheral areas of the oocytes only), suggests that this

is not the case in *A. allingi*. Moreover, Symbiodiniaceae were not observed in any

mature oocytes of other, healthy, colonies.

We hypothesize that such migration of Symbiodiniaceae into the coral oocyte

cytoplasm under severe bleaching conditions potentially enhances the survivorship of

the symbiont community, suggesting the existence of a possible mechanism of

exploitation of cytoplasmic materials by certain symbionts. An alternative hypothesis

is that the partially bleached hosts are in a process of absorbing oocyte lipids as an

emergency energetic source. Cell membranes have been disturbed, including the

symbiosome surrounding the host cells lining the exterior of the oocyte. This enables

symbionts to escape the host cell and invade the oocytes.

The Symbiodiniaceae community in corals is generally dominated by one or two main

symbiont types (van Oppen et al. 2005; Ziegler et al. 2017) but can also contain a large

number of rarer members of the family, which play important roles in the resilience of

the holobiont (Ziegler et al. 2018). Network theoretic modeling predicts that elevated

symbiont diversity can increase community stability in response to environmental

changes (Fabina et al. 2013). Moreover, the functional role of Symbiodiniaceae in the

holobiont is not defined by a single species but rather by the assemblages of symbionts

(Ziegler et al. 2018). The rare symbionts, however, associate in several manners with

the coral host (Knowlton and Rohwer 2003; Kirk et al. 2013; Baker et al. 2018), which

may include non-mutualistic species. Additionally, some Symbiodiniaceae may

transform their mutualistic relationship into a parasitic state during a stressed period, thus parasitizing their hosts for nutrition (Baker et al. 2018). The symbiont benefits are clearly not equal to those of the host, host and this partnership could be lost under times of stress to either the host or the symbionts. In addition to this novel behavior- the phenomenon of symbionts taking advantage of the stressed hosts, might also hold a key to some urgent questions regarding the coral-algae symbiosis breakdown and contribute to our understanding of the mechanisms involved in the bleaching phenomenon. The discovery that Symbiodiniaceae migrate into the oocytes of bleached parts of partially-bleached MCE corals, but not into the healthy (non-bleached) parts, reveals the existence of opportunistic demeanor of the symbionts on their coral host's gametes, which may result in lowered functionality of the affected host parts. From an evolutionary point of view, this behavior is highly beneficial to the symbionts' survival throughout the bleaching period, since expelled symbionts are not expected to survive for long in the environment (Hill and Ralph 2007); but could also be beneficial for the host's persistence and adaptation to bleaching. Selection processes towards bleaching-resistant capabilities at the gamete level of heathy oocytes will result, in the long run, in adapted populations of this coral.

Acknowledgements

We thank the Interuniversity Institute for Marine Sciences in Eilat for the logistical support, and Barbara Colorni for help with the histological work. The comments of two anonymous reviewers greatly improved the manuscript. This study was supported by the Israel Science Foundation (ISF) No. 1191/16 to YL and by the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie post-doctoral grant agreement No. 796025 to GE.

Compliance with ethical standards

All samples were collected and treated according to the Israeli Nature and Parks Authority permit no. 2011/38249.

Conflict of interest

On behalf of all authors, the corresponding author states that there is no conflict of interest.

References

Baird AH, Guest JR, Willis BL (2009) Systematic and biogeographical patterns in the reproductive biology of scleractinian corals. Annual Review of Ecology, Evolution, and Systematics 40:551-571

Baker DM, Freeman CJ, Wong JC, Fogel ML, Knowlton N (2018) Climate change promotes parasitism in a coral symbiosis. The ISME Journal 12:921-930

Baker EK, Puglise KA, Harris PT (2016) Mesophotic Coral Ecosystems: A Lifeboat for Coral Reefs? United Nations Environment Programme and GRID-Arendal

Bongaerts P, Ridgway T ,Sampayo E, Hoegh-Guldberg O (2010) Assessing the 'deep reef refugia'hypothesis: focus on Caribbean reefs. Coral reefs 29:309-327

Brown BE (1997) Coral bleaching: causes and consequences. Coral reefs 16:S129-S138

Carpenter KE, Abrar M, Aeby G, Aronson RB ,Banks S, Bruckner A, Chiriboga A, Cortés J, Delbeek JC, DeVantier L (2008) One-third of reef-building corals face elevated extinction risk from climate change and local impacts. Science 321:560-563

Eyal G, Tamir R, Kramer N, Eyal-Shaham L, Loya Y (2019) The Red Sea: Israel Mesophotic coral ecosystems. Springer, pp199-214

Eyal-Shaham L, Eyal G, Tamir R, Loya Y (2016) Reproduction, abundance and survivorship of two Alveopora spp. in the mesophotic reefs of Eilat, Red Sea. Scientific reports 6:20964

Eyal-Shaham L, Eyal G, Sakai K, Nozawa Y, Harii S, Sinniger F, Bronstein O, Ben-Zvi O, Shlesinger T, Loya Y (2019) Repetitive sex change in the stony coral Herpolitha limax across a wide geographic range. Scientific reports 9:1-12

Eyal-Shaham L, Eyal G, Ben-Zvi O, Sakai K, Harii S, Sinniger F, Hirose M, Cabaitan P, Bronstein O, Feldman B (2020) A unique reproductive strategy in the mushroom coral Fungia fungites. Coral Reefs:1-12

Fabina NS, Putnam HM, Franklin EC, Stat M, Gates RD (2013) Symbiotic specificity ,association patterns, and function determine community responses to global changes: defining critical research areas for coral-Symbiodinium symbioses. Global Change Biology 19:3306-3316

Fine M, Gildor H, Genin A (2013) A coral reef refuge in the Red Sea .Global Change Biology 19:3640-3647

Fitt W, Warner M (1995) Bleaching patterns of four species of Caribbean reef corals. The Biological Bulletin 189:298-307

Frade PR, Bongaerts P, Englebert N, Rogers A, Gonzalez-Rivero M, Hoegh-Guldberg O (2018) Deep reefs of the Great Barrier Reef offer limited thermal refuge during mass coral bleaching. Nature Communications 9:3447

Glynn P (1993) Coral reef bleaching: ecological perspectives. Coral reefs 12:1-17

Glynn PW (1996) Coral reef bleaching: facts, hypotheses and implications. Global Change Biology 2:495-509

Goreau TJ, Hayes RL (1994) Coral bleaching and ocean" hot spots". Ambio-Journal of Human
Environment Research and Management 23:176-180
Grottoli AG, Tchernov D, Winters G (2017) Physiological and Biogeochemical Responses of Super-Corals
to Thermal Stress from the Northern Gulf of Aqaba, Red Sea. Frontiers in Marine Science 4:215
Harrison PL (2011) Sexual reproduction of scleractinian corals Coral reefs: an ecosystem in transition.
Springer, pp59-85
Hill R, Ralph PJ (2007) Post-bleaching viability of expelled zooxanthellae from the scleractinian coral
Pocillopora damicornis. Marine Ecology Progress Series 352:137-144
Hoegh-Guldberg O (1999) Climate change, coral bleaching and the future of the world's coral reefs.
Marine and freshwater research 50:839-866
Hughes TP, Rodrigues MJ, Bellwood DR, Ceccarelli D, Hoegh-Guldberg O, McCook L, Moltschaniwskyj N,
Pratchett MS, Steneck RS, Willis B (2007) Phase shifts, herbivory, and the resilience of coral
reefs to climate change. Current biology 17:360-365
Hughes TP, Kerry JT, Álvarez-Noriega M, Álvarez-Romero JG, Anderson KD, Baird AH, Babcock RC, Beger
M, Bellwood DR, Berkelmans R (2017) Global warming and recurrent mass bleaching of corals.
Nature 543:373-377
Hughes TP, Anderson KD, Connolly SR, Heron SF, Kerry JT, Lough JM, Baird AH, Baum JK, Berumen ML,
Bridge TC (2018) Spatial and temporal patterns of mass bleaching of corals in the
Anthropocene. Science 359:80-83
Kirk N, Thornhill D, Kemp D, Fitt W, Santos S (201 (3Ubiquitous associations and a peak fall prevalence
between apicomplexan symbionts and reef corals in Florida and the Bahamas. Coral Reefs
32:847-858
Knowlton N (2001) The future of coral reefs. Proceedings of the National Academy of Sciences 98:5419-
5425
Knowlton N, Rohwer F (2003) Multispecies microbial mutualisms on coral reefs: the host as a habitat.
the american naturalist 162:S51-S62
Kondrashov AS (1988) Deleterious mutations and the evolution of sexual reproduction. Nature 336:435-
440
Krueger T ,Horwitz N, Bodin J, Giovani M-E, Escrig S, Meibom A, Fine M (2017) Common reef-building
coral in the Northern Red Sea resistant to elevated temperature and acidification. Royal
Society open science 4:170038
Kushmaro A, Loya Y, Fine M, Rosenberg E (1996) Bacterial infection and coral bleaching. Nature 380:396
LaJeunesse TC (2020) Zooxanthellae. Current Biology 30:R1110-R1113
Lesser M, Stat M, Gates R (2013) The endosymbiotic dinoflagellates (Symbiodinium sp.) of corals are
parasites and mutualists. Coral Reefs 32:603-611
Loya Y (2004) The coral reefs of Eilat—past, present and future: three decades of coral community
structure studies Coral Health and disease. Springer, pp1-34
Loya Y (2007) How to influence environmental decision makers? The case of Eilat) Red Sea) coral reefs.
Journal of Experimental Marine Biology and Ecology 344:35-53
Loya Y, Sakai K, Yamazato K, Nakano Y, Sambali H, van Woesik R (2001) Coral bleaching: the winners
and the losers. Ecology Letters 4:122-131
Muller-Parker G, D'elia CF, Cook CB (2015) Interactions between corals and their symbiotic algae Coral
Reefs in the anthropocene. Springer, pp99-116
Nir O, Gruber DF, Shemesh E, Glasser E, Tchernov D (2014) Seasonal mesophotic coral bleaching of
Stylophora pistillata in the Northern Red Sea. PloS one 9:e84968
Osman EO, Smith DJ, Ziegler M, Kürten B, Conrad C, El-Haddad KM, Voolstra CR, Suggett DJ (2018)
Thermal refugia against coral bleaching throughout the northern Red Sea. Global Change
Biology 24:e474-e484
Pandolfi JM, Bradbury RH ,Sala E, Hughes TP, Bjorndal KA, Cooke RG, McArdle D, McClenachan L,
Newman MJ, Paredes G (2003) Global trajectories of the long-term decline of coral reef
ecosystems. Science 301:955-958
Peng S-E, Moret A, Chang C, Mayfield AB, Ren Y-T, Chen W-NU, Giordano M, Chen C-S (2020) A shift
away from mutualism under food-deprived conditions in an anemone-dinoflagellate
association. PeerJ 8:e9745
Pinheiro HT, Eyal G, Shepherd B, Rocha LA (2019) Ecological insights from environmental disturbances
in mesophotic coral ecosystems. Ecosphere 10

R Development Core Team (2014) R: A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing, Vienna, Austria

Rocha LA, Pinheiro HT, Shepherd B, Papastamatiou YP, Luiz OJ, Pyle RL, Bongaerts P (2018) Mesophotic
coral ecosystems are threatened and ecologically distinct from shallow water reefs. Science
361:281-284

Rowan R (2004) Coral bleaching: thermal adaptation in reef coral symbionts. Nature 430:742-742

Rundle HD, Chenoweth SF, Blows MW (200 6The roles of natural and sexual selection during
adaptation to a novel environment. Evolution 60:2218-2225

Shlesinger T, Loya Y (2019) Sexual reproduction of scleractinian corals in mesophotic coral ecosystems
vs. shallow reefs Mesophotic Coral Ecosystems. Springer, pp653-666

Suggett DJ, Smith DJ (2020) Coral bleaching patterns are the outcome of complex biological and
environmental networking. Global Change Biology 26:68-79

Szmant A, Gassman N (1990) The effects of prolonged "bleaching" on the tissue biomass and
reproduction of the reef coral Montastrea annularis. Coral reefs 8:217-224

Trench R (1971) The physiology and biochemistry of zooxanthellae symbiotic with marine coelenterates
III. The effect of homogenates of host tissues on the excretion of photosynthetic products in
vitro by zooxanthellae from two marine coelenterates. Proceedings of the Royal Society of
London Series B Biological Sciences 177:251-264

van Oppen MJ, Mahiny AJ, Done TJ (2005) Geographic distribution of zooxanthella types in three coral
species on the Great Barrier Reef sampled after the 2002 bleaching event. Coral Reefs 24:482-
487

Wooldridge SA (2010) Is the coral-algae symbiosis really 'mutually beneficial'for the partners? BioEssays
32:615-625

Wooldridge SA (2013) Breakdown of the coral-algae symbiosis: towards formalising a linkage between
warm-water bleaching thresholds and the growth rate of the intracellular zooxanthellae.
Biogeosciences 10:1647-1658

Ziegler M, Eguíluz VM, Duarte CM, Voolstra CR (2018) Rare symbionts may contribute to the resilience
of coral–algal assemblages. The ISME Journal 12:161-172

Ziegler M, Arif C, Burt JA, Dobretsov S, Roder C, LaJeunesse TC, Voolstra CR (2017) Biogeography and
molecular diversity of coral symbionts in the genus Symbiodinium around the Arabian
Peninsula. Journal of Biogeography 44:674-686

Figure 1: Partially-bleached *Alveopora allingi*, with micrographs of gonadal development. (a)

general colony view at 60 m depth; (b) partial-bleaching within a single genet; (c) male and

female gonads from the non-bleached (pigmented) part of the colony; (d) male and female

gonads from the visually bleached part of the colony; (e) magnification of the marked area in

section c, showing stage 4 healthy mature oocytes; and (f) magnification of the marked area in

section (d), showing several stage 4 severely damaged oocytes with many Symbiodiniaceae in

the oocytes' cytoplasm. Scale bars as indicated above the scale-line. Oc: Oocyte, Sp:

Spermaries, N: nucleus, Z: Symbiodiniaceae.

Figure 2: Oocyte sizes of the mesophotic coral *Alveopora allingi* during a partial colony natural

bleaching event. (a) Significantly larger stage 4 oocytes in the bleached parts of partly bleached

colonies (t-test, T=544, P<0.001). Centerlines of the boxes represent the medians; box limits

indicate the 25th and 75th percentiles; whiskers extend 1.5 times the interquartile range from

the 25th and 75th percentiles, notches represent the 95% confidence interval for each median,

all data represented by dots. Two different genet pairs of bleached (n=84 oocytes) and non-

bleached (n=21 oocytes) parts were analyzed. (b) The difference in μm between median values

of the bleached parts compared to non-bleached parts. The difference in condition is calculated

from 1,000 bootstrap samples of the values, indicating the difference between the median

oocyte size of the bleached parts and that of the healthy parts. The horizontal black line indicates

the 95% confidence interval.