Copper intoxication in group B *Streptococcus* triggers transcriptional activation

of the *cop* operon that contributes to enhanced virulence during acute infection

Matthew J. Sullivan[a,b], Kelvin G. K. Goh[a,b], Dean Gosling[a,†], Lahiru Katupitiya[a], and

Glen C. Ulett[a,b #]

[a]School of Pharmacy and Medical Sciences, and [b]Menzies Health Institute Queensland,

Griffith University, Gold Coast, Queensland, Australia 4222.

[†]Current address: Pathology Queensland, Health Support Queensland, Department of

Health, Queensland Government, Southport, Queensland, Australia 4215.

Article Type: Research article

Running Title: Copper stress in group B *Streptococcus*

Word count: 6038, 198 (abstract)

Display Items: 7 Figures, 3 Tables (plus 5 Supplementary Figures)

Conflict Statement: All the authors do not have a commercial or other association that

might pose a competing financial interest in relation to the work described.

Correspondence and offprint requests: Professor Glen C. Ulett, School of Pharmacy

and Medical Sciences, and Menzies Health Institute Queensland, Griffith Health Centre,

Griffith University, Gold Coast, Queensland 4222, Email: g.ulett@griffith.edu.au, T: +61

7 56780765, F: +61 7 55632236

Abstract

Bacteria can utilize Copper (Cu) as a trace element to support cellular processes;

however, excess Cu can intoxicate bacteria. Here, we characterize the *cop* operon in

group B streptococcus (GBS), and establish its role in evasion of Cu intoxication and

the response to Cu stress on virulence. Growth of GBS mutants deficient in either the

copA Cu exporter, or the *copY* repressor, were severely compromised in Cu-stress

conditions. GBS survival of Cu stress reflected a mechanism of CopY de-repression of

the CopA efflux system. However, neither mutant was attenuated for intracellular

survival in macrophages. Analysis of global transcriptional responses to Cu by RNA-

sequencing revealed a stress signature encompassing homeostasis of multiple metals.

Genes induced by Cu stress included putative metal transporters for manganese import,

whereas a system for iron export was repressed. In addition, *copA* promoted the ability

of GBS to colonize the blood, liver and spleen of mice following disseminated infection.

Together, these findings show that GBS *copA* mediates resistance to Cu intoxication,

via regulation by the Cu-sensing transcriptional repressor, *copY*. Cu stress responses in

GBS reflect a transcriptional signature that heightens virulence and represents an

important part of the bacteria's ability to survive in different environments.

Keywords: Metallobiology; group B *Streptococcus*; Metal ions; Copper efflux; Bacterial

pathogenesis

Importance

Understanding how bacteria manage cellular levels of metal ions, such as copper, helps

to explain how microbial cells can survive in different stressful environments. We show

how the opportunistic pathogen group B Streptococcus (GBS) achieves homeostasis of

intracellular copper through the activities of the genes that comprise the cop operon,

and describe how this helps GBS survive in stressful environments, including in the

mammalian host during systemic disseminated infection.

Introduction

Copper (Cu) is the most reactive of the biologically-relevant first-row *d*-block transition metals. It can readily displace other metals, including Zinc (Zn), Nickel (Ni), Cobalt (Co), Iron (Fe) and Manganese (Mn) from metalloproteins in which these are bound (1, 2). In cells across all Kingdoms of life, Cu-dependent enzymes are pivotal to many essential processes that underpin physiologic biochemical reactions, owing to its ability to redox cycle between Cu(I) and Cu(II) states (3). When present in excess, however, Cu is hazardous to cellular processes and macromolecules due to effects of localized free-radical damage (4). A key role for Cu in bacterial cell biology encompasses the host-pathogen interface where a balance between Cu usage and avoidance of Cu intoxication resulting from host antimicrobial responses must be attained for bacteria to survive, as reviewed elsewhere (5). At this interface, human immune cells can mobilize Cu in a defence response to infection that culminates in exposure of intracellular bacteria to antimicrobial levels of Cu that kill the invading pathogen (6-9). Thus, bacterial responses to Cu stress and mechanisms of resistance to metal-ion intoxication have emerged as important facets of bacterial disease pathogenesis (10).

There are several mechanisms that enable bacteria to tolerate excess extracellular Cu, and these have been characterized in several pathogenic species. These mechanisms, which represent survival strategies against Cu intoxication (10) typically involve Cu-transporting P-type ATPases, exemplified by CopA, which detoxify Cu by exporting it from the bacterial cytosol (11-13). This mechanism has been described in gram-

negative and gram-positive bacteria, and among *Streptococcaceae* (14-16), including

pneumococci and *Streptococcus pyogenes* (17).

An opportunistic streptococcal pathogen of humans and animals for which resistance to

excess extracellular Cu has not been described is *Streptococcus agalactiae*, also known

as group B streptococcus (GBS). This organism, unlike other *Streptococcaceae*, is

associated with a comparatively broad host-range that encompasses humans, cattle

and fish (18). In humans, GBS is a major cause of invasive infection in infants <3

months of age (19), and causes a more diverse range of disease aetiologies compared

to other streptococci, including meningitis, pneumonia, skin and soft-tissue infections,

sepsis, arthritis, osteomyelitis, urinary tract infection, and endocarditis (19). GBS has

several virulence factors that enable survival in cytotoxic environments, such as acid

stress, oxidative stress, and during host colonization, as reviewed elsewhere (20).

Cellular tolerance to Cu stress and mechanisms of responding to and surviving Cu

intoxication have not been defined in GBS.

In this study, we characterise a system that enables Cu efflux in GBS, encompassing a

copA-homologue, which we show mediates control of Cu efflux in the bacteria. We

show this system affects survival, growth and virulence of GBS in the mammalian host.

Results

Transcriptomic profiling of GBS responses to extracellular Cu

To dissect the global response of GBS to extracellular Cu stress we performed RNA

sequencing (RNASeq) to define the complete primary transcriptional response to Cu.

RNASeq analysis of mid-log phase cells of GBS strain 874391 grown in the presence of

0.5 mM Cu (a sub-inhibitory level; see below) for 2.5h (compared to control cultures that

were grown without supplemental Cu) revealed a surprisingly modest Cu-responsive

transcriptome. The response comprised 18 transcriptional responses, defined as

significant based on $-2 \geq$ FC ≥ 2 (FDR < 0.05, *n*=4 biological replicates), of 11 up-

regulated transcripts and 7 down-regulated transcripts (Figure 1, Table 1). The most

significantly up-regulated were homologues of the *copY-copA-copZ* gene cluster (3.6-

4.0-fold), which encode the putative CopY (Cu-binding transcriptional repressor), CopA

(P-type ATPase Cu-efflux system) and CopZ (Cu-chaperone) proteins. To validate the

expression levels of selected targets identified in RNASeq, we targeted *pcl1*, *hvgA* and

copY in qRT-PCR assays, confirming near-identical fold-change values to RNASeq

analyses (Figure 1B). The products of *copY*, *copA* and *copZ* in GBS (Figure 2A) are

moderately conserved compared to other Lactobacillales, including *S. mutans, S.*

thermophilus, S. pyogenes, S. pneumoniae, and *Enterococcus* (Figure 2B, ranked by

CopA similarity) for which a homologous Cu management system is defined (9, 13-15).

Together, these findings provide transcriptional and comparative insights that support a

proposed model of Cu efflux in GBS (Figure 2C).

In addition to *copYAZ*, GBS up-regulated putative metal transporters for Mn import

(*mtsABC*, *mntH2*), and down-regulated an iron (Fe) export system *(fetAB;* Figure 1,

Table 1). In addition, we detected significant down-regulation of *pcl1*, that encodes a

putative membrane protein, and the gene for hypervirulence-associated factor, *hvgA*.

Roles of copA and copY in Cu-resistance in GBS

To functionally characterize two major elements of the Cu-responsive transcriptome in

GBS, we targeted *copA* and *copY* by generating isogenic deletion mutants and

comparing growth of these with the WT. In a nutrient-rich medium (Todd Hewitt Broth;

THB) supplemented with increasing amounts of Cu, we observed no growth inhibitory

effects of high Cu (up to 1.5 mM) for WT GBS; the lag phases, growth rates, and final

biomass yields were equivalent for the WT comparing different levels of Cu (Figure 3).

In contrast, growth of $\Delta copA$ GBS was severely attenuated by Cu stress $\geq$ 1 mM;

complementation of the *copA* mutation restored the phenotype to WT (Figure 3).

Deletion of *copY*, encoding a putative Cu-dependent repressor of *copA*, had no effect

on the growth rate or lag phase in THB supplemented with Cu but significantly affected

final biomass yield of cultures (*D* at 600nm at 18h; Supplementary Figure S1). Together,

this shows that *copA* confers cellular resistance to Cu stress in GBS in a manner that

supports bacterial growth in nutritive conditions.

Temporal- and Concentration-Dependent Bactericidal effects of Cu towards GBS

We examined the effects of Cu towards survival of GBS in nutrient limited conditions by

using a minimal Chemically-Defined Medium (CDM), in consideration of prior studies of

antibacterial activity in minimal media (21, 22), the influence of culture media (23, 24),

and possible cell-protective effects from buffering agents, *e.g.* glutathione (15). We

observed that the level of Cu required to inhibit GBS growth in CDM was markedly

lower than THB (Supplementary Figure S2); for example, the growth of $\Delta copA$ GBS was

significantly attenuated in CDM in the presence of 0.5 mM supplemental Cu, a level that

caused only slight inhibition of growth in THB (Figure 3). To more precisely define the

bactericidal effects of Cu towards GBS in CDM, we performed time-kill curves using 1-5

x 10^6 CFU/mL exposed to Cu concentrations ranging between 50 μM and 1 mM. We

observed potent killing effects that depended on both Cu concentration and time with $\geq$

0.5 mM Cu significantly killing WT GBS after 3h of exposure (Figure 4, WT); after 24h,

there was a ~230-fold reduction in CFU/mL at 0.5 mM Cu (0 mM Cu = 5.6 Log_{10}

CFU/mL, 0.5 mM CFU/mL = 3.2 Log_{10} CFU/mL), and no viable GBS remained in

cultures exposure to 1 mM Cu. The $\Delta copA$ strain was significantly more susceptible to

Cu-induced toxicity compared to the WT; Cu levels above $\geq$ 0.5 mM significantly

reduced the number of viable $\Delta copA$ GBS beginning as early as 1h exposure (Figure 4,

$\Delta copA$); the degree of bacterial killing was significant at the 24h timepoint for $\geq$ 0.5 mM

Cu (Figure 4). Interestingly, at the 6h timepoint, Cu at $\geq$ 0.5 mM did not affect viability of

$\Delta copY$ GBS, compared to the WT strain (Figure 4, $\Delta copY$), a hyper-resistance

phenotype that was not apparent at the 24 h timepoint. Together, these findings are

consistent with a role for CopA in resisting Cu stress, and CopY as Cu-dependent

repressor of *copA*.

Regulation of Cu efflux by CopY, and accumulation of metals during Cu stress

We next sought to examine the control of Cu-responses in GBS at the transcriptional

level. The capacity of Cu stress to induce expression of *copA* for Cu export was

examined by analysing *copA* expression by qRTPCR in GBS exposed to Cu

concentrations ranging from 0.25-1.5 mM Cu in THB. GBS significantly upregulated

copA in response to Cu (3.7-fold – 14.2-fold) in a manner that was titratable with the Cu

concentration (Figure 5A). To ascertain the role of CopY as a putative Cu-responsive

repressor of *copA* expression in GBS, we quantified *copA* mRNA transcripts in the

$\Delta copY$ mutant exposed to 0.5 mM Cu. This level of Cu was carefully chosen as sub-

inhibitory to enable comparisons independent of metabolic state and therefore limiting

any potential bias from possible discordant Cu stress between WT and mutants with

varied resistance phenotypes. Deletion of *copY* resulted in severe de-regulation of *copA*

expression, causing a ~207±45-fold increase of *copA* transcripts in the *copY*

background (Figure 5B). Thus, these data establish that GBS *copY* represses *copA* in

the absence of Cu, and are consistent with a model of CopY-mediated regulation of

copA that is responsive to intracellular accumulation of Cu.

To examine the impact of *copA* and *copY* mutations on accumulation of Cu within the

cell we used the equivalent conditions per RNASeq and qPCR assays, exposing GBS

to 0.5 mM Cu in THB and measuring total Cu content of cells compared to non-exposed

controls (Ctrl). Inductively coupled optical emission spectroscopy (ICP-OES)

demonstrated that standard THB contained 0.2±0.08 µM Cu, reflecting trace amounts in

the medium. In the absence of supplemental Cu, WT GBS limited intracellular Cu

content such that only 0.8±0.1 µg Cu. g dry weight^{-1} were detected in cultures grown in

THB. However, exposure of WT GBS to 0.5 mM Cu resulted in a significant increase in

intracellular Cu to 31.9±3.1 µg Cu. g dry weight^{-1} (Figure 5C). Strikingly, Δ*copY* GBS

exhibited significantly less cellular Cu upon exposure to Cu (6.7±0.4 µg Cu. g dry

weight^{-1}), consistent with the observation that transcription of the Cu-exporter CopA is

significantly elevated in this mutant. In addition, we noted significant accumulation of

cellular Cu in the Δ*copA* strain (109.6±1.9 µg Cu. g dry weight^{-1}), confirming a Cu-efflux

role for *copA*. Interestingly, we noted modest but significant increases in Mn and Fe in

the Δ*copA* strain, and Zn in the Δ*copY* strain due to Cu stress (Figure 5D-F).

To extend our findings comparing mutant strains to WT at a level of Cu that is sub-

inhibitory to the Δ*copA* strain, we next undertook experiments to investigate the impact

of severe Cu stress on WT GBS; by repeating metal accumulation analyses using cells

grown in 1.5 mM Cu. We detected modest but statistically significant accumulation of Fe

inside GBS cells exposed to 1.5 mM Cu, co-occurring with significantly higher Cu levels;

however, no difference in Mn or Zn content (Supplementary Figure S3). Fe status has

been reported to influence bacterial resistance to peroxide stress (25) and therefore we

also undertook experiments to investigate the consequence of Cu toxicity on resistance

to oxidative stress using assays of susceptibility to hydrogen peroxide (H_2O_2). These

experiments demonstrated a significant reduction in the viability of Δ*copA* GBS exposed

to H_2O_2, which was dependent on pre-exposure to Cu (Supplementary Figure S4).

Thus, GBS cells undergoing Cu stress are rendered significantly more susceptible to

oxidative stress.

Role of CopA in macrophage killing of GBS

To examine whether the CopA Cu efflux system in GBS supports survival of the bacteria

in phagocytes we performed antibiotic protection assays with human monocyte-derived

macrophage-like cells. Macrophages were infected with WT, Δ*copA* GBS or its

complemented strain for 1h, and antibiotics were added to kill extracellular bacteria.

Viable intracellular GBS were quantified at 1h post-antibiotic addition, and at 24h and

48h to assess intracellular survival. We performed assays comparing culture conditions

of standard RPMI medium to medium that was supplemented with 20 μM Cu to ensure

adequate availability of Cu for host cellular responses. These assays demonstrated

significant reductions in the numbers of viable GBS over the time course (24 to 48h);

however, we did not detect any significant differences between the numbers of WT and

Δ*copA* GBS recovered from the host cells at any time point (Figure 6). The numbers of

bacteria recovered comparing the WT and Δ*copA* mutant were equivalent regardless of

the presence of supplemental Cu in the medium. Thus, under these conditions, *copA*

does not contribute to the intracellular survival of GBS in human macrophage-like cells.

GBS CopA contributes to virulence **in vivo**

To examine the contribution of Cu efflux to GBS virulence, we used a murine model of

disseminated infection (26). In mice challenged with 10^7 GBS, we detected significantly

fewer Δ*copA* mutant in the liver (median of 3.5 vs 4.2 $\log_{10}$ CFU.g tissue^{-1}; P= 0.005),

spleen (median of 3.8 vs 4.1 $\log_{10}$ CFU.g tissue^{-1} P= 0.013) and blood (median of 1.4 vs

1.9 $\log_{10}$ CFU.mL^{-1}; P= 0.044) compared to the WT at 24h post-inoculation (Figure 7).

However, no significant differences were observed between the counts of the WT and

Δ*copA* mutant in several other tissues, including the brain, heart, lungs, and kidneys

(Supplementary Figure S5). These data support a modest but significant role for cellular

management of Cu via *copA* in GBS in supporting disseminated infection *in vivo*.

Materials and Methods

Bacterial strains, plasmids and growth conditions

GBS, *E. coli* and plasmids used are listed in Table 2. GBS was routinely grown in Todd-

Hewitt Broth (THB) or on TH agar (1.5% w/v). *E. coli* was grown in Lysogeny Broth (LB)

or on LB agar. Routine retrospective colony counts were performed by plating dilutions

of bacteria on tryptone soya agar containing 5% defibrinated horse blood (Thermo

Fisher Scientific). Media were supplemented with antibiotics (spectinomycin (Sp)

100μg/mL; chloramphenicol (Cm) 10 μg/mL), as indicated. Growth assays used 200μL

culture volumes in 96-well plates (Greiner) sealed using Breathe-Easy® membranes

(Sigma-Aldrich) and measured attenuance (D, at 600nm) using a ClarioSTAR

multimode plate reader (BMG Labtech) in Well Scan mode using a 3mm 5x5 scan

matrix with 5 flashes per scan point and path length correction of 5.88mm, with agitation

at 300rpm and recordings taken every 30min. Media for growth assays were THB and a

modified Chemically-Defined Medium (CDM) (27) (with 1g/L glucose, 0.11g/L pyruvate

and 50μg/L L-cysteine), supplemented with Cu (supplied as $CuSO_4$) as indicated. For

attenuance baseline correction, control wells without bacteria were included for Cu in

media alone.

DNA extraction and genetic modification of GBS

Plasmid DNA was isolated using miniprep kits (QIAGEN), with modifications for GBS as

described elsewhere (28). Deletions in *copA* (CHF17_00507 / CHF17_RS02570) and

copY (CHF17_00506 / CHF17_RS02565) were constructed by markerless allelic

exchange using pHY304aad9 as described previously (29). Plasmids and primers are

listed in Table 2 and Table 3, respectively. Mutants were validated by PCR using

primers external to the mutation site and DNA sequencing.

RNA extraction, qRTPCR

For Cu exposure experiments, 1mL of overnight THB cultures were back-diluted 1/100

in 100mL of THB (prewarmed at 37°C in 250mL Erlenmeyer flasks) supplemented with

0.25, 0.5, 1.0 or 1.5mM Cu. Cultures were grown shaking (200rpm) at 37°C; after

exactly 2.5h, 10-50mL volumes containing approximately 500 million mid-log bacteria

were harvested; RNA was preserved and isolated as described previously (30). RNA

quality was analysed by RNA LabChip using GX Touch (Perkin Elmer). RNA (1000ng)

was reverse-transcribed using Superscript IV according to manufacturer's instructions

(Life Technologies) and cDNA was diluted 1:50 in water prior to qPCR. Primers (Table

3) were designed using Primer3 Plus (31, 32) to quantify transcripts using Universal

SYBR Green Supermix (Bio-Rad) using a Quantstudio 6 Flex (Applied Biosystems)

system in accordance with MIQE guidelines (33). Standard curves were generated

using five-point serial dilutions of genomic DNA (5-fold) from WT GBS 874391 (34).

Expression ratios were calculated using C_T values and primer efficiencies as described

elsewhere (35) using *dnaN*, encoding DNA polymerase III β-subunit as housekeeper.

Whole bacterial cell metal content determination

Metal content in cells was determined as described (36) with minor modifications.

Cultures were prepared essentially as described for *RNA extraction, qRTPCR* with the

following modifications; THB medium was supplemented with 0.5 mM $CuSO_4$ or not

supplemented (Ctrl), and following exposure for 2.5h, bacteria were harvested by

centrifugation at 4122 x g at 4°C. Cell pellets were washed 3 times in PBS + 5mM

EDTA to remove extracellular metals, followed by 3 washes in PBS. Pelleted cells were

dried overnight at 80°C and resuspended in 1mL of 32.5% nitric acid and incubated at

95°C for 1h. The metal ion containing supernatant was collected by centrifugation

(14,000 x g, 30min) and diluted to a final concentration of 3.25% nitric acid for metal

content determination using inductively coupled plasma optical emission spectroscopy

(ICP-OES). ICP-OES was carried out on an Agilent 720 ICP-OES with axial torch,

OneNeb concentric nebulizer and Agilent single pass glass cyclone spray chamber. The

power was 1.4kW with 0.75L/min nebulizer gas, 15L/min plasma gas and 1.5L/min

auxiliary gas flow. Cu was analysed at 324.75nm, Zn at 213.85nm, Fe at 259.94nm and

Mn at 257.61nm with detection limits at <1.1ppm. The final quantity of each metal was

normalised using dry weight biomass of the cell pellet prior to nitric acid digestion,

expressed as $\mu g.g^{-1}$ dry weight. Baseline concentrations were determined to be 0.2 ±

0.08 μM Cu in THB medium, and 40 ± 4 nM Cu in CDM medium from at least three

independent assays.

Hydrogen Peroxide assay

Peroxide survival assays were based on prior studies (37, 38) with minor modifications

to encompass using GBS cells that had been pre-grown in conditions of Cu stress.

Overnight cultures of WT and mutants were grown in THB, cultures were back-diluted

1/100 into either fresh THB or THB supplemented with 0.5mM Cu and grown for exactly

2.5 h at 37°C with 200 rpm agitation. Bacteria were harvested by centrifugation at 4122

x g and washed twice in PBS and resuspended in assay buffer (0.1M sodium phosphate

buffer pH 7.5). This was prepared by combining 41 mL of 0.2M dibasic sodium

phosphate and 9 mL of 0.2M monobasic sodium phosphate in a volume of 200 mL.

GBS cells were diluted in assay buffer alone, or assay buffer containing 5 mM H_2O_2

(Sigma-Aldrich H1009), to a final concentration of ~$5x10^6$ CFU/mL. Cell suspensions +/-

H_2O_2 were then incubated for 1h at 37°C and survival was monitored by serial dilution

and plating for CFU/mL counts.

RNA sequencing and bioinformatics

Cultures were prepared as described above for *RNA extraction, qRTPCR* to compare

mid-log phase cells grown in THB + 0.5 mM Cu to THB without added Cu. RNase-free

DNase-treated RNA that passed Bioanalyzer 2100 (Agilent) analysis was used for RNA

sequencing (RNA-seq) using the Illumina NextSeq 500 platform. We used a Bacterial

Ribosomal RNA (rRNA) Depletion kit (Invitrogen) prior to library construction, and

TruSeq library generation kits (Illumina, San Diego, California). Library construction

consisted of random fragmentation of the RNA, and cDNA production using random

primers. The ends of the cDNA were repaired and A-tailed, and adaptors were ligated

for indexing (with up to 12 different barcodes per lane) during the sequencing runs. The

cDNA libraries were quantitated using qPCR in a Roche LightCycler 480 with the Kapa

Biosystems kit (Kapa Biosystems, Woburn, Massachusetts) prior to cluster generation.

Clusters were generated to yield approximately 725K–825K clusters/mm^2. Cluster

density and quality was determined during the run after the first base addition

parameters were assessed. We ran single-end 75–bp sequencing runs to align the

cDNA sequences to the reference genome. For data preprocessing and bioinformatics,

STAR (version 2.7.3a) was used (parameters used: --outReadsUnmapped Fastx --

outSAMtype BAM SortedByCoordinate --outSAMattributes All) to align the raw RNA

sequencing fastq reads to the WT *S. agalactiae* 874391 reference genome (34).

HTSeq-count, version 0.11.1 (parameters used: -r pos -t exon -i gene_id -a 10 -s no -f

bam), was used to estimate transcript abundances (39). DESeq2 was then used to

normalized and test for differential expression and regulation following their vignette.

Genes that met certain criteria (i.e. fold change of > ±2.0, q value (false discovery rate,

FDR of <0.05) were accepted as significantly altered (40). Raw and processed data

were deposited in Gene Expression Omnibus (accession no. GSE167895 for *S.*

agalactiae 874391 Cu condition; GSE167894 (*S. agalactiae* 874391 control condition).

Mammalian cell culture

U937 monocytes were differentiated into human monocyte-derived macrophages

(MDMs) as follows. Briefly, monocytes were seeded (5×10^5 per well such that 1×10^5

adhere) into the wells of a 96-well tissue culture-treated plate (Falcon) essentially as

described elsewhere (41, 42), except that U937 monocytes were differentiated by

exposure to 30ng/mL phorbol 12-myristate 13-acetate (PMA) for 48h and cells

subsequently rested in media without PMA for 72h to enhance morphological and

phenotypic markers of MDMs (43). A multiplicity of infection (MOI) of 100 bacteria:

macrophage for 1h was used in RPMI without antibiotics. Non-adherent bacteria were

removed by five washes of 200µL PBS using a Well Wash Versa (Thermo Scientific).

RPMI containing 250U/mL penicillin, streptomycin (Gibco) and 50µg/mL gentamicin

(Sigma-Aldrich) were used for antibiotic protection assays to kill extracellular bacteria as

described previously by incubating for 1h at 37°C in 5% CO_2 (42). Samples were

processed after 1h (time zero), 24h or 48h after infection, monolayers were washed five

times with 200µL PBS and lysed by brief exposure to 50µL of 0.25% trypsin EDTA

(Gibco) and 0.1% Triton-X-100 (10 min) prior to dilution with 150µL PBS and estimation

of CFU/mL by serial dilution and plate counts on agar. Additional assays that ran in

parallel were identical except that 20 µM Cu was added to RPMI culture media at all

stages post PMA-treatment of U937 cells.

Animals and Ethics statement

Virulence was tested using a mouse model of disseminated infection based on

intravenous challenge with 10^7 GBS (WT or ∆copA) as described elsewhere (26). This

study was carried out in accordance with the guidelines of the Australian National

Health and Medical Research Council. The Griffith University Animal Ethics Committee

reviewed and approved all experimental protocols for animal usage according to the

guidelines of the National Health and Medical Research Council (approval:

MSC/01/18/AEC).

Statistical methods

All statistical analyses used GraphPad Prism V8 and are defined in respective Figure

Legends. Statistical significance was accepted at P values of ≤0.05.

Discussion

Transcriptional and cellular responses of bacterial pathogens to metal ions including Cu can influence host-pathogen interactions and thereby play a role in disease pathogenesis (4). The role of Cu homeostasis and detoxification in the biology of GBS have not hitherto been characterized, and no reports of a Cu stress response in this important human and animal pathogen are published. The principal finding of this study is the establishment of a transcriptional and cellular connection between the response to Cu stress in GBS and survival of the bacteria in conditions of Cu toxicity; this connection is mediated through *copA* and controlled through *copY*, and enables the bacteria to resist killing via Cu-mediated intoxication. Additionally, this study establishes that the connection between Cu stress responses in GBS and bacterial survival promotes virulence in the host during systemic, disseminated infection. The new insights into gene function in GBS viewed through the lens of the Cu stress transcriptome, combined with the findings of enhanced virulence elucidate molecular mechanisms that underpin GBS survival of intoxicating conditions, including those likely to be encountered in the host.

The transcriptional remodelling that occurs in GBS in response to Cu stress, as defined in this study on a global level, comprises an intriguingly constrained subset of genes. These findings indicate a tightly controlled system of transcriptional responses to Cu in GBS. Interestingly, equivalent low numbers of target genes were identified in previous transcriptional analyses of other streptococci exposed to Cu stress (14-16). Our findings are consistent with these prior reports, and support the notion that these transcriptional responses function in housekeeping or homeostasis to set a low limit of Cu availability

in the cytoplasm (15). In GBS, *copA* is among the most strongly activated genes in the

transcriptional response to Cu stress, and the mutational analysis performed in this

study shows that *copA* is crucial for the bacteria to attain an essential Cu efflux response

during Cu stress. However, *copA* is only one of an assembly of genes engaged by GBS

during Cu stress and it is likely that other genes in the transcriptome contribute to Cu

detoxification via additional means. For example, we detected up-regulation of putative

metal transporters for Mn import (*mtsABC*, *mntH2*) along with concurrent down-

regulation of a system that encodes Fe export machinery *(fetAB)*. These transcriptional

insights are interesting because they hint at additional stress response mechanisms that

occur during Cu stress in GBS, which extend beyond CopA and which need elucidation.

Metal content analysis by ICP-OES showed that high Cu stress disrupts additional pools

of intracellular trace metal, including for example causing modest elevations in levels of

Fe along with major increases in Cu content. The consequences of these alterations in

metal content in GBS for cell activity need further examination.

Transcription of metal ion-import and export genes, including those above-mentioned, is

typically controlled by metal-dependent regulatory proteins termed metalloregulators that

sense metal ion-bioavailability and work to maintain cellular metal homeostasis (44). In

our study, we detected no major changes in the expression of genes predicted to encode

metalloregulators, such as for transport of Mn (*mntR/mtsR*), Fe (*fur*), Zn (*adcR/sczA*) and

for the sensing of peroxide (*perR*) in GBS undergoing Cu stress (25, 44). It would be of

interest to elucidate whether such regulators undergo mis-metallation in GBS during Cu

stress, which might occur due to excess Cu likely outcompeting Mn and Fe for binding

sites in proteins; it is conceivable that Cu may displace metals from proteins as

described in other studies (45); however, precisely if and how metalloregulators might

respond to mis-metallation in GBS, if this occurs, will need to be elucidated in future.

In *S. pyogenes*, Mn uptake is facilitated by *mtsABC* and is protective against peroxide

induced stress (25). Expression of *mtsABC* is up-regulated by Mn deficiency, and is

controlled by the MtsR regulator (25, 46). Mn import and Fe efflux are co-ordinated in

order to control the metalation of superoxide dismutase (25), which can use Mn or Fe at

its catalytic site in streptococci; noting that Mn uptake can be disrupted by Zn (36). The

consequences of Cu intoxication in relation to oxidative stress was explored using killing

assays with hydrogen peroxide. These revealed reduced viability in *copA*-deficient GBS,

but only in cells that had been pre-exposed to Cu. Taken with our observations that Cu

accumulates significantly in this strain following Cu intoxication (and to a lesser extent

Mn and Fe), these data hint at a role for oxidative damage as a killing mechanism. Fe

can produce hydroxyl radicals and Cu(II) can be oxidised to Cu(I) in reactions with

peroxide, both products of which are likely to damage bacterial cells. A system for the

dual import of Mn and Fe in GBS is encoded by *mntH* that is regulated by pH (47). In

some strains of GBS, including the hypervirulent ST17 lineage used in this study, two

homologues of MntH exist, encoded by *mntH* and *mntH2*. This study demonstrates up-

regulation of *mntH2*, but not *mntH*, in response to Cu, however the role of *mntH2* or its

regulation are undefined in GBS. In *Bacillus subtilis* MntR co-ordinates Mn import (and

efflux), including control of genes homologous to GBS *mntH* and *mntH2* that import Mn

(48). This may provide context for elevated *mntH2* expression in the presence of

modest Cu stress in our study, since Cu-bound MntR has a much lower affinity for

binding its target operator sequence (14580210). We also note that *mtsABC* and *mntH2*

are also up-regulated in concert with down-regulation of Fe-transporting *fetAB* in

response to Zn stress in GBS (49). Together, our data indicate that modulation of

mtsABC, *mntH2* and *fetAB* expression forms parts of a transcriptional signature of GBS

exposed to Cu or Zn stress.

We observed no significant transcriptional response of *nrdD* expression in WT GBS

exposed to Cu stress. In contrast, we detected a highly statistically significant, but

modestly fold-change (up-regulated) response of the *nrdFIA* genes, encoding aerobic

dNTP synthesis components (1.7-fold, FDR < 0.001; data not shown). These findings

might hint at a role for Mn in the GBS Cu stress response given NrdD encodes an Fe-

containing enzyme, whereas NrdF encodes a Mn-dependent pathway. The role of these

genes in the GBS response to Cu stress requires further study. Modulation of putative

Mn (*mtsABC*) and Fe (*fetAB*) transporters, as well as other proteins predicted to localise

to the membrane (*pcl1*) or cell surface (*hvgA*) in GBS in response to Cu stress takes

place in the absence of altered cellular Mn or Fe content in the WT strain. This leads us

to suggest that examining the roles of targets identified in this study in the context of Cu

transport is needed in concert with other metals (*e.g.,* Mn). Indeed, the 'dearth of

information' on precisely how Cu is transported into bacterial cells from the external

environment represents a central mystery for Cu trafficking in bacteria (50).

Our assays of bacterial growth *in vitro* in conditions of Cu stress demonstrate that GBS

deficient in *copA* cannot grow as efficiently compared to the WT in conditions of elevated

extracellular Cu. The differences noted in GBS growth between nutritionally rich (THB)

and limited (CDM) media possibly reflect relative quantities of compounds that confer a

protective advantage for survival during Cu stress, such as glutathione (15) or other

thiol-containing amino acids that may interact with free Cu ions in solution. Glutathione is

not included in CDM as a separate chemical constituent but the quantities of

methionine, cystine and cysteine are 30, 62.6 and 50 mg.L^{-1}. It is also possible that the

levels of Cu utilized in our assays are distinct to those in host niches. Importantly,

however, Cu exposure assays *in vitro* are almost certainly influenced by the compounds

present in the medium that likely affect levels of Cu that become inhibitory (15). For

example, buffering effects would likely occur in THB, which would affect the uptake of

Cu by the bacteria. Our rationale for use of THB to analyse the impact of Cu toxicity on

GBS is that this medium is a standard rich growth medium for GBS and is widely used

for studies of this organism. The challenges inherent in such a rich media, including for

example, buffering effects, led us to also examine the impact of Cu toxicity on GBS in

defined CDM, which revealed, expectedly, a heightened susceptibility of the bacteria to

the metal stress. Other similar studies have reported degrees of intoxication effects due

to metal stress in different growth media, and our findings are consistent with these

observations. Some of the Cu concentrations used in our study may be considered

supra-physiological (1-1.5 mM) but the magnitude of these concentrations is likely much

greater than the magnitude of 'free' Cu ions (not bound by other molecules) that would

be present and available to react biologically. The concentration ranges used in the

current study were informed by available literature along with measures of expression of

copA, since we and others have shown that transcriptional activity of *copA* is influenced

by cellular Cu. Thus, *copA* activity can be considered a *bona fide* reporter of free cellular

Cu; or a surrogate marker for exposure to Cu stress. Consistent with this notion, we

observed higher expression of *copA* (7.6 $\pm$ 2.3 fold up-regulated; not shown) in CDM

with 100 μM than in THB with 500 μM Cu (4.9 $\pm$ 0.5 fold up; Figure 5). The Cu

concentrations that are encountered by GBS at infection sites or in different intracellular

compartments in host cells are unknown. Cu is elevated in the host at sites infected with

S. pyogenes (15), including the blood and in infected skin lesions. For mycobacterial

phagosomes, investigators have reported Cu concentrations of 25μM, and in *M.*

tuberculosis-infected macrophages, intravacuolar concentrations of 0.4mM (51).

Notwithstanding considerations of possible physiological ranges of Cu, the Cu exposure

assays used in our study are beneficial to establish *bona fide* gene function and

bacterial responses to a defined Cu stress condition.

Our analysis of Cu content in GBS cells exposed to Cu stress in defined conditions

shows that GBS maintains low steady-state levels of cellular Cu, in the absence of

excess extracellular Cu. In supplying extracellular Cu in excess, we show the level of

cellular Cu content is increased. Our finding that *copY* functions to repress *copA* in the

absence of Cu is consistent with previous reports in other bacteria (52, 53). Disrupting

the genetic systems for Cu efflux in GBS, via mutation of the CopA exporter, or the

CopY regulator, reveals divergent phenotypes that stem from loss of export or

regulatory function, resulting in accumulation ($\Delta copA$) or reduction ($\Delta copY$) of cellular

Cu. These phenotypes will be of interest to dissect in terms of the role of CopY in GBS

biology in other models of infection and disease in the future. The results of the current

study indicate that GBS might have a higher intrinsic level of resistance to Cu compared

to other *Streptococcus* spp., including *S. pyogenes*. However, we did not directly

compare different species and differences in experimental approaches or background

solution, as our study demonstrates, might affect these assays. Thus, further work

characterising relative resistance between streptococci more directly would be of

interest to the field. The potential for Cu resistance mechanisms to confer heightened

Cu tolerance in the context of the host environment is unexplored for GBS.

Studies have demonstrated increased Cu levels in the blood of humans during bacterial

infection (54, 55), however most of the insight into Cu-driven antimicrobial responses is

derived from studies of mammalian cells infected *in vitro*. In macrophages, bioavailability

of Cu correlates with antibacterial activities (56), and Cu 'hot spot' formation mediates

antimicrobial responses against intracellular bacteria (7). The cellular consequences of

Cu stress to GBS*,* which likely encounters such stress in the host, remains undefined.

Our findings based on *in vitro* infection macrophages showed no attenuation of GBS

devoid of CopA in survival in host cells, which was surprising given the important role of

Cu management in survival of other bacteria inside macrophages (6), and epithelial

cells (12). Notably, however, intracellular survival of *Salmonella* deficient in *cueO*, which

encodes an enzyme required for resistance to Cu ions, was not impaired in murine

macrophages in a previous study (57), leading the authors to suggest multiple host

factors are involved in clearance of the bacteria. In addition, *copA*-deficient *S. pyogenes*

were not impaired for survival in assays with human neutrophils [15]. Our study is

consistent with these findings. Other researchers have described the limitations of *in*

vitro tissue culture monolayer assays for determining intracellular survival of bacteria in

the context of Cu homeostasis (58).

Despite negative findings in macrophage monolayer assays *in vitro*, systemic infection of

mice exposed a connection between the ability of GBS to generate a Cu management

response via CopA and bacterial virulence *in vivo*. Here, *copA* was essential for GBS to

fully colonize and survive in the blood, as well as in other tissues. In demonstrating a

significant attenuation of GBS deficient in CopA to be fully virulent in mice, we suggest

that Cu toxicity may represent a form of stress experienced by the bacteria *in vivo* during

systemic infection. In other bacteria, including *Pseudomonas aeruginosa* and *Listeria*

monocytogenes, compromised Cu transport leads to attenuation for colonization in

various infection models (58, 59). Attenuation of GBS for colonization of the blood, liver

and spleen indicates that Cu management in the bacterial cell is essential not only for

efficient survival of the bacteria in the bloodstream but also for colonization of highly

immunologically active tissues; *i.e.,* Kupffer cells and splenic lymphocytes for innate and

adaptive immune responses, respectively. It is plausible that Cu stress that might be

encountered by GBS *in vivo* could influence virulence factor function, such as the

hypervirulence associated adhesin, HvgA, which we have shown is down-regulated

under Cu stress. It would be of interest to analyse the effect of Cu transport deficiency

in GBS in other relevant models of infection, including in vaginal colonization (60).

Additional to defining the effects of metal homeostasis in GBS on the nature of infection

and disease caused by this organism, small-molecules probes might hold promise for

the identification of other molecular mechanisms of metal homeostasis in GBS, as

reported for Gram-positive bacteria (61).

In summary, this study shows that management of Cu export in GBS is essential for the

bacteria to survive in environments of Cu stress. Cu intoxication in GBS generates a

transcriptional signature that includes activation of the *cop* operon to confer bacterial

survival and virulence in stressful environments. The exact role for Cu ions as an

antibacterial response against GBS warrants further investigation.

Acknowledgements

We thank Michael Crowley and David Crossman of the Heflin Centre for Genomic

Science Core Laboratories, University of Alabama at Birmingham (Birmingham, AL) for

RNA sequencing. We also thank Ryan Stewart at the School of Environment Analytical

Chemistry Core Facility, Griffith University, for ICP-OES. This work was supported by a

Project Grant from the National Health and Medical Research Council (NHMRC)

Australia (APP1146820 to GCU).

References

1.   Waldron KJ, Rutherford JC, Ford D, Robinson NJ. 2009. Metalloproteins and

metal sensing. Nature 460:823-30.

2.   Irving H, Williams RJP. 1953. 637. The stability of transition-metal complexes.

Journal of the Chemical Society (Resumed) doi:10.1039/jr9530003192:3192-

3210.

3.   Festa RA, Thiele DJ. 2011. Copper: an essential metal in biology. Curr Biol

21:R877-83.

4.   Ladomersky E, Petris MJ. 2015. Copper tolerance and virulence in bacteria.

Metallomics 7:957-64.

5.   Djoko KY, Ong CL, Walker MJ, McEwan AG. 2015. The Role of Copper and Zinc

Toxicity in Innate Immune Defense against Bacterial Pathogens. J Biol Chem

290:18954-61.

6.   White C, Lee J, Kambe T, Fritsche K, Petris MJ. 2009. A role for the ATP7A

copper-transporting ATPase in macrophage bactericidal activity. J Biol Chem

284:33949-56.

7.   Achard Maud ES, Stafford Sian L, Bokil Nilesh J, Chartres J, Bernhardt Paul V,

Schembri Mark A, Sweet Matthew J, McEwan Alastair G. 2012. Copper

redistribution in murine macrophages in response to *Salmonella* infection.

Biochem J 444:51-57.

8.   Ladomersky E, Khan A, Shanbhag V, Cavet JS, Chan J, Weisman GA, Petris

MJ. 2017. Host and Pathogen Copper-Transporting P-Type ATPases Function

Antagonistically during *Salmonella* Infection. Infect Immun 85.

9.   Johnson MD, Kehl-Fie TE, Klein R, Kelly J, Burnham C, Mann B, Rosch JW.

2015. Role of copper efflux in pneumococcal pathogenesis and resistance to

macrophage-mediated immune clearance. Infect Immun 83:1684-94.

10.   Giachino A, Waldron KJ. 2020. Copper tolerance in bacteria requires the

activation of multiple accessory pathways. Mol Microbiol 114:377-390.

11.   Rensing C, Fan B, Sharma R, Mitra B, Rosen BP. 2000. CopA: An *Escherichia*

coli Cu(I)-translocating P-type ATPase. Proc Natl Acad Sci U S A 97:652-6.

12.   Djoko KY, Franiek JA, Edwards JL, Falsetta ML, Kidd SP, Potter AJ, Chen NH,

Apicella MA, Jennings MP, McEwan AG. 2012. Phenotypic characterization of a

copA mutant of *Neisseria gonorrhoeae* identifies a link between copper and

nitrosative stress. Infect Immun 80:1065-71.

13.   Solioz M, Stoyanov JV. 2003. Copper homeostasis in *Enterococcus hirae.* FEMS

Microbiol Rev 27:183-95.

14.   Johnson MD, Kehl-Fie TE, Rosch JW. 2015. Copper intoxication inhibits aerobic

nucleotide synthesis in *Streptococcus pneumoniae.* Metallomics 7:786-94.

15.   Stewart LJ, Ong CY, Zhang MM, Brouwer S, McIntyre L, Davies MR, Walker MJ,

McEwan AG, Waldron KJ, Djoko KY. 2020. Role of Glutathione in Buffering

Excess Intracellular Copper in *Streptococcus pyogenes.* mBio 11.

16.   Shafeeq S, Yesilkaya H, Kloosterman TG, Narayanan G, Wandel M, Andrew

PW, Kuipers OP, Morrissey JA. 2011. The *cop* operon is required for copper

homeostasis and contributes to virulence in *Streptococcus pneumoniae.* Mol

Microbiol 81:1255-70.

17. Facklam R. 2002. What happened to the streptococci: overview of taxonomic and

nomenclature changes. Clin Microbiol Rev 15:613-30.

18. Chen SL. 2019. Genomic Insights Into the Distribution and Evolution of Group B

Streptococcus. Frontiers in Microbiology 10:1447.

19. Edwards MS, Baker CJ. 2018. 119 - *Streptococcus agalactiae* (Group B

Streptococcus), p 723-729.e1. *In* Long SS, Prober CG, Fischer M (ed), Principles

and Practice of Pediatric Infectious Diseases (Fifth Edition)

doi:https://doi.org/10.1016/B978-0-323-40181-4.00119-5. Elsevier.

20. Lindahl G, Stalhammar-Carlemalm M, Areschoug T. 2005. Surface proteins of

Streptococcus agalactiae and related proteins in other bacterial pathogens. Clin

Microbiol Rev 18:102-27.

21. Tuomanen E, Cozens R, Tosch W, Zak O, Tomasz A. 1986. The rate of killing of

Escherichia coli by beta-lactam antibiotics is strictly proportional to the rate of

bacterial growth. J Gen Microbiol 132:1297-304.

22. Eng RH, Padberg FT, Smith SM, Tan EN, Cherubin CE. 1991. Bactericidal

effects of antibiotics on slowly growing and nongrowing bacteria. Antimicrob

Agents Chemother 35:1824-8.

23. Sanders S, Bartee D, Harrison MJ, Phillips PD, Koppisch AT, Freel Meyers CL.

2018. Growth medium-dependent antimicrobial activity of early stage MEP

pathway inhibitors. PLoS ONE 13:e0197638.

24. Peterson LR, Gerding DN, Hall WH, Schierl EA. 1978. Medium-dependent

variation in bactericidal activity of antibiotics against susceptible *Staphylococcus*

aureus. Antimicrob Agents Chemother 13:665-8.

25.   Turner AG, Djoko KY, Ong CY, Barnett TC, Walker MJ, McEwan AG. 2019.

632   Group A Streptococcus co-ordinates manganese import and iron efflux in

633   response to hydrogen peroxide stress. Biochem J 476:595-611.

634   26.   Sullivan MJ, Ulett GC. 2020. Evaluation of hematogenous spread and ascending

635   infection in the pathogenesis of acute pyelonephritis due to group B

636   streptococcus in mice. Microb Pathog 138:103796.

637   27.   Moulin P, Patron K, Cano C, Zorgani MA, Camiade E, Borezee-Durant E,

638   Rosenau A, Mereghetti L, Hiron A. 2016. The Adc/Lmb System Mediates Zinc

639   Acquisition in *Streptococcus agalactiae* and Contributes to Bacterial Growth and

640   Survival. J Bacteriol 198:3265-3277.

641   28.   Sullivan MJ, Ulett GC. 2018. Stable Expression of Modified Green Fluorescent

642   Protein in Group B Streptococci To Enable Visualization in Experimental

643   Systems. Appl Environ Microbiol 84.

644   29.   Ipe DS, Ben Zakour NL, Sullivan MJ, Beatson SA, Ulett KB, Benjamin WHJ,

645   Davies MR, Dando SJ, King NP, Cripps AW, Schembri MA, Dougan G, Ulett GC.

646   2015. Discovery and Characterization of Human-Urine Utilization by

647   Asymptomatic-Bacteriuria-Causing *Streptococcus agalactiae*. Infect Immun

648   84:307-19.

649   30.   Sullivan MJ, Leclercq SY, Ipe DS, Carey AJ, Smith JP, Voller N, Cripps AW, Ulett

650   GC. 2017. Effect of the *Streptococcus agalactiae* Virulence Regulator CovR on

651   the Pathogenesis of Urinary Tract Infection. J Infect Dis 215:475-483.

652   31.   Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, Rozen

653   SG. 2012. Primer3--new capabilities and interfaces. Nucleic Acids Res 40:e115.

654 32. Untergasser A, Nijveen H, Rao X, Bisseling T, Geurts R, Leunissen JA. 2007.

655   Primer3Plus, an enhanced web interface to Primer3. Nucleic Acids Res 35:W71-

656   4.

657 33. Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R,

658   Nolan T, Pfaffl MW, Shipley GL, Vandesompele J, Wittwer CT. 2009. The MIQE

659   guidelines: minimum information for publication of quantitative real-time PCR

660   experiments. Clin Chem 55:611-22.

661 34. Sullivan MJ, Forde BM, Prince DW, Ipe DS, Ben Zakour NL, Davies MR, Dougan

662   G, Beatson SA, Ulett GC. 2017. Complete Genome Sequence of Serotype III

663   *Streptococcus agalactiae* Sequence Type 17 Strain 874391. Genome

664   Announcements 5.

665 35. Pfaffl MW. 2001. A new mathematical model for relative quantification in real-

666   time RT-PCR. Nucleic Acids Res 29:e45.

667 36. Eijkelkamp BA, Morey JR, Ween MP, Ong CL, McEwan AG, Paton JC, McDevitt

668   CA. 2014. Extracellular zinc competitively inhibits manganese uptake and

669   compromises oxidative stress management in *Streptococcus pneumoniae*. PLoS

670   ONE 9:e89427.

671 37. Cumley NJ, Smith LM, Anthony M, May RC. 2012. The CovS/CovR acid

672   response regulator is required for intracellular survival of group B *Streptococcus*

673   in macrophages. Infect Immun 80:1650-61.

674 38. Korir ML, Laut C, Rogers LM, Plemmons JA, Aronoff DM, Manning SD. 2017.

675   Differing mechanisms of surviving phagosomal stress among group B

676   *Streptococcus* strains of varying genotypes. Virulence 8:924-937.

677   39.   Anders S, Pyl PT, Huber W. 2015. HTSeq--a Python framework to work with

678   high-throughput sequencing data. Bioinformatics 31:166-9.

679   40.   Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and

680   dispersion for RNA-seq data with DESeq2. Genome Biol 15:550.

681   41.   Acharya D, Sullivan MJ, Duell BL, Goh KGK, Katupitiya L, Gosling D, Chamoun

682   MN, Kakkanat A, Chattopadhyay D, Crowley M, Crossman DK, Schembri MA,

683   Ulett GC. 2019. Rapid Bladder Interleukin-10 Synthesis in Response to

684   Uropathogenic *Escherichia coli* Is Part of a Defense Strategy Triggered by the

685   Major Bacterial Flagellar Filament FliC and Contingent on TLR5. mSphere 4.

686   42.   Leclercq SY, Sullivan MJ, Ipe DS, Smith JP, Cripps AW, Ulett GC. 2016.

687   Pathogenesis of *Streptococcus* urinary tract infection depends on bacterial strain

688   and beta-hemolysin/cytolysin that mediates cytotoxicity, cytokine synthesis,

689   inflammation and virulence. Scientific Reports 6:29000.

690   43.   Valdes Lopez JF, Urcuqui-Inchima S. 2018. Synergism between phorbol-12-

691   myristate-13-acetate and vitamin D3 in the differentiation of U937 cells to

692   monocytes and macrophages. Morphologie 102:205-218.

693   44.   Price EE, Boyd JM. 2020. Genetic Regulation of Metal Ion Homeostasis in

694   *Staphylococcus aureus*. Trends Microbiol 28:821-831.

695   45.   Harvie DR, Andreini C, Cavallaro G, Meng W, Connolly BA, Yoshida K, Fujita Y,

696   Harwood CR, Radford DS, Tottey S, Cavet JS, Robinson NJ. 2006. Predicting

697   metals sensed by ArsR-SmtB repressors: allosteric interference by a non-effector

698   metal. Mol Microbiol 59:1341-56.

699   46.   Janulczyk R, Ricci S, Bjorck L. 2003. MtsABC is important for manganese and

700   iron transport, oxidative stress resistance, and virulence of *Streptococcus*

701   *pyogenes*. Infect Immun 71:2656-64.

702   47.   Shabayek S, Bauer R, Mauerer S, Mizaikoff B, Spellerberg B. 2016. A

703   streptococcal NRAMP homologue is crucial for the survival of *Streptococcus*

704   *agalactiae* under low pH conditions. Mol Microbiol 100:589-606.

705   48.   Huang X, Shin JH, Pinochet-Barros A, Su TT, Helmann JD. 2017. *Bacillus*

706   *subtilis* MntR coordinates the transcriptional regulation of manganese uptake and

707   efflux systems. Mol Microbiol 103:253-268.

708   49.   Sullivan MJ, Goh KGK, Ulett GC. 2021. Cellular Management of Zinc in Group B

709   Streptococcus Supports Bacterial Resistance against Metal Intoxication and

710   Promotes Disseminated Infection. mSphere 6.

711   50.   Hood MI, Skaar EP. 2012. Nutritional immunity: transition metals at the

712   pathogen-host interface. Nat Rev Microbiol 10:525-37.

713   51.   Wagner D, Maser J, Lai B, Cai Z, Barry CE, 3rd, Honer Zu Bentrup K, Russell

714   DG, Bermudez LE. 2005. Elemental analysis of *Mycobacterium avium*-,

715   *Mycobacterium tuberculosis*-, and *Mycobacterium smegmatis*-containing

716   phagosomes indicates pathogen-induced microenvironments within the host

717   cell's endosomal system. J Immunol 174:1491-500.

718   52.   Odermatt A, Solioz M. 1995. Two trans-acting metalloregulatory proteins

719   controlling expression of the copper-ATPases of *Enterococcus hirae*. J Biol

720   Chem 270:4349-54.

721   53.   Portmann R, Poulsen KR, Wimmer R, Solioz M. 2006. CopY-like copper

722   inducible repressors are putative 'winged helix' proteins. Biometals 19:61-70.

723   54.   Beisel WR, Pekarek RS, Wannemacher RW. 1974. The impact of infectious

724   disease on trace-element metabolism of the host., p 217. *In* Hoekstra WG, Suttie

725   JW, Ganther HE, Mertz W (ed), Trace element metabolism in animals, 2

726   University Park Press, Baltimore, MD.

727   55.   Fleming RE, Whitman IP, Gitlin JD. 1991. Induction of ceruloplasmin gene

728   expression in rat lung during inflammation and hyperoxia. Am J Physiol 260:L68-

729   74.

730   56.   Percival SS. 1998. Copper and immunity. Am J Clin Nutr 67:1064S-1068S.

731   57.   Achard ME, Tree JJ, Holden JA, Simpfendorfer KR, Wijburg OL, Strugnell RA,

732   Schembri MA, Sweet MJ, Jennings MP, McEwan AG. 2010. The multi-copper-ion

733   oxidase CueO of *Salmonella enterica* serovar Typhimurium is required for

734   systemic virulence. Infect Immun 78:2312-9.

735   58.   Francis MS, Thomas CJ. 1997. Mutants in the CtpA copper transporting P-type

736   ATPase reduce virulence of *Listeria monocytogenes*. Microb Pathog 22:67-78.

737   59.   Schwan WR, Warrener P, Keunz E, Stover CK, Folger KR. 2005. Mutations in

738   the *cueA* gene encoding a copper homeostasis P-type ATPase reduce the

739   pathogenicity of *Pseudomonas aeruginosa* in mice. Int J Med Microbiol 295:237-

740   42.

741   60.   Carey AJ, Tan CK, Mirza S, Irving-Rodgers H, Webb RI, Lam A, Ulett GC. 2014.

742   Infection and cellular defense dynamics in a novel 17beta-estradiol murine model

743   of chronic human group B streptococcus genital tract colonization reveal a role

for hemolysin in persistence and neutrophil accumulation. J Immunol 192:1718-

31.

61.   Juttukonda LJ, Beavers WN, Unsihuay D, Kim K, Pishchany G, Horning KJ,

Weiss A, Al-Tameemi H, Boyd JM, Sulikowski GA, Bowman AB, Skaar EP. 2020.

A Small-Molecule Modulator of Metal Homeostasis in Gram-Positive Pathogens.

mBio 11.

62.   Takahashi S, Nagano Y, Nagano N, Hayashi O, Taguchi F, Okuwaki Y. 1995.

Role of C5a-ase in group B streptococcal resistance to opsonophagocytic killing.

Infect Immun 63:4764-9.

63.   Neubert MJ, Dahlmann EA, Ambrose A, Johnson MDL. 2017. Copper Chaperone

CupA and Zinc Control CopY Regulation of the Pneumococcal *cop* Operon.

mSphere 2.

TABLE 1. Transcriptional signature of Cu intoxication in GBS

Locus_ tag[†]	Genbank accession	Label	Annotation	FC	FDR
02565	ASZ00809.1	*copY*	CopY/TcrY family copper transport repressor	4.1	1.09E-12
02575	ASZ00811.1	*copZ*	carbonate dehydratase	4.0	1.02E-37
02570	ASZ00810.1	*copA*	copper-translocating P-type ATPase	3.7	2.25E-35
10970	ASZ02392.1		peptidoglycan-binding protein LysM	3.4	6.13E-13
07965	ASZ01821.1	*mtsA*	metal ABC transporter substrate-binding protein	2.9	2.15E-12
07960	ASZ01820.1	*mtsB*	metal ABC transporter ATP-binding protein	2.6	1.02E-15
10190	ASZ02237.1	*mntH2*	divalent metal cation transporter	2.4	6.34E-09
10965	ASZ02391.1		transglycosylase	2.2	1.87E-06
09125	ASZ02032.1		CHAP domain-containing protein	2.1	1.58E-09
07955	ASZ01819.1	*mtsC*	metal ABC transporter permease	2.1	2.62E-10
10305	ASZ02261.1		PAP2 family protein	2.0	6.13E-13
08045	ASZ01834.1		glycosyltransferase family 2 protein	-2.0	9.51E-06
08200	ASZ01861.1		branched-chain amino acid ABC transporter substrate-binding protein	-2.3	1.98E-04
10430	ASZ02284.1	*hvgA*	pathogenicity protein	-2.3	2.18E-07
04720	ASZ01199.1	*pcl1*	membrane protein	-4.3	1.38E-09
04185	ASZ01095.1	*fetB*	iron export ABC transporter permease subunit, FetB	-4.9	1.39E-08
04180	ASZ01094.1	*fetA*	ABC transporter ATP-binding protein FetA	-4.9	4.58E-11

758   [†] denotes locus tag of *S. agalactiae* 874391, preceded by CHF17_RS; FC, Fold-

759   Change; FDR, False Discovery Rate

 TABLE 2. Bacterial strains and Plasmids

Bacteria		
Strains	**Characteristics**	**Source**
E. coli DH5α	*huA2 lac(Δ)U169 phoA glnV44 Φ80' lacZ(Δ)M15 gyrA96 recA1 relA1 endA1 thi-1 hsdR17*	Bethesda Research Labs
S. agalactiae 874391	Wild type, Sequence type-17, Serotype III strain, Vaginal isolate (Japan)	(62)
S. agalactiae GU2691	874391Δ*copA* (*copA⁻* mutant) Locus tag: CHF17_RS02570	This work
S. agalactiae GU2857	874391Δ*copY* (*copY* mutant) Locus tag: CHF17_RS02565	This work
S. agalactiae GU3121	GU2857 (*ΔcopA*) containing *copYAZ* complement construct pGU3112; Sp	This work
Plasmids		
pHY304aad9	*ori* (Ts); temperature-sensitive shuttle vector; Sp	(29)
pGU2650	pHY304*aad9*-derivative *copA*Δ construct; Sp	This work
pGU2847	pHY304*aad9*-derivative *copY*Δ construct; Sp	This work
pGU3112	pDL278 containing cloned *copYAZ* operon; Sp	This work

762 **TABLE 3. Primers**

Oligo Name	Sequence (5′ → 3′)	Usage
M13F	GTAAAACGACGGCCAG	Sequencing
M13R	CAGGAAACAGCTATGAC	Sequencing
copA-Up-F1	GAGAATTCAAGCGCAGAATGGGAAATTA	Cloning
copA-Up-R1	CTAATGTCACTAACCTGCCCCCACATTTGGCGATAAGGTTG	Cloning
copA-Down-F1	CCAATTTTCGTTTGTTGAACTAATTTTGGTGGTCCCTTACTCAA	Cloning
copA-Down-R1	GACTCGAGTTTCTAATAGCTCCACCACCAAA	Cloning
copA-Chk-F1	GACTCGAGGCATCAAGACTAGTTGAGAGCG	Sequencing
copA-Chk-R1	CACGTTCAAGTGTTTTGCCGTCGACGGTTGTATTCTGCTGTGCCC	Sequencing
copY-UpF1	GGGCACAGCAGAATACAACCGTCGACGGCAAAACACTTGAACGTG	Cloning
copY-UpR1	GAGAATTCGCTGGTGAGCGTAGTGTATGT	Cloning
copY-DownF1	GGGCACAGCAGAATACAACCGTCGACGGCAAAACACTTGAACGTG	Cloning
copY-DownR1	GAGAATTCGCTGGTGAGCGTAGTGTATGT	Cloning
copY-ChkF1	AGGTTGAGGAATAGCCTCTTG	Sequencing
copY-ChkR1	TCCAAAGGAACCATATAGTCTTCA	Sequencing
copYAZ_compF1	CAGTGGATCCTGAGAAATGTTTTTCGGTTTACAA	Cloning
copYAZ_compR1	CAGTGTCGACAAAAAGAAGTGAGAAACTGCTCAAA	Cloning
dnaN-1F	CAACAAGAAAGCCGTCCAAT	qPCR
dnaN-1R	TCTGTCGCAACAGCCTTAAA	qPCR
hvgA-3F	TTTGGAATGGTCAGGAAGGA	qPCR
hvgA-3R	GCTGGTTTTGCAGCTTCTGT	qPCR
copY-1F	CAGAAAGTTTGCCAGACCAA	qPCR
copY-1R	GCATTGACACTGACCAGGAA	qPCR
copA-1F	CCTTATCGCCAAACGTGATT	qPCR
copA-1R	ATTGCTAATTGGTGCCGTTC	qPCR
0928_1F	TTGCGCGTGAAGAGAAACTA	qPCR
0928_1R	CCATTGAGCATGTTCGTGAC	qPCR
0929_1F	GCCCAATCACCTTTTACGAA	qPCR
0929_1R	GCGGCAATTCCAACATATTC	qPCR
2108_1F	GCGCATAACCATCACTCAGA	qPCR
2108_1R	TTGGCAGTTTCTAAGGCAGTC	qPCR

FIG 1. Global transcriptomic analysis of GBS in response to Cu stress. Volcano plot

showing data from RNASeq analysis of WT GBS cultures exposed to 0.5 mM Cu

compared to non-exposed controls. Transcripts detected as up- or down-regulated in

response to Cu (n=4, >± 2-fold, FDR <0.05) are highlighted in red and blue,

respectively. Dotted lines show False discovery rate (FDR; q-value) and fold-change

cut-offs. Grey points indicate genes that were not significant changed in expression,

according to these analysis cut-offs. Selected genes are identified with black lines. B,

Validation of RNASeq data. Expression ratio (Fold-change) of *pcl1*, *hvgA* and *copY*

quantified by qRTPCR in THB medium containing 0.5 mM Cu compared to THB without

Cu. Ratios for qRTPCR were normalized using housekeeping *dnaN* and RNASeq ratios

were calculated using DESeq2. Bars show means and S.E.M from 4 independent

experiments.

FIG 2. Organisation of the *copY-copA-copZ* locus in GBS. A, *copY-copA-copZ* are

adjacent in the GBS genome and likely controlled by the promoter-proximal *copY,*

encoding a putative Cu-sensing repressor. Locus tags from the GBS 874391 genome

are indicated. B, Distribution of homologous *cop* genes in other Lactobacillales,

arranged by percentage identity of amino acid sequence to CopA of *S. agalactiae*. C,

Model of Cu efflux in GBS highlighting CopA as transmembrane Cu exporter,

transcriptionally repressed by CopY in the absence of Cu, which is likely inhibited by the

Cu-binding chaperone protein CopZ. Figure based on previous studies in other

lactobacillales (53, 63).

FIG 3. Growth curve analyses of GBS cultured in nutrient-rich THB medium (A), or in

THB supplemented with 0.5 mM Cu (B), 1.0 mM Cu (C) or 1.5 mM Cu (D), comparing

WT, $\Delta copA$ or $\Delta copY$ strains. Complementation of $copA$ mutation ($\Delta copA$+C) restored

growth of GBS to WT levels in conditions of Cu stress. Points and bars show mean and

S.E.M of several independent experiments (5 for WT, $\Delta copA$ and $\Delta copY$, 2 for

complemented strain) monitoring attenuance at 600nm.

FIG 4. Bactericidal effect of Cu on GBS viability. Time-kill assays comparing WT, $\Delta copA$

or $\Delta copY$ GBS (and complemented strain $\Delta copA$+C), incubated in CDM or in CDM

supplemented with 0.05, 0.1, 0.2, 0.5 or 1 mM Cu. Viable cells were quantified at 1h,

3h, 6h and 24h post incubation. [†] Viable Cell counts of 0 CFU/mL were assigned a value

of 1 to plot on $\log_{10}$ y-axes. Points and bars show mean and S.E.M of 4 independent

experiments. Data were analysed by One-way ANOVA with Holm Sidak Multiple

comparisons. Significance markers represent comparisons of the number of surviving

bacteria in conditions of the defined Cu concentration [Cu] vs the number of bacteria in

the non-exposed control (0 mM Cu) for the same time point and same strain (* $P < 0.05$).

FIG 5. Expression analysis of $copA$ and intracellular metal content in GBS strains. A,

Expression ratio (Fold-change) of $copA$ quantified by qRT-PCR in THB medium

containing 0.25, 0.5, 1.0 and 1.5 mM Cu, compared to THB without Cu. B, Relative

$copA$ transcripts were quantified in WT and $\Delta copY$ strains with and without Cu

supplementation (0.5 mM) to demonstrate de-regulation of $copA$ expression in the

$\Delta copY$ background. Intracellular accumulation of Cu (C), Mn (D), Fe (E) and Zn (F) was

compared with and without Cu supplementation (0.5 mM) in WT, Δ*copY* and Δ*copA*

strains. Ratios in A were calculated as described previously (35) using C_T values, primer

efficiencies and housekeeping *dnaN*. Bars show means and S.E.M from 3-4

independent experiments and compared by One-way ANOVA with Holm-Sidak multiple

comparisons (* $P < 0.05$; *** $P < 0.001$). Ctrl: Control condition (i.e., no supplemental Cu

in media). The ANOVA P value from comparisons of between strains was 0.06 for

Cellular Fe and 0.01 for Cellular Zn; subsequent pairwise Student's *t*-tests were used to

compare each strain in the Ctrl and Cu condition (* $P < 0.05$).

FIG 6. Interactions of GBS with human macrophage-like cells. Gentamicin protection

assays with WT and Δ*copA* strains in human (U937 monocyte-derived macrophage-like)

macrophages in media with and without 20 μM supplemental Cu. Surviving bacteria are

expressed as CFU/mL, indicating the number of intracellular bacteria at 1h after

antibiotic treatment, and at 24h and 48h post infection (h.p.i). Data are means and

S.E.M of 3-4 independent experiments.

FIG 7. Virulence of WT (grey circles) or Δ*copA* (blue diamonds) GBS in a mouse model

of disseminated infection. C57BL/6 mice (6-8 weeks old) were intravenously injected

with 10^7 bacteria; bacteremia and disseminated spread to liver and spleen were

monitored at 24h post infection. CFU were enumerated and counts were normalized

using tissue mass in g. Viable Cell counts of 0 CFU/mL were assigned a value of 1 to

enable visualisation on $\log_{10}$ y-axes. Lines and bars show median and interquartile

ranges and data are pooled from 2 independent experiments each containing n=10

mice compared using Mann-Whitney U-tests (*P < 0.05, **P < 0.01).

A Cu stress transcriptome in *S. agalactiae*

B Validation by qRTPCR