HIGHER VIRAL VIRULENCE SELECTS FOR FASTER EVOLUTION OF HOST RESISTANCE

CAROLIN C. WENDLING[1,2*]
JANINA LANGE[1]
HEIKO LIESEGANG[4]
ANJA PÖHLEIN[4]
BOYKE BUNK[5]
JELENA RAIKOV[1]
HENRY GÖHLICH[1]
OLIVIA ROTH[1†]
MICHAEL A. BROCKHURST[5†]

Addresses:

[1]GEOMAR Helmholtz Centre for Ocean Research Kiel, Marine Evolutionary Ecology, Düsternbrooker Weg 20, 24105 Kiel, Germany

[2]ETH Zürich, Institute of Integrative Biology, Universitätstrasse 16, CHN D 33, 8092 Zürich, Switzerland

[3]Georg-August-University Göttingen, department of genomic and applied microbiology, Grisebachstr 8 37077 Göttingen, Germany.

[4]Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures, Department Bioinformatics and Databases, Inhoffenstr. 7B, 38114 Braunschweig, Germany

[5]Division of Evolution and Genomic Sciences, University of Manchester, Dover Street, Manchester M13 9PT, UK

*Corresponding author:
E-mail: carolin.wendling@env.ethz.ch; Phone: +41 44 633 80 26

† shared last author

ABSTRACT

Infectious organisms can vary tremendously in their virulence. While the evolution of virulence and different levels thereof has received much attention over the past decades, the evolution of host resistance in response to different levels of virulence is far less understood and as of now mainly based on theoretical predictions. Increasing virulence can drive the evolution of host resistance if the costs of resistance are outweighed by the benefits of avoiding infection. Accordingly, we hypothesised that higher virulence will therefore select for faster resistance evolution, and ultimately shorter epidemics if parasites fail to overcome evolved host resistance. To test this hypothesis, we performed a bacteria-phage co-evolution experiment using two filamentous phages that differ in their virulence. Overall, we found two alternative defence strategies against filamentous phage infections: (1) immunity through super infection exclusion whereby the phage becomes part of the bacterial genome and protects its host against subsequent phage infections, and (2) resistance: surface-receptor modifications which prevent phages from entering the host cell. While immunity emerged rapidly and at a similar rate against both viruses, it did not persist long against the higher virulence virus, against which resistance evolved significantly faster compared to the lower virulence virus. The faster replacement of immunity by resistance in populations with higher virulence infections was driven by higher costs of being infected outweighing the fitness costs of resistance mutations. This resulted in a faster extinction of higher virulence viruses and implies that more virulent infections should cause shorter epidemics when viruses are unable to overcome evolved host resistance. Given that closely related viruses often vary substantially in their virulence, these findings suggest that resistance evolution will play a greater role in the epidemiological dynamics of higher virulence infections.

65   **Keywords**: virus, virulence, filamentous phages, experimental evolution, resistance

66   evolution

INTRODUCTION

Infectious organisms vary strikingly in their level of virulence and the resulting selection they impose on their host. While some human viruses such as Ebola or rabies can cause a deadly disease, others (including many cold viruses) often remain asymptomatic. Even closely related viruses, such as different strains of myxoma (Fenner and Marshall 1957) or corona viruses (Weiss and Leibowitz 2011), can differ greatly in virulence. The evolution of different levels of virulence has been suggested to result from a trade-off between parasite reproduction and transmission (Anderson and May 1982, May and Anderson 1983). Higher reproduction results in higher virulence but also in higher host morbidity and mortality and thus limited chances for transmission. Accordingly, virulence may be expected to peak at intermediate levels, maximising onward transmission. However, this pattern is not universal and selection for virulence is expected to vary according to a wide variety of epidemiological parameters (Alizon, Hurford et al. 2009).

The evolution of virulence has been studied extensively during the last two decades, both using selection experiments (Bull, Molineux et al. 1991, Turner, Cooper et al. 1998, Messenger, Molineux et al. 1999) and observations of parasites evolved in nature (Herre 1993, Ebert 1994). What remains open, however, is the question of how virulence will impact evolutionary trajectories of resistance in a host population, and how these trajectories change with different levels of virulence. Severe disease symptoms resulting from highly virulent infections will significantly reduce host fitness imposing strong selection for hosts to quickly acquire resistance, for instance through evolutionary changes. Thus, we expect benefits of host resistance relative to the costs of disease symptoms to be higher against highly virulent compared to low virulent infections. If this holds true, high virulence will lead to a stronger selection for and thus

faster evolution of host resistance. Ultimately this should lead to shorter epidemics if

parasites fail to overcome these evolved host resistances.

To explore how variation in virulence between closely related viruses influences

the dynamics of host resistance evolution, we designed a co-evolution experiment using

the model bacterium *Vibrio alginolyticus* K01M1 as a host and two closely related

versions of the filamentous *Vibrio* phage VALGΦ8, that differ in their replication rate

and hence in their virulence (Table 1)(Chibani, Hertel et al. 2020). Filamentous phages

(family *Inoviridae*), i.e., long, thin proteinaceous filaments which contain a circular

single-stranded DNA genome have been shown to be ideal model systems to study

virulence evolution (Bull, Molineux et al. 1991, Messenger, Molineux et al. 1999).

These phages can establish chronic infections whereby virions are continuously released

without lysis. Although filamentous phages do not kill their host, infections cause harm

by reducing host growth rates. This is because the host cell pays the metabolic costs

resulting from phage replication and through phage-encoded proteins inserted into the

bacterial membrane (Mai-Prochnow, Hui et al. 2015). Thus, the virulence of

filamentous phages, which can vary greatly across phage types (Rakonjac 2012), can be

directly quantified by measuring the reduction in bacterial growth rate caused by phage

infection.

By combining experimental evolution and whole genome sequencing, we show that

bacteria can defend themselves against these infections via two alternative strategies:

(1) immunity (i.e., infected bacteria became resistant to further infection) and (2)

resistance (surface receptor modifications). We show that immunity is faster than

resistance evolution and occurs at a similar rate in both treatments. In contrast, selection

for resistance evolution is stronger (because immunity is more costly), and this form of

resistance evolves more rapidly against high virulence compared to low virulence

phages. This resulted in faster extinction and ultimately shorter epidemics for high

virulence phage.

MATERIAL AND METHODS

(a) Strains and culture conditions

Experiments were conducted using the *Vibrio alginolyticus* strain K01M1 (Chibani,

Roth et al. 2020). K01M1 contains one integrated filamentous *Vibrio* phage VALGΦ6

(later called: resident K01M1Φ-phage throughout the manuscript), which replicates at a

very low frequency (Chibani, Hertel et al. 2020). Compared to other, closely related *V.*

alginolyticus strains, K01M1 is highly susceptible to infections by filamentous phages

(Wendling, Piecyk et al. 2017). For the selection experiment we used two different

versions of the filamentous *Vibrio* phage VALGΦ8 (Table 1), one integrative (isolated

from the host strain K04M5) and one episomal (isolated from the host strain K04M1).

While both phages have been shown to significantly reduce the growth of K01M1

(Wendling, Piecyk et al. 2017, Wendling, Goehlich et al. 2018), infections with the high

virulent phage impose a significantly stronger reduction in bacterial growth than

infections with the low virulent phage. All experiments were carried out in liquid

medium (Medium101: 0.5% (w/v) peptone, 0.3% (w/v) meat extract, 3.0% (w/v) NaCl

in MilliQ water) at 25° C in 30-ml microcosms containing 6 ml of medium with

constant shaking at 180 rpm.

Table 1 Bacteria and phages (including NCBI accession numbers) used in the present study.

Isolate	Accession Number(s)	Phages	Role in evolution experiment
V. alginolyticus K01M1	CP017889.1 CP017890.1	*Vibrio* phage VALGΦ6	Host strain during evolution experiment
V. alginolyticus K04M1	CP017891.1 CP017892.1	*Vibrio* phage VALGΦ6 *Vibrio* phage VALGΦ8 (episomal)	Donor of the episomal, low virulent phage
V. alginolyticus K04M5	CP017899.1 CP017900.1	*Vibrio* phage VALGΦ6 *Vibrio* phage VALGΦ8 (integrative on CR2)	Donor of the integrative, high virulent phage
Vibrio phage VALGΦ6	MN690600		Resident phage in ancestral K01M1
Vibrio phage VALGΦ8	MN719123		Co-evolving phage in evolution experiment

(b) Selection experiment

Six replicate populations were founded for each of three treatments from

independent clones of K01M1. Treatments comprised (a) a high virulent, integrative

version of the filamentous *Vibrio* phage VALGΦ8, later called VALGΦ8$_{K04M1}$, (b) a

low virulent, episomal version of the filamentous *Vibrio* phage VALGΦ8, later called

VALGΦ8$_{K04M5}$, and (c) no phage as control. Each population was established from 60 µl

of an independent overnight culture (5×10^8 CFU/ml). At the beginning of the

experiment, we inoculated phage-containing treatments with 300 µl of a 5×10^{10} PFU/ml

stock solution. Populations were propagated by transferring 1% to fresh medium every

24 hours for a total of 30 transfers. On transfer T0, T1, T2 followed by every other

transfer, phage and bacterial densities were determined, as described below and whole

population samples were frozen at -80° C at a final concentration of 33% glycerol. In

addition, on transfer T0, T1, T2, T6, followed by every sixth transfer 24 single colonies

were isolated at random from each population and stored at -80° C. Two populations

from the control treatment tested positive for virus infection, indicating contamination,

were excluded from later assays.

(c) Bacterial and phage densities

Bacterial densities: bacterial densities were determined by plating out 100 µl of a

dilution series ranging from 10^{-5} to 10^{-7} on *Vibrio* selective Thiosulfate Citrate Bile

Sucrose Agar (TCBS) plates (Fluka Analytica). Plates were incubated over night at 25°

C and the total amount of colonies was counted the following day.

Phage densities: Quantification of filamentous phages by standard spot assays is

often not possible (Rakonjac 2011). Instead of typical lytic plaques we mostly observed

opaque zonas of reduced growth. Thus, we used spectrometry to quantify phage

prevalence (http://www.abdesignlabs.com/technical-resources/bacteriophage-

spectrophotometry), which uses the constant relationship between the length of viral

DNA and the amount of the major coat protein VIII of filamentous phages, which,

together, are the main contributors of the absorption spectrum in the UV range. The

amount of phage particles per ml can be calculated according to the following formula:

170
$$phages/ml = \frac{(OD269 - OD320)*6e16}{bp},$$

where OD269 and OD320 stand for optical density at 269 and 320 nm and bp stands for

number of base pairs per phage.

This method is based on small-scale precipitation of phages by single PEG-

precipitation. After centrifuging 1500 µl of the phage containing overnight culture at

13,000 ×g for 2 min, 1200 µl of the supernatant was mixed with 300 µl PEG/NaCl 5×

and incubated on ice for 30 min. Afterwards phage particles were pelleted by two

rounds of centrifugation at 13,000 ×g for 2 min, resuspended in 120 µl TBS 1× and

incubated on ice. After one hour the suspension was cleaned by centrifugation at 13,000

×g for 1 min and absorbance was measured at 269 and 320 nm.

Quantification of filamentous phages using spectrometry is likely to be erroneous if

viral load is low. Therefore, we additionally quantified phage prevalence/ phage

extinction in each of the populations on every second transfer day by standard spot

assays with a serial dilution (10^{-1} to 10^{-6}) on the ancestral host (for details see

(Wendling, Piecyk et al. 2017)) and measured until which dilution the typical opaque

zones of reduced bacterial growth were visible.

(d) Measuring phage-resistance

We measured the rate of phage resistance evolution among bacteria against the

ancestral phage by determining the reduction in bacterial growth rate (RBG) imposed by

the phage, adapted from (Poullain, Gandon et al. 2008) with some modifications

according to (Goehlich, Roth et al. 2019). Twenty-four random colonies from each

population from transfer T0, T1, T2, T6, T12, T18, T24, and T30 were introduced into

96-well microtiter plates containing Medium101 at a concentration of 5×10^6 cells/ml

and inoculated with ~2.5×10^6 PFU/ml of each of the two ancestral phages used for the

selection experiment or without phage (control). Absorbance at 600 nm was measured

using an automated plate reader (TECAN infinite M200) at T0 and again after 20 hours

of static incubation at 25°C. The reduction in bacterial absorbance 'RBG' was

calculated according to the following formula:

199
$$RBG = \frac{OD600(t=20) - OD600(t=0)[Phage]}{OD600(t=20) - OD600(t=0)[Control]},$$

where OD stands for optical density at 600nm.

(e) Frequency of prophage carriage

On transfer T0, T1, T2, T6 followed by every sixth transfer we measured the

frequency of phage carriage of 24 random clones per population using standard PCR.

Primers (VALGΦ8_Forward TGGAAGTGCCAAGGTTTGGT, VALGΦ8_Revers

GAAGACCAGGTGGCGGTAAA) that specifically target the co-evolving *Vibrio*

phage VALGΦ8 have been designed using NCBI Primer-BLAST webpage

(httdol://www.ncbi.nlm.nih.gov/tools/primer-blast/). Note, while this primer-pair detects

the presence/ absence of *Vibrio* phage VALGΦ8, it does not confirm chromosomal

integration of the respective phage. Glycerol stocks were inoculated overnight (25°C,

180 rpm) in Medium 101 and subsequently diluted (1:10) in HPLC purified H_2O and

frozen at -80° C. One µl of this suspension was used as DNA template in the PCR

assay. Reaction comprised 1 µl Dream Tag Buffer, 0.1 µl Dream Tag DNA polymerase

(Thermo Scientific, USA), 4.9 µl H_2O, 1 µl dNTPs [5 mM] and 1 µl of each primer [50

µM]. The amplification program used consisted of: (i) 3 min at 95° C, (ii) 35 cycles of

45 sec at 95° C, 30 sec at 63° C, 45 sec at 72° C, (iii) 7 min at 72° C. Afterwards, 5 µl

of each reaction was mixed with 2 µl loading dye (10×) and loaded onto a 1.2% agarose

gel dissolved in 1×TAE gel buffer. GeneRuler Express DNA-ladder was used as size

marker. Agarose gels were run 15 min at 70 V in 0.5× TAE running buffer and

subsequently stained with ethidium bromide for 10 min. DNA was visualized using UV

light and documentation took place using Intas Gel iX20 Imager. Phage presence was

recorded positive if a PCR product of 1400 bp was visible.

For all subsequent assays, we randomly picked one phage-resistant clone with a

positive PCR product (later called: Φ-carrier) and one phage-resistant clone with a

negative PCR product (later called: mutant) from each phage-evolved population as

well as two randomly selected non-resistant clones from the control populations.

(f) Competition experiments

To determine differences in fitness between both resistance forms, we measured the

competitive fitness of Φ-carrier relative to mutants. Each competition culture was done

in triplicates as described in (Harrison, Guymer et al. 2015). In brief, overnight cultures

of both competing strains (of which one was labelled with a GFP-marker) were mixed

1:1 and 60 µl of this mixture was inoculated to 6 ml Medium 101 to initiate each

competitive culture. After 24 hours, fitness was estimated by means of flow cytometry

(FACS-Caliburm Becton & Dickinson, Heidelberg, GER), where absolute fluorescent

cells and non-fluorescent cells were calculated. Competitive fitness was estimated as the

ratio in Malthusian parameters (Lenski, Rose et al. 1991):

$W = \ln(abundance_{t=24}/abundance_{t=0})_{competitor1} / \ln(abundance_{t=24}/abundance_{t=0})_{competitor2}$

(g) Bacterial growth rate and phage production

To determine fitness parameters that could explain observed differences in

competitive fitness we additionally quantified bacterial growth rate (μ) by means of 24-

hour growth curves and phage production using PEG precipitation (as described in (c))

of the same clones used for the competition assays (i.e., one Φ-carrier and one mutant

from each phage-treated population and two random phage-susceptible clones from the

control populations plus the ancestor).

(h) Whole genome sequencing

We used a combination of long- and short read sequencing to obtain complete

genomes of the same clones from the assays above, i.e., one Φ-carrier and one mutant

from each phage-treated population and one random phage-susceptible clone from each

control population, which corresponds to six independently evolved clones per

treatment and resistance form. High molecular weight DNA was extracted from cell

pellets of overnight cultures following the protocol for gram negative bacteria from the

DNeasy Blood & Tissue Kit (Qiagen, Hilden, Germany). For long-read sequencing the

library was prepared using Pacific Bioscience protocol for SMRTbell^{TM} Libraries using

PacBio$^®$ Barcoded Adapters for Multiplex SMRT$^®$ Sequencing. To do so, DNA was

fragmented into 10kb fragments using g-tubes (Covaris). Samples were pooled during

library preparation aiming for equimolar pooling and library size was selected using

Ampure beads. The library was sequenced on a PacBio Sequel instrument using Sequel

Polymerase v3.9, SMRT cells v3 LR and Sequencing chemistry v3.0. Loading was

performed by diffusion. Two SMRT cells were sequenced (movie time: 600min, pre-

extension time: 240 min). Reads were demultiplexed using Barcoding pipeline on

SMRT Link (v6.0.0.47841, SMRT Link Analysis Services and GUI v6.0.0.47836) with

40 as a minimum barcode score.

Short-read sequencing was done on an Illumina 2500 platform and resulted in a

minimum average coverage of 88× per strain (coverage range was from 88× to 157×).

The reads were quality controlled using the program FastQC Version 0.11.5. High

quality reads were used for hybrid assemblies as well as for single nucleotide variation

analysis.

Genome assemblies were performed in two different ways: In a first approach

assemblies were performed using the Unicycler pipeline (v0.4.7) and the programs

within. Assemblies were performed as hybrid assemblies using short-read and long read

data in a short-read first approach. In brief: An initial assembly was performed with

short-read only using spades (v3.13.0) as provided within Unicycler. The resulting

contigs were co-assembled with long-read data using miniasm (v0.2-r168) and curated

using the racoon software. This step resulted in complete closed replicons. All long

reads were mapped and integrated into the contigs. All replicons were polished using

the Pilon software (1.22) to clear any small-scale assembly errors. Finally, all replicons

were rearranged according to the origin of replication. The assembly for the ancestral

K01M1 strain, as has been described in (Wendling, Piecyk et al. 2017) was performed

following the Hierarchical Genome Assembly Process (HGAP3) protocol, developed

for Pacific Biosciences Single Molecule Real-Time (SMRT) sequencing data (Chin,

Alexander et al. 2013). HGAP is available for use within PacBio's Secondary Analysis

Software SMRTPortal. Methodically, the longest subreads of a single SMRT Cell

(usually 25x genome coverage, e.g., 25 x 5 Mbp = 125 Mbp) are being chosen to be

error-corrected with "shorter" long reads in a process named preassembly. Hereby, a

length cut-off is computed automatically separating the "longer" reads (for genome

assembly) and the "shorter" reads (for error-correction). The level of error-correction is

being estimated with a per-read accuracy of 99%. Finally, error-corrected long read data

is being assembled with Celera Assembler 7.0.

(i) SNV analysis and reconstruction of infecting phages

All short-read sequences were mapped on a high quality closed reference genome

of *Vibrio alginolyticus* Strain K01M1 (Wendling et al., 2017) using Bowtie2

(Langmead & Salzberg, 2012). Single nucleotide variation (SNV) analysis was done

using the Breseq pipeline as described in Deatherage & Barrick, (Deatherage & Barrick,

2014). Assembly of reads were done using Spades (Bankevich et al., 2012). Whole

genome alignments have been calculated using the MAUVE aligner (Darling, Mau, &

Perna, 2010). Presence of infecting phage genomes were determined by assembling

NGS-reads that did not map on the K01M1 genome in a bowtie2 mapping using Spades.

The resulting contigs were annotated based on the review of Mai-Prochnow on

filamentous phages (Mai-Prochnow et al., 2015). The genomes of the evolved phages

were compared to the infecting phage genomes *Vibrio* phage VALGΦ8 as well as to the

genome of the resident prophage *Vibrio* phage VALGΦ6 from the challenged strain

K01M1 using BLAST and Easyfig 2.1 (Sullivan, Petty, & Beatson, 2011).

Coverage analysis of phage derived short reads were mapped against the complete

ancestral genome of *V. alginolyticus* K01M1 (NCBI accession numbers CP017889.1,

CP017890.1) as well as the phage genomes VALGΦ6 and VALGΦ8 using Bowtie2.

The resulting coverage data was visualized using Artemis (Version) and compared for

differences in coverage. Coverage analysis of Sequel long-reads was performed within

the "Base Modification Analysis" within SMRTlink 8 using the same reference as stated

above or additionally including the infecting phage. Coverage data was visualized

within SMRTlink.

(j) Statistical analyses

All statistics were performed in the R 3.1.2 statistical environment (Team 2011).

For all analysis aimed to compare the two different phage treatments to one another,

control populations (i.e., those that evolved without phages) were excluded. When

comparing temporal dynamics between phage-treatments, we excluded the starting

time-point T0, because these measurements were taken before phages were added to the

populations.

Bacteria and phage dynamics

Bacterial and phage densities were analysed over time using a generalized least

squares model to control for autocorrelation of residuals over time using the gls function

(package nlme) with phage treatment, transfer as categorical variable as well as their

interaction as fixed effect.

We considered phages to be prevalent in the population if opaque zones of reduced

growth were visible during standard spot assays. Phage prevalence was subsequently

quantified by a serial dilution, which were assigned with invers values (i.e., if reduced

growth zones were visible up to dilution of 10^{-6} we assigned to it a value of 7, whereas

if they were only visible on an undiluted spot, we assigned to it a value of 1, if no zone

of reduced growth was visible it was scored as 0). Phage extinction events across phage-

treatments were analysed using a log-rank test.

Measuring bacterial resistance

We observed a bimodal histogram on all RBG values with a local minimum at RBG

= 0.82 (Figure S1). Thus, we considered an infection as positive if RBG < 0.82. The

proportion of resistant clones per population as well as the proportion of clones that

tested positive for PCR (targeting the co-evolving phage) were analysed using a

generalized linear model with a binomial error distribution using the glm function

(package lme4) with phage treatment, transfer and their interaction as fixed effect.

Fitness effects

We determined differences in relative fitness between MSHA-mutants and phage-

carrier using a linear model with resistance mechanisms and GFP-label and the

interaction thereof as fixed effects. To determine differences in the amount of free

phages and in growth rates produced between ancestral strains and evolved strains and

between both resistance forms, we used Welch's pairwise *t*-tests with sequential

Bonferroni correction. We further performed a Pearson's correlation analysis to

determine whether phage production impacted bacterial growth rates.

RESULTS

Bacterial densities

We propagated six replicate populations of *Vibrio alginolyticus* K01M1 with or without either of two closely related filamentous phages that differed in their virulence — VALGΦ8$_{K04M5}$ (high virulence), VALGΦ8$_{K04M1}$ (low virulence) — for 30 serial transfers (~240 bacterial generations). Phages reduced bacterial densities in both phage-containing treatments by several orders of magnitude compared to no phage control populations (Figure 1a). However, the immediate reduction (measured 24 hours post infection [hpi]) in bacterial density was stronger in populations co-evolving with the high virulence phage than with the low virulence phage (Figure 1a), confirming that VALGΦ8$_{K04M5}$ is more virulent than VALGΦ8$_{K04M1}$. Over time, however, the densities of bacterial populations co-evolving with the high virulence phage recovered three times faster than populations co-evolving with the low virulence phage (significant phage:transfer interaction in gls-model: $F_{15,186}$=6.58, p<0.001, Figure 1a).

Phage densities

In both phage containing treatments, phages, that had initially been added at a titre of ~5 ×10^{10} PFU/ml, amplified massively during the first couple of bacterial generations and reached levels of 3.01×10^{12} PFU/ml (VALGΦ8$_{K04M5}$) 24 hpi and 2.83×10^{12} PFU/ml (VALGΦ8$_{K04M1}$) 48 hpi (Figure 1b), before densities decreased to levels comparable to phage-free populations, which stem from the resident phage VALGΦ6. These data suggest that the strong reduction in bacterial densities at the beginning of the experiment (Figure 1a) directly resulted from the production of viral particles (Figure 1b).

377   We further observed that phage extinction events differed significantly between

378   treatments (log-rank test: Chisq$_1$=4.9, p=0.03). High virulent phages went extinct in five

379   out of six populations after 12 transfers, whereas low virulent phages survived in four

380   out of six populations until transfer 28, before they finally went extinct on transfer 30

381   (Figure S2).

Figure 1 Population dynamics over 30 transfers of bacteria-phage co-evolution. (a) Bacteria in CFU/ml, (b) Phages in PFU/ml, the grey dashed line represents the detection limit for quantifying filamentous phages using spectrophotometry, (c) Fraction of phage-resistant clones (n=24), and (d) Fraction of Φ-carrier within phage-resistant clones. Fractions are based on 24 random clones per replicate population. In all panels, data are represented as mean of six replicate populations per treatments, error bars represent standard errors. Colours correspond to one of three phage-treatments, low virulent VALGΦ8$_{K04M1}$ (red), high virulent VALGΦ8$_{K04M5}$ (yellow), no phage (grey).

Modes of phage resistance selected by high and low virulence phage

To understand the evolutionary drivers behind these different selection dynamics

against both phages, we next isolated 24 random clones per population from seven

timepoints and quantified the fraction of phage-resistant clones per population. The

proportion of phage-resistant clones per population increased rapidly within the first 24

hours (Figure 1c). This was true for both phage treatments in which we observed almost

100% phage-resistant clones after 24 hours, and subsequently resistance levels remained

high in both treatments for the duration of the experiment.

Different mechanisms can confer resistance to filamentous vibriophages, such as

superinfection exclusion, (i.e. when the same phage is already present in the host cell

(Wendling, Piecyk et al. 2017)) or surface-receptor modifications, in particular at the

mannose-sensitive hemagglutinin (MSHA) type IV pilus, that prevents filamentous

vibriophages from entering the host (Jouravleva, McDonald et al. 1998). To determine

the underlying genetic basis of the resistance mechanism in the present study, we first

performed a PCR-based analysis to determine presence of newly acquired phages in

bacterial genomes, which confers resistance through superinfection exclusion. After 24

hours, Φ-carriers, were the dominating form among the resistant clones (Figure 1d),

suggesting that superinfection exclusion is a fast way to acquire phage resistance.

However, 48 hours post infection (i.e., on transfer 2) the proportion of Φ-carriers

declined, and did so significantly faster in populations co-evolving with the high

virulence, compared to the low virulence phage (Figure 1d, significant phage:transfer

interaction: $F_{6,60}=10.18$, p<0.001).

To test if the decline of Φ-carriers was driven by the invasion of surface receptor

modification resistance we used whole genome sequencing (WGS) of two randomly

chosen clones from each population (one PCR-positive and one PCR-negative clone) to

identify mutations. We observed no loci with mutations on chromosome 2 or the

plasmid pl9064, but on chromosome 1 we identified 12 loci with mutations that were

not present in clones from the control treatment, suggesting an association with phage-

mediated selection. Of these 12 loci, three were randomly distributed across PCR-

positive and PCR-negative clones while the remaining nine loci were exclusive to PCR-

negative clones suggesting a potential role in evolved phage resistance. Of these nine

loci, eight had substitutions, duplications, insertions or deletions in four different

proteins belonging to the MSHA type IV pilus operon (*mshL, mshE, mshG,*

K01M1_28150; Figure 2a/ Table S1). Of these, five caused severe frameshift mutations

that presumably have a high impact on the function of this protein. Overall mutated

MHSA-loci occurred in 8/12 PCR-negative clones, which suggests strongly parallel

evolution of phage resistance. The MSHA operon is highly conserved across *Vibrio*

clades (Figure 2b), suggesting that, similar to other vibrios (Jouravleva, McDonald et al.

1998), the MSHA type IV pilus plays an important role in resistance against the

filamentous *Vibrio* phage VALGΦ8. Note, a search of all assembled genomes for

CRISPR associated genes as well as for CRISPR array like repetitive sequence patterns

did not yield any results. All PCR-negative phage resistant clones are from here

onwards referred to as MSHA-mutants.

The genomic data also confirmed that clones with a positive PCR result (i.e., Φ-

carrier) did indeed contain the respective co-evolving phage. Interestingly, in all

sequenced PCR-positive clones, the co-evolving phage existed exclusively as an

episome, despite the K01M1 genome containing at least two known integration sites

((Chibani, Hertel et al. 2020) and Supplementary material Figure S3). Moreover, we

observed no genomic changes between ancestral and evolved phage genomes, nor at the

integration sites of the ancestral and evolved clones, suggesting that the failure to

integrate into the bacterial chromosome was not due to mutation either of the phage or

the integration site.

444
Figure 2 (A) Genetic loci on chromosome 1 under positive selection as indicated by parallel genomic
evolution in populations co-evolving with phages: left: Φ-carrier; middle: mutants; right zoom into
MSHA-operon region from mutants. Only loci which are not present in control populations are shown.
Concentric circles correspond to one clone isolated from either the VALGΦ8$_{K04M5}$ treatments (5 inner
circles, yellow) or the VALGΦ8$_{K04M1}$ treatment (six outer circles, red). Each coloured point corresponds
to one mutation event on the respective clone. HP corresponds to locus tag K01M1_28150. For more
detailed information on the underlying mutation see Table S1.
(B) Structure of the MSHA-operon and comparative genomics comprising MSHA operons from *V.*
alginolyticus FA2 (top), *V. alginolyticus* K01M1 (middle), and *V. cholerae* El Tor (bottom). Similarity
between regions is indicated by dark grey blocks, genes with detected mutations are marked in orange,
detected mutations are marked as arrows below *V. alginolyticus* K01M1.
456

Fitness effects of contrasting modes of resistance

Φ-carriers went to extinction in all six populations co-evolving with the high

virulence phage 12 days post infection but were able to persist until the end of the

experiment (i.e., transfer 30), albeit at very low frequencies, in five out of six

populations co-evolving with the low virulence phage. We hypothesised therefore that

the fitness benefit of MSHA-mutants relative to Φ-carriers would be higher for the high

virulence phage compared to the low virulence phage, because MSHA-mutants do not

suffer virulence from phage infection. This was confirmed in a separate pairwise

competition experiment, in which we quantified fitness advantages of MSHA-mutants

relative to each Φ-carrier (Figure S4). The fitness benefit of MSHA-mutants was higher

against bacteria carrying the high virulence phage compared to bacteria carrying the low

virulence phage (significant treatment term in linear model with treatment, GFP-label

and the interaction thereof as fixed factors: $F_{1,8}=18.63$, $p=0.003$, Table S2). These

results support the dynamics observed in the selection experiment, (i.e., VALGΦ8$_{K04M5}$-

carriers went extinct significantly faster than VALGΦ8$_{K04M1}$-carriers) and suggests

stronger selection for MSHA-mutants when coevolving with high virulence phage.

Bacterial population densities during the selection experiment were negatively

correlated with the number of Φ-carriers per population (Pearson's correlation without

zero inflation Φ-K04M1: $r=0.69$, $t_{21}=-4.38$, $p<0.001$, Φ-K04M5: $r=0.92$, $t_7=-6.29$,

$p<0.001$; Figure S5). This implies that, even though the majority of the clones in the

populations were resistant to infection, bacterial populations were not able to recover as

long as the dominating mechanism of phage-resistance was superinfection exclusion,

presumably due to virulence. To further quantify the parameters that influence the

fitness of MSHA-mutants relative to Φ-carriers, we measured the amount of free phages

and the absolute growth rate of selected clones. Specifically, we quantified differences

in phage production and tested if phage production impairs bacterial growth. We

observed an increase in phage production relative to the ancestor in Φ-carriers isolated

from populations coevolved with the high virulence phage (paired *t*-test: t_5=-4.31,

p=0.008) but not those coevolved with the low virulence phage (non-significant paired

t-test: $t_{4.2}$=-1.18, p=0.3, Figure S6a). Also, VALGΦ8$_{K04M5}$-carriers produced

significantly more free phages than VALGΦ8$_{K04M1}$-carriers (paired *t*-test: $t_{5.61}$=-3.36,

p=0.017).

Direct comparisons of growth rates among evolved clones revealed a significant

difference between both resistance forms with Φ-carriers growing on average slower

than MSHA-mutants (VALGΦ8$_{K04M5}$: paired *t*-test: $t_{6.61}$=-3.39, p=0.006;

VALGΦ8$_{K04M1}$: paired *t*-test: $t_{7.5}$=-3.32, p=0.01, Figure S6b). We further observed that

phage production significantly impaired bacterial growth (significant negative

correlation between the amount of produced phages and bacterial growth rate Figure 3).

Taken together these data suggest that superinfection exclusion buys time, offering

resistance to subsequent infection, but at the cost of suffering the virulence of being

infected; this resistance mechanism is replaced by surface receptor modification in

populations coevolving with both the high and the low virulence phage, but that

selection for resistance mutations is stronger with higher virulence.

Figure 3 Correlation between bacterial growth rate [μ] and production of free phages measured as

PFU/ml per clone (left VALGΦ8$_{K04M1}$, right: VALGΦ8$_{K04M5}$).

DISCUSSION

We hypothesised that selection for resistance evolution in viral infections will be

stronger for high virulent compared to low virulent viral infections. Resistance against

filamentous phages can emerge via two ways: (1) super-infection exclusion mediated by

the presence of the infecting phage in the bacterial host cell, or (2) molecular resistance

evolution, for instance surface-receptor modifications, which prevent phages from

entering host cells. Here, we show that upon phage infection, the first mechanism of

resistance acquisition was by super-infection exclusion, which occurred at a similar rate

in both treatments. However, over time, these phage-carrying clones were rapidly

replaced by phage resistant mutants. This replacement happened significantly faster in

high virulent compared to low virulent infections, suggesting that selection for

molecular resistance evolution is stronger in high virulent infection. Ultimately this

faster resistance evolution caused a faster extinction of phages, which were not able to

co-evolve in order to overcome evolved host-resistance, and a shorter epidemic.

At least two mutually non-exclusive scenarios can explain why mutants became the

dominant form in our experiments: (1) cost of resistance, and (2) stability of resistance.

While resistance acquisition is faster via super-infection exclusion than by molecular

changes, our data suggest that this process is less stable. That is because in episomal,

non-integrative phage infections some bacterial daughter cells will not inherit the

infecting phage due to spontaneous segregation-loss, rendering them once again

susceptible to subsequent infections. In contrast, molecular resistance evolution is more

stable and due to their higher fitness, those resistant clones can rapidly sweep through

the population and reach fixation.

The rapid extinction of phages and phage-carriers in our experiment raises the

questions, why filamentous phages are omnipresent in nature (Roux, Krupovic et al.

2019), including in environmental Vibrio strains closely related to our model strain

K01M1 (Chibani, Hertel et al. 2020). In other words, what selects for the presence of

filamentous phage-carriers in nature? In some cases filamentous phages can contribute

positively to the host's evolutionary fitness (Hay and Lithgow 2019) through lysogenic

conversion, i.e., by providing the bacterial host with new phenotypes induced by phage-

encoded proteins (Waldor and Mekalanos 1996). In certain environments this may be

beneficial enough to outweigh the costs, resulting in a successful chronic infection.

Among those, the most prominent example is CTXφ, which carries the genes for the

cholera toxin (Waldor and Mekalanos 1996). For the phage from our study, however,

we can only speculate, that its presence provides a fitness advantage in the natural

environment, which we did not capture in our laboratory experiments. These might

include for instance advantages during colonization (Davis and Waldor 2003), or

increased stress tolerance (Yu, Chen et al. 2015). The annotation and comparative

genomic analyses of VALGΦ8 based on the available information in today's databases,

does however not reveal any known-accessory traits that would support this hypothesis

(Chibani, Hertel et al. 2020). Answers might come from co-evolution experiments in

more natural environments, for instance inside the gut of marine animals. Such results

could then be used to predict the chances of establishing a persistent chronic infection

which we assume is likely to depend on ecological and evolutionary factors that

determine the net-benefit of the phage on its host (Bull, Molineux et al. 1991, Shapiro,

Williams et al. 2016, Shapiro and Turner 2018).

Ecological benefits of filamentous phages can arise during species interactions, e.g.,

by protecting hosts against infection by other phages (superinfection exclusion), by

acting as decoys for mammalian immunity (Sweere, Van Belleghem et al. 2019), or

from interactions with the abiotic environment e.g., through stress tolerance (Yu, Chen

et al. 2015). Evolutionary benefits include increased mutation supply, and lysogenic

conversion (Waldor and Mekalanos 1996). However, filamentous phages can also be

costly, for instance thorugh disruption of functional host genes following chromosomal

insertion or a fitness loss due to increased transcriptional load, in particular for phages

that reproduce at high frequencies. We predict that the net-benefit of a filamentous

phage on its host determines the co-evolutionary outcome of both species. If costs

associated with phage carriage exceed the benefits, we expect that phages will be

quickly lost from the population as we have seen in the present experiment. By contrast,

if phages increase their host's fitness we expect successful chronic infections.

Conclusion

Consistent with our hypothesis we could show that selection for resistance

evolution is stronger against high virulence compared to low virulence viral infections.

Accordingly, high virulence infections may be cleared faster leading to shorter

epidemics which may additionally limit the transmission success of highly virulent

viruses. However, since the virus in our study failed to co-evolve, this system should

not be used as a general model to study resistance evolution. Other outcomes in terms of

resistance evolution are to be expected, if viruses co-evolve with their hosts. It might be

possible that co-evolution of high virulent viruses is constrained as they have much less

time to acquire the necessary mutations that would allow them to overcome host

resistance. If this holds true, natural selection might favour low virulent variants, that

are able to persist longer inside an organism which not only increases their transmission

potential but also their time to co-evolve. Future work will focus on the impact of

virulence on the evolutionary potential during host-virus co-evolution.

Acknowledgements: We thank Katja Trübenbach, Veronique Merten, Silke-Mareike

Merten and Kim-Sara Wagner for their support in the laboratory.

Funding: This project was funded by two grants from the DFG [WE 5822/ 1-1], [WE

5822/ 1-2] within the priority programme SPP1819 given to CCW and OR and a DFG

grant within the Cluster of Excellence 80 "The Future Ocean" given to CCW.

References

Alizon, S., A. Hurford, N. Mideo and M. Van Baalen (2009). "Virulence evolution and
the trade-off hypothesis: history, current state of affairs and the future." J Evol Biol
22(2): 245-259.

Anderson, R. M. and R. M. May (1982). "Coevolution of hosts and parasites."
Parasitology **85 (Pt 2)**: 411-426.

Bull, J. J., I. J. Molineux and W. R. Rice (1991). "Selection of Benevolence in a Host-
Parasite System." Evolution **45**(4): 875-882.

Chibani, C. M., R. Hertel, M. Hoppert, H. Liesegang and C. C. Wendling (2020).
"Closely Related Vibrio alginolyticus Strains Encode an Identical Repertoire of
Caudovirales-Like Regions and Filamentous Phages." Viruses **12**(12).

Chibani, C. M., O. Roth, H. Liesegang and C. C. Wendling (2020). "Genomic
variation among closely related Vibrio alginolyticus strains is located on mobile
genetic elements." BMC Genomics **21**(1): 354.

Chin, C. S., D. H. Alexander, P. Marks, A. A. Klammer, J. Drake, C. Heiner, A. Clum,
A. Copeland, J. Huddleston, E. E. Eichler, S. W. Turner and J. Korlach (2013).
"Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing
data." Nat Methods **10**(6): 563-569.

Davis, B. M. and M. K. Waldor (2003). "Filamentous phages linked to virulence of
Vibrio cholerae." Curr Opin Microbiol **6**(1): 35-42.

Ebert, D. (1994). "Virulence and Local Adaptation of a Horizontally Transmitted
Parasite." Science **265**(5175): 1084-1086.

Fenner, F. and I. D. Marshall (1957). "A comparison of the virulence for European
rabbits (Oryctolagus cuniculus) of strains of myxoma virus recovered in the field in
Australia, Europe and America." J Hyg (Lond) **55**(2): 149-191.

Goehlich, H., O. Roth and C. C. Wendling (2019). "Filamentous phages reduce
bacterial growth in low salinities." R Soc Open Sci **6**(12): 191669.

Harrison, E., D. Guymer, A. J. Spiers, S. Paterson and M. A. Brockhurst (2015).
"Parallel Compensatory Evolution Stabilizes Plasmids across the Parasitism-
Mutualism Continuum." Current Biology **25**(15): 2034-2039.

Herre, E. A. (1993). "Population structure and the evolution of virulence in nematode
parasites of fig wasps." Science **259**(5100): 1442-1445.

Jouravleva, E. A., G. A. McDonald, J. W. Marsh, R. K. Taylor, M. Boesman-Finkelstein and R. A. Finkelstein (1998). "The Vibrio cholerae mannose-sensitive hemagglutinin is the receptor for a filamentous bacteriophage from V. cholerae O139." Infect Immun **66**(6): 2535-2539.

Lenski, R. E., M. R. Rose, S. C. Simpson and S. C. Tadler (1991). "Long-Term Experimental Evolution in Escherichia-Coli .1. Adaptation and Divergence during 2,000 Generations." American Naturalist **138**(6): 1315-1341.

Mai-Prochnow, A., J. G. Hui, S. Kjelleberg, J. Rakonjac, D. McDougald and S. A. Rice (2015). "'Big things in small packages: the genetics of filamentous phage and effects on fitness of their host'." FEMS Microbiol Rev.

May, R. M. and R. M. Anderson (1983). "Epidemiology and genetics in the coevolution of parasites and hosts." Proc R Soc Lond B Biol Sci **219**(1216): 281-313.

Messenger, S. L., I. J. Molineux and J. J. Bull (1999). "Virulence evolution in a virus obeys a trade-off." Proc Biol Sci **266**(1417): 397-404.

Poullain, V., S. Gandon, M. A. Brockhurst, A. Buckling and M. E. Hochberg (2008). "The evolution of specificity in evolving and coevolving antagonistic interactions between a bacteria and its phage." Evolution **62**(1): 1-11.

Rakonjac, J. (2012). Filamentous Bacteriophages: Biology and Applications eLS. Ltd: Chichester, John Wiley & Sons.

Roux, S., M. Krupovic, R. A. Daly, A. L. Borges, S. Nayfach, F. Schulz, A. Sharrar, P. B. Matheus Carnevali, J. F. Cheng, N. N. Ivanova, J. Bondy-Denomy, K. C. Wrighton, T. Woyke, A. Visel, N. C. Kyrpides and E. A. Eloe-Fadrosh (2019). "Cryptic inoviruses revealed as pervasive in bacteria and archaea across Earth's biomes." Nat Microbiol **4**(11): 1895-1906.

Shapiro, J. W. and P. E. Turner (2018). "Evolution of mutualism from parasitism in experimental virus populations." Evolution **72**(3): 707-712.

Shapiro, J. W., E. S. C. P. Williams and P. E. Turner (2016). "Evolution of parasitism and mutualism between filamentous phage M13 and Escherichia coli." Peerj **4**.

Sweere, J. M., J. D. Van Belleghem, H. Ishak, M. S. Bach, M. Popescu, V. Sunkari, G. Kaber, R. Manasherob, G. A. Suh, X. Cao, C. R. de Vries, D. N. Lam, P. L. Marshall, M. Birukova, E. Katznelson, D. V. Lazzareschi, S. Balaji, S. G. Keswani, T. R. Hawn, P. R. Secor and P. L. Bollyky (2019). "Bacteriophage trigger antiviral immunity and prevent clearance of bacterial infection." Science **363**(6434): 1416-+.

Team, R. D. C. (2011). " R: A language and environment for statistical computing. R Foundation for Statistical Computing." Vienna, Austria. ISBN 3-900051-07-0, URL http://www.R-project.org/. .

Turner, P. E., V. S. Cooper and R. E. Lenski (1998). "Tradeoff between Horizontal
and Vertical Modes of Transmission in Bacterial Plasmids." Evolution **52**(2): 315-329.

Waldor, M. K. and J. J. Mekalanos (1996). "Lysogenic conversion by a filamentous
phage encoding cholera toxin." Science **272**(5270): 1910-1914.

Weiss, S. R. and J. L. Leibowitz (2011). "Chapter 4 - Coronavirus Pathogenesis."
Advances in Virus Research **81**: 85-164.

Wendling, C. C., H. Goehlich and O. Roth (2018). "The structure of temperate phage-
bacteria infection networks changes with the phylogenetic distance of the host
bacteria." Biol Lett **14**(11).

Wendling, C. C., A. Piecyk, D. Refardt, C. Chibani, R. Hertel, H. Liesegang, B. Bunk,
J. Overmann and O. Roth (2017). "Tripartite species interaction: eukaryotic hosts
suffer more from phage susceptible than from phage resistant bacteria." BMC Evol
Biol **17**(98).

Yu, Z. C., X. L. Chen, Q. T. Shen, D. L. Zhao, B. L. Tang, H. N. Su, Z. Y. Wu, Q. L.
Qin, B. B. Xie, X. Y. Zhang, Y. Yu, B. C. Zhou, B. Chen and Y. Z. Zhang (2015).
"Filamentous phages prevalent in Pseudoalteromonas spp. confer properties
advantageous to host survival in Arctic sea ice." ISME J **9**(4): 871-881.
673