The temperature-regulated DEAD-box RNA helicase CrhR interactome:

Autoregulation and photosynthesis-related transcripts

Anzhela Migur[1,§], Florian Heyl[2], Janina Fuss[3,$], Afshan Srikumar[4], Bruno Huettel[3], Claudia

Steglich[1], Jogadhenu S. S. Prakash[4], Richard Reinhardt[3], Rolf Backofen[2], George W.

Owttrim[5], Wolfgang R. Hess[1*]

[1]Faculty of Biology, University of Freiburg, Schänzlestr. 1, D-79104 Freiburg, Germany

[2]Department of Computer Science, University of Freiburg, Georges-Koehler-Allee 106, D-

79110 Freiburg, Germany

[3]Max Planck-Genome-Centre Cologne, Carl-von-Linné-Weg 10, D-50829 Köln, Germany

[4]Department of Biotechnology & Bioinformatics, School of Life Sciences, University of

Hyderabad, Hyderabad, India

[5]Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada

T6G 2E9

Running title: CrhR DEAD-box RNA helicase interactome

Corresponding author: Wolfgang R. Hess; Phone: 0049-761-2032796; Fax: 0049-761-

2032745; Email: wolfgang.hess@biologie.uni-freiburg.de

§Current address: RNA Synthetic Biology Group, Helmholtz Institute for RNA-based

Infection Research (HIRI), Würzburg, Germany

$Current address: Institute of Clinical Molecular Biology, University of Kiel, Kiel, Germany

Email addresses:

anzhela.migur@helmholtz-hiri.de

j.fuss@ikmb.uni-kiel.de

reinhardt@mpipz.mpg.de

huettel@mpipz.mpg.de

claudia.steglich@biologie.uni-freiburg.de

backofen@informatik.uni-freiburg.de

heylf@informatik.uni-freiburg.de

syamsunderp@yahoo.com

afshan.sri17@gmail.com

gowttrim@ualberta.ca

wolfgang.hess@biologie.uni-freiburg.de

Date of submission: March 25, 2021

Number of tables and figures: 6

Total Word count: ~11,000

Word count from the start of the introduction to the end of the acknowledgements -
excluding materials and methods, references, tables and figure legends: 4698

Supplementary data: 1 figure, 5 tables

Conflicts of Interest: The authors declare the absence of conflicts of interest.

Highlight

The cyanobacterial DEAD-box RNA helicase CrhR binds mainly photosynthesis-associated and redox-controlled transcripts connecting its regulation, localization and phenotypes of mutants for the first time with a set of potential RNA targets.

Abstract

RNA helicases play crucial functions in RNA biology. In plants, RNA helicases are encoded by large gene families, performing roles in abiotic stress responses, development, the post-transcriptional regulation of gene expression as well as house-keeping functions. Several of these RNA helicases are targeted to the organelles, mitochondria and chloroplasts. Cyanobacteria are the direct evolutionary ancestors of plant chloroplasts. The cyanobacterium *Synechocystis* 6803 encodes a single DEAD-box RNA helicase, CrhR, that is induced by a range of abiotic stresses, including low temperature. Though the Δ*crhR* mutant exhibits a severe cold-sensitive phenotype, the physiological function(s) performed by CrhR have not been described. To identify transcripts interacting with CrhR, we performed RNA co-immunoprecipitation with extracts from a *Synechocystis crhR* deletion mutant expressing the FLAG-tagged native CrhR or a K57A mutated version with an anticipated enhanced RNA binding. The composition of the interactome was strikingly biased towards photosynthesis-associated and redox-controlled transcripts. A transcript highly enriched in all experiments was the *crhR* mRNA, suggesting an auto-regulatory molecular mechanism. The identified interactome explains the described physiological role of CrhR in response to the redox poise of the photosynthetic electron transport chain and characterizes CrhR as an enzyme with a diverse range of transcripts as molecular targets.

Keywords: chloroplasts, CrhR RNA helicase, cyanobacteria, gene expression regulation, photosynthesis, small regulatory RNA, redox regulation, RNA:RNA interaction

76 **Abbreviations:** co-IP, co-immunoprecipitation; CrhR, cyanobacterial RNA helicase

77 redox; ETC, electron transport chain, HLR1, high light regulatory 1; OCP, orange

78 carotenoid protein; sRNA, small regulatory RNA; TSS, transcriptional start site

Introduction

DEAD-box RNA helicases

The synthesis, maturation, modification and decay of RNA molecules and their interaction with each other and with other cellular components is central to the molecular basis of life. The largest family of enzymes involved in the metabolism of RNA molecules are enzymes belonging to the Superfamily I (SF1) and II (SF2) RNA helicases (Bourgeois *et al.*, 2016). DEAD-box RNA helicases, named after the conserved DEAD (Asp-Glu-Ala-Asp) amino acid motif in their core motif, form the largest and most complex group of SF2 RNA helicases (Jarmoskaite and Russell, 2011). DEAD-box RNA helicases are RNA-dependent ATPases (Rocak and Linder, 2004) characterized by the presence of twelve highly conserved motifs (Redder *et al.*, 2015). These motifs form the motor core of the helicase, which binds specifically adenine nucleotides and single-stranded RNA in a sequence-independent manner. The main function of these proteins is the conformational rearrangement of RNA by unwinding short double stranded regions. The unwinding reaction can be performed within one RNA molecule or between two duplex-forming RNAs. Besides the unwinding reaction, some RNA helicases are able to anneal single stranded RNAs (Chamot *et al.*, 2005; Yang and Jankowsky, 2005). In addition, some RNA helicases act as RNA clamps or facilitate RNA-protein complex dissociation without duplex unwinding (Jankowsky *et al.*, 2001; Fairman *et al.*, 2004).

Abiotic stresses and DEAD-box RNA helicases in plants and cyanobacteria

Cold stress, one of the most common stress conditions in nature, frequently induces the expression or activity of DEAD-box RNA helicases. At low temperature, RNA secondary structures are thermodynamically stabilized, which may interfere with their function. RNA helicases can rescue the RNA's functions by rearranging its secondary structures (Jones *et al.*, 1996). This role makes certain RNA helicases more relevant or even conditionally essential at low temperature. For instance, deletion of *csdA* or *srmB* in *Escherichia coli* leads to a cold-sensitive phenotype (Redder *et al.*, 2015). If all four DEAD-box RNA

helicase genes are deleted, *Bacillus subtilis* is not viable at low temperature (16°C),

although the strain grows well at 37°C (Lehnik-Habrink *et al.*, 2013).

Most organisms encode several DEAD-box RNA helicases possessing non-

complementary roles associated with a variety of physiological functions. Common

functions in all organisms involve ribosome biogenesis, RNA turnover and translation, and

responses to multiple stress conditions (Py *et al.*, 1996; de la Cruz *et al.*, 1999; Schneider

and Schwer, 2001; Rogers *et al.*, 2002; Macovei *et al.*, 2012). Plant genomes typically

possess larger and more diverse RNA helicase gene families than observed in other

systems (Linder and Owttrim, 2009). Hence, their relevance in RNA secondary structure

rearrangement under different environmental conditions or in plant development presents

interesting lines for investigation. For example, *Arabidopsis thaliana* encodes 58 DEAD-

box RNA helicases (Boudet *et al.*, 2001), many of which are essential as they are not

functionally complementary (Mingam *et al.*, 2004). RNA helicases in plants fulfil roles in

the defense against viruses (Wu and Nagy, 2020), abiotic stress responses to high salinity

(Capel *et al.*, 2020), low temperature (Lu *et al.*, 2020; Wang *et al.*, 2020), and in

development (Gong *et al.*, 2005). Interestingly, although not complementary, some plant

helicases are involved in the same processes, suggesting multiple, independent RNA

structure rearrangements are associated with a single physiological response (Huang *et*

al., 2016). In addition, several RNA helicases are targeted to the organelles, mitochondria

and chloroplasts (Matthes *et al.*, 2007; Nawaz and Kang, 2017; Nawaz *et al.*, 2018).

However, the expanded DEAD-box RNA helicase families make the precise functional

characterization of RNA helicases challenging in plants.

Cyanobacteria are the direct evolutionary ancestors of plant chloroplasts

(Mereschkowsky, 1905; Margulis, 1981; Martin and Kowallik, 1999; Ponce-Toledo *et al.*,

2017). The endosymbiosis of a cyanobacterium not only led to the chloroplast but also

had pivotal impact on the composition of the plant nuclear genome (Martin *et al.*, 2002).

Therefore, analysis of gene function in cyanobacteria is also informative at the higher plant

level.

The cyanobacterium *Synechocystis* encodes a single DEAD-box RNA helicase, CrhR

The unicellular cyanobacterium *Synechocystis* sp. PCC 6803 (from here: *Synechocystis*) encodes the DEAD-box RNA helicase CrhR, for cyanobacterial RNA helicase redox (Rosana *et al.*, 2012*a*). As defined by a 50 amino acids sequence motif, CrhR is the archetype protein of a new clade within the DEAD-box RNA helicase family (Whitford *et al.*, 2021). In contrast to the situation in plants and also in many other bacteria, CrhR (*crhR/slr0083*) is the only DEAD-box RNA helicase encoded in *Synechocystis* (Redder *et al.*, 2015). Functionally, CrhR can therefore be expected to be of particular importance as it likely performs multiple functions that from an evolutionary perspective could have been distributed to different members of the complex family of RNA helicases that we observe in plants today (Kiefer *et al.*, 2020). *crhR (slr0083)* was originally characterized as a salt and cold-shock inducible protein (Vinnemeier and Hagemann, 1999; Kujat and Owttrim, 2000). Cold stress-inducible RNA helicases were also investigated in the cyanobacterium *Anabaena* sp. PCC 7120 (Chamot *et al.*, 1999) and *Synechococcus* sp. WH 7803 (Gierga *et al.*, 2012). Detailed analysis indicated that a low temperature stabilization of both transcript and protein contribute to the observed low temperature induction of CrhR (Rosana *et al.*, 2012*a*) and that a *crhR* inactivation mutant was severely impaired morphologically and physiologically at lower but not at higher temperatures (Rosana *et al.*, 2012*b*). In addition to low temperature, *crhR* is induced by a range of abiotic stresses that reduce the electron transport chain, independent of temperature shift (Vinnemeier and Hagemann, 1999); Kujat and Owttrim, 2000; Ritter et al., 2020). The gene *crhR* (*slr0083*) is located in a dicistronic operon together with *rimO* (*slr0082*), whose putative protein product has 38% identity with RimO, a ribosomal protein S12 methylthiotransferase (UniProtKB P0AEI4), and 29% identity with the paralogous tRNA methylthiolase MiaB (Uni-ProtKB P0AEI1) from *E. coli* (Rosana *et al.*, 2020). The auto-regulated enhanced operon discoordination and processing of the *crhR* mRNA from the dicistronic operon RNA adds further complexity to its regulation (Rosana *et al.*, 2012; Rosana *et al.*, 2020). In addition, upon temperature upshift to non-permissive conditions, CrhR undergoes rapid repression via conditional proteolysis at the post-translational level

through an unknown mechanism (Tarassova *et al.*, 2014). In localization studies, CrhR

localized to the thylakoid membranes and co-sedimented with degradosome and

polysome complexes (Rosana *et al.*, 2016), consistent with the recent co-fractionation

analysis using Grad-seq (Riediger *et al.*, 2021).

Although the molecular effects of *crhR* deletion or inactivation were studied at both the

transcriptome (Prakash *et al.*, 2010; Georg *et al.*, 2019) and proteome level (Rowland *et*

al., 2011), the direct RNA targets of CrhR and interacting protein partners have not been

identified. To monitor transcriptome-wide binding of CrhR, we immuno-precipitated the

RNA species interacting with a FLAG-tagged version of the native RNA helicase

expressed in a $\Delta crhR$ background (subsequently called CrhR$_{WT}$). We performed UV cross-

linking *in vivo* with CrhR$_{WT}$ cultures grown at the permissive temperature of 30°C or

exposed to 20°C for 2 h (low temperature stress). A possible obstacle in the analysis of

RNA helicases can be their transient interaction with RNA molecules followed by rapid

ATP-dependent dissociation (Linder and Jankowsky, 2011). Therefore, we introduced a

K57A mutation located in the predicted Walker A ATP-binding motif I (TGTGKT) that is

conserved in CrhR (Tanner and Linder, 2001). Mutation of the conserved lysine or the last

threonine in this motif is known to interfere with the ATPase activity of DEAD-box RNA

helicases (Cordin *et al.*, 2006). Plasmid encoded K57A was conjugated into the $\Delta crhR$

background yielding strain CrhR$_{K57A}$. This strain was used for comparison applying UV

cross-linking *in vivo* at 30°C followed by co-immunoprecipitation (co-IP) and RNA

sequence analysis as for the CrhR$_{WT}$ strain.

Altogether, 98 RNA sequences were significantly enriched in at least one experiment.

Functional analysis of the specifically enriched RNAs indicated a striking preference for

CrhR interaction with transcripts associated with photosynthesis but also transcripts

associated with RNA metabolism, among them *crhR* itself. The enrichment of *crhR*–

transcripts with the tagged CrhR protein is consistent with an auto-regulatory mechanism

in which CrhR controls its own expression at the post-transcriptional level.

Broader implications were revealed by the potential RNA targets having previously been

characterized as controlled by the transcription factor RpaB (Riediger *et al.*, 2019), which

is of interest in conjunction with the known ETC redox poise regulation of *crhR* expression

(Kujat and Owttrim, 2000; Ritter *et al.*, 2020). Overall, the results connect the known
association of CrhR with both the thylakoid membrane and ribosomes and its cold-induced
and redox poise-controlled expression with a specific set of potential RNA targets
comprising a subset of the *Synechocystis* transcriptome.

Results

Physiological consequences of manipulation of CrhR sequence and abundance

To gain insight into the CrhR interactome, we performed immunoprecipitation and
isolation of tagged CrhR-RNA complexes from *in vivo* UV cross-linked *Synechocystis*
cells. A previously characterized helicase deletion mutant, Δ*crhR*, generated by
replacement of the *crhR* gene (*slr0083*) by a spectinomycin resistance gene (Prakash *et*
al., 2010), served as the genetic background in which three strains were constructed. One,
ectopically expressing the triple FLAG-tagged CrhR (CrhR$_{WT}$), and the other expressing
the triple FLAG-tagged CrhR(K57A) mutant (CrhR$_{K57A}$) from conjugative vector pVZ321
(**Fig. 1A**). The introduced K57A mutation should interfere with ATP binding and ATPase
activity of CrhR (Tanner and Linder, 2001) and therefore delay or reduce its capability to
dissociate from bound RNA. As a control, a third strain was constructed with pVZ321
carrying a short reading frame encoding a triple FLAG tag and identical promoter and
terminator sequences as the other two strains (Δ*crhR*$_{control}$). Three experiments were
performed with these strains (**Fig. 1B**). The proper regulation and expression of FLAG-
tagged CrhR proteins was confirmed by western blotting (**Fig. S1**).

We followed the growth of the three generated strains and wild type over 25 days at the
non-permissive temperature of 20°C. Growth of Δ*crhR*$_{control}$ was dramatically reduced
compared to wild type cells, while growth of CrhR$_{WT}$ was similar to the unmodified wild
type (**Fig. 2A**). Thus, ectopically transcribed, FLAG-tagged CrhR in the CrhR$_{WT}$ strain
complemented the *crhR* deletion at this temperature. Interestingly, complementation in
strain CrhR$_{K57A}$ led to a partial restoration of the cold-sensitive phenotype, i.e. growth of
CrhR$_{K57A}$ was reduced compared to the wild type but not to the same extent as of
Δ*crhR*$_{control}$ (**Fig. 2A**). In addition, a strong reduction in pigment content in CrhR$_{K57A}$ and in

Δ*crhR*_{control} was apparent compared to the other two strains (**Fig. 2B, C**). These variations

were congruent with previously identified changes in light-harvesting pigment composition

upon *crhR* inactivation (Rosana *et al.*, 2012*b*). We concluded that the FLAG tag did not

interfere with the enzymatic and regulatory functions of CrhR and therefore these lines

were suitable for more detailed analyses.

RNA co-immunoprecipitations

The RNA helicase CrhR performs unwinding and annealing of RNA (Chamot *et al.*, 2005),

reactions for which mainly transient interactions with RNA can be assumed including

binding to RNA followed by rapid dissociation. This makes the experimental mapping of

an RNA helicase interactome more challenging than for an RNA-binding protein.

Therefore, we performed covalent crosslinking *in vivo* to enrich RNA-CrhR complexes

sufficient for RNA and library preparation. $CrhR_{WT}$ and $CrhR_{K57A}$ cultures grown at the

standard growth temperature of 30°C were irradiated with 254 nm UV-C light at 500

mJ/cm^2 (Holmqvist *et al.*, 2016). To identify potentially different interacting partners, the

experiment was also performed on $CrhR_{WT}$ cultures exposed to cold stress at 20°C for 2

hours. In parallel, total transcriptome data were generated for all strains and conditions

(see overview in **Fig. 1B**). The total and mapped read numbers are listed in

Supplementary Material, Table S1. CrhR interacting RNAs were predicted with the peak

calling algorithm PEAKachu (Holmqvist *et al.*, 2016). In total, we obtained 98 different

peaks with statistically significant enrichment (padj < 0.05, log$_2$FC > 0) in the three

experiments corresponding to 64 different genes (**Supplementary Material, Tables S2**

to S4). CrhR interacting RNAs were most abundant in experiment 3 using $CrhR_{K57A}$ as

bait (75 peaks), while 28 and 16 peaks were identified in the experiments 1 (20°C) and 2

(30°C) using $CrhR_{WT}$. Hence, the assumption was confirmed that the K57A substitution

would stabilize CrhR-RNA interactions.

Interestingly, the co-IPs frequently identified only specific regions of RNA transcripts which

could provide insight into both the molecular mechanism and also the functional roles

CrhR is performing. Typical patterns of enriched transcripts are visualized in **Fig. 3**. The

predominant detected interactions were associated with segments containing the 5'-
untranslated region (5' UTR), start codon and a few codons into the reading frame (**Fig.**
3A). A related pattern also included the 5' UTR, start codon and initial codons but actually
matched the entire intergenic space between two genes in an operon (**Fig. 3B**). A
strikingly different pattern was observed in which multiple sharp peaks indicated a
multitude of interactions along the length of the dicistronic operon, as illustrated for the
psaA-psaB operon (**Fig. 3C**). Nevertheless, as observed for other target transcripts, the
most pronounced interactions corresponded to the 5' UTR, start codon and initial codons
of *psaA*, the first gene of this operon (**Fig. 3C**).

Transcripts encoding photosynthesis-associated proteins or with predicted roles
in redox regulation are enhanced in the CrhR interactome

Functional analysis over the complete dataset using GO terms (Ashburner *et al.*, 2000;
Gene Ontology Consortium, 2021) identified "photosynthetic membrane" (q_{adj} = 1.18E-
04), "thylakoid" (q_{adj} = 2.74E-04) and "photosystem" (q_{adj} = 2.74E-04) as the dominating
cellular components encoded by the enriched transcripts.

Accordingly, detailed analysis indicated that most of the enriched transcripts were
associated with photosynthesis (**Table 1;** for the exact coordinates of recovered
transcripts, enrichment factors and adjusted p values in each of the experiments, see
Supplementary Material, Tables S2 to S4). Several of these transcripts encode protein
components of the two photosystems, PSI and PSII. PSI proteins included *psaL/slr1655*
for PSI subunit XI, *slr1834/psaA* and *slr1835/psaB* encoding the P700 apoprotein subunits
Ia and A2 **(Fig. 3C)**, *ssr2831/psaE* encoding the PSI reaction center subunit IV.
Transcripts encoding PSII subunits included, *slr0906/psbB* for the CP47 reaction center
protein, *slr0927/psbD*, *slr1311/psbA2* and *smr0009* for the reaction center proteins D2,
D1 and PsbN.

Several enriched transcripts encoding proteins that are not directly components of PSI or
PSII but that are associated with photosynthesis were also detected. Transcripts in this
category included *slr0228/ftsH2* encoding the FtsH2 protease crucial for the replacement

of photodamaged D1 protein during the PSII repair cycle (Silva *et al.*, 2003; Komenda *et*
al., 2010), *ssl0020/petF* encoding the major plant-like ferredoxin Fed1 (Mazouni *et al.*,
2003)*, sll0199/petE* encoding plastocyanin and *ssr2016* encoding the *Synechocystis* Pgr5
(proton gradient regulation 5) homolog. In plants and cyanobacteria, Pgr5 contributes to
cyclic electron flow from PSI to the plastoquinone pool (Yeremenko *et al.*, 2005; Dann and
Leister, 2019), electron transfer that is associated with *crhR* expression (Kujat and
Owttrim, 2000; Ritter *et al.*, 2020). Another gene in this category was *trxA* (*slr0623*),
encoding thioredoxin A, the most abundant of the four thioredoxins in *Synechocystis*
(Hishiya *et al.*, 2008), and suggests CrhR is involved in maintaining intracellular redox
status by regulating protein expression associated with scavenging reactive oxygen
species (ROS).

Light harvesting is an integral aspect of photosynthesis, the primary contributor to which
are phycobilisomes in cyanobacteria. Among several mRNAs encoding phycobilisome
(PBS) proteins in the dataset (**Table 1**) was *apcC*/*ssr3383* encoding the PBS 7.8 kDa
linker polypeptide. The interacting RNA sequence encompassed 146 nt of the 5' UTR and
54 of the 67 *apcC* codons **(Fig. 4)**, importantly ending at the start of the region from which
the regulatory sRNA ApcZ originates. ApcZ targets the *ocp*/*slr1963* transcript encoding
the water-soluble orange carotenoid protein OCP (Zhan *et al.*, 2021). Again, interaction
of this transcript with CrhR is associated with redox homeostasis since OCP functions to
quench excess excitation energy absorbed by phycobilisomes, and also directly
scavenges singlet oxygen ROS ($^1O_2$) (Kerfeld *et al.*, 2003). Downstream of the
photosynthetic electron transport chain (ETC), CrhR interaction with the *slr0148* transcript
was detected (**Table 1**). The gene *slr0148* encodes the ferredoxin *fed5* (Angeleri *et al.*,
2018), homologs of which are widely distributed in cyanobacteria and plants (Cassier-
Chauvat and Chauvat, 2014). The Fed2 - Fed9 gene family functions to distribute
electrons to a range of metabolic pathways each of which are associated with
cyanobacterial tolerance to a different but overlapping range of environmental stresses
(Cassier-Chauvat and Chauvat, 2014). In addition, *slr0148* is the terminal gene of 5
consecutive genes *slr0144* to *slr0148,* all of whose mRNA fragments were enriched by
co-IP with CrhR$_{K57A}$ (**Table S4**).

In addition, we identified CrhR-interacting transcripts of genes more distantly related to
photosynthesis, among them *slr0374* **(Fig. 3C)***,* encoding the stress-responsive AAA+
protease protein Ycf46 (Singh and Sherman, 2002). Homologs of Ycf46 are highly
conserved in all cyanobacterial lineages and most algal chloroplast genomes. Ycf46
proteins are involved in the regulation of CO_2 utilization in photosynthesis but the precise
function is unknown (Jiang *et al.*, 2015). Transcript peaks were also recovered for *slr1674*,
encoding a DUF760-containing protein described to be associated with the thermal
acclimation of PSII (Rowland *et al.*, 2010).

Transcripts encoding proteins involved in housekeeping functions and RNA
metabolism as potential CrhR targets

Another category of transcripts enriched in the CrhR co-IPs were related to house-keeping
functions of the cell. Examples included *groEL1/slr2076* and *groEL2/sll0416* encoding the
60 kDa chaperonins 1 and 2 that match previous reports on the CrhR-dependent
upregulation of *groEL1* and *groEL2* in *Synechocystis* (Prakash *et al.*, 2010). Additionally,
transcripts encoding the ribosomal proteins, *rplU/slr1678* and *rpl26/ssr1604* for the 50S
ribosomal proteins L21 and L28, and *rps14/slr0628* and *rps1a/slr1356* for the 30S
ribosomal proteins S14 and S1a, belonging to four separate operons, were detected. Of
specific interest in relation to *crhR* expression, a transcript enriched specifically from cold
stressed cultures was *desC/sll0541* encoding the Δ9 fatty acid desaturase DesC was
identified as a potential CrhR target.

A regulatory gene of interest is *sigG/slr1545* encoding the Type 3 alternative sigma factor
SigG (Huckauf *et al.*, 2000), especially because it was recovered in all three experiments
suggesting a stable interaction with CrhR (**Fig. 3A**). SigG is associated with the oxidative
stress response caused by high light (Huckauf *et al.*, 2000) and is ectopically upregulated
in a Δ*rpaA* mutant in the dark (Köbler *et al.*, 2018). Furthermore, interacting transcripts
encoding enzymes associated with RNA turnover and metabolism were identified,
including *rnj/slr0551* encoding ribonuclease J, an enzyme possessing both endonuclease

and a robust 5'-exoribonuclease activity (Cavaiuolo *et al.*, 2020), *slr0053* encoding the

rRNA maturation RNase YbeY and, as an enzyme of RNA metabolism, *crhR* itself.

The *crhR* mRNA is a target for auto-regulation through CrhR

The *crhR* transcript was consistently enriched in all three *crhR* lines, irrespective of

temperature (**Fig. 5**). The enriched peak regions within the *crhR* transcript identify three

potential regions of preferred interaction: the major sequence overlaps the translational

start codon and extends 216 to 270 nt into the coding region, a second region from

approximately codon 125 to 219 and a third extending from approximately codon 350 to

450 of the 492 codon ORF.

To confirm CrhR-*crhR* mRNA interaction *in vitro,* 150 nt of the *crhR* transcript,

corresponding to the primary interacting region, was tested in an electrophoretic mobility

shift assay (EMSA) using His-tagged recombinant CrhR produced in *E. coli* (**Fig. 5B**).

Enhanced formation of a prominent retarded protein-RNA complex was directly

proportional to CrhR concentration. We conclude that the *crhR* mRNA is a preferred target

of CrhR and thus an auto-regulatory effect appears likely.

Discussion

Involvement of CrhR in a CrhR-dependent auto-regulatory negative feedback loop

The DEAD-box RNA helicase CrhR is one of the major cold shock proteins of

Synechocystis (Kujat and Owttrim, 2000; Prakash *et al.*, 2010; Rowland *et al.*, 2011;

Rosana *et al.*, 2012*a*,*b*; Sireesha *et al.*, 2012). Consistent with observations in many other

organisms that shift to lower temperature increases expression of DEAD-box RNA

helicases (Owttrim, 2013), we observed previously an increased *crhR* transcript

abundance after temperature downshift to 20°C, with a $\log_2$FC of 1.4 in the wild type and

2.6 for the 5' portion of the gene not deleted in a partial *crhR* inactivation mutant (Georg

et al., 2019). The higher induction in the partial *crhR* inactivation mutant pointed at a

possible feedback effect of CrhR that was active in the wild type but absent in the mutant.

Regulation of *crhR* expression is complex, involving transcriptional and post-
transcriptional effects (Rosana *et al.,* 2012*a*), and an auto-regulated operon
discoordination and processing mechanism was proposed (Rosana *et al.*, 2020).
Complicating the regulation, *crhR* is encoded in the *rimO*/*slr0082-crhR/slr0083* (*rimO-*
crhR) dicistronic operon (Rosana *et al.*, 2012*a*). However, in wild type *Synechocystis* cells,
accumulation of the full-length dicistronic transcript is essentially undetectable; *rimO* and
crhR mRNAs co-occur predominantly as monocistronic transcripts (Rosana *et al.*, 2012*a*,
2020). Interestingly, in the absence of functional CrhR, accumulation of the entire *rimO-*
crhR operon transcript is elevated compared to wild type cells (Rosana *et al.,* 2020). This
observation led to the assumption that CrhR may facilitate processing of the *rimO-crhR*
operon transcript, although CrhR is not necessary and sufficient for the processing activity
(Rosana et al., 2020). The discovery of an RNase E processing site ~138 nt upstream of
the *crhR* coding sequence then suggested that RNase E might play a role in processing
of the operon. We previously showed that the *rimO-crhR* dicistronic transcript was cleaved
by RNase E ~138 nt upstream of the *crhR* start codon *in vitro* (Rosana *et al.*, 2020).
Interestingly, both specific RNA sequence and structure were required for RNase E
substrate recognition and cleavage. Thus, we propose that CrhR may assist RNase E by
interacting with the *rimO-crhR* transcript and altering its structure to the one that favors
cleavage by RNase E. In this scenario, CrhR would auto-regulate its own expression by
enhancing processing of the *rimO-crhR* operon by RNase E. Moreover, CrhR could also
be involved in the further processing of monocistronic *crhR* (Rosana *et al.*, 2012*a*, 2020).
In support of this hypothesis, northern and microarray analysis previously indicated that
the absence of functional CrhR RNA helicase activity enhances stabilization of *crhR*
transcripts at 30 and 20°C (Georg *et al.*, 2019; Rosana *et al.*, 2020). From the degradation
constants, we concluded that at 20°C, alteration of RNA levels in the *crhR* helicase
deficient mutant could be largely explained by regulation occurring at post-transcriptional
level (Georg *et al.*, 2019).

In combination, these observations infer a CrhR-dependent negative feedback loop, in
which binding of CrhR to *crhR* transcripts leads to auto-regulation of expression.

The CrhR interactome *in vivo*

DEAD-box RNA helicases have been characterized as drivers and regulators of gene expression. For many DEAD-box RNA helicases, both the preferred targets as well as the mechanism of target recognition have remained unknown. This is relevant as the diversity of functions of DEAD-box RNA helicases and sequences of the auxiliary domains suggests that there is no single mechanism of specific target recognition or pathway association. Determination of the RNA helicase interactome is more challenging than for other RNA-interacting proteins since after binding, the helicase will rapidly modify the structure and dissociate. This potentially impedes the ability to co-IP interacting transcripts compared to RNA-binding proteins such as CsrA, Hfq or ProQ (Hör *et al.*, 2018). To mitigate this challenge, we employed the CrhR variant, CrhR$_{K57A}$, in which RNA residence time was predicted to be extended, since the Walker A ATP binding domain is inactivated.

This consideration was supported by the enhanced recovery of interacting transcripts from the CrhR$_{K57A}$ strain, i.e. experiment 3 (**Fig. 1**) suggesting that RNA binding was most efficient in the CrhR$_{K57A}$ strain. Moreover, the partial complementation of the Δ*crhR* phenotype by CrhR$_{K57A}$ (**Fig. 2**) supported the view that this variant had reduced RNA helicase enzymatic activity.

Functionally, the prime targets of CrhR interaction were consistently observed to include transcripts associated with photosynthesis, light harvesting or directly encoding core photosystem proteins (**Table 1**). A role in the biology of photosynthesis-relevant mRNAs would further be consistent with the demonstrated localization of ribosome-associated CrhR at the thylakoid membrane (Rosana *et al.*, 2016) and also the observation that *psbA* and *psaA* mRNAs are transported to the thylakoid membrane and possibly translated by thylakoid-associated ribosomes (Tyystjärvi *et al.*, 2001; Mahbub *et al.*, 2020). These findings point to a possible involvement of CrhR in physiological secondary structure rearrangements of the identified mRNAs that could influence translation on thylakoid membrane associated ribosomes.

Since *Synechocystis* encodes only a single DEAD-box RNA helicase, it is likely that CrhR operates in multiple pathways. These roles could be associated with different stresses,

independent of temperature, as observed for the *Staphylococcus aureus* helicase, CsdA

(Oun *et al.*, 2013; Khemici *et al.*, 2020). Transcripts recovered for ribosomal proteins and

ribonucleases support this idea. Indeed, a role in translation regulation is supported by

the observation that CrhR interacts with the *rps1a/slr1356* transcript encoding Rps1a. In

previous analyses, Rps1a sedimented away from the majority of ribosomal proteins

(Riediger *et al.*, 2021) and was reported to perform a role in the Shine-Dalgarno-

independent initiation of translation (Mutsuda and Sugiura, 2006; Nakagawa *et al.*, 2010),

another process which may hence be related to CrhR helicase activity.

Another intriguing example for the possible involvement of CrhR in functionally relevant

secondary structure re-arrangements was observed for *apcC/ssr3383* where the

recovered reads terminated in the region from which the regulatory sRNA ApcZ originates.

ApcZ is an sRNA, which targets the *ocp/slr1963* mRNA encoding the water-soluble

orange carotenoid protein (OCP) by inhibiting its translation (Zhan *et al.*, 2021). If

expressed, OCP directly senses light intensity and induces thermal energy dissipation

under stress conditions (Muzzopappa and Kirilovsky, 2020). Therefore, its expression is

tightly controlled and, in this context, ApcZ can inhibit or delay the production of OCP.

This regulation is also relevant under nitrogen starvation conditions, when ApcZ is induced

from a transcription start site (TSS) that is activated by the transcription factor NtcA (Zhan

et al., 2021). However, ApcZ is also detectable as a transcript under other conditions

where it derives from processing of the *apcABC* mRNA. Interestingly, the co-IP coverage

included the first 30 nt of ApcZ, matching precisely the upstream portion of the helical

structure located at the 5' end of ApcZ (**Fig. 4**). Therefore, it is tempting to speculate that

CrhR interaction and helicase activity is either required for i) ApcZ processing, ii) to open

the ApcZ structure for interaction, or possibly iii) to facilitate ApcZ-*ocp* duplex formation.

These functions are supported by the observation that the *ocp* mRNA was identified as a

potential CrhR target and the ability of CrhR to catalyze both RNA duplex unwinding and

annealing (Chamot *et al.*, 2005).

459 **Overlaps in the regulon controlled by RpaB and the set of transcripts interacting**

460 **with CrhR**

461 Functionally, it was interesting that several of the genes whose transcripts interacted with

462 CrhR were previously predicted or demonstrated to play a role in redox signaling or are

463 regulated in a redox-dependent manner. This applies first and foremost to *crhR* itself

464 (Kujat and Owttrim, 2000; Ritter *et al.*, 2020). Another is thioredoxin A, the homolog of

465 plant m-type thioredoxin and therefore sometimes also called TrxM, which plays a major

466 role in redox regulation in *Synechocystis* (Hishiya *et al.*, 2008). A protein interacting with

467 TrxM *in vitro* is the redox-responsive transcription factor RpaB (Kadowaki *et al.*, 2015).

468 RpaB recognizes the "high light regulatory 1'' (HLR1) promoter element, a pair of imperfect

469 8-nt long direct repeats (G/T)TTACA(T/A) (T/A) separated by two random nucleotides

470 (Eriksson *et al.*, 2000; Kappell and van Waasbergen, 2007). Binding of RpaB to HLR1

471 promoter motifs yields one of two different outcomes, repression or activation under low

472 light depending on the distance between the location of these elements and the TSS

473 (reviewed by Riediger *et al.*, 2018). RpaB is the response regulator (RR) of the histidine

474 kinase Hik33 in a two-component signal transduction system that controls >150 target

475 promoters in *Synechocystis*, regulating expression of diverse genes associated with

476 photosynthesis (Riediger *et al.*, 2019). Thus, there is a striking correspondence that

477 approximately two thirds of the transcripts recovered in our CrhR-co-IP analysis belong to

478 the RpaB regulon. Identification of RpaB regulon members as CrhR-interacting transcripts

479 therefore suggests a scenario in which CrhR contributes to the post-translational

480 regulation of RpaB regulon expression. Together, these observations support the function

481 of CrhR as an important post-transcriptional regulatory element within the network of

482 redox-dependent signaling and regulation.

483

484 ## Materials and methods

485 **Bacterial strains**

486 The *Synechocystis* Δ*crhR* mutant (Prakash *et al.*, 2010) was used for the construction of

487 three different strains. To establish ectopic expression of tagged CrhR, the triple FLAG-

tag was introduced at the 5'-end of the wild type *crhR*/*slr0083* gene by PCR in three steps.

First, *crhR* was amplified with the primers 3xFL-CrhR-F1 and 3xFL-CrhR-R (see **Table**

S5 for the sequences of oligonucleotide primers used in this work), re-amplified with

primers 3xFL-CrhR-F2 and 3xFL-CrhR-R, followed by another reamplification using the

primers 3xFL-CrhR-F3 and 3xFL-CrhR-R. The resulting amplicon 3xFLAG-*crhR* was

digested with *Nde*I and *Xba*I and ligated into pJET1.2. The copper inducible promoter,

P_{petE} (Zhang *et al.*, 1992) was amplified with the primers petE-FP and petE-RP and cloned

into pJET1.2 via Gibson assembly. The *rrnB* terminator was amplified from pBAD using

primers rrnB-TT-F and rrnB-TT-R and ligated into pJET1.2 via *Sac*I and *Xba*I restriction

sites yielding pJET1.2::P*petE*::3xFLAG-*crhR*::*rrnB*T.

To introduce the K57A mutation into *crhR*, primers crhR(K57A)_inverse_fw and

crhR(K57A)_inverse_rv were used in an inverse PCR to generate $CrhR_{K57A}$. Both the

pJET1.2::P*petE*::3xFLAG-*crhR*::*rrnB*T and $CrhR_{K57A}$.constructs contain the P_{petE} promoter

followed by the triple FLAG tag translationally fused to the *crhR* ORF followed by the *rrnB*

terminator. To eliminate the *crhR* gene and create the negative control $\Delta crhR_{control}$, the

vector pJET1.2::P*petE*::3xFLAG-*crhR*::*rrnB*T was inverse PCR amplified with primers

petE-3xF-rrnb_neg_ctrl_FP and petE-3xF-rrnb_neg_ctrl_RP carrying *Sac*I restriction

sites on the 5′ overhangs, cleaved with *Sac*I and self-ligated.

For cloning of the constructs into pVZ321 (Zinchenko *et al.*, 1999), the constructs were

amplified from pJET1.2 using the primers petE_foR-pVZ_FP and rrnB_foR-pVZ_RP

carrying *Eco*RI restriction sites. The amplified constructs were inserted into the *Eco*RI

restriction site of pVZ321. This insertion resulted in disruption of the chloramphenicol

resistance cassette of pVZ321, so that the final constructs pVZ321::P*petE*::3xFLAG-

crhR::*rrnB*T (strain $CrhR_{WT}$), pVZ321::P*petE*::3xFLAG-*crhR*(K57A)::*rrnB*T (strain $CrhR_{K57A}$)

and pVZ321::P*petE*::3xFLAG::*rrnB*T (strain $\Delta crhR_{control}$) possessed only kanamycin

resistance. The resulting pVZ321 plasmids were transferred into the $\Delta crhR$ *Synechocystis*

strain (Prakash *et al.*, 2010) via triparental mating with *E. coli* J53/RP4 and TOP10F'

(Scholz *et al.*, 2013). Exconjugants were selected on agar plates containing 20 µg/ml

spectinomycin (Sp) and 50 µg/ml kanamycin (Km) in BG-11 (Rippka *et al.*, 1979).

Culture conditions

E. coli strains were grown in liquid LB medium (10 g l^{-1} bacto-tryptone, 5 g l^{-1} bacto-yeast
extract, 10 g l^{-1} NaCl), with continuous agitation or on agar-solidified (1.5% [w/v] bacto
agar) LB, supplemented with appropriate antibiotics, at 37°C.

Synechocystis strains were cultivated in the presence of appropriate antibiotics either in
Erlenmeyer flasks in BG-11 medium (Rippka *et al.*, 1979) (experiments shown in **Fig. 2**)
or in 100 ml two-tier vessel CellDEG cultivators (CellDEG GmbH (Bähr *et al.*, 2016)) at
the indicated temperatures in fresh water organisms (FWO) medium with shaking. FWO
medium consists of 50 mM $NaNO_3$, 15 mM KNO_3, 2 mM $MgSO_4$, 0.5 mM $CaCl_2$, 0.025
mM H_3BO_3, 0.15 mM $FeCl_3$/Na_2EDTA, 1.6 mM KH_2PO_4, 2.4 mM K_2HPO_4, 10 mM
$NaHCO_3$, 10 nM $MnCl_2$, 1 nM $ZnSO_4$, 2 nM Na_2MoO_4, 2 nM $CuSO_4$, 30 pM $CoCl_2$.

High CO_2 concentrations were delivered to the flasks from a carbonate buffer (1.447 M
K_2CO_3 and 12.486 M $KHCO_3$) through a highly gas-permeable polypropylene membrane.
The CellDEG cultivators were inoculated at an OD_{750} ~0.8 and a light intensity of 150 µmol
photons m^{-2} s^{-1}. Light intensity was adjusted to culture density; therefore, 24 h after the
start of inoculation light intensity was increased to 300 µmol photons m^{-2} s^{-1} and to 500
µmol photons m^{-2} s^{-1} after another 24 h. To induce gene expression from the Cu^{2+}-
responsive promoter P$_{petE}$ in the CellDEG system, 6 µM $CuSO_4$ was added to the medium.
The cells were harvested at OD_{750} ~20, which was reached upon 72 hours of cultivation.

FLAG-tag affinity purification, UV crosslinking and RNA co-IP

Synechocystis cultures were grown in CellDEG cultivators to an OD_{750} ~20 at 30°C. For
cold stress, cultures were exposed to 20°C for 2 h. The cells were placed in a 20 x 20 cm
Petri dish and irradiated three times on ice with 500 mJ UV-C light (254 nm) in a
Stratalinker® 2400. The treated cells were harvested by centrifugation (4,000 x g, 10 min,
4°C). Cell pellets were resuspended in precooled TBS buffer (50 mM Tris-HCl (pH 7.5)),
containing cOmplete™ Protease Inhibitor Cocktail (Roche), washed once with TBS and
centrifuged again.

The washed and cross-linked cells were suspended in TBS buffer and 0.5 volume of glass

beads were added. Cells were disrupted mechanically applying 5 cycles at 6,500 rpm with

10 s breaks on ice between the cycles in a Precellys® 24 homogenizer. The cell lysate

was separated by centrifugation (13,000 x g, 15 min, 4°C) into two fractions, one

containing membrane proteins and the other containing soluble proteins. The latter was

incubated with 50 µl of Anti-FLAG® M2 magnetic beads (Sigma Aldrich) to allow binding

of FLAG-tagged proteins at 4°C for 4 hours. Subsequently, beads were washed twice in

TBS-containing cOmplete™ Protease Inhibitor Cocktail followed by a wash step that

removed the non-crosslinked RNA by incubation in 500 µl RNA elution buffer (25 mM Tris-

HCl (pH7.5), 2 M NaCl) for 15 min at room temperature. The crosslinked RNA was

released by digestion of the FLAG-tagged protein with 20 µg of proteinase K for 10 min at

room temperature, purified using the RNA Clean & Concentrator-5 Kit (Zymo Research)

and utilized for the generation of cDNA libraries for Illumina sequencing as described

below. All co-immunoprecipitation experiments were performed in biological duplicates.

RNA preparation

Synechocystis cell pellets were disrupted by incubation for 15 min at 65 °C in PGTX as

described by Pinto *et al.* (2009). Chloroform/isoamyl alcohol (IAA) (24:1) extraction was

performed for 10 min at room temperature followed by centrifugation at 3,270 x g for 15

min at room temperature for phase separation. The chloroform/IAA extraction was

repeated on the aqueous phase, followed by isopropanol precipitation. Precipitated RNA

was pelleted by centrifugation (13,000 x g, 30 min, 4°C), washed with 70% ethanol, air-

dried and resuspended in nuclease-free water. RNA concentration was determined on a

NanoDrop ND-1000 spectrophotometer. RNA purity and quality was evaluated on a

Fragment Analyzer parallel capillary electrophoresis system (Agilent).

Preparation of cDNA libraries

To initiate cDNA library preparation, 0.3-5 µg of total RNA was treated with 4 U of TURBO

DNase (Life Technologies) in 2 consecutive incubation steps, each at 37°C for 15 min as

indicated by the manufacturer. Next, RNA clean-up and separation into small (17-200 nt)

and large (>200 nt) RNA fractions was performed using RNA Clean & Concentrator

columns (Zymo Research). The large RNA fraction was fragmented according to the

protocol by Pfeifer-Sancar *et al.* (2013). After RNA Clean & Concentrator column

purification, both fractions were combined and all subsequent steps performed as

previously described (Pfeifer-Sancar *et al.*, 2013) except that the clean-up of RNA 5'-

pholyphosphatase (Epicentre)-treated samples was performed by Clean & Concentrator

column purification. For RNA adapter ligation, the UGA linker (Supplementary **Table S5**)

was used. After RNA adapter ligation and cDNA synthesis, the samples were excised

from 2% agarose gels and purified using the NucleoSpin Gel and PCR Clean-up kit

(Macherey-Nagel) using NTC buffer for solubilization of the gel slices. After PCR

amplification, residual primers were removed by adding 10 µl of ExoSAP-IT (USB) to 50

µl of PCR, and with samples incubated for 15 min at 37°C followed by heat inactivation of

the enzyme for 15 min at 85°C. The samples were cleaned-up using a NucleoSpin Gel

and PCR Clean-up kit. The quality of RNA and DNA was analyzed on a Fragment

Analyzer parallel capillary electrophoresis system (Agilent). Sequencing of libraries was

performed on a HiSeq 3000 sequencer with a 2 x 150 bp read mode.

Bioinformatics analyses

Analysis of the raw reads from the RNA sequencing was performed using the Galaxy web

platform (https://usegalaxy.eu/ (Afgan *et al.*, 2018)). Illumina paired-end reads were

trimmed, adapter sequences and reads shorter than 14 nt filtered out using cutadapt 1.16

(Martin, 2011). The remaining reads were mapped to the chromosome and plasmids of

Synechocystis using bowtie2 version 2.3.4.3 with the following selected parameters for

paired-end reads: -I 0 -X 500 --fr --no-mixed --no-discordant --very-sensitive(Langmead

and Salzberg, 2012). Peak calling of the mapped reads was performed by PEAKachu

0.1.0.2 with the parameters: --pairwise_replicates --norm_method deseq --mad_multiplier

2.0 –fc_cutoff 1 --padj_treshold 0.05. The total number of reads as well as those that

mapped to the *Synechocystis* genome are listed in **Supplementary Material, Table S1**.

The transcriptome data have been deposited in the SRA database under the accession
numbers SAMN14615974 to SAMN14615989 and SAMN14651279 to SAMN14651286.

Generation of recombinant CrhR

To obtain recombinant, C-terminal His-tagged CrhR, the *crhR* gene was amplified from
the constructs pVZ321::P*petE*::3xFLAG-*crhR*::*rrnB*T with primers pQE70_crhR-fw and
pQE70_crhR-rv (**Table S5**). The pQE70 vector was amplified with primers pQE_GIBSOn-
fw and pQE_GIBSOn-rv. The fragments were mixed and transformed into *E. coli* Top10
F' using AQUA cloning (Beyer *et al.*, 2015) yielding strain pQE70:crhR-6xHis. For
recombinant protein expression, the construct pQE70::*crhR*-6xHis was transformed into
E. coli M15. Overnight cultures were diluted 1:100 in fresh LB medium supplemented with
50 µg/ml ampicillin, and grown to an OD_{600} of 0.7 at 37°C. Protein expression was induced
by adding IPTG to 1 mM final concentration. Three hours after induction, cells were
harvested by centrifugation at 6,000 x g for 10 min at room temperature. Cell pellets were
resuspended in lysis buffer (50 mM NaH_2PO_4 (pH 8), 1 M NaCl, 10% glycerol, 15 mM
imidazole, cOmplete™ Protease Inhibitor Cocktail (Roche)) and lysed using the One Shot
constant cell disruption system (Constant Systems Limited, U.K.) at 2.4 kbar. Cell lysates
were cleared by centrifugation at 13,000 g for 30 min at 4°C and the lysate was filtered
through 0.45 µm Supor-450 filters (Pall). Recombinant proteins were immobilized on a
HiTrap Talon crude 1 ml column (GE Healthcare), equilibrated with buffer A (50 mM
NaH_2PO_4 (pH 8), 500 mM NaCl), and eluted with elution buffer B (50 mM NaH_2PO_4 (pH
8), 500 mM imidazole, 500 mM NaCl). The protein concentration was calculated with a
Bradford assay using bovine serum albumin as the standard.

Cy3 labelling of RNA

To initiate Cy3 RNA labelling for EMSA analysis, two volumes of 6 mM KIO_4 were added
to 4 µg RNA and incubated at room temperature in the dark for 1 h. The RNA was
precipitated at -20°C for 1 h by addition of one volume of ethylene glycol/H_2O (1:1), 2.9
volumes of 100% ethanol, and 0.1 volume of 3.3 M NaCl and pelleted at 13,000 x g for 30

min at 4°C. The RNA pellet was washed with 70 % ethanol, air dried, and resuspended in

10 µl of 50 mM Cy3 dye dissolved in DMSO (Thermo Fisher Scientific) and incubated at

37°C in the dark for 2 h. Two volumes of 0.1 M Tris-HCl (pH 7.5) were added followed by

the addition of 20 mM freshly prepared $NaBH_4$ and the reaction mix incubated at 4°C in

the dark for 30 min. The reaction was quenched and RNA precipitated by ethanol

precipitation.

Electrophoretic mobility shift assays

Binding of different amounts of recombinant CrhR (0.1 to 5 pmol) to 0.2 pmol of Cy3

labelled RNA was performed in 20 mM HEPES-KOH (pH 8.3) buffer containing 3 mM

$MgCl_2$, 1 mM DTT, 500 µg/ml BSA. As a substrate competitor, 1 µg of LightShift Poly

(dIdC) (Thermo Fischer Scientific) was included in each assay. The reaction was

incubated at room temperature for 15 min prior to separation on 2% agarose-TAE gels.

The signals were visualized with a Laser Scanner Typhoon FLA 9500 (GE Healthcare)

using a green light laser at a wavelength of 532 nm and a Cy3 filter (LPG, DGR1, BPG1.

Acknowledgements

We thank the Freiburg Galaxy Team for maintaining this great bioinformatics resource.

Author contributions

WRH designed the study, CS established the cDNA library preparation and supported the

bioinformatic analysis; JF, BH and RR sequenced cDNA libraries; AM, FH, RB and WRH

analyzed data; AS and JSSP constructed the *Synechocystis* strains. All other experiments

were carried out by AM. AM, GWO and WRH wrote the paper with contributions from all

authors.

FUNDING

This work was supported by grants from the Baden-Wuerttemberg Foundation BWST_NCRNA_008, by the German Research Foundation (DFG) - 322977937/GRK2344 and by the Federal Ministry of Education and Research (BMBF) program RNAProNet, grant 031L0164B to WRH and RB, by the EU ITN "Photo.COMM" to support the stay of AS in the lab of WRH, a Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery Grant, grant 171319 to GWO and a Department of Biotechnology (DBT) of India grant BT/PR13616/BRB/10/774/2010 to JSSP.

References

Afgan E, Baker D, Batut B, *et al.* 2018. The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. Nucleic Acids Research **46**, W537–W544.

Angeleri M, Zorina A, Aro E-M, Battchikova N. 2018. Interplay of SpkG kinase and the Slr0151 protein in the phosphorylation of ferredoxin 5 in *Synechocystis* sp. strain PCC 6803. FEBS letters **592**, 411–421.

Ashburner M, Ball CA, Blake JA, *et al.* 2000. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. Nature Genetics **25**, 25–29.

Bähr L, Wüstenberg A, Ehwald R. 2016. Two-tier vessel for photoautotrophic high-density cultures. Journal of Applied Phycology **28**, 783–793.

Beyer HM, Gonschorek P, Samodelov SL, Meier M, Weber W, Zurbriggen MD. 2015. AQUA cloning: A versatile and simple enzyme-free cloning approach. PLOS One **10**, e0137652.

Boudet N, Aubourg S, Toffano-Nioche C, Kreis M, Lecharny A. 2001. Evolution of intron/exon structure of DEAD helicase family genes in Arabidopsis, Caenorhabditis, and Drosophila. Genome Research **11**, 2101–2114.

Bourgeois CF, Mortreux F, Auboeuf D. 2016. The multiple functions of RNA helicases as drivers and regulators of gene expression. Nature Reviews Molecular Cell Biology **17**, 426–438.

Capel C, Albaladejo I, Egea I, *et al.* 2020. The res (restored cell structure by salinity) tomato mutant reveals the role of the DEAD-box RNA helicase SlDEAD39 in plant development and salt response. Plant, Cell & Environment **43**, 1722–1739.

Cassier-Chauvat C, Chauvat F. 2014. Function and regulation of ferredoxins in the cyanobacterium, *Synechocystis* PCC6803: recent advances. Life **4**, 666–680.

Cavaiuolo M, Chagneau C, Laalami S, Putzer H. 2020. Impact of RNase E and RNase J on global mRNA metabolism in the cyanobacterium *Synechocystis* PCC6803. Frontiers in Microbiology **11**, 1055.

Chamot D, Colvin KR, Kujat-Choy SL, Owttrim GW. 2005. RNA structural rearrangement via unwinding and annealing by the cyanobacterial RNA helicase, CrhR. The Journal of Biological Chemistry **280**, 2036–2044.

Chamot D, Magee WC, Yu E, Owttrim GW. 1999. A cold shock-induced cyanobacterial RNA helicase. Journal of Bacteriology **181**, 1728–1732.

Cordin O, Banroques J, Tanner NK, Linder P. 2006. The DEAD-box protein family of RNA helicases. Gene **367**, 17–37.

de la Cruz J, Kressler D, Linder P. 1999. Unwinding RNA in Saccharomyces cerevisiae: DEAD-box proteins and related families. Trends in Biochemical Sciences **24**, 192–198.

Dann M, Leister D. 2019. Evidence that cyanobacterial Sll1217 functions analogously to PGRL1 in enhancing PGR5-dependent cyclic electron flow. Nature Communications **10**, 5299.

Eriksson J, Salih GF, Ghebramedhin H, Jansson C. 2000. Deletion mutagenesis of the 5′ *psbA2* region in *Synechocystis* 6803: Identification of a putative cis element involved in photoregulation. Molecular Cell Biology Research Communications **3**, 292–298.

Fairman ME, Maroney PA, Wang W, Bowers HA, Gollnick P, Nilsen TW, Jankowsky E. 2004. Protein displacement by DExH/D 'RNA helicases' without duplex unwinding. Science **304**, 730–734.

Gene Ontology Consortium. 2021. The Gene Ontology resource: enriching a GOld mine. Nucleic Acids Research **49**, D325–D334.

Georg J, Rosana ARR, Chamot D, Migur A, Hess WR, Owttrim GW. 2019. Inactivation of the RNA helicase CrhR impacts a specific subset of the transcriptome in the cyanobacterium *Synechocystis* sp. PCC 6803. RNA Biology **16**, 1205–1214.

Gierga G, Voss B, Hess WR. 2012. Non-coding RNAs in marine *Synechococcus* and their regulation under environmentally relevant stress conditions. The ISME Journal **6**, 1544–1557.

Gong Z, Dong C-H, Lee H, Zhu J, Xiong L, Gong D, Stevenson B, Zhu J-K. 2005. A DEAD box RNA helicase is essential for mRNA export and important for development and stress responses in *Arabidopsis*. The Plant Cell **17**, 256–267.

Gruber AR, Lorenz R, Bernhart SH, Neuböck R, Hofacker IL. 2008. The Vienna RNA websuite. Nucleic Acids Research **36**, W70–W74.

Hihara Y, Sonoike K, Kanehisa M, Ikeuchi M. 2003. DNA microarray analysis of redox-responsive genes in the genome of the cyanobacterium *Synechocystis* sp. strain PCC 6803. Journal of Bacteriology **185**, 1719–1725.

Hishiya S, Hatakeyama W, Mizota Y, Hosoya-Matsuda N, Motohashi K, Ikeuchi M, Hisabori T. 2008. Binary reducing equivalent pathways using NADPH-thioredoxin reductase and ferredoxin-thioredoxin reductase in the cyanobacterium *Synechocystis* sp. strain PCC 6803. Plant & Cell Physiology **49**, 11–18.

Holmqvist E, Wright PR, Li L, Bischler T, Barquist L, Reinhardt R, Backofen R, Vogel J. 2016. Global RNA recognition patterns of post-transcriptional regulators Hfq and CsrA revealed by UV crosslinking in vivo. The EMBO Journal **35**, 991–1011.

Hör J, Gorski SA, Vogel J. 2018. Bacterial RNA biology on a genome scale. Molecular Cell **70**, 785–799.

Huang C-K, Shen Y-L, Huang L-F, Wu S-J, Yeh C-H, Lu C-A. 2016. The DEAD-box RNA helicase AtRH7/PRH75 participates in pre-rRNA processing, plant development and cold tolerance in *Arabidopsis*. Plant and Cell Physiology **57**, 174–191.

Huckauf J, Nomura C, Forchhammer K, Hagemann M. 2000. Stress responses of *Synechocystis* sp. strain PCC 6803 mutants impaired in genes encoding putative alternative sigma factors. Microbiology **146**, 2877–2889.

Jankowsky E, Gross C, Shuman S, Pyle A. 2001. Active disruption of an RNA-protein interaction by a DExH/D RNA helicase. Science **291**, 121–125.

Jarmoskaite I, Russell R. 2011. DEAD-box proteins as RNA helicases and chaperones: DEAD-box proteins. Wiley Interdisciplinary Reviews: RNA **2**, 135–152.

Jiang H-B, Song W-Y, Cheng H-M, Qiu B-S. 2015. The hypothetical protein Ycf46 is involved in regulation of CO2 utilization in the cyanobacterium *Synechocystis* sp. PCC 6803. Planta **241**, 145–155.

Jones PG, Mitta M, Kim Y, Jiang W, Inouye M. 1996. Cold shock induces a major ribosomal-associated protein that unwinds double-stranded RNA in E*scherichia coli*. Proceedings of the National Academy of Sciences of the USA **93**, 76–80.

Kadowaki T, Nishiyama Y, Hisabori T, Hihara Y. 2015. Identification of OmpR-family response regulators interacting with thioredoxin in the cyanobacterium *Synechocystis* sp. PCC 6803. PloS ONE **10**, e0119107.

Kappell AD, van Waasbergen LG. 2007. The response regulator RpaB binds the high light regulatory 1 sequence upstream of the high-light-inducible *hliB* gene from the cyanobacterium *Synechocystis* PCC 6803. Archives of Microbiology **187**, 337–342.

Kerfeld CA, Sawaya MR, Brahmandam V, Cascio D, Ho KK, Trevithick-Sutton CC, Krogmann DW, Yeates TO. 2003. The crystal structure of a cyanobacterial water-soluble carotenoid binding protein. Structure **11**, 55–65.

Khemici V, Prados J, Petrignani B, Di Nolfi B, Bergé E, Manzano C, Giraud C, Linder P. 2020. The DEAD-box RNA helicase CshA is required for fatty acid homeostasis in *Staphylococcus aureus*. PLoS Genetics **16**, e1008779.

Kiefer M, Nauerth BH, Volkert C, Ibberson D, Loreth A, Schmidt A. 2020. Gene function rather than reproductive mode drives the evolution of RNA helicases in sexual and apomictic *Boechera*. Genome Biology and Evolution **12**, 656–673.

Köbler C, Schultz S-J, Kopp D, Voigt K, Wilde A. 2018. The role of the *Synechocystis* sp. PCC 6803 homolog of the circadian clock output regulator RpaA in day-night transitions. Molecular Microbiology **110**, 847–861.

Komenda J, Knoppová J, Krynická V, Nixon PJ, Tichý M. 2010. Role of FtsH2 in the repair of Photosystem II in mutants of the cyanobacterium *Synechocystis* PCC 6803 with impaired assembly or stability of the CaMn(4) cluster. Biochimica et Biophysica Acta **1797**, 566–575.

Kujat SL, Owttrim GW. 2000. Redox-regulated RNA helicase expression. Plant Physiology **124**, 703–714.

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nature Methods **9**, 357–359.

Lehnik-Habrink M, Rempeters L, Kovacs AT, Wrede C, Baierlein C, Krebber H, Kuipers OP, Stülke J. 2013. DEAD-box RNA helicases in *Bacillus subtilis* have multiple functions and act independently from each other. Journal of Bacteriology **195**, 534–544.

Linder P, Jankowsky E. 2011. From unwinding to clamping — the DEAD box RNA helicase family. Nature Reviews Molecular Cell Biology **12**, 505–516.

Linder P, Owttrim GW. 2009. Plant RNA helicases: linking aberrant and silencing RNA. Trends in Plant Science **14**, 344–352.

Lu C-A, Huang C-K, Huang W-S, Huang T-S, Liu H-Y, Chen Y-F. 2020. DEAD-box RNA helicase 42 plays a critical role in pre-mRNA splicing under cold stress. Plant Physiology **182**, 255–271.

Macovei A, Vaid N, Tula S, Tuteja N. 2012. A new DEAD-box helicase ATP-binding protein (OsABP) from rice is responsive to abiotic stress. Plant Signaling & Behavior **7**, 1138–1143.

Mahbub M, Hemm L, Yang Y, *et al.* 2020. mRNA localization, reaction centre biogenesis and thylakoid membrane targeting in cyanobacteria. Nature Plants **6**, 1179–1191.

Margulis L. 1981. *Symbiosis in cell evolution: Life and its environment on the early earth*. W.H.Freeman & Co Ltd.

Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet.journal **17**, 10.

Martin W, Kowallik K. 1999. Annotated English translation of Mereschkowsky's 1905 paper 'Über Natur und Ursprung der Chromatophoren im Pflanzenreiche'. European Journal of Phycology **34**, 287–295.

Martin W, Rujan T, Richly E, Hansen A, Cornelsen S, Lins T, Leister D, Stoebe B, Hasegawa M, Penny D. 2002. Evolutionary analysis of *Arabidopsis*, cyanobacterial, and chloroplast genomes reveals plastid phylogeny and thousands of cyanobacterial genes in the nucleus. Proceedings of the National Academy of Sciences USA **99**, 12246–12251.

Matthes A, Schmidt-Gattung S, Köhler D, Forner J, Wildum S, Raabe M, Urlaub H, Binder S. 2007. Two DEAD-box proteins may be part of RNA-dependent high-molecular-mass protein complexes in *Arabidopsis* mitochondria. Plant Physiology **145**, 1637–1646.

Mazouni K, Domain F, Chauvat F, Cassier-Chauvat C. 2003. Expression and regulation of the crucial plant-like ferredoxin of cyanobacteria. Molecular Microbiology **49**, 1019–1029.

Mereschkowsky C. 1905. Über Natur und Ursprung der Chromatophoren im Pflanzenreiche. Biol. Centralbl. **25**, 593–604.

Mingam A, Toffano-Nioche C, Brunaud V, Boudet N, Kreis M, Lecharny A. 2004. DEAD-box RNA helicases in *Arabidopsis thaliana*: establishing a link between quantitative expression, gene structure and evolution of a family of genes: AtRH expression by real-time Q-PCR. Plant Biotechnology Journal **2**, 401–415.

Mutsuda M, Sugiura M. 2006. Translation initiation of cyanobacterial *rbcS* mRNAs requires the 38-kDa ribosomal protein S1 but not the Shine-Dalgarno sequence: development of a cyanobacterial *in vitro* translation system. The Journal of Biological Chemistry **281**, 38314–38321.

Muzzopappa F, Kirilovsky D. 2020. Changing color for photoprotection: The Orange Carotenoid Protein. Trends in Plant Science **25**, 92–104.

Nakagawa S, Niimura Y, Miura K, Gojobori T. 2010. Dynamic evolution of translation initiation mechanisms in prokaryotes. Proceedings of the National Academy of Sciences of the USA **107**, 6382–6387.

Nawaz G, Kang H. 2017. Chloroplast-or mitochondria-targeted DEAD-box RNA helicases play essential roles in organellar RNA metabolism and abiotic stress responses. Frontiers in Plant Science **8**, 871.

Nawaz G, Lee K, Park SJ, Kim Y-O, Kang H. 2018. A chloroplast-targeted cabbage DEAD-box RNA helicase BrRH22 confers abiotic stress tolerance to transgenic Arabidopsis plants by affecting translation of chloroplast transcripts. Plant Physiology and Biochemistry: PPB **127**, 336–342.

Oun S, Redder P, Didier J-P, François P, Corvaglia A-R, Buttazzoni E, Giraud C, Girard M, Schrenzel J, Linder P. 2013. The CshA DEAD-box RNA helicase is important for quorum sensing control in *Staphylococcus aureus*. RNA Biology **10**, 157–165.

Owttrim GW. 2013. RNA helicases. RNA Biology **10**, 96–110.

Pfeifer-Sancar K, Mentz A, Rückert C, Kalinowski J. 2013. Comprehensive analysis of the *Corynebacterium glutamicum* transcriptome using an improved RNAseq technique. BMC Genomics **14**, 888.

Pinto F, Thapper A, Sontheim W, Lindblad P. 2009. Analysis of current and alternative phenol based RNA extraction methodologies for cyanobacteria. BMC Molecular Biology **10**, 79.

Ponce-Toledo RI, Deschamps P, López-García P, Zivanovic Y, Benzerara K, Moreira D. 2017. An early-branching freshwater cyanobacterium at the origin of plastids. Current Biology **27**, 386–391.

Prakash JSS, Krishna PS, Sirisha K, Kanesaki Y, Suzuki I, Shivaji S, Murata N. 2010. An RNA helicase, CrhR, regulates the low-temperature-inducible expression of heat-shock genes *groES, groEL1* and *groEL2* in *Synechocystis* sp. PCC 6803. Microbiology **156**, 442–451.

Py B, Higgins CF, Krisch HM, Carpousis AJ. 1996. A DEAD-box RNA helicase in the *Escherichia coli* RNA degradosome. Nature **381**, 169–172.

Redder P, Hausmann S, Khemici V, Yasrebi H, Linder P. 2015. Bacterial versatility requires DEAD-box RNA helicases. FEMS Microbiology Reviews **39**, 392–412.

Riediger M, Hihara Y, Hess WR. 2018. From cyanobacteria and algae to land plants: The RpaB/Ycf27 regulatory network in transition. Perspectives in Phycology **5**, 13–25.

Riediger M, Kadowaki T, Nagayama R, Georg J, Hihara Y, Hess WR. 2019. Biocomputational analyses and experimental validation identify the regulon controlled by the redox-responsive transcription factor RpaB. iScience **15**, 316–331.

Riediger M, Spät P, Bilger R, Voigt K, Maček B, Hess WR. 2021. Analysis of a photosynthetic cyanobacterium rich in internal membrane systems via gradient profiling by sequencing (Grad-seq). The Plant Cell, Epub ahead of press.

Rippka R, Deruelles J, Waterbury JB, Herdman M, Stanier RY. 1979. Generic assignments, strain histories and properties of pure cultures of cyanobacteria. Journal of General Microbiology **111**, 1–61.

Ritter SPA, Lewis AC, Vincent SL, Lo LL, Cunha APA, Chamot D, Ensminger I, Espie GS, Owttrim GW. 2020. Evidence for convergent sensing of multiple abiotic stresses in cyanobacteria. Biochimica et Biophysica Acta (BBA) - General Subjects **1864**, 129462.

Rocak S, Linder P. 2004. DEAD-box proteins: the driving forces behind RNA metabolism. Nature Reviews Molecular Cell Biology **5**, 232–241.

Rogers GW, Komar AA, Merrick WC. 2002. eIF4A: the godfather of the DEAD box helicases. Progress in Nucleic Acid Research and Molecular Biology **72**, 307–331.

Rosana ARR, Chamot D, Owttrim GW. 2012*a*. Autoregulation of RNA helicase expression in response to temperature stress in *Synechocystis* sp. PCC 6803. PLoS One **7**, e48683.

Rosana ARR, Ventakesh M, Chamot D, Patterson-Fortin LM, Tarassova O, Espie GS, Owttrim GW. 2012*b*. Inactivation of a low temperature-induced RNA helicase in *Synechocystis* sp. PCC 6803: Physiological and morphological consequences. Plant and Cell Physiology **53**, 646–658.

Rosana ARR, Whitford DS, Fahlman RP, Owttrim GW. 2016. Cyanobacterial RNA helicase CrhR localizes to the thylakoid membrane region and cosediments with degradosome and polysome complexes in *Synechocystis* sp. strain PCC 6803. Journal of Bacteriology **198**, 2089–2099.

Rosana ARR, Whitford DS, Migur A, Steglich C, Kujat-Choy SL, Hess WR, Owttrim GW. 2020. RNA helicase–regulated processing of the *Synechocystis rimO-crhR* operon results in differential cistron expression and accumulation of two sRNAs. Journal of Biological Chemistry **19**, 6372–6386.

Rowland JG, Pang X, Suzuki I, Murata N, Simon WJ, Slabas AR. 2010. Identification of components associated with thermal acclimation of photosystem II in *Synechocystis* sp. PCC6803. PLoS One **5**, e10511.

Rowland JG, Simon WJ, Prakash JSS, Slabas AR. 2011. Proteomics reveals a role for the RNA helicase *crhR* in the modulation of multiple metabolic pathways during cold acclimation of *Synechocystis* sp. PCC6803. Journal of Proteome Research **10**, 3674–3689.

Schneider S, Schwer B. 2001. Functional domains of the yeast splicing factor Prp22p. The Journal of Biological Chemistry **276**, 21184–21191.

Scholz I, Lange SJ, Hein S, Hess WR, Backofen R. 2013. CRISPR-Cas systems in the cyanobacterium *Synechocystis* sp. PCC6803 exhibit distinct processing pathways involving at least two Cas6 and a Cmr2 protein. PLoS One **8**, e56470.

Silva P, Thompson E, Bailey S, Kruse O, Mullineaux CW, Robinson C, Mann NH, Nixon PJ. 2003. FtsH is involved in the early stages of repair of photosystem II in *Synechocystis* sp PCC 6803. The Plant Cell **15**, 2152–2164.

Singh AK, Sherman LA. 2002. Characterization of a stress-responsive operon in the cyanobacterium *Synechocystis* sp. strain PCC 6803. Gene **297**, 11–19.

Sireesha K, Radharani B, Krishna PS, Sreedhar N, Subramanyam R, Mohanty P, Prakash JSS. 2012. RNA helicase, CrhR is indispensable for the energy redistribution and the regulation of photosystem stoichiometry at low temperature in *Synechocystis* sp. PCC6803. Biochimica et Biophysica Acta (BBA) - Bioenergetics **1817**, 1525–1536.

Tanner NK, Linder P. 2001. DExD/H box RNA helicases: from generic motors to specific dissociation functions. Molecular Cell **8**, 251–262.

Tarassova OS, Chamot D, Owttrim GW. 2014. Conditional, temperature-induced proteolytic regulation of cyanobacterial RNA helicase expression. Journal of Bacteriology **196**, 1560–1568.

Tyystjärvi T, Herranen M, Aro EM. 2001. Regulation of translation elongation in cyanobacteria: membrane targeting of the ribosome nascent-chain complexes controls the synthesis of D1 protein. Molecular Microbiology **40**, 476–484.

Vinnemeier J, Hagemann M. 1999. Identification of salt-regulated genes in the genome of the cyanobacterium *Synechocystis* sp. strain PCC 6803 by subtractive RNA hybridization. Archives of Microbiology **172**, 377–386.

Wang B, Chai H, Zhong Y, Shen Y, Yang W, Chen J, Xin Z, Shi H. 2020. The DEAD-box RNA helicase SHI2 functions in repression of salt-inducible genes and regulation of cold-inducible gene splicing. Journal of Experimental Botany **71**, 1598–1613.

Whitford DS, Whitman BT, Owttrim GW. 2021. Genera specific distribution of DEAD-box RNA helicases in cyanobacteria. Microbial Genomics, Epub ahead of press.

Wu C-Y, Nagy PD. 2020. Role reversal of functional identity in host factors: Dissecting features affecting pro-viral versus antiviral functions of cellular DEAD-box helicases in tombusvirus replication. PLoS Pathogens **16**, e1008990.

Yang Q, Jankowsky E. 2005. ATP- and ADP-dependent modulation of RNA unwinding and strand annealing activities by the DEAD-box protein DED1. Biochemistry **44**, 13591–13601.

Yeremenko N, Jeanjean R, Prommeenate P, Krasikov V, Nixon PJ, Vermaas WFJ, Havaux M, Matthijs HCP. 2005. Open reading frame *ssr2016* is required for antimycin A-sensitive photosystem I-driven cyclic electron flow in the cyanobacterium *Synechocystis* sp. PCC 6803. Plant & Cell Physiology **46**, 1433–1436.

Zhan J, Steglich C, Scholz I, Hess WR, Kirilovsky D. 2021. Inverse regulation of light harvesting and photoprotection Is mediated by a 3'end-derived sRNA in cyanobacteria. The Plant Cell, Epub ahead of press.

Zhang L, McSpadden B, Pakrasi HB, Whitmarsh J. 1992. Copper-mediated regulation of cytochrome c553 and plastocyanin in the cyanobacterium *Synechocystis* 6803. The Journal of Biological Chemistry **267**, 19054–19059.

Zinchenko VV, Piven IV, Melnik VA, Shestakov SV. 1999. Vectors for the complementation analysis of cyanobacterial mutants. Russ J Genet. **35**, 228–32.

Tables

Table 1. Selected transcripts recovered in CrhR co-IPs. Assignment to one of the three experiments (**Fig.** 1) is given by the numbers in the first column, followed by gene IDs according to Genbank file NC_000911 and localization of the peak within the coding region (CDS) or UTR. Annotation of the interacting RNAs and their transcription regulation, either activated (A) or repressed (R) by the transcription factor RpaB under low light (Riediger *et al.*, 2019) or with "redox", if they were classified as redox-responsive genes (Hihara *et al.*, 2003; Ritter *et al.*, 2020). Detailed information on peak locations, enrichment factors and adjusted p values generated by the PEAKachu algorithm (Holmqvist *et al.*, 2016) are given in **Tables S2** to **S4**.

E	ID	Annotation	RpaB/ Redox
1,2,3	slr1834 5'UTR, CDS	PsaA, PSI P700 apoprotein subunit Ia **Fig. 3C**	A,R
1,3	slr1655 5'UTR, CDS	PSI subunit XI, PsaL	A
3	slr0927 CDS, 3'UTR	PSII protein PsbD/D2	R
1	smr0009 CDS	photosystem II reaction center protein PsbN	R
3	slr0906 CDS	PSII reaction center protein PsbB/CP47	R
3	slr1311 5'UTR, CDS, 3'UTR	PSII reaction center protein PsbA/D1	R
3	sll1867 5'UTR, CDS	PSII reaction center protein PsbA/D1	R
1	slr1674 5'UTR, CDS	protection of PSII from thermal damage	R, redox
3	slr0144 to slr0148 CDS	operon of PSII assembly proteins, Fed5	A
3	slr0342 5'UTR, CDS	PetB, cytochrome b_6	R
1	ssl0020 5'UTR, CDS	ferredoxin I, PetF	R
2,3	sll0199 5'UTR, CDS	plastocyanin, PetE	
3	slr2067 5'UTR, CDS	allophycocyanin alpha subunit ApcA	
3	slr1986 CDS	allophycocyanin beta subunit ApcB	
3	ssr3383 5'UTR, CDS	allophycocyanin-associated linker ApcC, **Fig. 4**	
3	slr0335 5'UTR, CDS	PBS LCM core-membrane linker ApcE	A
3	slr2051 5'UTR, CDS 3'UTR	PBS rod-core linker CpcG1	
3	slr0009 to slr0012 CDS	RuBisCO operon *rbcLXS*	
1,2,3	slr0373-slr0374 CDS	Ycf46, CO_2 utilization, **Fig. 3B**	redox
1	ssr2016 CDS	Pgr5, soluble electron acceptor	R
1,2	slr0228 CDS	FtsH2 protease	R
1	slr1963 CDS	Water-soluble carotenoid protein OCP	redox
1,3	slr0623 5'UTR, CDS	thioredoxin A	
1,2,3	slr1545 5'UTR, CDS	Extracytoplasmic function (ECF) sigma factor SigG, **Fig. 3A**	A
1,2,3	slr0083 5'UTR, CDS	DEAD-box RNA helicase CrhR, **Fig. 5**	redox

1	slr0923 5'UTR, CDS	YCF65; 30S ribosomal protein 3 PSRP-3	R
1	slr1638 5'UTR, CDS	Hypothetical protein	R
1,2	slr0551 CDS	ribonuclease J	redox
1	sll0541 5'UTR	*desC/des9*, Acyl-CoA desaturase	A
1	sll0006 5'UTR, CDS	putative aminotransferase AspC	A
1,3	slr1841 CDS	Porin, major outer membrane protein	R
1	slr0915 CDS, 3'UTR; *trnL* intron; 6803t34 exon 2; slr0917 5'UTR	group I intron, tRNA(Leu)[UAA] exon 2, endonuclease; *bioF*, 8-amino-7-oxononanoate synthase	R
3	6803t34 exon 1, *trnL* intron, slr0915 5'UTR, CDS	tRNA(Leu)[UAA] exon 1, group I intron, intron endonuclease	R

Figures

Fig. 1. Overview on the experimental flow. (A) Schematic representation of the cyanobacterial strains created for the chosen strategy. A deletion mutant, Δ*crhR*, previously generated by replacement of the *crhR* gene (*slr0083*) by a spectinomycin (Sp^R) resistance gene by homologous recombination (Prakash *et al*., 2010) served as the initial platform. In this background, strains were engineered that express the native form of CrhR (CrhR$_{WT}$) or a K57A substitution (CrhR$_{K57A}$), both translationally fused to an N-terminal triple FLAG tag in the conjugative vector pVZ321. A strain expressing only the 3xFLAG-tag was constructed as a control (Δ*crhR*$_{control}$). **(B)** Experimental design. Experiment 1: co-IP of the UV-crosslinked RNA-CrhR$_{WT}$ complexes was performed from *Synechocystis* CrhR$_{WT}$ cultures exposed to cold stress at 20°C for 2 h or, Experiment 2, grown at 30°C. Experiment 3: co-IP of UV-crosslinked RNA-CrhR$_{K57A}$ complexes was performed from cultures grown at 30°C. In parallel, to control for non-specific enrichment, an RNA mock co-IP was performed with the Δ*crhR*$_{control}$ strain. Cultures in Experiments 1 to 3 were grown in FWO medium using the CellDEG system. Each experiment was performed in biological duplicates and cDNA libraries were prepared from the enriched RNA and from the total RNA of all respective cultures omitting the co-IP step.

Fig. 2. Phenotypic characterization of wild type, CrhR$_{WT}$, CrhR$_{K57A}$ and the Δ$crhR$$_{control}$ strains. (A) Growth of the wild type (WT), CrhR$_{WT}$, CrhR$_{K57A}$ and Δ$crhR$$_{control}$ strains was determined spectrophotometrically by measuring the OD$_{750}$ of cultures grown at 20°C in FWO medium. The experiments were performed in biological triplicates and standard deviations are indicated. **(B)** Pigmentation in the wild type (WT), CrhR$_{WT}$, CrhR$_{K57A}$ and Δ$crhR$$_{control}$ cultures at an OD$_{750}$ of 0.52 grown at 20°C in BG-11 medium. Spectra were normalized to 750 nm. **(C)** Visual appearance of the four strains at the indicated OD$_{750}$. Strains were grown in the presence of antibiotics and 2 μM Cu^{2+}.

Fig. 3. Peak pattern of transcripts enriched in CrhR co-IPs. (A) The region containing the 5' UTR, start codon and first 72 to 90 codons were enriched for gene *slr1545* encoding the RNA polymerase sigma factor SigG. This peak was significant in all three experiments, using CrhR$_{K57A}$ or CrhR$_{WT}$ as bait proteins at the permissive temperature of 30°C (read coverage in red and brown) as well as with CrhR$_{WT}$ at the non-permissive temperature of 20°C (coverage in blue). **(B)** Reads

yielding a peak in the middle of the first gene and extending into the entire intergenic space between two genes and including the 5' UTR, start codon and first 72 to 90 codons of the second gene were recovered for the dicistronic operon *slr0373* and *slr0374*. **(C)** Multiple sharp peaks indicating a multitude of interactions between CrhR and the dicistronic *psaAB* mRNA. *psaAB* encode the PSI P700 apoprotein subunits Ia and A2. In all three panels, the coverage from reads recovered from CrhR co-IP experiments is shown (red, brown and blue) together with the coverage from RNA-Seq analyses of the respective total transcriptomes in shades of grey as indicated. The numbers along the x-axis correspond to chromosomal nucleotide positions in *Synechocystis* (Genbank accession number NC_000911.1), while the normalized numbers of reads are given on the y-axis.

Fig. 4. CrhR and *apcC* transcript interaction. Reads recovered from CrhR co-IP experiments cover the 5' UTR and almost the entire coding sequence of *apcC/ssr3383* that encodes the 7.8 kDa allophycocyanin-associated linker polypeptide ApcC. Only the reads recovered in experiment 3 using CrhR$_{K57A}$ as bait protein were significantly enriched (red) but an enrichment was also observed in CrhR$_{WT}$ cells grown at the permissive temperature 30°C (brown). The sRNA ApcZ originates from within the *apcC* coding sequence (boxed). The 5' end of ApcZ was previously mapped to position 1431853, 46 nt upstream of the *apcC* stop codon (Zhan *et al.*, 2021). The secondary structure of ApcZ is shown in the inset. The interaction decreases in the region from which ApcZ originates, colored light green in both the transcript and in the corresponding segment of ApcZ secondary structure where it corresponds to half of the first helix of the predicted secondary structure (nt 1 to 30). The RNAfold algorithm was used for the secondary structure prediction (Gruber *et al.*, 2008).

Fig. 5. CrhR interaction with its own transcript *in vivo* and *in vitro*. (A) Enrichment of *crhR* mRNA in the UV-crosslinking RNA co-IP from *Synechocysti*s CrhR$_{WT}$ grown at 20°C or 30°C (experiments 1 and 2). The colored plots show the read coverage of the enriched RNAs, as indicated. **(B)** EMSA showing binding of recombinant His-tagged CrhR to the *crhR* mRNA. For *crhR* transcript synthesis, a 150 nt DNA fragment starting from the start codon of the *crhR/slr0083* ORF was amplified with primers EMSA_CrhR-T7_Fw (carrying the T7 promoter sequence followed by two Gs) and EMSA_CrhR-T7_Rv (**Table S5**), matching to the genomic positions 2887644 to 2887793. The resulting 150 nt transcript was Cy3 labelled. Binding of 0.2 pmol of the Cy3 labelled transcripts with the indicated amounts of purified recombinant His-tagged CrhR protein was performed in the presence of poly dI-dC (1 µg).