ANALYSIS OF CHD-7 DEFECTIVE DAUER NEMATODES IMPLICATES COLLAGEN MISREGULATION IN CHARGE SYNDROME FEATURES

Diego Martín Jofré[1][*], Dane Kristian Hoffman[2][*][§], Ailen S. Cervino[3], McKenzie Grundy[2], Sijung Yun[4], Francis Raj Gandhi Amrit[5], Donna Beer Stolz[6], Esteban Salvatore[7], Fabiana Alejandra Rossi[8], Arjumand Ghazi[5,9], M. Cecilia Cirio[3], Judith L. Yanowitz[2,10][#], Daniel Hochbaum[1][#]

[*]authors contributed equally

[#] Co-corresponding authors

[1] Departamento de Biodiversidad y Biología Experimental. FCEN. Universidad de Buenos Aires. Buenos Aires, Argentina.
[2] Magee-Womens Research Institute, Pittsburgh, PA, USA
[3] Instituto de Fisiología, Biología Molecular y Neurociencias (IFIBYNE-UBA-CONICET), Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires (UBA). Buenos Aires, Argentina.
Institute
[4] Laboratory of Molecular Biology, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, Maryland, USA.
[5] Department of Pediatrics, University of Pittsburgh School of Medicine. Pittsburgh, PA, USA.
[6] Center for Biologic Imaging, University of Pittsburgh Medical School. Pittsburgh, PA, USA.
[7] mAbxience S.A.U. Buenos Aires, Argentina.
[8] Instituto de Investigaciones en Medicina Traslacional (IIMT) - CONICET, Universidad Austral. Pilar, Buenos Aires, Argentina.
[9] Departments of Developmental Biology and Cell Biology & Physiology, University of Pittsburgh School of Medicine. Pittsburgh, PA, USA.
[10] Departments of Obstetrics, Gynecology & Reproductive Sciences; Developmental Biology; Microbiology and Molecular Genetics; and the Hillman Cancer Center, University of Pittsburgh, Pittsburgh, PA, USA.

[§]Current Address: Department of Cancer & Cell Biology. Baylor College of Medicine. Houston, Texas, USA.

*Correspondence: Correspondence should be addressed to

Dr. Daniel Hochbaum
Ciudad Universitaria. Intendente Guiraldes 2160, 4 piso. Laboratorio 79
CABA, Buenos Aires
C1428BGA
Argentina
hochbaumd@bg.fcen.uba.ar
+54 (11) 5285 8658
51
Dr. Judith Yanowitz
Magee-Womens Research Institute
204 Craft Avenue
Pittsburgh, PA
15213
United States of America
58
59

60   **Conflict of Interest**

61   The authors declare no conflicts of interest

63
64

ABSTRACT

CHARGE syndrome is a complex developmental disorder caused by mutations in the chromodomain helicase DNA-binding protein7 (CHD7) and characterized by retarded growth and malformations in the heart and nervous system. However, despite the public health relevance of this disorder, relevant targets of CHD7 that relate to disease pathology are still poorly understood. Here we report that *chd-7*, the nematode ortholog of CHD7, is required for dauer morphogenesis, lifespan determination, and stress response. Genetic epistasis placed *chd-7* in the TGF-β pathway. Consistent with our discoveries, we found *chd-7* to be allelic to *scd-3*, a previously identified dauer suppressor from the TGF-β pathway. Interestingly, DAF-12 transcriptionally upregulated *chd*-7, which is necessary to repress *daf-9* for execution of the dauer program. Transcriptomic analysis comparing *chd-7*–defective and normal dauers showed enrichment of collagen genes, consistent with a conserved role for the TGF-β pathway in expression of the extracellular matrix. To validate a conserved function for *chd-7* in vertebrates, we used *Xenopus laevis* embryos, an established model to study craniofacial development. Morpholino mediated knockdown of Chd7 led to embryonic lethality, a reduction in *col2a1* mRNA levels and craniofacial defects in tadpoles. Both lethality and malformations were partially rescued in Chd7-depleted embryos by over-expression of *col2a1*. We suggest that pathogenic features of CHARGE syndrome caused by Chd7 mutations, such as craniofacial malformations, result from the reduction of collagen levels. These studies establish *C. elegans* as an amenable animal model to study the etiology of the developmental defects associated with pathogenic Chd7.

Keywords: CHARGE Syndrome, *C. elegans*, dauer, longevity, TGF-β, Col2a1

INTRODUCTION

CHARGE syndrome is a rare and severe neurodevelopmental disorder that affects the neural tube and neural crest cell derivatives, leading to hypogonadism, heart defects and craniofacial anomalies among other features (1). Inactivating mutations in *CHD7* (chromodomain-helicase-DNA binding 7) are the predominant cause of CHARGE, accounting for greater than 90% of the cases (2). *CHD7* is also mutated in Kallmann syndrome, a milder neurodevelopmental disorder with features overlapping with CHARGE, including impaired olfaction and hypogonadism (3). Exome sequencing studies in patients with autism spectrum disorders (ASDs) identified recurrent disruptive mutations in *CHD8* (4). The CHD proteins comprise a highly conserved family of SNF2-related ATP-dependent chromatin remodelers that are involved in chromatin remodeling and transcriptional regulation (5). Despite the public health relevance of these cognitive disorders, functionally relevant targets of CHD7/8 that relate to disease pathology are still poorly understood. The development of fly, fish and mouse models of CHARGE has abetted in characterization of associated dysfunction, but our understanding of the underlying pathology of CHARGE is still incomplete (6–11). CHD-7 is the ortholog of human CHD7 and CHD8 in *Caenorhabditis elegans*. It contains an ATPase/SNF2 domain and two chromodomains for nucleosome interaction and is the only worm homolog of the Class III CHD proteins, which contain a signature BRK domain (Brahma and Kismet domain) (Figure 1E).

When *C. elegans* encounter crowding, starvation, or high temperature during early development, worms can halt reproductive programs to enter an alternative larval stage known as dauer. Dauers are long-lived, highly stress resistant, and exhibit altered motility and metabolism (12–15). Upon return to normal growth conditions, the larvae exit dauer and develop into fertile adults. Study of dauer formation mutants has provided fundamental insights into pathways affecting longevity, neurodevelopment, metabolism, autophagy and neurodegeneration (12, 15–18).

The DAF-2/insulin/IGF-1 (IIS) signaling pathways controls the dauer entry decision by coupling external cues with neuroendocrine signaling (19). In favorable

conditions, DAF-2 activity initiates a conserved kinase cascade, leading to

phosphorylation and inhibition of the transcription factor DAF-16/FOXO. In harsh

environments, a decrease in the activity of DAF-2 and downstream components of

the pathway leads to activation of DAF-16 and causes animals to arrest as dauers

(20, 21). In addition to the DAF-2 pathway, DAF-7/ TGFβ also regulates dauer

development (22). When worms sense suitable conditions for reproductive

development, ASI neurosensory cells secrete DAF-7, which binds to DAF-1/4

receptors leading to activation and phosphorylation of the R-SMAD complex DAF-

8/14, promoting reproductive programs and inhibiting the pro-dauer complex

composed of the SMAD protein DAF-3 and repressor DAF-5. Conversely, absence

of DAF-7 leads to activation of the DAF-3/DAF-5 complex to promote dauer entry

(23, 24).

DAF-7/TGF-β like and DAF-2/IIS pathway were initially described as parallel

pathways to regulate dauer entry (25), but recent observations suggest a strong

positive feedback between these pathways for dauer entry and longevity (15–20).

First, decreased signaling through the TGF-β pathway leads to differential

expression of many DAF-16 regulated genes with functions in longevity and dauer

entry, such as SOD-3 or insulin peptides. This cross-activation of target genes may

be important to amplify weak signals from each sensory pathway to make an all-or-

none decision to enter dauer (26–28). Second, the longevity of *daf-2* mutants can be

blocked or enhanced by *daf-5* and *daf-3*, respectively, suggesting that transcriptional

components of the TGF-β can modulate IIS-dependent longevity genes. Third, for

dauer development, *daf-16* is epistatic to *daf-7/8/14 daf-c* mutants (28, 29). Lastly,

both signaling pathways converge on *daf-9* and *daf-12* to integrate outputs for

diapause entry (30, 31). Central to dauer entry decision is the nuclear receptor DAF-

12, which integrates IIS and TGF-β signaling pathways (32). (33) In chromatin

immunoprecipitation (ChIP)-chip studies, we identified *chd-7* as a DAF-12 target

gene whose loss caused defective execution of dauer morphogenesis programs.

We show that while CHD-7 can modulate multiple IIS-associated processes,

including *daf-2(e1370)* longevity and immunity. Our epistatic experiments place *chd-*

7 in the TGF-β/DAF-7 pathway. Whole genome mRNA expression profiling of partial

dauers show that *chd-7* mutants fail to repress *daf-9* expression during dauer

development, preventing developmental arrest of the constitutive dauer mutant *daf-*

7(e1372). In addition, we found collagens to be the most differentially regulated

genes, consistent with a known role for the TGFβ pathway in regulating components

of the extracellular matrix (34).To validate a conserved function for *chd-7* in

vertebrates and study the relevance of our results for CHARGE etiology, we used

Xenopus laevis. Disruption of *Chd7* function in *Xenopus* embryos results in

craniofacial defects that mimic CHD-dependent pathological phenotypes (35, 36).

We demonstrate that Chd7 regulates expression of the collagen type-II, alpha 1,

col2a1, the main collagen protein of cartilage (37). Interestingly, craniofacial

malformations and embryonic lethality due to *Chd7* knockdown can be rescued by

col2a1 expression. These findings suggest a conserved function of *Chd7*/*chd-7* in

regulation of extracellular matrix components and raise the intriguing possibility that

defects in collagen expression may contribute to the craniofacial defects seen in

CHD7/8-dependent syndromes.

RESULTS

***chd-7* functions in development of the dauer larva.**

Previously, we identified ~3000 potential DAF-12 target genes by ChIP-chip (32).

We reasoned that among these genes would be novel regulators of dauer

development. To identify such genes, we took advantage of the temperature-

sensitive, constitutive dauer (Daf-c) allele of *daf-2(e1371)*. At the non-permissive

temperature of 25°, all the offspring enter the dauer phase and become SDS

resistant (20). We reasoned that targets of DAF-12 that function in dauer formation

should suppress the Daf-c phenotype of *daf*-2 and produce either defective, SDS-

sensitive "partial" dauers or reproductive adults. Screening for RNAi clones that led

to development of SDS-sensitive offspring in the more severe *daf-2(e1370)* mutant

background, we discovered that *chd-7* RNAi produced arrested, SDS sensitive

larvae, indicative of the partial dauer phenotype (38). While typically the dauer larva

has an axial ratio (length:width) of nearly 30:1 (20), the partial dauers resulting from

chd-7(RNAi) exhibited a significant reduction of these proportions and appeared to

have defects in radial constriction of the dauer cuticle (Figure 1A and B). By using a

transcriptional reporter, we confirmed a strong upregulation of *chd-7* by DAF-12

(Figure 1C and D).

To further validate our screen, we crossed *daf-2(e1370)* worms with three *chd-7*

deletion alleles available from the Nematode Knockout Consortia (39). Allele *chd-*

7(tm6139) contains a 594bp deletion that generates a frame shift and premature

stop codon, eliminating all known protein domains (Figure 1E). As shown in Figure

1D, SDS-sensitive dauers are obtained when double mutants are grown at 25°C,

validating our screen. Comparison of two partial deletion alleles uncovered a critical

role for the BRK domain in dauer formation. The *chd-7(gk290)* allele contains an

859bp deletion that spans the BRK domain and introduces a frameshift that

eliminates the last 356aa. The *chd-7(gk306)* deletion is slightly more C-terminal,

truncating the protein immediately after the BRK domain (Figure 1E). When crossed

into *daf-2(e1370)* worms, *chd-7(gk306)* develops normal, SDS-resistant dauer

larvae, whereas *chd-7(gk290)* formed partial dauers, that are sensitive to detergent

(Figure 1F). Importantly, a translational reporter expressing CHD-7::GFP (Figure

1F) rescued the "partial dauer" phenotypes observed in *chd-7(gk290);daf-2(e1370)*

and *chd-7(tm6139);daf-2(e1370)* mutants.

The dauer larva is characterized by a slim physique because of a reduction in the

volume of epidermal tissues, including the hypodermis, seam cells and XXX cells. In

addition, the epidermal tissues produce the dauer cuticle, which confers protection

against external damage and dehydration. Because of this cuticle stiffness,

hypodermal seam cells fuse to produce the alae, which are bilateral ridges that

facilitate body motion. Dauers also switch their metabolism to accumulate lipids to

survive for longer periods (40). To further characterize the nature of the defects in

the *chd-7*-induced partial dauers, we first studied the cuticle by scanning electro

microscopy (SEM) and the adherent junction-associated protein marker AJM-

1::GFP. As shown in Figure 1E, partial dauers fail to fuse seam cells, leading to a

defective dauer alae. Next we used Oil Red O to examine fat storage and found that

unlike other partial dauer mutants (17), the *chd-7;daf-2* abnormal dauers did not

exhibit fat-storage deficiencies (Supplementary Figure 1). During the dauer stage,

the germ cells remains quiescent to protect their reproductive capacity (41). We

observed that germ line in *chd-7* mutant dauers was substantially larger than in

dauers and appeared to arrest with an L3-like germ line morphology (Figure 1F).

Therefore, we conclude that major morphological changes that occur during dauer

larvae formation fail to be executed in *chd-7* mutants including radial constriction of

the body, SDS resistance, a developmentally-arrested germ line and presence of

dauer alae.

chd-7 is required for glp-1(e2144) and daf-2(e1370) longevity

In addition to dauer development, the IIS pathway also regulates longevity (12).

Thus, we sought to investigate whether *chd-7* also has roles in the determination of

lifespan. First, we compared survival of wild-type (N2) worms with two *chd-7* alleles

and found that the dauer-suppressor allele *chd-7(gk290)* significantly shortens

lifespan, whereas *chd-7(gk306)* had only a marginal effect on longevity (Figure 2A).

Surprisingly, a CHD-7::GFP rescue transgene also reduced N2 lifespan (Figure 2B),

suggesting that *chd-7* copy number can influence longevity. We then analyzed how

chd-7 affects longevity of IIS mutants. Remarkably, the dauer suppressor allele *chd-*

7(gk290), but not *chd-7(gk306),* shortened the lifespan extension of daf-2 mutants to

an extent comparable with the null allele of *daf-16,* the key IIS downstream target

(Figure 2C) (12). Furthermore, *daf-2* longevity is fully restored by CHD-7::GFP

(Figure 2D).

To determine if the effects on lifespan were specific to the IIS pathway, we also

assayed the contribution of *chd-7* to the longevity of germ cell-less animals. *glp-*

1(e2144) animals are long-lived due to the absence of proliferative germ cells (42).

This lifespan extension is dependent on *chd-7* as *chd-7(gk290);glp-1(e2144)* double

mutants have a mean lifespan significantly shorter than *glp-1(e2144)* single mutants

(Figure 2E).

In addition to longevity, IIS reduction also enhances resistance against multiple

stressors including pathogen attack and starvation. We asked if chd-7 inactivation

also impaired daf-2 mutant's immunoresistance. As shown in Figure 2F, *chd-7*

mutants repressed the increased survival of *daf-2* worms upon exposure to the

opportunistic pathogen *Pseudomonas aeruginosa* strain PA14 (43). In contrast, *chd-*

7 has a modest effect on survival of wild-type worms' pathogen resistance.

Furthermore, the *chd-7(gk290)* allele reduces the survival of starved *daf-2(e1370)* L1

larvae (Supplemental Figure 3) (44). Taken together, these results suggest that *chd-*

7 mediates the increased lifespan of at least two longevity paradigms (IIS mutants

and germ cell-less animals) as well as the survival response to pathogens and

starvation.

CHD-7 belongs to the TGF-β pathway

Genetic analyses implicated *chd-7* as a dauer defective mutant (Daf-d) in the IIS

signaling pathway (Figure 1). We sought to understand whether *chd-7* has also roles in

the TGF-β dauer pathway (25, 26). While *chd-7(gk290); daf-2(e1370)* and *chd-*

7(tm6139);daf-2(e1370) arrest as partial dauers at the restrictive temperature of 25°C,

double mutants with *daf-7(e1372)* totally bypass the dauer developmental arrest to

become fertile adults (Figure 3A). Interestingly, *daf-7*-dependent arrest is recapitulated

at higher temperatures (26.5°C), suggesting that *chd-7* has a Hid (high temperature-

induced dauer formation) phenotype (45). As expected for a putative transcriptional

regulator, epistasis analysis place CHD-7 downstream the receptor DAF-1 and the R-

Smad DAF-14 (Figure 3B).

In addition to dauer formation, the TGF-β signaling pathway also regulates body size

and male tale development, mainly through the DBL-1 ligand (46, 47). We compared

the length of *chd-7* young adults and found them to be significantly shorter than N2

(Figure 3C). We also observed that males carrying the severe loss of function allele

chd-7(tm6139) fail to mate with *fog-2* mutants (Figure 3D) due to defects in male tail

development (not shown). This male infertility is rescued by the CHD-7::GFP transgene.
279

More than twenty years ago, in a screen for suppressors of dauer formation within the
TGF-β pathway, *Inoue et al* identified three complementation groups (29). We noticed
that one of these, *scd-3* (suppressor of constitutive dauer 3), was located between *unc-*
11 and *dpy-5* on chromosome I, in the same genetic region as *chd-7*. *scd-3(sa253)*
phenotypes include low brood size, Egl (egg-laying defects), Dpy (weak short body
shape) and Mab (Male abnormal), phenotypes shared with *chd-7*. To determine if *chd-7*
and *scd-3* are allelic, we sequenced *scd-3(sa253)* and found a single G/A mutation in
exon 8 of *chd-7*, that introduces a premature STOP codon at position Q2422,
eliminating the BRK domain (Figure 1E).
289

Transcriptomics analysis identifies collagens as targets of CHD-7
To gain further insight into the role of CHD-7 in dauer development, we performed RNA-
Seq on *daf-2(e1370)* dauers and *chd-7(gk290);daf-2(e1370)* partial dauers. We
analyzed differentially expressed genes (DEGs) and found decreased expression of 28
genes and increased expression of 56 genes in double mutants (adjusted p-value by
Benjamini Hochberg's False Discovery Rate (FDR)<0.05) (Figure 4A). Among the latter
group, we found the *daf-9* cytochrome p450, which integrates inputs from TGF-β and
insulin/IGF-I pathways to regulate *daf-12* activity (48). While RNA-seq reads for this
gene were marginally detected in *daf-2* dauers, partial dauers exhibited a significantly
increased number of reads (Figure 4B). Because *chd-7* loss promotes *daf-9*
upregulation in partial dauers (Figure 3A), we hypothesized that expression levels must
be critical in the decision to develop as fertile adults or to enter diapause and that the
increased expression of *daf-9* in the *chd-7;daf-2* double mutants was preventing full
execution of the dauer program. Consistent with this hypothesis, we found that
depletion of *daf-9* in *chd7;daf-7* animals restored dauer entry at the non-permissive
temperature of 25°C (Figure 4C). Thus, the inability of *chd-7* mutants to fully repress
daf-9 is sufficient to inhibit the developmental arrest of TGF-β mutants.
307

Transcriptomics analysis showed that 10% of the DEGs are collagens (*col-103*, *col-50*,

dpy-2, *col-184*, *col-141*, *col-142*, *col-42* and *dpy-9*), which are structural components of

the cuticle. This is not surprising due to the conserved role of the TGF-β pathway in

regulation of the extracellular matrix (ECM) components, such as fibronectins and

collagens (34). Recently, Liu *et al* performed transcriptomic analysis of TGF-β mutants

during diapause entry and found both *daf-9* downregulation and misexpression of

multiple collagens, supporting our finding that CHD-7 functions within the TGF-β

signaling pathway (27).

Chd7 regulates *col2a1* during *Xenopus* embryogenesis

The partial dauer larvae that arise from knockdown of *chd-7* are short and thin and lack

the dauer alae, indicative of cuticle defects (Figure 1G). Type-II collagen is an ECM

protein conserved in all multicellular animals, which forms fibrils (49) and have

fundamental roles in development and tissues homeostasis (50). In vertebrates, the

fibrillar type-II collagen is the major structural protein of cartilage and plays a prominent

role in cranial development in multiple organisms (37, 51). Collagen, type-II, alpha 1

(Col2a1) is the major component of the cartilage matrix having a structural function and

being an important extracellular signaling molecule for regulation of chondrocyte

proliferation, metabolism and differentiation (52–54). The African frog *Xenopus laevis* is

a well-established model to study vertebrate facial disorders which often arise from

defects in neural crest development and migration (36). In *Xenopus* embryos, prior

studies established that Chd7 regulates neural crest specification and migration and its

depletion recapitulates the craniofacial defects seen in CHARGE patients (35). To

investigate a conserved mechanism of action for Chd7 regulating collagen in

vertebrates, we used a previously validated morpholino to induce Chd7 knockdown

(*chd7*-MO) (35). In *Xenopus,* Col2a1 is essential for normal development of the skeleton

and its expression is restricted to the cartilaginous skeleton of the embryo and adult frog

(55). We first analyzed *col2a1* expression by RT-qPCR in *chd7*-MO injected embryos

and observed a significant reduction upon morpholino injection (Figure 5A). For targeted

disruption of Chd7 function, we injected the dorsal-animal (D1) blastomeres of 8-cell

stage embryos fated to contribute to the dorsal anterior structures. *In situ* hybridization

of unilaterally injected embryos confirmed a reduction of *col2a1* expression in the

branchial arches (55) upon *Chd7* depletion (Figure 5B). The injection of *chd7*-MO into

both D1 blastomeres of 8-cell stage embryos with doses between 1.25 ng and 5 ng

showed 10-15% embryonic mortality. In contrast, 10 ng led to more than 50% lethality

(data not shown and Figure 5C). We sought to investigate if mortality was related to

collagen deficits and conducted rescue experiments co-expressing *Xenopus col2a1*

mRNA. As shown in Figure 5C, co-injection of *col2a1* mRNA improves embryo survival

relative to those injected with *chd7*-MO alone. We then asked if ectopic *col2a1* mRNAs

could overcome craniofacial defects associated with *chd7* loss-of-function. We

examined the eye size and eye distance as microthalmia and midline defects are often

associated with CHARGE syndrome (56) and are recapitulated in *Xenopus* embryos

(35). Both eye size and eye distance are reduced in Chd7-depleted embryos and

partially rescued by *col2a1* mRNA expression (Figure 5D and 5E). Therefore,

expression of *col2a1* ameliorates the phenotypes associated with pathogenic Chd7

suggesting that collagen is an important target of this helicase.

DISCUSSION

We initially identified *chd-7* in chromatin immunoprecipitation (ChIP)-chip studies of the

nuclear receptor DAF-12 (32), a transcription factor regulating worm aging,

development and dauer formation (57), and found it to be critical for *daf-2* mediated

dauer development. Comparison of the *chd-7* alleles *gk290* and *gk306* shows a critical

role for the BRK domain in dauer development and longevity. In CHARGE patients,

deletions or mutations within the BRK domains of Chd7 are sufficient to elicit all the

features characteristic of the disease underscoring the importance of this domain (58).

The phenotypic differences between the worm alleles highlight the potential of the worm

to delimit functional domains of CHD-7 that contribute to disease pathology. In mice,

homozygous mutations in *Chd7* lead to embryonic lethality at E10.5, in part because

Chd7 is necessary for early brain development (35). In worms, the presumptive null

alleles c*hd-7(tm6139)* and *scd-3(sa253)* were viable but showed reproductive defects,

such as improper gonad proliferation and migration, reduced fecundity, male tail defects

and hermaphrodite vulval defects (Supplemental Figure 5 and (29)) indicating that *C.*

elegans are more able to tolerate loss of CHD-7 than mice or humans. Thus, our

studies establish *C. elegans* as an amenable animal model to study developmental

defects and postnatal features observed in pathogenic Chd7.

Both a *chd-7* mutation and CHD-7 overexpression shortened the lifespan of otherwise

wild-type worms suggesting that the CHD-7 levels must be tightly regulated to ensure

proper development. Of note, CHD7 is the most commonly amplified gene in tumors

amongst the CHD superfamiliy members, and its overexpression is associated with

aggressive subtypes of breast cancer, high tumor grade and poor prognosis (59). Thus,

in humans, as well as worms, there are gain-of-function phenotypes associated with

Chd7/*chd-7* overexpression. While outside the scope of this work, a CHD-7

overexpression strain could be instrumental for modeling CHD7-dependent tumor

progression.

Genetic epistasis analyses place CHD-7 downstream of the TGFβ-like DAF-7, the type I

receptor DAF-1, and the R-SMAD DAF-14 placing CHD-7 at the level of the Co-Smad

DAF-3 and the Sno/Ski repressor DAF-5 which are also Daf-d. 4. While *daf-3* and *daf-5*

completely suppress dauer formation in a *daf-7* background at 25°, both genes fail to

suppress dauer formation in TGF-β mutants at 27°C (13, 38) supporting a role for CHD-

7 at the DAF-3/DAF-5 step in the pathway. While DAF-3 and DAF-5 work together to

control dauer entry, they have opposite effects on *daf-2*-induced longevity: *daf-3*

enhances *daf-2(e1370)* longevity while *daf-5* mutations suppresses it (28). We observed

that *chd-7* suppresses *daf-2* longevity (Figure 2C), like *daf-5(e1386)*. Together these

data suggest that CHD-7 either regulates *daf-5* expression or directly interacts with

DAF-5 to regulate downstream target genes or both. In mice, CHD7 was shown to

physically interact with SMAD1 and form a transcriptional complex with SMAD4, the

mammalian ortholog of DAF-3 (60). We therefore speculate that DAF-3, DAF-5, and

CHD-7 may be in ternary complex that regulates *daf-9* expression for dauer entry

(Figure 4C). Interactome mapping of the TGF-β pathway also identified SWSN-1, a

SWI/SNF subunit component of the BAF complex, as a physical interactor of DAF-3

(61). CHD7 interacts with human and Xenopus PBAF (polybromo- and BRG1-

associated factor-containing complex) to control neural crest gene expression (35). In

worms, both *swsn-1* and *chd-7* fail to develop normal dauers in *daf-2* and *daf-7* mutants

(Supplemental Figure 4 and (62)). Thus, we envision that CHD-7 may work together

with the BAF complex and DAF-3/DAF-5 to control gene expression of targets genes

critical for dauer formation.

DAF-9 activity is necessary and sufficient for the decision to enter diapause; reduced

activity of TGF-β and insulin/IGF-I pathways leads to *daf-9* repression and dauer

development. Conversely, *daf-9* expression in the hypodermis is sufficient to inhibit

diapause, driving reproductive programs of *daf-7* mutants (27, 30). Ectopic *daf-9*

expression drives reproductive programs in the weak *daf-2(e1368)* allele, but only

partially suppresses *daf-2(e1370)* diapause animals which arrest at as L3 or early

L4, (30). Based on these observations, we speculate that *daf-9* misexpression

explains a subset of the *chd-7* phenotypes observed herein, including the partial

dauer phenotype, gonad migration defects, and vulval protrusions, all of which

overlap with published *daf-9* phenotypes (30, 31, 63). Interestingly, *daf-9* is both

upstream and downstream of *daf-12* for dauer formation *(30, 31)*. We show here that

DAF-12 regulates *chd-7* expression and that CHD-7 in turn regulates *daf-9*

expression. Therefore, we propose that CHD-7 belongs to the transcriptional

complex that regulates the feedback loop between *daf-12* and *daf-9*. Moreover, we

found increased expression of the *eak-4* paralog F14H8.2 in partial dauers (Figure

4A). High activity of this gene may prevent dauer entry by increasing DAF-9 activity

(64).

Among the differentially expressed genes, we note the presence of several G protein

coupled receptors (GPCRs), which could be candidate chemoreceptors for dauer

pheromone or environmental cues and may be required for either dauer entry or

maintenance (42). Consistent with this, we observed that the *chd-7* partial dauers

resume reproductive development significantly faster than normal dauers when

returned to 16°C. Consistent with our transcriptomic studies, *Liu et al.* analyzed the

expression differences between L2/L3 larvae and TGF-β mutants undergoing

diapause and found enrichment of collagen genes, GPCRs and *daf-9*, further

validating our placement of *CHD-7* in the TGF-β dauer signaling pathway (27).

The expression profiles of adult TGF-β and IIS mutants show a significant

overlapping and co-regulation of many DAF-16 target genes. Indeed, the DAF-16-

associated element (DAE), is found overrepresented in TGF-β regulated genes (26).

We analyzed DAF-16 (L4) and CHD-7 (YA) CHIP-seq data generated by the

Modencode project (65) and found both transcriptional regulators to share a

significant number of genes (Supplemental Figure 5). This, along with our

observation that chd-7 is necessary of lifespan extension of daf-2 mutants, indicates

that *chd-7* modulates longevity by regulation of the IIS pathway.

In *C. elegans*, the TGF-β family pathway has diversified and has unique ligands and

effectors for the control of dauer induction and aspects of somatic development.

Mutations in the TGF-β ligand *dbl-1* and its downstream components cause a small

body size (Sma phenotype) (47, 66). Consistent with *chd-7* being a component of the

TGF-β signaling pathway, *chd-7(gk290)* and *chd-7(tm6139)* are significantly shorter

than wild-type worms. This phenotype is also observed in *scd-3* mutants (29). The *dbl-1*

pathway mutants have defects in male tale development (47), a phenotype shared in

scd-3 (29) and *chd-7* mutant animals (data not shown and (29)). Therefore, our results

suggest that CHD-7 acts as a regulator of both the *daf-7* and *dbl-1* branches of the

TGF-β pathway. The short body length of *dbl-1(ev580)* animals is partially rescued by

mutations in the collagen gene *lon-3*, a component of the *C. elegans* cuticle.

Interestingly, *lon-3* expression depends on the *dbl-1* pathway, suggesting that in worms

and in vertebrates, the TGF-β pathway regulates extracellular matrix components, such

as collagens (34, 67).

Craniofacial anomalies seen in CHARGE patients involve tissues deriving from cells of

the neural crest, including craniofacial cartilage and bone, heart, ears and eyes (56, 68).

Xenopus laevis is an established model to understand fundamental questions of

craniofacial biology (36). *chd7* knockdown in frogs and more recently demonstrated in

zebrafish (11, 35), leads to defects in neural crest specification and migration, which

was recently recapitulated in human embryonic stem cells (hESCs) (69). Our RNA-seq

analysis revealed that CHD-7 regulates expression of genes from the cuticle, a modified

extracellular matrix (70). Here, we show a conserved role for Chd7 in regulating *col2a1*

expression, a type II collagen and the major component of cartilage (51). While

additional studies are required, we speculate that Chd7 could regulate *col2a1* in a

complex with the transcription factor Sox10 (71, 72) or through the TGF-β signaling

pathway (34, 73). Supporting the latter mechanism, it was demonstrated recently that

TGF-β regulates *col2a1* expression in chondrocytes (74). Taken together, our data

supports a model in which collagen misexpression by pathogenic Chd7 leads to

craniofacial defects and embryonic lethality.

METHODS

***C. elegans* strains**

All strains were grown and maintained on standard nematode growth medium (NGM). In

Argentina (Hochbaum lab), plates were supplemented with 0.1mg/ml streptomycin and

100U/ml Nystatin using either *E. coli* OP50 or OP50-1 strains as the food source. In

Pittsburgh (Yanowitz lab), strains were grown on standard NGM seeded with OP50.

Temperature-sensitive strains were maintained at 16°C and shifted to the non-

permissive temperature of 25°C to induce dauer formation, unless otherwise indicated.

All other strains were maintained at 20°C. All *daf-7* strains were maintained at low

density to prevent dauer induction at 16°C. Dauer studies were performed in both the

Yanowitz and Hochbaum laboratories to validate results. All *chd-7* alleles were

outcrossed at least 5 times prior to use.

Strains utilized in this study are listed in Supplementary Table 1. Standard genetic

crosses were used to make double or triple mutants. The presence of mutant alleles

was confirmed a) by the *daf-c* phenotypes in animals heterozygous for additional

mutations and b) by PCR and/or sequencing for all additional mutations. Primers and
PCR conditions used in this study are shown in Supplemental Table 2.
496

RNAi screen for dauer suppressors

498 All RNAi clones were picked from the Ahringer bacterial feeding library (75). These *E.*
499 *coli* clones were seeded on NGM plates supplemented with 1mM of IPTG (Isopropyl β-
500 D-1-thiogalactopyranoside) and 0.1μg/ml ampicillin, and used for inducing RNAi by the
501 feeding method (76).
502

GL228 [*rrf-3(pk1426)*] II;*daf-2(e1371)* III] eggs were placed in 24-well RNAi plates
seeded with bacteria expressing the dsRNA of interest. Worms were maintained for 5
days at 15ºC until adulthood, then were transferred to an identical 24-well plate to lay
eggs for 5 hours. Adults were removed and the eggs were incubated at 25°C for 4 days
to allow formation of dauers. *daf-16(RNAi)* and the empty vector were used as controls.
Proper dauer formation was assessed by observation in a dissecting microscope and by
1% SDS resistance. RNAi clones that caused abnormal dauer phenotypes were
validated in *daf-2(e1370)* worms. Identity of the dsRNA was confirmed by sequencing
(Macrogen, Korea).
512

daf-9 suppression assays

For preparation of *daf-9*(RNAi) plates, bacterial cultures were grown overnight in LB
with 10 μg/ml tetracycline and 50 μg/ml carbenicillin and induced with 4 mM IPTG
(Isopropyl *β*-d-1-thiogalactopyranoside) for four hours. Cultures were spun down,
suspended in 1/10 volume and 30 μl of bacteria were seeded on 3 cm NGM plates
made with 1 mM IPTG and 50 μg/ml carbenicillin. Plates were grown overnight at 25°C
and stored at 4°C for no more than 2 weeks prior to use.
520

521 L4 stage animals were placed on 3 cm RNAi plates. Two worms per plate were used for
522 *daf-7(e1372)*, while three worms were used for *daf-7; chd-7(gk290)* and *daf-7; chd-*
523 *7(tm6139)*. After 72 hours, the adults were removed and plates were replaced at 25° for
524 2-3 days. The total number of dauers, L4s, and adults were then assessed.

SDS survival assay

Young adults were transferred to seeded plates and permitted to lay for 5 days at 25°C. The arrested progeny were then washed off plates with M9 (22 mM KH_2PO_4, 42mM Na_2HPO_4, 85.5 mM NaCl, 1mM $MgSO_4$) into 15 ml glass conical tubes. Collected animals were washed 2-3 times with M9 and the excess liquid was aspirated off. Animals were then treated with 2 ml of 1% SDS for 30 minutes on a nutator at 25 °C. Following incubation, the samples were washed 3 times with M9 and any excess liquid was aspirated off. Animals were aliquoted to 5 seeded plates with 50-70 worms per plate and allowed to recover at 16°C. The recovered animals were then quantified, and the percentage recovered was calculated. This was repeated 3 times for each strain tested. Comparisons were performed using the χ-squared Test in R; P values were corrected using the Bonferroni method for multiple comparisons.

fog-2 **mating assay**

To determine if males were capable of siring offspring, 8 males from each strain tested were plated with 4 *fog-2(q71)* adult females on seeded 10 cm plates**.** After 24h, *fog-2* females were transferred to new plates and within 48h the proportion of fertile females were scored. Assay was repeated two times.

Lifespan assays

All lifespan experiments were conducted by transferring 1 day-old adults from 15°C to 20°C for the remainder of the lifespan assay. NGM plates were seeded with *E. coli* OP50-1 or, for experiments involving gene silencing, with HT115::DE3. ~150 L4 hermaphrodites were transferred to 5 plates per experiment. Every 48h, animals were scored as alive, dead or censored (animals that exploded, died from bagging or dried out at the edges of the plates). Animals were considered dead when they did not respond to a soft touch to the head with a pick. To prevent the progeny from interfering with the assay, adults were transferred to fresh plates every 48h until egg production ceased. For *glp-1(e2144)* assays, eggs were kept at 20°C for 4h and then transferred at 25.5°C for 72h to induce sterility and switched to 20°C for the remainder of the

556 experiment. Lifespan data were analyzed using the Kaplan-Meier method. Statistics
557 were calculated using the Mantel-Cox nonparametric log-rank method using OASIS2
558 (77).
559

L1 survival
Eggs were obtained by of gravid adults (Worms are mixed by inversion in 5 ml bleach
solution, 3.5 ml ddH20, 1.0 ml 5% NaOCl, 0.5 ml 5 N KOH. After the worms have lysed
and eggs are released, the eggs were centrifuged, the bleach solution aspirated, and
eggs were washed withM9 buffer). After extensive washing, eggs were suspending in
Eggs were suspended in 5ml of sterile M9 with 0.1µg/ml of streptomycin and left to
hatch at 20°C under gentle shaking for 16h. To normalize population density, resulting
L1 larvae were diluted to obtain 1 larva/µl in 5ml of M9 and kept at 20°C under constant
agitation for the remainder of the experiment. Every 48h, a 100µl aliquot of L1 larvae
were spotted onto a seeded NGM plate and then incubated at 20°C for 72h. Animals
were scored as alive that developed beyond the L2 stage. Percentage of the population
alive was normalized to day 1 L1 larvae.
572

Pathogen resistance assay
The pathogenic bacterial strain used in this study was *Pseudomonas aeruginosa* (strain
PA14). This strain was streaked from a frozen stock onto a LB agar plate, incubated at
37°C overnight and then kept at 4°C (shelf-life of one week). For survival assay, a single
PA14 colony was inoculated in King's broth and incubated at 37°C overnight with
shaking. 20µl of this culture was seeded onto slow killing NGM plates (containing 0.35%
peptone instead of 0.25%) and incubated for 24h at 37°C. The plates were then left to
sit at room temperature for 24hrs. The following day, 150 L4 hermaphrodites grown at
15°C were distributed onto 5 PA14 plates and incubated at 25°C. Survival was
monitored at intervals of 6h to 12h and live, dead and censored animals were recorded.
Data and statistics were analyzed using the Kaplan-Meier method as described in the
section "Lifespan assays".
585

Oil-Red-O (ORO) staining

Dauers grown at 25°C for 5 days were stained for lipids using ORO, as previously

described (78). Animals were mounted on a 4% agarose pad and observed in a Zeiss

Axioplan brightfield microscope equipped with a Micropublisher 3.3 camera (Q

Imaging). Image J (NIH) was used to quantify the amount of lipids in each animal. At

least 20 animals of each strain were quantified. Statistical differences were determined

by Student's t test.

594   Microscopy and fluorescence imaging

For imaging dauers and adults, worms were immobilized in a 25mM sodium azide

solution and imaged at 10x and 20x magnification. Images were collected using a Zeiss

Axioplan Imaging Microscope with a DIC system and Zeiss Plan-Neofluar 10x and 20x

objectives lens. Images were acquired with a Micropublisher 3.3 camera (Q Imaging).

ImageJ (NIH) software was used to quantify worm size.

For imaging *pCHD-7::mCherry* fluorescence, worms were immobilized with 1mM

levamisole on fresh 2% agarose pads and imaged immediately using a Nikon A1r

confocal microscope equipped with a 40x PLAN APO oil objective. ImageJ (NIH)

software was used to quantify fluorescence intensity.

For imaging gonads in *daf-2(e1370)* and mutants, young adults were transferred to

seeded plates and permitted to lay eggs for 5 days at 25°C to form dauers, then

collected and fixed with Carnoy's solution (75µl EtOH, 37.5µl Acetate, 12.5µl

Chloroform) and stained with 5mg/ml DAPI (4',6-diamidino-2-phenylindole) in PBS.

Animals were imaged as 0.5µm Z-stacks with the Nikon A1r Confocal Microscope with

40x and 60x plan APO oil objectives.

For imaging *ajm-1::GFP*, young adults were transferred to seeded plates and permitted

to lay eggs for 5 days at 25°C to form dauers. The arrested progeny was then washed

off plates with M9 into 15 ml glass conical tubes. Collected animals were washed 2-3

times with M9 and the excess liquid was aspirated off. Animals were then immobilized

with levamisole and the seam cells were imaged as Z-stacks by confocal microscopy as

described immediately above.

For scanning electron microscopy, worms were fixed by immersing in 2.5%

glutaraldehyde in PBS for several hours. Worms were washed 3X in PBS then post

fixed in aqueous OsO 4 for 1h, then washed 3x in PBS, dehydrated through a graded

series of ethanols (30-100%) then chemically dried with 2 x 10 min incubations in

hexamethyldisilazane. Dried worms were sprinkled onto copper double stick tape on

aluminum stubs, sputter coated with 3.5 nm gold-palladium alloy then evaluated on a

JEOL JEM 6335F microscope at 5 kV.

Library preparation and RNA-seq

daf-2(e1370) and *chd-7(gk290);daf-2(e1370)*, synchronized eggs were kept at 25°C for

10 days and resulting dauers were collected and frozen. Total RNA was extracted with

TRIzol (Invitrogen) following the kit´s protocol. cDNA library was prepared with NEBNext

Ultra II RNA library prep kit for Illumina (New England Biolabs), and the sequencing

carried out using Illumina's HiSeq-2500 sequencer with single-end mode and read

length of 50 bp. Five replicates for *daf-2(e1370)* vs. *chd-7(gk290);daf-2(e1370)* were

sequenced. For data assessment, a quality control with FastQC software (version

0.11.5) was used. First the raw reads that aligned against the *E. coli* genome (K12

genome) were removed. The remaining sequences were aligned against the reference

genome of *C. elegans* WS260 using STAR (version 2.5.4a). The number of mapped

reads to genes was counted using Htseq (version 0.9.1). Finally, the DEGs were

determined using DESeq2 (version 1.20.0) with a cutoff of 0.05 on False Discovery

Rate (FDR). R version 3.5.0 (2018-04-23) and Bioconductor version 3.7 with

BiocInstaller version 1.30.0 were used. Heatmaps were generated using pheatmap

package (version 1.0.12) with hierarchical clustering on the rows with the default

options.

Genomic Sequencing of *scd-3*

A 6cm plate replete with *scd-3* gravid animals was washed with 1ml M9 and collected in

a glass conical. Worms were washed extensively, then placed on the nutator for 1 hour

to allow remaining gut bacteria to be passed into the medium. Worms were again

washed 3-4 times and settled by gravity. The worm pellet was transferred to 1.5ml

Eppendorf and genomic DNA was prepared according to the manufacturer's protocol

with the Purelink Genomic DNA Kit (Invitrogen) except that after addition of digestion

buffer, worms were pulverized with a microfuge hand dounce prior to incubation at

55°C. Sequencing was performed by Psomagen, Qiagen CLC Genomics Workbench

was used to align the DNA against WBcel235 and view the variant.

Xenopus laevis embryo manipulation and microinjections

Xenopus embryos were obtained by natural mating. Adult frogs' reproductive behavior

was induced by injection of human chorionic gonadotropin hormone. Eggs were

collected, de-jellied in 3% cysteine (pH 8.0), maintained in 0.1 X Marc's Modified

Ringer's (MMR) solution and staged according to Nieuwkoop and Faber (79). The

embryos were placed in 3% ficoll prepared in 1 X MMR for microinjection. Chd7

morpholino (*chd7*-MO: 5′-AACTCATCATGCCAGGGTCTGCCAT-3′) (Gene Tools, LLC)

specificity has been previously characterized (Bajpai 2010). The cDNA of *Xenopus*

laevis Col2a1 was amplified by PCR from pCMV-Sport 6-*col2a1* (Dharmacon) with

primers M13F and M13R (see Supplemental Table 2). The PCR fragment was digested

with *EcoR*V and *Not*I and cloned into pCS2+ previously digested with *Stu*I and *Not*I.

Capped mRNAs for *col2a1* were transcribed *in vitro* with SP6 using the mMessage

mMachine kit (Ambion) following linearization with *Not*I. *Chd7*-MO and *col2a1* mRNA

were injected into both D1 blastomeres of 8-cell staged embryos (80) for lethality and

morphometrics analysis. *Chd7*-MO was injected into one D1 blastomeres of 8-cell

staged embryos for analysis of *col2a1* expression.

Whole-mount *in situ* hybridization was carried out as previously described (81). pCMV-

Sport 6-*col2a1* (Dharmacon) construct was linearized with *Sal*I and transcribed with T7

for antisense probe synthesis. Images of whole embryos were collected with a Leica

DFC420 camera attached to a Leica L2 stereoscope. Morphometics analysis were done
on fixed embryos using ImageJ software (https://fiji.sc/).
679

RNA preparation and RT-qPCR
For RNA extraction, 2-cell stage embryos were injected into both blastomeres with 5 ng
of *chd7*-MO, collected at stage 23 and snap-frozen for later processing. Six embryos
were combined for each treatment and RNA was isolated using RNAzol (Sigma-
Aldrich). RNA concentration of each sample was measured using a Nanodrop
spectrophotometer, and 250 ng of RNA were reverse transcribed using qScript cDNA
SuperMix (QuantaBio). Quantitative PCR (qPCR) was preformed using the Forget-Me-
Not qPCR Master Mix (Biotium) with a Biorad iCycler thermocycler. Amplification was
performed using oligonucleotides designed with PerlPrimer software for *col2a1*
(Forward 5′TCCCTGTTGATGTTGAAGCC3′; Reverse 5′-
CAATAGTCACCGCTCTTCCA-3′ and ODC primers have been previously described
(82) (Forward 5′CAAAGCTTGTTCTACGCATAGCA3′; Reverse
5′GGTGGCACCAAATTTCACACT 3′). The relative expression of *col2a1* was
normalized to *ODC* expression and determined via the method previously described
(83).
695

Ethics Statement
Xenopus laevis experiments were carried out in strict accordance with the
recommendations in the Guide for the Care and Use of Laboratory Animals of the NIH
and also the ARRIVE guidelines. The animal care protocol was approved by the
Comisión Institucional para el Cuidado y Uso de Animales de Laboratorio (CICUAL) of
the School of Applied and Natural Sciences, University of Buenos Aires, Argentina
(Protocol #64).

Acknowledgments

The authors would like to thank Bruno Moretti, Hernan Grecco and Mario Rossi for experimental support. We also thank the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET). Diego Martín Jofré is a CONCIET fellow and Daniel Hochbaum is a career investigator from CONICET. DH laboratory was supported by the Agencia Nacional de Promoción Científica y Tecnológica of Argentina (PICT-2016-0269) and Conicet (PIP 1122015 0100731 CO). Funding for this work was also provided by grants from the CHARGE Syndrome Foundation (JLY and DH), The Company of Biologists (MF), NIGMS (R01GM104007 JLY), and NIA (R01AG051659 AG). This work was also supported, in part, by the Intramural Research Program of the NIH and the National Institute of Diabetes and Digestive and Kidney Diseases (SY).

Author roles and contributions

Conceived and designed the experiments: DMJ, DKH, ASC, FAG, ES, MCC, JLY, DH. Performed the experiments: DMJ, DKH, ASC, MG, FRGA, DBS, ES, FAR, DH. Analyzed the data: DMJ, DKH, ASC, SY, FRGA, AG, MCC, JLY, DH. Drafted the manuscript: JLY,DH. Reviewed and edited the manuscript: DMJ, DKH, AG, MCC, JLY, DH.

REFERENCES

1.   L. E. L. M. Vissers, *et al.*, Mutations in a new member of the chromodomain gene family cause CHARGE syndrome. *Nature genetics* **36**, 955–7 (2004).

2.   R. Balasubramanian, *et al.*, Functionally compromised CHD7 alleles in patients with isolated GnRH deficiency. *Proc Natl Acad Sci U S A* **111**, 17953–17958 (2014).

3.   H.-G. Kim, *et al.*, Mutations in CHD7, encoding a chromatin-remodeling protein, cause idiopathic hypogonadotropic hypogonadism and Kallmann syndrome. *Am. J. Hum. Genet.* **83**, 511–519 (2008).

4.   R. Bernier, *et al.*, Disruptive CHD8 Mutations Define a Subtype of Autism Early in Development. *Cell* **158**, 263–276 (2014).

5.   C. G. A. Marfella, A. N. Imbalzano, The Chd family of chromatin remodelers. *Mutation research* **618**, 30–40 (2007).

6.   S. A. Patten, *et al.*, Role of Chd7 in Zebrafish: A Model for CHARGE Syndrome. *PLoS One* **7** (2012).

7.   E. A. Bosman, *et al.*, Multiple mutations in mouse Chd7 provide models for CHARGE syndrome. *Hum Mol Genet* **14**, 3463–3476 (2005).

8.   C. Tian, *et al.*, Otitis media in a new mouse model for CHARGE syndrome with a deletion in the Chd7 gene. *PloS one* **7**, e34944 (2012).

9.   G. Daubresse, *et al.*, The Drosophila kismet gene is related to chromatin-remodeling factors and is required for both segmentation and segment identity. *Development* **126**, 1175–1187 (1999).

10.   D. J. Melicharek, L. C. Ramirez, S. Singh, R. Thompson, D. R. Marenda, Kismet/CHD7 regulates axon morphology, memory and locomotion in a Drosophila model of CHARGE syndrome. *Hum Mol Genet* **19**, 4253–4264 (2010).

11.   Z. Asad, *et al.*, Rescue of neural crest-derived phenotypes in a zebrafish CHARGE model by Sox10 downregulation. *Hum Mol Genet* **25**, 3539–3554 (2016).

12.   C. Kenyon, J. Chang, E. Gensch, A. Rudner, R. Tabtiang, A C. elegans mutant that lives twice as long as wild type. *Nature* **366**, 461–464 (1993).

13.   P. L. Larsen, Aging and resistance to oxidative damage in Caenorhabditis elegans. *Proc Natl Acad Sci U S A* **90**, 8905–8909 (1993).

14. M. M. Gaglia, C. Kenyon, Stimulation of Movement in a Quiescent, Hibernation-
Like Form of Caenorhabditis elegans by Dopamine Signaling. *J Neurosci* **29**, 7302–
7314 (2009).

15. J. Wang, S. K. Kim, Global analysis of dauer gene expression in Caenorhabditis
elegans. *Development* **130**, 1621–1634 (2003).

16. E. Cohen, J. Bieschke, R. M. Perciavalle, J. W. Kelly, A. Dillin, Opposing
activities protect against age-onset proteotoxicity. *Science* **313**, 1604–1610 (2006).

17. A. Meléndez, *et al.*, Autophagy genes are essential for dauer development and
life-span extension in C. elegans. *Science* **301**, 1387–1391 (2003).

18. R. Christensen, L. de la Torre-Ubieta, A. Bonni, D. A. Colón-Ramos, A conserved
PTEN/FOXO pathway regulates neuronal morphology during C. elegans development.
Development **138**, 5257–5267 (2011).

19. C. A. Wolkow, K. D. Kimura, M.-S. Lee, G. Ruvkun, Regulation of C. elegans
Life-Span by Insulinlike Signaling in the Nervous System. *Science* **290**, 147–150 (2000).

20. R. C. Cassada, R. L. Russell, The dauerlarva, a post-embryonic developmental
variant of the nematode Caenorhabditis elegans. *Dev Biol* **46**, 326–342 (1975).

21. D. L. Riddle, P. S. Albert, "Genetic and Environmental Regulation of Dauer Larva
Development" in *C. Elegans II*, 2nd Ed., D. L. Riddle, T. Blumenthal, B. J. Meyer, J. R.
Priess, Eds. (Cold Spring Harbor Laboratory Press, 1997) (September 25, 2017).

22. P. Ren, *et al.*, Control of C. elegans larval development by neuronal expression
of a TGF-beta homolog. *Science* **274**, 1389–1391 (1996).

23. D. Park, A. Estevez, D. L. Riddle, Antagonistic Smad transcription factors control
the dauer/non-dauer switch in C. elegans. *Development* **137**, 477–485 (2010).

24. L. S. da Graca, DAF-5 is a Ski oncoprotein homolog that functions in a neuronal
TGF pathway to regulate C. elegans dauer development. *Development* **131**, 435–446
(2003).

25. J. H. Thomas, D. A. Birnby, J. J. Vowels, Evidence for parallel processing of
sensory information controlling dauer formation in Caenorhabditis elegans. *Genetics*
134, 1105–1117 (1993).

26. W. M. Shaw, S. Luo, J. Landis, J. Ashraf, C. T. Murphy, The C. elegans TGF-β
Dauer Pathway Regulates Longevity via Insulin Signaling. *Current Biology* **17**, 1635–

1645 (2007).

27.   T. Liu, K. K. Zimmerman, G. I. Patterson, Regulation of signaling genes by

TGFbeta during entry into dauer diapause in C. elegans. *BMC Dev Biol* **4**, 11 (2004).

28.   S. D. Narasimhan, *et al.*, PDP-1 Links the TGF-β and IIS Pathways to Regulate

Longevity, Development, and Metabolism. *PLoS Genet* **7** (2011).

29.   T. Inoue, J. H. Thomas, Suppressors of transforming growth factor-beta pathway

mutants in the Caenorhabditis elegans dauer formation pathway. *Genetics* **156**, 1035–

1046 (2000).

30.   B. Gerisch, A. Antebi, Hormonal signals produced by DAF-9/cytochrome P450

regulate C. elegans dauer diapause in response to environmental cues. *Development*

131, 1765–1776 (2004).

31.   H. Y. Mak, G. Ruvkun, Intercellular signaling of reproductive development by the

C. elegans DAF-9 cytochrome P450. *Development* **131**, 1777–1786 (2004).

32.   D. Hochbaum, *et al.*, DAF-12 regulates a connected network of genes to ensure

robust developmental decisions. *PLoS Genet* **7**, e1002179 (2011).

33.   N. Fielenbach, A. Antebi, C. elegans dauer formation and the molecular basis of

plasticity. *Genes Dev* **22**, 2149–2165 (2008).

34.   R. A. Ignotz, J. Massagué, Transforming growth factor-beta stimulates the

expression of fibronectin and collagen and their incorporation into the extracellular

matrix. *J. Biol. Chem.* **261**, 4337–4345 (1986).

35.   R. Bajpai, *et al.*, CHD7 cooperates with PBAF to control multipotent neural crest

formation. *Nature* **463**, 958–962 (2010).

36.   A. Dubey, J.-P. Saint-Jeannet, Modeling Human Craniofacial Disorders in

Xenopus. *Curr Pathobiol Rep* **5**, 79–92 (2017).

37.   D. W. Seufert, J. Hanken, M. W. Klymkowsky, Type II collagen distribution during

cranial development in Xenopus laevis. *Anat. Embryol.* **189**, 81–89 (1994).

38.   P. S. Albert, D. L. Riddle, Mutants of Caenorhabditis elegans that form dauer-like

larvae. *Dev Biol* **126**, 270–293 (1988).

39.   E. S. Haag, Dial-a-mutant: web-based knockout collections for model organisms.

Biol. Cell **99**, 343–347 (2007).

40. ,   Dauer Epithelium (March 24, 2021).

41. P. Kadekar, R. Roy, AMPK regulates germline stem cell quiescence and integrity

through an endogenous small RNA pathway. *PLoS Biol* **17** (2019).

42. H. Hsin, C. Kenyon, Signals from the reproductive system regulate the lifespan of

C. elegans. *Nature* **399**, 362–366 (1999).

43. D. A. Garsin, *et al.*, Long-lived C. elegans daf-2 mutants are resistant to bacterial

pathogens. *Science* **300**, 1921 (2003).

44. I. Lee, A. Hendrix, J. Kim, J. Yoshimoto, Y.-J. You, Metabolic rate regulates L1

longevity in C. elegans. *PLoS One* **7**, e44720 (2012).

45. M. Ailion, J. H. Thomas, Dauer Formation Induced by High Temperatures in

Caenorhabditis elegans. *Genetics* **156**, 1047–1067 (2000).

46. C. Savage, *et al.*, Caenorhabditis elegans genes sma-2, sma-3, and sma-4

define a conserved family of transforming growth factor beta pathway components. *Proc*

Natl Acad Sci U S A **93**, 790–794 (1996).

47. Y. Suzuki, *et al.*, A BMP homolog acts as a dose-dependent regulator of body

size and male tail patterning in Caenorhabditis elegans. *Development* **126**, 241–250

(1999).

48. B. Gerisch, C. Weitzel, C. Kober-Eisermann, V. Rottiers, A. Antebi, A hormonal

signaling pathway influencing C. elegans metabolism, reproductive development, and

life span. *Dev Cell* **1**, 841–851 (2001).

49. R. P. Boot-Handford, D. S. Tuckwell, Fibrillar collagen: the key to vertebrate

evolution? A tale of molecular incest. *Bioessays* **25**, 142–151 (2003).

50. W. R. Wilcox, Connective Tissue and Its Heritable Disorders: Molecular, Genetic,

and Medical Aspects,. *Am J Hum Genet* **72**, 503–504 (2003).

51. A. Wood, D. E. Ashhurst, A. Corbett, P. Thorogood, The transient expression of

type II collagen at tissue interfaces during mammalian craniofacial development.

Development **111**, 955–968 (1991).

52. N. Garamszegi, *et al.*, Extracellular matrix-induced transforming growth factor-

beta receptor signaling dynamics. *Oncogene* **29**, 2368–2380 (2010).

53. W. Xin, J. Heilig, M. Paulsson, F. Zaucke, Collagen II regulates chondroycte

integrin expression profile and differentiation. *Connect Tissue Res* **56**, 307–314 (2015).

54. , Differential effect of ECM molecules on re-expression of cartilaginous markers in

near quiescent human chondrocytes - Chiu - 2011 - Journal of Cellular Physiology -
Wiley Online Library (March 24, 2021).

55. R. Kerney, B. K. Hall, J. Hanken, Regulatory elements of Xenopus col2a1 drive
cartilaginous gene expression in transgenic frogs. *Int J Dev Biol* **54**, 141–150 (2010).

56. J. R. Siebert, J. M. Graham, C. MacDonald, Pathologic features of the CHARGE
association: Support for involvement of the neural crest. *Teratology* **31**, 331–336 (1985).

57. A. Antebi, W. H. Yeh, D. Tait, E. M. Hedgecock, D. L. Riddle, daf-12 encodes a
nuclear receptor that regulates the dauer diapause and developmental age in C.
elegans. *Genes Dev* **14**, 1512–1527 (2000).

58. C. L. Hale, A. N. Niederriter, G. E. Green, D. M. Martin, Atypical phenotypes
associated with pathogenic CHD7 variants and a proposal for broadening CHARGE
syndrome clinical diagnostic criteria. *Am. J. Med. Genet. A* **170**, 344–354 (2016).

59. X. Chu, *et al.*, Genotranscriptomic meta-analysis of the CHD family chromatin
remodelers in human cancers – initial evidence of an oncogenic role for CHD7. *Mol*
Oncol **11**, 1348–1360 (2017).

60. Y. Liu, *et al.*, CHD7 interacts with BMP R-SMADs to epigenetically regulate
cardiogenesis in mice. *Human Molecular Genetics* **23**, 2145–2156 (2014).

61. M. Tewari, *et al.*, Systematic Interactome Mapping and Genetic Perturbation
Analysis of a C. elegans TGF-$_{\beta}$ Signaling Network. *Molecular Cell*, 14.

62. C. G. Riedel, *et al.*, DAF-16 employs the chromatin remodeller SWI/SNF to
promote stress resistance and longevity. *Nat. Cell Biol.* **15**, 491–501 (2013).

63. K. Jia, P. S. Albert, D. L. Riddle, DAF-9, a cytochrome P450 regulating C.
elegans larval development and adult longevity. *Development* **129**, 221–231 (2002).

64. P. J. Hu, J. Xu, G. Ruvkun, Two membrane-associated tyrosine phosphatase
homologs potentiate C. elegans AKT-1/PKB signaling. *PLoS Genet.* **2**, e99 (2006).

65. S. E. Celniker, *et al.*, Unlocking the secrets of the genome. *Nature* **459**, 927–930
(2009).

66. C. Savage, *et al.*, Caenorhabditis elegans genes sma-2, sma-3, and sma-4
define a conserved family of transforming growth factor β pathway components.
Developmental Biology, 5 (1996).

67. Y. Suzuki, G. A. Morris, M. Han, W. B. Wood, A Cuticle Collagen Encoded by the

lon-3 Gene May Be a Target of TGF-β Signaling in Determining Caenorhabditis elegans

Body Shape. *Genetics* **162**, 1631–1639 (2002).

68.   X. Huang, J.-P. Saint-Jeannet, Induction of the neural crest and the opportunities

of life on the edge. *Developmental Biology* **275**, 1–11 (2004).

69.   H. Okuno, *et al.*, CHARGE syndrome modeling using patient-iPSCs reveals

defective migration of neural crest cells harboring CHD7 mutations. *Elife* **6** (2017).

70.   J. M. Kramer, "Extracellular Matrix" in *C. Elegans II*, 2nd Ed., D. L. Riddle, T.

Blumenthal, B. J. Meyer, J. R. Priess, Eds. (Cold Spring Harbor Laboratory Press,

1997) (March 22, 2021).

71.   T. Suzuki, D. Sakai, N. Osumi, H. Wada, Y. Wakamatsu, Sox genes regulate

type 2 collagen expression in avian neural crest cells. *Dev Growth Differ* **48**, 477–486

(2006).

72.   D. He, *et al.*, Chd7 cooperates with Sox10 and regulates the onset of CNS

myelination and remyelination. *Nature Neuroscience* **19**, 678–689 (2016).

73.   A. B. Roberts, B. K. McCune, M. B. Sporn, TGF-β: Regulation of extracellular

matrix. *Kidney International* **41**, 557–559 (1992).

74.   A. Tekari, R. Luginbuehl, W. Hofstetter, R. J. Egli, Transforming Growth Factor

Beta Signaling Is Essential for the Autonomous Formation of Cartilage-Like Tissue by

Expanded Chondrocytes. *PLOS ONE* **10**, e0120857 (2015).

75.   R. S. Kamath, J. Ahringer, Genome-wide RNAi screening in Caenorhabditis

elegans. *Methods* **30**, 313–321 (2003).

76.   L. Timmons, D. L. Court, A. Fire, Ingestion of bacterially expressed dsRNAs can

produce specific and potent genetic interference in Caenorhabditis elegans. *Gene* **263**,

103–112 (2001).

77.   S. K. Han, *et al.*, OASIS 2: Online application for survival analysis 2 with features

for the analysis of maximal lifespan and healthspan in aging research. *Oncotarget* **7**,

56147–56152 (2016).

78.   R. Ratnappan, *et al.*, Germline Signals Deploy NHR-49 to Modulate Fatty-Acid β-

Oxidation and Desaturation in Somatic Tissues of C. elegans. *PLoS Genetics* **10**

(2014).

79. ,   Normal Table of Xenopus Laevis (Daudin) | Taylor & Francis Group (March 23,

2021).

80. ,   Blastomeres show differential fate changes in 8-cell Xenopus laevis embryos

that are rotated 90° before first cleavage - Huang - 1998 - Development, Growth &

Differentiation - Wiley Online Library (March 23, 2021).

81.   V. Gawantka, *et al.*, Gene expression screening in Xenopus identifies molecular

pathways, predicts gene function and provides a global view of embryonic patterning.

Mechanisms of Development **77**, 95–141 (1998).

82.   Q. Cao, Y. Shen, W. Zheng, H. Liu, C. Liu, Tcf7l1 promotes transcription of

Kruppel-like factor 4 during Xenopus embryogenesis. *J Biomed Res* **32**, 215–221

(2018).

83.   M. W. Pfaffl, A new mathematical model for relative quantification in real-time

RT-PCR. *Nucleic Acids Res* **29**, e45 (2001).

Figure 1: *chd-7*, a CHD superfamily member and a *daf-12* controlled gene, regulates *daf-2* dauer formation. A) *chd-7(RNAi)* causes a partial dauer phenotype in *daf-2(e1370)*. Representative DIC photomicrographs of *daf-2(e1370)* normal dauers fed on vector control RNAi at 25°C or the partial dauer phenotype when worms are fed on *chd-7(RNAi)*. **B)** Quantification of the axial ratio. Three biological replicates were scored with 16-21 individual dauers per replicate. Horizontal black lines represent mean with SD. *p<0.05. **C)** *daf-12* regulates *chd-7* expression. Representative images of *tag-192::H1-wCherry* or *tag-192::H1-wCherry; daf-12(rh61rh411)* worms in L2/L3 stage. **D)** Relative expression of the

translational reporter (n>10/strain). Unpaired t test, ****p<0.0001. **E)** *C.elegans chd-7* gene and proteins. Top: *chd-7* genomic region. UTR and exons shown as bars; introns by lines. In red, available *chd-7* deletional alleles (data obtained from CGC and NBRP). Signature domains in CHD proteins**:** two N-terminal chromodomains for interaction with a variety of chromatin components (green), a SNF-2 like domain with ATPase activity (yellow) and a Helicase domain (blue). The Class III subfamily is defined by a BRK domain (purple). Bottom images depict the predicted protein isoforms of *C. elegans* CHD-7, human CHD7 and human CHD8. **F)** *chd-7*(*gk290*);*daf-2* and *chd-7*(*tm6139*);*daf-2* develop as SDS sensitive dauer larvae*.* Each dot represents one plate with at least 40 arrested animals per plate and a minimum of 725 total animals per strain tested. Horizontal black lines represent mean percentage with SD. Statistical significance was calculated using Chi-squared test with Bonferroni correction for multiple comparisons. ****p<0.0001. **G)** *chd-7* partial dauers fail to develop the dauer alae. Top row, representative photomicrographs of *daf-2(e1370)* dauers or partial dauers expressing the *ajm-1*::GFP reporter. Arrowheads indicate seam cell junctions. Bottom row, SEM images showing alae details. **H)** Dauer germ cells in *chd-7* mutants keep proliferating. (Top to Bottom) *daf-2(e1370), chd-7(gk290);daf-2, chd-7(tm6139);daf-2.* Representative z-projections for dauer animals stained with DAPI (green). Arrowheads denote the left and right most distal end of the gonad.

Figure 2: *chd-7* affects longevity and immune response. A-E) *chd-7* promotes longevity in *wild-type*, *daf-2(e1370)* and *glp-1(e2144)* mutants. Mean survival days on OP50-1 **A)** WT (18.12), *chd-7(gk290)* (14.91), *chd-7(gk306)* (16.53) **B)** WT (17.85), CHD-7::GFP (14.35) **C)** *daf-2(e1370)* (35.58), *chd-7(gk306);daf-2* (37.18), *chd-7(gk290);daf-2* (17.47), *daf-16(mu86);daf-2(e1370)* (18.85). **D)** *daf-2(e1370)* (27.9), *chd-7(gk290);daf-2;CHD-7::GFP* (30.84), *chd-7(gk290);daf-2* (19.13), *daf-16(mu86);daf-2* (14.48). **E)** *glp-1(e2144)* (30.4), *chd-7(gk290);glp-1* (28.37), *glp-1;daf-12(rh61rh411)* (25.99). Number of animals and other data from replicate found in Supplementary Table 2. **F)** *chd-7* mediates the immune response against the opportunistic bacteria *P. aeruginos*a. Mean lifespan in hours (m) ± standard error of the mean (SEM). 'n' is the number of animals analyzed/total number in experiment. N2 (m = 52.11 ± 0.96, n = 94/162), *chd-7(gk290)* (m = 47.45 ± 1.05, n = 54.130/), *chd-7(tm6139)* (m = 44.47 ± 1.57, n = 56/80), *daf-2(e1370)* (m = 105.46 ± 5.61, n = 65/115), *chd-7(gk290);daf-2* (m = 86.74 ± 4.21, n = 103/120) and *chd-7(tm6139);daf-2* (m = 79.13 ± 4.9, n = 50/60). In A-F) survival data analyzed using Kaplan−Meier test. *p<0.05 (*), ***p <0.001 and ****p <0.0001. Asterisk´s color represents the strain of reference.

Figure 3: CHD-7 belongs to the TGF-β signaling pathway. A-B) Dauer larva formation of *daf-c;chd-7* mutants. 7 individual L4's were grown on 7 plates at the specified temperature for 1 week and the number of arrested and non-arrested progeny was scored. Each dot represents a plate with at least one progeny and minimum 44 totals animals quantified for each strain. Horizontal black lines represent mean percentage with SD. Statistical significance was calculated using Chi-squared test with Bonferroni correction for multiple comparisons. ****p<0.0001. **C)** CHD-7 regulates body size. Synchronized worms were grown at 20°C until adulthood and length of at least 16 individuals was scored. Statistical significance was calculated using one-way Anova. **p<0.01, ***p<0.001. **D)** *chd-7(tm6139)* males do not mate. 8 male animals from each strain tested were plated with 4 *fog-2* females on 10 cm plates. After 24 hrs *fog-2* females were transferred to new plates and within 48h the proportion of fertile females were scored. Assay was repeated two times.

Figure 4: RNA-Seq analysis of transcriptome changes in dauers induced by *chd-7(gk290)* mutant. A) Heat map of individual expression values of all 84 differentially expressed genes (DEGs, cutoff of 0.05 on FDR). Prior to heat map calculation, DEGs were determined using DESeq2 (version 1.20.0). The color scale represents row z-score. Hierarchical clustering of the DEGs is represented by dendrograms at the left. **B)** Partial dauers exhibit an increased number of reads of *daf-9. daf-9* transcripts per million (TPM) were generated and summarized to gene level using Qiagen's CLC Genomics

Workbench Software. Bars represent the SD and Mean. **C)** *daf-9* knockdown in *chd-7*;*daf-7(e1372)* animals rescues dauer arrest at 25°C. 2-3 young L4's were plated on to freshly seeded plates with either *daf-9* or L4440 empty vector RNAi and allowed to lay eggs. After 72h, the adults were removed and the proportion of progeny that arrested as dauers was calculated. Each dot represents a plate of at least 34 animals and a minimum of 598 total animals per strain tested. Horizontal black lines represent mean with SD. Statistical significant differences were determined by one-way ANOVA. **D)** *daf-9* RNAi rescues arrest in *chd-7*;*daf-7(e1372)* animals but leads to incomplete dauers. Arrested animals grown at 25°C on *daf-9* RNAi plates were treated with 1% SDS for 30 min and survival was scored. Each dot represents a plate of at least 30 animals and a minimum of 159 total animals per strain tested. Horizontal black lines represent mean with SD. Statistical significant differences were determined by one-way ANOVA. **E)** Mechanism of action for *chd-7* in dauer development. In dauering conditions, CHD-7 forms a transcriptional complex with DAF-3/DAF-5 to repress *daf-9* expression. In *chd-7* mutants, *daf-9* expression prevents dauer formation driven by reduced activity of both DAF-7/TGF-β and DAF-2/IIS signaling pathways, presumably by binding dafachronic acids (DA) and subsequent activation of DAF-12.

Figure 5: Expression of *col2a1* rescues Chd-7 knock-down in *Xenopus* embryos. A) Expression levels of *col2a1* mRNA are reduced in *chd7*-MO injected embryos. 2-cell staged embryos were injected into both blastomeres with 5 ng of *chd7*-MO. Error bars indicate standard error from two repeats of the PCR reaction with different biological samples. Statistical significant differences were determined by a one-tailed paired t-test (* $p < 0.05$). **B)** *Col2a1* expression is reduced in the branchyal arches of Chd7-depleted embryos. *In situ* hybridization of stage 35 embryos for *col2a1*, lateral views. 8-cell staged embryos were injected into one D1 blastomeres with 2.5 ng of *chd7*-MO. Left and right panels are control and injected sides of the same embryo, respectively. Reduced staining of the branchyal arches (asterisks) was observed in 56% of the injected embryos (N = 1, 22) (N = number of experiments, number of embryos). ov: otic vesicle. Representative embryo is shown. **C)** Lethality in Chd-7-depleted embryos is rescued by over-expression of *col2a1* in *Xenopus*. Bar graph with the percentage of dead embryos by stage 45. Higher dosage of *chd7*-MO causes an increase percentage in the number of dead embryos and co-injection of *col2a1* mRNA reduces embryonic lethality. 8-cell staged embryos were injected into both D1 blastomeres with *chd7*-MO and/or *col2a1* mRNA at the indicated doses and scored for survival at stage 45. Uninjected (N = 3, 84); *chd7*-MO 5 ng (N = 2, 64); *chd7*-MO 5 ng + *col2a1* mRNA 50 pg (N = 2, 61); *chd7*-MO 10 ng (N = 1, 33); *chd7*-MO 10 ng + *col2a1* mRNA 50 pg (N = 1, 31) and *col2a1* mRNA 50 pg (N = 2, 58). (N = number of experiments, number of embryos). **D)** Quantification of the eye area of *Xenopus* stage 45 embryos. 8-cell staged embryos were injected into both D1 blastomeres with *chd7*-MO or co-injected with *chd7*-MO and *col2a1* mRNA at the indicated doses. Means and standard deviation are indicated. Each dot represents a single embryo. Uninjected (N = 1, 27); *chd7*-MO 5 ng (N = 1, 24) and *chd7*-MO 5 ng + *col2a1* mRNA 50 pg (N = 1, 33). Statistical significant differences were determined by a one-way ANOVA and means between groups were compared using Tukey's multiple comparisons test (** $p < 0.01$). **E)** Quantification of the eye distance of *Xenopus* stage 45 embryos. 8-cell staged embryos were injected into both D1 blastomeres with *chd7*-MO or co-injected with *chd7*-MO and *col2a1* mRNA at the indicated doses. Means and standard deviation are indicated. Each dot represents a single embryo. Uninjected (N = 1, 27); *chd7*-MO 5 ng (N = 2,

35) and *chd7*-MO 5 ng + *col2a1* mRNA 50 pg (N = 1, 33). Statistical significant differences were determined by a one-way ANOVA and means between groups were compared using Tukey's multiple comparisons test (**** $P < 0.0001$, #$P < 0.05$). * represents the comparison to the uninjected group and # represents the comparison to the *chd7*-MO injected group.