Analysis of CHD-7 defective dauer nematodes implicates collagen misregulation in CHARGE

syndrome features.

Diego M. Jofré[1*], Dane K. Hoffman[2*§], Ailen S. Cervino[3], McKenzie Grundy[2], Sijung Yun[4], Francis RG.

Amrit[5], Donna B. Stolz[6], Esteban Salvatore[7], Fabiana A. Rossi[8], Arjumand Ghazi[5,9], M. Cecilia Cirio[3],

Judith L. Yanowitz[2,10#], Daniel Hochbaum[1#]

*Authors contributed equally.

#Co-corresponding authors.

[1] Departamento de Biodiversidad y Biología Experimental, Facultad de Ciencias Exactas y

Naturales, Universidad de Buenos Aires (UBA). Buenos Aires, Argentina.

[2] Magee-Womens Research Institute. Pittsburgh, PA, USA.

[3] Instituto de Fisiología, Biología Molecular y Neurociencias (IFIBYNE-UBA-CONICET), Facultad de

Ciencias Exactas y Naturales, Universidad de Buenos Aires (UBA). Buenos Aires, Argentina.

[4] Laboratory of Molecular Biology, National Institute of Diabetes and Digestive and Kidney Diseases,

National Institutes of Health. Bethesda, Maryland, USA.

[5] Department of Pediatrics, University of Pittsburgh School of Medicine. Pittsburgh, PA, USA.

[6] Center for Biologic Imaging, University of Pittsburgh Medical School. Pittsburgh, PA, USA.

[7] mAbxience S.A.U. Buenos Aires, Argentina.

[8] Instituto de Investigaciones en Medicina Traslacional (IIMT) - CONICET, Universidad Austral. Pilar,

Buenos Aires, Argentina.

[9] Departments of Developmental Biology and Cell Biology & Physiology, University of Pittsburgh School of

Medicine. Pittsburgh, PA, USA.

[10] Departments of Obstetrics, Gynecology & Reproductive Sciences; Developmental Biology; Microbiology

and Molecular Genetics at he Hillman Cancer Center, University of Pittsburgh. Pittsburgh, PA, USA.

§Current Address: Department of Cancer & Cell Biology. Baylor College of Medicine. Houston, Texas,

USA.

*Correspondence: Correspondence should be addressed to

Dr. Daniel Hochbaum

Ciudad Universitaria. Intendente Guiraldes 2160, 4 piso. Laboratorio 79

CABA, Buenos Aires.

C1428BGA

Argentina

hochbaumd@bg.fcen.uba.ar

Dr. Judith L. Yanowitz

Magee-Womens Research Institute

204 Craft Avenue

Pittsburgh, PA

15213

United States of America

yanowitzjl@mwri.magee.edu

Author Contributions

Conceived and designed the experiments: DMJ, DKH, ASC, FAG, ES, MCC, JLY, DH. Performed the

experiments: DMJ, DKH, ASC, MG, FRGA, DBS, ES, FAR, DH. Analyzed the data: DMJ, DKH, ASC, SY,

FRGA, AG, MCC, JLY, DH. Drafted the manuscript: MCC, JLY, DH. Reviewed and edited the manuscript:

DMJ, DKH, AG, MCC, JLY, DH.

Conflict of Interest

The authors declare no conflicts of interest

Classification: Biological Sciences, Developmental Biology.

Keywords: CHARGE Syndrome, *C. elegans*, dauer, TGF-β, Col2a1

This PDF file includes:

Main Text

Figures 1 to 5

Supplementary figures 1-6 and Supplementary Tables S1-S4

ABSTRACT

CHARGE syndrome is a complex developmental disorder caused by mutations in the

chromodomain helicase DNA-binding protein7 (CHD7) and characterized by retarded growth and

malformations in the heart and nervous system. Despite the public health relevance of this disorder,

relevant targets of CHD7 that relate to disease pathology are still poorly understood. Here we report that

chd-7, the nematode ortholog of Chd7, is required for dauer morphogenesis, lifespan determination, and

stress response. Consistent with our discoveries, we found *chd-7* to be allelic to *scd-3*, a previously

identified dauer suppressor from the TGF-β pathway. Notably, DAF-12 promoted *chd*-7 expression, which

is necessary to repress *daf-9* for execution of the dauer program. Transcriptomic analysis comparing *chd-*

7–defective and normal dauers showed enrichment of collagen genes, consistent with a conserved role

for the TGF-β pathway in formation of the extracellular matrix. To validate a conserved function for *chd-7*

in vertebrates, we used *Xenopus laevis* embryos, an established model to study craniofacial

development. Morpholino mediated knockdown of Chd7 led to a reduction in *col2a1* mRNA levels. Both

embryonic lethality and craniofacial defects in Chd7-depleted tadpoles were partially rescued by over-

expression of *col2a1* mRNA. We suggest that pathogenic features of CHARGE syndrome caused by

Chd7 mutations, such as craniofacial malformations, result from the reduction of collagen levels, implying

that the extracellular matrix might represent a critical target of Chd7 in CHARGE development.

SIGNIFICANCE STATEMENT

CHARGE Syndrome is a complex developmental disorder caused by mutations in the

chromodomain helicase DNA-binding protein-7 (CHD7). Unfortunately, the cellular events that lead to

CHARGE syndrome are still poorly understood. In *C. elegans*, we identified *chd-7* in a screen for

suppressors of dauer formation, an alternative larval stage that develops in response to sensory signals

of a harsh environment. We found that *chd-7* regulates expression of collagens, which constitute the

worm's cuticle, a specialized extracellular matrix. In frog's embryos, we show that Chd7 inhibition leads to

poor Col2a1, which is necessary and sufficient to exhibit CHARGE features. These studies establish *C.*

elegans as an amenable animal model to study the etiology of the developmental defects associated with

pathogenic Chd7.

INTRODUCTION

CHARGE syndrome is a rare and severe neurodevelopmental disorder that affects the neural tube and neural crest cell derivatives, leading to hypogonadism, heart defects and craniofacial anomalies among other features (1). Inactivating mutations in *CHD7* (chromodomain-helicase-DNA binding 7) are the predominant cause of CHARGE, accounting for greater than 90% of the cases (2). *CHD7* is also mutated in Kallmann syndrome, a milder neurodevelopmental disorder with features overlapping with CHARGE, including impaired olfaction and hypogonadism (3). Exome sequencing studies in patients with autism spectrum disorders (ASDs) identified recurrent disruptive mutations in the related gene *CHD8* (4). The CHD proteins comprise a highly conserved family of SNF2-related ATP-dependent chromatin remodelers that are involved in chromatin remodeling and transcriptional regulation (5). Despite the public health relevance of these cognitive disorders, the mechanism of disease pathology due to mutations in CHD7/8 is poorly understood. The development of fly, fish and mouse models of CHARGE has abetted in characterization of associated dysfunction, but our understanding of the underlying pathology of CHARGE is still incomplete (6–11). CHD-7 is the ortholog of human CHD7 and CHD8 in *Caenorhabditis elegans* and has functions in habituation learning, normal locomotion, body size and fecundity (12, 13). It contains a conserved ATPase/SNF2 domain and two chromodomains for nucleosome interaction. Being the only worm homolog of the Class III CHD family, it contains a signature BRK domain (Brahma and Kismet domain) (Figure 1E).

When *C. elegans* encounter crowding, starvation or high temperature during early development, worms can halt reproductive programs to enter an alternative larval stage known as dauer. Dauers are long-lived, highly stress resistant and exhibit altered motility and metabolism (14–17). Upon return to normal growth conditions, the larvae exit dauer and develop into fertile adults. Study of dauer formation mutants has provided fundamental insights into pathways affecting longevity, neurodevelopment, metabolism, autophagy and neurodegeneration (14, 17–20).

The DAF-2/insulin/IGF- signaling 1 (IIS) pathway controls the dauer entry decision by coupling external cues with neuroendocrine signaling (21). In favorable conditions, DAF-2 activity initiates a conserved kinase cascade, leading to phosphorylation and inhibition of the transcription factor DAF-16/FOXO. In harsh environments, a decrease in the activity of DAF-2 and downstream components of the pathway leads to activation of DAF-16 and causes animals to arrest as dauers (22, 23). In addition to the DAF-2 pathway, DAF-7/ TGF-β signaling also regulates dauer development (24). When worms sense suitable conditions for reproductive development, ASI neurosensory cells secrete DAF-7, which binds to DAF-1/4 receptors leading to activation and phosphorylation of the R-SMAD complex DAF-8/14, promoting reproductive programs and inhibiting the pro-dauer complex composed of the SMAD protein DAF-3 and repressor DAF-5. Conversely, absence of DAF-7 leads to activation of the DAF-3/DAF-5 complex to promote dauer entry (25, 26).

DAF-7/TGF-β and DAF-2/IIS pathway were initially described as parallel pathways to regulate dauer entry

(27), but recent observations suggest a strong positive feedback between these pathways for dauer entry

and longevity (15–20). First, decreased signaling through the TGF-β pathway leads to differential

expression of many DAF-16 regulated genes with functions in longevity and dauer entry, such as SOD-3

or insulin peptides. This cross-activation of target genes may be important to amplify weak signals from

each sensory pathway to make an all-or-none decision to enter dauer (28–30). Second, the longevity of

daf-2 mutants can be blocked or enhanced by *daf-5* and *daf-3*, respectively, suggesting that

transcriptional components of the TGF-β can modulate IIS-dependent longevity genes. Third, for dauer

development, *daf-16* is epistatic to *daf-7/8/14 daf-c* mutants (30, 31). Lastly, both signaling pathways

converge on *daf-9* and *daf-12* to integrate outputs for diapause entry (32, 33). Indeed, *daf-9* expression

levels is critical for both entering and exiting diapause (17, 32).

In chromatin immunoprecipitation (ChIP)-chip studies, we identified *chd*-7 as a DAF-12 target

gene whose loss caused defective execution of dauer morphogenesis programs (34). We show that while

CHD-7 can modulate multiple IIS-associated processes, including *daf-2(e1370)* dauer formation,

longevity and immunity, epistatic experiments place *chd-7* in the DAF-7/TGF-β pathway. Whole genome

mRNA expression profiling of partial dauers show that *chd-7* mutants fail to repress *daf-9* during dauer

development, preventing developmental arrest of the constitutive dauer mutant *daf-7(e1372)*. In addition,

we found collagens to be among the most differentially regulated genes, consistent with a known role for

the TGF-β pathway in regulating components of the extracellular matrix (35).To validate a conserved

function for *chd-7* in vertebrates and study the relevance of our results for CHARGE etiology, we used

Xenopus laevis. Disruption of Chd7 function in *Xenopus* embryos results in craniofacial defects that mimic

CHD-dependent pathological phenotypes (36, 37). We demonstrate that Chd7 regulates expression of

the collagen type-II, alpha 1, *col2a1*, the main collagen protein of cartilage (38). Interestingly, craniofacial

malformations and embryonic lethality due to *chd7* knockdown can be rescued by *col2a1* expression.

These findings suggest a conserved function of *chd7*/*chd-7* in regulation of extracellular matrix

components and raise the intriguing possibility that defects in collagen expression may contribute to the

craniofacial defects seen in CHD7/8-dependent syndromes.

RESULTS

***chd-7* functions in development of the dauer larva**

Previously, we identified ~3000 potential DAF-12 target genes by ChIP-chip (34). We reasoned

that among these genes would be novel regulators of dauer development. To identify such genes, we

took advantage of the temperature-sensitive, constitutive dauer (Daf-c) allele *daf-2(e1371)*. At the non-

permissive temperature of 25°C, all the offspring enter the dauer phase and become SDS resistant (22).

We expected that inactivating targets of DAF-12 that function in dauer formation should suppress the Daf-

c phenotype of *daf-*2 and produce either defective, SDS-sensitive "partial" dauers or reproductive adults.

To confirm these results, we assayed suppression of the more severe *daf-2(e1370)* mutation and

discovered that *chd-7(RNAi)* produced arrested, SDS-sensitive larvae, indicative of the partial dauer

phenotype (39). As observed in Figure 1A and B, the axial ratio (length/width) of the partial dauers

resulting from *chd-7(RNAi)* exhibited a significant reduction of these proportions and appeared to have

defects in radial constriction of the dauer cuticle. By using a *chd-7* transcriptional reporter

(WBStrain00033709), we observed a substantial decrease in *chd-7* expression in the *daf-12(rh61rh411)*

background (Figure 1C and D). Thus, we infer that DAF-12 binds to the *chd-7* promoter (34) and

upregulates its expression.

To further validate our screen, we crossed *daf-2(e1370)* mutants with three *chd-7* deletion alleles

available from the Nematode Knockout Consortia (40). Allele *chd-7(tm6139)* contains a 594bp deletion

that generates a frame shift and premature stop codon, eliminating all known protein domains (Figure

1E). As shown in Figure 1F, SDS-sensitive dauers were obtained when double mutants are grown at

25°C, validating our screen. Comparison of two partial deletion alleles uncovered a critical role for the

BRK domain in dauer formation. The *chd-7(gk290)* allele contains a 859bp deletion that spans the BRK

domain and introduces a frameshift that eliminates the last 356aa. The *chd-7(gk306)* deletion is slightly

more C-terminal, truncating the protein immediately after the BRK domain (Figure 1E). When crossed into

daf-2(e1370) worms, *chd-7(gk306)* developed normal, SDS-resistant dauer larvae, whereas *chd-7(gk290)*

formed partial dauers that were sensitive to detergent (Figure 1F). Importantly, a functional transgene

expressing GFP-tagged CHD-7 protein (CHD-7::GFP) rescued the partial dauer phenotypes observed in

chd-7(gk290);daf-2(e1370) and *chd-7(tm6139);daf-2(e1370)* mutants (Figure 1F).

The dauer larva is characterized by a slim physique because of a reduction in the volume of

ectodermal tissues, including the hypodermis, seam cells and pharyngeal cells. In addition, the

hypodermis produces the dauer cuticle, which confers protection against external damage and

dehydration. During development, the seam cells fuse and the multinucleate cells produce the alae-

bilateral ridges in the cuticle that facilitate body motion. Dauer larvae also switch their metabolism to

accumulate lipids to survive for longer periods. To further characterize the nature of the defects in the

chd-7-induced partial dauers, we first studied the seam cells fusions with the adherens junction-

associated protein marker AJM-1::GFP and the morphology of the cuticle with scanning electron

microscopy (SEM). In dauering preparation, animals store fat in their intestinal and hypodermal cells,

which is critical to survive during hibernation (41). As shown in Figure 1G, partial dauers exhibited defects

in seam cell fusion and defective dauer alae formation. Next we used the lipid-labeling dye, Oil Red O

(ORO), to examine fat storage and found that, unlike other partial dauer mutants (19), the *chd-7;daf-2*

abnormal dauers did not exhibit fat-storage deficiencies (Supplemental Figure 1).

During dauer, the global developmental arrest also impacts the germ cells, slowing their divisions

and finally remaining quiescent (42). We observed that the germ line in *chd-7* mutant dauers was

substantially larger than in control dauers and appeared to arrest with a germline morphology resembling

that seen in L3 larval stage (Figure 1H). Therefore, we conclude that major morphological changes that

occur during dauer formation fail to be executed in *chd-7* mutants, including radial constriction of the

body, formation of an SDS-resistant cuticle with dauer alae, and developmental arrest of the germ line.

chd-7 is required for longevity and immunoresistance induced by IIS inactivation and germline

removal

In addition to dauer development, the IIS pathway also regulates longevity (14). Hence, we

sought to investigate whether *chd-7* also has roles in the determination of lifespan. First, we compared

survival of wild-type (N2) worms with two *chd-7* alleles and found that the dauer-defective allele *chd-*

7(gk290) significantly shortened lifespan, whereas *chd-7(gk306)* had only a marginal effect on longevity

(Figure 2A). Surprisingly, the CHD-7::GFP rescue transgene also reduced N2 lifespan (Figure 2B),

suggesting that *chd-7* copy number can influence longevity. We then analyzed how *chd-7* affects

longevity of IIS mutants. Remarkably, the dauer suppressor allele *chd-7(gk290),* but not *chd-7(gk306),*

shortened the lifespan extension of *daf-2* mutants to an extent comparable with the null allele of *daf-16,*

the key IIS downstream target (Figure 2C) (14). Furthermore, *daf-2* longevity was fully restored by CHD-

7::GFP (Figure 2D).

To determine if the effects on lifespan were specific to the IIS pathway, we assayed the

contribution of *chd-7* to the longevity induced by germ cell-less mutations, a longevity paradigm that

operates in parallel to IIS. Temperature-sensitive *glp-1(e2144)* animals are sterile and long-lived at non-

permissive temperatures (43). This lifespan extension was dependent on *chd-7*, as *chd-7(gk290);glp-*

1(e2144) double mutants had a mean lifespan significantly shorter than *glp-1(e2144)* single mutants

(Figure 2E). The impact was similar to that produced by absence of the nuclear receptor *daf-12* (Figure

2C), which is strictly necessary for germless longevity (43).

In addition to longevity, IIS reduction also enhances resistance against multiple stressors

including pathogen attack and starvation. We asked if *chd-7* inactivation impaired *daf-2*

immunoresistance too. As shown in Figure 2F, *chd-7* mutants repressed the increased survival of *daf-2*

worms upon exposure to the human opportunistic pathogen, *Pseudomonas aeruginosa* strain PA14 (44).

In contrast, *chd-7* had a modest effect on survival of wild-type worms' pathogen resistance. Furthermore,

the *chd-7(gk290)* and *chd-7(tm6139)* alleles reduced the survival of *daf-2(e1370)* L1 larvae subjected to

starvation stress (Supplemental Figure 2) (45). Taken together, these results suggest that *chd-7* mediates

the increased lifespan of at least two longevity paradigms (IIS mutants and germ cell-less animals), as

well as the response to pathogens and starvation.

CHD-7 is a member of the TGF-β pathway

Since our genetic analyses implicated *chd-7* as a dauer defective mutant (Daf-d) in the IIS

signaling pathway (Figure 1), we sought to understand whether it also had roles in the TGF-β dauer

pathway (27, 28). As shown in Figure 3A, at the restrictive temperature of 25°C, *chd-7;daf-7(e1372)*

double mutants bypassed the dauer arrest to become fertile adults. For *daf-5*, which acts as a dauer

suppressor of the TGF-β pathway, the Daf-d phenotype is observed at 25°C, but at a slightly higher

temperature, suppression is incomplete (26). Thus, we asked whether *chd-7* also prevented dauer arrest

at higher temperatures. As shown in Figure 3A, *daf-7*-dependent arrest was recapitulated at 26.5°C,

suggesting that *chd-7* has a Hid (high temperature-induced dauer formation) phenotype (46). As expected

for a putative transcriptional regulator, epistasis analysis placed CHD-7 downstream of the receptor DAF-

1 and the R-Smad DAF-14 (Figure 3B).

In addition to dauer formation, the TGF-β signaling pathway also regulates body size and male

tale development, mainly through the ligand DBL-1 (47, 48). To test if *chd-7* impacted these processes as

well, we measured the length of *chd-7* young adults and found them to be significantly shorter than N2

control animals (Figure 3C). We also observed that males carrying the severe loss-of-function allele *chd-*

7(tm6139) failed to mate with *fog-2* mutant females (Figure 3D) due to defects in male tail development

(not shown). This male infertility was rescued by the CHD-7::GFP transgene (Figure 3D).

More than twenty years ago, in a screen for suppressors of dauer formation within the TGF-β

pathway, *Inoue et al.* identified three complementation groups (31). We noticed that one of these, *scd-3*

(suppressor of constitutive dauer-3), was located between *unc-11* and *dpy-5* on chromosome I, in the

same genetic region as *chd-7*. Features of *scd-3(sa253)* worms include low brood size, egg-laying

defects (Egl), short body size (Dpy) and male abnormal defects (Mab), all of which were phenotypes

shared with *chd-7* as described above. In addition, improper gonad migration is a common phenotype of

scd-3 and *chd-7* mutants (Supplemental Figure 4). To determine if *chd-7* and *scd-3* are allelic, we

sequenced *scd-3(sa253)* and found a single G/A mutation in exon 8 of the *chd-7* locus, that introduces a

premature STOP codon at position Q2422, eliminating the BRK domain of CHD-7 protein, confirming that

chd-7 is *scd-3* (Figure 1E).

Transcriptomics analysis identifies *daf-9* and collagens as targets of CHD-7

To gain further insight into the role of CHD-7 in dauer development, we performed RNA-Seq. We

compared the transcriptomes of *daf-2(e1370)* dauers and *chd-7(gk290);daf-2(e1370)* partial dauers.

Differentially expressed genes (DEGs) analysis revealed that decreased expression of 28 genes and

increased expression of 56 genes in the double mutants (Figure 4A and Supplemental Table 3). Among

the latter group, we found *daf-9,* encoding the cytochrome p450 that integrates inputs from TGF-β and

insulin/IGF-II pathways to regulate DAF-12 activity during dauer development (49). We confirmed the

increased expression of *daf-9* upon *chd-7* loss by RT-qPCR (Figure 4B). We hypothesized that *daf-9*

expression levels must be critical in the decision to develop as either fertile adults or to enter diapause,

and therefore the increased expression of *daf-9* in the *chd-7;daf-2* double mutants may be preventing full

execution of the dauer program. Consistent with this hypothesis, we found that depletion of *daf-9* in

chd7;daf-7 animals restored the dauer arrest phenotype at the non-permissive temperature of 25°C

(Figure 4C). However, these animals still appeared to be detergent-sensitive partial dauers (Figure 4D),

since *daf-9* itself is required for proper dauer morphogenesis (39). Thus, we posit that the inability of *chd-*

7 mutants to fully repress *daf-9* may be sufficient to activate DAF-12 and inhibit dauer formation in both

the TGF-β and IIS pathways (Figure 4E).

The partial dauer larvae that arise from knockdown of *chd-7* are short and thick and lack dauer

alae, indicative of cuticle defects (Figures 1A and 1G). Thus, we reasoned that *chd-7* might also regulate

target genes that are required for dauer morphogenesis. Further analysis of our transcriptomic data

showed that 10% of the DEGs were collagens (*col-103*, *col-50*, *dpy-2*, *col-184*, *col-141*, *col-142*, *col-42*

and *dpy-9*), which are structural components of the cuticle. All these collagens showed increased

expression in the *chd-7*-mutant dauer larvae (Figure 4A). FPKM data from Modencode libraries indicates

that each of these collagens show very low expression during dauer and are expressed during various

stages of reproductive development, suggesting that repression of the genes must be important for dauer

development (50). Thus, our results suggest that CHD-7 is required for repression of dauer-specific

collagens. These results are consistent with growing evidence that the TGF-β pathway is a major

regulator of collagen deposition (35, 51). In fact, it was recently reported that two of our target genes, *col-*

141 and *col-142*, contain SMAD-binding elements (SBEs) and their expression is directly regulated by the

TGF-β pathway to determine body size (52).

In addition to the transcriptomic analysis of dauers, we analyzed CHD-7's CHIP-seq data

generated by the ModEncode project in young adults (Supplemental Table 4). Because the expression

profiles of adult TGF-β and IIS mutants show significant overlap and co-regulation of many DAF-16 target

genes (28), we compared this analysis with DAF-16 CHIP-seq data also from ModEncode in L4 larvae.

To our surprise, we found that both transcriptional regulators share a significant number of genes

(Supplemental Figure 5). Along with our observation that *chd-7* is necessary for lifespan extension of *daf-*

2 mutants, these data indicate that *chd-7* could modulate longevity by regulation of the IIS pathway,

possibly through direct regulation of target genes.

Chd7 regulates *col2a1* during *Xenopus* embryogenesis

Type-II collagen is an extracellular matrix (ECM) protein conserved in all multicellular animals,

which forms fibrils (53) and has fundamental roles in development and tissues homeostasis (54). In

vertebrates, the fibrillar type-II collagen is the major structural protein of cartilage and plays a prominent

role in cranial development in multiple organisms (38, 55). Collagen, type-II, alpha 1 (Col2a1) is the major

component of the cartilage matrix having a structural function and being an important extracellular

signaling molecule for regulation of chondrocyte proliferation, metabolism, and differentiation (56–58).

The African frog *Xenopus laevis* is a well-established model to study vertebrate facial disorders which

often arise from defects in neural crest development and migration (37). In *Xenopus* embryos, prior

studies established that Chd7 regulates neural crest specification and migration and its depletion

recapitulates the craniofacial defects seen in CHARGE patients (36). To investigate whether Chd7 has a

conserved mechanism of action and regulates collagen expression in vertebrates, we used a previously

validated morpholino to induce *Xenopus* Chd7 knockdown (*chd7*-MO) (36). In *Xenopus,* Col2a1 is

essential for normal development of the skeleton and its expression is restricted to the cartilaginous

skeleton of the tadpole and adult frog (59). In *chd7*-MO injected embryos, RT-qPCR revealed a significant

reduction of *col2a1* mRNAs as compared to control-injected animals (Figure 5A).

For targeted disruption of Chd7 function, we injected the dorsal-animal (D1) blastomeres of 8-cell

stage embryos fated to contribute to the dorsal anterior structures. *In situ* hybridization of unilaterally

injected embryos showed alterations in *col2a1* expression in the branchial arches and/or the otic vesicle

(ear vesicle) (59) upon *Chd7* depletion relative to embryos injected with a control morpholino (St-MO)

(Figure 5B). The injection of *chd7*-MO into both D1 blastomeres of 8-cell stage embryos with doses

between 1.25 ng and 2.5 ng showed 10-20% embryonic mortality (Figure 5C and data not shown). The

higher doses of 5 ng and 10 ng led to >50% lethality (Figure 5C). We next sought to investigate if the

mortality was related to collagen deficits and therefore conducted rescue experiments by co-expressing

Xenopus col2a1 mRNA with the morpholino. As shown in Figure 5C, coinjection of *col2a1* mRNA

substantially improved (~50%) embryo survival relative to the injection of *chd7*-MO alone. We then asked

if ectopic *col2a1* mRNAs could suppress the craniofacial defects associated with *chd7* loss-of-function.

Initially, we analyzed the gross morphology of the surviving stage 45 tadpoles and observed a high

incidence of craniofacial malformations (83%) in Chd7-depleted animals (Supplemental Figure 6A and

6C) and a significant reduction upon *col2a1* mRNA coinjection (43%) (Supplemental Figure 6B). Next, we

examined eye size and eye distance, since microthalmia and midline defects are often associated with

CHARGE syndrome (60) and are recapitulated in *Xenopus* embryos (36). Both eye size and distance

between eyes were reduced in Chd7-depleted tadpoles and were partially rescued by *col2a1* mRNA

expression (Figure 5D and 5E). Therefore, expression of *col2a1* ameliorated the phenotypes associated

with pathogenic Chd7 suggesting that collagen is a conserved and important target of this protein.

DISCUSSION

We initially identified *chd-7* in chromatin immunoprecipitation (ChIP)-chip studies of the nuclear

receptor DAF-12 (34), a transcription factor regulating worm aging, development, and dauer formation

(61), and found it to be critical for *daf-2* mediated dauer development. Comparison of the *chd-7* alleles

gk290 and *gk306* showed a critical role for the BRK domain in dauer development and longevity (Figures

1F-H, 2A and 2C). In CHARGE patients, deletions or mutations within the BRK domains of CHD7 are

sufficient to elicit all the features characteristic of the disease, underscoring the importance of this domain

(62). The phenotypic differences between the worm alleles highlight the potential of the worm to delimit

functional domains of CHD-7 that contribute to disease pathology. In mice, homozygous mutations in

Chd7 lead to embryonic lethality at E10.5, in part because *Chd7* is necessary for early brain development

(36). In worms, the presumptive null allele c*hd-7(tm6139)* and *scd-3(sa253)* were viable but showed

reproductive defects, such as improper gonad proliferation and migration, reduced fecundity, male tail

defects and hermaphrodite vulval defects (31), indicating that *C. elegans* are more able to tolerate loss of

CHD-7 than mice or humans. Thus, our studies establish *C. elegans* as an animal model to study the

mechanisms underlying the developmental defects observed in pathogenic Chd7.

Both a *chd-7* mutation and *chd-7* overexpression shortened the lifespan of otherwise wild-type

worms, suggesting that CHD-7 proteins levels must be tightly regulated to ensure proper development. Of

note, Chd7 is the most commonly amplified gene in tumors amongst the CHD superfamiliy members, and

its overexpression is associated with aggressive subtypes of breast cancer and poor prognosis (63).

Thus, in humans, as well as worms, gain-of-function phenotypes are associated with Chd7/*chd-7*

overexpression.

Our genetic epistasis analyses placed CHD-7 downstream of the TGF-β-like DAF-7, the type I

receptor DAF-1, and the R-SMAD DAF-14 placing CHD-7 at the level of the Co-Smad DAF-3 and the

Sno/Ski repressor DAF-5 (64). While *daf-3* and *daf-5* completely suppress dauer

formation in a *daf-7* background at 25°C, *daf-5* fails to suppress dauer formation in TGF-β mutants at

27°C (26), supporting a role for CHD-7 at the DAF-3/DAF-5 step in the pathway (Figures 3A and 3B).

Interestingly, these transcriptional regulators have opposite effects on *daf-2*-induced longevity: *daf-3*

enhances *daf-2(e1370)* longevity while *daf-5* mutations suppresses it (30). We observed that *chd-7*

suppresses *daf-2* longevity (Figure 2C), like *daf-5(e1386)*. Together these data suggest that CHD-7 either

regulates *daf-5* expression or directly interacts with DAF-5 to regulate downstream target genes, or both.

In mice, CHD7 was shown to physically interact with SMAD1 and form a transcriptional complex with

SMAD4, the mammalian ortholog of DAF-3 (65). We therefore speculate that DAF-3, DAF-5, and CHD-7

may be in ternary complex that regulates *daf-9* expression for dauer entry (Figure 4C). Interactome

mapping of the TGF-β pathway also identified SWSN-1, a SWI/SNF subunit component of the BAF

complex, as a physical interactor of DAF-3 (66). CHD7 interacts with human and *Xenopus* PBAF

(polybromo- and BRG1-associated factor-containing complex) to control neural crest gene expression

(36). In worms, both *swsn-1* and *chd-7* fail to develop normal dauers in *daf-2* and *daf-7* mutants

(Supplemental Figure 3 and (67)). Thus, we envision that CHD-7 may work together with the BAF

complex and DAF-3/DAF-5 to control gene expression of target genes critical for dauer formation.

DAF-9 activity is necessary and sufficient for the decision to enter diapause: reduced activity of

TGF-β and IIS pathways leads to *daf-9* repression and dauer development. Conversely, *daf-9* expression

in the hypodermis is sufficient to inhibit diapause, driving reproductive programs in *daf-7* mutants (29, 32).

Ectopic *daf-9* expression also drives reproductive programs in the weak *daf-2(e1368)* allele, but only

partially suppresses *daf-2(e1370)* diapause leading to arrest as L3 or early L4 larvae (32). Based on

these observations, we speculate that *daf-9* misexpression explains a subset of the *chd-7* phenotypes

observed herein, including the partial dauer phenotype, gonad migration defects, and vulval protrusions,

all of which overlap with published *daf-9* phenotypes (32, 33, 68). Interestingly, *daf-9* is both upstream

and downstream of *daf-12* for dauer formation *(32, 33)*. Our data demonstrated that DAF-12 regulates

chd-7 expression and CHD-7 in turn regulates *daf-9* expression. Therefore, we propose that CHD-7

belongs to the transcriptional complex that regulates the feedback loop between *daf-12* and *daf-9*.

Among the differentially expressed genes, we noted the presence of several G protein coupled

receptors (GPCRs), which could be candidate chemoreceptors for dauer pheromone or environmental

cues and thus may be required for either dauer entry or maintenance (42). Consistent with our

transcriptomic studies, *Liu et al.* analyzed the expression differences between L2/L3 larvae and TGF-β

mutants undergoing diapause and found enrichment of collagen genes, GPCRs and *daf-9*, further

validating our placement of *CHD-7* in the TGF-β dauer signaling pathway (29). In a more recent paper,

also was established that GPCR gene expression increases significantly during dauer commitment (69).

In *C. elegans*, the TGF-β family pathway has diversified and has unique ligands and effectors for

the control of dauer induction and aspects of somatic development. Mutations in the TGF-β ligand *dbl-1*

and its downstream components cause a small body size (Sma phenotype) (47, 48). Consistent with *chd-*

7 being a component of the TGF-β signaling pathway, *chd-7(gk290)* and *chd-7(tm6139)* are significantly

shorter than wild-type worms (Figure 3C). This phenotype is also observed in *scd-3* mutants (31). The

dbl-1 pathway mutants have defects in male tale development (48), a phenotype shared in *scd-3* (31) and

chd-7 mutant animals (data not shown and (31)). Therefore, our results suggest that CHD-7 acts as a

regulator of both the *daf-7* and *dbl-1* branches of the TGF-β pathway.

Craniofacial anomalies seen in CHARGE patients involve tissues deriving from cells of the neural

crest, including craniofacial cartilage and bone, heart, ears and eyes (60, 70). *Xenopus laevis* is an

established model to understand fundamental questions of craniofacial biology (37). *Chd7* knockdown in

frogs and in zebrafish (11, 36), leads to defects in neural crest specification and migration, which was

recently been recapitulated in human embryonic stem cells (hESCs) (71). Our RNA-seq analysis revealed

that CHD-7 regulates expression of genes from the cuticle, a modified extracellular matrix (72). Here, we

show a role for Chd7 in regulating *col2a1* expression, a type II collagen and the major component of

cartilage (55). Interestingly, regulation of *col2a1* by Chd7 is also observed in zebrafish (73). While

additional studies are required, we speculate that Chd7 could regulate *col2a1* in a complex with the

transcription factor Sox10 (74, 75) or through the TGF-β signaling pathway (35, 76). Supporting the latter

mechanism, it was demonstrated in chondrocytes that TGF-β regulates *col2a1* expression (77, 78). Taken

together, our data supports a model in which collagen misexpression by pathogenic Chd7 leads to
craniofacial defects and embryonic lethality.
428
429 **MATERIALS and METHODS**
430
C. elegans **strains**
All strains were grown and maintained on standard nematode growth medium (NGM). In
Argentina (Hochbaum lab), plates were supplemented with 0.1 mg/ml streptomycin and 100 U/ml Nystatin
using the *E. coli* OP50-1 strain as the food source. In Pittsburgh (Yanowitz lab), strains were grown on
standard NGM seeded with OP50. Temperature-sensitive strains were maintained at 16°C and shifted to
the non-permissive temperature of 25°C to induce dauer formation, unless otherwise indicated. All other
strains were maintained at 20°C. All *daf-7* strains were maintained at low density to prevent dauer
induction at 16°C. Dauer studies were performed in both the Yanowitz and Hochbaum laboratories to
validate results. All *chd-7* alleles were outcrossed at least 5 times prior to use.
Strains utilized in this study are listed in Supplemental Table 1. Standard genetic crosses were
used to make double or triple mutants. The presence of mutant alleles was confirmed a) by the *daf-c*
phenotypes in animals heterozygous for additional mutations and b) by PCR and/or sequencing for all
additional mutations.
444
RNAi screen for dauer suppressors
All RNAi clones were picked from the Ahringer bacterial feeding library (79). These *E. coli* clones
were seeded on NGM plates supplemented with 1 mM of IPTG (Isopropyl β-D-1-thiogalactopyranoside)
and 0.1 μg/ml ampicillin, and used for inducing RNAi by the feeding method (80).
GL228 [*rrf-3(pk1426)*] II;*daf-2(e1371)* III] eggs were placed in 24-well RNAi plates seeded with
bacteria expressing the dsRNA of interest. Worms were maintained for 5 days at 15ºC until adulthood,
then were transferred to an identical 24-well plate to lay eggs for 5 h. Adults were removed and the eggs
were incubated at 25°C for 4 days to allow formation of dauers. *daf-16(RNAi)* and the empty vector were
used as controls. Proper dauer formation was assessed by observation in a dissecting microscope and by
1% SDS resistance. RNAi clones that caused abnormal dauer phenotypes were validated in *daf-2(e1370)*
worms. Identity of the dsRNA was confirmed by sequencing (Macrogen, Korea).
456
daf-9 **suppression assays**
For preparation of *daf-9*(RNAi) plates, bacterial cultures were grown overnight in LB with 10 μg/ml
tetracycline and 50 μg/ml carbenicillin and induced with 4 mM IPTG for 4 h. Cultures were spun down,
suspended in 1/10 volume and 30 μl of bacteria were seeded on 3 cm NGM plates made with 1 mM IPTG

and 50 µg/ml carbenicillin. Plates were grown overnight at 25°C and stored at 4°C for no more than 2
weeks prior to use.
L4 stage animals were placed on 3 cm RNAi plates. Two worms per plate were used for *daf-*
7(e1372), while three worms were used for *daf-7;chd-7(gk290)* and *daf-7;chd-7(tm6139)*. After 72 h, the
adults were removed and plates were replaced at 25°C for 2-3 days. The total number of dauers, L4s,
and adults were then assessed.
467

SDS survival assay
Young adults were transferred to seeded plates and permitted to lay eggs for 5 days at 25°C. The
arrested progeny were then washed off plates with M9 (22 mM KH_2PO_4, 42 mM Na_2HPO_4, 85.5 mM
NaCl, 1 mM $MgSO_4$) into 15 ml glass conical tubes. Collected animals were washed 2-3 times with M9
and the excess liquid was aspirated off. Animals were then treated with 2 ml of 1% SDS for 30 min on a
nutator at 25ºC. Following incubation, the samples were washed 3 times with M9 and any excess liquid
was aspirated off. Animals were aliquoted to 5 seeded plates with 50-70 worms per plate and allowed to
recover at 16°C. The recovered animals were then quantified, and the percentage recovered was
calculated. This was repeated 3 times for each strain tested. Comparisons were performed using the χ-
squared Test in R; P values were corrected using the Bonferroni method for multiple comparisons.
478

479 *fog-2* **mating assay**
480   To determine if males were capable of siring offspring, 8 males from each strain tested were
481 plated with 4 *fog-2(q71)* adult females on seeded 10 cm plates. After 24 h, *fog-2* females were transferred
482 to new plates and within 48 h the proportion of fertile females were scored. Assay was repeated two
483 times.
484

Lifespan assays
All lifespan experiments were conducted by transferring 1 day-old adults from 15°C to 20°C for
the remainder of the lifespan assay. NGM plates were seeded with *E. coli* OP50-1. ~150 L4
hermaphrodites were transferred to 5 plates per experiment. Every 48 h, animals were scored as alive,
dead or censored (animals that exploded, died from bagging or dried out at the edges of the plates).
Animals were considered dead when they did not respond to a soft touch to the head with a pick. To
prevent the progeny from interfering with the assay, adults were transferred to fresh plates every 48 h
until egg production ceased. For *glp-1(e2144)* assays, eggs were kept at 20°C for 4 h and then
transferred at 25.5°C for 72 h to induce sterility and switched to 20°C for the remainder of the experiment.
Lifespan data were analyzed using the Kaplan-Meier method. Statistics were calculated using the Mantel-
Cox nonparametric log-rank method using OASIS2 (81).
496

Pathogen resistance assay

The pathogenic bacterial strain used in this study was *Pseudomonas aeruginosa* (strain PA14). This strain was streaked from a frozen stock onto an LB agar plate, incubated at 37°C overnight and then kept at 4°C (shelf-life of one week). For survival assay, a single PA14 colony was inoculated in King's broth and incubated at 37°C overnight with shaking. 20 µl of this culture was seeded onto slow killing NGM plates (containing 0.35% peptone instead of 0.25%) and incubated for 24 h at 37°C. The plates were then left to sit at room temperature for 24 h. The following day, 150 L4 hermaphrodites grown at 15°C were distributed onto five PA14 plates and incubated at 25°C. Survival was monitored at intervals of 6-12 h and live, dead and censored animals were recorded. Data and statistics were analyzed using the Kaplan-Meier method as described in the section "Lifespan assays".

Microscopy and fluorescence imaging

For imaging dauers and adults, worms were immobilized in a 25 mM sodium azide solution on fresh 4% agarose pads and imaged at 10x and 20x magnification. Images were collected using a Zeiss Axioplan Imaging Microscope with a DIC system and Zeiss Plan-Neofluar 10x and 20x objectives lens. Images were acquired with a Micropublisher 3.3 camera (Q Imaging). ImageJ (NIH) software was used to quantify worm size.

For imaging *pCHD-7::mCherry* fluorescence, worms were immobilized with 1 mM levamisole on fresh 2% agarose pads and imaged immediately using a Nikon A1r confocal microscope equipped with a 40x PLAN APO oil objective. ImageJ (NIH) software was used to quantify fluorescence intensity.

For imaging gonads in *daf-2(e1370)* and mutants, young adults were transferred to seeded plates and permitted to lay eggs for 5 days at 25°C to form dauers, then collected and fixed with Carnoy's solution (75 µl EtOH, 37.5 µl Acetate, 12.5 µl Chloroform) and stained with 5 mg/ml DAPI (4′,6-diamidino-2-phenylindole) in PBS. Animals were imaged as 0.5 µm Z-stacks with the Nikon A1r Confocal Microscope with 40x and 60x plan APO oil objectives.

For imaging *ajm-1::GFP*, young adults were transferred to seeded plates and allowed to lay eggs for 5 days at 25°C to form dauers. The arrested progeny was then washed off plates with M9 into 15 ml glass conical tubes. Collected animals were washed 2-3 times with M9 and the excess liquid was aspirated off. Animals were then immobilized with levamisole and the seam cells were imaged as Z-stacks by confocal microscopy as described immediately above.

For scanning electron microscopy, worms were fixed by immersing in 2.5% glutaraldehyde in PBS for several hours. Worms were washed 3X in PBS then post fixed in aqueous OsO_4 for 1h, then washed 3x in PBS, dehydrated through a graded series of ethanols (30-100%) then chemically dried with 2x 10 min incubations in hexamethyldisilazane. Dried worms were sprinkled onto copper double stick tape on aluminum stubs, sputter coated with 3.5 nm gold-palladium alloy then evaluated on a JEOL JEM 6335F microscope at 5 kV.

Library preparation and RNA-seq

daf-2(e1370) and *chd-7(gk290);daf-2(e1370)*, synchronized eggs were kept at 25°C for 10 days and resulting dauers were collected and frozen. Total RNA was extracted with TRIzol (Invitrogen) following the kit´s protocol. cDNA library was prepared with NEBNext Ultra II RNA library prep kit for Illumina (New England Biolabs), and the sequencing carried out using Illumina's HiSeq-2500 sequencer with single-end mode and read length of 50 bp. Five replicates for *daf-2(e1370)* vs. *chd-7(gk290);daf-2(e1370)* were sequenced. For data assessment, a quality control with FastQC software (version 0.11.5) was used. First the raw reads that aligned against the *E. coli* genome (K12 genome) were removed. The remaining sequences were aligned against the reference genome of *C. elegans* WS260 using STAR (version 2.5.4a). The number of mapped reads to genes was counted using Htseq (version 0.9.1). Finally, the DEGs were determined using DESeq2 (version 1.20.0) with a cutoff of 0.05 on False Discovery Rate (FDR). R version 3.5.0 (2018-04-23) and Bioconductor version 3.7 with BiocInstaller version 1.30.0 were used. Heatmaps were generated using pheatmap package (version 1.0.12) with hierarchical clustering on the rows with the default options.

Genomic Sequencing of *scd-3*

A 6 cm plate replete with *scd-3* gravid animals was washed with 1 ml M9 and collected in a glass conical tube. Worms were washed extensively and then placed on a nutator for 1 h to allow remaining gut bacteria to be passed into the medium. Worms were again washed 3-4 times and settled by gravity. The worm pellet was transferred to 1.5 ml tubes and genomic DNA was prepared according to the manufacturer's protocol with the Purelink Genomic DNA Kit (Invitrogen) except that after addition of digestion buffer, worms were pulverized with a microfuge hand dounce prior to incubation at 55°C. Sequencing was performed by Psomagen, Qiagen CLC Genomics Workbench was used to align the DNA against WBcel235 and view the variant.

Oil-Red-O (ORO) staining

Dauers grown at 25°C for 5 days were stained for lipids using ORO, as previously described (82). Animals were mounted on a 4% agarose pad and observed in a Zeiss Axioplan brightfield microscope equipped with a Micropublisher 3.3 camera (Q Imaging). Image J (NIH) was used to quantify the amount of lipids in each animal. At least 20 animals of each strain were quantified. Statistical differences was determined by Student's t-test.

L1 survival

Experiments were done at 20°C. Eggs were obtained by bleaching of gravid adults and kept under gentle shaking in 5ml sterile M9 supplemented with 0.1µg/ml streptomycin to hatch (16h). To

normalize population density, resulting L1 larvae were diluted to obtain 1 larva/µl in 5ml of M9 and kept

under constant agitation for the remainder of the experiment. Every 48h, a 100µl aliquot of L1 larvae were

spotted onto a NGM plate and then incubated for 72h. Animals were scored as alive if developed beyond

the L2 stage. Percentage of the population alive was normalized to day 1 seeded L1 larvae.

Gonad staining

Late L4/Young Adult animals from the relevant strains were collected and fixed with Carnoy's

solution (75 µl EtOH, 37.5 µl Acetate, 12.5 µl Chloroform) and stained with 5 mg/ml DAPI (4′,6-diamidino-

2-phenylindole) in PBS. Animals were imaged as 0.5 µm Z-stacks with the Nikon A1r Confocal

Microscope with 20x objective.

ChIP-seq analysis

Data from CHD-7 and DAF-16 ChIP-seq generated by ModEncode project was analyzed.

Reference for CHD-7-eGFP: https://www.encodeproject.org/experiments/ENCSR010MNU/. Reference for

DAF-16-eGFP: https://www.encodeproject.org/experiments/ENCSR946AUI/. Peaks were downloaded in

ce10 and annotated using Homer software. Gene lists from the peak calling were generated and used to

compare CHD-7 and DAF-16.

***Xenopus laevis* embryo manipulation and microinjections**

Xenopus embryos were obtained by natural mating. Adult frogs' reproductive behavior was

induced by injection of human chorionic gonadotropin hormone. Eggs were collected, de-jellied in 3%

cysteine (pH 8.0), maintained in 0.1 X Marc's Modified Ringer's (MMR) solution and staged according to

Nieuwkoop and Faber (83). The embryos were placed in 3% ficoll prepared in 1 X MMR for

microinjection. Chd7 morpholino (*chd7*-MO: 5′-AACTCATCATGCCAGGGTCTGCCAT-3′) specificity has

been previously characterized (36). *Chd7*-MO and Standard Control morpholino (St-MO) were provided

by Gene Tools, LLC. The cDNA of *X. laevis col2a1* was amplified by PCR from pCMV-Sport 6-*col2a1*

(Dharmacon) with primers M13F and M13R. The PCR fragment was digested with *EcoR*V and *Not*I and

cloned into pCS2+ previously digested with *Stu*I and *Not*I. Capped mRNAs for *col2a1* were transcribed *in*

vitro with SP6 using the mMessage mMachine kit (Ambion) following linearization with *Not*I. *Chd7*-MO

and *col2a1* mRNA were injected into both D1 blastomeres of 8-cell staged embryos (82) for lethality and

morphometrics analysis. *Chd7*-MO was injected into one D1 blastomeres of 8-cell staged embryos for

analysis of *col2a1* expression.

Whole-mount *in situ* hybridization was carried out as previously described (85). pCMV-Sport 6-

col2a1 (Dharmacon) construct was linearized with *Sal*I and transcribed with T7 for antisense probe

synthesis. Morphometrics analyses were done on fixed tadpoles using ImageJ (NIH) software.

Morphometric measurements were normalized to the mean of the uninjected group in order to compare

between independent experiments. For cartilage staining, stage 45 tadpoles were fixed with MEMFA for

24 hours at 4°C, dehydrated into 100% ethanol and stained in 0.01% Alcian blue 8GX in 70%

ethanol/30% glacial acetic acid for three nights. Distaining was done in 100% ethanol followed by

rehydration in 2% KOH. Animals were cleared in graded glycerol in 2% KOH and skulls were dissected

under stereoscope. Images of whole embryos and skulls were collected with a Leica DFC420 camera

attached to a Leica L2 stereoscope.

RNA preparation and RT-qPCR

For *Xenopus* RNA extraction, 2-cell staged embryos were injected into both blastomeres with 5

ng or 10 ng of *chd7*-MO, collected at stage 23 and snap-frozen for later processing. Six embryos were

combined for each treatment. For worm RNA extraction, dauers and partial dauers were developed for a

week at 25°C. For all the samples, RNA was isolated using RNAzol (Molecular Research). RNA

concentration of each sample was measured using a Nanodrop spectrophotometer and 250 ng of RNA

was reverse transcribed using qScript cDNA SuperMix (QuantaBio). Quantitative PCR (qPCR) was

preformed using the Forget-Me-Not qPCR Master Mix (Biotium) with a BioRad iCycler thermocycler.

Amplification was performed using oligonucleotides designed with PerlPrimer software for *col2a1*

(Forward 5′ TCCCTGTTGATGTTGAAGCC 3′; Reverse 5′CAATAGTCACCGCTCTTCCA 3′ and ODC

primers have been previously described (86) (Forward 5′CAAAGCTTGTTCTACGCATAGCA 3′; Reverse

5′GGTGGCACCAAATTTCACACT 3′). The relative expression of *col2a1* was normalized

to *ODC* expression and determined via the method previously described (87). *daf-9* was amplified with

specific oligos (Forward 5′ATTCCCCACAAAACAATCGAAGAAT 3′; Reverse 5′

GAGATTCAAACACGTTTGGATCG 3′) and expression was normalized to the housekeeping gene *cdc-42*

(Forward 5′CTGCTGGACAGGAAGATTACG 3′; Reverse 5′ CTGGGACATTCTCGAATGAAG 3′).

Ethics Statement

Xenopus laevis experiments were carried out in strict accordance with the recommendations in

the Guide for the Care and Use of Laboratory Animals of the NIH and also the ARRIVE guidelines. The

animal care protocol was approved by the Comisión Institucional para el Cuidado y Uso de Animales de

Laboratorio (CICUAL) of the School of Applied and Natural Sciences, University of Buenos Aires,

Argentina (Protocol #64).

ACKNOWLEDGMENTS

The authors would like to thank Bruno Moretti, Hernan Grecco, Mario Rossi, and Julie Kocherzat

for experimental support. DMJ and ASC were supported by the Consejo Nacional de Investigaciones

Científicas y Técnicas (CONICET) Doctoral Fellowship Program. DH laboratory was supported by the

Agencia Nacional de Promoción Científica y Tecnológica of Argentina (PICT-2016-0269) and CONICET

(PIP 1122015 0100731 CO). MCC laboratory was supported by the Agencia Nacional de Promoción

Científica y Tecnológica of Argentina (PICT-2013-0381). Funding for this work was also provided by

grants from the CHARGE Syndrome Foundation (JLY and DH), The Company of Biologists (DMJ),

NIGMS (R01GM104007 to JLY), and NIA (R01AG051659 to AG). This work was supported in part by the

Intramural Research Program of the NIH and the National Institute of Diabetes and Digestive and Kidney

Diseases (SY).

REFERENCES

1.   L. E. L. M. Vissers, *et al.*, Mutations in a new member of the chromodomain gene family cause

CHARGE syndrome. *Nature genetics* **36**, 955–7 (2004).

2.   R. Balasubramanian, *et al.*, Functionally compromised CHD7 alleles in patients with isolated

GnRH deficiency. *Proc Natl Acad Sci U S A* **111**, 17953–17958 (2014).

3.   H.-G. Kim, *et al.*, Mutations in CHD7, encoding a chromatin-remodeling protein, cause idiopathic

hypogonadotropic hypogonadism and Kallmann syndrome. *Am. J. Hum. Genet.* **83**, 511–519 (2008).

4.   R. Bernier, *et al.*, Disruptive CHD8 Mutations Define a Subtype of Autism Early in Development.

Cell **158**, 263–276 (2014).

5.   C. G. A. Marfella, A. N. Imbalzano, The Chd family of chromatin remodelers. *Mutation research*

618, 30–40 (2007).

6.   S. A. Patten, *et al.*, Role of Chd7 in Zebrafish: A Model for CHARGE Syndrome. *PLoS One* **7**

(2012).

7.   E. A. Bosman, *et al.*, Multiple mutations in mouse Chd7 provide models for CHARGE syndrome.

Hum Mol Genet **14**, 3463–3476 (2005).

8.   C. Tian, *et al.*, Otitis media in a new mouse model for CHARGE syndrome with a deletion in the

Chd7 gene. *PloS one* **7**, e34944 (2012).

9.   G. Daubresse, *et al.*, The Drosophila kismet gene is related to chromatin-remodeling factors and

is required for both segmentation and segment identity. *Development* **126**, 1175–1187 (1999).

10.   D. J. Melicharek, L. C. Ramirez, S. Singh, R. Thompson, D. R. Marenda, Kismet/CHD7 regulates

axon morphology, memory and locomotion in a Drosophila model of CHARGE syndrome. *Hum Mol Genet*

19, 4253–4264 (2010).

11.   Z. Asad, *et al.*, Rescue of neural crest-derived phenotypes in a zebrafish CHARGE model by

Sox10 downregulation. *Hum Mol Genet* **25**, 3539–3554 (2016).

12.   T. A. McDiarmid, *et al.*, Systematic phenomics analysis of autism-associated genes reveals

parallel networks underlying reversible impairments in habituation. *PNAS* **117**, 656–667 (2020).

13.   W.-R. Wong, *et al.*, Autism-associated missense genetic variants impact locomotion and

neurodevelopment in Caenorhabditis elegans. *Hum Mol Genet* **28**, 2271–2281 (2019).

14.   C. Kenyon, J. Chang, E. Gensch, A. Rudner, R. Tabtiang, A C. elegans mutant that lives twice as
long as wild type. *Nature* **366**, 461–464 (1993).

15.   P. L. Larsen, Aging and resistance to oxidative damage in Caenorhabditis elegans. *Proc Natl*
Acad Sci U S A **90**, 8905–8909 (1993).

16.   M. M. Gaglia, C. Kenyon, Stimulation of Movement in a Quiescent, Hibernation-Like Form of
Caenorhabditis elegans by Dopamine Signaling. *J Neurosci* **29**, 7302–7314 (2009).

17.   J. Wang, S. K. Kim, Global analysis of dauer gene expression in Caenorhabditis elegans.
Development **130**, 1621–1634 (2003).

18.   E. Cohen, J. Bieschke, R. M. Perciavalle, J. W. Kelly, A. Dillin, Opposing activities protect against
age-onset proteotoxicity. *Science* **313**, 1604–1610 (2006).

19.   A. Meléndez, *et al.*, Autophagy genes are essential for dauer development and life-span
extension in C. elegans. *Science* **301**, 1387–1391 (2003).

20.   R. Christensen, L. de la Torre-Ubieta, A. Bonni, D. A. Colón-Ramos, A conserved PTEN/FOXO
pathway regulates neuronal morphology during C. elegans development. *Development* **138**, 5257–5267
(2011).

21.   C. A. Wolkow, K. D. Kimura, M.-S. Lee, G. Ruvkun, Regulation of C. elegans Life-Span by
Insulinlike Signaling in the Nervous System. *Science* **290**, 147–150 (2000).

22.   R. C. Cassada, R. L. Russell, The dauerlarva, a post-embryonic developmental variant of the
nematode Caenorhabditis elegans. *Dev Biol* **46**, 326–342 (1975).

23.   D. L. Riddle, P. S. Albert, "Genetic and Environmental Regulation of Dauer Larva Development"
in *C. Elegans II*, 2nd Ed., D. L. Riddle, T. Blumenthal, B. J. Meyer, J. R. Priess, Eds. (Cold Spring Harbor
Laboratory Press, 1997) (September 25, 2017).

24.   P. Ren, *et al.*, Control of C. elegans larval development by neuronal expression of a TGF-beta
homolog. *Science* **274**, 1389–1391 (1996).

25.   D. Park, A. Estevez, D. L. Riddle, Antagonistic Smad transcription factors control the dauer/non-
dauer switch in C. elegans. *Development* **137**, 477–485 (2010).

26.   L. S. da Graca, DAF-5 is a Ski oncoprotein homolog that functions in a neuronal TGF pathway to
regulate C. elegans dauer development. *Development* **131**, 435–446 (2003).

27.   J. H. Thomas, D. A. Birnby, J. J. Vowels, Evidence for parallel processing of sensory information
controlling dauer formation in Caenorhabditis elegans. *Genetics* **134**, 1105–1117 (1993).

28.   W. M. Shaw, S. Luo, J. Landis, J. Ashraf, C. T. Murphy, The C. elegans TGF-β Dauer Pathway
Regulates Longevity via Insulin Signaling. *Current Biology* **17**, 1635–1645 (2007).

29.   T. Liu, K. K. Zimmerman, G. I. Patterson, Regulation of signaling genes by TGFbeta during entry
into dauer diapause in C. elegans. *BMC Dev Biol* **4**, 11 (2004).

30.   S. D. Narasimhan, *et al.*, PDP-1 Links the TGF-β and IIS Pathways to Regulate Longevity,
Development, and Metabolism. *PLoS Genet* **7** (2011).

31. T. Inoue, J. H. Thomas, Suppressors of transforming growth factor-beta pathway mutants in the

Caenorhabditis elegans dauer formation pathway. *Genetics* **156**, 1035–1046 (2000).

32. B. Gerisch, A. Antebi, Hormonal signals produced by DAF-9/cytochrome P450 regulate C.

elegans dauer diapause in response to environmental cues. *Development* **131**, 1765–1776 (2004).

33. H. Y. Mak, G. Ruvkun, Intercellular signaling of reproductive development by the C. elegans DAF-

9 cytochrome P450. *Development* **131**, 1777–1786 (2004).

34. D. Hochbaum, *et al.*, DAF-12 regulates a connected network of genes to ensure robust

developmental decisions. *PLoS Genet* **7**, e1002179 (2011).

35. R. A. Ignotz, J. Massagué, Transforming growth factor-beta stimulates the expression of

fibronectin and collagen and their incorporation into the extracellular matrix. *J. Biol. Chem.* **261**, 4337–

4345 (1986).

36. R. Bajpai, *et al.*, CHD7 cooperates with PBAF to control multipotent neural crest formation.

Nature **463**, 958–962 (2010).

37. A. Dubey, J.-P. Saint-Jeannet, Modeling Human Craniofacial Disorders in Xenopus. *Curr*

Pathobiol Rep **5**, 79–92 (2017).

38. D. W. Seufert, J. Hanken, M. W. Klymkowsky, Type II collagen distribution during cranial

development in Xenopus laevis. *Anat. Embryol.* **189**, 81–89 (1994).

39. P. S. Albert, D. L. Riddle, Mutants of Caenorhabditis elegans that form dauer-like larvae. *Dev Biol*

126, 270–293 (1988).

40. E. S. Haag, Dial-a-mutant: web-based knockout collections for model organisms. *Biol. Cell* **99**,

343–347 (2007).

41. P. Narbonne, R. Roy, Caenorhabditis elegans dauers need LKB1/AMPK to ration lipid reserves

and ensure long-term survival. *Nature* **457**, 210–214 (2009).

42. P. Kadekar, R. Roy, AMPK regulates germline stem cell quiescence and integrity through an

endogenous small RNA pathway. *PLoS Biol* **17** (2019).

43. H. Hsin, C. Kenyon, Signals from the reproductive system regulate the lifespan of C. elegans.

Nature **399**, 362–366 (1999).

44. D. A. Garsin, *et al.*, Long-lived C. elegans daf-2 mutants are resistant to bacterial pathogens.

Science **300**, 1921 (2003).

45. I. Lee, A. Hendrix, J. Kim, J. Yoshimoto, Y.-J. You, Metabolic rate regulates L1 longevity in C.

elegans. *PLoS One* **7**, e44720 (2012).

46. M. Ailion, J. H. Thomas, Dauer Formation Induced by High Temperatures in Caenorhabditis

elegans. *Genetics* **156**, 1047–1067 (2000).

47. C. Savage, *et al.*, Caenorhabditis elegans genes sma-2, sma-3, and sma-4 define a conserved

family of transforming growth factor beta pathway components. *Proc Natl Acad Sci U S A* **93**, 790–794

(1996).

48.   Y. Suzuki, *et al.*, A BMP homolog acts as a dose-dependent regulator of body size and male tail
patterning in Caenorhabditis elegans. *Development* **126**, 241–250 (1999).

49.   B. Gerisch, C. Weitzel, C. Kober-Eisermann, V. Rottiers, A. Antebi, A hormonal signaling pathway
influencing C. elegans metabolism, reproductive development, and life span. *Dev Cell* **1**, 841–851 (2001).

50.   M. B. Gerstein, *et al.*, Integrative analysis of the Caenorhabditis elegans genome by the
modENCODE project. *Science* **330**, 1775–1787 (2010).

51.   N. A. Noble, J. R. Harper, W. A. Border, In vivo interactions of TGF-beta and extracellular matrix.
Prog Growth Factor Res **4**, 369–382 (1992).

52.   U. Madaan, *et al.*, BMP Signaling Determines Body Size via Transcriptional Regulation of
Collagen Genes in Caenorhabditis elegans. *Genetics* **210**, 1355–1367 (2018).

53.   R. P. Boot-Handford, D. S. Tuckwell, Fibrillar collagen: the key to vertebrate evolution? A tale of
molecular incest. *Bioessays* **25**, 142–151 (2003).

54.   W. R. Wilcox, Connective Tissue and Its Heritable Disorders: Molecular, Genetic, and Medical
Aspects,. *Am J Hum Genet* **72**, 503–504 (2003).

55.   A. Wood, D. E. Ashhurst, A. Corbett, P. Thorogood, The transient expression of type II collagen
at tissue interfaces during mammalian craniofacial development. *Development* **111**, 955–968 (1991).

56.   N. Garamszegi, *et al.*, Extracellular matrix-induced transforming growth factor-beta receptor
signaling dynamics. *Oncogene* **29**, 2368–2380 (2010).

57.   W. Xin, J. Heilig, M. Paulsson, F. Zaucke, Collagen II regulates chondroycte integrin expression
profile and differentiation. *Connect Tissue Res* **56**, 307–314 (2015).

58.   L.-H. Chiu, *et al.*, Differential effect of ECM molecules on re-expression of cartilaginous markers
in near quiescent human chondrocytes. *J Cell Physiol* **226**, 1981–1988 (2011).

59.   R. Kerney, B. K. Hall, J. Hanken, Regulatory elements of Xenopus col2a1 drive cartilaginous
gene expression in transgenic frogs. *Int J Dev Biol* **54**, 141–150 (2010).

60.   J. R. Siebert, J. M. Graham, C. MacDonald, Pathologic features of the CHARGE association:
Support for involvement of the neural crest. *Teratology* **31**, 331–336 (1985).

61.   A. Antebi, W. H. Yeh, D. Tait, E. M. Hedgecock, D. L. Riddle, daf-12 encodes a nuclear receptor
that regulates the dauer diapause and developmental age in C. elegans. *Genes Dev* **14**, 1512–1527
(2000).

62.   C. L. Hale, A. N. Niederriter, G. E. Green, D. M. Martin, Atypical phenotypes associated with
pathogenic CHD7 variants and a proposal for broadening CHARGE syndrome clinical diagnostic criteria.
Am. J. Med. Genet. A **170**, 344–354 (2016).

63.   X. Chu, *et al.*, Genotranscriptomic meta-analysis of the CHD family chromatin remodelers in
human cancers – initial evidence of an oncogenic role for CHD7. *Mol Oncol* **11**, 1348–1360 (2017).

64.   M. Murakami, M. Koga, Y. Ohshima, DAF-7/TGF-β expression required for the normal larval
development in C. elegans is controlled by a presumed guanylyl cyclase DAF-11. *Mechanisms of*

Development **109**, 27–35 (2001).

65.   Y. Liu, *et al.*, CHD7 interacts with BMP R-SMADs to epigenetically regulate cardiogenesis in

mice. *Human Molecular Genetics* **23**, 2145–2156 (2014).

66.   M. Tewari, *et al.*, Systematic Interactome Mapping and Genetic Perturbation Analysis of a C.

elegans TGF-ᴺᴸ Signaling Network. *Molecular Cell*, 14.

67.   C. G. Riedel, *et al.*, DAF-16 employs the chromatin remodeller SWI/SNF to promote stress

resistance and longevity. *Nat. Cell Biol.* **15**, 491–501 (2013).

68.   K. Jia, P. S. Albert, D. L. Riddle, DAF-9, a cytochrome P450 regulating C. elegans larval

development and adult longevity. *Development* **129**, 221–231 (2002).

69.   J. S. Lee, *et al.*, FMRFamide-like peptides expand the behavioral repertoire of a densely

connected nervous system. *PNAS* **114**, E10726–E10735 (2017).

70.   X. Huang, J.-P. Saint-Jeannet, Induction of the neural crest and the opportunities of life on the

edge. *Developmental Biology* **275**, 1–11 (2004).

71.   H. Okuno, *et al.*, CHARGE syndrome modeling using patient-iPSCs reveals defective migration of

neural crest cells harboring CHD7 mutations. *Elife* **6** (2017).

72.   J. M. Kramer, "Extracellular Matrix" in *C. Elegans II*, 2nd Ed., D. L. Riddle, T. Blumenthal, B. J.

Meyer, J. R. Priess, Eds. (Cold Spring Harbor Laboratory Press, 1997) (March 22, 2021).

73.   M. Breuer, M. Rummler, C. Zaouter, B. M. Willie, S. A. Patten, Abnormal craniofacial and spinal

bone development with col2a1a depletion in a zebrafish model of CHARGE syndrome. *bioRxiv*,

2020.07.10.197533 (2020).

74.   T. Suzuki, D. Sakai, N. Osumi, H. Wada, Y. Wakamatsu, Sox genes regulate type 2 collagen

expression in avian neural crest cells. *Dev Growth Differ* **48**, 477–486 (2006).

75.   D. He, *et al.*, Chd7 cooperates with Sox10 and regulates the onset of CNS myelination and

remyelination. *Nature Neuroscience* **19**, 678–689 (2016).

76.   A. B. Roberts, B. K. McCune, M. B. Sporn, TGF-β: Regulation of extracellular matrix. *Kidney*

International **41**, 557–559 (1992).

77.   A. Tekari, R. Luginbuehl, W. Hofstetter, R. J. Egli, Transforming Growth Factor Beta Signaling Is

Essential for the Autonomous Formation of Cartilage-Like Tissue by Expanded Chondrocytes. *PLOS*

ONE **10**, e0120857 (2015).

78.   C. Chadjichristos, *et al.*, Down-regulation of human type II collagen gene expression by

transforming growth factor-beta 1 (TGF-beta 1) in articular chondrocytes involves SP3/SP1 ratio. *J Biol*

Chem **277**, 43903–43917 (2002).

79.   R. S. Kamath, J. Ahringer, Genome-wide RNAi screening in Caenorhabditis elegans. *Methods*

30, 313–321 (2003).

80.   L. Timmons, D. L. Court, A. Fire, Ingestion of bacterially expressed dsRNAs can produce specific

and potent genetic interference in Caenorhabditis elegans. *Gene* **263**, 103–112 (2001).

81.   S. K. Han, *et al.*, OASIS 2: Online application for survival analysis 2 with features for the analysis
of maximal lifespan and healthspan in aging research. *Oncotarget* **7**, 56147–56152 (2016).

82.   R. Ratnappan, *et al.*, Germline Signals Deploy NHR-49 to Modulate Fatty-Acid β-Oxidation and
Desaturation in Somatic Tissues of C. elegans. *PLoS Genetics* **10** (2014).

83.   Nieuwkoop, P.D., Faber, J., *Normal Table of Xenopus Laevis (Daudin) | Taylor & Francis Group*
(Garland Publishing, 1994).

84.   S. Huang, K. E. Johnson, H. Z. Wang, Blastomeres show differential fate changes in 8-cell
Xenopus laevis embryos that are rotated 90 degrees before first cleavage. *Dev Growth Differ* **40**, 189–
198 (1998).

85.   V. Gawantka, *et al.*, Gene expression screening in Xenopus identifies molecular pathways,
predicts gene function and provides a global view of embryonic patterning. *Mechanisms of Development*
77, 95–141 (1998).

86.   Q. Cao, Y. Shen, W. Zheng, H. Liu, C. Liu, Tcf7l1 promotes transcription of Kruppel-like factor 4
during Xenopus embryogenesis. *J Biomed Res* **32**, 215–221 (2018).

87.   M. W. Pfaffl, A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic
Acids Res* **29**, e45 (2001).

Jofré, DM and Hoffman, D *et al*, Figure 1

Figure 1: The DAF-12 regulated target *chd-7* is required for proper dauer morphogenesis

A) *chd-7(RNAi)* causes a partial dauer phenotype in *daf-2(e1370).* Representative DIC photomicrographs

of normal and partial dauers from *daf-2(e1370)* exposed to Control (L4440) or *chd-7* dsRNA, respectively.

B) Quantification of axial ratio of *daf-2(e1370);*Control(RNAi) and *daf-2(e1370);chd-7(RNAi)* dauers.

Three biological replicates were scored (n=16-21/replicate). Horizontal black lines represent mean with

SD. Unpaired t test, *p<0.05.

C) *daf-12* regulates *chd-7* expression. Representative images of *chd-7* transcriptional reporter, *tag-*

192::H1-wCherry or *tag-192::H1-wCherry;daf-12(rh61rh411)* worms at L2/L3 stage.

D) Relative expression of the translational reporter (n>10/strain). Unpaired t test, ****p<0.0001.

E) *C. elegans chd-*7 gene and protein. Top: *chd-7* genomic region. UTR and exons shown as bars;

introns by lines. In red, available *chd-7* deletional alleles (data obtained from CGC and NBRP). Bottom

images depict the predicted protein isoforms of *C. elegans* CHD-7, human CHD7 and human CHD8.

Signature domains in CHD proteins: two N-terminal chromodomains for interaction with a variety of

chromatin components (green), a SNF-2 like domain with ATPase activity (yellow) and a Helicase domain

(blue). The Class III subfamily is defined by a BRK domain (purple).

F) *chd-7*(*gk290*);*daf-2(e1370)* and *chd-7*(*tm6139*);*daf-2(e1370)* develop as SDS-sensitive dauer larvae*.*

n>725 animals/strain tested. Bars and horizontal black lines represent mean percentage with SD. Chi-

squared test with Bonferroni correction for multiple comparisons. ****p<0.0001. * represents the

comparison to the *daf-2(e1370)* strain

G) *chd-7* partial dauers fail to develop the dauer alae. Top row, representative photomicrographs of *daf-*

2(e1370) dauers or *chd-7;daf-2(e1370)* partial dauers expressing the *ajm-1*::GFP reporter to delineate the

seam cell borders (arrowheads mark a subset of junctions). Bottom row, SEM images of *daf-2(e1370)*

dauers or *chd-7;daf-2(e1370)* partial dauers (arrowheads mark alae details).

H) Germ cells in *chd-7;daf-2(e1370)* mutants overproliferate and arrest as L3-like germ cells*.*

Representative Z-projections *daf-2(e1370)* dauers or *chd-7;daf-2(e1370)* partial dauers stained with DAPI

(green). Arrowheads denote the anterior and posterior ends of the developing gonad arms.

Jofré, DM and Hoffman, D *et al*, Figure 2

Figure 2: *chd-7* affects longevity and response to pathogen.

A-E) *chd-7* promotes longevity in wild-type, *daf-2(e1370)* and *glp-1(e2144)* mutants. Mean survival days

on OP50-1; survival data analyzed using Kaplan−Meier test.

A) WT (18.12 days), *chd-7(gk290)* (14.91), *chd-7(gk306)* (16.53). *p<0.05 and ****p<0.0001 compared to

the wild-type, N2 strain.

B) WT (17.85), CHD-7::GFP (14.35). ****p<0.0001 compared to the wild-type, N2 strain.

C) *daf-2(e1370)* (35.58)*, chd-7(gk306);daf-2(e1370)* (37.18)*, chd-7(gk290);daf-2(e1370)* (17.47)*, daf-*

16(mu86);daf-2(e1370) (18.85)*. ****,^{††††}*p<0.0001. *represents comparison to *daf-2(e1370)*; ^{†}represents

comparison to *chd-7(gk306);daf-2(e1370)*.

D) *daf-2(e1370)* (27.9)*, chd-7(gk290);daf-2(e1370);CHD-7::GFP* (30.84)*, chd-7(gk290);daf-2(e1370)*

(19.13)*, daf-16(mu86);daf-2(e1370)* (14.48)*. ****,^{††††}*p<0.0001. *represents comparison to *daf-2(e1370);*

^{†}represents comparison to *chd-7(gk290);daf-2(e1370);CHD-7::GFP*.

E) *glp-1(e2144)* (30.4)*, chd-7(gk290);glp-1(e2144)* (28.37), *glp-1(e2144);daf-12(rh61rh411)* (25.99). ****p

<0.0001 and ^{†}p<0.05. *comparison to *glp-1(e2144)*; ^{†}comparison to *glp-1(e2144);daf-12(rh61rh411)*.

Details of number of animals and other data from replicates found in Supplemental Table 2.

F) *chd-7* mediates the response against the opportunistic bacteria *P. aeruginosa*. Mean lifespan in hours

(m) ± standard error of the mean (SEM). 'n' is the number of animals analyzed/total number in

experiment. WT (m = 52.11 ± 0.96, n = 94/162), *chd-7(gk290)* (m = 47.45 ± 1.05, n = 54.130/), *chd-*

7(tm6139) (m = 44.47 ± 1.57, n = 56/80), *daf-2(e1370)* (m = 105.46 ± 5.61, n = 65/115), *chd-*

7(gk290);daf-2(e1370) (m = 86.74 ± 4.21, n = 103/120) and *chd-7(tm6139);daf-2(e1370)* (m = 79.13 ±

4.9, n = 50/60). *,^{†}p<0.05; ***,^{†††}p <0.001 and ****p <0.0001. *comparison to N2; ^{†}comparison to *daf-*

2(e1370). Survival data analyzed using Kaplan−Meier test.

Genotype	% Reproductive *fog-2* females	SD
fog-2 (-males)	0	0
fog-2 (+males)	100	0
chd-7(gk290);him-5	100	0
chd-7(tm6139);him-5	0	0
chd-7(tm6139);him-5;chd-7::eGFP	87.5	17.67

Jofré, DM and Hoffman, D *et al*, Figure 3

Figure 3: CHD-7 belongs to the TGF-β signaling pathway.

A-B) Loss of *chd-7* suppresses dauer arrest of TGF-β pathway mutants at 25°C but not 26.5°C. 7 L4's

were plated individually and grown at the specified temperature for 1 week when arrested and non-

arrested progeny were scored. Bars and horizontal black lines represent mean percentage with SD.

Statistical significance was calculated using Chi-squared test with Bonferroni correction for multiple

comparisons. ****,[††††]p<0.0001 and £££ p <0.001.

A) Quantification of dauer arrest in *chd-7;daf-7(e1372)* mutants. * comparison to *daf-7(e1372)* grown at

25°C and [†]comparison to *daf-7(e1372)* at 26.5°C.

B) Dauer arrest in TGF-β pathway mutant backgrounds. *comparison to *daf-1(m213)* grown at 25°C,

[†]comparison to *daf-14(m77)* at 25°C; [£]comparison to *daf-1(m213)* at 26.5°C.

C) *chd-7* regulates body size. Body length of day 1 adults at 20°C (n>16). One-way Anova, **p<0.01,

***p<0.001 compared to wild-type, N2 strain.

D) *chd-7(tm6139)* males do not mate. 8 males from each strain tested were plated with 4 *fog-2* females

on 10 cm plates. After 24 hrs, *fog-2* females were transferred to new plates and within 48h the proportion

of fertile females were scored. Assay was repeated twice.

Jofré, DM and Hoffman, D *et al*, Figure 4

Figure 4: RNA-Seq analysis of transcriptome changes in *chd-7(gk290)* mutant dauers.

A) Heat map of expression values for the 84 differentially expressed genes (DEGs, cutoff of 0.05 on

FDR). DEGs were determined using DESeq2 (version 1.20.0). The color scale represents row z-score.

Hierarchical clustering of the DEGs is represented by dendrograms at the left.

B) Expression levels of *daf-9* mRNA are increased in *chd-7(gk290);daf-2(e1370)* partial dauers. Error

bars indicate standard error from three repeats with different biological samples. Two-tailed unpaired t-

test. **p <0.01.

C) *daf-9* knockdown in *chd-7;daf-7(e1372)* animals rescues dauer arrest at 25°C. 2-3 young L4's were

plated on to freshly seeded plates with either *daf-9* or Control empty vector RNAi and allowed to lay eggs.

After 72h, the adults were removed and the proportion of progeny that arrested as dauers was calculated.

Each dot represents a plate of at least 34 animals (n>598 total animals/ strain). Bars and horizontal black

lines represent mean percentage with SD. One-way ANOVA.

D) *daf-9* RNAi rescues arrest in *chd-7;daf-7(e1372)* animals but leads to partial dauers. Arrested animals

grown at 25°C on *daf-9* RNAi plates were treated with 1% SDS for 30 min and survival was scored.

n>159 total animals/ strain Each dot represents a plate of at least 30 animals and a minimum of 159 total

animals per strain tested. Bars and horizontal black lines represent mean percentage with SD. One-way

ANOVA, ****p <0.0001, compared to *daf-7(e1372)*.

E) Proposed mechanism of action for *chd-7* in dauer development. Under dauer-inducing conditions,

CHD-7 forms a transcriptional complex with DAF-3/DAF-5 to repress *daf-9* expression. In *chd-7* mutants,

daf-9 expression prevents dauer formation driven by reduced activity of both DAF-7/TGF-β and DAF-2/IIS

signaling pathways, presumably by binding dafachronic acids (DA) and subsequent activation of DAF-12.

Jofré, DM and Hoffman, D *et al*, Figure 5

Figure 5: Expression of *col2a1* rescues Chd-7 knock-down in *Xenopus* embryos.

A) Expression levels of *col2a1* mRNA are reduced in *chd7*-MO injected embryos. Both blastomeres of 2-cell staged embryos were injected with 5 or 10 ng of *chd7*-MO. Error bars indicate standard error from two repeats of the PCR reaction with different biological samples. One-tailed paired t-test, ***$p < 0.001$.

B) *Col2a1* expression domain is altered in the branchyal arches (ba) and otic vesicle (ov) of Chd7-depleted embryos. *In situ* hybridization of stage (St.) 35 embryos for *col2a1*. Lateral views, anterior to the left. One D1 blastomeres of 8-cell staged embryos was injected with 2.5 ng of St-MO or *chd7*-MO (N = 2; 42 and N = 3; 71, respectively) (N = number of experiments; number of embryos). Top and bottom panels are control and injected sides of the same representative *chd7*-MO injected embryo, respectively. e: eye. The graph is a quantification of the results. Reduced *col2a1* staining was observed in 18% of St-MO and 77% of *chd7*-MO injected embryos. Data on graph is presented as means with standard error. Fisher's exact test (****$p < 0.0001$). * represents comparison to St-MO group.

C) Lethality in Chd-7-depleted embryos is rescued by over-expression of *col2a1*. Graph showing the percentage of dead embryos. Both D1 blastomeres of 8-cell staged embryos were injected as indicated in the graph and scored for survival at St. 45. Data on graph is presented as means with standard error. Fisher's exact test ($^{†}p < 0.05$, **$p < 0.01$, $^{\&\&\&}p < 0.001$, ****$^{,††††,\#\#\#\#}$ $p < 0.0001$). * represents comparison to uninjected group, † represents comparison to St-MO group, $^{\#,\&}$ represent comparison to *chd7*-MO 5 ng and 10 ng, respectively.

D-E) Craniofacial morphometric analysis of *Xenopus* tadpoles at St. 45. Embryos were injected as indicated in panel C. Each dot represents a single embryo. Means and standard deviation are indicated. One-way ANOVA and Tukey's multiple comparisons test. D) Quantification of the eye size (top left panel). ***$^{,†††}p < 0.001$, ****$p < 0.0001$. E) Quantification of the eye distance (top left panel). $^{††}p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$. Lateral views, anterior to the left. * represents comparison to uninjected group and † represents comparison to *chd7*-MO 5 ng injected group. N = number of experiments, n = number of embryos.

SUPPLEMENTAL FIGURES

A

B

Figure S1: *chd-7* does not affects fat storage. Dauers of the shown genotypes were grown at 25°C for 5 days prior to lipid staining with Oil Red O (ORO). A. Representative photomicrographs of ORO-strained worms. B. Quantification of total area of ORO staining/ worm (Image J, see methods) reveals no significant differences in fat accumulation between control and *chd-7* mutants. Three biological replicates were scored with at least 16 individual dauers per replicate. Horizontal black lines represent mean with SD. P>0.05, Student's t test.

Figure S2: Loss of *chd-7* reduces L1 starvation survival in *daf-2(e1370)* mutants. L1 animals hatched from bleached egg preps into sterile M9 with 0.1µg/ml of streptomycin were diluted to a density of 1 larva/µl and kept at 20°C with constant agitation. Every 48h, 100µl aliquot was spotted onto a seeded NGM plate, and scored 72h later for development beyond the L2 larval stage. Percentage of the population alive was obtained by comparing the initial number of worms at t=0. Each dot represents at least 50 L1 worms.

Figure S3*: **swsn-1** and **swsn-8** share dauer suppression phenotypes with **chd-7**. *daf-7(e1372)* L1 synchronized larvae were placed on the indicated RNAi bacteria and grown at 25°C for 4 days to induce dauer formation. Dauer were identified after 4-5 days based on morphology and their resistance to 1% SDS for 15 min. Common SWI/SNF subunits and BAF or PBAF subclasses are indicated on top. Lines above bars represent standard error of the mean (SEM) from three independent experiments. At least 60 animals were assayed for each RNAi. One-way ANOVA. ****$p < 0.0001$. *comparison to the control, L4440 RNAi.

Figure S4: c*hd-7* mutants show altered gonad proliferation and migration. Whole mount fixation and DAPI (green) staining of late L4/young adult animals from the relevant strains. RNAi sample was obtained by growing *daf-2(e1370)* for two generations on *chd-7* dsRNA-producing E.Coli.

Figure S5: ChIP-seq analysis of CHD-7 and DAF-16 from ModEncode. Comparison of gene lists from peak calling of CHD-7::GFP (young adult) and DAF-16::GFP (L4) using Homer software. A) Venn diagram of ChIP peaks for CHD-7 and DAF-16. Overlap was defined as sharing at least one base in common. B) Common genes shared by CHD-7::GFP and DAF-16::GFP peaks. CHIP-seq data was obtained from publicly available data from the ModEncode project.

Figure S6: Craniofacial defects of *Chd7* knock-down *Xenopus* tadpoles. A) Gross morphology of surviving tadpoles at St. 45. Embryos were injected into both D1 blastomeres at the 8-cell stage with St-MO (5 ng) and *chd7*-MO (5 ng) as indicated. Tadpoles position is anterior to the left. Embryos are representative. B) Quantification of *Xenopus* tadpoles with craniofacial defects at stage 45. Embryos were injected as indicated in A and in the graph. Data on graph is presented as means with standard error. Fisher's exact test (**$p< 0.01$, ****,††††,####$p< 0.0001$). *comparison to uninjected group, † comparison to St-MO group, # comparison to *chd7*-MO group. N = number of experiments, n = number of embryos. C) Examples of Alcian blue-stained craniofacial skeletal elements from St. 45 tadpoles injected with St-MO (5 ng) and *chd7*-MO (5 ng). M: Meckel's, CH: ceratohyle and CB: ceratobranchial cartilages. *chd7*-MO injected tadpoles presented a high incidence of gross head malformations, reduced head and eyes sizes, and abnormal development of craniofacial cartilages.