1

Differential viral RNA methylation contributes to pathogen blocking in *Wolbachia*-colonized arthropods

Tamanash Bhattacharya[1,2], Liewei Yan[3], Hani Zaher[3] Irene L.G. Newton[1]*, Richard W. Hardy[1]*

[1]Department of Biology, Indiana University, Bloomington, IN, USA
[2]Basic Sciences Division, Fred Hutchinson Cancer Research Center, Seattle, WA, USA
[3]Department of Biology, Washington University, St. Louis, MO, USA

*Corresponding authors: ILGN, RWH

Email: irnewton@indiana.edu, rwhardy@indiana.edu

Tamanash Bhattacharya: 0000-0002-8129-7568
Irene L.G. Newton: 0000-0002-7118-0374
Richard W. Hardy: 0000-0001-6912-6291

Abstract

Arthropod endosymbiont *Wolbachia pipientis* is part of a global biocontrol strategy to reduce the replication of mosquito-borne RNA viruses such as alphaviruses. We previously demonstrated the importance of a host cytosine methyltransferase, DNMT2, in *Drosophila* and viral RNA as a cellular target during pathogen-blocking*.* Here we report on the role of DNMT2 in *Wolbachia*-induced alphavirus inhibition in *Aedes* species*.* Expression of DNMT2 in mosquito tissues, including the salivary glands, is elevated upon virus infection. Notably, this is suppressed in *Wolbachia*-colonized animals, coincident with reduced virus replication and decreased infectivity of progeny virus. Ectopic expression of DNMT2 in cultured *Aedes* cells is proviral, increasing progeny virus infectivity, and this effect of DNMT2 on virus replication and infectivity is dependent on its methyltransferase activity. Finally, examining the effects of *Wolbachia* on modifications of viral RNA by LC-MS show a decrease in the amount of 5-methylcytosine modification consistent with the down-regulation of DNMT2 in *Wolbachia* colonized mosquito cells and animals. Collectively, our findings support the conclusion that disruption of 5-methylcytosine modification of viral RNA is a vital mechanism operative in pathogen blocking. These data also emphasize the essential role of epitranscriptomic modifications in regulating fundamental alphavirus replication and transmission processes.

Introduction

Viruses are remarkably adept at using a limited set of viral factors to replicate in vastly different host cell environments. This ability is vital for the success of zoonotic arboviruses, which encounter physiologically and ecologically distinct invertebrate and vertebrate hosts during transmission. As these viruses oscillate between vertebrate and arthropod hosts, the progeny virions reared in one host cell context are primed for the next, predicating successful transmission. However, arbovirus transmission events are influenced by many host-specific biotic and abiotic factors (2-6). Recent studies have identified the vector microbiome as a critical biotic factor influencing arbovirus transmission (5, 6). One notable member of this microbial population is the arthropod endosymbiont *Wolbachia pipientis,* which dramatically impacts the transmission of multiple zoonotic arboviruses, a phenomenon termed "pathogen-blocking" (PB) (7-15). *Wolbachia* is transmitted transovarially and induces a wide range of reproductive manipulations in its host (16, 17). For example, *Wolbachia's* presence results in sperm-egg incompatibility between colonized and non-colonized individuals, mediated by a bacterially-encoded toxin-antitoxin system (17). This phenomenon, known as cytoplasmic incompatibility (CI), allows *Wolbachia* to be inherited at a rate higher than Mendelian inheritance, like a natural gene drive. Over the last

decade, scientists have leveraged this property to deploy *Wolbachia* as a novel vector control
agent with the aim of either suppressing or replacing the local mosquito population (18). Recent
data suggest that *Wolbachia* release programs significantly reduce the transmission of Dengue
virus (DENV) in endemic regions across 11 territories in Asia and Latin America (19, 20).
Remarkably, however, despite its success, the underlying cellular mechanism of pathogen-
blocking remains unidentified.

We recently showed that viral RNA is a cellular target of *Wolbachia*-mediated inhibition and that
loss in progeny virus infectivity occurs at the level of the encapsidated virion RNA, which is
compromised in its ability to replicate in naïve vertebrate cells (1). Mosquito-derived viruses,
reared in the presence of *Wolbachia,* are less infectious when seeded in either mosquito or
vertebrate cells. These results suggest a transgenerational mechanism by which *Wolbachia* limits
virus dissemination within the mosquito and subsequent transmission into vertebrates (1). We,
therefore, speculated that factor(s) regulating pathogen blocking likely target the viral plus sense
RNA genome, a feature shared between all viruses susceptible to *Wolbachia*-mediated inhibition.
Notably, the presence of *Wolbachia* reduces the infectivity of the encapsidated virion RNA in
mammalian cells (1). This observation alone suggests one or more RNA-targeting factors may be
responsible for compromising viral RNA replication in *Wolbachia*-colonized arthropod cells, as
well as in mammalian cells, which are devoid of *Wolbachia.* Prior work has implicated mosquito
exonuclease in pathogen-blocking, which is in line with the reduced half-life of incoming viral
RNAs in *Wolbachia*-colonized cells (1, 21). While faster degradation of viral RNA explains the
observed reduction in virus replication in arthropod cells, it does not explain reduced replication
in mammalian cells. In light of these findings, we sought to focus our attention on the RNA cytosine
methyltransferase DNMT2. Our prior study demonstrated that DNMT2 is essential for pathogen-
blocking in fruit flies (9). As an RNA modifying protein, DNMT2 does not directly antagonize viral
RNA replication but instead influences the cellular fate of its target(s). Both arthropods and
mammals encode proteins capable of interpreting epitranscriptomic signatures on different RNA
species, so we hypothesized that DNMT2-mediated modifications to the viral RNA in *Wolbachia*-
colonized arthropods impact virus replication in arthropod and mammalian cells.

To test this hypothesis, we investigated whether DNMT2 is essential for *Wolbachia*-mediated
pathogen blocking in mosquitoes. Additionally, we ask whether this MTase is functionally crucial
to virus regulation in the absence of *Wolbachia.* Given DNMT2's biological role as a cellular RNA
cytosine methyltransferase, we further examined the possibility of m5C modification of viral RNA
in mosquito cells and whether viral RNA is differentially modified in the presence and absence of
Wolbachia in mosquito cells. We find that *Wolbachia* and viruses differentially influence MTase
expression in mosquitoes. Specifically, the presence of the virus leads to elevated MTase
expression, which is proviral in mosquito cells. In contrast, the presence of *Wolbachia*
downregulates MTase levels to seemingly disrupt this proviral state, contributing to virus inhibition
as well as reduced progeny virus infectivity. Furthermore, the proviral effect is dependent on the
catalytic activity of DNMT2. Finally, as a consequence of this downregulation and DNMT2's role
as an RNA cytosine MTase, we show that the presence of *Wolbachia* in cells results in a reduced
abundance of 5-methylcytosine (m5C) modification of progeny viral RNA. These changes imply
that m5C modifications play a role in regulating viral RNA infectivity in mammalian cells. In
summary, our findings highlight a previously underappreciated role of RNA methylation in
alphavirus replication, with important implications for virus dissemination and transmission.
Overall, our results indicate a role of the viral epitranscriptome as regulatory signatures capable
of influencing the transmission of other arboviruses.

Results

Virus and *Wolbachia* differentially modulate *Aedes* DNMT2 expression

Wolbachia in *Aedes* mosquitoes is associated with reduced DNMT2 (*AMt2*) expression (22). We,
therefore, examined the expression of *AMt2* in *w*AlbB-colonized *Aedes aegypti* mosquitoes. We
chose to assess *in vivo AMt2* expression changes in whole mosquitos (which would give us a
sense of the *Mt2* environment encountered by disseminating viruses) or in dissected salivary
glands (the tissue important for transmission to the vertebrate host).

We measured *AMt2* expression in female *Aedes aegypti* mosquitoes colonized with and without
*Wolbachia (w*AlbB) five days post eclosion, forty-eight hours following bloodmeals with and
without Sindbis virus (SINV) (Fig 1B). The presence of both endosymbiont and virus was
associated with altered *AMt2* expression (Two-way ANOVA, $p < 0.0001$), with a nearly 100-fold
increase in *AMt2* levels in *Wolbachia*-free mosquitoes that received an infectious virus-containing
blood meal (Fig 1B. W-/V- compared to W-/V+; Two-way ANOVA, $p < 0.0001$). In contrast, we
found *Wolbachia* to reduce *AMt2* expression in mock-infected individuals (W+/V-) by
approximately 5-fold (Fig 1B, W-/V- compared to W+/V-). Importantly, we also observed low *AMt2*
expression in *Wolbachia*-colonized mosquitoes post-infectious (V+) bloodmeal*,* indicating that
Wolbachia prevents virus-induced stimulation of *AMt2* expression and that these levels are
maintained during infection. Virus replication in *Wolbachia*-colonized mosquitoes, therefore,
occurs in a low *AMt2* environment (Fig 1B. W-/V- compared to W+/V+). This pattern of reduced
AMt2 expression was also observed *ex vivo* in cultured *Aedes albopictus*-derived mosquito cells
colonized with both a native (*w*AlbB strain in Aa23 cells) and a non-native *Wolbachia* (*w*Mel in
RML12 cells) strain (Fig S1).

Fig 1. Virus and *Wolbachia* each differentially modulate expression of the RNA methyltransferase gene DNMT2 in mosquitoes. (A) Schematic of virus dissemination (white arrows) within mosquito tissues and transmission into vertebrate host (black arrow). (B) *AMt2* expression was measured 5 days post bloodmeal in whole female mosquitoes using qRT-PCR with and without SINV. Error bars represent standard error of mean (SEM) of biological replicates. Two-way ANOVA with Tukey's post-hoc test of multivariate comparisons. (C) *AMt2* expression measured in dissected salivary gland tissues collected from female mosquitoes with and *Wolbachia*-free 5 days post bloodmeal with or without SINV. Error bars represent standard error of mean (SEM) of biological replicates. Two-way ANOVA with Tukey's post-hoc test of multivariate comparisons. (D) Viral RNA levels were quantified in dissected salivary gland tissues with and *Wolbachia*-free using qRT-PCR at 5 days post infectious blood meal with SINV. Unpaired, student's t-test, error bars represent standard error of mean (SEM) of biological replicates. For all panels: ****$P < 0.0001$; ***$P < 0.001$; **$P < 0.01$; ns = non-significant.

We next quantified *AMt2* expression in isolated salivary gland tissues from five-day-old *Aedes*
aegypti mosquitoes colonized with or *Wolbachia*-free *(w*AlbB), post bloodmeal with (V+) or without
(V-) SINV (Fig 1C). *AMt2* expression in the salivary gland tissues post-infectious bloodmeal was
elevated nearly 100-fold, similar to the increase induced by the virus in whole mosquitoes (Fig
1C. W-/V- compared to W-/V+). As before, the presence of *Wolbachia* alone was associated with
lower *AMt2* expression. However, this difference was not statistically significant (Fig 1C. W-/V-
compared to W+/V-). Importantly, however, *Wolbachia* did prevent SINV-induced *AMt2*
upregulation, reducing it 100-1000 fold (Fig 1C. W-/V+ compared to W+/V+). Under these
conditions, we also observed a significant 2 to 3 log$_{10}$ reduction in viral RNA in the salivary gland

tissues (Fig 1D). It should be noted that our observations regarding the effect of *Wolbachia*
(*w*AlbB) or SINV on *AMt2* expression are analogous to previous reports that describe differential
AMt2 expression in the presence of the flavivirus DENV-2 and *Wolbachia* (*w*Mel) in *Aedes aegypti*
mosquitoes (22).

DNMT2 promotes virus infection in mosquito cells

The positive correlation between *AMt2* expression and SINV genome replication in *Aedes*
mosquitoes (Fig 1B-D) led us to examine whether there is a functional consequence of elevated
MTase expression on virus infection in these insects. We therefore ectopically expressed *AMt2*
and assessed its effect on virus infection in cultured *Aedes albopictus* cells (Fig 2A), using
azacytidine-immunoprecipitation (AZA-IP) to determine whether viral RNA in the cell is a direct
DNMT2 target. *Wolbachia*-free *Aedes albopictus* (C710) cells were transfected with an epitope-
tagged *AMt2* expression vector (FLAG-*AMt2*) or control vector (FLAG-empty) for 48 hours before
infection with SINV at an MOI of 10. After 24 hours post-infection, cells were labeled with a
cytosine analog, 5-Azacytidine (5-AZAC), for 18 hours to incorporate the label into newly
synthesized cellular and viral RNA. We reasoned that if mosquito DNMT2 directly targets viral
RNAs for methylation, the presence of 5-AZAC in the RNA should covalently trap the enzyme
forming a stable m5C-DNMT2-viral RNA complex, allowing co-immunoprecipitation of the RNA-
protein complexes using anti-FLAG antibody (23). Targeted quantitative RT-PCR analyses of total
immunoprecipitated RNA revealed enrichment of SINV RNA relative to a control host RNA
transcript (GAPDH), confirming that viral RNA is indeed a direct MTase target in these cells (Fig
2B).

We next assessed the effect of elevated *AMt2* expression on RNA virus infection by measuring
the output of infectious progeny viruses following infection of cells expressing FLAG-*AMt2*.
Ectopic MTase expression resulted in a four-fold increase in SINV titer, further supporting the
positive *in vivo* correlation between *AMt2* expression and virus replication observed previously
(Fig 2C). We also observed a concomitant increase in the per-particle infectivity of viruses upon
assaying them on vertebrate baby hamster kidney fibroblast cells, as evidenced by higher specific
infectivity ratios (Fig 2D). Together these results support the idea of *Aedes* DNMT2 being a
proviral factor exploited by the virus to enhance its replication and transmission in the mosquito
vector.

Fig 2. Overexpressing *AMt2* in mosquito cells improve progeny virus infectivity. (A) Western Blot of *Aedes albopictus* (C710) cells transfected with expression vector constructs with (FLAG-AMt2) or without (FLAG-empty) *AMt2.* Cytoplasmic lysates of cells were collected 48 hours post transfection and probed with anti-FLAG and anti-β actin antibodies. (B) Relative levels of SINV RNA recovered following AZA-IP of AMt2 in C710 cells was quantified using qRT-PCR. *Wolbachia*-free C710 mosquito cells were transfected with expression vectors FLAG-empty or FLAG-AMt2 for 48 hours prior to infection with SINV at MOI of 10. Cells were treated for approximately 18h with 5 μM 5-Azacytidine to covalently trap AMt2 with its target cellular RNA prior to RNA immunoprecipitation using anti-FLAG antibody. The horizontal dotted line represents the threshold set at 1 ($\log_{10}$). Unpaired two-tailed t-test with Welch's correction, p = 0.0004, t = 4.216, df = 20 (C) Infectious progeny (PFU/mL) SINV produced from mosquito cells *Wolbachia*-free expressing either FLAG-empty (w/o Wolb) or FLAG-AMt2 (w/o Wolb + AMt2). Cells were transfected 48 hours prior to infection with SINV at MOI of 10. Infectious progeny viruses collected from supernatants 48 hours post-infection were quantified using plaque assays on BHK-21 cells. Unpaired two-tailed t-test with Welch's correction, p = 0.0002, t = 5.404, df = 11.81. (D) Specific Infectivity Ratios of progeny SINV were calculated as described earlier (1). Unpaired two-tailed t-test with Welch's correction, p = 0.0084, t = 3.911, df = 5.820. For all panels error bars represent standard error of mean (SEM) of biological replicates and **P < 0.01; ****P < 0.0001.

AMt2 expression is reduced in the presence of *Wolbachia.* To test whether virus restriction *in vivo*
is a consequence of reduced DNMT2 activity, we measured virus replication in mosquito cells
following pharmacological inhibition of DNMT2. Structural homology of DNMT2 to other members
of the DNA MTase family has allowed it to retain its DNA binding ability *in vitro.* However*,* they
are canonically known to methylate tRNA molecules (24)*.* Furthermore, DNMT2 is known to
methylate RNA substrates by a different mechanism than canonical RNA methyltransferases.
This mechanism of action makes DNMT2 susceptible to the action of DNA methyltransferase

inhibitors ribo- (5-Azacytidine or 5-AZAC) or deoxyribo- (Deoxy-5-Azacytidine or DAC5) while
ensuring that the function of other RNA methyltransferases in the cell remain unperturbed (25,
26). We reasoned that pretreatment of mosquito cells with either MTase inhibitor should reduce
cellular DNMT2 activity and consequently restrict alphavirus replication. Pretreating *Wolbachia*-
free C710 cells with RNA MTase inhibitor 5-AZAC prior to infection reduced SINV RNA replication
approximately 5-fold at 24 hours post-infection (Fig 3A). Virus titer was also reduced
approximately 10-fold (Fig 3B). Finally, MTase inhibition also negatively influenced SINV per-
particle infectivity, as evidenced by a 50-fold reduction in SI ratio (Fig 3C). Similar results were
obtained for related alphavirus, Chikungunya virus (CHIKV, Fig S2A,B).

Using our previously published live-cell imaging system, we used a fluorescently-tagged CHIKV
reporter virus (CHIKV-mKate) to examine the effect of deoxyribo- MTase inhibitor DAC5 on virus
replication in *Wolbachia*-free and *Wolbachia*-colonized *Aedes albopictus* cells using the Incucyte
live-cell imaging platform (1). As before, fluorescent protein expression was used as a proxy of
virus replication in cells with (DAC5) and without (DMSO) inhibitor pretreatment. Virus replication
was measured over 50 hours by quantifying mean virus-encoded red fluorescent reporter (mKate)
expression observed over four distinct fields of view taken per well every 2-hours (Fig 3D, S2C).

Fig 3. Pharmacological inhibition of mosquito DNMT2 reduces virus replication and per-particle infectivity. Inhibition of mosquito DNMT2 in *Wolbachia*-free *Aedes albopictus* derived C710 cells was carried out using MTase inhibitors 5-Azacytidine (5-AZAC). Dimethyl-sulfoxide (DMSO) was used as the negative control. In each case, cells were pretreated with 5 µM inhibitors overnight prior to infections with SINV at MOI of 10. Cell lysates and supernatants were harvested at 24 hours post infection to quantify cellular viral RNA levels and infectious titer, respectively. (A) Levels of SINV RNA in mosquito cells treated with MTase inhibitor 5-AZAC were determined using quantitative RT-PCR. Unpaired two-tailed t-test with Welch's correction, SINV: p = 0.0012, t = 6.742, df = 4.892. (B) Infectious SINV titers produced from mosquito cells treated with MTase inhibitor 5-AZAC were determined using plaque assays on BHK-21 cells. Unpaired two-tailed t-test with Welch's correction, SINV: p = 0.0339, t = 4.541, df = 2.322. (C) Specific infectivity ratios of progeny SINV was calculated as the ratio of infectious plaque forming units (B) over total viral genome copies present in collected cell supernatants as quantified by qRT-PCR. Unpaired two-tailed t-test with Welch's correction, SINV: p = 0.0002, t = 12.59, df = 3.946. Error bars represent standard error of mean (SEM) of three independent experimental replicates. (D) CHIKV expressing mKate fluorescent reporter protein was grown in C710 *Aedes albopictus* cells in the presence (W+ virus) or absence (W- virus) of *Wolbachia (*strain *w*Stri*)*. These progeny viruses were then used to infect C710 cells without and with *Wolbachia (*strain *w*Stri) pretreated with MTase inhibitor DAC5 (black data points) or DMSO (blue data points for *Wolbachia* free cells, red data points for *Wolbachia* colonized cells) synchronously at a MOI of 1 particle/cell. Virus growth in cells was measured in real time by imaging and quantifying the number of red cells expressing the virus encoded mKate protein forty-eight hours post infection, using live cell imaging. Shape of data points represent the origin of virus used to initiate infection; circles represent viruses derived from W- cells, boxes represent viruses derived from W+ cells. Three-way ANOVA with Tukey's post hoc test for multivariate comparisons. For all panels error bars represent standard error of mean (SEM) of independent experimental replicates (n=3). *P < 0.05; **P < 0.01; ****P < 0.0001, ns = non-significant.

In line with previous observations, viruses derived from *Wolbachia*-colonized cells (W+ virus, we
will refer to viruses derived from *Wolbachia*-colonized cells as W+ and their counterparts, derived
from *Wolbachia*-free cells as W- virus), produced under low *AMt2* conditions, are less infectious
on W- cells, limiting their dissemination (1). Notably, this phenotype is particularly pronounced
when W+ viruses encounter cells also colonized with the endosymbiont, presumably exhibiting
reduced *AMt2* expression (Fig 3D, S2D-E). To further investigate the importance of DNMT2
activity in *Wolbachia*-mediated virus inhibition we treated cultured *Aedes albopictus* (C710) cells
with DAC5, a DNMT2 inhibitor. We predicted that replication kinetics of W+ viruses in inhibitor-
treated *Wolbachia*-free (W-) cells should phenocopy kinetics of W+ virus replication in *Wolbachia*-
colonized (W+) cells. Additionally, the kinetics of W- virus replication in inhibitor-treated
Wolbachia-free (W-) cells should phenocopy W- virus replication in *Wolbachia*-colonized (W+)
cells (1). Three-way ANOVA was used to determine the effect of MTase inhibitor (DAC5), progeny
virus type (derived from producer cells with or *Wolbachia*-free*),* and/or time on virus replication in
recipient cells (assaying cells with or *Wolbachia*-free*)*. In the presence of MTase inhibitor,
replication of both W- and W+ viruses was reduced throughout infection, with a greater decrease
in the replication of W+ viruses relative to W- viruses phenocopying the replication of W+ viruses
in *Wolbachia*-colonized cells. Finally, replication of W- viruses in the presence of inhibitor was
comparable to that of W+ viruses in mock-treated *Wolbachia*-free cells. We observed no
synergistic effect of virus source and MTase inhibitor on virus replication in *Wolbachia*-colonized
cells, likely due to low mean reporter activity. Altogether, these results demonstrate that
alphavirus replication is negatively impacted by perturbed DNMT2 activity in either producer (W+
virus) or recipient mosquito cells (W- + DAC5 cells or W+ cells) and that the effect is compounded
when both co-occur (W+ virus in W- + DAC5 cells or W+ virus in W+ cells).

Ectopic DNMT2 expression rescues alphaviruses from *Wolbachia*-mediated inhibition

Evidence gathered indicates that *AMt2* downregulation is responsible for pathogen blocking in
mosquitoes. Therefore, we ectopically overexpressed *AMt2* in *Wolbachia*-colonized mosquito
cells to alleviate virus inhibition, including disruption of viral RNA synthesis and progeny virus
infectivity (Fig 4A). We observed a significant reduction in viral RNA levels in *Wolbachia*-colonized
cells relative to *Wolbachia*-free cells (Fig 4B). Interestingly, expression of FLAG_*Amt2* increased
SINV RNA levels 70-fold in *Wolbachia*-colonized cells compared to cells carrying FLAG-empty
vector, restoring virus RNA synthesis; One-way ANOVA Holm-Sidak's multiple comparisons test,
w/ Wolb vs. w/ Wolb + AMt2, p < 0.0001 (Fig 4B). Additionally, we observed a significant
improvement in per-particle infectivity of progeny viruses derived from *Wolbachia*-colonized cells
ectopically expressing *AMt2* (W+ AMt2+ virus, Fig 4C). Therefore, both phenotypes of pathogen
blocking were abrogated upon *AMt2* over-expression. Given that endosymbiont titers can
influence the degree of virus inhibition, we checked whether altering *AMt2* levels significantly
impacted *Wolbachia* titer in cells. Quantitative PCR was used to measure relative *Wolbachia* titer
in cells transfected with FLAG-AMt2 or FLAG-empty. However, no changes in endosymbiont titer
were observed following ectopic *AMt2* expression (Fig 4D). Therefore, changes in *Wolbachia* titer
do not explain the loss of pathogen blocking.

Fig 4. ***AMt2* overexpression in *Wolbachia*-colonized cells rescues virus from endosymbiont-mediated inhibition.** C710 cells with *Wolbachia* were transfected with expression vectors FLAG-empty (w/ Wolb) or FLAG-AMt2 (w/ Wolb + AMt2) for 48 hours prior to infection with SINV-nLuc at MOI of 10. *Wolbachia* -free cells expressing FLAG-empty (w/o Wolb) were used as a positive control. (A) Schematic of experimental workflow. (B) Viral genome replication in C710 cells was quantified using qRT-PCR using extracted total RNA from infected cell lysates. One-way ANOVA with Tukey's post-hoc test of multivariate comparison. (C) Specific Infectivity Ratios of progeny viruses produced from the aforementioned infection was calculated as described earlier (1). Briefly, infectious progeny viruses collected from supernatants 48 hours post infection were quantified using plaque assays on BHK-21 cells, while total number of progeny virus particles was quantified via qRT-PCR of viral genome copies released into the supernatant. Error bars represent standard error of mean (SEM). One-way ANOVA with Tukey's post-hoc test of multivariate comparison, w/ Wolb vs w/ Wolb + AMt2, p = 0.0003, w/o Wolb vs w/ Wolb, p < 0.0001. (D) C710 mosquito cells with *Wolbachia* were transfected with expression vectors FLAG-empty (w/ Wolb) or FLAG-AMt2 (w/ Wolb + AMt2) for 48 hours prior to quantification of endosymbiont titer via quantitative PCR using DNA from extracted cell lysates. Error bars represent standard error of mean (SEM). Unpaired, student's t-test, p = 0.1316, t = 1.794, df = 5.097. Statistically non-significant values are indicated by ns. (E) Progeny viruses were used to synchronously infect naïve BHK-21 cells at equivalent MOIs of 5 particles/cell. Cell lysates were collected at indicated times post infection and luciferase activity (RLU), was used as a proxy for viral replication. Two-way ANOVA with Tukey's post-hoc test of multivariate comparison, Time: p < 0.0001, *Wolbachia/AMt2*: p = 0.0003, Time x *Wolbachia/AMt2*: p < 0.0001. (F) Approximately 10^5 copies (determined using qRT-PCR) each of virion encapsidated RNA extracted from the aforementioned W+, W+ AMt2 and W- viruses were transfected into naïve BHK-21 cells and infectious titer was determined by the counting the number of plaques produced after 72 hours post transfection. Numbers above bars refer to the proportion of samples that formed quantifiable plaque-forming units on BHK-21 cells. One-way ANOVA with Tukey's post-hoc test of multivariate comparison. (G) 10^5 copies each of virion encapsidated RNA extracted from the W+, W+ AMt2 and W- viruses were transfected into naïve BHK-21 cells and luciferase activity (RLU) was used as a proxy for viral replication at 9 hours post-transfection. Numbers above bars refer to the proportion of samples that produced luciferase signal above background levels, indicated by the dotted line. One-way ANOVA with Tukey's post-hoc test of multivariate comparison, w/ Wolb vs w/ Wolb + AMt2: p < 0.00001, w/o Wolb vs w/ Wolb: p < 0.0001; w/o Wolb vs w/ Wolb + AMt2: p = 0.991. For all panels error bars represent standard error of mean (SEM). *P < 0.05; **P < 0.01; ****P < 0.0001.

Reduced per-particle infectivity of viruses occurring in the presence of *Wolbachia* (W+ viruses) is
associated with reduced replication kinetics of these viruses in vertebrate cells and reduced
infectivity of the encapsidated W+ virion RNA (1). As *AMt2* overexpression in *Wolbachia*-
colonized mosquito cells rescued viral RNA synthesis and progeny virus infectivity, we examined
the ability of progeny viruses derived from *Wolbachia*-colonized cells overexpressing *AMt2* (W+
derived AMt2+ cells) to replicate in vertebrate cells. We used SINV encoding a luciferase reporter
for these experiments, allowing viral replication kinetics to determine the following synchronous
infection of three progeny virus types: W- derived virus, W+ derived virus, and W+ derived AMt2+
virus. Replication of W+ derived AMt2+ viruses was significantly higher on a per-particle basis
relative to W+ derived and, interestingly, W- derived viruses. This could be due to higher ectopic
AMt2 expression relative to what is induced natively during virus infection, implying perhaps a
dose-dependent effect (Fig 4E). We then examined whether ectopic *AMt2* expression caused
changes in the infectivity of the encapsidated virion RNA itself. Based on results from Fig 3D, we
hypothesized that ectopic *AMt2* expression in *Wolbachia*-colonized cells should restore virion
RNA infectivity. Indeed, following transfection of virion RNA into vertebrate BHK-21 cells, W+
derived virus RNA was largely non-infectious in contrast to RNA derived from viruses derived
from W- and W+ AMt2+ cells (Fig 4F). Restored infectivity of W+ AMt2+ derived viral RNA was
also validated using the luciferase-based virus replication assay (Fig 4G).

As demonstrated in Fig 2B, DNMT2 possesses the ability to bind viral RNA in mosquito cells.
However, this alone does not indicate whether its MTase activity is essential for its proviral role.
Broadly, DNMT2 comprises a catalytic domain and a target recognition domain responsible for
RNA binding (27, 28). It is, therefore, possible that DNMT2's regulatory role is independent of its
MTase activity. To determine the importance of catalytic activity, we overexpressed a catalytically-
inactive mutant of *AMt2*, replacing the highly conserved cysteine residue (C78) present in the
motif IV region with a glycine (*AMt2* C78G, Fig 5A) in *Wolbachia*-colonized mosquito cells and
asked whether this allele is capable of relieving pathogen blocking. Our data show *AMt2*-mediated
rescue of SINV RNA synthesis and infectivity depends on its MTase activity as expression of the
C78G mutant failed to rescue virus from *Wolbachia*-mediated inhibition (Fig 5B). We observed no
improvement in SINV infectivity under these conditions (Fig 5C). Similar results were obtained
from experiments carried out using CHIKV, where expression of wild-type *AMt2*, but not *AMt2*-
C78G, resulted in improved virus titer (Fig 5D). However, while expression of *AMt2* increased
CHIKV specific infectivity compared to expression of *AMt2-C78G* this increase was not
statistically significant (Fig 5E). Based on these results, we conclude that the MTase *AMt2*
promotes virus infection in mosquitoes and that lower *AMt2* expression in the presence of
Wolbachia contributes to virus restriction and that MTase activity of DNMT2 is required for proviral
function.

Fig 5: Catalytically inactive DNMT2 is unable to rescue *Wolbachia*-mediated virus inhibition in mosquito cells. (A) Multiple sequence alignment of the Motif IV region of DNMT2 derived from dipterans that are known to be colonized with native or non-native *Wolbachia.* The conserved catalytic cysteine (C) residue, depicted in red on the consensus sequence at the top, was mutated to a glycine (G) to abolish MTase activity of mosquito AMt2. Expression of the catalytic mutant (AMt2 C78G) was determined at 48 hours post transfection using Western Blot. C710 mosquito cells transfected with expression vector constructs with (FLAG-AMt2 C78G) or without (FLAG-empty) *AMt2.* Cytoplasmic lysates of cells were collected 48 hours post transfection and probed with anti-FLAG antibody. The non-specific band appearing below the desired size was used as a loading control. (B and C) C710 mosquito cells with *Wolbachia* were transfected with expression vectors FLAG-empty (w/ Wolb) or FLAG-C78G AMt2 (w/ Wolb + AMt2 C78G) for 48 hours prior to infection with SINV at MOI of 10. Infectious virus and Specific Infectivity (SI) of progeny viruses produced after 72 hours post infection were quantified as before. Unpaired t-test with Welch's correction, p = 0.1734, t = 1.920, df = 2.396 (B), p = 0.4544, t = 0.8291, df = 3.937 (C). (D and E) C710 mosquito cells with *Wolbachia* were transfected with expression vectors FLAG-empty (w/ Wolb), FLAG-AMt2 (w/ Wolb + AMt2) or FLAG-C78G AMt2 (w/ Wolb + AMt2 C78G) for 48 hours prior to infection with CHIKV-mKate at MOI of 10. Infectious virus (D) and Specific Infectivity (E) of progeny viruses produced after 72 hours post infection were quantified as before. One-way ANOVA followed by Tukey's post hoc test for multivariate comparisons for (D), w/ Wolb vs w/ Wolb + AMt2, p = 0.0009, w/ Wolb vs w/ Wolb + AMt2 C78G p = 0.2694, w/ Wolb + AMt2 vs w/ Wolb + AMt2 C78G p < 0.0040. Error bars represent standard error of mean (SEM) of biological replicates. **P < 0.01; ***P < 0.001, ns = non-significant.

 DNMT2 orthologs from mosquitoes and fruit flies regulate virus infection differentially in their respective hosts

 The proviral role of the *AMt2* is intriguing, given the previously described antiviral role for the corresponding fruit fly ortholog, *Mt2* (9, 29). Interestingly, in our previous study, we observed that knocking down *Mt2* led to increased progeny virus infectivity. Therefore, we reasoned that ectopic

expression of *Mt2* should reduce Sindbis virus infectivity (Fig S3). As with mosquito *AMt2,* we
asked whether this involved direct targeting of viral RNA. AZA-IP of epitope-tagged Mt2 confirmed
direct interactions between viral RNA and fly DNMT2 in *Wolbachia*-free *Drosophila melanogaster-*
derived JW18 cells, which showed a 10-fold enrichment in SINV RNA-binding relative to a control
host transcript (18S) (Fig S3A, B).

In contrast to the proviral effect of mosquito *AMt2*, ectopic *Mt2* expression significantly reduced
infectivity of progeny SINV and CHIKV (W- derived Mt2+ virus) relative to those produced from
cells expressing the control vector (W- derived virus), confirming our previous findings (Fig S3C).
As with *AMt2,* we assessed whether reduced infectivity of W- derived Mt2+ viruses was due to
their inability to replicate in vertebrate cells. Indeed, results from our luciferase reporter based
viral replication assay revealed significantly reduced replication of W- Mt2+ derived viruses
relative to W- derived viruses in vertebrate BHK-21 cells, similar to the behavior observed for W+
derived viruses (Fig S3D). Finally, we quantified the infectivity of virion encapsidated RNA from
W- derived Mt2+ SINV and CHIKV viruses by measuring the number of plaque-forming units
generated following transfection into vertebrate BHK-21 cells. For both SINV and CHIKV, the
infectivity of virion encapsidated RNA was reduced for W+ viruses. Notably, this was phenocopied
by virion RNA isolated from W- Mt2+ derived SINV and CHIKV (Fig S3E,F).

Similar to mosquito *AMt2,* fly *Mt2's* ability to regulate virus fitness also rely on its catalytic activity,
as expressing a catalytically inactive mutant (*Mt2* C78A) was unable to restrict the production of
infectious virus and per-particle infectivity of SINV (Fig S4). Taken together, these results suggest
that progeny virus/virion RNA infectivity is reduced in fly cells under conditions where MTase
expression is elevated natively in the presence of *Wolbachia* (W+ virus) or artificially (W- Mt2+
virus).

The presence of *Wolbachia* in mosquito cells is associated with altered viral RNA
methylation

That ectopic *AMt2* expression in *Wolbachia*-colonized *Aedes albopictus* cells can restore the
infectivity of SINV progeny virion RNA implies two important things; (i) Sindbis virion RNA carries
5-methylcytosine (m5C) modifications, and (ii) that altered *AMt2* expression in the presence of
Wolbachia is associated with changes in the overall m5C content of the virion RNA. To directly
determine if virus RNA is modified differentially in the presence of *Wolbachia*, we subjected virion
RNA isolated from progeny SINV produced from *Aedes albopictus* cells colonized with (W+ virus)
and without (W- virus) *Wolbachia* to liquid chromatography tandem mass spectrometry (LC-
MS/MS) analyses (Fig 6A). We chose to focus our efforts on identifying the presence of 5-
methylcytosine (m5C) and 6-methyladenosine (m6A) residues on the viral genome for our present
analyses. We examined m6A due to recent reports highlighting the importance of this modification
in regulating RNA virus replication (30, 31). A potential complication for these analyses is the
presence of residue(s) of similar mass to charge ratio(s) to m5C, such as m3C. However, as
shown in Fig S5A, we observed distinct distribution of the individual m3C and m5C peaks in the
spectral output, demonstrating our ability to distinguish between these two bases (Fig S5A).

Figure 6: Presence of *Wolbachia* is associated with altered virion RNA methylation. (A) RNA isolated from progeny viruses derived from mosquito cells colonized with (W+ virus) or without (W- virus) *Wolbachia* were subjected to LC-MS/MS analyses to determine their nucleotide content. (B) Normalized 5-methyl cytosine (M5C) content of RNA isolated from W- and W+ viruses represented as a ratio of total unmodified cytosine content. Unpaired two-tailed t-test, p = 0.0013, t = 8.080, df = 4 (C) Normalized 6-methyl adenosine (M6A) content of RNA isolated from W- and W+ viruses represented as a ratio of total unmodified adenosine content. Unpaired two-tailed t-test, p = 0.666, t = 0.4643, df = 4. Error bars represent standard error of mean (SEM) of three independent virus preps from each cell type. **P < 0.01; ns = non-significant.

LC-MS/MS analyses of RNA purified from virion RNA derived from *Wolbachia* free (W-) and
Wolbachia-colonized (W+) cells demonstrated W+ virion RNA to contain, on average, more than
2-fold fewer m5C residues compared to W- virion RNA across three independent virus preps from
each cell type (Fig 6B). Notably, both W+ and W- virion RNA was determined to consist of
comparable levels of m6A residues across all biological replicates (Figure 6C). In addition, we
observed no significant changes in the overall m3C content between W+ and W- virion RNA (Fig
S5B). It should be noted that while we did not observe changes in the overall abundance of m6A
and m3C residues between W+ and W- virion RNA, it is unclear whether the presence of
Wolbachia leads to altered distribution of m6A and/or m3C modifications in the context of the
overall SINV RNA sequence. Finally, we used LC-MS/MS analyses to quantify viral Type-0 (7-
methyl-GpppNp or M7G) cap structures present in W- and W+ virion RNA to estimate relative
ratios of capped versus non-capped virus progeny produced in the presence or absence of
Wolbachia. While there were no statistically significant differences present between the respective

W- and W+ sample means (t-test: p=0.999), we found M7G content to vary significantly among
W+ virion RNA replicates (F-test: p=0.0088), indicating either that the ratios of capped vs. non-
capped viruses vary significantly within virus populations derived from *Wolbachia*-colonized cells,
or that viral RNAs produced under these conditions carry varying amounts of internal M7G
signatures (Fig S5C).

These data are consistent with: (i) DNMT2 being an essential host factor in mosquito cells for
efficient virus replication and transmission, (ii) this proviral effect being exert through m5C
modification of the viral genomic RNA, and (iii) a mechanism of *Wolbachia*-mediated pathogen
blocking being the reduction of DNMT2 expression.

Discussion

Virus inhibition in *Wolbachia*-colonized arthropods is associated with two distinct features
independent of any particular host-*Wolbachia* strain combination; (i) reduced genome replication
of the +ssRNA viruses in *Wolbachia*-colonized cells, and (ii) reduced per-particle infectivity of
progeny +ssRNA viruses produced under these conditions (1). While these shared attributes
constitute a subset of several virus inhibition phenotypes, it indicates the existence of a conserved
cellular mechanism of restriction. In our previous study, we used the prototype alphavirus, Sindbis
as our +ssRNA virus model to uncover an essential role of the fruit fly RNA cytosine
methyltransferase (MTase) gene *Mt2* (DNMT2) as an essential host determinant of *Wolbachia*-
mediated pathogen blocking (9). Furthermore, loss of *Drosophila* DNMT2 is associated with a
loss in virus inhibition by *Wolbachia* and increased progeny virus infectivity in mammalian cells,
suggesting that DNMT2 might regulate these two aspects of alphavirus replication. These findings
thus led us to ask the following question in our present study: Is DNMT2 a conserved host
determinant of *Wolbachia*-mediated +ssRNA virus inhibition between fruit flies and mosquitoes?

MTase expression in adult *Aedes aegypti* mosquitoes is distinctly altered in the presence of both
virus and *Wolbachia* and in opposite directions (Fig 1). DNMT2 expression is elevated in the body
of mosquitoes following an infectious bloodmeal. Notably, this pattern is observed in the salivary
gland tissues, representing the final site of virus production in the vector prior to transmission to
a vertebrate. We show that this is beneficial to the virus, as ectopic MTase expression in cultured,
Wolbachia-free *Aedes albopictus* mosquito cells promotes virus replication and, importantly,
progeny virus infectivity. This has also been reported for DENV-2 infection in *w*Mel-colonized
Aedes aegypti mosquitoes (22). We observed that baseline MTase activity is required for virus
replication and spread in *Aedes* cells (Fig S2). Furthermore, the extent to which virus replication
is affected by MTase inhibitors depends on the virus source, with viruses produced from
Wolbachia-colonized cells (W+ viruses) being more susceptible to MTase inhibition.

This outcome phenocopies the scenario in which virus spread is most restricted under conditions
where both producer and target mosquito cells are colonized with *Wolbachia* (1). In line with these
findings, our data indicate a decrease in MTase expression occurs in the presence of *Wolbachia*
(Fig 1). This observation is in line with previous reports (22). Thus, our collective data support a
model in which endosymbiont-dependent inhibition of MTase expression and catalytic MTase
function contribute to reduced virus replication and per-particle infectivity in mosquitos (Fig 2-4).
This consequently limits virus dissemination within the vector and transmission to a vertebrate
host (Fig 2-4). Given that our results are coherent with prior reports involving a different RNA virus
and *Wolbachia* strain, the interaction between virus, *Wolbachia,* and host MTase expression is
likely independent of any particular virus-host-*Wolbachia* combination, representing a conserved
feature of pathogen blocking in the native *Aedes* vector.

Our data demonstrate an interaction between DNMT2 orthologs from *Aedes albopictus* and
Drosophila melanogaster and viral RNA (Fig 5B, Fig S4B) (1). However, it remains to be seen
whether these interactions are analogous to DNMT2-DCV RNA interactions in *Drosophila,* where
the MTase-viral RNA binding occurs specifically at structured viral Internal Ribosomal Entry Sites
(IRES) (29). Additionally, it is unclear if DNMT2 interactions in the mosquito cell are specific for
viral RNA or whether it extends to host transcripts. Future studies involving PAR-CLIP-sequencing
of immunoprecipitated DNMT2-RNA complexes should allow the identification and mapping of
distinct DNMT2-binding motifs and/or structural elements within viral and host RNA species.
Nevertheless, sites of DNMT2 recruitment to viral RNA, and viral and host proteins are required
for recruitment remain unidentified. Assuming that the proviral role of *Aedes* DNMT2 involves the
addition of m5C signatures to specific residues on the viral genome, it seems likely that viral co-
factor(s) are required for specificity. *Drosophila* DNMT2 is antiviral in fruit flies, however
Drosophila is not a natural host for alphaviruses, and it is likely that the adaptation of the virus to
the natural vector has facilitated an appropriate proviral interaction with *Aedes* DNMT2 or an
associated *Aedes*-specific host factors(s) absent in *Drosophila melanogaster* (32).

The m5C content of virion RNA produced from *Wolbachia*-colonized cells (W+ viruses) is
significantly reduced relative to cells without the symbiont (Fig 6B) (1), consistent with DNMT2s
role as a cytosine MTase. Incidentally, this finding follows reports dating back several decades
describing the occurrence of m5C residues within intracellular SINV RNA (33). Relative
abundance of methylated to total cytosine residues on SINV virion RNA derived from *Wolbachia*-
free mosquito cells is 15:10000, accounting for cytosine content in the viral genome we predict
there to be 4 to 5 m5C signatures per encapsidated virion RNA genome produced in *Aedes*
albopictus cells. Our observation supports the involvement of these intracellular m5C signatures
in alphavirus genome replication that W+ virion RNA, which are presumably hypomethylated, are
less infectious on a per-genome basis (Fig 4E-F) (34). Indeed, based on our data, we can infer
that these m5C modifications regulate alphavirus infection across multiple hosts and, thus, by
extension, aspects of the virus transmission cycle. It should also be noted that while methylated
nucleotide residues like m6A and m5C occur on RNA virus genomes at higher rates than those
present in cellular RNA species, our results do not exclude the possibility of other RNA
modifications, as well as differential modification of host RNA species and playing a role in
regulating virus replication and transmission. This may be of particular consequence given recent
evidence of altered m6A modification of specific cellular transcripts during flavivirus infection in
vertebrate cells (31, 35, 36).

Alphaviruses derived from mosquito cells are more infectious on vertebrate cells on a per-particle
basis than vertebrate cell-derived viruses and vice versa (37). This carries the implication that
progeny viruses originating from one cell type may possess intrinsic properties that can confer a
fitness advantage while infecting a destination host cell type, altering their infectivity on these
destination cells on a per-particle level. As to what such properties may represent, current
evidence points towards differences in virus structure, such as differential sialation or
glycosylation of viral glycoproteins impacting host receptor-binding and/or differences in the
encapsidated cargo, e.g., packaging of host ribosomal components (38-40). Furthermore, as our
results suggest, another property that might confer unique cell-type-specific advantages to viruses
is the differential modification of the virion RNA. Indeed, recent evidence shows that modifications
like N[6]-methyladenosine (m6A) and 5-methylcytosine (m5C) can regulate viral RNA functionality
in the cell (30, 41). Therefore, it is possible that such modifications also influence the infectivity of
progeny viruses produced from said cells. However, how these modifications affect virus
replication in a cell is still an open question. Indeed, information regarding the functional
consequence of m5C or other RNA modifications on viral RNA is limited. Thus, while we may
draw certain conclusions based on our current knowledge of known eukaryotic RNA
modifications, the potential implications of arbovirus RNA methylation may be broader than we

are currently able to anticipate (42). We can hypothesize that differential viral methylation may
alter host responses to infection, in that depending on the host or cell type, as well as the genomic
context of methylation, presence or absence of m5C may either allow detection by and/or provide
a mechanism of escape from RNA-binding proteins (e.g., Dicer, RIG-I, MDA5, TLRs, APOBEC3)
involved in virus restriction or non-self RNA recognition that trigger downstream immune signaling
and interferon production (43). Differential modifications of viral RNA may thus also regulate
different cytological outcomes of arboviruses infection of arthropod and vertebrate cells i.e.
persistence versus cell death.

It remains to be seen whether or not one or more of these situations occur during pathogen
blocking and if W- and W+ viruses trigger differential innate immune responses in vertebrate cells.
Based on our data, we propose a model in which our current estimates of m5C residues on W-
viruses represent the "wild-type" epitranscriptome of mosquito-derived alphavirus. In naïve
vertebrate cells, the presence of these signatures allows viruses to replicate efficiently following
successful evasion of host innate immunity. In contrast, m5C hypomethylation of W+ viruses
renders them more susceptible to host-induced restriction, thus impacting their ability to
propagate. Aside from heightened immune susceptibility, the decreased fitness of
hypomethylated W+ viruses could also result from reduced incoming viral RNA stability and/or
translation.

Given that pharmacological inhibition of MTase activity impacts virus spread in mosquito cells, it
is likely that W+ virus hypomethylation also influences dissemination in arthropod cells (1).
However, it is also possible that other factors contribute to the reduced fitness of W+ viruses. In
particular, our LC-MS/MS analyses suggest increased heterogeneity in m7G moiety abundance
on W+ virion RNA, indicating either difference in abundance of internal m7G methylation
signatures or a potential imbalance in viral RNA capping in the presence of *Wolbachia.* In addition,
past work has shown that SINV populations derived from different hosts vary with regard to the
ratios of capped and non-capped SINV RNA (44). Despite being important for alphavirus
replication, non-capped SINV RNA alone are compromised in their ability to undergo translation,
are more susceptible to RNA decay machineries, and have been shown to induce elevated innate
immune response, all of which might contribute to the observed loss in infectivity.

Finally, the data presented here implicating epitranscriptomic regulation of alphaviruses unlocks
multiple avenues of investigation, which include, but are not limited to the following. First, it is
crucial to determine the genomic context of m5C and other RNA modifications on viral RNA for
different hosts, cell types, and infection timeline, which may be achieved by long-read, direct RNA
sequencing from virus-infected cells. Doing so would allow sequence-specific mapping of these
signatures and help address whether virus infection is regulated solely via targeting viral RNA by
cellular MTases. Furthermore, deriving mapping information might inform us whether
modifications are directed to specific RNA elements that result in spatiotemporal changes in RNA
structure and altered base-pairing, thus regulating virus RNA polymerase fidelity and/or
translation in the cell. Additional areas of inquiry involve identifying cellular pathways responsible
for determining the fate of W+ viruses and characterizing the functional consequences of
abolishing highly conserved m5C residues on the viral RNA. This would allow further exploration
into the effect of these signatures on RNA stability, gene expression, and/or packaging across
arthropod and vertebrate cells. Lastly, unlike m6A-modifications, little is known regarding how
m5C signatures are interpreted, i.e., how they are "read," "maintained," and "erased," in
mammalian and, to an even lesser extent, in arthropod cells (42). Promising candidates include
m5C-binding "reader" proteins ALYREF and YBX1, which function alongside the known cellular
m5C MTase NSUN2 to influence mRNA nuclear transport and stability (45, 46). Following
approaches described in recent studies, identification of these RNA-binding proteins, either viral

or host-derived, may be achieved via affinity-based immunoprecipitation of viral RNA and form
the basis of future studies (47).

Like most other RNA viruses, Alphaviruses are limited in their coding capacity and are known to
alter their genome structure under various cellular conditions to regulate aspects of their
replication as a way to maximize viral genome functionality. Echoing this idea, the findings
presented in this study add to our understanding of regulatory mechanisms adopted by these
viruses to successfully navigate within and transition between vertebrate and arthropod host
species.

Materials and Methods

Insect and Mammalian Cell Culture

RML12 *Aedes albopictus* cells with and *Wolbachia*-free *w*Mel was grown at 24 ºC in Schneider's
insect media (Sigma-Aldrich) supplemented with 10% heat-inactivated fetal bovine serum
(Corning), 1% each of L-Glutamine (Corning), non-essential amino acids (Corning), and penicillin-
streptomycin-antimycotic (Corning). C710 *Aedes albopictus* cells with and *Wolbachia*-free were
grown at 27 ºC under 5% ambient CO_2 in 1X Minimal Essential Medium (Corning) supplemented
with 5% heat-inactivated fetal bovine serum (Corning), 1% each of L-Glutamine (Corning), non-
essential amino acids (Corning) and penicillin-streptomycin-antimycotic (Corning). Vertebrate
baby hamster kidney fibroblast or BHK-21 cells were grown at 37 ºC under 5% ambient CO_2 in
1X Minimal Essential Medium (Corning) supplemented with 10% heat-inactivated fetal bovine
serum (Corning), 1% each of L-Glutamine (Corning), non-essential amino acids (Corning) and
penicillin-streptomycin-antimycotic (Corning). JW18 *Drosophila melanogaster* cells with and
without *Wolbachia w*Mel were grown at 24 ºC in Shields, and Sang M3 insect media (Sigma-
Aldrich) supplemented with 10% heat-inactivated fetal bovine serum, 1% each of L-Glutamine
(Corning), non-essential amino acids (Corning), and penicillin-streptomycin-antimycotic
(Corning).

Mosquito rearing and blood meals

Aedes aegypti mosquitoes either -infected and -uninfected with *Wolbachia* (*w*AlbB strain)
(generously provided by Dr. Zhiyong Xi, Michigan State University, USA), were reared in an insect
incubator (Percival Model I-36VL, Perry, IA, USA) at 28 °C and 75% humidity with 12 h light/dark
cycle. Four to six-day-old, mated female mosquitoes were allowed to feed for one hour on
approximately 10^8 PFUs of SINV (TE12-untagged) containing citrated rabbit blood (Fisher
Scientific DRB030) supplemented with 1mM ATP (VWR), and 10% sucrose using a Hemotek
artificial blood-feeding system (Hemotek, UK) maintained under a constant temperature of 37 °C.
Engorged mosquitoes were then isolated and reared at 28 °C in the presence of male mosquitoes.
For harvesting whole tissues, mosquitoes were harvested 5-7 days post blood meal before being
snap-frozen in liquid nitrogen and stored at −80 °C before further processing. For salivary gland
dissections, mosquitoes were kept immobilized on ice before dissection. Collected salivary gland
tissues were washed three times in a cold, sterile saline solution (1XPBS) before being snap-
frozen in liquid nitrogen and stored at −80 °C before further processing. Three salivary glands
were pooled to create each biological replicate. Samples for qPCR and qRT-PCR were
homogenized in TRiZOL (Sigma Aldrich) reagent and further processed for nucleic acid
extractions using manufacturer's protocols.

Virion RNA extraction and transfection

Virion encapsidated RNA was extracted from viruses (SINV-nLuc) were purified over a 27%
sucrose cushion using TRiZOL reagent (Sigma Aldrich) using the manufacturer's protocol. Post
extraction, RNAs were DNase (RQ1 RNase-free DNase, NEB) treated according to the
manufacturer's protocol to remove cellular contaminants, and viral RNA copies were quantified

via quantitative RT-PCR using primers probing for SINV nsP1 and E1 genomic regions (Table
S1) and a standard curve comprised of linearized SINV infectious clone containing the full-length
viral genome. To determine infectivity or replication kinetics of Sindbis virion RNA, equal copies
of virion isolated RNA (10^5 copies), quantified using qRT-PCR, were transfected into BHK-21 cells
in serum-free Opti-MEM (Gibco). Transfection was carried out for 6 hours before the transfection
inoculum was removed, and overlay was applied. Cells were fixed post-transfection using 10%
(v/v) formaldehyde and stained with crystal violet to visualize plaques. To maximize the production
of infectious units, equal mass (1 µg) of virion (SINV-nLuc) isolated RNA derived from JW18 fly
cells was transfected into BHK-21 cells. Transfection was carried out for 6 hours before the
transfection inoculum was removed, and overlay was applied. Cells were fixed 48 (SINV) or 72
(CHIKV) hours post-transfection using 10% (v/v) formaldehyde and stained with crystal violet to
visualize plaque-forming units.

Viral replication assays

The viral genome and sub-genome translation were quantified using cellular lysates following
synchronized infections with reporter viruses (SINV-nLuc) or transfections with virion-derived
RNA from the aforementioned viruses. At indicated times post-infection, samples were collected
and homogenized in 1X Cell Culture Lysis Reagent (Promega). In addition, samples were mixed
with NanoGlo luciferase reagent (Promega), incubated at room temperature for three minutes
before luminescence was recorded using a Synergy H1 microplate reader (BioTech instruments).

Virus infection in cells and progeny virus production

Virus stocks were generated from RML12, C710, or JW18 cells, either with or without *Wolbachia*
or overexpressing DNMT2 orthologs by infecting naïve cells with the virus at an MOI of 10. In all
cases, serum-free media was used for downstream virus purification. Media containing virus was
collected five days post-infection for alphaviruses SINV (SINV-nLuc, TE12-untagged, TE3'2J-
GFP, and TE3'2J-mCherry) and CHIKV (CHIKV18125-capsid-mKate). Virus stocks were
subsequently purified and concentrated by ultracentrifugation (43K for 2.5 h) over a 27% (w/v)
sucrose cushion dissolved in HNE buffer. Viral pellets were stored and aliquoted in HNE buffer
before being used for all subsequent experiments.

DNMT2 overexpression in arthropod cells

Aedes albopictus AMt2 coding region was subcloned into PCR 2.1 TOPO vector (Invitrogen) by
PCR amplification of cDNA generated using reverse transcribed from total cellular RNA isolated
from C636 *Aedes albopictus* cells using Protoscript II RT (NEB) and oligo-dT primers (IDT). The
coding region was validated via sequencing before being cloned into the pAFW expression vector
(1111) (Gateway Vector Resources, DGRC), downstream of and in-frame with the 3X FLAG tag
using the native restriction sites *AgeI* and *NheI* (NEB). Expression of both FLAG-tagged
AaDNMT2 in mosquito cells was confirmed using qRT-PCR and Western Blots using an anti-
FLAG monoclonal antibody (SAB4301135 - Sigma-Aldrich, 1:3000 dilution in 2% Milk in 1X TBS
+ 1% Tween-20) (Fig 4A). In addition, catalytic MTase mutant of *AMt2* (*AMt2*-C78G*)* was
generated via site-directed mutagenesis (NEB, Q5 Site-Directed Mutagenesis Kit), using primers
listed in the primer table (Table S1). *Drosophila Mt2* (FBgn0028707) cDNA clone (GM14972)
obtained from DGRC (https://dgrc.bio.indiana.edu/) was cloned into the pAFW expression vector
(1111) with an engineered *SalI* site (Gateway Vector Resources, DGRC) downstream of and in-
frame with the 3X FLAG tag using Gibson assembly (HiFi DNA assembly mix, NEB). Expression
of FLAG-tagged DNMT2 in fly cells was confirmed using qRT-PCR and Western Blots using an
anti-FLAG monoclonal antibody (SAB4301135 - Sigma-Aldrich, 1:3000 dilution in 2% Milk in 1X
TBS + 1% Tween-20). Catalytically inactive *Mt2* (*Mt2* C78A) variant was generated via site-
directed mutagenesis (NEB, Q5 Site-Directed Mutagenesis Kit) using primers listed in the primer
table (Table S1).

Immunoprecipitation of DNMT2-viral RNA complexes

JW18 fly cells and C710 mosquito cells were transfected with expression vectors FLAG-Mt2 and FLAG-AMt2 respectively for 48 hours before infection with SINV at MOI of 10. Control cells were transfected with the empty vector plasmid FLAG-empty. In addition, cells were treated for approximately 18h with 5 µM 5-Azacytidine to covalently trap Mt2 or AMt2 with its target cellular RNA before RNA immunoprecipitation using an anti-FLAG antibody following manufacturer's protocols (SAB4301135 - Sigma-Aldrich, 1:100 dilution) (23).

Real-time quantitative RT-PCR analyses

Following total RNA extraction using TRiZOL reagent, cDNA was synthesized using MMuLV Reverse Transcriptase (NEB) with random hexamer primers (Integrated DNA Technologies). Negative (no RT) controls were performed for each target. Quantitative RT-PCR analyses were performed using Brilliant III SYBR Green QPCR master mix (Bioline) with gene-specific primers according to the manufacturer's protocol and the Applied Bioscience StepOnePlus qRT-PCR machine (Life Technologies). The expression levels were normalized to the endogenous 18S rRNA expression using the delta-delta comparative threshold method (ΔΔCT). Fold changes were determined using the comparative threshold cycle (CT) method (Table S1). Efficiencies for primer sets used in this study have been validated in our previous study (1).

DNMT2 inhibition in mosquito cells

Inhibition of *Aedes* DNMT2 activity in C710 cells was achieved using RNA and DNA cytosine methyltransferase inhibitors, 5-aza-cytidine (5-AZAC, Sigma-Aldrich) and 5-deoxy-azacytidine (DAC-5, Sigma-Aldrich). In each case, *Aedes albopictus* C710 cells were treated overnight with media containing either 5µM inhibitor diluted in Dimethyl sulfoxide (DMSO) or DMSO alone. Due to the poor stability of 5-AZAC, media containing fresh inhibitors was added every day post-infection (48).

Live cell imaging

Live-cell imaging experiments were carried out using a setup similar to our previous study (1). The growth of fluorescent reporter viruses in *Aedes albopictus* (C710) cells was monitored using Incucyte live-cell analysis system (Essen Biosciences, USA). *Aedes albopictus* C710 cells were grown under standard conditions as described earlier under 5% ambient CO_2 at 27 ºC. Cells were plated to 75-80% confluency in 96-well plates to separate adjacent cells and preserve cell shape for optimal automated cell counting. To synchronously infect cells virus was adsorbed at 4°C. Post adsorption, cell monolayers were extensively washed with cold 1XPBS to remove any unbound virus particles, followed by the addition of warm media (37ºC) to initialize virus internalization and infection. Cells per well were imaged and averaged across four distinct fields of view, each placed in one-quarter of the well every 2 hours throughout the infection. Total fluorescence generated by cells expressing the red fluorescent reporter mKate was calculated and normalized by the cell number for every sample. A manual threshold was set to minimize background signal via automated background correction at the time of data collection. Following the acquisition, data were analyzed in real-time using the native Incucyte ® Base Analysis Software.

Quantification of RNA modification by LC-MS/MS

Total RNA (3-7µg) was digested by nuclease P1 (10 Units) at 50°C for 16 hr. Additional Tris pH 7.5 was then added to a final concentration of 100 mM to adjust, followed by the addition of calf intestinal alkaline phosphatase (CIP, NEB, 2Units). The mixture was incubated at 37°C for 1 hour to convert nucleotide 5'-monophosphates to their respective nucleosides. Next, 10µl of RNA samples were diluted to 30 µL and filtered (0.22 µm pore size). 10µL of the sample was used for LC-MS/MS. Briefly, nucleosides were separated on a C18 column (Zorbax Eclipse Plus C18

column, 2.1 x 50mm, 1.8 Micron) paired with an Agilent 6490 QQQ triple-quadrupole LC mass spectrometer using multiple-reaction monitoring in positive-ion mode. The nucleosides were quantified using the retention time of the pure standards and the nucleoside to base ion mass transitions of 268.1 to 136 (A), 244.1 to 112 (C), 284.2 to 152 (G), 258 to 126 (m3C and m5C), 282.1 to 150 (m6A), 298 to 166 (m7G). Standard calibration curves were generated for each nucleoside by fitting the signal intensities against concentrations of pure-nucleoside preparations. The curves were used to determine the concentration of the respective nucleoside in the sample. The A, G, and C standards were purchased from ACROS ORGANICS; m5C was purchased from BioVision; m7G, m1G, and m3C were purchased from Carbosynth, m6G and m6A were purchased from Berry's Associates, and m1A was from Cayman Chemical Company. The modification level on the nucleosides was calculated as the ratio of modified: unmodified.

Data availability

The full Incucyte dataset is available in the form of an excel sheet labeled Dataset S1.

Statistical analyses of experimental data

All statistical analyses were conducted using GraphPad Prism 8 (GraphPad Software Inc., San Diego, CA).

Graphics

Graphical assets made in © BioRender - biorender.com.

Acknowledgments

We thank members of Hardy, Newton, Danthi, Mukhopadhyay, and Patton Labs for critical proofreading of the manuscript and for fostering ideas through helpful discussions. In addition, we thank Dr. David Mackenzie-Liu for generating the original cloning vector with the SacI restriction site. Finally, we thank our colleagues outside of IU for their generosity in sharing reagents, including Dr. Horacio Frydman, Boston University, for providing us Aa23 and C710 *Aedes albopictus* cells, Dr. William Sullivan for providing us with JW18 *Drosophila melanogaster* cells, and Dr. Zhiyong Xi, Michigan State University for providing us *Aedes aegypti* mosquitoes. This work was supported by NSF award to ILGN and RWH (MTM2025389), NIH R01 to ILGN (R01AI144430), NIH R21 to RWH (R21AI153785).

References

1.   Bhattacharya T, Newton ILG, Hardy RW. Viral RNA is a target for Wolbachia-mediated pathogen blocking. PLoS Pathog. 2020;16(6):e1008513.

2.   Díaz-Sánchez S, Hernández-Jarguín A, Torina A, Fernández de Mera I, Estrada-Peña A, Villar M, et al. Biotic and abiotic factors shape the microbiota of wild-caught populations of the arbovirus vector Culicoides imicola. Insect molecular biology. 2018;27(6):847-61.

3.   Bellone R, Failloux A-B. Temperature in shaping mosquito-borne viruses transmission. Frontiers in Microbiology. 2020;11:2388.

4.   Tabachnick WJ. Ecological effects on arbovirus-mosquito cycles of transmission. Current opinion in virology. 2016;21:124-31.

5.   Hegde S, Rasgon JL, Hughes GL. The microbiome modulates arbovirus transmission in mosquitoes. Current opinion in virology. 2015;15:97-102.

6.   Cirimotich CM, Ramirez JL, Dimopoulos G. Native microbiota shape insect vector competence for human pathogens. Cell host & microbe. 2011;10(4):307-10.

7.   Lindsey ARI, Bhattacharya T, Hardy RW, Newton ILG. Wolbachia and virus alter the host
transcriptome at the interface of nucleotide metabolism pathways. bioRxiv.
2020:2020.06.18.160317.
8.   Lindsey ARI, Bhattacharya T, Newton ILG, Hardy RW. Conflict in the Intracellular Lives of
Endosymbionts and Viruses: A Mechanistic Look at Wolbachia-Mediated Pathogen-blocking.
Viruses. 2018;10(4).
9.   Bhattacharya T, Newton IL, Hardy RW. Wolbachia elevates host methyltransferase
expression to block an RNA virus early during infection. PLoS pathogens. 2017;13(6):e1006427.
10.   Bhattacharya T, Newton ILG. Mi Casa es Su Casa: how an intracellular symbiont
manipulates host biology. Environ Microbiol. 2017.
11.   Schultz M, Tan A, Gray C, Isern S, Michael S, Frydman HM, et al. Wolbachia wStri blocks
Zika virus growth at two independent stages of viral replication. MBio. 2018;9(3).
12.   Kaur R, Shropshire JD, Cross KL, Leigh B, Mansueto AJ, Stewart V, et al. Living in the
endosymbiotic world of Wolbachia: A centennial review. Cell Host & Microbe. 2021.
13.   Caragata EP, Dutra HLC, Moreira LA. Inhibition of Zika virus by Wolbachia in Aedes aegypti.
Microbial cell. 2016;3(7):293.
14.   Rainey SM, Martinez J, McFarlane M, Juneja P, Sarkies P, Lulla A, et al. Wolbachia Blocks
Viral Genome Replication Early in Infection without a Transcriptional Response by the
Endosymbiont or Host Small RNA Pathways. PLoS Pathog. 2016;12(4):e1005536.
15.   Geoghegan V, Stainton K, Rainey SM, Ant TH, Dowle AA, Larson T, et al. Perturbed
cholesterol and vesicular trafficking associated with dengue blocking in Wolbachia-infected
Aedes aegypti cells. Nature communications. 2017;8(1):1-10.
16.   Zabalou S, Riegler M, Theodorakopoulou M, Stauffer C, Savakis C, Bourtzis K. Wolbachia-
induced cytoplasmic incompatibility as a means for insect pest population control. Proceedings
of the National Academy of Sciences. 2004;101(42):15042-5.
17.   Beckmann JF, Ronau JA, Hochstrasser M. A Wolbachia deubiquitylating enzyme induces
cytoplasmic incompatibility. Nature microbiology. 2017;2(5):1-7.
18.   Caragata EP, Dutra HL, Sucupira PH, Ferreira AG, Moreira LA. Wolbachia as translational
science: controlling mosquito-borne pathogens. Trends in Parasitology. 2021.
19.   Indriani C, Tantowijoyo W, Rancès E, Andari B, Prabowo E, Yusdi D, et al. Reduced dengue
incidence following deployments of Wolbachia-infected Aedes aegypti in Yogyakarta, Indonesia:
a quasi-experimental trial using controlled interrupted time series analysis. Gates open research.
2020;4.
20.   Nazni WA, Hoffmann AA, NoorAfizah A, Cheong YL, Mancini MV, Golding N, et al.
Establishment of Wolbachia strain wAlbB in Malaysian populations of Aedes aegypti for dengue
control. Current biology. 2019;29(24):4241-8. e5.
21.   Thomas S, Verma J, Woolfit M, O'Neill SL. Wolbachia-mediated virus blocking in mosquito
cells is dependent on XRN1-mediated viral RNA degradation and influenced by viral replication
rate. PLoS pathogens. 2018;14(3):e1006879.
22.   Zhang G, Hussain M, O'Neill SL, Asgari S. Wolbachia uses a host microRNA to regulate
transcripts of a methyltransferase, contributing to dengue virus inhibition in Aedes aegypti. Proc
Natl Acad Sci U S A. 2013;110(25):10276-81.

23.   Khoddami V, Cairns BR. Transcriptome-wide target profiling of RNA cytosine methyltransferases using the mechanism-based enrichment procedure Aza-IP. Nat Protoc. 2014;9(2):337-61.

24.   Goll MG, Kirpekar F, Maggert KA, Yoder JA, Hsieh CL, Zhang X, et al. Methylation of tRNAAsp by the DNA methyltransferase homolog Dnmt2. Science. 2006;311(5759):395-8.

25.   Singh V, Sharma P, Capalash N. DNA methyltransferase-1 inhibitors as epigenetic therapy for cancer. Current cancer drug targets. 2013;13(4):379-99.

26.   Pleyer L, Greil R. Digging deep into "dirty" drugs–modulation of the methylation machinery. Drug metabolism reviews. 2015;47(2):252-79.

27.   Jeltsch A, Ehrenhofer-Murray A, Jurkowski TP, Lyko F, Reuter G, Ankri S, et al. Mechanism and biological role of Dnmt2 in nucleic acid methylation. RNA biology. 2017;14(9):1108-23.

28.   Jurkowski TP, Meusburger M, Phalke S, Helm M, Nellen W, Reuter G, et al. Human DNMT2 methylates tRNAAsp molecules using a DNA methyltransferase-like catalytic mechanism. Rna. 2008;14(8):1663-70.

29.   Durdevic Z, Hanna K, Gold B, Pollex T, Cherry S, Lyko F, et al. Efficient RNA virus control in Drosophila requires the RNA methyltransferase Dnmt2. EMBO reports. 2013;14(3):269-75.

30.   Gokhale NS, McIntyre ABR, McFadden MJ, Roder AE, Kennedy EM, Gandara JA, et al. N6-Methyladenosine in Flaviviridae Viral RNA Genomes Regulates Infection. Cell Host Microbe. 2016;20(5):654-65.

31.   McIntyre W, Netzband R, Bonenfant G, Biegel JM, Miller C, Fuchs G, et al. Positive-sense RNA viruses reveal the complexity and dynamics of the cellular and viral epitranscriptomes during infection. Nucleic Acids Res. 2018;46(11):5776-91.

32.   Bhattacharya T, Rice DW, Hardy RW, Newton I. Adaptive evolution in DNMT2 supports its role in the dipteran immune response. bioRxiv. 2020.

33.   Dubin DT, Stollar V. Methylation of Sindbis virus "26S" messenger RNA. Biochem Biophys Res Commun. 1975;66(4):1373-9.

34.   Frey TK, Gard DL, Strauss JH. Replication of Sindbis virus. VII. Location of 5-methyl cytidine residues in virus-specific RNA. Virology. 1978;89(2):450-60.

35.   Gokhale NS, McIntyre ABR, Mattocks MD, Holley CL, Lazear HM, Mason CE, et al. Altered m(6)A Modification of Specific Cellular Transcripts Affects Flaviviridae Infection. Mol Cell. 2020;77(3):542-55.e8.

36.   Tsai K, Cullen BR. Epigenetic and epitranscriptomic regulation of viral replication. Nat Rev Microbiol. 2020:1-12.

37.   Mackenzie-Liu D, Sokoloski KJ, Purdy S, Hardy RW. Encapsidated Host Factors in Alphavirus Particles Influence Midgut Infection of Aedes aegypti. Viruses. 2018;10(5).

38.   Sokoloski KJ, Snyder AJ, Liu NH, Hayes CA, Mukhopadhyay S, Hardy RW. Encapsidation of host-derived factors correlates with enhanced infectivity of Sindbis virus. J Virol. 2013;87(22):12216-26.

39.   Dunbar CA, Rayaprolu V, Wang JC, Brown CJ, Leishman E, Jones-Burrage S, et al. Dissecting the Components of Sindbis Virus from Arthropod and Vertebrate Hosts: Implications for Infectivity Differences. ACS Infect Dis. 2019;5(6):892-902.

40.   Hsieh P, Rosner MR, Robbins PW. Host-dependent variation of asparagine-linked oligosaccharides at individual glycosylation sites of Sindbis virus glycoproteins. J Biol Chem. 1983;258(4):2548-54.

41.   Courtney DG, Tsai K, Bogerd HP, Kennedy EM, Law BA, Emery A, et al. Epitranscriptomic
Addition of m(5)C to HIV-1 Transcripts Regulates Viral Gene Expression. Cell Host Microbe.
2019;26(2):217-27.e6.
42.   Netzband R, Pager CT. Epitranscriptomic marks: Emerging modulators of RNA virus gene
expression. Wiley Interdiscip Rev RNA. 2020;11(3):e1576.
43.   Durbin AF, Wang C, Marcotrigiano J, Gehrke L. RNAs Containing Modified Nucleotides Fail
To Trigger RIG-I Conformational Changes for Innate Immune Signaling. mBio. 2016;7(5).
44.   Sokoloski K, Haist K, Morrison T, Mukhopadhyay S, Hardy R. Noncapped alphavirus
genomic RNAs and their role during infection. Journal of virology. 2015;89(11):6080-92.
45.   Chen X, Li A, Sun BF, Yang Y, Han YN, Yuan X, et al. 5-methylcytosine promotes
pathogenesis of bladder cancer through stabilizing mRNAs. Nat Cell Biol. 2019;21(8):978-90.
46.   Yang X, Yang Y, Sun BF, Chen YS, Xu JW, Lai WY, et al. 5-methylcytosine promotes mRNA
export - NSUN2 as the methyltransferase and ALYREF as an m(5)C reader. Cell Res.
2017;27(5):606-25.
47.   Garcia-Moreno M, Järvelin AI, Castello A. Unconventional RNA-binding proteins step into
the virus–host battlefront. Wiley Interdisciplinary Reviews: RNA. 2018;9(6):e1498.
48.   Israili ZH, Vogler WR, Mingioli ES, Pirkle JL, Smithwick RW, Goldstein JH. The disposition
and pharmacokinetics in humans of 5-azacytidine administered intravenously as a bolus or by
continuous infusion. Cancer Res. 1976;36(4):1453-61.

754