Title - Rescue of aberrant huntingtin palmitoylation ameliorates mutant

huntingtin-induced toxicity

Authors – Fanny L. Lemarié[a], Nicholas S. Caron[a], Shaun S. Sanders[a,b], Mandi E.

Schmidt[a], Yen T.N. Nguyen[a], Yun Ko[a], Xiaohong Xu[d,e], Mahmoud A. Pouladi[a,d], Dale D.O.

Martin[a,c], Michael R. Hayden[a]*

[a] Centre for Molecular Medicine and Therapeutics, BC Children's Hospital Research Institute, University of
British Columbia, Vancouver, Canada

[b] Department of Molecular and Cellular Biology, University of Guelph, Guelph, Canada

[c] Department of Biology· University of Waterloo, Waterloo, Canada

[d] Translational Laboratory in Genetic Medicine, Department of Physiology, Yong Loo Lin School of
Medicine, National University of Singapore, Singapore

[e] Department of Neurology and Stroke Center, The First Affiliated Hospital, Jinan University, Guangzhou,
China

*To whom correspondence should be addressed (mrh@cmmt.ubc.ca)
Centre for Molecular Medicine and Therapeutics
BC Children's Hospital Research Institute
University of British Columbia
3025-950 West 28th Ave
Vancouver, BC Canada
V5Z 4H4
604-875-3535
mrh@cmmt.ubc.ca

Highlights:

• Palmitoylation of mHTT is reduced in multiple transgenic HD mouse models

• HTT palmitoylation decreases with increasing polyQ length in HD patient cells

• mHTT-lowering in mouse brains does not rescue aberrant palmitoylation

• mHTT palmitoylation in HD patient-derived cells can be rescued via APT inhibition

• Promoting palmitoylation reduces mHTT aggregation and cytotoxicity *in vitro*

Abstract: Huntington disease (HD) is a neurodegenerative disorder caused by a CAG

expansion in the *HTT* gene that codes for an elongated polyglutamine tract in the

huntingtin (HTT) protein. HTT is subject to multiple posttranslational modifications (PTMs)

that regulate its cellular function. Mutating specific PTM sites within mutant HTT (mHTT)

in HD mouse models can modulate disease phenotypes, highlighting the key role of HTT

PTMs in the pathogenesis of HD. These findings have led to increased interest in

developing small molecules to modulate HTT PTMs in order to decrease mHTT toxicity.

However, the therapeutic efficacy of pharmacological modulation of HTT PTMs in

preclinical HD models remains unknown. HTT is palmitoylated at cysteine 214 by the

huntingtin-interacting protein 14 (HIP14 or ZDHHC17) and 14-like (HIP14L or ZDHHC13)

acyltransferases. Here, we assessed if HTT palmitoylation should be regarded as a

therapeutic target to treat HD by (1) investigating palmitoylation dysregulation in

humanized rodent and human HD model systems, (2) measuring the impact of mHTT-

lowering therapy on brain palmitoylation, and (3) evaluating if HTT palmitoylation can be

modulated by pharmacological intervention. We show that palmitoylation of mHTT and

some HIP14/HIP14L-substrates is decreased early in multiple HD mouse models, and

that aging further reduces HTT palmitoylation. Lowering mHTT in the brain of YAC128

mice is not sufficient to rescue aberrant palmitoylation. However, we demonstrate that

mHTT palmitoylation can be normalized in COS-7 cells, in YAC128 cortico-striatal primary

neurons and HD patient-derived lymphoblasts using an acyl-protein thioesterase (APT)

inhibitor. Moreover, we show that modulating palmitoylation reduces mHTT aggregation

and mHTT-induced cytotoxicity in COS-7 cells and YAC128 neurons.

Keywords: Palmitoylation, S-acylation, huntingtin, Huntington disease, ZDHHC17, acyl-

protein thioesterase, neuroprotection

Abbreviations: ABE, acyl-biotin exchange assay; APT, acyl-protein thioesterase; ASO,

antisense oligonucleotide; HD, Huntington disease; hiPSC, human induced pluripotent

stem cell; HIP14, huntingtin-interacting protein 14 (ZDHHC17); HIP14L, huntingtin-

interacting protein 14-like (ZDHHC13); HTT, huntingtin (human); Htt, huntingtin (mouse);

ICV, intracerebroventricular; LNA, locked nucleic acid; mHTT, mutant huntingtin; NEM,

N-ethylmaleimide; PalmB, palmostatin B; PAT, palmitoyl acyltransferase; PTM, post-

translational modification

1. Introduction

Huntington disease (HD) is an inherited neurodegenerative disorder characterized by involuntary movements, behavioural changes and cognitive impairment. There are currently no therapies available to influence its inexorable progression (Caron et al., 2018; Evans et al., 2013; Fisher and Hayden, 2014; Myers, 2004). The genetic cause of HD is an expanded CAG-repeat tract (>35 repeats) in the *HTT* gene leading to an abnormally long polyglutamine (polyQ) tract in the huntingtin (HTT) protein.

HTT interacts with numerous protein partners and is involved in many cellular processes, including endocytosis, vesicle/organelle transport and recycling, autophagy, and DNA transcription (Maiuri et al., 2017; Ochaba et al., 2014; Saudou and Humbert, 2016). HTT is subjected to numerous post-translational modifications (PTMs), including phosphorylation (Aiken et al., 2009; Humbert et al., 2002; Schilling et al., 2006), acetylation (Jeong et al., 2009), ubiquitination (Kalchman et al., 1996), SUMOylation (Steffan et al., 2004), proteolysis (Graham et al., 2006), and fatty acylation (palmitoylation and myristoylation) (D. D. O. Martin et al., 2018; Martin et al., 2014; Yanai et al., 2006). These PTMs regulate subcellular localization, protein-protein interactions, aggregation and clearance which modulates the toxicity of mutant HTT (mHTT) (Atwal et al., 2007; Cariulo et al., 2017; Caron et al., 2013; Kratter et al., 2016; Lontay et al., 2020).

In the presence of the HD mutation, levels of several key PTMs are significantly altered *in vitro* and *in vivo* (Cariulo et al., 2020, 2017; Warby et al., 2005). Introducing mutations at specific PTM sites of mHTT has been shown to modulate disease phenotypes in various HD models. Blocking proteolysis of mHTT at aspartic acid 586 (D586) prevents striatal neurodegeneration in the YAC128 HD mouse model (Graham et al., 2006).

Reducing phosphorylation at serine residues 13/16 or 421 (S13/16A, S421A) in the

BACHD mouse model and HD human induced pluripotent stem cell (hiPSC)-derived

neural cells exacerbates disease phenotypes. In contrast, mimicking phosphorylation at

these residues (S13/16D, S421D) prevents neurodegeneration (Gu et al., 2009; Kratter

et al., 2016; Xu et al., 2020). These findings highlight the key role of HTT PTMs in the

pathogenesis of HD and have stimulated interest in the development of drugs to modulate

HTT PTMs in order to decrease mHTT levels and toxicity. To date, most research has

focused on normalizing HTT phosphorylation at S13 and S16, and preventing mHTT

proteolysis (Aharony et al., 2015; Atwal et al., 2011; Bowie et al., 2018).

Palmitoylation refers to the addition of a palmitic acid (C16:0) onto a cysteine

residue via a thioester bond. This process is catalyzed by a family of palmitoyl

acyltransferases (PATs). Palmitoylation increases the hydrophobicity of proteins, and

therefore plays a key role in protein trafficking, stability, membrane association and

protein-protein interactions (Fukata and Fukata, 2010). It is the most common protein-

lipid modification in the brain (Smotrys and Linder, 2004). Most importantly, it is reversible

and allows for dynamic regulation of protein functions at the synapse, thus playing an

important role in neurodevelopmental processes and in neuronal survival (Globa and

Bamji, 2017; Hubalkova et al., 2021; Matt et al., 2019; Mukai et al., 2008). Disruption of

protein palmitoylation has been implicated in the pathogenesis of several neurological

disorders, including Alzheimer's disease, schizophrenia and intellectual disability (Cho

and Park, 2016; Sanders et al., 2015b).

We first described that HTT is palmitoylated at cysteine 214 (C214), a reaction catalyzed

by the palmitoyl acyltransferases ZDHHC17 and ZDHHC13, also known as huntingtin-

interacting protein 14 and 14-like (HIP14 and HIP14L) (Yanai et al., 2006). In the presence

of the HD mutation, mHTT was found to be less palmitoylated in the brain of the YAC128

HD mouse model (Yanai et al., 2006). Transient expression of mHTT carrying a

palmitoylation-resistant mutation (C214 to serine, C214S) in immortalized cell lines and

primary neurons led to increased aggregates and nuclear inclusion formation, increased

cell death and increased susceptibility to excitotoxicity (Yanai et al., 2006).

Autopalmitoylation of HIP14 and HIP14L, and palmitoylation of many of their synaptic

substrates were also downregulated in the YAC128 HD mouse brains or in the brains of

mice lacking one *Htt* allele (*Htt*$^{+/-}$) (Huang et al., 2011; Singaraja et al., 2011). Moreover,

mouse models deficient in either HIP14 (*Hip14*$^{-/-}$) or HIP14L (*Hip14l*$^{-/-}$) develop

neuropathological deficits that are reminiscent of HD (Sutton et al., 2013), and loss of

both *Hip14* and *Hip14l* leads to embryonic lethality (Sanders et al., 2016, 2015a).

Collectively, these findings link defects in fatty acylation with neuronal dysfunction and

neurodegeneration in HD. However, the effect of palmitoylation on HTT function and the

role that loss of HTT palmitoylation plays in the pathogenesis of HD are still not fully

elucidated.

In this study, we investigate whether palmitoylation is dysregulated in HD mouse

models and human HD samples. We also evaluate whether pharmacological modulation

of mHTT palmitoylation represents a potential therapeutic target for the treatment of HD.

2. Materials and methods

2.1. Materials

2.1.1. Reagents and chemicals

Table 1 lists the reagents and chemicals used for this study along with the application, manufacturer and catalogue number.

2.1.2. Antibodies

Table 2 displays the list of antibodies used along with the application, concentrations and experimental conditions.

2.1.3. Experimental models

HD mouse lines - Heterozygous YAC128 (line 53), homozygous YAC128 (line 55), homozygous YAC18 (line 60), BACHD, humanized Hu18/18 and Hu97/18, humanized Hu21/21 and Hu128/21 mice were maintained on the FVB/N background. The description of the HD mouse lines used is displayed in **Table 3**. The YAC128 model expresses two mouse Htt alleles and the full length human mHTT transgene (Slow et al., 2003). The BACHD model expresses two mouse Htt alleles and the full length human mHTT transgene (Gray et al., 2008). The two humanized Hu97/18 and Hu128/21 models express only full length human WT and mHTT transgenes (Southwell et al., 2017, 2013). Experiments were performed according to protocols approved by the University of British Columbia Animal Care Committee (experimental protocol 2016/2020 #A20-0107, breeding protocol #A20-0233). Homozygous YAC128 and YAC18 mice were used for the neuronal culture, to improve culture health by avoiding an overnight genotyping step. For

all cohorts, approximately equal numbers of male and female mice were used. All mice

were sacrificed using CO_2 asphyxiation in accordance with UBC SOP #I012.

Cell line culture – Control and HD patient-derived lymphoblastoid cell lines (**Table 4**)

were obtained from the NIGMS Human Genetic Cell Repository at the Coriell Institute for

Medical Research. All cell lines were maintained in RPMI 1640 medium containing 10%

FBS, 2 mM GlutaMAX, 10 units/mL of penicillin and 10 µg/mL of streptomycin. Cells were

grown in T-25 flasks incubated in an upright position with vented caps at 37°C and 5%

CO_2. Cell density was kept between 2.10^5 and 5.10^5 cells/mL of media.

COS-7 cells were grown in DMEM medium supplemented with 10% FBS, 1 mM sodium

pyruvate, 2 mM GlutaMAX, 10 units/mL of penicillin and 10 µg/mL of streptomycin. Cells

were grown in 10-cm plates at 37°C and 5% CO_2, and were split using a 0.25% trypsin

solution (1:10 dilution) when cells were ~90% confluent by microscopy.

Human pluripotent stem cells (hPSCs) lines listed in **Table 5** were cultured on Matrigel-

coated plates in mTeSR-1 complete medium supplemented with 10 units/mL of penicillin

and 10 µg/mL of streptomycin at 37°C and 5% CO_2, and were split with ReLeSR.

Neuronal culture - Timed pregnancies were set up by mating homozygous YAC18 (line

60) or YAC128 (line 55) mice. At E16.5, embryos were removed from pregnant females.

Cortical and striatal tissues attached were dissected in ice-cold HBSS, gently dissociated

with a P1000 pipette (×2-3 times) and briefly spun. Tissues were digested in 0.05%

trypsin-EDTA for 8 min at 37°C until addition of 10% FBS in neurobasal (NB) medium, as

previously described (Schmidt et al., 2020). Cells were centrifuged, further dissociated by

mechanical trituration in NB medium (supplemented with B27, GlutaMAX, penicillin and

streptomycin) and DNase (0.08 mg/mL). Neurons were plated in 6-well Nunclon Delta

surface plates at a density of 1.10^6 cells/well in 2 mL of complete NB medium. Prior to

plating, 6-well plates were coated with sterile-filtered 50 µg/mL of poly-D-lysine

hydrobromide for 2 h at room temperature and washed four times with sterile water. All

cultures were regularly supplemented with fresh complete NB medium (20% well volume).

2.2. Methods

2.2.1. Plasmid transfection

COS-7 cells were seeded in 6-well or 96-well plates overnight at 37°C and 5% CO_2. The

next day, cells were transfected with cDNA plasmid constructs encoding for 15Q and

mutant 128Q N-terminus HTT_{1-548} (N548), HTT_{1-1212} (N1212) and full-length HTT_{1-3144}

(HTT), or 17Q and 138Q HTT_{1-588}-YFP (N588) using X-tremeGENE 9 DNA transfection

reagent (ratio 3:1, reagent (µL): DNA (µg)). Cells were treated or harvested 18 to 24h

post-transfection.

2.2.2. Acyl-protein thioesterase (APT) inhibition with palmostatin B

Inhibition of APTs in immortalized cell lines

At 18-24h post-transfection, COS-7 cells were deprived of lipid for 1 h in DMEM (without

methionine or cysteine) supplemented with 5% Charcoal stripped FBS, subsequently

treated with DMSO or 15 µM Palmostatin B (PalmB) (diluted in DMSO at 1000X on

InSolution) while concurrently metabolically labelled with 100 µM alkyne-palmitate for 6h.

Bio-orthogonal click chemistry of alkyne-palmitate was performed as subsequently

described in section §2.2.4.

Inhibition of APTs in primary neurons

Primary neurons from cortex and striatum of E16.5 mouse embryos (homozygous YAC18

and YAC128 mouse lines) were cultured as previously described in section §2.1.3. At

DIV10, neuronal cultures were treated with DMSO or with 15 µM PalmB (InSolution) for

6h. The ABE assay was performed on neuronal lysates as described in section §2.2.4.

Inhibition of APTs in control and HD patient-derived lymphoblastoid lines

Lymphoblasts were cultured as previously described in section §2.1.3. Cells were seeded

at a density of 600,000 cells/2 mL/well in 6 well plates. The next day, cells were treated

with DMSO or 10 µM PalmB for 4h. The ABE assay was performed on lymphoblasts as

described in section §2.2.4.

2.2.3. Intracerebroventricular HTT antisense oligonucleotide infusion

A fully phosphorothioate modified gapmer ASO with 5' locked nucleic acid (LNA)

modifications in each wing, matching a sequence in intron 22 of the *HTT* gene (5'-

TAATACGTAAGTGTCACAA-3', custom synthesized by Qiagen) was delivered by

intracerebroventricular (ICV) injection as previously described (Southwell et al., 2015).

Five-month-old YAC128 (line 53) mice were anesthetized using 3% isoflurane and

secured on a stereotaxic frame. Mice were given pre-operative subcutaneous

buprenorphine (0.1 mg/kg) and 0.25% bupivacaine at the site of incision. A midline

incision was made in the scalp to expose the skull. Then, a 50 µL Hamilton syringe with

a 26-gauge needle was oriented to the bregma and moved 0.3 mm anterior, 1 mm lateral,

and 3 mm ventral. An injection volume of 10 µL was delivered into the lateral ventricle at

a rate of 5 µL/min. Mice were either injected with 50 µg ASO in sterile PBS (n=7) or PBS

only (n=6). The needle was left in place for 2 min after injection, before being slowly

retracted. The scalp was closed with sutures and Vetbond tissue adhesive. Animals were

then placed in a recovery cage with a heating pad, hydrogel, and food on the cage floor.

Mice were given post-operative buprenorphine as needed and monitored for 72h. Brains

were collected a month later, when mHTT levels have been shown to be lowered by 70

to 90% (Southwell et al., 2015).

2.2.4. Palmitoylation assays

Acyl-biotin exchange (ABE) assay

Mouse brain - ABE assays were performed on whole or half frozen brain samples from

YAC128 (line 53) and WT littermate controls, BACHD and WT littermate controls,

Hu97/18 and Hu18/18, Hu128/21 and Hu21/21, as previously described (Drisdel and

Green, 2004; Huang et al., 2009). Briefly, brains were harvested and immediately snap-

frozen in liquid N_2 and stored at -80°C until being processed. Frozen brains were

homogenized on ice in lysis buffer (150 mM NaCl, 50 mM Tris, 5 mM EDTA, 0.1% SDS,

1% triton X-100, pH 7.4) with 100 mM NEM (4 mL lysis buffer/brain hemisphere).

Homogenates were sonicated for 15 sec at 30% power to shear DNA and the insoluble

material was removed by centrifugation at 20,000 × g for 15 min at +4°C. Protein

concentrations in lysates were assessed by DC protein assay. HTT, PSD95, and GAD65

were immunoprecipitated (IP) from brain lysates (300 μL) by overnight incubation with

Protein G Dynabeads or PureProteome protein A/G, and appropriate antibodies for

immunoprecipitation (**Table 2**). Beads were then washed and split into two and treated

either with neutral pH hydroxylamine (HAM+) in lysis buffer or just lysis buffer (HAM-) for

2h for HTT and 1h for the other three proteins at room temperature. Following HAM

treatment, beads were washed and treated with 2.5 μM EZ-Link BMCC-Biotin in pH 6.2

lysis buffer for 1h at 4°C. At the end of the BMCC-Biotin treatment, beads were washed

and heated at 70°C for 10 min with NuPAGE LDS sample buffer and 100 mM DTT to

elute proteins.

Control and HD-patient derived lymphoblastoid and hiPSC lines – Lymphoblastoid aggregates in suspension in the media were dissociated by trituration and centrifuged for 5 min at 500 × g at room temperature. HiPSCs were washed with DMEM/F12, lifted from plated using ReLeSR and spun at 500 × g for 5 min at +4°C. Supernatant was discarded and cell pellets were frozen until being processed, as previously described for mouse brain samples. HTT was immunoprecipitated from cell lysates by overnight incubation with Protein G Dynabeads and HTT (MAB2166) antibodies (approx. 250 µg protein and 2 µL MAB2166 for 20 µL beads).

Bio-orthogonal labelling assay using alkyne-palmitate

Immortalized cell lines - cells transiently transfected with cDNA HTT constructs were deprived of lipid during 1 h incubation in DMEM supplemented with 5% charcoal stripped FBS 18h post-transfection and subsequently metabolically labelled with 100 µM alkyne-palmitate for 6 h. Cells were lifted from plates in PBS using cell scrapers, pelleted by centrifugation (500 × g, 5 min, +4°C) and subsequently lysed in RIPA (150 mM $NaCl_2$, 5 mM M EDTA, 50 mM Tris, 1% NP-40, 0.5% sodium deoxycholate, 0.1% SDS in dH_2O, complete protease inhibitors cocktails) or SDP lysis (50 mM Tris pH 8.0, 150 mM NaCl, 1% Igepal, 40 mM NaF, 1 mM PMSF, 2 mM sodium vanadate, complete protease inhibitors cocktails) buffer for 10 min on ice, 5 min at 4°C on rotator and were centrifuged (16,000 × g, 10 min, 4°C). Bio-orthogonal click chemistry of alkyne-palmitate was performed, as previously described (Yap et al., 2010) on cell lysates immunoprecipitated with anti-HTT antibodies (BKP1 or MAB2166). HTT immunoprecipitates were adjusted to 1% SDS and incubated with 100 mM TBTA, 1 mM $CuSO_4$, 1 mM TCEP and 100 mM azido-biotin at +37°C in darkness for 1 h.

2.2.5. Western blot analysis

Brain, COS-7 and HD patient-derived hiPSC and lymphoblastoid supernatants from HTT assays were run on NuPAGE™ 3-8% Tris-Acetate gradient protein gels in Tris-Acetate SDS Running Buffer (50 mM Tricine, 50 mM Tris Base, 0.1% SDS, pH 8.24) or homemade 8% low bis acrylamide (1:200 bis-acrylamide:acrylamide) gels in SDS-Tris-glycine running buffer (25 mM Tris base, 190 mM glycine, 3.5 mM SDS) (Carroll et al., 2011) to resolve full length HTT alleles with small ΔQs. Brain lysates from the other proteins (SNAP25, PSD95 and GAD65) were run on homemade tris-glycine gels (7-12%) in SDS-Tris-glycine running buffer. Protein were transferred to 0.45 μm nitrocellulose. Membranes were blocked with 5% milk in Phosphate-Buffered Saline (PBS) or 3-5% BSA in Tris-Buffered Saline (TBS) supplemented with 0.1-0.5% Tween-20 (T). Primary antibody dilutions of HTT (D7F7 or MAB2166), SNAP25 polyclonal, PSD95 monoclonal, and GAD65 polyclonal in 3-5% BSA PBST or TBST were applied to the immunoblots at room temperature for 1h or overnight at +4°C (**Table 2**). Membranes were washed 4 × 5 min in PBST or TBST. The appropriate secondary antibodies and Alexa Fluor 680 conjugated streptavidin were applied in 3-5% BSA PBST or TBST for 1 to 2h at room temperature. Membranes were washed 4 × 5 min in PBST or TBST and imaged using the LI-COR Odyssey Infrared Imaging System (Li-Cor Biosciences). Densitometry was quantified using the LiCor Image Studio Lite software and median signal intensity following background subtraction was used for analysis. Palmitoylation was analyzed as a ratio of HAM+ palmitoylation signal to total immunoprecipitated protein signal and normalized to the control.

2.2.6. Aggregation and cytotoxicity assays

Filter retardation assay

Protein aggregation of insoluble HTT_{1-588}-YFP transiently expressed in COS-7 cells was

measured using the filter-trap retardation assay, as previously described (Dale D. O.

Martin et al., 2018; Wanker et al., 1999). After 72 h of transfection, COS-7 cells expressing

the indicated HTT_{1-588}-YFP constructs and treated with 15 µM PalmB for 48h were lysed

in lysis buffer (50 mM Tris pH 8.8, 100 mM NaCl, 5 mM $MgCl_2$, 0.5% Igepal CA630, 1 mM

EDTA with 1X cOmplete protease inhibitor cocktail and 25 unit/mL benzonase fresh)

(~150 µL/cell pellet) for 30 min on ice without vortexing or spinning (only pipetting to

homogenize). Protein concentrations in lysates were assessed by DC protein assay and

adjusted to 1 µg/µL in a total volume of 50 µL. Samples (50 µg of proteins) were denatured

in a 1:1 volume of denaturation buffer (2% SDS and 50 mM DTT) at 98°C for 10 min.

Denatured samples were loaded in duplicate over a 0.2 µm cellulose acetate (OE66)

membrane positioned in the filter retardation apparatus (Whatman, The Convertible

filtration manifold system cat. Series 1055). The membrane was washed with a 0.2% SDS

solution once prior to loading the samples, and twice after (100 µL/wash). The membrane

was blocked in 3% BSA TBST (0.1% Tween 20) for an hour and probed by standard

immunoblotting analysis with primary mouse anti-HTT (MAB2166) and rabbit anti-GFP

antibodies, followed by secondary antibodies and detection using the LI-COR Odyssey

immunoblotting system. The quantity of aggregates retained on the filter expressed was

measured with Image Studio Lite and calculated relatively to the PBS control group

(arbitrarily set at 1) and technical replicates were averaged.

LDH activity cytotoxicity assay

Cytotoxicity was assessed in COS-7 transiently expressing human HTT or YAC128

neurons after 6 and 24h of PalmB treatment, using the CYQUANT LDH cytotoxicity assay

kit, as per the manufacturer's guidelines. Lactate dehydrogenase (LDH) is a well-defined

and reliable indicator of cytotoxicity. Extracellular LDH was here quantified by a coupled

enzymatic reaction in which LDH catalyzes the conversion of lactate to pyruvate via NAD+

reduction to NADH. This can be measured by using an excitation of 560 nm and an

emission of 590 nm. A cell density of 10,000 cells/well was chosen after experimental

optimization. Fluorescence was read on an EnSpire 2300 Multilabel plate reader from

PerkinElmer using the EnSpire manager software. Cytotoxicity (in %) was calculated

using the formula:

$$\% \, Cytotoxicity = \left(\frac{Compound \; treated \; LDH \; activity^1 - Spontaneous \; LDH \; activity^2}{Maximum \; LDH \; activity^3 - Spontaneous \; LDH \; activity^2}\right) \times 100$$

[1] fluorescence in transfected COS-7 cells or YAC128 neurons treated with PalmB (x%),

[2] fluorescence in untransfected COS-7 cells untreated or YAC128 neurons treated with DMSO (0%),

[3] fluorescence in untransfected COS-7 cells or YAC128 neurons treated with 10X lysis buffer (100%).

2.2.7. Statistical Analysis

GraphPad Prism 9 was used for all statistical analysis and graph preparation. Figures

were generated in Adobe illustrator 2020. All data are presented as mean ± SEM.

Biological replicates (n) are displayed on each graph, or indicated in the captions.

Student's t-test or 1- or 2-way ANOVA statistical tests with post-hoc analysis (Tukey,

Dunnett, Sidak or Bonferroni) were used for all experiments. In some cases, t-tests were

performed between specific groups after an ANOVA test when a planned comparison

was established during the experimental design process. Significance for these tests is

noted with the "#" symbol instead of "*".

3. Results

3.1. Palmitoylation levels of HTT and HIP14/HIP14L substrates are decreased in the BACHD and in two humanized HD mouse models

We first wanted to assess in a low throughput, high confidence manner how the presence of mHTT impacts protein palmitoylation in multiple mouse models of HD, including BACHD, YAC128, Hu97/18 and Hu128/21 (**Table 3**). We therefore compared levels of palmitoylation of HTT, HIP14 and/or HIP14L substrates 25 kDa synaptosomal-associated protein (SNAP25), 95 kDa post-synaptic density protein (PSD95), and 65 kDa glutamate decarboxylase (GAD65) in the brains of 6-month-old HD mouse lines and their respective control line (**Figure 1**).

In the BACHD model (**Figure 1.B**), the palmitoylation level of mHTT Q97 was significantly lower (70%) than palmitoylation of mouse Htt Q7 in the WT or HD brains (1-way ANOVA (1WA): $p<0.0001$; Tukey's post-test (Tk): $p<0.0001$ for Q97 vs. Q7 in WT or HD brains). In the humanized Hu97/18 and Hu18/18 brains (**Figure 1.C** - 1WA: $p=0.0002$), the palmitoylation level of mHTT Q97 was significantly lower than wtHTT Q18 in the Hu97/18 (50%; Tk: $p=0.0002$) and Hu18/18 (40%; Tk: $p=0.0023$) brains. In the Hu128/21 and Hu21/21 brains (**Figure 1.D** - 1WA: $p<0.0001$), the palmitoylation level of mHTT Q128 was significantly lower than wtHTT Q21 in the Hu128/21 (20%; Tk: $p=0.0004$) and Hu21/21 (30%; Tk: <0.0001) brains. In the YAC128 model (**Figure 1.A** - 1WA: $p<0.0001$), palmitoylation of mHTT Q128 was significantly decreased compared to endogenous mouse Htt Q7 expressed in the YAC128 brain (60%; Tk: $p<0.0001$) and in the WT littermate control brain (70%; Tk: $p<0.0001$), consistent with the 80% decrease of mHTT Q128 compared to wtHTT Q18 in the YAC18 brain (Yanai et al., 2006).

HIP14 (ZDHHC17) and HIP14L (ZDHHC13) show substrate specificity for

neuronal proteins, including SNAP25, PSD95, and GAD65 (Huang et al., 2004), which all

play important roles in synaptic transmission. We therefore assessed palmitoylation levels

of these proteins in all four HD mouse models. SNAP25 palmitoylation was decreased by

30% in the YAC128 brain (**Figure 1.E** - Student's test, p=0.031), 40% in the BACHD brain

(**Figure 1.F** - p=0.0017), 45% in the Hu97/18 brain (**Figure 1.G** - p=0.041) and 20% in

the Hu128/21 brain (**Figure 1.H** - p=0.029) compared to their respective controls. PSD95

palmitoylation was also significantly decreased in the YAC128 brain (**Figure 1.I** - 35%;

p<0.0001) and the BACHD brain (**Figure 1.J** - 25%; p=0.020) compared to WT littermate

control brains. PSD95 palmitoylation showed a trend towards being reduced in the

Hu97/18 brain (**Figure 1.K** - 23%; p=0.10), but was unchanged in the Hu128/21 brain

(**Figure 1.L** - p=0.98) compared to controls. In contrast, palmitoylation of GAD65 was

unchanged in the YAC128, BACHD and Hu97/18 mouse models (**Figure 1.M-N-O**), but

was decreased by 15% in the brain of Hu128/21 (**Figure 1.P** - p=0.043) compared to the

Hu21/21 mice.

Our data shows that palmitoylation levels of HTT and various HIP14/14L

substrates are reproducibly reduced in all four HD mouse models. The absence of

endogenous Htt in the humanized HD brain does not exacerbate protein palmitoylation

dysregulation but may reveal subtle differences in palmitoylation dysregulation for GAD65

and PSD95.

3.2. Palmitoylation of mutant HTT is decreased early in the brain of the humanized HD mouse model, Hu128/21

Our next goal was to investigate if aging affects palmitoylation of HTT in the brain of humanized Hu128/21 HD mice. Palmitoylation of wtHTT and mHTT in the brain of Hu21/21 and Hu128/21 mice at 1, 3, 6 and 12 months of age was measured using the ABE assay (**Figure 2.A-B-C-D**). At all the ages investigated, the palmitoylation level of mHTT was significantly lower compared to wtHTT in either the Hu21/21 brain (30-40%) or the Hu128/21 brain (20-40%) (1WA: $p<0.01$ at 1 and 3 months, $p<0.001$ at 6 and 12 months). Notably, the decreased palmitoylation of mHTT was most pronounced at 6 and 12 months of age (**Figure 2.C and D**).

We then directly compared palmitoylation levels of wt and mHTT at different ages in Hu21/21 (**Figure 2.E**) or Hu128/21 brain lysates (**Figure 2.F**) and found that palmitoylation levels of wtHTT (1WA, $p=0.0022$) and mHTT (2-way ANOVA (2WA), $p<0.0001$) significantly declined with age, reaching a 30% decrease at 12 months compared to wt and mHTT palmitoylation levels at 1 month. This decline of HTT palmitoylation with age was observed for both wt and mHTT protein, independent of the presence of the HD mutation in the mouse brain. Taken together, our data demonstrates that palmitoylation of HTT is decreased with age and further decreased in the presence of the HD mutation as early as 1 month of age in the humanized Hu128/21 HD mouse model.

3.3. HTT palmitoylation levels in human lymphoblasts decrease with increasing polyQ tract length

To further characterize palmitoylation dysregulation in HD, we assessed the impact of the polyQ tract length on HTT palmitoylation in control and HD patient lymphoblasts (**Figure 3.A, Table 4**). In the control lymphoblasts, wtHTT polyQ tracts were approximately 17Qs in length. In the HD lymphoblasts, we stratified samples into two sub-groups where mHTT polyQ tract lengths were either approximately 44Q or above 59Q. The wtHTT protein in the HD lines had polyQ tracts with 17Qs.

We quantified the palmitoylation levels of wt or mHTT separately (**Figure 3.B**) or HTT (wt and mHTT together) global palmitoylation level (**Figure 3.C**) and observed that mHTT and global HTT palmitoylation levels were significantly decreased with increased polyQ tract length (1WA: (**B**) p=0.027, Tk: p=0.016 for mHTT (>59Q) vs. wtHTT (17Q) and p=0.057 for mHTT (>59Q) vs. wtHTT (~44Q); (**C**) p=0.037, Tk: p=0.031 for total HTT(>59Q) vs. total HTT (17Q)). The exponential relationship between the global palmitoylation level of HTT (wt and mHTT together) and the polyQ tract length (high polyQ) in each lymphoblastoid line is displayed in **Figure 3.D** and supports a negative linear relationship between the two parameters (correlation, p(two-tailed) =0.018).

Palmitoylation levels were also investigated in isogenic HD human pluripotent stem cells (hPSCs) previously generated (Ooi et al., 2019; Xu et al., 2017) using the ABE assay (**Figure 3.E**). The exponential relationship between the global palmitoylation level of HTT (WT and mHTT together) and the polyQ tract length (high polyQ) in each hPSC line (**Figure 3.F**) suggests a negative linear relationship between palmitoylation and polyQ length without reaching statistical significance (correlation, p(two-tailed) =0.14).

Our findings in patient-derived cell lines further validate the impact of the HD

mutation on HTT palmitoylation level and suggest that palmitoylation levels of wt and

mHTT decreases with increasing polyQ length.

3.4 ASO-mediated mHTT lowering in the brain of the YAC128 HD mouse model does

not rescue palmitoylation of HIP14 and HIP14L synaptic substrates

We next wanted to evaluate the impact of mHTT lowering in the CNS on

palmitoylation and function of synaptic proteins. Five-month-old YAC128 mice received

intracerebroventricular (ICV) injection of either PBS (n=6) or an ASO targeting human

HTT (HTT-ASO; n=7) (Southwell et al., 2015). Brains were collected 1 month later, and

palmitoylation of HTT, SNAP25, PSD95 and GAD65 was measured using the ABE assay

(**Figure 4**). As expected, mHTT levels were lowered by approximately 90%, whereas

levels of mouse Htt remained unchanged (**Figure 4.A**).

We measured a 30% reduction of mHTT palmitoylation compared to mouse Htt

(2WA: $p<0.0001$, HD mutation, $p<0.0001$) in the PBS controls (Tk: $p=0.049$) and a 60%

reduction in the ASO treated animals (Tk: $p<0.0001$), but the ASO treatment did not

change Htt and mHTT palmitoylation levels (**Figure 4.B**). ASO treatment also did not

significantly alter the palmitoylation level of SNAP25 (**Figure 4.C**), PSD95 (**Figure 4.D**)

and GAD65 (**Figure 4.E**). Taken together, mHTT lowering in the brain of the YAC128

mouse did not rescue aberrant palmitoylation of the proteins examined.

3.5. Inhibition of APT increases palmitoylation levels of HTT in COS-7 cells, YAC128 cortico-striatal primary neurons and HD patient-derived lymphoblasts

Since ASO-mediated mHTT lowering *in vivo* did not rescue the reduced palmitoylation observed in YAC128 brains, we hypothesized that promoting this modification pharmacologically may represent a relevant therapeutic approach for HD. Therefore, our next objective was to assess if HTT palmitoylation levels can be upregulated in various HD models using alternative approaches. To test this, we chose to reduce protein depalmitoylation via the inhibition of the APT enzymes that catalyze thioester hydrolysis of palmitoylated cysteine residues (**Figure 5**). We chose the well characterized and membrane-permeable chiral acyl-β-lactone Palmostatin B (PalmB) (IC_{50} = 5.4 nM) that inhibits the APT1 and APT2 enzymes (Dekker et al., 2010; Vujic et al., 2016). PalmB has also been shown to partially inhibits the activity of FASN, PNPLA6, and ABHD proteins (Lin and Conibear, 2015a).

The impact of PalmB treatment on mHTT palmitoylation levels in COS-7 cells transiently expressing N-terminal fragments HTT_{1-548} (N548), HTT_{1-1212} (N1212) or full length HTT_{1-3144} 18Q (wt) or 128Q was assessed by metabolic labelling with palmitic acid alkyne followed by bioorthogonal labelling (click chemistry) (**Figure 5.A**). The PalmB concentration of 15 µM was selected after testing a wide range of concentrations (0, 10, 15, 20 and 50 µM) in our experimental models. Palmitoylation levels of N548, N1212 and full length HTT were significantly decreased in the presence of the HD mutation, with a stronger effect for shorter fragments (2WA: HD mutation, p<0.0001 for N548 and N1212, and p=0.0002 for full length HTT). Treatment of COS-7 cells with 15 µM PalmB for 6h significantly increased the palmitoylation levels of 15Q and 128Q N548, N1212 and full

length HTT compared to the respective DMSO controls (2WA: PalmB effect: p<0.0001 for

N548 and full length HTT, and p=0.0007 for N1212). Notably, palmitoylation levels of full

length mHTT were restored to those of wtHTT following treatment with PalmB.

We next used primary cortico-striatal neuronal cultures from YAC18 and YAC128 mice to

evaluate the impact of PalmB treatment on HTT palmitoylation levels (**Figure 5.B**). As

expected, the palmitoylation level of mHTT was significantly decreased compared to

mouse Htt and wtHTT (2WA: HD mutation, p<0.0001). Treatment with PalmB significantly

increased human HTT palmitoylation (2WA: PalmB effect, p=0.032). In the YAC128

neuronal cultures, wt and mHTT palmitoylation was significantly increased by the PalmB

treatment compared to the DMSO control by approximately 20% (Bonferroni's test (Bf):

p=0.041 and p=0.028). In the YAC18 neuronal culture, the effect of PalmB on the

palmitoylation levels of mouse Htt and human wtHTT (not distinguishable with our

protocol) was weaker (Bf: p=0.54).

Finally, we explored the impact of PalmB treatment on HTT palmitoylation in control and

HD-patient derived lymphoblasts (**Figure 5.C**). Notably, treatment with PalmB

significantly increased HTT palmitoylation in this model (2WA: PalmB effect, p=0.0071),

rescuing mHTT palmitoylation levels to wtHTT palmitoylation level.

475   Inhibiting the activity of the APT enzymes with PalmB treatment significantly

476   increased full length HTT palmitoylation levels in immortalized cell lines, primary YAC128

477   neuronal cultures and HD patient-derived cells. Therefore, the impact of the HD mutation

478   on mHTT palmitoylation is reversible and can be modulated using small molecules.

3.6. Inhibition of APTs with PalmB treatment reduces mHTT-induced cytotoxicity in COS-7 cells and YAC128 primary neurons

Our last objective was to investigate if increasing mHTT palmitoylation levels through APT inhibition would impact mHTT inclusion formation and/or mHTT-induced cell toxicity in an *in vitro* HD model. We measured the impact of PalmB treatment in COS-7 cells transfected with mutant 138Q HTT_{1-588}-YFP (138Q-N588-YFP) on the quantity of insoluble mHTT inclusions by filter retardation assay (**Figure 6**). We confirmed that PalmB treatment increased 138Q-N588-YFP palmitoylation level (by 30%) compared to the DMSO treated control (p=0.029) (**Figure 6.A**), and that the amount of insoluble HTT inclusions were significantly higher when 138Q-N588-YFP is expressed compared to 17Q-N588-YFP (p<0.0001) (**Figure 6.B**) consistent with a previous study (D. D. O. Martin et al., 2018). Treatment with PalmB for 48h significantly decreased the formation of insoluble inclusions by 20-22% (p=0.038 and 0.0065 with HTT or GFP antibodies) (**Figure 6.C**).

The impact of APT inhibition on cytotoxicity was measured using the LDH activity assay in COS-7 cells transiently expressing 15Q or $128Q$-HTT_{1-3144}. First, we confirmed that the expression of full length 128Q-HTT increased cytotoxicity (relative to untransfected cells) compared to 15Q-HTT (p=0.019) (**Figure 6.D**). Then, we measured the impact of PalmB treatment for 6 or 24h on cytotoxicity in COS-7 expressing 128Q-HTT (**Figure 6.E**). Negative cytotoxicity values correspond to a protective effect of the treatment. Treatment with PalmB for 6 and 24h significantly reduced cytotoxicity in COS-7 cells expressing 128Q-HTT relatively to DMSO controls (2WA: PalmB effect, p<0.0001). The effect of PalmB on cytotoxicity compared to the DMSO control was the strongest at

a concentration of 30 µM, but was also significant at lower concentrations (Dunnett's test:

p<0.05).

The impact of PalmB treatment on cytotoxicity was also assessed in YAC128 cortico-

striatal primary neurons (**Figure 6.F**). Cytotoxicity was significantly reduced by the PalmB

treatment compared to neuron treated with DMSO (2WA: PalmB effect, p=0.0014). The

effect of PalmB against cytotoxicity compared to the DMSO treated control was the

strongest with 10 µM and 15 µM after 6 and 24h or treatment, respectively (Dunnett's

test: p=0.048 and 0.0098).

Altogether, the data shows that inhibiting of APTs with PalmB reduces mHTT

inclusion levels and mHTT-induced cytotoxicity in COS-7 cells and YAC128 cortico-

striatal primary neurons.

4. Discussion

Targeting the causative agent of HD, by lowering levels of the toxic mHTT protein,

is one of the therapeutic strategies for the treatment of this disease (Caron et al., 2018).

This can be achieved by suppressing transcription of *HTT*, reducing the synthesis of the

toxic protein or increasing its degradation in cells. Promising strategies to promote mHTT

degradation use mHTT-LC3 linker compounds (Li et al., 2019) or proteolysis-targeting

chimera (PROTAC) based approaches (Harding and Tong, 2018). Another approach has

been to target specific PTMs of HTT that are dysregulated in presence of the HD mutation

in order to promote neuronal survival (Aharony et al., 2015; Bowie et al., 2018; Di Pardo

et al., 2012; Ehrnhoefer et al., 2011; Gu et al., 2009; Kratter et al., 2016).

The challenge with this strategy lies in the identification of enzymes adding or

removing PTMs with a high specificity for HTT. Most kinases, phosphatases or caspases

have a large variety of substrates. Developing therapeutically relevant pharmacological

modulators of PTM requires that they be developed within the context of well-

characterized PTM-related signaling systems. However, there are still many knowledge

gaps regarding the role of PTM dysregulation on mHTT toxicity in human tissues that will

need to be addressed to develop effective therapies.

A potential PTM candidate is palmitoylation of HTT at cysteine 214, catalyzed by

HIP14 (ZDHHC17) and HIP14L (ZDHHC13) (Yanai et al., 2006). Here, we showed that

palmitoylation levels of HTT and various HIP14/14L substrates were reproducibly reduced

in a variety of HD models, including humanized HD mouse models, HD patient-derived

lymphoblast and isogenic hiPSCs (**Figures 1 and 3**). Palmitoylation of HTT was

decreased as early as 1 month of age in the HD humanized mouse model and

palmitoylation levels were further decreased with aging (**Figure 2**). Lowering mHTT levels

in the YAC128 mouse brain was not sufficient to rescue palmitoylation levels of synaptic

HIP14/14L substrates (**Figure 4**). Treatment of COS-7 cells, YAC128 cortico-striatal

neurons and in HD patient derived lymphoblast with Palm B, a small molecule that targets

mainly APTs 1 and 2, normalized palmitoylation of full length mHTT (**Figure 5**). Finally,

modulating mHTT palmitoylation was beneficial against mHTT-induced cytotoxicity

(**Figure 6**).

Our first aim was to investigate how the HD mutation impacts HTT palmitoylation

in multiple mouse and human HD models to address model biases and technical

limitations previously encountered and to obtain a more accurate representation of

palmitoylation dysregulation that may occur in HD patients. The palmitoylation of mHTT

was previously shown to be reduced by 50% in COS-7 cells expressing 15Q and 128Q

N548 and metabolically labelled with [^{3}H]-palmitate, and by 80% in the YAC128 HD

mouse brain compared to wtHTT in the YAC18 brain (a different mouse line than YAC128

expressing endogenous mouse 7Q-Htt and human 18Q-HTT) by IP of human HTT (Yanai

et al., 2006) followed by the earliest version of the ABE assay (Drisdel and Green, 2004),

which now has been optimized to yield higher sensitivity (Brigidi and Bamji, 2013).

A potential confounding factor of using the YAC128 and BACHD mouse models is the

presence of two endogenous mouse Htt alleles in addition to the mHTT transgene, which

could compensate for the effect of mHTT on HIP14 and HIP14L activity. To take this into

consideration, the two humanized HD mouse models, Hu97/18 and Hu128/21, which

harbor a mutant and a wtHTT human transgene with no mouse Htt complement, were

also used to investigate palmitoylation in a model that more closely models the human

condition. In all four mouse models of HD, mHTT palmitoylation was reproducibly and

consistently decreased both compared to littermate control mice and compared to the WT

protein within the same mouse (**Figure 1**).

To assess HTT palmitoylation in a more physiologically relevant system, we used

control and HD patient-derived lymphoblasts. We observed that palmitoylation of HTT is

decreased with increasing number of polyQ repeats (**Figure 3.A-D**). This is, to our

knowledge, the first demonstration of aberrant palmitoylation of mHTT in a HD human

cell model. This supports the finding that decreased palmitoylation of mHTT in the brains

of a variety of HD mouse models is not an artifact due to lower affinity of mouse HIP14/14L

(or potentially other PATs) for the human huntingtin protein, but a robust biochemical

phenotype. Previous studies have looked at palmitoylation dysregulation in HD models

with extreme polyQ lengths (>90) rarely seen in the clinic (Huang et al., 2011; Yanai et

al., 2006). Our results now suggest that aberrant palmitoylation of HTT occurs within a

range of clinically-relevant polyQ-repeat lengths found in HD patients. We also provide

the first evidence that mHTT palmitoylation levels are inversely correlated with polyQ

repeat length. The data generated in two sets of isogenic control and HD (IsoHD) hPSC

lines (**Figure 3.E-F**) suggests that the differences of palmitoylation level are not due to

additional factors such as genetic variants beside the *HTT* gene.

Our next objective was to investigate the earliest timepoint at which palmitoylation

is altered in the humanized HD mouse model, Hu128/21 (**Figure 2**). We showed that

palmitoylation of mHTT was decreased very early in the whole brain (at 1 month), before

onset of behavioural and neuropathological HD-like phenotypes (Southwell et al., 2017),

making it one of the earliest phenotypes observed in this model. This downregulation of

mHTT palmitoylation is maintained at 3, 6 and 12 months in the brain of Hu128/21.

Additional studies are required to determine if mHTT is hypo-palmitoylated during

embryonic development.

We report here for the first time that aging led to a decrease of both wtHTT and

mHTT palmitoylation. This suggests that the presence of the HD mutation has the same

impact as aging on HTT palmitoylation levels. The combined impact of aging and the HD

mutation may be additive in reducing palmitoylation levels. The relationship between

palmitoylation levels of proteins in aging have not been extensively investigated (Zamzow

et al., 2019).

Therapeutics that lower toxic mHTT have shown promise in preclinical studies in

HD rodents (Kordasiewicz et al., 2012; Southwell et al., 2018) and are currently being

evaluated in clinical trials for HD (NCT04120493, NCT04000594 and NCT03225833). We

have previously shown that reducing levels of Htt in *Htt*[+/−] mice and FVB neurons treated

with HTT-ASO (~95% HTT knockdown) leads to reduced palmitoylation of HIP14 and

HIP14 substrates such as SNAP25 and GluR1 (Huang et al., 2011). However, the impact

of lowering mHTT on palmitoylation of HIP14 and HIP14L substrates has not been

investigated. Our data from YAC128 animals treated with an ASO against mHTT at 5

months of age and assessed at 6 months of age (**Figure 4;** ~90% reduction of mHTT

levels in the whole brain) suggests that mHTT lowering for 1 month is not sufficient to

rescue palmitoylation alterations in HD. This suggest that specific palmitoylation-targeted

therapies may be necessary. Additional studies to investigate the impact of ASO treatment

at a younger age, or for a longer duration are needed.

Aberrant palmitoylation causes or contributes to a large variety of disorders such

as Alzheimer's disease , schizophrenia, mental retardation and Batten Disease

(Bhattacharyya et al., 2013; Cho and Park, 2016; Hofmann and Lu, 2014; Ko and Dixon,

2018). The concept of developing therapeutically relevant pharmacological modulators of

palmitoylation to promote or reduce palmitoylation of specific proteins has been emerging

over the past years (Chavda et al., 2014; Fraser et al., 2019). In this study, we observed

robust reduction of HTT palmitoylation in a variety of HD models. HTT is dynamically

palmitoylated by HIP14 and 14L and de-palmitoylated by APTs 1 and 2 (also LYPA1 and

2) (Lin and Conibear, 2015b; Yanai et al., 2006). We have previously shown that

increasing mHTT palmitoylation levels *in vitro* by overexpression of HIP14 may be

protective (Yanai et al., 2006), but this approach is not easily translatable *in vivo*. Inhibition

of APTs with the membrane-permeable chiral acyl-β-lactone PalmB (Davda and Martin,

2014; Dekker et al., 2010; Vujic et al., 2016) was shown to increase HTT_{1-548}

palmitoylation levels in COS-7 cells (Lin and Conibear, 2015b). Here, we demonstrate

that PalmB can be used to promote palmitoylation of full length mHTT in multiple models

(**Figure 5**).

This study serves as a proof-of-concept that mHTT palmitoylation can be

increased, and even normalized to wtHTT levels, using a small molecule. Furthermore,

we show that increasing palmitoylation of mHTT exerts beneficial effects on aggregation

and cytotoxicity (**Figure 6**). Further studies evaluating the impact of promoting HTT

palmitoylation on well characterized HD phenotypes will help elucidate the potential

mechanism of action leading to the observed protective effects. PalmB inhibits APT 1 and

2 that depalmitoylate a large variety of protein substrates, and also targets FASN,

PNPLA6, and ABHD proteins (Lin and Conibear, 2015a). Testing additional APT inhibitors

(Vujic et al., 2016) or developing new ones specifically disrupting the interaction between

APTs and HTT or HIP14 to evaluate the impact of promoting HTT palmitoylation in

preclinical models will be critical.

Protective effects of promoting palmitoylation of HTT and other substrates may be

mediated by increased mHTT clearance, or reduced proteolytic cleavage of mHTT by

caspase 6 (CASP6) at D586. We have indeed recently shown that CASP6 is

palmitoylated by HIP14, resulting in CASP6 inactivation (Lontay et al., 2020; Skotte et al.,

2017) which would be expected to reduce cleavage and potentially provide beneficial

effects.

Acknowledgments & Funding

The authors would like to thank Mark Wang, Xiaofan Qu and Qingwen Xia for their

technical support, Dr. Amirah Ali, Dr. Philip Ly, Hailey Findlay-Black, Jennifer Collins, Dr.

Niels Skotte, Dr. Amber Southwell and Stephanie Bortnick for their support on the project.

FLL was funded by a Canadian Institutes of Health Research (CIHR) Postdoctoral

Fellowship. NSC was funded by the Ripples of Hope Pfizer Trainee Award, a Canadian

Institutes for Health Research Postdoctoral Fellowship, and the Huntington Disease

Society of America Berman/Topper Family HD Career Development Fellowship. MES

was supported by a Vanier Canada Graduate Scholarship and a University of British

Columbia 4-Year Graduate Fellowship. MAP is Michael Smith Foundation for Health

Research Scholar. Project operational support for MRH provided by a CIHR Foundation

grant (FDN-154278) and by the CHDI Foundation.

Conflict of Interest

650 The authors have no conflict of interest to report.

651   **BIBLIOGRAPHY**

652   Aharony, I., Ehrnhoefer, D.E., Shruster, A., Qiu, X., Franciosi, S., Hayden, M.R., Offen,
653   D., 2015. A Huntingtin-based peptide inhibitor of caspase-6 provides protection
654   from mutant Huntingtin-induced motor and behavioral deficits. Hum Mol Genet
655   24, 2604–2614. https://doi.org/10.1093/hmg/ddv023
656   Aiken, C.T., Steffan, J.S., Guerrero, C.M., Khashwji, H., Lukacsovich, T., Simmons, D.,
657   Purcell, J.M., Menhaji, K., Zhu, Y.-Z., Green, K., Laferla, F., Huang, L.,
658   Thompson, L.M., Marsh, J.L., 2009. Phosphorylation of threonine 3: implications
659   for Huntingtin aggregation and neurotoxicity. J. Biol. Chem. 284, 29427–29436.
660   https://doi.org/10.1074/jbc.M109.013193
661   Atwal, R.S., Desmond, C.R., Caron, N., Maiuri, T., Xia, J., Sipione, S., Truant, R., 2011.
662   Kinase inhibitors modulate huntingtin cell localization and toxicity. Nat. Chem.
663   Biol. 7, 453–460. https://doi.org/10.1038/nchembio.582
664   Atwal, R.S., Xia, J., Pinchev, D., Taylor, J., Epand, R.M., Truant, R., 2007. Huntingtin
665   has a membrane association signal that can modulate huntingtin aggregation,
666   nuclear entry and toxicity. Hum. Mol. Genet. 16, 2600–2615.
667   https://doi.org/10.1093/hmg/ddm217
668   Bhattacharyya, R., Barren, C., Kovacs, D.M., 2013. Palmitoylation of Amyloid Precursor
669   Protein Regulates Amyloidogenic Processing in Lipid Rafts. J Neurosci 33,
670   11169–11183. https://doi.org/10.1523/JNEUROSCI.4704-12.2013
671   Bowie, L.E., Maiuri, T., Alpaugh, M., Gabriel, M., Arbez, N., Galleguillos, D., Hung,
672   C.L.K., Patel, S., Xia, J., Hertz, N.T., Ross, C.A., Litchfield, D.W., Sipione, S.,
673   Truant, R., 2018. N6-Furfuryladenine is protective in Huntington's disease
674   models by signaling huntingtin phosphorylation. PNAS 115, E7081–E7090.
675   https://doi.org/10.1073/pnas.1801772115
676   Brigidi, G.S., Bamji, S.X., 2013. Detection of protein palmitoylation in cultured
677   hippocampal neurons by immunoprecipitation and acyl-biotin exchange (ABE). J
678   Vis Exp. https://doi.org/10.3791/50031
679   Cariulo, C., Azzollini, L., Verani, M., Martufi, P., Boggio, R., Chiki, A., Deguire, S.M.,
680   Cherubini, M., Gines, S., Marsh, J.L., Conforti, P., Cattaneo, E., Santimone, I.,
681   Squitieri, F., Lashuel, H.A., Petricca, L., Caricasole, A., 2017. Phosphorylation of
682   huntingtin at residue T3 is decreased in Huntington's disease and modulates
683   mutant huntingtin protein conformation. PNAS 114, E10809–E10818.
684   https://doi.org/10.1073/pnas.1705372114
685   Cariulo, C., Verani, M., Martufi, P., Ingenito, R., Finotto, M., Deguire, S.M., Lavery, D.J.,
686   Toledo-Sherman, L., Lee, R., Doherty, E.M., Vogt, T.F., Dominguez, C., Lashuel,
687   H.A., Petricca, L., Caricasole, A., 2020. Ultrasensitive quantitative measurement
688   of huntingtin phosphorylation at residue S13. Biochem. Biophys. Res. Commun.
689   521, 549–554. https://doi.org/10.1016/j.bbrc.2019.09.097
690   Caron, N.S., Desmond, C.R., Xia, J., Truant, R., 2013. Polyglutamine domain flexibility
691   mediates the proximity between flanking sequences in huntingtin. Proceedings of
692   the National Academy of Sciences of the United States of America 110, 14610–
693   14615. https://doi.org/10.1073/pnas.1301342110
694   Caron, N.S., Dorsey, E.R., Hayden, M.R., 2018. Therapeutic approaches to Huntington
695   disease: from the bench to the clinic. Nat Rev Drug Discov 17, 729–750.
696   https://doi.org/10.1038/nrd.2018.133

Carroll, J.B., Warby, S.C., Southwell, A.L., Doty, C.N., Greenlee, S., Skotte, N., Hung,
G., Bennett, C.F., Freier, S.M., Hayden, M.R., 2011. Potent and selective
antisense oligonucleotides targeting single-nucleotide polymorphisms in the
Huntington disease gene / allele-specific silencing of mutant huntingtin. Mol Ther
19, 2178–2185. https://doi.org/10.1038/mt.2011.201
Chavda, B., Arnott, J.A., Planey, S.L., 2014. Targeting protein palmitoylation: selective
inhibitors and implications in disease. Expert Opin Drug Discov 9, 1005–1019.
https://doi.org/10.1517/17460441.2014.933802
Cho, E., Park, M., 2016. Palmitoylation in Alzheimer's disease and other
neurodegenerative diseases. Pharmacol. Res. 111, 133–151.
https://doi.org/10.1016/j.phrs.2016.06.008
Davda, D., Martin, B.R., 2014. Acyl protein thioesterase inhibitors as probes of dynamic
S-palmitoylation. Medchemcomm 5, 268–276.
https://doi.org/10.1039/C3MD00333G
Dekker, F.J., Rocks, O., Vartak, N., Menninger, S., Hedberg, C., Balamurugan, R.,
Wetzel, S., Renner, S., Gerauer, M., Schölermann, B., Rusch, M., Kramer, J.W.,
Rauh, D., Coates, G.W., Brunsveld, L., Bastiaens, P.I.H., Waldmann, H., 2010.
Small-molecule inhibition of APT1 affects Ras localization and signaling. Nat.
Chem. Biol. 6, 449–456. https://doi.org/10.1038/nchembio.362
Di Pardo, A., Maglione, V., Alpaugh, M., Horkey, M., Atwal, R.S., Sassone, J.,
Ciammola, A., Steffan, J.S., Fouad, K., Truant, R., Sipione, S., 2012. Ganglioside
GM1 induces phosphorylation of mutant huntingtin and restores normal motor
behavior in Huntington disease mice. Proc. Natl. Acad. Sci. U.S.A. 109, 3528–
3533. https://doi.org/10.1073/pnas.1114502109
Drisdel, R.C., Green, W.N., 2004. Labeling and quantifying sites of protein
palmitoylation. BioTechniques 36, 276–285.
Ehrnhoefer, D.E., Sutton, L., Hayden, M.R., 2011. Small changes, big impact:
posttranslational modifications and function of huntingtin in Huntington disease.
Neuroscientist 17, 475–492. https://doi.org/10.1177/1073858410390378
Evans, S.J.W., Douglas, I., Rawlins, M.D., Wexler, N.S., Tabrizi, S.J., Smeeth, L., 2013.
Prevalence of adult Huntington's disease in the UK based on diagnoses recorded
in general practice records. J. Neurol. Neurosurg. Psychiatry 84, 1156–1160.
https://doi.org/10.1136/jnnp-2012-304636
Fisher, E.R., Hayden, M.R., 2014. Multisource ascertainment of Huntington disease in
Canada: prevalence and population at risk. Mov. Disord. 29, 105–114.
https://doi.org/10.1002/mds.25717
Fraser, N.J., Howie, J., Wypijewski, K.J., Fuller, W., 2019. Therapeutic targeting of
protein S-acylation for the treatment of disease. Biochemical Society
Transactions 48, 281–290. https://doi.org/10.1042/BST20190707
Fukata, Y., Fukata, M., 2010. Protein palmitoylation in neuronal development and
synaptic plasticity. Nat. Rev. Neurosci. 11, 161–175.
https://doi.org/10.1038/nrn2788
Globa, A.K., Bamji, S.X., 2017. Protein palmitoylation in the development and plasticity
of neuronal connections. Curr Opin Neurobiol 45, 210–220.
https://doi.org/10.1016/j.conb.2017.02.016

Graham, R.K., Deng, Y., Slow, E.J., Haigh, B., Bissada, N., Lu, G., Pearson, J.,
Shehadeh, J., Bertram, L., Murphy, Z., Warby, S.C., Doty, C.N., Roy, S.,
Wellington, C.L., Leavitt, B.R., Raymond, L.A., Nicholson, D.W., Hayden, M.R.,
2006. Cleavage at the caspase-6 site is required for neuronal dysfunction and
degeneration due to mutant huntingtin. Cell 125, 1179–1191.
https://doi.org/10.1016/j.cell.2006.04.026
Gray, M., Shirasaki, D.I., Cepeda, C., André, V.M., Wilburn, B., Lu, X.-H., Tao, J.,
Yamazaki, I., Li, S.-H., Sun, Y.E., Li, X.-J., Levine, M.S., Yang, X.W., 2008. Full-
length human mutant huntingtin with a stable polyglutamine repeat can elicit
progressive and selective neuropathogenesis in BACHD mice. J. Neurosci. 28,
6182–6195. https://doi.org/10.1523/JNEUROSCI.0857-08.2008
Gu, X., Greiner, E.R., Mishra, R., Kodali, R., Osmand, A., Finkbeiner, S., Steffan, J.S.,
Thompson, L.M., Wetzel, R., Yang, X.W., 2009. Serines 13 and 16 are critical
determinants of full-length human mutant huntingtin induced disease
pathogenesis in HD mice. Neuron 64, 828–840.
https://doi.org/10.1016/j.neuron.2009.11.020
Harding, R.J., Tong, Y., 2018. Proteostasis in Huntington's disease: disease
mechanisms and therapeutic opportunities. Acta Pharmacol Sin 39, 754–769.
https://doi.org/10.1038/aps.2018.11
Hofmann, S., Lu, J.-Y., 2014. Molecular basis of infantile Batten disease: impact of
disordered palmitoylated protein degradation on the nervous system (466.1). The
FASEB Journal 28, 466.1. https://doi.org/10.1096/fasebj.28.1_supplement.466.1
Huang, K., Sanders, S., Singaraja, R., Orban, P., Cijsouw, T., Arstikaitis, P., Yanai, A.,
Hayden, M.R., El-Husseini, A., 2009. Neuronal palmitoyl acyl transferases exhibit
distinct substrate specificity. FASEB J. 23, 2605–2615.
https://doi.org/10.1096/fj.08-127399
Huang, K., Sanders, S.S., Kang, R., Carroll, J.B., Sutton, L., Wan, J., Singaraja, R.,
Young, F.B., Liu, L., El-Husseini, A., Davis, N.G., Hayden, M.R., 2011. Wild-type
HTT modulates the enzymatic activity of the neuronal palmitoyl transferase
HIP14. Hum Mol Genet 20, 3356–3365. https://doi.org/10.1093/hmg/ddr242
Huang, K., Yanai, A., Kang, R., Arstikaitis, P., Singaraja, R.R., Metzler, M., Mullard, A.,
Haigh, B., Gauthier-Campbell, C., Gutekunst, C.-A., Hayden, M.R., El-Husseini,
A., 2004. Huntingtin-Interacting Protein HIP14 Is a Palmitoyl Transferase
Involved in Palmitoylation and Trafficking of Multiple Neuronal Proteins. Neuron
44, 977–986. https://doi.org/10.1016/j.neuron.2004.11.027
Hubalkova, P., Ladislav, M., Vyklicky, V., Smejkalova, T., Krausova, B.H., Kysilov, B.,
Krusek, J., Naimová, Z., Korinek, M., Chodounska, H., Kudova, E., Cerny, J.,
Vyklicky, L., 2021. Palmitoylation Controls NMDA Receptor Function and Steroid
Sensitivity. J. Neurosci. 41, 2119–2134.
https://doi.org/10.1523/JNEUROSCI.2654-20.2021
Humbert, S., Bryson, E.A., Cordelières, F.P., Connors, N.C., Datta, S.R., Finkbeiner, S.,
Greenberg, M.E., Saudou, F., 2002. The IGF-1/Akt pathway is neuroprotective in
Huntington's disease and involves Huntingtin phosphorylation by Akt. Dev. Cell
2, 831–837.
Jeong, H., Then, F., Melia, T.J., Mazzulli, J.R., Cui, L., Savas, J.N., Voisine, C.,
Paganetti, P., Tanese, N., Hart, A.C., Yamamoto, A., Krainc, D., 2009.

Acetylation Targets Mutant Huntingtin to Autophagosomes for Degradation. Cell
137, 60–72. https://doi.org/10.1016/j.cell.2009.03.018

Kalchman, M.A., Graham, R.K., Xia, G., Koide, H.B., Hodgson, J.G., Graham, K.C.,
Goldberg, Y.P., Gietz, R.D., Pickart, C.M., Hayden, M.R., 1996. Huntingtin is
ubiquitinated and interacts with a specific ubiquitin-conjugating enzyme. J. Biol.
Chem. 271, 19385–19394. https://doi.org/10.1074/jbc.271.32.19385

Kalchman, M.A., Koide, H.B., McCutcheon, K., Graham, R.K., Nichol, K., Nishiyama, K.,
Kazemi-Esfarjani, P., Lynn, F.C., Wellington, C., Metzler, M., Goldberg, Y.P.,
Kanazawa, I., Gietz, R.D., Hayden, M.R., 1997. HIP1 , a human homologue of S.
cerevisiae Sla2p , interacts with membrane-associated huntingtin in the brain.
Nature Genetics 16, 44–53. https://doi.org/10.1038/ng0597-44

Ko, P., Dixon, S.J., 2018. Protein palmitoylation and cancer. EMBO Rep 19.
https://doi.org/10.15252/embr.201846666

Kordasiewicz, H.B., Stanek, L.M., Wancewicz, E.V., Mazur, C., McAlonis, M.M., Pytel,
K.A., Artates, J.W., Weiss, A., Cheng, S.H., Shihabuddin, L.S., Hung, G.,
Bennett, C.F., Cleveland, D.W., 2012. Sustained therapeutic reversal of
Huntington's disease by transient repression of huntingtin synthesis. Neuron 74,
1031–1044. https://doi.org/10.1016/j.neuron.2012.05.009

Kratter, I.H., Zahed, H., Lau, A., Tsvetkov, A.S., Daub, A.C., Weiberth, K.F., Gu, X.,
Saudou, F., Humbert, S., Yang, X.W., Osmand, A., Steffan, J.S., Masliah, E.,
Finkbeiner, S., 2016. Serine 421 regulates mutant huntingtin toxicity and
clearance in mice. J Clin Invest 126, 3585–3597.
https://doi.org/10.1172/JCI80339

Li, Z., Wang, C., Wang, Z., Zhu, C., Li, J., Sha, T., Ma, L., Gao, C., Yang, Y., Sun, Y.,
Wang, J., Sun, X., Lu, C., Difiglia, M., Mei, Y., Ding, C., Luo, S., Dang, Y., Ding,
Y., Fei, Y., Lu, B., 2019. Allele-selective lowering of mutant HTT protein by HTT-
LC3 linker compounds. Nature 575, 203–209. https://doi.org/10.1038/s41586-
019-1722-1

Lin, D.T.S., Conibear, E., 2015a. ABHD17 proteins are novel protein depalmitoylases
that regulate N-Ras palmitate turnover and subcellular localization. eLife 4,
e11306. https://doi.org/10.7554/eLife.11306

Lin, D.T.S., Conibear, E., 2015b. ABHD17 proteins are novel protein depalmitoylases
that regulate N-Ras palmitate turnover and subcellular localization. Elife 4,
e11306. https://doi.org/10.7554/eLife.11306

Lontay, B., Kiss, A., Virág, L., Tar, K., 2020. How Do Post-Translational Modifications
Influence the Pathomechanistic Landscape of Huntington's Disease? A
Comprehensive Review. International Journal of Molecular Sciences 21, 4282.
https://doi.org/10.3390/ijms21124282

Maiuri, T., Mocle, A.J., Hung, C.L., Xia, J., van Roon-Mom, W.M.C., Truant, R., 2017.
Huntingtin is a scaffolding protein in the ATM oxidative DNA damage response
complex. Hum. Mol. Genet. 26, 395–406. https://doi.org/10.1093/hmg/ddw395

Martin, D.D.O., Heit, R.J., Yap, M.C., Davidson, M.W., Hayden, M.R., Berthiaume, L.G.,
2014. Identification of a post-translationally myristoylated autophagy-inducing
domain released by caspase cleavage of huntingtin. Hum. Mol. Genet. 23, 3166–
3179. https://doi.org/10.1093/hmg/ddu027

Martin, D. D. O., Kay, C., Collins, J.A., Nguyen, Y.T., Slama, R.A., Hayden, M.R., 2018.
A human huntingtin SNP alters post-translational modification and pathogenic
proteolysis of the protein causing Huntington disease. Sci Rep 8, 8096.
https://doi.org/10.1038/s41598-018-25903-w
Martin, Dale D. O., Schmidt, M.E., Nguyen, Y.T., Lazic, N., Hayden, M.R., 2018.
Identification of a novel caspase cleavage site in huntingtin that regulates mutant
huntingtin clearance. The FASEB Journal 33, 3190–3197.
https://doi.org/10.1096/fj.201701510RRR
Matt, L., Kim, K., Chowdhury, D., Hell, J.W., 2019. Role of Palmitoylation of
Postsynaptic Proteins in Promoting Synaptic Plasticity. Front. Mol. Neurosci. 12.
https://doi.org/10.3389/fnmol.2019.00008
Mukai, J., Dhilla, A., Drew, L.J., Stark, K.L., Cao, L., MacDermott, A.B., Karayiorgou, M.,
Gogos, J.A., 2008. Palmitoylation-dependent neurodevelopmental deficits in a
mouse model of 22q11 microdeletion. Nature Neuroscience 11, 1302–1310.
https://doi.org/10.1038/nn.2204
Myers, R.H., 2004. Huntington's disease genetics. NeuroRx 1, 255–262.
https://doi.org/10.1602/neurorx.1.2.255
Ochaba, J., Lukacsovich, T., Csikos, G., Zheng, S., Margulis, J., Salazar, L., Mao, K.,
Lau, A.L., Yeung, S.Y., Humbert, S., Saudou, F., Klionsky, D.J., Finkbeiner, S.,
Zeitlin, S.O., Marsh, J.L., Housman, D.E., Thompson, L.M., Steffan, J.S., 2014.
Potential function for the Huntingtin protein as a scaffold for selective autophagy.
Proc. Natl. Acad. Sci. U.S.A. 111, 16889–16894.
https://doi.org/10.1073/pnas.1420103111
Ooi, J., Langley, S.R., Xu, X., Utami, K.H., Sim, B., Huang, Y., Harmston, N.P., Tay,
Y.L., Ziaei, A., Zeng, R., Low, D., Aminkeng, F., Sobota, R.M., Ginhoux, F.,
Petretto, E., Pouladi, M.A., 2019. Unbiased Profiling of Isogenic Huntington
Disease hPSC-Derived CNS and Peripheral Cells Reveals Strong Cell-Type
Specificity of CAG Length Effects. Cell Rep 26, 2494-2508.e7.
https://doi.org/10.1016/j.celrep.2019.02.008
Sanders, S.S., Hou, J., Sutton, L.M., Garside, V.C., Mui, K.K.N., Singaraja, R.R.,
Hayden, M.R., Hoodless, P.A., 2015a. Huntingtin interacting proteins 14 and 14-
like are required for chorioallantoic fusion during early placental development.
Developmental Biology 397, 257–266.
https://doi.org/10.1016/j.ydbio.2014.11.018
Sanders, S.S., Martin, D.D.O., Butland, S.L., Lavallée-Adam, M., Calzolari, D., Kay, C.,
Iii, J.R.Y., Hayden, M.R., 2015b. Curation of the Mammalian Palmitoylome
Indicates a Pivotal Role for Palmitoylation in Diseases and Disorders of the
Nervous System and Cancers. PLOS Computational Biology 11, e1004405.
https://doi.org/10.1371/journal.pcbi.1004405
Sanders, S.S., Parsons, M.P., Mui, K.K.N., Southwell, A.L., Franciosi, S., Cheung, D.,
Waltl, S., Raymond, L.A., Hayden, M.R., 2016. Sudden death due to paralysis
and synaptic and behavioral deficits when Hip14/Zdhhc17 is deleted in adult
mice. BMC Biol. 14, 108. https://doi.org/10.1186/s12915-016-0333-7
Saudou, F., Humbert, S., 2016. The Biology of Huntingtin. Neuron 89, 910–926.
https://doi.org/10.1016/j.neuron.2016.02.003

Schilling, B., Gafni, J., Torcassi, C., Cong, X., Row, R.H., LaFevre-Bernt, M.A., Cusack,
M.P., Ratovitski, T., Hirschhorn, R., Ross, C.A., Gibson, B.W., Ellerby, L.M.,
2006. Huntingtin phosphorylation sites mapped by mass spectrometry.
Modulation of cleavage and toxicity. J. Biol. Chem. 281, 23686–23697.
https://doi.org/10.1074/jbc.M513507200
Schmidt, M.E., Caron, N.S., Aly, A.E., Lemarié, F.L., Dal Cengio, L., Ko, Y., Lazic, N.,
Anderson, L., Nguyen, B., Raymond, L.A., Hayden, M.R., 2020. DAPK1
Promotes Extrasynaptic GluN2B Phosphorylation and Striatal Spine Instability in
the YAC128 Mouse Model of Huntington Disease. Front. Cell. Neurosci. 14.
https://doi.org/10.3389/fncel.2020.590569
Singaraja, R.R., Huang, K., Sanders, S.S., Milnerwood, A.J., Hines, R., Lerch, J.P.,
Franciosi, S., Drisdel, R.C., Vaid, K., Young, F.B., Doty, C., Wan, J., Bissada, N.,
Henkelman, R.M., Green, W.N., Davis, N.G., Raymond, L.A., Hayden, M.R.,
2011. Altered palmitoylation and neuropathological deficits in mice lacking
HIP14. Hum. Mol. Genet. 20, 3899–3909. https://doi.org/10.1093/hmg/ddr308
Skotte, N.H., Sanders, S.S., Singaraja, R.R., Ehrnhoefer, D.E., Vaid, K., Qiu, X.,
Kannan, S., Verma, C., Hayden, M.R., 2017. Palmitoylation of caspase-6 by
HIP14 regulates its activation. Cell Death & Differentiation 24, 433–444.
https://doi.org/10.1038/cdd.2016.139
Slow, E.J., van Raamsdonk, J., Rogers, D., Coleman, S.H., Graham, R.K., Deng, Y.,
Oh, R., Bissada, N., Hossain, S.M., Yang, Y.-Z., Li, X.-J., Simpson, E.M.,
Gutekunst, C.-A., Leavitt, B.R., Hayden, M.R., 2003. Selective striatal neuronal
loss in a YAC128 mouse model of Huntington disease. Hum Mol Genet 12,
1555–1567. https://doi.org/10.1093/hmg/ddg169
Smotrys, J.E., Linder, M.E., 2004. Palmitoylation of Intracellular Signaling Proteins:
Regulation and Function. Annual Review of Biochemistry 73, 559–587.
https://doi.org/10.1146/annurev.biochem.73.011303.073954
Southwell, A.L., Kordasiewicz, H.B., Langbehn, D., Skotte, N.H., Parsons, M.P.,
Villanueva, E.B., Caron, N.S., Østergaard, M.E., Anderson, L.M., Xie, Y., Cengio,
L.D., Findlay-Black, H., Doty, C.N., Fitsimmons, B., Swayze, E.E., Seth, P.P.,
Raymond, L.A., Frank Bennett, C., Hayden, M.R., 2018. Huntingtin suppression
restores cognitive function in a mouse model of Huntington's disease. Sci Transl
Med 10. https://doi.org/10.1126/scitranslmed.aar3959
Southwell, A.L., Skotte, N.H., Villanueva, E.B., Østergaard, M.E., Gu, X., Kordasiewicz,
H.B., Kay, C., Cheung, D., Xie, Y., Waltl, S., Dal Cengio, L., Findlay-Black, H.,
Doty, C.N., Petoukhov, E., Iworima, D., Slama, R., Ooi, J., Pouladi, M.A., Yang,
X.W., Swayze, E.E., Seth, P.P., Hayden, M.R., 2017. A novel humanized mouse
model of Huntington disease for preclinical development of therapeutics targeting
mutant huntingtin alleles. Hum. Mol. Genet. 26, 1115–1132.
https://doi.org/10.1093/hmg/ddx021
Southwell, A.L., Smith, S.E.P., Davis, T.R., Caron, N.S., Villanueva, E.B., Xie, Y.,
Collins, J.A., Ye, M.L., Sturrock, A., Leavitt, B.R., Schrum, A.G., Hayden, M.R.,
2015. Ultrasensitive measurement of huntingtin protein in cerebrospinal fluid
demonstrates increase with Huntington disease stage and decrease following
brain huntingtin suppression. Sci Rep 5, 12166.
https://doi.org/10.1038/srep12166

Southwell, A.L., Warby, S.C., Carroll, J.B., Doty, C.N., Skotte, N.H., Zhang, W.,
Villanueva, E.B., Kovalik, V., Xie, Y., Pouladi, M.A., Collins, J.A., Yang, X.W.,
Franciosi, S., Hayden, M.R., 2013. A fully humanized transgenic mouse model of
Huntington disease. Hum. Mol. Genet. 22, 18–34.
https://doi.org/10.1093/hmg/dds397
Steffan, J.S., Agrawal, N., Pallos, J., Rockabrand, E., Trotman, L.C., Slepko, N., Illes,
K., Lukacsovich, T., Zhu, Y.-Z., Cattaneo, E., Pandolfi, P.P., Thompson, L.M.,
Marsh, J.L., 2004. SUMO modification of Huntingtin and Huntington's disease
pathology. Science 304, 100–104. https://doi.org/10.1126/science.1092194
Sutton, L.M., Sanders, S.S., Butland, S.L., Singaraja, R.R., Franciosi, S., Southwell,
A.L., Doty, C.N., Schmidt, M.E., Mui, K.K.N., Kovalik, V., Young, F.B., Zhang, W.,
Hayden, M.R., 2013. Hip14l-deficient mice develop neuropathological and
behavioural features of Huntington disease. Hum. Mol. Genet. 22, 452–465.
https://doi.org/10.1093/hmg/dds441
Vujic, I., Sanlorenzo, M., Esteve-Puig, R., Vujic, M., Kwong, A., Tsumura, A., Murphy,
R., Moy, A., Posch, C., Monshi, B., Rappersberger, K., Ortiz-Urda, S., 2016. Acyl
protein thioesterase 1 and 2 (APT-1, APT-2) inhibitors palmostatin B, ML348 and
ML349 have different effects on NRAS mutant melanoma cells. Oncotarget 7,
7297–7306. https://doi.org/10.18632/oncotarget.6907
Wanker, E.E., Scherzinger, E., Heiser, V., Sittler, A., Eickhoff, H., Lehrach, H., 1999.
Membrane filter assay for detection of amyloid-like polyglutamine-containing
protein aggregates. Meth. Enzymol. 309, 375–386.
Warby, S.C., Chan, E.Y., Metzler, M., Gan, L., Singaraja, R.R., Crocker, S.F.,
Robertson, H.A., Hayden, M.R., 2005. Huntingtin phosphorylation on serine 421
is significantly reduced in the striatum and by polyglutamine expansion in vivo.
Hum Mol Genet 14, 1569–1577. https://doi.org/10.1093/hmg/ddi165
Xu, X., Ng, B., Sim, B., Radulescu, C.I., Yusof, N.A.B.M., Goh, W.I., Lin, S., Lim, J.S.Y.,
Cha, Y., Kusko, R., Kay, C., Ratovitski, T., Ross, C., Hayden, M.R., Wright, G.,
Pouladi, M.A., 2020. pS421 huntingtin modulates mitochondrial phenotypes and
confers neuroprotection in an HD hiPSC model. Cell Death & Disease 11, 1–12.
https://doi.org/10.1038/s41419-020-02983-z
Xu, X., Tay, Y., Sim, B., Yoon, S.-I., Huang, Y., Ooi, J., Utami, K.H., Ziaei, A., Ng, B.,
Radulescu, C., Low, D., Ng, A.Y.J., Loh, M., Venkatesh, B., Ginhoux, F.,
Augustine, G.J., Pouladi, M.A., 2017. Reversal of Phenotypic Abnormalities by
CRISPR/Cas9-Mediated Gene Correction in Huntington Disease Patient-Derived
Induced Pluripotent Stem Cells. Stem Cell Reports 8, 619–633.
https://doi.org/10.1016/j.stemcr.2017.01.022
Yanai, A., Huang, K., Kang, R., Singaraja, R.R., Arstikaitis, P., Gan, L., Orban, P.C.,
Mullard, A., Cowan, C.M., Raymond, L.A., Drisdel, R.C., Green, W.N.,
Ravikumar, B., Rubinsztein, D.C., El-Husseini, A., Hayden, M.R., 2006.
Palmitoylation of huntingtin by HIP14 is essential for its trafficking and function.
Nat. Neurosci. 9, 824–831. https://doi.org/10.1038/nn1702
Yap, M.C., Kostiuk, M.A., Martin, D.D.O., Perinpanayagam, M.A., Hak, P.G., Siddam,
A., Majjigapu, J.R., Rajaiah, G., Keller, B.O., Prescher, J.A., Wu, P., Bertozzi,
C.R., Falck, J.R., Berthiaume, L.G., 2010. Rapid and selective detection of fatty

acylated proteins using omega-alkynyl-fatty acids and click chemistry. J. Lipid
Res. 51, 1566–1580. https://doi.org/10.1194/jlr.D002790
Zamzow, D.R., Elias, V., Acosta, V.A., Escobedo, E., Magnusson, K.R., 2019. Higher
Levels of Protein Palmitoylation in the Frontal Cortex across Aging Were
Associated with Reference Memory and Executive Function Declines. eNeuro 6.
https://doi.org/10.1523/ENEURO.0310-18.2019

 Figure 1 – Palmitoylation levels of mutant HTT, SNAP25, PSD95 and GAD65 are
reduced in the BACHD, and humanized Hu97/18 and Hu128/21 mouse models of
HD.

Brain lysates from 6-month-old YAC128 and littermate controls (WT), BACHD and WT,
humanized Hu97/18 and Hu18/18, Hu128/21 and Hu21/21 mice were subjected to the
ABE assay. The palmitoylation signal (HAM+) is shown in the top panels for each mouse
line and protein, and the total protein immunoprecipitated is presented in the
corresponding bottom panel. The negative control (HAM-) treatment is displayed on the
right of each set of blots. In the huntingtin (HTT) blots (**A-D**), the lower band corresponds
to Htt or wtHTT, and the upper band to mHTT. The palmitoylation levels for HTT (**A-D**),
SNAP25 (**E-H**), PSD95 (**I-L**) and GAD65 (**M-P**) were calculated as a ratio of palmitoylation
over total protein signal. Mutant HTT was quantified separately from Htt or wtHTT and is
represented by dark grey bars on the graphs, while Htt/wtHTT from the same HD brain is
in light grey.

Statistical analysis: 1-way ANOVA: (**A**) YAC128 and (**B**) BACHD, $p<0.0001$; (**C**) Hu97/18,
$p=0.0002$; (**D**) Hu128/21, $p=0.0013$. Stars indicate Tukey's multiple comparisons test
results. Student's T-test: (**E-P**) results are indicated by stars or p-values on the graphs.

Figure 2 - Palmitoylation of mutant HTT is decreased early in the brain of the humanized Hu128/21 mouse model of HD.

HTT palmitoylation at (**A**) 1, (**B**) 3, (**C**) 6 and (**D**) 12 months measured using the ABE assay in brain lysates from Hu21/21 (WT) and Hu128/21 (HD) mice. The palmitoylation signal (HAM+) is shown in the top panels for each time point, and the total protein immunoprecipitated is presented in the corresponding bottom panel. The negative control (HAM-) treatment is displayed on the right of each set of blots. Wildtype wtHTT (Q21) is the lower band and mHTT (Q128) the upper one. The quantification of 4-9 mice of each genotype is shown and was calculated as a ratio of HTT palmitoylation signal over total HTT and expressed relatively to wtHTT in the Hu21/21 brain. In the Hu128/21 brain, mHTT was quantified separately from wtHTT and is represented by dark grey bars while wtHTT from the same HD mice is represented by light bars.

Effect of aging (1, 3, 6 and 12 months) on the palmitoylation levels of wt and mHTT in
lysates from Hu21/21 (**E**) and Hu128/21 (**F**) brains. Total brain lysates at different ages
were subjected to the ABE assay and run on the same gel for each genotype (n=3-4).
Overlay of both channel (palm-HTT in red and total HTT in green) are displayed in the
bottom panels. Palmitoylation levels of wt and mHTT brain are expressed relatively to
wtHTT palmitoylation at 1 month.

<u>Statistical analysis</u>: 1-way ANOVA: (**A**) p=0.0022; (**B**) p=0.0094; (**C**) p<0.0001; (**D**)
p=0.0002. Stars indicate Tukey's multiple comparisons test results. (**E**) 1-way ANOVA:
p=0.0022. Stars indicate Dunnett's test results compared to 1 month. (**F**) 2-way ANOVA:
interaction, p=0.58; age and allele effect, p<0.0001. Stars indicate Dunnett's test results
compared to wtHTT (Q21) or mHTT (Q128) at 1 month, and therefore only show the effect
of age and not the effect of the HD mutation.

1019
1020

Figure 3 - HTT palmitoylation levels decrease with increasing polyQ tract length in
human lymphoblasts.

(**A**) HTT (wt and mHTT) palmitoylation levels in control and HD-patient derived
lymphoblasts with increasing polyQ tract length (17 to 71). Proteins were resolved using
8% low bis or 3-8% tris-acetate gels. Palmitoylation is shown in the top panels (HAM+),
and the total protein immunoprecipitated with anti-HTT (D7F7) antibodies are shown in
the corresponding bottom panels. The negative control (HAM-) is shown below. Each lane
corresponds to a lymphoblastoid line from a control or HD patients, and represents a
biological replicate (n=4-5 for each group).

(**B**) Quantification of wt or mHTT palmitoylation levels in control and HD lymphoblasts in
the control (17Q), HD (~44Q) and HD (>59Q) groups after protein separation using 8%

low bis gel. The quantification is shown and is calculated as a ratio of HTT palmitoylation
signal over total HTT protein and expressed relatively to wtHTT palmitoylation in the
control group (17Q).
(**C**) Quantification of HTT (wt and mHTT together) global palmitoylation levels measured
in control (17Q), HD (~44Q) and HD (>59Q) lymphoblasts. Global palmitoylation levels of
HTT was calculated as a ratio of wt and mHTT palmitoylation signal over total wt and
mHTT protein and expressed relatively to wtHTT in the control (17Q) group. Mutant HTT
was quantified with wtHTT, as it is challenging to resolve full length HTT alleles with small
ΔQs with commonly used 3-8% tris-acetate gel.

(**D**) Exponential relationship between HTT global palmitoylation levels and the polyQ tract
length (high polyQ) in lymphoblasts.

(**E**) HTT (wt and mutant) palmitoylation levels in HD hPSCs and isogenic controls (Ooi et
al., 2019; Xu et al., 2017) with increasing polyQ tract length (18 to 180). Proteins were
resolved using 3-8% tris-acetate gels. The white arrowhead indicates an unidentified
band only detected with streptavidin. The hPSC parental lines used to generate HD or
control hPSC lines are annotated with an asterisk.

(**F**) Exponential relationship between HTT global palmitoylation level and the polyQ tract
length (high polyQ) in hPSCs.

<u>Statistical analysis</u>: 1-way ANOVA: (**B**) p=0.027; (**C**) p=0.037. Stars indicate Tukey's
multiple comparisons test results. (**D**) Variance explained by polyQ tract length, R^2=0.57.
Pearson r=-0.75; p=0.018. (**F**) Variance explained by polyQ tract length, R^2=0.43. Pearson
r=-0.65; p=0.15.

1054

<u>Figure 4</u> – ASO-mediated mHTT lowering does not rescue aberrant palmitoylation of HIP14/HIP14L synaptic substrates in the brain of the YAC128 mouse model.

Five-month-old YAC128 mice received intracerebroventricular (ICV) injection of either PBS (n=6) or an ASO targeting human HTT (HTT-ASO; n=7), and brains were collected 1 month later.

(**A**) Protein from brain lysates were separated using 3-8% tris-acetate gradient gel and probed with HTT (MAB2166) and calnexin antibodies. HTT levels were quantified by densitometry and normalized to calnexin, and relative levels of mHTT are shown.

Palmitoylation of HTT (**B**), SNAP25 (**C**), PSD95 (**D**) and GAD65 (**E**) was measured using
the ABE assay. The palmitoylation signal (HAM+) is shown in the top panels for each
protein, and the total protein immunoprecipitated is presented in the panels below. The
negative control (HAM-) is displayed on the right of each set of blots. Overlay of both
channel (palmitoylated protein in red and total protein in green) are displayed in the
bottom panels. In the HTT blots (**B**), the lower band corresponds to mouse Htt, and the
upper band to mHTT. The palmitoylation levels were calculated as a ratio of palmitoylation
over total protein signal.

<u>Statistical analysis</u>: (**B**) 2-way ANOVA: interaction, $p=0.059$; allele, $p<0.0001$, ASO
treatment, $p=0.48$. Stars indicate Tukey's multiple comparisons test results between PBS
and ASO groups. Student's T-test: (**A**, **C**, **D**, **E**) results are indicated by p-values or stars
on the graphs.

A COS-7 cells exogenously expressing wt and mHTT

B WT and HD cortico-striatal primary neurons

C Control and HD patient-derived lymphoblasts

1075

<u>Figure 5</u> - **Inhibition of APTs with Palmostatin B increases palmitoylation level of**
HTT in COS-7 cells, YAC128 primary neurons and HD-patient derived lymphoblasts.

(**A**) COS-7 cells were transfected with the N-terminal fragments HTT_{1-548} (N548), HTT_{1-}
$_{1212}$ (N1212) and full length wt (15Q) or mutant (128Q) HTT_{1-3144} constructs. Cells were
treated with DMSO or 15 µM PalmB for 6h while concurrently labelled with 100 µM alkyne-
palmitate. Bio-orthogonal click chemistry of alkyne-palmitate was performed on lysates
immunoprecipitated with anti-HTT (MAB2166) antibodies. Palmitoylated and total HTT
were detected by immunoblots using streptavidin and anti-HTT (HD46) antibodies,
respectively. Overlay of both channel (palm in red and total protein in green) are displayed
in the bottom panels. Palmitoylation was calculated as a ratio of palmitoylation signal over
total HTT protein and are expressed relative to wtHTT palmitoylation in the DMSO control
group.

(**B**) YAC18 and YAC128 cortico-striatal primary neurons were treated at DIV10 with
DMSO or 15 µM PalmB for 6h. Cell lysates were subjected to ABE assay to measure
palmitoylated HTT. HTT was immunoprecipitated with anti-HTT (MAB2166) antibodies
and detected with D7F7 for Htt (mouse) or HTT (human) and streptavidin for palmitoylated
protein. HTT palmitoylation level was calculated as a ratio of palmitoylation signal over
total HTT protein and expressed relative to mHTT palmitoylation in the DMSO control
group.

(**C**) Control and HD-patient derived lymphoblastoid lines were treated at with DMSO or
10 µM PalmB for 4h, and lysates were subjected to the ABE assay. HTT was
immunoprecipitated with anti-HTT (MAB2166) antibodies and detected with D7F7 for total
HTT and streptavidin for palmitoylated HTT. HTT palmitoylation level was calculated as a
ratio of palmitoylation signal over total HTT protein and expressed relative to wtHTT
palmitoylation level in the DMSO treated control group.

<u>Statistical analysis</u>: 2-way ANOVA: (**A**) N548, interaction, p=0.0042; HD mutation,
p<0.0001; PalmB, <0.0001. N1212, interaction, p=0.0075; HD mutation, p<0.0001;
PalmB, p=0.0007. Full length HTT, interaction, p=0.65; HD mutation, p=0.0002; PalmB,
p<0.0001. (**B**) Interaction, p=0.86; HD mutation, p<0.0001; PalmB, p=0.0320. (**C**)
Interaction, p=0.90; HD mutation, p=0.43; PalmB, p=0.0071. Stars indicate Bonferroni's
multiple comparisons test results and hashtag t-test results.

1107

Figure 6 - Inhibition of APTs with Palmostatin B reduces mHTT insoluble inclusions
in COS-7 cells and mHTT-induced cytotoxicity in COS-7 cells and YAC128 cortico-
striatal primary neurons.

COS-7 cells were transfected at t=0h with 138Q-HTT$_{1-588}$-YFP (138Q-N588-YFP)
construct, treated with 15 µM PalmB at t=24h and 48h post-transfection and harvested at
t=72h.

(**A**) Cell lysates were subjected to ABE assay to measure palmitoylated HTT. HTT was
immunoprecipitated with anti-GFP antibodies and detected with MAB2166 for HTT and
streptavidin for palmitoylated protein. HTT palmitoylation level was calculated as a ratio
of palmitoylation signal over total HTT protein and expressed relative to palmitoylation in
the DMSO control group.

(**B-C**) Insoluble mHTT inclusions were monitored by filter retardation assay and
immunodetection with anti-HTT (MAB2166) or GFP antibodies. The amount of
aggregates retained on the filter is expressed relative to (**B**) 17Q-N588-YFP or (**C**) to the
DMSO treated group.

(**D**) Expression of full length 128Q-HTT in COS-7 cells increased cytotoxicity compared
to 15Q-HTT expression. COS-7 cells seeded in 96-well pates were transfected with WT
(15Q) or mutant (128Q) full length HTT for 24h. LDH activity in conditioned media was
measured using the CyQUANT cell proliferation assay. Cytotoxicity (%) is expressed
relative to untreated and non-transfected cells.

(**E**) Cytotoxicity in COS-7 cells transfected for 24h with full length 128Q-HTT and treated
with DMSO or PalmB (InSolution at 10, 15, 20, 30 and 100 µM) for 6 or 24h was assessed
using the LDH activity assay (n=3). Cytotoxicity (%) is expressed relative to untreated and
non-transfected cells. Negative cytotoxicity values correspond to a protective effect.

(**F**) Cytotoxicity in YAC128 cortico-striatal primary neurons at DIV15 treated with DMSO
or PalmB (InSolution at 5, 10, 15, 20, 50 and 100 µM) for 6 or 24h, using the LDH activity
assay (n=4). Cytotoxicity (%) is expressed relatively to YAC128 neurons treated with
DMSO. Negative cytotoxicity corresponds to a protective effect. Negative cytotoxicity
values correspond to a protective effect.

<u>Statistical analysis</u>: Student's T-test (**A**) p=0.029, (**B**) p<0.0001, (**C**) MAB2166, p=0.038
and GFP, p=0.0065. (**D**) p=0.019. 2-way ANOVA: (**E**) Interaction, p=0.90, PalmB effect,
p<0.0001, Treatment duration, p=0.0033; (**F**) Interaction, p=0.97; PalmB, p=0.0014;
Treatment duration, p=0.29. Stars indicate Dunnett's test results compared to respective
DMSO control.

Table 1 – List of reagents and chemicals used and application

1143

	Reagent	Company	Catalogue #
Cell Culture Medium	DMEM without methionine and cystine	ThermoFisher Scientific	21013024
	DMEM/F12	ThermoFisher Scientific	11320033
	Dulbecco's Modified Eagle (DMEM)	ThermoFisher Scientific	11960044
	Neurobasal (NB)	ThermoFisher Scientific	21103049
	RPMI 1640	ThermoFisher Scientific	11875093
	mTeSR™1	STEMCELL Technologies	05850
Cell Culture Reagents	APT1 Inhibitor, palmostatin B	MilliporeSigma	178501
	B27 supplement	ThermoFisher Scientific	17504044
	Corning™ Matrigel™ hESC-Qualified Matrix	Fisher Scientific	354277
	Charcoal stripped FBS	ThermoFisher Scientific	12676029
	Dimethyl sulfoxide (DMSO)	MilliporeSigma	D2650
	DNase I, RNase-free	ThermoFisher Scientific	EN0525
	Fetal Bovine Serum	ThermoFisher Scientific	12483020
	GlutaMAX	ThermoFisher Scientific	35050061
	Hank's balanced salt solution (HBSS) (10X)	ThermoFisher Scientific	14060040
	InSolution™ APT1 Inhibitor, palmostatin B	MilliporeSigma	508738
	Nunc™ EasYFlask™ 25cm^2	ThermoFisher Scientific	156367
	Penicillin-Streptomycin (10,000 U/mL)	ThermoFisher Scientific	15140122
	Phosphate-buffered saline (PBS) (10X)	ThermoFisher Scientific	70011044
	Poly-D-lysine hydrobromide	MilliporeSigma	P7886
	ReLeSR™	STEMCELL Technologies	05872
	Sodium pyruvate	ThermoFisher Scientific	11360070
	Trypsin-EDTA (0.25%), phenol red	ThermoFisher Scientific	25200056
	Trypsin-EDTA (0.05%), phenol red	ThermoFisher Scientific	25300
	X-tremeGENE™ 9 DNA Reagent	MilliporeSigma	6365809001
	6-well, Nunclon delta surface plates	ThermoFisher Scientific	140685
	96-Well, Nunclon Delta-Treated plate	ThermoFisher Scientific	167425
Palmitoylation Assays	Biotin Azide	ThermoFisher Scientific	B10184
	Click Tag™ Palmitic Acid Alkyne	Cayman Chemical	13266
	EZ-Link BMCC-Biotin	ThermoFisher Scientific	21900
	Hydroxylamine hydrochloride	MilliporeSigma	255580
	N-ethylmaleimide (NEM)	MilliporeSigma	E3876
	Triton X-100	MilliporeSigma	T9284
	Tris(benzyltriazolylmethyl)amine (TBTA)	MilliporeSigma	678937
	Tris-carboxyethylphosphine (TCEP)	MilliporeSigma	C4706
	CYQUANT LDH cytotoxicity assay kit	ThermoFisher Scientific	C20303

Benzonase® Nuclease	MilliporeSigma	E1014-25KU
Cellulose acetate (OE66) membrane	MilliporeSigma	WHA10404180
Protran WB membranes, nitrocellulose	MilliporeSigma	GE10600000
Dithiothreitol (DTT)	MilliporeSigma	3483-12-3
Dynabeads™ Protein G	ThermoFisher Scientific	10003D
Gel Cassettes, mini, 1.5 mm	ThermoFisher Scientific	NC2015
NuPAGE LDS Sample Buffer (4X)	ThermoFisher Scientific	NP0007
Invitrogen UltraPure TEMED	ThermoFisher Scientific	15524010
NuPAGE Novex 3-8% Tris-Acetate gels	ThermoFisher Scientific	EA0375
Protein Assay Reagent A	Bio-Rad	5000113
Protein Assay Reagent B	Bio-Rad	5000114
PureProteome magnetic beads	MilliporeSigma	LSKMAGAG02
Roche complete Protease Inhibitor Cocktail	MilliporeSigma	11836145001
Sodium dodecyl sulfate (SDS)	ThermoFisher Scientific	BP166-500
2% Bis Solution	Bio-Rad	1610142
30% Acrylamide/Bis Solution, 29:1	Bio-Rad	1610156
40% Acrylamide Solution	Bio-Rad	1610140

Immunoblotting

1144

   Table 2 – List of antibodies used and application

Target (clone)	Species	Type	Supplier	Catalogue AB Registry Reference	Application	Dilution	Volume[*]
HTT (1HU-4C8)	Mouse	Monoclonal	Sigma-Aldrich	MAB2166 AB_2123255	WB	1:1,000	-
					IP	-	2 µL
HTT (D7F7)	Rabbit	Monoclonal	Cell Signaling Technology	5656 AB_10827977	WB	1:1,000	-
HTT (BKP1/HD46)	Rabbit	Polyclonal	in-house	(Kalchman et al., 1997)	IP	-	2 µL
					WB	1:400	-
HTT	Rabbit	Polyclonal	Sigma-Aldrich	H7540 AB_1840946	WB	1:1,000	-
					IP	-	2 µL
Streptavidin	-	Conjugated to AF-680	Invitrogen	S-32358	WB	1:5,000	-
PSD95 (7E3-1B8)	Mouse	Monoclonal	Invitrogen	MA1-046 AB_2092361	WB	1:1,000	-
PSD95	Rabbit	Polyclonal	in-house	-	IP	-	5 µL
GAD65	Rabbit	Polyclonal	Sigma-Aldrich	ABN101	WB	1:1,000	-
	Mouse	Monoclonal	DSHB	GAD-6 AB_2314499	IP	-	3 µL
SNAP25	Rabbit	Polyclonal	Synaptic Systems	111-002 AB_887790	WB	1:1,000	-
SNAP25 (SMI 81)	Mouse	Monoclonal	BioLegend	836303 AB_2715864	IP	-	3 µL
GFP	Rabbit	Polyclonal	Eusera	EU2	WB	1:5,000	-
					IP	-	0.75 µL
Mouse IgM (µ chain specific)	goat	-	Rockland Immuno-chemicals	92632210	WB (secondary)	1:5,000	-
Rabbit IgG	goat	-		926-32211	WB (secondary)	1:5,000	-

1146   [*]Volume of antibodies used for 20 µL of Dynabeads and 200 µg – 1 mg proteins

1147 **Table 3 – HD mouse lines used for palmitoylation assays**

Mouse model	Line	Reference	Promoter	Transgene product	CAG repeat length	Endogenous product
YAC128	53	(Slow et al., 2003)	Human HTT promoter and RE (24 kb upstream, 117 kb downstream)	full length human mHTT	128	2 copies of 7Q Htt
YAC18	60			2 copies of full length human wtHTT	18	2 copies of 7QnHtt
YAC128	55			2 copies of full length human mHTT	128	2 copies of 7Q Htt
BACHD	LA	(Gray et al., 2008)	Human HTT promoter and RE (20 kb upstream, 50 kb downstream)	full length human mHTT	97	2 copies of 7Q Htt
Hu18/18	KLAY	(Southwell et al., 2013)	Human HTT promoter and RE (20 kb upstream, 50 kb downstream)	full length human wtHTT	18	-
Hu97/18	KLAY			full length human WT and mHTT	97/18	-
Hu21/21	KT53	(Southwell et al., 2017)	Human HTT promoter and RE (24 kb upstream, 117 kb downstream)	full length human wtHTT	21	-
Hu128/21	KT53			full length human wt and mHTT	128/21	-

1148

1149

1150 **Table 4 – Lymphoblastoid cell lines from Coriell Biorepositories**

Catalog ID	Haplotype U/L	Status	Race	Age (years)	Sex	Family	CAG Lower	CAG Upper
GM07029		Control		47	Male	N/a	15	17
GM12716		Control		N/a	Male	1358	15	19
GM02160		Control	Caucasian	21	Male	109	16	17
GM02170		Control		22	Female	111	17	17
GM03863		Control		24	Female	268	17	17
GM03867	N/a	HD		26	Female	268	21	43
GM04023	C/A2	HD		28	Female	587	18	44
GM02176	A1/C	HD		26	Male	110	18	44
GM02150	A2/C	HD	Caucasian	26	Female	108	18	45
GM03620	A1/C	HD		29	Female	110	18	59
GM04724	A3/C	HD		19	Female	691	15	67
GM13509	A3/C	HD		N/a	Female	2706	15	66
GM04282	A3/A4	HD		20	Male	690	17	71

1151

1152

1153

1154   **Table 5 – Isogenic HD patient-derived hiPSCs lines**

Cell type	Line	Reference	CAG repeat length
iPSCs	Parental control line (H9)	(Ooi et al., 2019)	30
	IsoHD lines		45
			65
			81
	Parental HD line (CAG180)	(Xu et al., 2017)	180
	Corrected line (HD-C#2)		18

1155