Identifying key federal, state, and private lands strategies for achieving 30x30 in the US

Lindsay Dreiss[1*], Jacob Malcom[1]

[1]Center for Conservation Innovation, Defenders of Wildlife, Washington, DC 20036, USA

* Correspondence:
Corresponding Author
lrosa@defenders.org

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial

relationships that could be construed as a potential conflict of interest.

Acknowledgement

We thank T Niederman for her thoughtful review of this manuscript. We also thank USGS for

making PADUS, C. Soto-Navarro for making above- and below-ground carbon, and NatureServe and

Esri for making the MoBI layers available to streamline general analyses.

Data Availability

All data used in analyses are publicly available.

Abstract

Addressing the current biodiversity crisis will require transformative changes to social, political, and

economic structures. One science-based recommendation is protecting 30% of the Earth's terrestrial

and marine systems by 2030, "30x30". Here we analyze the current spatial patterns of imperiled

species biodiversity and carbon stores in the U.S. relative to protected areas to help conservationists

and decision makers understand the starting point on the path to achieving 30x30. Multi-scale

analyses demonstrate that 30x30 is numerically achievable nationally, but high spatial heterogeneity

highlights the need for tailored approaches from a mix of authorities at federal, regional, and state

scales. Critically, current land protections rarely overlap with areas essential for conserving imperiled

species biodiversity and mitigating climate change. We discuss this baseline relative to key policy

considerations for making practical, substantive progress toward the goal.

Keywords: biodiversity conservation, climate mitigation, protected areas, endangered species

Introduction

According to the international conservation community, goals for conserving biodiversity cannot be

met given current trajectories of environmental degradation and without transformative changes

across economic, social, political, and technological spheres (IPBES Secretariat 2019). Restoration

and maintenance of quality habitat through a more extensive global protected areas (PAs) network

(see Aichi Target 11; SCBD 2010) is considered essential for conservation. Yet the global

community has fallen short of 2020 targets for PA coverage at a time when threats to biodiversity –

foremost, habitat conversion – are at an all-time high (Powers and Jetz 2019).

Recognizing the critical role that PAs have in conserving biodiversity and mitigating climate change

impacts has led to increased interest in adopting new national and international conservation targets.

The Global Deal for Nature, a science-driven plan to sustain biodiversity and address climate change,

calls for at least 30% of Earth to be formally protected by 2030 ("30x30"; Dinerstein *et al*. 2019).

The Convention on Biological Diversity has drafted a post-2020 global framework that includes

30x30 as a steppingstone toward a 2050 Vision for Biodiversity (SCBD 2020). In the United States,

the number of proposed policy measures aligning with a 30x30 framework is on the rise. As of early

2021, this includes a federal executive order and a California state executive order (Exec Order No

14008 2021, CA Executive Order N-82-20). These efforts provide an opportunity to integrate

biodiversity and climate agendas and promote land protections that can maximize biodiversity

conservation and minimize carbon loss at multiple scales.

Past research on setting PA priorities for biodiversity (Kulberg *et al*. 2019; Belote *et al*. 2017;

Jenkins *et al*. 2015) and natural climate strategies (Soto-Navarro *et al*. 2020; Stralberg *et al*. 2020)

provide a useful foundation for addressing this need, but do not align with the policy tools and units

at which federal and state decision-makers govern. To achieve 30x30 in a way that most benefits

biodiversity and climate, policy makers need guidance on how to operationalize these targets.

Here we synthesize biodiversity and ecosystem carbon data with policy-relevant land protections at

multiple scales to provide a baseline assessment for charting a path to 30x30 in the U.S. We use the

Protected Areas Database of the U.S. (PADUS; U.S. Geological Survey 2020) to spatially define

areas that are currently managed for biodiversity conservation (GAP 1 & 2) and that may be in need

of additional protections (GAP 3 & 4) (Panel 1). We compare GAP classified lands with lands rich in

imperiled species and carbon. These results will help conservationists and decision-makers plan and

take critical next steps to operationalize 30x30-in doing so, addressing some of the greatest

conservation challenges faced by the United States and the world. some of the greatest conservation

challenges faced by the United States and the world.

Methods

Datasets

Terrestrial imperiled species richness and rarity-weighted richness from the Map of Biodiversity

Importance project database (NatureServe 2020) are based on habitat suitability models for 2,216

species and 11 taxa. Species richness is useful for understanding where ranges of the most species

overlap a single location, but could be a poor indicator of diversity because it does not account for

complementarity among sites. Alternatively, rarity-weighted richness – where species are assigned a

value inversely proportional to the size of their range - may be a better method for maximizing the

representation of species (Albuquerque and Gregory 2017). Therefore, we take into account the

number of species, proportional abundance of species presence, and the occurrence of rare species of

conservation priority. Total ecosystem carbon is based on a map of global above- and below-ground

carbon stored in biomass and soil in tonnes C per hectare (Soto-Navarro *et al*. 2020).

Data on protected areas are from the PADUS v2.1 database (U.S. Geological Survey 2020). The layer

was flattened prior to spatial overlay analysis to avoid overlaps in land units. This was done so in a

way to give preference to the lowest GAP code present in any single location (i.e., greatest

protection). U.S. terrestrial boundaries reflect all states and territories from the U.S. Geological

Survey.

<u>Analyses</u>

We used spatial overlay analysis to describe the extent to which the PA network covers U.S. lands

and seas as well as areas of high imperiled species biodiversity and ecosystem carbon (see SI for

marine results). Areas considered hotspots of biodiversity or carbon richness were in the 90th

percentile of the distribution of values for the respective metric. We used the associated Gap Analysis

Program (GAP) codes, which are specific to the management intent to conserve biodiversity. GAP 1

and 2 areas are managed in ways typically consistent with conservation. GAP 3 areas are governed

under multiple-use mandates and GAP 4 areas lack any conservation mandates (see Panel 1).

Additionally, we summarized statistics by lands management (e.g., manager type and name attribute)

and state or territory. We used ArcPro v. 2.7.1 (ESRI, USA) to produce maps and run analyses. Maps

use the Albers Equal Area Conic, Alaska Albers, and Old Hawaiian UTM Zone 4 projections.

Results

Twelve percent of lands within the U.S. and its territories are generally managed consistently with

biodiversity conservation goals (i.e., GAP 1 and 2). Up to 29.8% of U.S. lands and territories are

managed for either biodiversity conservation or multiple uses (i.e., GAP 1, 2, and 3). This leaves a

large majority of the U.S. lacking any known protections from land conversion (70.2%; i.e., GAP 4).

Areas managed for biodiversity (85.7% of GAP 1 and 2 areas) and for multiple uses (85.6% of GAP

3 lands; Figure 1a) fall largely on federally managed lands. Protecting 30% of lands could nearly be

achieved at the national scale if new conservation-based mandates were applied to all federally

managed GAP 1-3 areas (27.7%).

At the state-level, Alaska is the only state to have at least 30% of its territory managed for

conservation (Figure 1b). Twelve western states - California, Nevada, Washington, Idaho, Oregon,

Hawaii, Utah, Wyoming, Arizona, Colorado, Montana and New Mexico - would achieve the 30%

numerical target if GAP 3 areas were similarly managed. Generally, federal agencies manage the

majority of GAP 3 lands in these states. In contrast, the 11 states with greater state-level authority are

located in the Northwest and Midwest (Table S1).

Imperiled species biodiversity hotspots and GAP 1 or 2 areas rarely overlap, with only 7% of

hotspots covered by GAP 1 and 2 lands (Figure 2). Similarly, 12.8% of highest potential areas fall

within GAP 1 and 2 lands. While including GAP 3 lands significantly increases the coverage for

biodiversity hotspots (from 7% to 20%; Figure 2), 80% of the most diverse areas would still remain

unprotected because they fall on GAP 4 or otherwise unprotected private lands. Nearly two thirds of

the top carbon-rich areas fall in GAP 4 areas. While rarity-weighted richness exhibits similar broad

spatial patterns to raw species richness (Figures 1 and 2), there is significantly greater overlap with

key GAP lands: 32.6% of rarity hotspots are covered by GAP 1 and 2 areas and an additional 20.5%

by GAP 3 lands.

As expected, the goals of protecting areas of high biodiversity and areas of high carbon mitigation

potential are not completely decoupled: 22.3% of the top quantile of species richness locations

(25.6% of rarity-weighted richness hotspots) are also very carbon rich. These percentages nearly

double when assessing overlap with the top 30% of carbon rich areas. Unfortunately, few areas that

meet both needs are managed in ways to help sustain them: 17.4% of areas that serve as hotspots for

biodiversity and carbon fall on GAP 1 or 2 lands (<0.01% of CONUS).

A 30x30 Typology

To guide operationalization of the 30x30 framework, we use GAP categories as proxies for policy

options and biodiversity and ecosystem carbon as representative of the underlying goals of 30x30

(Figure 2 legend):

Well-sited: Areas with long-term conservation management mandates (GAP 1-2 coverage)

and high biodiversity and/or ecosystem carbon, where areas are effectively placed to achieve

greater biodiversity conservation and climate mitigation. Priority actions for well-sited areas

include maintaining existing protections and expanding protections outward in a way that

ensures landscape connectivity.

High Priority: Areas of overlap between weaker or short-term mandates (GAP 3-4) and

greater biodiversity and/or ecosystem carbon, where new PAs or more protective policies

would do the most to protect biodiversity and mitigate climate. These are the areas of greatest

opportunity, where efforts to expand PAs would have especially high returns on investment.

Well-protected: Areas with long-term conservation mandates but relatively low local

imperiled species diversity and/or ecosystem carbon potential, indicating lower return on

conservation goals than other areas might provide. Despite their lower biodiversity and

carbon stocks, these areas may serve as anchors for expanding protections to key adjacent

areas.

Limited Value: Areas with weaker or short-term mandates and low levels of biodiversity

and/or ecosystem carbon potential, indicating a low return on protections and an advantage to

site PAs elsewhere. These are the lowest priority areas for 30x30. However, it is important to

note that basing decisions solely on current habitat and species ranges may discount the

necessity for future habitat recovery and restoration given climate change (Lawler *et al*.

2013).

Discussion

While the basic numerical accomplishment of protecting 30x30 is feasible at the national scale given

the current extent of the PAs network, prioritizing biodiversity protection and climate change

mitigation presents challenges and opportunities. In addition, high spatial variability in the

distribution of PA designations and will require tailored approaches across regions (Figure 1).

An option for more rapidly reaching 30% includes establishing additional protections on GAP 3

lands. For example, nearly 30% can be achieved at the national level if regulatory changes to GAP 3

areas emphasized biodiversity protection over other uses. Because the majority of the PA network

(GAP 1-3) is managed by federal agencies, action at the federal level may be essential to reach

numerical goals. Land management laws like the National Forest Management Act and the Federal

Lands Planning and Management Act afford species and habitats protections, but the effectiveness of

protections may vary by ownership (Eichenwald *et al*. 2020). Moreover, federal agencies can provide

necessary leadership and coordination across jurisdictions to better ensure representation of all

natural ecosystems (Dinerstein *et al*. 2019).

Importantly, focusing on key federal lands alone would ignore the substantive goals of 30x30: 80%

of biodiversity hotspots would still lack significant place-based protections if GAP 3 federal lands

were converted. This result highlights the alarming reality that focusing strictly on the numerical goal

of 30x30 could lead to outcomes contrary to intent, with new PAs being established in areas with low

biodiversity or carbon mitigation potential (Barnes *et al.* 2018). Regions with few public lands will

face trade-offs between siting new areas based on biodiversity need or opportunity (e.g., isolated and

sparsely populated; Baldi *et al.* 2017). GAP 4 areas are, by far, the most extensive, but would require

more effort and investment from decision-makers to acquire land and/or establish biodiversity

protections as priorities. Current federal conservation incentive programs, such as Farm Bill

programs and those administered by the U.S. Fish and Wildlife Service are inadequate to address the

need. As such, there must be significant efforts to advance conservation on private lands in key parts

of the country.

In addition to federal and private lands, understanding state variation in biodiversity and protections

is vital as important legal, social, and policy mechanisms operate at this level. We note twelve

western states would achieve the 30% numerical target if GAP 3 mandates were strengthened.

However, many areas of these states harbor few biodiversity or carbon-rich hotspots. In contrast, the

11 states where the state manages the majority of GAP 3 lands have higher biodiversity on average,

but GAP 3 areas may not overlap with hotspots or be enough to significantly lessen the disparities in

current PA coverage and the 30% target. State wildlife conservation programs employ state wildlife

action plans that have the potential to advance conservation (e.g., Michalak and Lerner 2008), but are

woefully underfunded (H. Res. 3742 2019). New state-level programs for public and private lands

conservation can complement federal programs and create a more complete, multi-level solution.

Finally, we acknowledge the strong need for several additional considerations not accounted for in

this work. In addition to the primary focus on biodiversity and climate, pursuing 30x30 will require

addressing issues related to economic, political, and social constraints. For example, many High

Priority areas for siting new protections also tend to be GAP 4 regions with higher human

disturbance. This elevates the importance of restoration efforts (see, e.g., Strassburg *et al.* 2020) and

relative habitat condition could be integrated into analyses to avoid conserving areas with negligible

conservation benefits. If properly planned, PAs in these areas may also provide opportunities for

improving human health, well-being, and equitable access to nature. Goals to ensure a healthy

environment for all communities have long been ignored or discounted in protected areas

designations (Wood *et al*. 2018), in part because these topics are not well studied (Ussery *et al*.

2016). Further research and planning are essential to ensuring access to quality nature for all.

This analysis is meant as a tool to aid decision-makers and not as a fully comprehensive plan. Our

analyses are national in scope and intended to identify broad patterns to frame the national

discussion; as such, local and domain-specific details are likely to vary. Additionally, focusing on

values at the national scale means that entire ecosystems important to representing local species

assemblages and key ecosystem services are not included as high priority. A stratified approach may

ensure that all native ecosystems are represented in the expanding PA network. Second, we are using

models of current imperiled species distributions to infer the general patterns of protections, some of

which may shift with global climate change (see Elsen *et al*. 2020). Future local, regional and

continental scale analyses can help inform which areas need long-term protections. Finally, the GAP

classification definitions may not account for substantive protections observed on the ground. For

example, Department of Defense installations represent 20 million acres of GAP 4 land (some with

high imperiled species diversity; Stein *et al*. 2008) and have Integrated Natural Resource

Management Plans that address some biodiversity concerns. There is also a great need for

incorporation of Tribal knowledge as Tribal areas have some of the lowest rates of habitat

modification, yet much of the over 56 million acres held in trust by the Bureau of Indian Affairs are

not well documented in PADUS (Vincent *et al*. 2017). These and similar situations highlight the

nuances of inferring protection status.

Achieving 30x30 to help protect biodiversity and address the climate crisis in the U.S. is feasible but

will require partnerships with nongovernmental landowners and across levels of government. Our

analysis recognizes that the approaches and policy tools for doing so will vary considerably

throughout the country. The key to operationalizing 30x30 will be planning beyond the numerical

target for a protected areas network that can be established in a way that ensures a long-term

commitment to biodiversity and climate. By doing so, the U.S. can continue to lead the way globally

in protecting nature for its own sake and for our health and well-being.

References

Albuquerque FS and Gregory A. 2017. The geography of hotspots of rarity-weighted richness of

birds and their coverage by Natura 2000. *PLoS One* **12**: e0174179.Baldi G, Texeira M, Martin OA,

et al. 2017. Opportunities drive the global distribution of protected areas. *PeerJ* **5:** e2989.

Barnes MD, Glew ., Wyborn C, *et al*. 2018. Prevent perverse outcomes from global protected area

policy. *Nat Ecol Evol* **2**: 759–762.

Belote R T, Dietz MS, Jenkins CN, *et al*. 2017. Wild, connected, and diverse: building a more

resilient system of protected areas. *Ecol Appl* **27**: 1050-1056.

Dietz MS, Belote RT, Gage J, *et al*. 2020. An assessment of vulnerable wildlife, their habitats, and

protected areas in the contiguous United States. *Biol Conserv* **248**: 108646.

Dinerstein E, Vynne C, Sala E, *et al*. 2019. A global deal for nature: guiding principles, milestones,

and targets. *Sci Adv* **5**: EAAW2869.

Elsen PR, Monahan WB, Dougherty ER, Merenlender AM. 2020. Keeping pace with climate change

in global terrestrial protected areas. *Sci Adv* **6**: eaay0814.

House Resolution 3742. 2019. Recovering America's Wildlife Act.

https://www.congress.gov/bill/116th-congress/house-bill/3742 Viewed February 21, 2021.

Lawler JJ, Ruesch AS, Olden JD, *et al*. 2013. Projected climate-driven faunal movement routes. *Ecol*

Letters **16**: 1014-1022.

Michalak J and Lerner J. 2008. Linking conservation and land use planning: using the State Wildlife

Action Plans to protect wildlife from urbanization. In: Transportation Land Use, Planning, and

Air Quality. Presented at the Transportation Land Use, Planning, and Air Quality Congress 2007,

American Society of Civil Engineers, Orlando, Florida, United States, pp. 32–40.

Jenkins CN, Van Houtan KS, Pimm SL, *et al*. 2015. US protected lands mismatch biodiversity

priorities. *PNAS* **112**: 5081–5086.

Kulberg P, Di Minin E, Moilane A. 2019. Using key biodiversity areas to guide effective expansion

of the global protected areas network. *Glob Ecol Conserv* **20**: e00768.

Powers RP and Jetz W. 2019. Global habitat loss and extinction risk of terrestrial vertebrates under

future land-use-change scenarios. *Nat Clim Chang* **9:** 323–329.

Richards R and Lee-Ashley M. 2020. The Race for Nature. Available at: https://ampr.gs/3f0sBk0

Viewed July 4, 2020.

Secretariat of the Convention on Biological Diversity. 2020. Zero draft of the Post-2020 Global

Biodiversity Framework.

https://www.cbd.int/doc/c/efb0/1f84/a892b98d2982a829962b6371/wg2020-02-03-en.pdf Viewed

February 19, 2021.

Secretariat of the Convention on Biological Diversity. 2010. COP-10 Decision X/2. Secretariat of the

convention on biological diversity. Viewed February 19, 2021.

Soto-Navarro C, Ravilious C, Arnell A, *et al*. 2020. Mapping co-benefits for carbon storage and

biodiversity to inform conservation policy and action. *Phil Trans Royal Soc B* doi:

10.1098/rstb.2019.0128

Stein BA, Scott C and Benton N. 2008. Federal lands and endangered species: the role of military

and other federal lands in sustaining biodiversity. *BioSci* **58**: 339-347.

Stralberg D, Carroll C and Nielsen SE. 2020. Toward a climate-informed North American protected

areas network: Incorporating climate-change refugia and corridors in conservation planning.

Conserv Lett **13:** ee12712.

Strassburg, B. B. N., Iribarrem, A., Beyer, H. L., Cordeiro, C. L., Crouzeilles, R., Jakovac, C. C., et

al. 2020. Global priority areas for ecosystem restoration. Nature, 1–6. doi:10.1038/s41586-020-

2784-9.

U.S. Geological Survey (USGS) Gap Analysis Project (GAP). 2020. Protected Areas Database of the

United States (PAD-US): U.S. Geological Survey data release. doi: 10.5066/P955KPLE

Ussery EN, Yngve L, Merriam D, *et al*. 2016. The national public health tracking network access to

parks indicator: a national county-level measure of park proximity. *J Park Recreat Admi* **34**: 52–

63.

Vincent C, Hanson LA, and Argueta C. 2017. Federal Land Ownership: Overview and Data.

https://fas.org/sgp/crs/misc/R42346.pdf. Viewed February 19, 2021.

Wood E, Harsant A, Dallimer M, *et al*. 2018. Not All Green Space Is Created Equal: Biodiversity

Predicts Psychological Restorative Benefits From Urban Green Space. *Front Psychol* **9:** doi:

10.3389/fpsyg.2018.02320

313   FIGURES

Panel 1. PADUS and GAP

We use the USGS Gap Analysis Project's GAP status code from the Protected Areas Database of the U.S. (PADUS) to characterize current protections and future possibilities. GAP codes are specific to the management intent to conserve biodiversity and largely coincide with legal requirements for protections: GAP 1 and 2 areas are managed in ways typically consistent with conservation (e.g., national wilderness and national wildlife refuges); GAP 3 areas are governed under multiple-use mandates (e.g., wildlife, forestry and mining); and GAP 4 areas lack any conservation mandates).

Approximately 39% of terrestrial and over 73% of marine territories are included in PADUS, but only GAP 1 and 2 are typically considered protected (e.g., Richards 2018). However, GAP 3 may have particular potential to advance biodiversity and climate protections more quickly through administrative mechanisms (Figure 1).

GAP 1	An area having permanent protection from conversion of natural land cover and mandated management plan to maintain a natural state.
GAP 2	Permanent protections and a mandated plan maintain a primarily natural state, but management practices may degrade quality.
GAP 3	Permanent protections in majority, but some areas subject to extractive uses. Protections for federally listed species exist throughout.
GAP 4	The area generally allows conversion to unnatural land cover throughout or management intent is unknown.

Management Level	Example Protections
Federal	National parks, national forests, national conservation lands, national marine sanctuaries, national recreation areas, national estuarine research reserves, national trails, national wild and scenic rivers, national wilderness areas and national wildlife refuges
State	Parks, wilderness, wildlife management areas
Local	Tribal, regional, county or city parks and public lands
Private	Easements: agricultural, conservation, forest stewardship, cultural area, ranch, recreation or education

The GAP status map highlights the high spatial variation in areas currently managed for biodiversity conservation (GAP 1 and 2) in the U.S. and the reality that unprotected areas (GAP 3 and 4) dominate most of the country.

Figure Legends:

Panel 1

Figure 1. Proportion of U.S. Territory lands in PADUS by GAP status code and manager type,

showing that (A) federal agencies manage much of the protected areas network and that (B) most

state have few protections for biodiversity as they fall well-short of achieving 30x30 given existing

and potential (GAP 1-3) protections.

Figure 2. Combining protected area coverage to locations of A) imperiled species richness, B) rarity-

weighted richness and C) ecosystem carbon (tonnes C per hectare) show significant areas of

mismatch that need to be addressed. The blue (y-axis) component of the bivariate color ramp

signifies protections under GAP categories 1-4 while the yellow (x-axis) component signifies the

corresponding 30x30 variable based on quantile intervals where the top 10% of values are in yellows

(imperiled species richness from the Map of Biodiversity Importance program [NatureServe 2020] or

ecosystem carbon from Soto-Navarro et al. 2020). Resolution is 1km2. Bar charts represent the

overlaps of different percentiles of biodiversity or carbon values with GAP categories (green color

ramp). Biodiversity and carbon hotspots were considered as locations with values in the top 90[th]

percentile of the distribution.