Title: Purkinje Cell Activity in Medial and Lateral Cerebellum During Suppression of Voluntary Eye Movements in Rhesus Macaques

Abbreviated title: Cerebellar activity during antisaccades

Eric Avila[1,2#], Nico A. Flierman[1,2#], Peter J. Holland[3], Pieter R. Roelfsema[1,4,5], Maarten A. Frens[2], Aleksandra Badura[1,2*] and Chris I. De Zeeuw[1,2*]

[1] Netherlands Institute for Neuroscience, Amsterdam, 1105 BA, The Netherlands

[2] Department of Neuroscience, Erasmus MC, Rotterdam, 3015 CN, The Netherlands

[3] School of Psychology, University of Birmingham, Birmingham, B15 2TT, UK

[4] Department of Integrative Neurophysiology, VU University, Amsterdam, 1081 HV, The Netherlands

[5] Department of Psychiatry, Academic Medical Centre, Amsterdam, 1105 AZ, The Netherlands

[#] These authors contributed equally to this work.

EA current address: Center for Neural Science, New York University, New York, 10003, USA.

[*]**Correspondence**:

Aleksandra Badura (a.badura@erasmusmc.nl)

Department of Neuroscience, Erasmus MC

Wytemaweg 80, 3015 CN Rotterdam

tel: 0031-(0)10 7043589

Chris I De Zeeuw (c.de.zeeuw@nin.knaw.nl)

Netherlands Institute for Neuroscience

Meibergdreef 47,1105 BA, Amsterdam

tel: 0031-(0)20- 5665500

Number of pages: 40

Number of figures: 5

Abstract: 246 words.

Introduction: 626 words.

Discussion: 1440 words.

Competing interests

The authors declare no competing interests.

Acknowledgements

We thank Kor Brandsma and Anneke Ditewig for their excellent biotechnical assistance; Beerend

Winkelman for providing the MATLAB code for spike sorting; Chris van der Togt for providing

assistance with stimulus presentation; Mario Negrello for providing the MATLAB code for

saccade detection; Geert Springeling, Peter Thier, Peter W. Dicke, Ruud Smith for their help with

anatomical scans and 3D model building; and Kaushik J. Lakshminarasimhan for helpful

comments in the analysis. This research was supported by FP7-C7 European Commission, the

Marie Curie Initial Training Network ITN-GA-2009-238214, ENW-Klein, ZonMw

(VIDI/917.18.380,2018), Netherlands Organization for Scientific Research, European Research

Council (Advanced Grant and Proof of Concept Grant), Medical NeuroDelta Programme,

Topsector Life Sciences & Health (Innovative Neurotechnology for Society or INTENSE), and

the Albinism Vriendenfonds Netherlands Institute for Neuroscience.

Abstract

Consciously suppressing responses to distracting external stimuli helps us to reach our goals. This

volitional inhibition of a specific behavior is supposed to be mainly mediated by the cerebral

cortex; however recent evidence supports the involvement of the cerebellum in this process. It is

currently not known whether different parts of the cerebellar cortex play differential or synergistic

roles in planning and execution of this behavior. Here, we measured Purkinje cell (PC) responses

in the medial and lateral cerebellum in two rhesus macaques during the antisaccade task. In this

task the non-human primates were instructed to make a saccadic eye movement away from a target,

rather than towards it. Our data shows that the cerebellum plays an important role not only during

execution of the antisaccades, but also during the volitional inhibition of eye movements towards

the target. Simple Spike (SS) modulation during the instruction and execution period of pro- and

antisaccades is prominent in PCs of both medial and lateral cerebellum. However, only the SS

activity in the lateral cerebellar cortex contains information about trial identity and shows a

stronger reciprocal interaction with complex spikes. Moreover, SS activity of different PC groups

modulate bidirectionally in both regions, but the PCs that show facilitating and suppressive activity

are predominantly associated with instruction and execution, respectively. These findings show

that different cerebellar regions and PC groups contribute to goal-directed behavior and volitional

inhibition, but with different propensities, highlighting the rich repertoire of cerebellar control in

executive functions.

Keywords: Flexible behavior, simple spikes, complex spikes, executive control, antisaccade, non-

human primates

Significance Statement

The antisaccade task is commonly used in research and clinical evaluation as a test of conscious

and flexible control of behavior. It requires volitional suppression of prosaccades, a function that

has been attributed to the neocortex. However, recent findings indicate that cerebellum also

contributes to this behavior. We recorded from neurons in the medial and lateral cerebellum to

evaluate their responses in this task. We found that both regions significantly modulated their

activity during this task, but only cells in the lateral cerebellum encoded the stimulus identity in

each trial. These results indicate that the cerebellum actively contributes to the control of flexible

behavior and that lateral and medial cerebellum play different roles during volitional eye

movements.

Introduction

In a dynamic environment, volitional control of behavior is necessary to make flexible, well-adapted choices. This often requires selective suppression of responses to external stimuli, a hallmark of conscious, executive control (Diamond, 2013). In a laboratory setting, we can measure this complex behavior using the antisaccade task. It requires the subjects to refrain from looking at a suddenly appearing target and instead execute a saccade to the (unmarked) mirror position of that target (Everling and Fischer, 1998; Mitchell et al., 2002; Munoz and Everling, 2004). Similar to other volitional movements, many regions in the cerebral cortex have been identified as crucial for the correct execution of the antisaccade task (Funahashi et al., 1993; Schlag-Rey et al., 1997; Gottlieb and Goldberg, 1999; Bunge et al., 2005; Hakvoort Schwerdtfeger et al., 2012; Everling and Johnston, 2013; Cutsuridis et al., 2014). Given its reciprocal connections with the relevant neocortical regions (Kelly and Strick, 2003), the cerebellum may form an additional hub in this voluntary motor control circuitry (Tanaka et al., 2003; Peterburs et al., 2012; Brunamonti et al., 2014; Kunimatsu et al., 2016; Dacre et al., 2019). This possibility is corroborated by recent findings that the cerebellum participates in movement planning (Ashmore and Sommer, 2013; Giovannucci et al., 2017; Deverett et al., 2018; Gao et al., 2018; Kostadinov et al., 2019). However, to what extent different parts of the cerebellum contribute to the planning and execution of the antisaccades, and complex movements in general, is unclear (Miall et al., 1993; Thach, 2007; Ito, 2008; Gao et al., 2018, 2019; Chabrol et al., 2019; De Schutter, 2019).

Several studies have indicated that execution of simple or reflexive movements may be controlled by Purkinje cells (PCs) in the medial part of the cerebellum, whereas complex

behaviors, which require instruction, inhibition and planning, might be controlled by more lateral

parts (Strick et al., 2009; Caligiore et al., 2017; Chabrol et al., 2019; De Schutter, 2019;

Sendhilnathan et al., 2020). Others however, advocate that both simple and complex movements

can be controlled by the same group of PCs (Gao et al., 2018, 2019), the location of which in the

cerebellar cortex may be determined by the PC-to-effector pathway (Voogd et al., 2012). Settling

these questions would require monitoring of spiking activity of PCs in both medial and lateral

cerebellar areas during a motor task that includes the execution of both simple and complex forms

of movements related to the same effector.

Here, we investigated the hypotheses that (1) PCs in the cerebellum play a role in the

volitional inhibition of stimulus-driven eye movements during the antisaccade task; and (2) that

PCs in medial and lateral cerebellum differentially contribute to pro- and antisaccades. More

specifically, we set out to study modulation of randomly selected saccade-related PCs in the medial

(oculomotor vermis, OMV) and lateral (crus-I/II) cerebellum of non-human primates (macaca

mulatta) during the generation of both pro- and antisaccades (**Fig. 1**). We found that Simple Spike

(SS) modulation during the instruction and execution periods of pro- and antisaccade trials was

prominent in PCs in both cerebellar regions. However, whereas SS activity in the lateral

cerebellum contained information about the trial identity, we were not able to detect this in the

vermis. In addition, Complex Spike (CS) activity showed a higher level of reciprocity with respect

to SS activity in the lateral cerebellum. Although the PC activity modulated bidirectionally in both

cerebellar regions, the PCs in the lateral cerebellum that facilitated their activity, so called upbound

PCs (De Zeeuw, 2021), contributed most prominently to the instruction of antisaccade trials.

Together, our data are adding to the growing body of evidence for a role of the cerebellum in

executive control and are inciting the field to explore cerebellar activity under conditions where

response inhibition is engaged.

Materials and Methods

Animals

All procedures complied with the NIH Guide for the Care and Use of Laboratory Animals

(National Institutes of Health, Bethesda, Maryland), and were approved by the institutional animal

care and use committee of the Royal Netherlands Academy of Arts and Sciences

(AVD8010020184587). Two adult male non-human primates (NHPs, Macaca mulatta), Mi and

Mo, were used in this study.

Surgical procedures

Animals were prepared for eye movement and awake, extracellular single unit recordings

in the cerebellum using surgical and electrophysiological techniques in a two-step procedure

(Roelfsema et al., 2012). Under general anesthesia induced with ketamine (15 mg/kg, i.m.) and

maintained under intubation by ventilating with a mixture of 70% N2O and 30% O2, supplemented

with 0.8% isoflurane, fentanyl (0.005 mg/kg, i.v.), and midazolam (0.5 mg/kg • h, i.v.), we first

implanted a titanium head holder to painlessly immobilize the NHP's head. Four months later,

once the NHPs had mastered the behavioral task (see below), a custom-made 40 mm chamber was

implanted under the same anesthesia conditions as described above to gain access to the cerebellum

with a 25 angle (**Fig. 1B**). Animals recovered for at least 21 days before training was resumed.

Behavioral task

Animals were trained to perform a randomized interleaving pro- and antisaccade task in 8

different directions (cardinal and diagonal directions), with amplitudes between 5° and 14° (**Fig.**

1A). All recordings were conducted in complete darkness. During training and experiments, NHPs

were seated in a primate chair (Crist Instrument, USA) with their head restrained at 100 cm from

a screen with a resolution of 1,024 x 768 pixels. Visual stimuli were presented by a CRT-Projector

Marquee 9500 LC (VDC Display Systems, Florida, USA) with a refresh rate of 100 Hz. Binocular

vision was unrestricted. A trial started when the animal fixated on a red or green fixation point at

the center of the screen for a random time between 300-500 ms. Next, a red target appeared in one

of 8 different target locations. After 100 ms the fixation point changed to gray and the NHP was

allowed to perform either a prosaccade toward the target (when fixation point was red) or

antisaccade in the opposite direction with the same amplitude as the target, (when fixation point

was green) within 500 ms (in the figures colors of fixation point and target were changed to blue

and magenta to accommodate colorblind readers). The animals received liquid reward if they

performed a correct saccade (within 6° of the (anti-)target) and maintained fixation for 100 ms.

Within a single block the 8 (targets) * 2 (pro/anti) possible configurations were presented in a

random order.

Recordings

Position of the right eye was recorded with an infrared video-based eye tracker (iViewX

Hi-Speed Primate, SMI GmbH, Germany) at a sampling rate of 350 Hz. The eye tracker was

calibrated before every recording session by having the animal look at 1° target grid consisting of

9 points (one at the center of the screen and 8 points 10° apart) to adjust the offset of X and Y

position by hand. Pre and post-surgical MRI-images were used to build a 3D model of the skull

and cerebellum for anatomical localization of cerebellar oculomotor vermis (lobules VI and VII)

and lateral cerebellum (crus I/II; **Fig. 1B** (Sendhilnathan et al., 2020). Single-unit recordings were

obtained using tungsten glass-coated electrodes (1-2 MΩ, Alpha Omega Engineering, Nazareth,

Israel) through a 23-gauge guide tube, which was inserted only through the dura. A motorized

microdriver (Alpha Omega Engineering, Nazareth, Israel) with a 1-mm spaced grid was used to

introduce the electrode and map the recording sites with a maximum resolution of 0.25 mm.

Data Analysis Software

All analyses were performed off-line using custom scripts written in MATLAB

(Mathworks, Natick, MA, USA).

Eye movement analysis

Eye position was sampled at 350 Hz with an infrared video eye tracker which tracked the

pupil center of mass. Noise was reduced using a finite impulse response filter and a savitzky-golay

filter (20 ms window). Eye velocity and acceleration traces were created by differentiating the

signal. Eye acceleration was further processed using a median filter. Saccade onset and offset were

detected using an adaptive threshold based on 6 s.d. of the noise during fixation from the recording

session of that day as described in Nyström and Holmqvist (2010) (Sendhilnathan et al., 2020).

Trials with reaction times less than 80 ms were excluded, as these were thought to represent

anticipatory actions.

Electrophysiological analysis and statistics: Simple spikes (SSs)

To perform statistical analysis, neurons were required to have at least 5 recorded trials per

direction as well as per pro- or antisaccadic condition. Only neurons with significant saccade-

related SS activity and only correct trials were incorporated in the analyses. The latter choice was

based on the fact that both NHPs Mo and Mi exhibited an expert level performance reaching >90%

correct responses, rendering insufficient statistical power to analyze the incorrect trials. A neuron

was considered to be saccade-related when SS firing during the baseline period (defined as the

window 400 to 50 ms before trial onset in the intertrial interval), was significantly different from

the saccade execution period activity (defined as the 150 ms time-window after saccade onset) for

at least one of the 8 directions (Wilcoxon rank test; $p < 0.05$). We then computed the instantaneous

SS firing rate of the neurons using a continuous spike density function (SDF) generated by

convolving the spike train with a Gaussian function of $\sigma = 50$ ms width and averaging all the

individual spike density functions (Sendhilnathan et al., 2020). We compared the number of spikes

for all pro- and antisaccade trials of each PC using a Kolmogorov-Smirnov test, and subsequently

determined whether the activity of a PC was significantly different between a pro- and antisaccade

with a *p* value < 0.05. The characteristics of the saccade-related activity, as previously reported in

PC recordings, was quite heterogeneous. On that account, we categorized the PC activity as

facilitation, if the activity increased after instruction or saccade onset, or as *suppression,* if the

activity decreased. For the instruction period, we compared the mean firing rate of each cell from

a window 400 ms before instruction onset to a period of 300 ms at the end of the instruction period.

For the saccade period we compared the mean firing rate of each cell from a window of 150 ms

before saccade onset to a window from saccade onset to 150 ms after saccade onset (Wilcoxon

signed rank, *p* value of < 0.05).

To determine the correlation between kinematic parameters and neuronal activity we

determined Pearson's correlation coefficient (*r*) for all neurons in the two tasks. Modulation ratios

for both areas were obtained by computing the ratio between the response during pro- and

antisaccades.

Changes in firing rate during the instruction period were computed by taking the

differences between the mean firing rates during the baseline period separately for pro- and

antisaccade trials. For the saccade period, we compared the mean activity 150 ms before the

saccade onset to a period 150 ms after saccade onset.

We performed a demixed principal component analysis as described by Kobak et al. (2016)

to decompose the population activity into individual components and extract the dependence of

the PCs on the two stimulus conditions. The dPCA data analysis tool code is available at

http://github.com/machenslab/dPCA (Kobak et al., 2016). Furthermore, all standard settings were

used; data were aligned to instruction offset and binned in 100 ms bins. For determining cross-

validated classification accuracies a 100 iterations were used.

Electrophysiological analysis and statistics: Complex spikes (CSs)

Cells were included into the analysis for complex spikes if they had a CS firing rate of

between 0.5 and 2 hz over the entire recording session and at least 5 trials for every of the 8 pro-

and antisaccade directions. To determine if the CS modulated to one of the task epochs, we

determined whether the CS rate significantly exceeded baseline levels during either the instruction

or the saccade window (50 - 350 ms after onset of the instruction and -150 to +150 ms around the

onset of the saccade). Baseline firing was calculated for a period of 500 ms of inter-trial activity,

and the modulation was considered significant if firing rate exceeded $\pm$ 3 standard deviations from

baseline.

Reciprocity of modulation between SS and CS rates was determined by finding the peak

of the modulation of CS and SS responses during the same 300 ms windows (50 - 350 ms after

instruction onset or -150 - 150 ms around the saccades). We first calculated the peak firing rate

changes for SS and CS from baseline (ΔCS, ΔSS), and next determined the SS-CS interaction, i.e.

reciprocity (ΔCS*ΔSS). Subsequently, we used a linear regression model to find out whether there

was an association between reciprocity in the instruction period and reciprocity during the saccade

period.

Results

Pro- and antisaccades and related PC activity in medial and lateral cerebellum

We trained two adult, male rhesus macaques (referred to as NHPs Mo and Mi) to perform

a randomized, interleaved pro- and antisaccade task. In the prosaccade trials, the NHPs performed

a (pro) saccade to a single visual target in one out of eight different locations separated 45° from

each other. Instead, in antisaccade trials, the NHPs were trained to suppress the prepotent saccade

to a visible target and perform a saccadic eye movement in the opposite direction to an unmarked,

mirror position (**Fig. 1A**).

A trial started with the appearance of the central fixation point (instruction period), the

color of which determined the trial condition. The colors red and green were the instruction cues

for prosaccade and antisaccade movements, respectively (these colors are presented as magenta

and blue in the figures to accommodate color-blind readers). Following the instruction period (300-

500 ms), a target appeared randomly at one of the eight different locations, while NHPs maintained

fixation on the central point. After 100 ms the central fixation point turned gray, serving as the

"go-cue". Animals had 500 ms to execute the pro- or antisaccade eye movement (saccade period)

before the trial was aborted. This task design temporally separates the initial neural representation

of the stimulus (300-500 ms long instruction period) from the subsequent movement execution

(saccade period) (Amador et al., 1998).

Following the training period, we performed single-unit electrophysiological recordings.

We recorded randomly from 90 PCs in medial and 72 PCs in lateral (Crus-I/II) cerebellum during

122 behavioral sessions (**Fig. 1B**). We removed trials with reaction times shorter than 100 ms

following the "go-cue" to avoid contamination with data from "anticipatory" saccades; these

reflect a visual, involuntary reflex towards a novel stimulus in the environment and do not

represent correctly planned saccades (Goldring and Fischer, 1997; Coe and Munoz, 2017). We

only considered correct trials for further analysis, as we did not have enough power to analyze

error trials due to the high performance of both NHPs ($94.1 \pm 5.6\%$ and $94.4 \pm 5.3\%$ for Mo and

Mi, respectively). Eye movements during antisaccades exhibited longer reaction times, lower peak

velocities, larger amplitudes, and longer durations than prosaccades (**Fig. 1C**).

Figure 1. Task, recording location, saccade kinematics and related PC firing characteristics. *A*, A scheme representing the different periods of the task. Trial condition was indicated by the color of the fixation point. After a random delay between 300-500 ms (i.e., instruction period, dashed symbol) a target appeared in one of 8 locations, while NHPs were required to maintain fixation for 100 ms. Next, the fixation target switched to gray, and NHPs were given 500 ms to execute a saccade (i.e., saccade period) towards the target (prosaccade) or the mirror position of the target (antisaccade). *B, left,* CT-based 3D-reconstruction of the skull of Mi; *bottom*, coronal MRI section showing the localization of an electrode in the left lateral cerebellum; *right,* overview of the localization of recordings sites in the medial (vermal lobules VIc and VII) and lateral cerebellum (crus-I/II, left hemisphere). *C*, Features of kinematics of all prosaccades (magenta; monkey Mo $n = 6114$, monkey Mi $n = 6670$) and antisaccades (blue; Mo $n = 4114$, Mi $n = 6305$) for all trials over the 122 recording sessions. Antisaccades were characterized by longer reaction times (pro- mean 178.4 ms $\pm$ 70.6 s.d.; anti- mean 236.6 ms $\pm$ 72.7 s.d.; $p < 0.0001$), lower peak velocities (pro- mean 311.8°/s $\pm$ 102.6 s.d.; anti- mean 253.4°/s $\pm$ 103.1 s.d.; $p < 0.0001$), bigger amplitudes (pro- mean 5.8° $\pm$ 2.3 s.d.; anti- mean 9.5° $\pm$ 4.1 s.d.; $p < 0.0001$), and longer durations (pro- mean 34.7 ms $\pm$ 7.9 s.d.; anti- mean 69.8 ms $\pm$ 24 s.d.; $p < 0.0001$). Triangles in magenta and purple indicate means of pro- and antisaccade movement parameters, respectively. *D*, Scatter plots with corrected PC activity after a linear regression model fitted to amplitude or peak velocity during all pro- and antisaccades in medial (left) and lateral (right) cerebellum. *E*, Violin plots showing the mean average simple spike firing rate for all cells during prosaccades (magenta) and antisaccades (blue) for medial (left) and lateral (right) cerebellum. Dashed lines depict the mean value for the population. The asterisks

show significance levels for a Wilcoxon rank sum test for the comparison between medial and lateral cerebellum in each condition (pro medial vs. lateral $p = 0.04$; anti medial vs. lateral $p = 0.03$).

A neuron was considered to be saccade-related if SS activity during the beginning of the

saccade period (i.e., from saccade onset to 150 ms after saccade onset) was significantly different

from the intertrial interval period (see **Methods**). Approximately half of the cells recorded in both

areas fulfilled this criterion (n = 40 (44%) for the medial, and n = 38 (53%) for the lateral

cerebellum). All subsequent analyses were performed only for these cells unless noted otherwise

(n = 78 PCs). The average number of trials per cell in the medial cerebellum was 75 ± 49 and 71

± 47 for pro- and antisaccades, respectively, while in the lateral cerebellum these numbers were

80 ± 52 and 76 ± 50, respectively.

To assess whether there was a correlation between saccade parameters and PC SS firing

rate, we regressed the firing rates in the two recording areas to amplitude or peak velocity and

compared the corrected firing rate using the regression coefficient (ß) between pro- and

antisaccade (corrected firing = firing rate/regression coefficient). We found no significant

correlations between SS firing rate and saccade amplitude or saccade peak velocity in neither the

medial nor the lateral cerebellum (**Fig. 1D**). This allowed us to directly compare pro- and

antisaccade trials between the two areas by pooling the mean firing rates of each cell in each

condition. SS activity during saccade execution was significantly higher in the lateral cerebellum

compared to that in the medial cerebellum for both pro- and antisaccade trials (mean and s.d.,

medial pro 66.7 ± 30 spks/s vs. lateral pro 69.3 ± 39 spks/s, $p = 0.04$; medial anti 68 ± 36 spks/s

vs. lateral anti 71.8 ± 39 spks/s, $p = 0.03$; Wilcoxon rank sum test) (**Fig. 1E**).

PCs in both medial and lateral cerebellum modulate their activity during the task but with different characteristics

The activity profile of the PCs followed roughly one of two patterns, they either increased or decreased their activity following saccade onset. Accordingly, we classified these cells as *facilitation* or *suppression* PCs, respectively (**Fig. 2**). The activity profile of the PCs was the same during pro- and antisaccades in both areas, meaning that if one PC increased its firing rate after saccade execution for prosaccades, it also increased it for antisaccades. This was true for all PCs except for two in the lateral cerebellum, which showed a mixed firing behavior; these two cells were included in their respective category for further analyses (e.g., as a *facilitation* cell in prosaccade trials and a *suppression* cell in antisaccade trials).

We next assessed SS activity during the instruction period (for details, see **Methods**). Given the variable duration of the instruction period, which ranged between 300 and 500 ms, we analyzed the last 300 ms of this epoch, before target onset (**Fig. 1A**). Here, we also encountered diverse neural responses in both regions. We classified the activity of these neurons in the same way as for the saccade period, depending on whether they increased (*facilitation* PCs), or decreased (*suppression* PCs) their SS firing during the instruction period compared to the activity during the intertrial interval. The medial compared to the lateral cerebellum had overall less *facilitation* PCs (Medial pro 10%, medial anti ~11%, Lateral pro ~17%, Lateral anti ~18%) and *suppression* PCs (Medial pro ~11%, Medial anti ~7%, Lateral pro ~14%, Lateral anti ~15%) during the instruction of pro- and antisaccade trials (**Fig. 2B-F**). The remaining cells had no change in firing rates during the instruction period (medial: ~61%, lateral 36%).

Figure 2. PCs in both medial and lateral cerebellum start modulating their activity during the instruction period. *A top*, Eye trace for an example trial during saccade onset for a pro- (left) and antisaccade (right). Dashed line indicates the time of the "go" cue. Dark and light magenta and blue show the x and y position (horizontal and vertical components) of the eye trace for pro- and antisaccades respectively; ***bottom***, traces of activity recorded from a PC in medial cerebellum for a single trial aligned to the eye trace indicated above. Note that the traces include many simple spikes (SSs; smaller spikes) and only a few complex spikes (CSs, bigger spikes). *B*, *left* Average response time-course for one example neuron aligned to the end of the instruction period (target onset) for prosaccades (magenta) and antisaccades (blue) for medial cerebellum. Dashed line shows the baseline mean. Shaded regions denote s.e.m. Gray shaded areas show time after the instruction period when the target appears, but the animals must keep fixating at the central dot; *right* Average response time-course for one example neuron aligned to saccade onset. Dashed line shows the baseline mean. **C,** Time-course of normalized responses averaged for all neurons in medial cerebellum that either increased their firing (*facilitation* cells, solid line) or decreased it (*suppression* cells, dashed lines) aligned to target onset (left) or saccade onset (right). Statistical comparisons after saccade onset, facilitation pro vs anti: $p = 1.61 \times 10^{-7}$, suppression pro vs anti: $p = 5.3 \times 10^{-23}$. **D,** Same as in (a) but for lateral cerebellum. *E*, Same as in (b) but for lateral cerebellum. *F*, Same as in (c) for lateral cerebellum (facilitation pro vs anti: $p = 1.08 \times 10^{-17}$, suppression pro vs anti: $p = 4.4 \times 10^{-9}$). Prosaccades are denoted in magenta and antisaccades in blue for all panels.

To relate PC activity during the instruction period with the subsequent saccade period, we

next looked at the timing of the SS activity of *facilitation* and *suppression* PCs between medial

and lateral cerebellum pooling the pro- and antisaccade conditions (**Fig. 2**). We found that the

latencies of the *suppression* PCs in the lateral cerebellum, measured at the trough after saccade

onset, were significantly longer than those in the medial cerebellum (mean ± s.d. Lateral 133 ± 40

ms vs. Medial 89 ± 68 ms, $p = 0.01$; Wilcoxon rank sum, **Fig. 2C** and **F**). In contrast, the *facilitation*

PCs showed similar timing of the peak activity in the medial and lateral cerebellum (mean ± s.d.

Medial 105 ± 41 ms vs. Lateral pro 84 ± 48 ms; $p = 0.4$; Wilcoxon rank sum). There were no

differences between prosaccades and antisaccades in this respect, neither for *suppression* PCs

(mean ± s.d. Medial pro 87 ± 70 ms vs. Medial anti 93 ± 67 ms, $p = 0.83$; Lateral pro 139 ± 40 ms

vs. Lateral anti 127 ± 42 s; $p = 0.1$; Wilcoxon Rank sum test), nor for *facilitation* PCs (mean ± s.d.

Medial pro 105 ± 41 ms vs. Medial anti 104 ± 33 ms, $p = 0.2$; Lateral pro 84 ± 48 ms vs. Lateral

anti 106 ± 43 ms, $p = 0.9$; Wilcoxon Rank sum test).

Together, these results suggest that PCs in the lateral cerebellum are more prominently

involved in the instruction of planned saccades than those in the medial cerebellum, and that the

course of SS responses of *suppression* PCs in the lateral cerebellum during the subsequent saccade

period is delayed with respect to that in the medial cerebellum.

PCs in lateral cerebellum show a stronger modulation during the instruction period

We observed that in some PCs the SS modulation ramped throughout the interval between

the instruction and the saccade epoch (**Fig. 2B-E** and **Fig. 3A**). We quantified the ramping of SS

activity by fitting a linear model to the firing rate during the last 300 ms of the instruction period,

and we classified PCs as ramping cells when R^2 was higher than 0.75. We found that ramping

modulation was quite rare in the medial cerebellum and more prevalent in the lateral cerebellum

(**Fig. 3B**); this held true for both prosaccade and antisaccade trials (Medial *n* (pro) = 3, *n* (anti) =

2; Lateral n (pro) = 11, n (anti) = 10; both comparisons p = 0.02; χ^2 test). Overall, we found more

PCs that significantly changed their firing from baseline during the instruction period in the lateral

cerebellum than in the medial cerebellum (Medial vs Lateral, p = 0.03; Mann-Whitney test),

suggesting a more prominent role for PC modulation in the lateral cerebellum during instruction.

Figure 3. PCs in the lateral cerebellum exhibit stronger modulation than PCs in the medial cerebellum during the instruction period. *A*, Explanation of slope calculations used for panel (b); *left*, Time-course of responses during prosaccades (magenta) and antisaccades (blue) for an example neuron in the lateral cerebellum aligned to "go" cue (vertical dashed line at zero); *right*, activity of the same neuron, but now for an antisaccade trial also aligned to the "go" cue . Solid vertical line shows the end of the instruction period, shaded area is when the target appears but the NHPs must keep fixating. A linear regression model was fitted (orange diagonal solid and green dotted line, for pro- and antisaccades, respectively) to the firing rate 300 ms before the end of the instruction period to calculate the slopes (see **Methods**). ***B*,** Number of cells with significant slope modulation in the medial and lateral cerebellum (χ^2 test for proportions) during pro- and antisaccades (p = 0.02 for both comparisons). ***C*,** Scatter plots of maximum simple spike firing rate change of PCs during the instruction period vs. that during the saccade

period (in spks/s) for pro- and antisaccades in medial cerebellum. *D*, Same as in (c) but for lateral cerebellum. *E*, Difference in change in absolute firing rate between pro- and antisaccades (abs antisaccade SS activity- abs prosaccade SS activity) for *facilitation* and *suppression* neurons in medial cerebellum. *F,* Same as in (e) but for lateral cerebellum. *G*, Cumulative distribution of the SS modulation ratio for medial and lateral cerebellum for *facilitation* and *suppressive* PCs. Insets show *p*-values for the comparison between medial and lateral cerebellum (two sample K-S test).

When we investigated the changes in the firing rate (modulation, **Methods**) for each PC

during the instruction and saccade period within the medial and lateral cerebellum, we found that

in the lateral cerebellum, the maximum change in the firing rate was significantly lower during the

instruction period than during the saccade period, for both pro- and antisaccade trials (Lateral pro

$p = 0.002$, anti $p = 0.01$; Wilcoxon rank sum test) (**Fig. 3C**). In the medial cerebellum only the

prosaccade trails showed this trend (Medial pro $p = 0.03$, anti $p = 0.07$, Wilcoxon rank sum test)

(**Fig. 3D**).

Further examination of the differences in responses during pro- and antisaccades showed

that *facilitation* and *suppression* cells in the lateral cerebellum show significant encoding of SS

modulation for trial conditions. We calculated a modulation ratio between the responses during

the saccade period in the two types of trials, defined as the ratio between the response during pro-

and antisaccades (**Fig. 3G**, **Methods**). Values of 1 indicate that both responses are the same, values

higher than 1 indicate that prosaccade activity was higher, and below 1 that antisaccade activity

was higher. Medial PCs had somewhat different responses when comparing pro- and antisaccade

activity in contrast to lateral PCs (mean of *facilitation* PCs in medial cerebellum 0.91 ± 0.1 s.d.

versus that in lateral cerebellum 0.88 ± 0.3, $p = 0.03$; mean of *suppression* PCs in medial

cerebellum 1.16 ± 0.2 versus that in lateral cerebellum 1 ± 0.2, $p = 8.02$ x 10^{-04}; one-tailed K-S

test). To summarize, PCs in both medial and lateral cerebellum modulate their activity during

instruction of pro- and antisaccades, with a relatively bigger role for the *facilitation* cells during

the instruction of antisaccades. Yet, PCs in the lateral cerebellum differ from those in the medial

cerebellum in that they have a relatively stronger modulation in both the instruction and saccade

periods as well as for the pro- vs antisaccade conditions.

Differences between facilitation and suppression cells that occur in both medial and lateral

cerebellum

In addition to the observed differences across the recorded regions, we identified features

that were shared across the medial and lateral cerebellum. First, we observed that *facilitation* and

suppression PCs in medial and the lateral cerebellum operate at different baseline frequencies,

which is in line with the upbound and downbound modules of PCs described for the cerebellar

cortex of rodents (De Zeeuw, 2021). Since these modules have been hypothesized to operate at a

relatively low and high baseline SS firing frequency, respectively, so as to leave ample room for

increases and decreases during epochs of stimulation and modulation, we investigated whether the

suppression and *facilitation* cells differed in this respect (see also (Sun et al., 2017; Herzfeld et al.,

2018; Soetedjo et al., 2019). We found that *suppression* cells started at a relatively high baseline

SS firing frequency in both the medial and lateral cerebellum (62.43 ± 3.76 spks/s and 69.6 ± 6.12

spks/s, respectively), and *facilitation* cells started at a significantly lower baseline SS firing

frequency in both regions (56.36 ± 3.65 spks/s and 54.72 ± 3.76 spks/s, $p = 0.02$; one-tailed

Wilcoxon rank-sum test comparison between all *facilitation* vs *suppression* cells pooled in both

areas).

Next, we found that whereas the saccade period contained more *suppression* PCs, during

the instruction period the contribution of the *facilitation* cells appeared most prominent,

particularly during that of the antisaccade trials. When we subtracted the SS modulation during

the prosaccade trails from the activity during the antisaccade trials (antisaccade SS activity -

prosaccade SS activity), *facilitation* PCs in both medial and lateral cerebellum showed a larger

change in firing rate just before the target onset during the instruction period (**Fig. 3E, F**). These

changes were significantly less pronounced for *suppression* cells (*facilitation* PCs versus

suppression PCs in medial cerebellum p = 4.4x10-6; *facilitation* PCs versus *suppression* PCs in

lateral cerebellum p = 0.004; Wilcoxon Rank sum test). Thus, *facilitation* and *suppression* cells

can be characterized by several features that are common for medial and lateral cerebellum.

The lateral but not the medial cerebellum contains information about stimulus identity during the

instruction period

To extract the features of the population activity related to pro- and antisaccades we applied

a linear dimensionality reduction technique: demixed principal component analysis (dPCA,

(Kobak et al., 2016). dPCA demixes the neural activity into activity related to the task components

that capture the most variance in the data. The dPCA was used to investigate to what extent the

neurons are tuned to two stimulus conditions indicated by the color of the fixation point (i.e. the

pro- or antisaccades), or whether their activity is condition-independent. In the condition-

independent group, dPCA components show task related activity modulations that cannot be

assigned to one of the stimulus conditions. As explained above, due to the high behavioral

performance of both NHPs, our dataset lacked the power to create another category with decision-

related components in correct/incorrect trials. dPCA was applied to the entire population of

recorded cells in the medial (n = 90) and the lateral cerebellum (n = 72). We included cells which

had at least 5 trials in pro- and antisaccade conditions. The results were cross-validated to measure

time-dependent classification accuracy and a shuffling procedure was applied to assess whether

classification accuracy was significantly above chance level (see **Methods**).

In the medial cerebellum, the overall variance explained by the dPCA components was

mostly captured by the first 3 components, which were independent of the stimulus condition, and

together they explained 66.9 % of variance, (**Fig. 4A**). These components all have strong activity

after the end of the instruction, and thus around the time the animal initiates the saccade (**Fig. 4B**).

Components 4, 5 and 7 were selected as containing the strongest activity differences in relation to

the stimulus conditions (**Fig. 4B**, bottom row and **Fig. 4C**). To determine significance, each

component was used as a linear decoder to classify each of the stimulus conditions (see **Methods**).

No components could reliably be used to classify the stimulus condition in the medial cerebellum.

Figure 4. Demixed PCA indicates that PCs in the lateral cerebellum encode stimulus identity. *A,* Cumulative fraction of variance explained by dPCA in the medial cerebellum. Dotted line indicates an estimate of the fraction of variance in the data explained by the dPCA. *B,* Projections of the PSTH's of all medial cerebellar PCs onto the most prominent decoding axes; *top*, first three condition-independent components; *bottom*, first three stimulus-related (pro/anti) components. Percentages denote variance captured by each component. Time from instruction offset. *C,* Variance of the individual demixed principal components for the medial cerebellum. Each bar shows the proportion of total variance. Gray bars depict condition-independent variance, cyan bars indicate stimulus-related (pro/anti) variance. Pie chart shows the total signal variance for stimulus-related and condition-independent components. *D-F,* Same as in (A-C) but for the lateral cerebellum, orange horizontal bars in *E* indicate time during which the stimulus can be decoded from the population firing rate.

Similar to the medial cerebellum, components 1, 2 and 3 of the lateral cerebellum captured

most of the variance in the condition independent categories (56.4%, **Fig 4E**). These components

also showed a strong modulation after instruction offset. In contrast to the medial cerebellum, two

of these components modulated during most of the instruction period (**Fig. 4F).** Stimulus related

components showed significant tuning for pro- and/or antisaccades in components 5 and 7.

Notably, in component 5, stimulus condition classification was already possible almost 200 ms

before the end of the instruction period. In the lateral cerebellum the variance in this component

could almost be entirely attributed to the stimulus conditions (**Fig. 4F**). Indeed, in component 5,

the time period in which the stimulus condition could be decoded from the firing rates was broad,

bridging the instruction period to the saccade. This appears to be in line with the prominent

ramping modulation described in **Figure 3**. These results show that PCs in the lateral, but not in

the medial, cerebellum contain reliable information about the trial identity during the instruction

period.

Complex spike responses of PCs in the lateral cerebellum show reciprocal activity with simple

spike responses

In a subset of recorded PCs, we were able to reliably isolate and analyze CS responses

throughout all trials. We restricted the CS analysis only to cells with average CS firing rates higher

than 0.5 Hz over the duration of the whole recording (Medial n = 11, Lateral n = 18). Similar to

the SS analyses, we examined the CS responses separately in the medial and lateral cerebellum

during the instruction and saccade periods (**Fig. 5**). The average CS activities during the instruction

(1.10 Hz ± 0.47 s.d., n = 11) and saccade (1.11 Hz ± 0.50 s.d., n = 11) period in the medial

cerebellum were not significantly different from those in the lateral cerebellum (1.15 Hz ± 0.33

s.d., n = 18 and 1.13 Hz ± 0.52 s.d., n = 18, respectively; $p = 0.87$ and $p = 0.36$ for comparison

medial versus lateral during instruction and saccade period, respectively; Wilcoxon rank sum).

Figure 5. Complex spike and simple spike interaction during pro- and antisaccade trials. *A*, Top, average CS rate of PCs in medial cerebellum during prosaccades (magenta) and antisaccades (blue) aligned to the onset of the instruction (left) and saccade (right) period. Bottom panels show the associated SS responses. Shaded areas represent s.e.m. *B*, Same as *A*, but for neurons recorded in the lateral cerebellum. *C*, Relation of reciprocity between CS and SS responses during instruction and saccade period. To quantify the reciprocity, we calculated the maximum change in CS firing rate in the windows of interest (instruction and saccade period), the associated maximum change in SS rate in a window of 300 ms around the peak of CS activity, and subsequently the product of the two (Δ CS rate x Δ SS rate). Note that if a positive modulation of CSs is associated with a negative modulation of SSs, the reciprocity measure presented here will give a negative value. If CSs and SSs show simultaneous positive modulations, and thus no reciprocity, this will result in a positive value. Every dot represents one cell.

One outlier was removed from the medial antisaccade group (unfilled blue dot). The coefficients of determination (R^2) are: medial pro 0.30, medial anti 0.01. **D**, same as C but of lateral neurons, coefficients of determination: lateral pro 0.59, and lateral anti 0.48. Shading indicates 95% confidence interval of the regression lines.

Next, we quantified the interaction between CS and SS responses during the instruction

period (**Fig. 5A** and **B,** left panels). For every PC of both medial and lateral cerebellum we

calculated the maximum change in CS firing rate from the baseline in the instruction window and

the maximum change in SS firing rate from baseline in the 300 ms window around the time of the

peak CS rate. This was also done for the saccade period for the pro- and antisaccade trials

separately. Change in CS firing from the baseline was multiplied with change in SS firing from

baseline to establish their interdependence, i.e., reciprocity (Badura et al., 2013). To determine

whether reciprocity was dependent on both epochs, linear regression was applied on the reciprocity

values in the instruction and saccade windows (**Fig 5C, D)**. Lateral cerebellar PCs had a positive

association with reciprocity during the instruction and the saccade period, whereas medial

cerebellum did not (coefficients of determination (R^2) : lateral pro 0.59, and lateral anti 0.48, medial

pro 0.30, medial anti 0.01).

Discussion

The antisaccade task has been used to investigate flexible behavioral control, where a correct suppression of a prosaccade followed by a response to the opposite direction forms a feature of executive control (Mokler and Fischer, 1999; Mitchell et al., 2002). We found evidence for cerebellar involvement in volitional control of this behavior in both medial (i.e., OMV) and lateral (i.e., crus I/II) cerebellar cortex by recording the activity of PCs during pro- and antisaccades in NHPs. Our findings add to the growing body of evidence pointing to an important contribution of the cerebellum to cognitive processes in general and particularly voluntary behavior. We demonstrated that PCs in both areas modulate their activity depending on the trial type, and that SS responses of populations of PCs in the lateral cerebellum are able to discriminate the type of trial before the eye movement takes place. Even though the lateral cerebellum is not traditionally viewed as a saccade-control area, our findings support early reports where saccades are elicited following electrical stimulation to lateral regions of the hemispheres, especially in crus I/II (Ron and Robinson, 1973).

Antisaccades exhibit kinematic properties that differ from prosaccades, a result consistent with previous reports in human and non-human primates (Amador et al., 1998; Munoz and Everling, 2004). Even so, the preferential modulations of SS responses in the PCs were not correlated to any of those kinematic changes. This suggests that the observed activity was not merely an efference copy of the motor command but may also reflect the preparatory or planning phase of the voluntary movement in which the PC firing is determined by the cue. This is in line with recent findings in cerebellar atrophy patients who show impairments when making goal-oriented movements (Piu et al., 2019).

Various parameters of PC activity differed between medial and lateral cerebellum. SS

activity of PCs in the lateral cerebellum (1) displayed more signal separation/selectivity for pro-

and antisaccades during the instruction period, suggesting that the population contains information

about the trial identity, (2) showed a greater firing rate during the execution of saccades. Likewise,

their CS activity showed a higher level of reciprocity with respect to simple spike activity, during

both the instruction and execution period. Together, these data indicate that different modules of

different cerebellar regions can contribute to the same complex behavior, yet with different

propensities.

PC modulation during instruction period

Studies from cerebellar patients suggest that the cerebellum might not be involved in the

initiation of saccades, but in the correct and accurate execution of them (Kornhuber, 1971; Thier

et al., 2002). Our results indicate that PCs in the lateral cerebellum encode information about the

trial identity. The dPCA analysis in Crus I/II PCs exhibited significantly different SS activity

between both conditions, showing that these contain sufficient information to distinguish a pro-

from an antisaccade. This is supported by a previous study showing that downstream neurons in

the dentate nucleus modulate during preparation of antisaccades and that inactivation thereof

promotes antisaccade errors (Kunimatsu et al., 2016). This instruction-related activity in

antisaccade trials may be relayed to brainstem nuclei that are engaged in curtailing reflexive

saccades (Everling et al., 1999). The persistent SS modulation of PCs in lateral cerebellum upon

cue presentation suggests that it updates an internal model for well-timed activity during motor

planning, which may be relayed to the prefrontal cortex (Gao et al., 2018).

Correct decoding of a visual stimulus is an essential component of the pro- and antisaccade

task. Visual responses in the cerebellum have been reported in several areas including the floccular

complex, crus I/II and the inferior semilunar lobe (VII/VIII) (Marple-Horvat and Stein, 1990).

Dentate nucleus neurons have activity bridging the period from task instructions to motor

execution (Kunimatsu et al., 2016), which could be derived from the same population of PC's as

we recorded in the lateral cerebellum. Our results on the relatively prominent CS - SS reciprocity

in lateral cerebellum raise the possibility that CS responses play a similar role in decoding visual

stimuli for the preparation of ensuing movements.

Contemporary models of antisaccade decision-making require evidence accumulation to

reach a decision boundary in the frontal cortex (Munoz and Everling, 2004). PCs receive diverse

sensory, motor and cognitive information through their parallel fiber inputs, which may facilitate

efficient evidence accumulation. Classical models of cerebellar function indicate that climbing

fiber mediated plasticity in the molecular layer determines which parallel fiber inputs are turned

into SS modulation (Gao et al., 2012). The PCs that we measured in the lateral cerebellum carry

both SS and CS signals during the instruction period of the antisaccade task. We hypothesize that

through the pre-learned associations of parallel fiber and climbing fiber signals, relevant evidence

from the stimulus (i.e., dot color in instruction) for the selection of the required action (i.e., pro-

or antisaccade) can be rapidly relayed from the lateral cerebellum to the frontal saccade areas. In

cerebellar patients, latencies for antisaccades are prolonged, indicating a less efficient decision-

making process in absence of cerebellar input (Piu et al., 2019).

PC modulation during saccade execution

Single PCs modulated their SS activity during the execution of pro- and antisaccades in

both medial and lateral cerebellum, but only lateral PCs were able to successfully discriminate the

type of trial. Despite the heterogeneity of the responses, neurons that facilitated or suppressed their

response after saccade onset exhibited differential activity during pro- and antisaccade trials (**Fig.**

3). As a population, PC responses were different when executing a saccade towards a visible target

than when using an internally generated goal. This is particularly interesting for two reasons: First,

medial cerebellum has always been identified as a motor and timing controller for the different

types of saccades, interacting with cortical and brainstem brain areas (McElligott and Keller, 1984;

Noda and Fujikado, 1987; Thier and Möck, 2006; Herzfeld et al., 2015, 2018), while the role of

the lateral cerebellum in saccade control has remained under debate. Indeed, lesions of lateral

cerebellum can delay onset of saccades between 10 and 60 ms and result in variable hypo- and

hypermetria (Ohki et al., 2009), but the contribution of Purkinje cell activity in the lateral

cerebellum to proactive control of saccades and flexible behavioral in general has not been

conclusively proven. Second, differential cerebellar activity during antisaccades implies that it

contains neural signatures to modulate preparation and execution by encoding information about

the current context of the upcoming action (saccade), which is necessary for the fast classification

of the response as correct or erroneous. This finding is in line with the impact of focal cerebellar

lesions in humans on the changes in potentials observed during performance monitoring in an

antisaccade task (Peterburs et al., 2012).

Cerebellar modules operate in parallel

The data on *suppression* PCs in the medial cerebellum showing a relatively early SS

modulation during execution of both pro- and antisaccades as well as those on *facilitation* PCs in

the lateral cerebellum showing a prominent modulation at the end of the instruction of antisaccades

indicate that different modules of different cerebellar regions can simultaneously contribute to the

same complex behavior, yet with different particularities. A similar conclusion was drawn from a

recent study on delay eyeblink conditioning (Wang et al., 2020). Even though this form of

conditioning is classically considered to be controlled solely by modules in lobule simplex of the

lateral cerebellum, Wang and colleagues have shown that modules of the medial cerebellum are

equally essential, yet also contributing in a slightly differential fashion, possibly regulating mainly

muscle tone. Similar to the current pro- and antisaccade tasks, during eyeblink conditioning the

medial and lateral cerebellum also both engage *facilitation (i.e., upbound)* and *suppression (i.e.,*

downbound) cells, and they also operate at relatively low and high baseline firing frequencies,

respectively (ten Brinke et al., 2015; Wang et al., 2020). Our current finding that *facilitation* and

suppression cells appear to play a more dominant role during the instruction and execution of the

saccade task, respectively, further highlights the differential functional relevance of the upbound

and downbound modules (De Zeeuw, 2021).

Previous reports of activity in the dentate nucleus, which is the main target of PCs in the

lateral cerebellum, have shown mixed evidence regarding the direction of modulation during

saccade-related behavior (Ashmore and Sommer, 2013; Kunimatsu et al., 2018). Kunimatsu and

colleagues (2018) found only upward modulating units during self-initiated saccades, whereas

Ashmore and Sommer (2013) revealed groups of both facilitating and suppressing cells during

delayed saccades. Possibly, these differences reflect differences in the behavioral paradigms and/or

downstream pathways involved (De Zeeuw, 2021). In our dataset, the modulation exhibited by

PCs is bidirectional in both medial and lateral cerebellum with populations of both *facilitation* and

suppression cells.

References

Amador N, Schlag-Rey M, Schlag J (1998) Primate antisaccades. I. Behavioral

characteristics. J Neurophysiol 80:1775–1786.

Ashmore RC, Sommer MA (2013) Delay activity of saccade-related neurons in the caudal

dentate nucleus of the macaque cerebellum. J Neurophysiol 109:2129–2144.

Badura A, Schonewille M, Voges K, Galliano E, Renier N, Gao Z, Witter L, Hoebeek FE,

Chedotal A, De Zeeuw CI (2013) Climbing fiber input shapes reciprocity of Purkinje cell

firing. Neuron 78:700–713.

Brunamonti E, Chiricozzi FR, Clausi S, Olivito G, Giusti MA, Molinari M, Ferraina S,

Leggio M (2014) Cerebellar damage impairs executive control and monitoring of

movement generation. PLoS One 9:e85997.

Bunge SA, Wallis JD, Parker A, Brass M, Crone EA, Hoshi E, Sakai K (2005) Neural

circuitry underlying rule use in humans and nonhuman primates. J Neurosci 25:10347–

10350.

Caligiore D, Pezzulo G, Baldassarre G, Bostan AC, Strick PL, Doya K, Helmich RC, Dirkx

M, Houk J, Jorntell H, Lago-Rodriguez A, Galea JM, Miall RC, Popa T, Kishore A,

Verschure PF, Zucca R, Herreros I (2017) Consensus Paper: Towards a Systems-Level

View of Cerebellar Function: the Interplay Between Cerebellum, Basal Ganglia, and

Cortex. Cerebellum 16:203–229.

Chabrol FP, Blot A, Mrsic-Flogel TD (2019) Cerebellar Contribution to Preparatory

Activity in Motor Neocortex. Neuron 103:506–519.e4.

Coe BC, Munoz DP (2017) Mechanisms of saccade suppression revealed in the anti-

saccade task. Philos Trans R Soc Lond B Biol Sci 372 Available at:

http://dx.doi.org/10.1098/rstb.2016.0192.

Cutsuridis V, Kumari V, Ettinger U (2014) Antisaccade performance in schizophrenia: a

neural model of decision making in the superior colliculus. Front Neurosci 8:13.

Dacre J, Colligan M, Ammer J, Schiemann J, Clarke T, Chamosa-Pino V, Claudi F, Alex

Harston J, Eleftheriou C, Pakan JMP, Huang C-C, Hantman A, Rochefort NL, Duguid I

(2019) Cerebellar-recipient motor thalamus drives behavioral context-specific movement

initiation. bioRxiv:802124 Available at: https://www.biorxiv.org/content/10.1101/802124v1

[Accessed October 23, 2019].

De Schutter E (2019) Fallacies of Mice Experiments. Neuroinformatics 17:181–183.

Deverett B, Koay SA, Oostland M, Wang SS-H (2018) Cerebellar involvement in an

evidence-accumulation decision-making task. Elife 7 Available at:

http://dx.doi.org/10.7554/eLife.36781.

De Zeeuw CI (2021) Bidirectional learning in upbound and downbound microzones of the

cerebellum. Nat Rev Neurosci 22:92–110.

Diamond A (2013) Executive functions. Annu Rev Psychol 64:135–168.

Everling S, Dorris MC, Munoz DP (1999) Neuronal Activity in Monkey Superior

Colliculus during an Antisaccade Task. In: Current Oculomotor Research: Physiological

and Psychological Aspects (Becker W, Deubel H, Mergner T, eds), pp 17–23. Boston, MA:

Springer US.

Everling S, Fischer B (1998) The antisaccade: a review of basic research and clinical

studies. Neuropsychologia 36:885–899.

Everling S, Johnston K (2013) Control of the superior colliculus by the lateral prefrontal

cortex. Philos Trans R Soc Lond B Biol Sci 368:20130068.

Funahashi S, Chafee MV, Goldman-Rakic PS (1993) Prefrontal neuronal activity in rhesus

monkeys performing a delayed anti-saccade task. Nature 365:753–756.

Gao Z, Davis C, Thomas AM, Economo MN, Abrego AM, Svoboda K, De Zeeuw CI, Li N

(2018) A cortico-cerebellar loop for motor planning. Nature 563:113–116.

Gao Z, Thomas AM, Economo MN, Abrego AM, Svoboda K, De Zeeuw CI, Li N (2019)

Response to "Fallacies of Mice Experiments." Neuroinformatics 17:475–478 Available at:

http://dx.doi.org/10.1007/s12021-019-09433-y.

Gao Z, van Beugen BJ, De Zeeuw CI (2012) Distributed synergistic plasticity and

cerebellar learning. Nat Rev Neurosci 13:619–635.

Giovannucci A, Badura A, Deverett B, Najafi F, Pereira TD, Gao Z, Ozden I, Kloth AD,

Pnevmatikakis E, Paninski L, De Zeeuw CI, Medina JF, Wang SS (2017) Cerebellar

granule cells acquire a widespread predictive feedback signal during motor learning. Nat

Neurosci 20:727–734.

Goldring J, Fischer B (1997) Reaction times of vertical prosaccades and antisaccades in gap

and overlap tasks. Exp Brain Res 113:88–103.

Gottlieb J, Goldberg ME (1999) Activity of neurons in the lateral intraparietal area of the

monkey during an antisaccade task. Nat Neurosci 2:906–912.

Hakvoort Schwerdtfeger RM, Alahyane N, Brien DC, Coe BC, Stroman PW, Munoz DP

(2012) Preparatory neural networks are impaired in adults with attention-

deficit/hyperactivity disorder during the antisaccade task. Neuroimage Clin 2:63–78.

Herzfeld DJ, Kojima Y, Soetedjo R, Shadmehr R (2015) Encoding of action by the Purkinje

cells of the cerebellum. Nature 526:439–442.

Herzfeld DJ, Kojima Y, Soetedjo R, Shadmehr R (2018) Encoding of error and learning to

correct that error by the Purkinje cells of the cerebellum. Nat Neurosci 21:736–743.

Ito M (2008) Control of mental activities by internal models in the cerebellum. Nat Rev

Neurosci 9:304–313.

Kelly RM, Strick PL (2003) Cerebellar loops with motor cortex and prefrontal cortex of a

nonhuman primate. J Neurosci 23:8432–8444.

Kobak D, Brendel W, Constantinidis C, Feierstein CE, Kepecs A, Mainen ZF, Qi X-L,

Romo R, Uchida N, Machens CK (2016) Demixed principal component analysis of neural

population data. Elife 5 Available at: http://dx.doi.org/10.7554/eLife.10989.

Kornhuber HH (1971) Motor functions of cerebellum and basal ganglia: the

cerebellocortical saccadic (ballistic) clock, the cerebellonuclear hold regulator, and the basal

ganglia ramp (voluntary speed smooth movement) generator. Kybernetik 8:157–162.

Kostadinov D, Beau M, Pozo MB, Häusser M (2019) Predictive and reactive reward signals

conveyed by climbing fiber inputs to cerebellar Purkinje cells. Nat Neurosci 22:950–962.

Kunimatsu J, Suzuki TW, Ohmae S, Tanaka M (2018) Different contributions of

preparatory activity in the basal ganglia and cerebellum for self-timing. Elife 7 Available at:

http://dx.doi.org/10.7554/eLife.35676.

Kunimatsu J, Suzuki TW, Tanaka M (2016) Implications of Lateral Cerebellum in Proactive

Control of Saccades. Journal of Neuroscience 36:7066–7074.

Marple-Horvat DE, Stein JF (1990) Neuronal activity in the lateral cerebellum of trained

monkeys, related to visual stimuli or to eye movements. J Physiol 428:595–614.

McElligott JG, Keller EL (1984) Cerebellar vermis involvement in monkey saccadic eye

movements: microstimulation. Exp Neurol 86:543–558.

Miall RC, Weir DJ, Wolpert DM, Stein JF (1993) Is the cerebellum a smith predictor? J

Mot Behav 25:203–216.

Mitchell JP, Macrae CN, Gilchrist ID (2002) Working memory and the suppression of

reflexive saccades. J Cogn Neurosci 14:95–103.

Mokler A, Fischer B (1999) The recognition and correction of involuntary prosaccades in

an antisaccade task. Exp Brain Res 125:511–516.

Munoz DP, Everling S (2004) Look away: the anti-saccade task and the voluntary control of

eye movement. Nat Rev Neurosci 5:218–228.

Noda H, Fujikado T (1987) Involvement of Purkinje cells in evoking saccadic eye

movements by microstimulation of the posterior cerebellar vermis of monkeys. J

Neurophysiol 57:1247–1261.

Ohki M, Kitazawa H, Hiramatsu T, Kaga K, Kitamura T, Yamada J, Nagao S (2009) Role

of primate cerebellar hemisphere in voluntary eye movement control revealed by lesion

effects. J Neurophysiol 101:934–947.

Peterburs J, Gajda K, Koch B, Schwarz M, Hoffmann K-P, Daum I, Bellebaum C (2012)

Cerebellar lesions alter performance monitoring on the antisaccade task--an event-related

potentials study. Neuropsychologia 50:379–389.

Piu P, Pretegiani E, Rosini F, Serchi V, Zaino D, Chiantini T, Rufa A (2019) The

cerebellum improves the precision of antisaccades by a latency-duration trade-off. Prog

Brain Res 249:125–139.

Ron S, Robinson DA (1973) Eye movements evoked by cerebellar stimulation in the alert

monkey. J Neurophysiol 36:1004–1022.

Schlag-Rey M, Amador N, Sanchez H, Schlag J (1997) Antisaccade performance predicted

by neuronal activity in the supplementary eye field. Nature 390:398–401.

Sendhilnathan N, Semework M, Goldberg ME, Ipata AE (2020) Neural Correlates of

Reinforcement Learning in Mid-lateral Cerebellum. Neuron 106:188–198.e5.

Soetedjo R, Kojima Y, Fuchs AF (2019) How cerebellar motor learning keeps saccades

accurate. J Neurophysiol 121:2153–2162.

Strick PL, Dum RP, Fiez JA (2009) Cerebellum and nonmotor function. Annu Rev

Neurosci 32:413–434.

Sun Z, Smilgin A, Junker M, Dicke PW, Thier P (2017) The same oculomotor vermal

Purkinje cells encode the different kinematics of saccades and of smooth pursuit eye

movements. Sci Rep 7:40613.

Tanaka H, Harada M, Arai M, Hirata K (2003) Cognitive dysfunction in cortical cerebellar

atrophy correlates with impairment of the inhibitory system. Neuropsychobiology 47:206–

211.

ten Brinke MM, Boele HJ, Spanke JK, Potters JW, Kornysheva K, Wulff P, Ac IJ,

Koekkoek SK, De Zeeuw CI (2015) Evolving Models of Pavlovian Conditioning:

Cerebellar Cortical Dynamics in Awake Behaving Mice. Cell Rep 13:1977–1988.

Thach WT (2007) On the mechanism of cerebellar contributions to cognition. Cerebellum

6:163–167.

Thier P, Dicke PW, Haas R, Thielert C-D, Catz N (2002) The role of the oculomotor vermis

in the control of saccadic eye movements. Ann N Y Acad Sci 978:50–62.

Thier P, Möck M (2006) The oculomotor role of the pontine nuclei and the nucleus

reticularis tegmenti pontis. Prog Brain Res 151:293–320.

Voogd J, Schraa-Tam CK, van der Geest JN, De Zeeuw CI (2012) Visuomotor cerebellum

in human and nonhuman primates. Cerebellum 11:392–410.

Wang X, Yu S-Y, Ren Z, De Zeeuw CI, Gao Z (2020) A FN-MdV pathway and its role in

cerebellar multimodular control of sensorimotor behavior. Nat Commun 11:6050.

Author contributions

CIDZ, AB, MF, NF and EA designed the study and analysis. EA and PR performed surgeries on the animals. EA and NF executed the experiments and EA, NF, AB and PJH analyzed the data. EA, NF, AB and CIDZ wrote the first draft. All authors edited the manuscript.

Data and code availability

The data that support the findings of this study and code for analysis of in vivo eye movement and Purkinje cell recordings are available from the corresponding author upon reasonable request.