Mechanism of a COVID-19 nanoparticle vaccine candidate that elicits a broadly

neutralizing antibody response to SARS-CoV-2 variants

Yi-Nan Zhang[1], Jennifer Paynter[1], Tatiana Fourfouris[1], Cindy Sou[1], Timothy Ngo[1], Linling He[1],

and Jiang Zhu[1, 2, *]

[1]Department of Integrative Structural and Computational Biology, [2]Department of Immunology

and Microbiology, The Scripps Research Institute, La Jolla, California 92037, USA

*Corresponding author

JZ: Phone +1 (858) 784-8157; Email: jiang@scripps.edu

KEYWORDS

Coronavirus disease 2019 (COVID-19); severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2); variant of concern (VOC); vaccine; lymph node; germinal center; broadly neutralizing antibody (bNAb).

ABSTRACT (150 words)

Vaccines against severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) are essential for combating the coronavirus disease 2019 (COVID-19) pandemic. Neutralizing antibody responses to the original Wuhan-Hu-1 strain that were generated during infection and vaccination showed lower effectiveness against variants of concern. Here, we demonstrated that mouse plasma induced by protein nanoparticles that present rationally designed S2GΔHR2 spikes can neutralize the B.1.1.7, B.1.351, and P.1 variants with comparable titers. The mechanism of nanoparticle vaccine-induced immunity was examined in mice for an I3-01v9 60-mer that presents 20 stabilized spikes. Compared with the soluble spike, this nanoparticle showed 6-fold longer retention, 4-fold greater presentation on follicular dendritic cell dendrites, and 5-fold higher germinal center reactions in lymph node follicles. Intact nanoparticles in lymph node tissues were visualized by transmission electron microscopy. In conclusion, spike-presenting protein nanoparticles that induce robust long-lived germinal centers may provide a vaccine solution for emerging SARS-CoV-2 variants.

ONE-SENTENCE SUMMARY (125 characters)

With prolonged lymph node retention and robust germinal centers, nanoparticles elicit neutralizing antibodies to diverse SARS-CoV-2 variants.

The COVID-19 pandemic has led to more than 124 million infection cases and 2.7 million deaths

globally. It has been reported that the human antibody responses to SARS-CoV-2 spike antigens

in COVID-19 patients can be sustained for several months after infection (*1-5*). However,

recently identified variants of concern (VOCs) exhibit higher transmissibility and resistance to

prior immunity as SARS-CoV-2 continues to adapt to the human host (*6, 7*). One such variant,

B.1.1.7, emerged from southeast England in October 2020 and accounted for two-thirds of new

infections in London in December 2020, with a higher transmission rate (43-90%) and risk of

mortality (32-104%) than previously circulating strains (*8-11*). Other variants, such as B.1.351

and P.1, also became prevalent in the Eastern Cape, Western Cape, and KwaZulu-Natal

provinces in South Africa and in Manaus, Brazil, respectively (*7, 12-14*). The rise of SARS-

CoV-2 VOCs and their rapid spread worldwide will likely result in more infection cases,

hospitalizations, and potentially more deaths, further straining healthcare resources (*14*).

To date, seven COVID-19 vaccines have been approved for emergency use in humans,

with more than 100 candidates being assessed in various phases of clinical trials (*15*). With the

exception of inactivated whole-virion vaccines, diverse platforms have been used to deliver the

recombinant SARS-CoV-2 spike, such as mRNA-encapsulating liposomes (e.g., BNT162b2 and

mRNA-1273), adenovirus vectors (e.g., ChAdOx1 nCoV-19 [AZD1222], CTII-nCoV, Sputnik

V, and Ad26.COV2.S), and nanoparticles (e.g., NVX-CoV2373). These vaccines demonstrated

65-96% efficacies in Phase 3 trials, with reduced morbidity and mortality associated with

COVID-19 disease (*16-20*). However, a notable loss of vaccine efficacy against new SARS-

CoV-2 variants was reported because of spike mutations in the receptor-binding domain (RBD;

e.g., K417N, E484K, and N501Y), N-terminal domain (NTD; e.g., L18F, D80A, D215G, and

Δ242-244), and other regions that are critical to spike stability (e.g., D614G) (*7, 21-28*). Among

circulating VOCs, the B.1.351 lineage appeared to be most resistant to neutralization by

convalescent plasma (9.4-fold) and vaccine sera (10.3- to 12.4-fold) (*29*), while a lesser degree

of reduction was observed for an early variant, B.1.1.7 (*30, 31*). Based on these findings, it was

suggested that vaccines would need to be updated periodically to maintain protection against

rapidly evolving SARS-CoV-2 (*24, 32, 33*). This raises the concern that herd immunity may not

be achievable with current vaccines and highlights the necessity of developing vaccines that can

elicit a broadly neutralizing antibody (bNAb) response to SARS-CoV-2 variants (*24, 28*). As

previously reported (*34-40*), the production of a bNAb response relies on long-lived germinal

center (GC) reactions to activate precursor B cells, stimulate affinity maturation, and form long-

term immune memory. Antigen retention and presentation within lymph node follicles are key to

the induction of long-lived GC reactions (*34, 36*) and thus should be a main criterion in the

development of bNAb-producing vaccines.

We previously investigated the cause of SARS-CoV-2 spike metastability and rationally

designed the S2GΔHR2 spike, which was displayed on three self-assembling protein

nanoparticle (SApNP) platforms, including ferritin (FR) 24-mer and multilayered E2p and I3-

01v9 60-mers, as COVID-19 vaccine candidates (*41*) (**Fig. 1A**). Notably, the I3-01v9 SApNP

that presents 20 stabilized spikes induced a potent NAb response against both SARS-CoV-1 and

SARS-CoV-2, in addition to critically needed T cell responses (*41*). Here, we first examined the

neutralizing activity of mouse plasma induced by these rationally designed vaccine constructs

against representative SARS-CoV-2 variants. We then investigated how SApNPs "behave" in

lymph nodes and induce GCs in the mouse model by characterizing vaccine delivery and

immunological responses at the intraorgan, intracellular, and intercellular levels. Our findings

suggest that a spike-presenting protein nanoparticle vaccine may confer broad protection against

SARS-CoV-2 VOCs.

We first assessed the neutralizing activity of polyclonal plasma elicited by various spike

and SApNP vaccine formulations in pseudoparticle (pp) neutralization assays (*42*). Mouse

plasma at week 5 after intraperitoneal (i.p.) injections of adjuvanted vaccine antigens (50 μg)

from our previous study were analyzed against the original SARS-CoV-2 strain, Wuhan-Hu-1, as

a baseline for comparison (**Fig. 1B**). The soluble S2P$_{ECTO}$ spike elicited the lowest 50%

inhibitory dilution (ID$_{50}$) titers, whereas the soluble S2GΔHR2 spike induced a stronger NAb

response, showing a 7.1-fold higher (or 8.1 times) average ID$_{50}$ titer, which did not reach

statistical significance because of within-group variation. All three spike SApNP vaccines

elicited superior neutralizing antibody responses compared with the soluble S2P$_{ECTO}$ spike (*41,*

43). Notably, the I3-01v9 SApNP vaccine showed the most potent NAb response, with an

average ID$_{50}$ titer of 2090, which was 8.1-fold higher than the soluble S2P$_{ECTO}$ spike. Despite the

differences in ID$_{50}$ titers, the overall pattern remained the same as observed in our previous study

(*41*). These differences can be attributed to the inherent variation of pseudovirus neutralization

assays (*42*) (see Methods).Next, we assessed neutralizing activity against three major SARS-

CoV-2 variants (**Fig. 1C**, **fig. S1A, B**). The I3-01v9 SApNP induced a stronger NAb response

against three VOCs, with 0.5-fold (B.1.1.7), 0.8-fold (B.1.351), and 1.8-fold (P.1) higher ID$_{50}$

titers compared with the Wuhan-Hu-1 strain (**Fig. 1C**). These results were confirmed in a second

experiment, in which mice were intraperitoneally immunized with low doses (5 and 15 μg) of I3-

01v9 SApNP. Remarkably, all three SARS-CoV-2 VOC-pps were neutralized by mouse plasma,

with ID$_{50}$ titers comparable to the high-dose group (45 μg; **fig. S1C, D**). To examine whether

routes of injection affect the elicitation of NAb responses against VOCs, we performed a third

experiment, in which a low dose (3.3 μg) of adjuvanted antigen was intradermally administered

into four footpads (i.e., 0.8 μg/footpad). The large (~55-60 nm) E2p and I3-01v9 SApNPs that

present 20 S2GΔHR2 spikes yielded higher ID_{50} titers than the soluble S2GΔHR2 spike (**Fig.**

1D, **fig. S1E, F**), whereas a significant reduction of neutralizing titers against the variants was

observed for mouse plasma from the S2GΔHR2 group (**Fig. 1E**, **fig. S1E, F**), suggesting that

nanoparticle display is critical for eliciting a bNAb response. The E2p and I3-01v9 SApNP

groups exhibited comparable or stronger NAb responses against three variants relative to the

original Wuhan-Hu-1 strain (**Fig. 1E**). A panel of human NAbs was previously used to validate

the SARS-CoV-2-pp neutralization assays, in addition to evaluating the antigenicity of rationally

designed spikes and SApNPs (*41*). This panel of human NAbs was tested against SARS-CoV-2-

pps that carry various VOC spikes (**Fig. 1F**, **fig. S1G**). Lower potency, measured by the 50%

inhibitory concentration (IC_{50}), was observed for all human NAbs against B.1.351 and P.1

variants, with the exception of NAb S309, which was identified from a SARS-CoV-1 patient

(*44*). This finding is consistent with previous reports on convalescent patient plasma (*29-31*). As

a negative control, mouse plasma induced by the S2GΔHR2-presenting I3-01v9 SApNP was

tested against pseudoviruses carrying the murine leukemia virus (MLV) Env, or MLV-pps.

Nonspecific neutralization was not detected (**fig. S1H, I**). Altogether, these results demonstrated

a clear advantage of spike-presenting SApNPs in eliciting a strong bNAb response against

SARS-CoV-2 VOCs compared with soluble spikes.

Next, we studied *in vivo* behaviors of the S2GΔHR2 spike and two large SApNPs to

understand vaccine-induced immunity and why nanoparticles outperform soluble spikes in terms

of bNAb elicitation. We first examined the distribution of S2GΔHR2-presenting I3-01v9

SApNPs in mouse lymph nodes via footpad injection (10 μg). The mice were sacrificed 12 h

after single-dose (**Fig. 2A**) and prime-boost (**Fig. 2B**) regimens. The axillary, brachial, and

popliteal sentinel lymph nodes were isolated for histological analysis. The lymph node tissues

were stained with the human anti-spike antibody P2B-2F6 (*45*) to characterize SARS-CoV-2

spikes presented on the I3-01v9 SApNPs. Consistent with our previous study (*46*), SApNPs

accumulated in lymph node follicles, regardless of the number of doses. SApNPs were

sequestrated in the center of lymph node follicles after a single-dose (**Fig. 2A,** images on the left,

schematics on the right), but were located along the outer layer of expanded lymph node follicles

after the second injection due to preexisting humoral immunity (i.e., GC reactions) that was

induced by the first dose (**Fig. 2B**, images on the left, schematics on the right). Overall, the

majority of SApNPs accumulated in lymph node follicles, while their distribution differed

slightly depending on the doses.

In this context, we examined patterns of trafficking and lymph node follicle retention for

soluble S2GΔHR2 spike *vs.* the S2GΔHR2-presenting E2p and I3-01v9 SApNPs. To facilitate

this analysis, the mice were sacrificed 2 h to 8 weeks after a single dose (**Fig. 2C**) and 2 h to 5

weeks after the boost (**Fig. 2D**). The injection dose was normalized to the total amount of protein

(10 μg) per injection into each footpad (40 μg/mouse). As shown in **Fig. 2C**, the S2GΔHR2

spikes that trafficked into lymph node follicles at 2 h cleared within 48 h. In contrast, the two

large SApNPs accumulated in the subcapsular sinus at 2 h and then trafficked into follicles 12 h

after the single-dose injection. Remarkably, I3-01v9 SApNPs remained detectable in lymph node

follicles after 2 weeks, suggesting 6-fold longer retention than the S2GΔHR2 spike (**Fig. 2C**).

The results for these protein nanoparticles are thus consistent with the pattern of size dependency

that was observed for ovalbumin-conjugated gold nanoparticles in our previous study (*46*), in

which small (5-15 nm) nanoparticles cleared shortly after the injection, whereas large (50-100

nm) nanoparticles were retained in lymph node follicles for weeks. Similar patterns of antigen

retention were observed after the second injection, although the boost appeared to exert a more

positive effect on the soluble spike, which could be detected in lymph node follicles at 48 h (**Fig.**

2D). Nonetheless, prolonged retention was observed for both E2p and I3-01v9 SApNPs 2 weeks

after the boost injection. Overall, the multivalent display of S2GΔHR2 spikes on the I3-01v9

SApNP resulted in 325- and 4-fold greater accumulation in lymph node follicles compared with

the soluble spike 48 h after the single-dose **(Fig. 2E)** and prime-boost **(Fig. 2F)** injections,

respectively. These findings reveal the advantage of a leading vaccine candidate identified in our

previous study, S2GΔHR2-10GS-I3-01v9-L7P (*41*), in terms of spike antigen retention in lymph

node follicles.

Next, we determined the involvement of resident cells in antigen retention in lymph node

follicles. Recent studies reported that follicular dendritic cells (FDCs) are resident stromal cells

in follicles and can retain immune complexes, virus-like particles, virus, and bacteria (*46-51*).

Follicular dendritic cells are key to naive antigen retention, GC initiation and maintenance, and B

cell affinity maturation (*39, 52-55*). Here, we hypothesized that FDCs comprise the major cell

population in lymph node follicles that retain SARS-CoV-2 spikes and spike-presenting

SApNPs. To test this hypothesis, we administered vaccines via footpad injections and collected

mouse lymph nodes at the peak of accumulation (12 h) after single-dose (**Fig. 3A**) and prime-

boost (**Fig. 3B**) injections. Lymph node tissue samples were stained with the anti-spike antibody

P2B-2F6 (*45*) for the spike antigen and anti-CD21 and CD169 antibodies for FDCs and

subcapsular sinus macrophages, respectively. The S2GΔHR2 spike and SApNP (E2p or I3-01v9)

signals colocalized with FDC ($CD21^+$) networks in lymph node follicles (**Fig. 3A, B**). This result

confirmed the critical role of FDC networks in mediating vaccine retention in lymph node

follicles.

The induction of potent bNAb responses by spike-presenting SApNPs (**Fig. 1**) suggests

the effective activation of naïve B cells and subsequent recalls by crosslinking B cell receptors

(*56-58*). However, still unclear are the manners in which FDC networks present SApNPs to B

cells to trigger such a bNAb response. Here, we analyzed the interface between FDC networks

and B cells by transmission electron microscopy (TEM). Briefly, fresh lymph nodes were

isolated and directly immersed in fixative. The processed tissue samples were sectioned and

stained on copper grids for TEM analysis. We first determined whether SApNPs, such as the

S2GΔHR2-presenting I3-01v9 SApNP, remain intact *in vivo* (**fig. S2**). Mouse lymph nodes were

isolated 2 h after the injection of a high dose (50 μg) of the non-adjuvanted I3-01v9 SApNP. The

TEM images revealed that round-shape granules corresponding to intact SApNP aligned on the

macrophage surface or inside endolysosomes of the macrophage in a lymph node (**fig. S2**). We

next studied the relative location between FDCs and I3-01v9 SApNPs and how FDCs present

SApNPs to B cells. Mouse lymph nodes were collected 2, 12, and 48 h after a single-dose (50

μg) and 12 h after the boost of the I3-01v9 SApNP vaccine. The FDCs exhibited the

characteristic morphology of long dendrites that surrounded and interacted with B cells in lymph

node follicles (**Fig. 3C**, **fig. S3**). While few I3-01v9 SApNPs were observed on FDC dendrites at

2 h (**fig. S3D**), notably more nanoparticles migrated to and aligned on FDC dendrites at 12 and

48 h (**Fig. 3C**, **figs. S3A-C**, yellow arrows). The TEM images indicated that FDCs can present

many SApNPs to neighboring B cells in this "hugging mode", in which their long dendrites

brace B cells to maximize interactions between multivalently displayed spikes and B cell

receptors. These results demonstrated the intrinsic nature of FDCs as a reservoir for the

sequestration, retention, and presentation of virus-like particles, or SApNPs with similar

molecular traits, to initiate GC reactions.

Lastly, we investigated whether the prolonged retention of S2GΔHR2-presenting E2p and

I3-01v9 SApNPs induce more robust GCs in lymph node follicles than the soluble S2GΔHR2

spike. Immunohistological analysis was performed to characterize GC B cells (GL7$^+$) and T

follicular helper (T$_{fh}$) cells (CD4$^+$Bcl6$^+$). For the I3-01v9 SApNP, 2 weeks after immunization,

we observed robust GCs in lymph node B cell follicles (B220$^+$) with well-formed dark zone

(DZ) and light zone (LZ) compartments, which contain GC B cells, FDCs, and T$_{fh}$ cells (*35, 37,*

59-62) (**Fig. 4A**). We then extended the analysis to the S2GΔHR2 spike and spike-presenting

SApNPs 2, 5, and 8 weeks after the single-dose injection (**Fig. 4B, fig. S4A-C**) and 2 and 5

weeks after the boost (**Fig. 4C, fig. S4D, E**). Two semi-quantitative metrics, the GC/FDC ratio

(i.e., whether GC formation is associated with an FDC network, %) and GC size (i.e., occupied

area), were used in this analysis. Overall, the soluble S2GΔHR2 spike and both large SApNPs

induced robust GCs 2 weeks after immunization (**Fig. 4B, fig. S4A**). The E2p and I3-01v9

SApNPs that present 20 S2GΔHR2 spikes induced robust, long-lived GCs, whereas the spike

alone failed to sustain robust GCs at week 8 with either the single-dose (**Fig. 4B, D**) or prime-

boost (**Fig. 4C, E**) injections. The I3-01v9 SApNP generated larger GCs than the soluble spike,

2.0-fold larger after the single-dose (**Fig. 4B, D**) and 2.4-fold larger after the boost (**Fig. 4C, E**),

measured at week 8.

We further characterized GC reactions by flow cytometry. Fresh mouse lymph nodes

were disaggregated into a single cell suspension and stained with an antibody cocktail to quantify

GC B cells and T$_{fh}$ cells (**fig. S5A**). The results were consistent with the immunohistological

analysis, in which all spike-based vaccine antigens, including the S2GΔHR2 spike and SApNPs, showed robust GCs at week 2 after the injection that declined over time, as measured at weeks 5 and 8 (**Fig. 4F**). The E2p and I3-01v9 SApNPs generated a larger population of GC B cells than both the S2P$_{ECTO}$ and S2GΔHR2 spikes at week 2 (**fig. S5B, C**). Although the boost dose had little impact on the frequency of GC B cells and T$_{fh}$ cells, it appeared to extend GC formation within lymph nodes (**Fig. 4F, G**), which may promote B cell development toward bNAbs. Notably, the GC B cell and T$_{fh}$ cell populations elicited by the soluble S2GΔHR2 spike were barely detectable 5 weeks after immunization (**Figs. 4F, G**). This result was reminiscent of a recent study of an mRNA vaccine, in which GC reactions diminished to baseline levels at week 4 after a single-dose injection (*63*). The S2GΔHR2-presenting I3-01v9 SApNP generated 3.7/5.2-fold more GC B cells and 3.7/4.4-fold more T$_{fh}$ cells than the soluble S2GΔHR2 spike after one/two-dose immunization at week 8 (**Fig. 4G**). Therefore, SApNPs that were retained on FDC dendrites could present SARS-CoV-2 neutralizing epitopes to enable more effective B cell recognition than the soluble spike, and consequently induce more robust and long-lived GCs in lymph nodes.

To end the COVID-19 pandemic, vaccines need to effectively block current and emerging SARS-Cov-2 variants that evade NAb responses by mutating key epitopes on the viral spike (*24*). Such vaccines must induce long-lasting GCs to facilitate the development of bNAbs. Effective vaccine retention and presentation are critical for inducing and sustaining GC reactions, which in turn promote the proliferation and affinity maturation of antigen-specific B cells. Here, we found that the S2GΔHR2-presenting I3-01v9 SApNP, a vaccine candidate from our recent study (*41*), elicited 6-fold longer retention and 4-fold greater accumulation in lymph node follicles than the stabilized S2GΔHR2 spike alone with a prime-boost regimen (**Fig. 2D, F**).

This can be attributed to the intrinsic physiological properties of lymph nodes that mediate

vaccine trafficking and retention in follicles in a size-dependent manner, which would favor

retaining large (> 50 nm) virus-like nanoparticles (*46-48, 64, 65*). Supporting this notion are the

images of retained I3-01v9 SApNPs aligned on long FDC dendrites, suggesting that such protein

nanoparticles can present spike antigens to B cells for rapid initiation and then sustain GC

reactions in lymph node follicles for an extended period of time (**Fig. 3, 4**). Specifically, the

S2GΔHR2-presenting I3-01v9 SApNP generated 2.4-fold larger GCs and greater numbers of GC

B cells (5.2-fold) and T_{fh} cells (4.4-fold) than the soluble S2GΔHR2 spike with the prime-boost

regimen (**Fig. 4**). The findings in this study provide quantitative evidence that spike-presenting

SApNP vaccines are uniquely suited for inducing long-lived robust GCs in lymph node follicles.

Our analyses also provide potential explanations for the mechanism by which the S2GΔHR2-

presenting I3-01v9 SApNP can elicit a more effective bNAb response than the soluble spike (**Fig.

1B)** (*32*). Further investigation at the monoclonal level will reveal how bNAbs recognize

conserved epitopes and facilitate the rational design of SARS-CoV-2 spike antigens (*24*).

Superior NAb responses have been reported for other COVID-19 vaccine candidates that take

advantage of particulate display (*66-78*). Importantly, our S2GΔHR2-presenting I3-01v9 SApNP

vaccine elicited bNAb responses to three SARS-CoV-2 VOCs (**Fig. 1**), thus meeting a major

criterion for next-generation COVID-19 vaccines.

Protein vaccines have well-established records of both safety and effectiveness (*79-82*),

but they have yet to be deployed in the global campaign against the COVID-19 pandemic.

Through formulation with adjuvants to further enhance immune responses (*83-88*), protein

vaccines can be used either alone or as a booster for currently used nucleic acid vaccines. One

such protein vaccine, NVX-CoV2373 (full-length S2P spikes formulated with Matrix-M^{TM}

adjuvant), showed 96.4% efficacy in human trials (*89*). In our studies, we formulated the spike-

presenting E2p and I3-01v9 SApNPs with the AddaVax and aluminum phosphate adjuvants (*41*),

which induced stronger GC reactions than non-adjuvanted SApNPs at week 2 (**fig. S6**). Further

development of these SApNP vaccines may benefit from a systematic analysis of adjuvants that

target diverse immune pathways. An in-depth understanding of how different vaccine platforms

(e.g., inactivated virions, mRNAs, viral vectors, and nanoparticles) behave *in vivo*, such as

trafficking and retention in lymph nodes, interactions with immune cells, antigen processing and

presentation, and GC reactions and NAb responses, will provide insights into vaccine-induced

immunity (*82, 90-92*). This knowledge will in turn accelerate vaccine development against

SARS-CoV-2 and other emerging pathogens.

MATERIALS AND METHODS

SARS-CoV-2 spike and SApNP vaccine antigens

The design, expression, and purification of a stabilized SARS-CoV-2 spike, S2GΔHR2, and

three SApNPs that present either 8 or 20 S2GΔHR2 spikes were described in our recent study

(*41*). Briefly, the spike gene of the SARS-CoV-2 isolate Wuhan-Hu-1 (GenBank accession no.

MN908947) was modified to include the mutations ^{682}GSAGSV^{687} and K986G/V987G, in

addition to truncation of the HR2 stalk (ΔE1150-Q1208). The viral capsid protein SHP (Protein

Data Bank: 1TD0) was added as a C-terminal trimerization motif to stabilize the S2GΔHR2

trimer, resulting in a soluble S2GΔHR2-5GS-1TD0 spike (*41*). The S2GΔHR2 spike was

genetically fused to FR, multilayered E2p, and multilayered I3-01v9 with 5GS, 5GS, and 10GS

linkers, respectively, resulting in three S2GΔHR2-presenting SApNPs (*41*). An S2P$_{ECTO}$-5GS-

1TD0 spike construct that contained the mutations ^{682}GSAGSV^{687} and K986G/V987G but

without HR2 deletion (*41)* was included for comparison. All of the vaccine antigens were

transiently expressed in ExpiCHO cells and purified by a CR3022 antibody column and size-

exclusion chromatography (SEC) on a Superose 6 10/300 GL column as described previously

(*41*).

Animal immunization and sample collection

Similar immunization protocols were reported in our previous vaccine studies (*41, 93-95*).

Briefly, Institutional Animal Care and Use Committee (IACUC) guidelines were followed for all

of the animal studies. BALB/c mice (6 weeks old) were purchased from The Jackson Laboratory

and kept in ventilated cages in environmentally controlled rooms at The Scripps Research

Institute. The mouse studies were conducted according to Association for the Assessment and

Accreditation of Laboratory Animal Care guidelines, and the protocols were approved by the

IACUC. For the immunogenicity study, the mice were intraperitoneally immunized at either

weeks 0 or 3 with 200 μl of antigen/adjuvant mix containing 5-50 μg of vaccine antigen and 100

μl of adjuvant (*41*) or immunized at weeks 0 and 3 with 20 μl of antigen/adjuvant mix containing

0.8 μg of vaccine antigen per injection (3.3 μg/mouse) and 10 μl of adjuvant per injection (four

footpads were injected with a total of 80 μl [20 μl/footpad]) via intradermal footpad injections.

For the mechanism study of vaccine trafficking, retention, and induced GCs, the mice were

immunized at weeks 0 and 3 with 20 μl of antigen/adjuvant mix containing 10 μg of vaccine

antigen and 10 μl of adjuvant per injection. To visualize S2GΔHR2-presenting I3-01v9 SApNPs

in lymph node tissues using TEM, the mice were immunized at weeks 0 and 3 with 70 μl of

antigen/adjuvant mix containing 50 μg of vaccine antigen and 20 μl of adjuvant per injection.

Vaccines were administered into intradermal footpads of mice using a 29-gauge insulin needle

under 3% isoflurane anesthesia with oxygen. Blood was drawn from the maxillary/facial vein

into an ethylenediaminetetraacetic acid (EDTA)-coated tube 2 weeks after each immunization.

Blood plasma was isolated after centrifugation at 14000 rotations per minute (rpm) for 10 min.

Plasma was heated and inactivated at 56°C for 30 min. The supernatant was then collected after

centrifugation at 8000 rpm for 10 min. Plasma was used for the neutralization assay to determine

neutralizing antibody responses. The axillary, brachial, and popliteal sentinel lymph nodes were

collected at the end timepoint for further analysis.

SARS-CoV-2 pseudovirus neutralization assay

The SARS-CoV-2-pp neutralization assays were described in our previous study (*41*). Briefly,

SARS-CoV-2-pps were generated by the co-transfection of HEK293T cells with the HIV-1

pNL4-3.lucR-E- plasmid (obtained from the National Institutes of Health AIDS reagent program;

https://www.aidsreagent.org/) and the expression plasmid encoding the S gene of various SARS-

CoV-2 strains, including three variants: B.1.1.7, B.1.351, and P.1 (GISAID accession no.

EPI_ISL_601443, EPI_ISL_678597, and EPI_ISL_792680, respectively). The HEK293T-

hACE2 cell line (catalog no. NR-52511) and pcDNA3.1(-) vector containing the S gene of the

SARS-CoV-2 isolate Wuhan-Hu-1 (catalog no. NR52420) were requested from the BEI

Resources (https://www.beiresources.org/) on September 23, 2020 and used in the pseudovirus

neutralization assays (*42*). Based on sequence alignment, spike mutations were incorporated into

the S gene of the original Wuhan-Hu-1 isolate (catalog no. NR52420) to create respective

expression plasmids for B.1.1.7, B.1.351, and P.1. SARS-CoV-2-pp neutralization by immunized

mouse plasma and human monoclonal antibodies (mAbs) was performed according to a

previously described protocol (*41*). Using the same co-transfection expression system as

described above for the SARS-CoV-2-pps, we produced pseudoviruses carrying the murine

leukemia virus (MLV) Env, MLV-pps, for use as a negative control (*41*). Percent neutralization

data were analyzed using GraphPad Prism 9.0.2 software. ID_{50}/IC_{50} values were calculated using

constraints for the percent neutralization (0-100%), whereas unconstrained neutralization plots

were shown in Fig 1 and fig S1.

Histology, immunostaining, and imaging

The mice were sacrificed 2 h to 8 weeks after a single-dose and 2 h to 5 weeks after the boost

immunization. The axillary, brachial, and popliteal sentinel lymph nodes were isolated for

histological analysis. Fresh lymph nodes were rapidly merged into frozen section compound

(VWR International, catalog no. 95057-838) in a plastic cryomold (Tissue-Tek at VWR, catalog

no. 4565) using liquid nitrogen to preserve the antigens on the cell membrane and spike. Lymph

node samples were stored at -80°C and sent to the Centre for Phenogenomics on dry ice for

further sample processing. Tissue sections (8 μm) were cut on a cryostat (Cryostar NX70) and

collected on charged slides. Sections were post-fixed in 10% neutral buffered formalin and

permeabilized in phosphate-buffered saline containing 0.5% Triton X-100 before

immunostaining. Protein Block (Agilent) was used to block nonspecific antibody binding before

incubating the sections with primary antibody overnight at 4°C. After washing in TBST, the

sections were incubated in fluorophore-conjugated secondary antibodies for 1 h at room

temperature. Lymph node tissue sections were stained with human anti-spike antibody P2B-2F6

(*45*) (1:50) and biotinylated goat anti-human secondary antibody (Abcam, catalog no. ab7152,

1:300), followed by streptavidin-horseradish peroxidase reagent (Vectastain Elite ABC-HRP Kit,

Vector, catalog no. PK-6100) then DAB (ImmPACT DAB, Vector, catalog no. SK-4105) to

study the distribution and retention of the S2GΔHR2 spike alone and S2GΔHR2-presenting E2p

and I3-01v9 SApNPs. For immunofluorescent staining, tissue sections were stained for FDCs

using anti-CD21 antibody (Abcam, catalog no. ab75985, 1:1800) followed by anti-rabbit

secondary antibody conjugated with Alexa Fluor 555 (Thermo Fisher, catalog no. A21428;

1:200), for B cells using anti-B220 antibody (eBioscience, catalog no. 14-0452-82, 1:100)

followed by anti-rat secondary antibody conjugated with Alexa Fluor 674 (Thermo Fisher,

catalog no. A21247; 1:200), and for subcapsular sinus macrophages using anti-sialoadhesin

(CD169) antibody (Abcam, catalog no. ab53443, 1:600) followed by anti-rat secondary antibody

conjugated with Alexa Fluor 488 (Abcam, catalog no. ab150165; 1:200). Germinal center B cells

were labeled using rat anti-GL7 antibody (FITC; BioLegend, catalog no. 144604, 1:250). T

Follicular helper cells were labeled using CD4 antibody (Biolegend, catalog no. 100402, 1:100)

followed by anti-rat secondary antibody conjugated with Alexa Fluor 488 (Abcam, catalog no.

ab150165; 1:1000) and Bcl6 antibody (Abcam, catalog no. ab220092, 1:300) followed by anti-

rabbit secondary antibody conjugated with Alexa Fluor 555 (Thermo Fisher, catalog no. A21428;

1:1000). Nuclei were then counterstained with DAPI (Sigma-Aldrich, catalog no. D9542, 100

ng/ml). The stained tissue sections were scanned using an Olympus VS-120 slide scanner and

imaged using a Hamamatsu ORCA-R2 C10600 digital camera for all bright-field and fluorescent

images. The bright-field images of stained S2GΔHR2 spike and S2GΔHR2-presenting SApNPs

in lymph node follicles and fluorescent images of GCs were quantified using ImageJ software

(National Institutes of Health) (*96*).

Electron microscopy analysis of protein nanoparticles and lymph node tissues

Electron microscopy (EM) analysis was performed by the Core Microscopy Facility at The

Scripps Research Institute. For negative-staining EM analysis of protein nanoparticles, the

S2GΔHR2-10GS-I3-01v9-L7P SApNP samples were prepared at the concentration of 0.01

mg/ml. Carbon-coated copper grids (400 mesh) were glow-discharged and 10 µL of each sample

was adsorbed for 2 min. Excess sample was wicked away and grids were negatively stained with

2% uranyl formate for 2 min. Excess stain was wicked away and the grids were allowed to dry.

For EM analysis of mouse tissues, the lymph nodes were dissected from each animal and

immersed in oxygenated 2.5% glutaraldehyde and 4% paraformaldehyde in 0.1M Na cacodylate

buffer (pH 7.4) fixative overnight at 4ºC (*97*). After washing in 0.1 M sodium cacodylate buffer,

the tissue samples were post-fixed in buffered 1% osmium tetroxide and 1.5% potassium

ferrocyanide for 1-1.5 h at 4°C, rinsed in the same buffer, and then stained *en bloc* with 0.5%

uranyl acetate overnight at 4°C. The tissue samples were washed in double-distilled H_2O and

dehydrated through a graded series of ethanol followed by acetone, infiltrated with LX-112

(Ladd) epoxy resin, and polymerized at 60°C. Ultrathin lymph node sections (at 70-nm

thickness) were prepared for imaging. Samples were analyzed at 80 kV with a Talos L120C

transmission electron microscope (Thermo Fisher) and images were acquired with a CETA 16M

CMOS camera.

Lymph node disaggregation, cell staining, and flow cytometry

Germinal center reactions, including the percentage of GC B cells (GL7[+]B220[+]) and T follicular

helper cells (CD3[+]CD4[+]CXCR5[+]PD1[+]), and the number of GC B cells and T follicular helper

cells were studied by flow cytometry (**fig. S5A**). The mice were sacrificed 2, 5, and 8 weeks after

a single-dose and 2 and 5 weeks after the boost immunization. Fresh axillary, brachial, and

popliteal sentinel lymph nodes were collected and mechanically disaggregated. These lymph

node samples were merged in enzyme digestion solution containing 958 µl of Hanks' balanced

salt solution (HBSS) buffer (Thermo Fisher Scientific, catalog no. 14185052), 40 µl of 10 mg/ml

collagenase IV (Sigma-Aldrich, catalog no. C5138), and 2 μl of 10 mg/ml of DNase (Roche,

catalog no. 10104159001) in an Eppendorf tube. After incubation at 37°C for 30 min, lymph

node samples were filtered through a 70 μm cell strainer and spun down at $400 \times g$ for 10 min.

The supernatant was discarded, and the cell pellet was resuspended in HBSS blocking solution

containing 0.5% (w/v) bovine serum albumin and 2 mM EDTA. The nonspecific binding of Fc

receptors was blocked using anti-CD16/32 antibody (BioLegend, catalog no. 101302) on ice for

30 min. Cocktail antibodies, Zombie NIR live/dead stain (BioLegend, catalog no. 423106),

Brilliant Violet 510 anti-mouse/human CD45R/B220 antibody (BioLegend, catalog no. 103247),

FITC anti-mouse CD3 antibody (BioLegend, catalog no. 100204), Alexa Fluor 700 anti-mouse

CD4 antibody (BioLegend, catalog no. 100536), PE anti-mouse/human GL7 antibody

(BioLegend, catalog no. 144608), Brilliant Violet 605 anti-mouse CD95 (Fas) antibody

(BioLegend, catalog no. 152612), Brilliant Violet 421 anti-mouse CD185 (CXCR5) antibody

(BioLegend, catalog no. 145511), and PE/Cyanine7 anti-mouse CD279 (PD-1) antibody

(BioLegend, catalog no. 135216) were then mixed with the cells and placed on ice for 30 min.

After washing the cell with HBSS blocking solution after antibody staining, the samples were

fixed using 1.6% paraformaldehyde (Thermo Fisher Scientific, catalog no. 28906) in HBSS on

ice for 30 min. The cell samples were stored in HBSS blocking solution for the flow cytometry

study. Sample events were acquired by a 5-laser BD Biosciences LSR II analytical flow

cytometer with BD FACS Diva 6 software at the Core Facility of The Scripps Research Institute.

The data were further processed using FlowJo 10 software.

Statistical analysis

Data were collected from 4-7 mice per group. All of the statistical analyses were performed and

graphs were generated using GraphPad Prism 6.01 software. For the antibody analysis,

comparisons between different vaccine groups were performed using two-tailed unpaired

Student's *t*-test. Comparisons of neutralizing antibody titers against SARS-CoV-2 variants were

performed using the same plasma samples and analyzed using two-tailed paired Student's *t*-test.

For the vaccine accumulation and GC study, comparisons between different vaccine groups were

performed using one-way analysis of variance (ANOVA) followed by Tukey's multiple

comparison *post hoc* test. Statistical significance was determined as ns: not significant, $*p <$

0.05, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$.

SUPPLEMENTARY MATERIALS

Supplementary material for this article is available at http://xxx/xxx/xxx.

fig. S1. Spike and spike-presenting SApNP vaccine-induced neutralizing antibody responses

against SARS-CoV-2 variants of concern (VOCs).

fig. S2. SARS-CoV-2 spike-presenting SApNP interaction with macrophages in a lymph node.

fig. S3. TEM images of SARS-CoV-2 spike-presenting I3-01v9 SApNP interaction with FDCs in

a lymph node.

fig. S4. Immunohistological analysis of SARS-CoV-2 spike/spike-presenting SApNP vaccine-

induced GCs.

fig. S5. Flow cytometry analysis of SARS-CoV-2 spike/spike-presenting SApNP vaccine-

induced GCs.

fig. S6. Adjuvant effect on SARS-CoV-2 spike/spike-presenting SApNP vaccine-induced GCs.

References

1. J. M. Dan *et al.*, Immunological memory to SARS-CoV-2 assessed for up to 8 months after infection. *Science* **371**, eabf4063 (2021).
2. B. Isho *et al.*, Persistence of serum and saliva antibody responses to SARS-CoV-2 spike antigens in COVID-19 patients. *Sci. Immunol.* **5**, eabe5511 (2020).
3. A. S. Iyer *et al.*, Persistence and decay of human antibody responses to the receptor binding domain of SARS-CoV-2 spike protein in COVID-19 patients. *Sci. Immunol.* **5**, eabe0367 (2020).
4. Y. Chen *et al.*, Quick COVID-19 healers sustain anti-SARS-CoV-2 antibody production. *Cell* **183**, 1496-1507.e1416 (2020).
5. S. Marot *et al.*, Rapid decline of neutralizing antibodies against SARS-CoV-2 among infected healthcare workers. *Nat. Commun.* **12**, 844 (2021).
6. R. Burioni, E. J. Topol, Assessing the human immune response to SARS-CoV-2 variants. *Nat. Med.*, Published Ahead-of-Print (2021).
7. H. Tegally *et al.*, Emergence of a SARS-CoV-2 variant of concern with mutations in spike glycoprotein. *Nature*, Published Ahead-of-Print (2021).
8. R. Challen *et al.*, Risk of mortality in patients infected with SARS-CoV-2 variant of concern 202012/1: matched cohort study. *BMJ* **372**, n579 (2021).
9. T. Kirby, New variant of SARS-CoV-2 in UK causes surge of COVID-19. *Lancet Respir. Med.* **9**, e20-e21 (2021).
10. N. G. Davies *et al.*, Estimated transmissibility and impact of SARS-CoV-2 lineage B.1.1.7 in England. *Science*, eabg3055 (2021).
11. N. L. Washington *et al.*, Genomic epidemiology identifies emergence and rapid transmission of SARS-CoV-2 B.1.1.7 in the United States. *medRxiv*, 2021.2002.2006.21251159 (2021).
12. E. C. Sabino *et al.*, Resurgence of COVID-19 in Manaus, Brazil, despite high seroprevalence. *Lancet* **397**, 452-455 (2021).
13. N. R. Faria *et al.*, Genomics and epidemiology of a novel SARS-CoV-2 lineage in Manaus, Brazil. *medRxiv*, 2021.2002.2026.21252554 (2021).
14. J. R. Mascola, B. S. Graham, A. S. Fauci, SARS-CoV-2 viral variants—tackling a moving target. *JAMA*, Published Ahead-of-Print (2021).
15. C. Zimmer, J. Corum, W. S.-L., in *Coronavirus vaccine tracker*. (https://www.nytimes.com/interactive/2020/science/coronavirus-vaccine-tracker.html).
16. N. Dagan *et al.*, BNT162b2 mRNA Covid-19 vaccine in a nationwide mass vaccination setting. *N. Engl. J. Med.*, Published Ahead-of-Print (2021).
17. L. R. Baden *et al.*, Efficacy and safety of the mRNA-1273 SARS-CoV-2 vaccine. *N. Engl. J. Med.* **384**, 403-416 (2020).
18. T. C. Williams, W. A. Burgers, SARS-CoV-2 evolution and vaccines: cause for concern? *Lancet Respir. Med.*, Published Ahead-of-Print (2021).
19. D. Y. Logunov *et al.*, Safety and efficacy of an rAd26 and rAd5 vector-based heterologous prime-boost COVID-19 vaccine: an interim analysis of a randomised controlled phase 3 trial in Russia. *Lancet* **397**, 671-681 (2021).
20. F. P. Polack *et al.*, Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine. *N. Engl. J. Med.* **383**, 2603-2615 (2020).

21.   Q. Li *et al.*, The impact of mutations in SARS-CoV-2 spike on viral infectivity and
antigenicity. *Cell* **182**, 1284-1294.e1289 (2020).
22.   C. Rees-Spear *et al.*, The impact of spike mutations on SARS-CoV-2 neutralization.
bioRxiv, 2021.2001.2015.426849 (2021).
23.   C. K. Wibmer *et al.*, SARS-CoV-2 501Y.V2 escapes neutralization by South African
COVID-19 donor plasma. *bioRxiv*, 2021.2001.2018.427166 (2021).
24.   D. R. Burton, E. J. Topol, Variant-proof vaccines - invest now for the next pandemic.
Nature **590**, 386-388 (2021).
25.   M. Voysey *et al.*, Safety and efficacy of the ChAdOx1 nCoV-19 vaccine (AZD1222)
against SARS-CoV-2: an interim analysis of four randomised controlled trials in Brazil,
South Africa, and the UK. *Lancet* **397**, 99-111 (2021).
26.   E. Andreano *et al.*, SARS-CoV-2 escape in vitro from a highly neutralizing COVID-19
convalescent plasma. *bioRxiv*, 2020.2012.2028.424451 (2020).
27.   M. Hoffmann *et al.*, SARS-CoV-2 variants B.1.351 and P.1 escape from neutralizing
antibodies. *Cell*, Published Ahead-of-Print (2021).
28.   W. F. Garcia-Beltran *et al.*, Multiple SARS-CoV-2 variants escape neutralization by
vaccine-induced humoral immunity. *Cell*, Published Ahead-of-Print (2021).
29.   P. Wang *et al.*, Antibody resistance of SARS-CoV-2 variants B.1.351 and B.1.1.7.
bioRxiv, 2021.2001.2025.428137 (2021).
30.   P. Supasa *et al.*, Reduced neutralization of SARS-CoV-2 B.1.1.7 variant by convalescent
and vaccine sera. *Cell*, Published Ahead-of-Print (2021).
31.   D. A. Collier *et al.*, SARS-CoV-2 B.1.1.7 sensitivity to mRNA vaccine-elicited,
convalescent and monoclonal antibodies. *medRxiv*, 2021.2001.2019.21249840 (2021).
32.   Z. Wang *et al.*, mRNA vaccine-elicited antibodies to SARS-CoV-2 and circulating
variants. *Nature*, Published Ahead-of-Print (2021).
33.   Q. Li *et al.*, No higher infectivity but immune escape of SARS-CoV-2 501Y.V2 variants.
Cell, Published Ahead-of-Print (2021).
34.   A. Singh, Eliciting B cell immunity against infectious diseases using nanovaccines. *Nat.*
Nanotechnol. **16**, 16-24 (2021).
35.   G. D. Victora, M. C. Nussenzweig, Germinal centers. *Annu. Rev. Immunol.* **30**, 429-457
(2012).
36.   R. Rappuoli, Glycoconjugate vaccines: principles and mechanisms. *Sci. Transl. Med.* **10**,
eaat4615 (2018).
37.   J. G. Cyster, C. D. C. Allen, B cell responses: cell Interaction dynamics and decisions.
Cell **177**, 524-540 (2019).
38.   M. McHeyzer-Williams, S. Okitsu, N. Wang, L. McHeyzer-Williams, Molecular
programming of B cell memory. *Nat. Rev. Immunol.* **12**, 24-34 (2012).
39.   M. Akkaya, K. Kwak, S. K. Pierce, B cell memory: building two walls of protection
against pathogens. *Nat. Rev. Immunol.* **20**, 229-238 (2020).
40.   N. S. De Silva, U. Klein, Dynamics of B cells in germinal centres. *Nat. Rev. Immunol.* **15**,
137-148 (2015).
41.   L. He *et al.*, Single-component, self-assembling, protien nanoparticles presenting the
receptor binding domain and stabilized spike as SARS-CoV-2 vaccine candidates. *Sci.*
Adv. **7**, eabf1591 (2021).
42.   K. H. D. Crawford *et al.*, Protocol and reagents for pseudotyping lentiviral particles with
SARS-CoV-2 spike protein for neutralization assays. *Viruses* **12**, 513 (2020).

43.   D. Wrapp *et al.*, Cryo-EM structure of the 2019-nCoV spike in the prefusion
541   conformation. *Science* **367**, 1260-1263 (2020).

542   44.   D. Pinto *et al.*, Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV
543   antibody. *Nature* **583**, 290-295 (2020).

544   45.   B. Ju *et al.*, Human neutralizing antibodies elicited by SARS-CoV-2 infection. *Nature*
545   **584**, 115-119 (2020).

546   46.   Y.-N. Zhang *et al.*, Nanoparticle size influences antigen retention and presentation in
547   lymph node follicles for humoral immunity. *Nano Lett.* **19**, 7226-7235 (2019).

548   47.   B. A. Heesters, R. C. Myers, M. C. Carroll, Follicular dendritic cells: dynamic antigen
549   libraries. *Nat. Rev. Immunol.* **14**, 495-504 (2014).

550   48.   J. G. Cyster, B cell follicles and antigen encounters of the third kind. *Nat. Immunol.* **11**,
551   989-996 (2010).

552   49.   F. D. Batista, N. E. Harwood, The who, how and where of antigen presentation to B cells.
553   *Nat. Rev. Immunol.* **9**, 15-27 (2009).

554   50.   M. Kuka, M. Iannacone, Viral subversion of B cell responses within secondary lymphoid
555   organs. *Nat. Rev. Immunol.* **18**, 255-265 (2018).

556   51.   R. Rappuoli, D. Serruto, Self-assembling nanoparticles usher in a new era of vaccine
557   design. *Cell* **176**, 1245-1247 (2019).

558   52.   C. D. C. Allen, T. Okada, J. G. Cyster, Germinal-center organization and cellular
559   dynamics. *Immunity* **27**, 190-202 (2007).

560   53.   C. D. C. Allen, J. G. Cyster, Follicular dendritic cell networks of primary follicles and
561   germinal centers: Phenotype and function. *Semin. Immunol.* **20**, 14-25 (2008).

562   54.   L. Mesin, J. Ersching, Gabriel D. Victora, Germinal center B cell dynamics. *Immunity* **45**,
563   471-482 (2016).

564   55.   N. Baumgarth, The shaping of a B cell pool maximally responsive to infections. *Annu.*
565   *Rev. Immunol.* **39**, (2021).

566   56.   J. López-Sagaseta, E. Malito, R. Rappuoli, M. J. Bottomley, Self-assembling protein
567   nanoparticles in the design of vaccines. *Comput. Struct. Biotechnol. J.* **14**, 58-68 (2016).

568   57.   D. J. Irvine, M. C. Hanson, K. Rakhra, T. Tokatlian, Synthetic nanoparticles for vaccines
569   and immunotherapy. *Chem. Rev.* **115**, 11109-11146 (2015).

570   58.   D. J. Irvine, M. A. Swartz, G. L. Szeto, Engineering synthetic vaccines using cues from
571   natural immunity. *Nat. Mater.* **12**, 978-990 (2013).

572   59.   S. Crotty, Follicular helper CD4 T cells (TFH). *Annu. Rev. Immunol.* **29**, 621-663 (2011).

573   60.   A K Szakal, a. M H Kosco, J. G. Tew, Microanatomy of lymphoid tissue during humoral
574   immune responses: structure function relationships. *Annu. Rev. Immunol.* **7**, 91-109
575   (1989).

576   61.   J. Merkenschlager *et al.*, Dynamic regulation of TFH selection during the germinal centre
577   reaction. *Nature* **591**, 458-463 (2021).

578   62.   Z. Shulman *et al.*, T follicular helper cell dynamics in germinal centers. *Science* **341**,
579   673-677 (2013).

580   63.   K. Lederer *et al.*, SARS-CoV-2 mRNA vaccines foster potent antigen-specific germinal
581   center responses associated with neutralizing antibody generation. *Immunity* **53**, 1281-
582   1295.e1285 (2020).

583   64.   T. Tokatlian *et al.*, Innate immune recognition of glycans targets HIV nanoparticle
584   immunogens to germinal centers. *Science* **363**, 649-654 (2019).

65. S. N. Mueller, S. Tian, J. M. DeSimone, Rapid and persistent delivery of antigen by
lymph node targeting PRINT nanoparticle vaccine carrier to promote humoral immunity.
Mol. Pharm. **12**, 1356-1365 (2015).
66. B. Zhang *et al.*, A platform incorporating trimeric antigens into self-assembling
nanoparticles reveals SARS-CoV-2-spike nanoparticles to elicit substantially higher
neutralizing responses than spike alone. *Sci. Rep.* **10**, 18149 (2020).
67. A. C. Walls *et al.*, Elicitation of potent neutralizing antibody responses by designed
protein nanoparticle vaccines for SARS-CoV-2. *Cell* **183**, 1367-1382.e1317 (2020).
68. A. E. Powell *et al.*, A single immunization with spike-functionalized ferritin vaccines
elicits neutralizing antibody responses against SARS-CoV-2 in mice. *ACS Cent. Sci.* **7**,
183-199 (2021).
69. P. J. M. Brouwer *et al.*, Two-component spike nanoparticle vaccine protects macaques
from SARS-CoV-2 infection. *Cell* **184**, 1188-1200.e1119 (2021).
70. K. S. Park *et al.*, Lipid-based vaccine nanoparticles for induction of humoral immune
responses against HIV-1 and SARS-CoV-2. *J. Control. Release* **330**, 529-539 (2021).
71. Y.-F. Kang *et al.*, Rapid development of SARS-CoV-2 spike protein receptor-binding
domain self-assembled nanoparticle vaccine candidates. *ACS Nano* **15**, 2738-2752
(2021).
72. J. H. Lam *et al.*, Next generation vaccine platform: polymersomes as stable nanocarriers
for a highly immunogenic and durable SARS-CoV-2 spike protein subunit vaccine.
bioRxiv, 2021.2001.2024.427729 (2021).
73. A. A. Cohen *et al.*, Mosaic nanoparticles elicit cross-reactive immune responses to
zoonotic coronaviruses in mice. *Science* **371**, 735-741 (2021).
74. W. C. Huang *et al.*, SARS-CoV-2 RBD neutralizing antibody induction is enhanced by
particulate vaccination. *Adv. Mater.* **32**, e2005637 (2020).
75. T. K. Tan *et al.*, A COVID-19 vaccine candidate using SpyCatcher multimerization of
the SARS-CoV-2 spike protein receptor-binding domain induces potent neutralising
antibody responses. *Nat. Commun.* **12**, 542 (2021).
76. D. Lainšček *et al.*, Immune response to vaccine candidates based on different types of
nanoscaffolded RBD domain of the SARS-CoV-2 spike protein. *bioRxiv*,
2020.2008.2028.244269 (2020).
77. X. Ma *et al.*, Nanoparticle vaccines based on the receptor binding domain (RBD) and
heptad repeat (HR) of SARS-CoV-2 elicit robust protective immune responses. *Immunity*
53, 1315-1330.e1319 (2020).
78. N. C. Dalvie *et al.*, Engineered SARS-CoV-2 receptor binding domain improves
immunogenicity in mice and elicits protective immunity in hamsters. *bioRxiv*,
2021.2003.2003.433558 (2021).
79. R. Rappuoli *et al.*, Vaccinology in the post–COVID-19 era. *Proc. Natl. Acad. Sci. U.S.A.*
118, e2020368118 (2021).
80. M. Jeyanathan *et al.*, Immunological considerations for COVID-19 vaccine strategies.
Nat. Rev. Immunol. **20**, 615-632 (2020).
81. L. Corey, J. R. Mascola, A. S. Fauci, F. S. Collins, A strategic approach to COVID-19
vaccine R&D. *Science* **368**, 948-950 (2020).
82. L. DeFrancesco, Whither COVID-19 vaccines? *Nat. Biotechnol.* **38**, 1132-1145 (2020).

83. P. S. Arunachalam *et al.*, Adjuvanting a subunit SARS-CoV-2 nanoparticle vaccine to
induce protective immunity in non-human primates. *bioRxiv*, 2021.2002.2010.430696
(2021).
84. P. J. Hotez, D. B. Corry, U. Strych, M. E. Bottazzi, COVID-19 vaccines: neutralizing
antibodies and the alum advantage. *Nat. Rev. Immunol.* **20**, 399-400 (2020).
85. S. P. Kasturi *et al.*, Programming the magnitude and persistence of antibody responses
with innate immunity. *Nature* **470**, 543-547 (2011).
86. S. G. Reed, M. T. Orr, C. B. Fox, Key roles of adjuvants in modern vaccines. *Nat. Med.*
19, 1597-1608 (2013).
87. T.-Y. Kuo *et al.*, Development of CpG-adjuvanted stable prefusion SARS-CoV-2 spike
antigen as a subunit vaccine against COVID-19. *Sci. Rep.* **10**, 20085 (2020).
88. J.-H. Tian *et al.*, SARS-CoV-2 spike glycoprotein vaccine candidate NVX-CoV2373
immunogenicity in baboons and protection in mice. *Nat. Commun.* **12**, 372 (2021).
89. C. Keech *et al.*, Phase 1–2 trial of a SARS-CoV-2 recombinant spike protein nanoparticle
vaccine. *N. Engl. J. Med.* **383**, 2320-2332 (2020).
90. I. Quast, D. Tarlinton, B cell memory: understanding COVID-19. *Immunity* **54**, 205-210
(2021).
91. D. J. Irvine, B. J. Read, Shaping humoral immunity to vaccines through antigen-
displaying nanoparticles. *Curr. Opin. Immunol.* **65**, 1-6 (2020).
92. M. D. Shin *et al.*, COVID-19 vaccine development and a potential nanomaterial path
forward. *Nat. Nanotechnol.* **15**, 646-655 (2020).
93. L. He *et al.*, Proof of concept for rational design of hepatitis C virus E2 core nanoparticle
vaccines. *Sci. Adv.* **6**, eaaz6225 (2020).
94. L. He *et al.*, Single-component multilayered self-assembling nanoparticles presenting
rationally designed glycoprotein trimers as Ebola virus vaccines. *bioRxiv*,
2020.2008.2022.262634 (2020).
95. L. He *et al.*, HIV-1 vaccine design through minimizing envelope metastability. *Sci. Adv.*
4, aau6769 (2018).
96. C. A. Schneider, W. S. Rasband, K. W. Eliceiri, NIH Image to ImageJ: 25 years of image
analysis. *Nat. Methods* **9**, 671-675 (2012).
97. T. J. Johnson, Glutaraldehyde fixation chemistry: oxygen-consuming reactions. *Eur. J.*
Cell Biol. **45**, 160-169 (1987).
661

Acknowledgements

Funding: This work was funded by National Institutes of Health grants AI137472, AI139092 (to

J.Z.), Ufovax/SFP-2018-0416, Ufovax/SFP-2018-1013, and Ufovax/SFP-2020-0111 (to J.Z.).

Y.-N.Z. thanks the Natural Sciences and Engineering Research Council of Canada (NSERC) for

a postdoctoral fellowship. We thank V. Bradaschia, Kyle Duffin, and M. Ganguly at the Centre

for Phenogenomics for their expertise in histology and immunostaining. We acknowledge the

expert assistance of S. Henderson, K. Vanderpool, and T. Fassel at the Core Microscopy Facility

at The Scripps Research Institute. We thank A. Saluk, B. Seegers, and B. Monteverde at the

Flow Cytometry Core Facility of The Scripps Research Institute for their expertise in flow

cytometry. The authors thank M. Arends for proofreading the manuscript. **Author**

contributions: Project design by Y.-N.Z., L.H., and J.Z. SARS-CoV-2 variant plasmid design

and processing by L.H. and C.S. Antigen production, purification, and basic characterization by

T.N., T.F., and L.H. Antibody and mouse plasma neutralization by J.P., T.F., and L.H. Mouse

immunization, plasma collection, lymph node isolation, immunohistology, TEM, and flow

cytometry by Y.-N.Z. Manuscript written by Y.-N.Z., L.H., and J.Z. All authors commented on

the manuscript. This is manuscript number 30082 from The Scripps Research Institute.

Competing interests: The authors declare no competing interests. **Data and material**

availability: All data are available in the main text or in the supplementary materials. Additional

data related to this paper may be requested from the corresponding author.

 Figure Legends

Fig. 1. SARS-CoV-2 SApNP vaccines induce broadly neutralizing antibody responses to

three variants of concern. (**A**) Molecular surface representations of vaccine constructs,

including two spikes (S2P$_{ECTO}$-5GS-1TD0 and S2GΔHR2-5GS-1TD0) and three spike-

presenting SApNPs (S2GΔHR2-5GS-ferritin (FR), S2GΔHR2-5GS-E2p-LD4-PADRE (E2p-

L4P), and S2GΔHR2-10GS-I3-01v9-LD7-PADRE (I3-01v9-L7P)). Representative negative-

stain EM (nsEM) image of S2GΔHR2-10GS- I3-01v9-L7P SApNPs is shown on the right. (**B**)

Neutralization of the original Wuhan-Hu-1 strain by mouse plasma induced by 5 different

vaccines at week 5 after two intraperitoneal injections. ID$_{50}$ titers derived from SARS-CoV-2-pp

neutralization assays are plotted, with average ID$_{50}$ values labeled on the plots. (**C**) Mouse

plasma neutralization against the original Wuhan-Hu-1 strain and three variants, B.1.1.7, B1.351,

and P.1, at week 5 after two intraperitoneal injections of the adjuvanted S2GΔHR2-10GS-I3-

01v9-L7P vaccine. Left panels 1-4: percent neutralization plots of individual mice against 4

SARS-CoV-2 strains; Right panel: ID$_{50}$ plot. In (B) and (C), the plasma samples were generated

in the previous study (*41*), where mice were immunized with 50 μg of adjuvanted vaccine

antigen. (**D**) Neutralization of the original Wuhan-Hu-1 strain by mouse plasma induced the

S2GΔHR2 spike and two large S2GΔHR2-presenting SApNPs. Vaccines were administered via

footpad injections (0.8 μg/injection, for a total of 3.3 μg/mouse). (**E**) Mouse plasma

neutralization against the original Wuhan-Hu-1 strain and three variants, B.1.1.7, B1.351, and

P.1, at week 5 after two footpad injections of the S2GΔHR2 spike and two large S2GΔHR2-

presenting SApNPs. In (B)-(E), the ID$_{50}$ values are plotted as mean ± SEM. The data were

analyzed using two-tailed unpaired Student's *t*-test for comparison between different vaccine

groups or two-tailed paired Student's *t*-test for comparison of ID$_{50}$ titers against SARS-Cov-2

variants using the same plasma samples from a mouse. $*p < 0.05$, $**p < 0.01$. (**F**) Neutralization

of four SARS-CoV-2 strains by human monoclonal antibodies including CR3022, B38, CB6,

S309, CC12.1, CC12.3, C105, and P2B-2F6. IC_{50} values are listed and color-coded (white: no

neutralization; green to red: low to high). The IC_{50} values were calculated with the

%neutralization range constrained within 0.0-100.0%.

Fig. 2. SARS-CoV-2 SApNP vaccines induce long-term lymph node follicle retention. (**A**, **B**)

S2GΔHR2-presenting I3-01v9 SApNP vaccine distribution in a lymph node 12 h after (**A**) a

single-dose or (**B**) prime-boost footpad injections (10 μg/injection, 40 μg/mouse). Schematic

illustration of SApNPs in lymph node follicles is shown. (**C**, **D**) Histological images of the

S2GΔHR2 spike and S2GΔHR2-presenting E2p and I3-01 SApNP vaccine trafficking and

retention in lymph node follicles 2 h to 8 weeks after (**C**) a single-dose or (**D**) prime-boost

injections. (**E**, **F**) Quantification of vaccine accumulation in lymph node follicles 48 h after (**E**) a

single-dose or (**F**) prime-boost injections. Data were collected from more than 10 lymph node

follicles ($n = 3\text{-}4$ mice/group). The data points are expressed as mean $\pm$ SD. The data were

analyzed using one-way ANOVA followed by Tukey's multiple comparison *post hoc* test. $**p <$

0.01, $***p < 0.001$, $****p < 0.0001$.

Fig. 3. SARS-CoV-2 SApNP vaccines interact with follicular dendritic cells (FDCs) and are

presented on FDC dendrites to B cells. (**A**, **B**) S2GΔHR2 spike and S2GΔHR2-presenting E2p

and I3-01 SApNP vaccine interaction with FDC networks in lymph node follicles 12 h after (**A**)

a single-dose or (**B**) prime-boost injections (10 μg/injection, 40 μg/mouse). Vaccine antigens (the

S2GΔHR2 spike and S2GΔHR2-presenting E2p and I3-01 SApNPs) colocalized with FDC

networks. Immunostaining is color-coded (Green: CD21; Red: CD169; White: anti-spike). (**C**)

Representative TEM images of an FDC surrounded by multiple B cells. S2GΔHR2-presenting

I3-01 SApNPs (yellow arrows) presented on FDC dendrites.

Fig. 4. SARS-CoV-2 SApNP vaccines induce robust long-lived germinal centers. (**A**) Top:

Representative immunohistological images of germinal centers at week 2 after a single-dose

injection of the S2GΔHR2-presenting I3-01 SApNP vaccine (10 μg/injection, 40 μg/mouse).

Bottom: germinal center B cells (GL7$^+$, red) adjacent to FDCs (CD21$^+$, green) in lymph node

follicles (left), and T_{fh} cells in the light zone (LZ) of germinal centers (right). (**B**, **C**)

Quantification of germinal center reactions using immunofluorescent images: GC/FDC ratio and

sizes of germinal centers 2, 5, and 8 weeks after (**B**) a single-dose or (**C**) prime-boost injections

(n = 4-7 mice/group). The GC/FDC ratio is defined as whether germinal center formation is

associated with an FDC network (%). Representative immunohistological images of germinal

centers in mice immunized using S2GΔHR2 spike or S2GΔHR2-presenting E2p and I3-01

SApNP vaccines at week 8 after (**D**) a single-dose or (**E**) prime-boost injections. (**F**, **G**)

Quantification of germinal center reactions using flow cytometry: percentage and number of

germinal center B cells and T follicular helper cells 2, 5, and 8 weeks after (**F**) a single-dose or

(**G**) prime-boost injections. The data points are shown as mean ± SD. The data were analyzed

using one-way ANOVA followed by Tukey's multiple comparison *post hoc* test for each

timepoint. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

Figure 1

A

S2P$_{ECTO}$-5GS-1TD0

S2GΔHR2-5GS-1TD0

S2GΔHR2-5GS-FR

S2GΔHR2-5GS-E2p-LD4-PADRE (L4P)

S2GΔHR2-10GS-I3-01v9-LD7-PADRE (L7P)

50 nm

B Mouse plasma neutralization against Wuhan-Hu-1

C Mouse plasma neutralization against SARS-CoV-2 variants

Wuhan-Hu-1

B.1.1.7

B.1.351

P.1

S2GΔHR2-10GS-I3-01v9-L7P

D Mouse plasma neutralization against Wuhan-Hu-1

E Mouse plasma neutralization against SARS-CoV-2 variants

S2GΔHR2-5GS-1TD0

S2GΔHR2-5GS-E2p-L4P

S2GΔHR2-10GS-I3-01v9-L7P

F Human monoclonal antibody neutralization against SARS-CoV-2 variants

mAb	Wuhan-Hu-1	B.1.1.7	B.1.351	P.1
CR3022	>10 µg/ml	>10 µg/ml	>10 µg/ml	>10 µg/ml
B38	0.700	>10 µg/ml	>10 µg/ml	>10 µg/ml
CB6	0.036	0.160	>10 µg/ml	>10 µg/ml
S309	8.222	3.217	0.148	0.107
CC12.1	0.049	0.205	>10 µg/ml	8.326
CC12.3	0.034	0.064	>10 µg/ml	>10 µg/ml
C105	0.412	0.452	>10 µg/ml	>10 µg/ml
P2B-2F6	0.837	0.352	>10 µg/ml	>10 µg/ml

Figure 2

A Single-dose - 12h

S2GΔHR2 spike | E2p SApNP | I3-01v9 SApNP | CD169/CD21 | Spike (anti-spike Abs)

B Prime-boost - 3w + 12h

S2GΔHR2 spike | E2p SApNP | I3-01v9 SApNP | CD169/CD21 | Spike (anti-spike Abs)

C I3-01v9 SApNPs (yellow arrow) are aligned on FDC dendrites

Figure 4