Mechanism of a COVID-19 nanoparticle vaccine candidate that elicits a broadly

neutralizing antibody response to SARS-CoV-2 variants

Yi-Nan Zhang[1], Jennifer Paynter[1], Cindy Sou[1], Tatiana Fourfouris[1], Ying Wang[3,4], Ciril

Abraham[3], Timothy Ngo[1], Yi Zhang[3,4], Linling He[1], and Jiang Zhu[1, 2, *]

[1]Department of Integrative Structural and Computational Biology, [2]Department of Immunology

and Microbiology, The Scripps Research Institute, La Jolla, California 92037, USA

[3]Fels Institute for Cancer Research and Molecular Biology, and [4]Department of Microbiology

and Immunology, Temple University, Philadelphia, Pennsylvania 19140, USA.

*Corresponding author

JZ: Phone +1 (858) 784-8157; Email: jiang@scripps.edu

KEYWORDS

ABSTRACT (150 words)

Vaccines that induce potent neutralizing antibody (NAb) responses against emerging variants of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) are essential for combating the coronavirus disease 2019 (COVID-19) pandemic. We demonstrated that mouse plasma induced by self-assembling protein nanoparticles (SApNPs) that present 20 rationally designed S2GΔHR2 spikes of the ancestral Wuhan-Hu-1 strain can neutralize the B.1.1.7, B.1.351, P.1, and B.1.617 variants with the same potency. The adjuvant effect on vaccine-induced immunity was investigated by testing 16 formulations for the multilayered I3-01v9 SApNP. Using single-cell sorting, monoclonal antibodies (mAbs) with diverse neutralization breadth and potency were isolated from mice immunized with the receptor binding domain (RBD), S2GΔHR2 spike, and SApNP vaccines. The mechanism of vaccine-induced immunity was examined in mice. Compared with the soluble spike, the I3-01v9 SApNP showed 6-fold longer retention, 4-fold greater presentation on follicular dendritic cell dendrites, and 5-fold stronger germinal center reactions in lymph node follicles.

ONE-SENTENCE SUMMARY (125 characters)

With a well-defined mechanism, spike nanoparticle vaccines can effectively counter SARS-CoV-2 variants.

INTRODUCTION

The COVID-19 pandemic has led to more than 188 million infection cases and 4 million deaths globally. Antibody responses to SARS-CoV-2 spike antigens can be sustained for several months in most COVID-19 patients after infection (*1-4*). However, recently identified variants of concern (VOCs) exhibit higher transmissibility and resistance to prior immunity as SARS-CoV-2 continues to adapt to the human host (*5, 6*). One such variant, B.1.1.7 (WHO classification: Alpha), emerged from southeast England in October 2020 and accounted for two-thirds of new infections in London in December 2020, with a higher transmission rate (43-90%) and risk of mortality (32-104%) than previously circulating strains (*7, 8*). Other variants, such as B.1.351 (Beta) and P.1 (Gamma), also became prevalent in three provinces in South Africa and Manaus, Brazil, respectively (*6, 9, 10*). The B.1.617.2 (Delta) variant, which was initially identified in India, is becoming a dominant strain in many countries (*11, 12*) and responsible for the majority of new COVID-19 cases. This variant was found to be ~60% more transmissible than the highly infectious B.1.1.7 variant (*12*). The rise of SARS-CoV-2 VOCs and their rapid spread worldwide result in more infection cases, hospitalizations, and potentially more deaths, further straining healthcare resources (*10*).

To date, eight COVID-19 vaccines have been approved for emergency use in humans, with more than 90 candidates assessed in various phases of clinical trials (*13*). With the exception of inactivated whole-virion vaccines, diverse platforms have been used to deliver the recombinant SARS-CoV-2 spike, such as mRNA-encapsulating liposomes (e.g., BNT162b2 and mRNA-1273), adenovirus vectors (e.g., ChAdOx1 nCoV-19 [AZD1222], CTII-nCoV, Sputnik V, and Ad26.COV2.S), and micelle-attached spikes (e.g., NVX-CoV2373). These vaccines demonstrated 65-96% efficacy in Phase 3 trials, with lower morbidity and mortality associated

with COVID-19 disease (*14-19*). However, a notable loss of vaccine efficacy against new SARS-

CoV-2 variants was reported, likely caused by spike mutations in the receptor-binding domain

(RBD; e.g., K417N, E484K, and N501Y), N-terminal domain (NTD; e.g., L18F, D80A, D215G,

and Δ242-244), and other regions that are critical to spike stability and function (e.g., D614G and

P681R) (*6, 11, 20-25*). Among circulating VOCs, the B.1.351 lineage appeared to be most

resistant to neutralization by convalescent plasma (9.4-fold) and vaccine sera (10.3- to 12.4-fold)

(*26*), whereas a lesser degree of reduction was observed for an early variant, B.1.1.7 (*27-29*).

Based on these findings, it was suggested that vaccines would need to be updated periodically to

maintain protection against rapidly evolving SARS-CoV-2 (*30-32*). However, in a recent study,

convalescent sera from B.1.351 or P.1-infected individuals showed a more visible reduction of

B.1.617.2 neutralization than convalescent sera from individuals infected with early pandemic

strains (*33*). Together, these issues raise the concern that herd immunity may be difficult to

achieve, highlighting the necessity of developing vaccines that can elicit a broadly neutralizing

antibody (bNAb) response to current and emerging variants (*25, 31*). As previously reported (*34-

38*), the production of a bNAb response relies on long-lived germinal center (GC) reactions to

activate precursor B cells, stimulate affinity maturation, and form long-term immune memory. In

particular, antigen retention and presentation within lymph node follicles are key to the induction

of long-lived GC reactions (*34, 36, 39*) and should be considered in the development of bNAb-

producing vaccines (*40*).

We previously investigated the cause of SARS-CoV-2 spike metastability and rationally

designed the S2GΔHR2 spike, which was displayed on three self-assembling protein

nanoparticle (SApNP) platforms, including ferritin (FR) 24-mer and multilayered E2p and I3-

01v9 60-mers, as COVID-19 vaccine candidates (*41*). In the present study, we investigated the

vaccine-induced NAb response to SARS-CoV-2 VOCs and mechanism by which SApNP

vaccines (e.g., I3-01v9) generate such a response. We first examined the neutralizing activity of

mouse plasma from our previous study (*41*) against four representative SARS-CoV-2 variants,

B.1.1.7, B.1.351, P.1, and B.1.617$_{Rec}$, which was derived from an early analysis of the B.1.617

lineage (*11*) and shares key spike mutations with VOC B.1.617.2. Mouse plasma induced by the

S2GΔHR2 spike-presenting I3-01v9 SApNP potently neutralized all four variants with

comparable titers to the wildtype strain, Wuhan-Hu-1. When a different injection route was

tested in mouse immunization, E2p and I3-01v9 SApNPs sustained neutralizing titers against the

four variants, even at a low dosage of 3.3 μg, whereas a significant reduction of plasma

neutralization was observed for the soluble spike. Next, we examined the adjuvant effect on

vaccine-induced humoral and T-cell responses for the I3-01v9 SApNP. While detectable plasma

neutralization was observed for the non-adjuvanted I3-01v9 group, conventional adjuvants, such

as aluminum hydroxide (AH) and phosphate (AP), boosted the titers by 8.6- to 11.3-fold (or 9.6

to 12.3 times). Adjuvants that target the stimulator of interferon genes (STING) and Toll-like

receptor 9 (TLR9) pathways enhanced neutralization by 21- to 35-fold, alone or combined with

AP, in addition to a Th1-biased cellular response. We then performed antigen-specific single-cell

sorting and isolated 20 monoclonal antibodies (mAbs) from RBD, spike, and I3-01v9 SApNP-

immunized mice. These mAbs were derived from diverse B cell lineages, of which some

neutralized the wildtype Wuhan-Hu-1 strain and four variants with equivalent potency. Lastly,

we investigated how SApNPs behave in lymph nodes and induce GCs by characterizing vaccine

delivery and immunological responses at the intraorgan, intracellular, and intercellular levels in

mice. The I3-01v9 SApNP showed 6-fold longer retention, 4-fold greater presentation on

follicular dendritic cell (DC) dendrites, and 5-fold higher GC reactions than the soluble spike.

Intact SApNPs in lymph node tissues could be visualized by transmission electron microscopy

(TEM). Our study thus demonstrates that a spike-presenting SApNP vaccine derived from the

"ancestral" SARS-CoV-2 strain may confer broad protection against emerging variants.

RESULTS

Spike and SApNP vaccine-induced neutralizing responses to SARS-CoV-2 variants.

We previously demonstrated that the rationally designed S2GΔHR2 spike was more

immunogenic than the S2P spike (*42*), and SApNPs displaying 8-20 spikes outperformed soluble

spikes in NAb elicitation (*41*) (**Fig. 1A**). Notably, the I3-01v9 SApNP that presents 20

S2GΔHR2 spikes induced a potent NAb response to both SARS-CoV-1 and SARS-CoV-2, as

well as critically needed T-cell responses (*41*). Because SARS-CoV-1 shares only modest

sequence similarity (~73% in the RBD) with SARS-CoV-2, we hypothesized that our vaccines

would protect against emerging variants that are much more closely related to the ancestral

SARS-CoV-2 strain, Wuhan-Hu-1.

We first assessed the neutralizing activity of polyclonal plasma induced by various spike

and SApNP vaccine formulations from our previous study (*41*) against the wildtype SARS-CoV-

2 strain, Wuhan-Hu-1, as a baseline for comparison (**Fig. 1B**). Mouse plasma collected at week 5

after two intraperitoneal (i.p.) injections of adjuvanted vaccine antigens (50 μg) was analyzed in

pseudoparticle (pp) neutralization assays (*43*). The soluble S2P$_{ECTO}$ spike elicited the lowest 50%

inhibitory dilution (ID$_{50}$) titers, whereas the soluble S2GΔHR2 spike increased neutralization

with a 7.1-fold higher average ID$_{50}$ titer, which did not reach statistical significance because of

within-group variation. All three spike-presenting SApNPs elicited superior neutralizing

responses than the soluble S2P$_{ECTO}$ spike (*41*). Notably, the I3-01v9 SApNP achieved the highest

potency, with an average ID$_{50}$ titer of 2090, which was 8.1-fold higher than the soluble S2P$_{ECTO}$

spike. Despite differences in ID_{50} titers, the overall pattern remained the same as reported in our

previous study (*41*). The differences might be attributable to the inherent variation of

pseudovirus assays (*43, 44*). We then assessed plasma neutralization against four major SARS-

CoV-2 variants (**Fig. 1C**, **fig. S1A, B**). The I3-01v9 SApNP induced a stronger neutralizing

response against variants, with 0.5-fold (B.1.1.7), 0.8-fold (B.1.351), 1.8-fold (P.1), and 1.0-fold

(B.1.617$_{Rec}$) higher (or 1.5-2.8 times) ID_{50} titers compared with the wildtype strain (**Fig. 1C**).

Altogether, these results confirmed our hypothesis and highlighted the advantages of spike-

presenting SApNPs.

Next, we examined the influence of injection dosage and route on the plasma neutralizing

response to various SARS-CoV-2 strains. To this end, we performed a mouse study, in which

three groups of mice were immunized with 5, 15, and 45 μg of the I3-01v9 SApNP three times

via i.p. injection. Remarkably, all four variants were neutralized by mouse plasma with

comparable ID_{50} titers observed across dose groups (**fig. S1C, D**). To examine whether routes of

injection affect the plasma neutralizing response against variants, we performed another mouse

study, in which a low dose (3.3 μg) of adjuvanted antigen was intradermally administered into

four footpads (i.e., 0.8 μg/footpad). At week 5, the large (~55-60 nm) E2p and I3-01v9 SApNPs

that present 20 S2GΔHR2 spikes yielded higher ID_{50} titers against the wildtype strain than the

soluble S2GΔHR2 spike (**Fig. 1D**, **fig. S1E, F**), whereas a notable reduction of ID_{50} titers against

the variants was noted for mouse plasma from the S2GΔHR2 group (**Fig. 1E**, **fig. S1E, F**),

suggesting that multivalent display is critical for eliciting a broad neutralizing response. Overall,

the E2p and I3-01v9 SApNP groups exhibited similar or slightly stronger plasma neutralization

against the four variants relative to the wildtype strain, Wuhan-Hu-1 (**Fig. 1E**). Lastly, we

assessed longevity of the low-dose vaccination-induced neutralizing response by testing week-26

plasma against Wuhan-Hu-1 (**Fig. 1F**, **fig. S1G, H**). It is noteworthy that ID_{50} titers at week 26

were at the same level as week 5, suggesting a long-lasting protective humoral immunity. In our

previous study, a panel of human NAbs was used to evaluate antigenicity of the stabilized

S2GΔHR2 spike and SApNPs and validate the SARS-CoV-2-pp neutralization assays (*41*). Here,

this antibody panel was tested against SARS-CoV-2-pps that carry spikes of the wildtype strain

and the four variants (**Fig. 1G**, **fig. S1I**). Lower potency against the B.1.351 and P.1 variants,

measured by the 50% inhibitory concentration (IC_{50}), was observed for all human NAbs, with the

exception of NAb S309, which was identified from a SARS-CoV-1 patient (*45*). This finding is

consistent with recent reports on convalescent patient plasma (*26-28*). Interestingly, most human

NAbs remained effective against $B.1.617_{Rec}$ showing a similar pattern to the wildtype Wuhan-

Hu-1 strain and B.1.1.7 variant, consistent with the results of a recent cohort analysis of

convalescent sera from individuals infected with early VOCs against a rising B.1.617 (*33*). As a

negative control, mouse plasma induced by the S2GΔHR2-presenting I3-01v9 SApNP was tested

against pseudoviruses carrying the murine leukemia virus (MLV) envelope glycoprotein (Env),

or MLV-pps. Nonspecific MLV-pp neutralization was not detected for plasma samples produced

in two independent immunization experiments (**fig. S1J, K**).

Altogether, our results demonstrate that spike-presenting SApNPs are more advantageous

than soluble spikes in eliciting a strong neutralizing response to diverse SARS-CoV-2 variants.

In our previous study, soluble SARS-CoV-2 spikes induced a more effective neutralizing

response to SARS-CoV-1 than a scaffolded SARS-CoV-2 RBD trimer (*41*). Recently, a two-

component RBD-NP vaccine showed reduced serum neutralization of variants bearing the

E484K mutation (*46*). It is plausible that both the nanoparticle (NP) platform (one-component

SApNP *vs.* two-component NP) and the antigen type (spike *vs.* RBD) contribute to vaccine

breadth.

Adjuvant effect on vaccine-induced neutralizing antibody and T-cell responses.

Innate immunity plays an important role in regulating adaptive immunity, including humoral and

cellular immune responses (*47-49*). Adjuvant-formulated vaccines have been shown to recruit

and activate innate immune cells more effectively at injection sites and local lymph nodes (*50-*

52). Among commonly used adjuvants, AH and AP create depots for the recruitment and

activation of antigen-presenting cells (APCs) at injection sites and sentinel lymph nodes (*53, 54*),

whereas oil-in-water emulsions such as MF59 promote antigen retention and APC stimulation in

lymph nodes (*55*). Pattern recognition receptor (PRR) agonists (e.g., STING, TLR3, TLR4,

TLR7/8, and TLR9 agonists) stimulate APCs at injection sites and nearby lymph nodes (*47, 52,*

56-59). Macrophage inhibitors (e.g., clodronate liposomes, termed CL) directly stimulate B cells

or inhibit antigen sequestration by subcapsular sinus macrophages, thus resulting in more

effective GC simulation in lymph nodes (*60*). Adjuvant combinations may generate a synergistic

immune response by simultaneously activating multiple pathways (*52, 57*).

To examine the effect of innate signaling pathways on SApNP-induced immune

responses, we tested 16 adjuvant formulations in a systematic study (**Fig. 2A**), in which mice

were immunized with the adjuvanted I3-01v9 SApNP (20 μg) via intradermal injections in four

footpads (i.e., 5 μg/footpad). We first tested mouse plasma neutralization against the wildtype

Wuhan-Hu-1 stain. Mouse plasma at week 2 after a single dose was analyzed in SARS-CoV-2-

pp assays, with most groups showing negligible or borderline ID_{50} titers (**fig. S2A, B**). We then

analyzed mouse plasma at week 5 after two injections (**Fig. 2B, fig. S2C, D**). The non-

adjuvanted group showed detectable neutralization after two doses, with an average ID_{50} titer of

160, which was used as a baseline in this analysis. By comparison, conventional adjuvants, such

as AH, AP, and AddaVax, increased ID_{50} titers by 10.3-, 7.6-, and 12.5-fold, respectively. The

macrophage inhibitor CL boosted plasma neutralization by merely 1.6-fold relative to the non-

adjuvanted group. Adjuvants that target various PRRs exhibited differential effects on plasma

neutralization, increasing ID_{50} titers by 1.1- to 34.2-fold. Notably, STING and CpG (TLR9)

substantially enhanced neutralizing titers, whereas TLR3, TLR4, and TLR7/8 agonists only

exerted a modest effect. In most cases, adjuvants combined with AP further boosted plasma

neutralizing activity. For example, when TLR4 and TLR7/8 agonists were mixed with AP, a 3.1-

fold increase in ID_{50} titers was observed, suggesting a synergistic effect of stimulating multiple

immune pathways. Overall, STING and CpG, either alone or combined with AP, showed plasma

neutralization superior to that of any other adjuvant or adjuvant mix, increasing ID_{50} titers by 21

to 34-fold compared with the non-adjuvanted group. This is consistent with the results of the S-

Trimer (SCB-2019), which, when formulated with CpG 1018 (TLR9 agonist) and alum

adjuvants, induced potent NAb responses in nonhuman primates and human trials (*61, 62*).

Mouse plasma at week 8 showed further increases in ID_{50} titers (1 to 3-fold) for most adjuvant

groups (**Fig. 2C, fig. S2E, F**). Lastly, we examined mouse plasma at week 5 from the STING

and CpG groups against the B.1.1.7, B.1.351, P.1, and B.1.617$_{rec}$ variants (**Fig. 2D, fig. S2G, H**).

Both adjuvant groups exhibited potent neutralizing responses to the four variants, with ID_{50} titers

comparable to the wildtype strain.

We previously demonstrated that the AP-formulated I3-01v9 SApNP induces interferon-γ

(IFN-γ)-producing CD4$^+$ T helper 1 (Th1) cells and IFN-γ/interleukin-4 (IL-4) double-positive

memory CD4$^+$ T cells (*41*). Given the superior plasma neutralizing response observed for STING

and CpG, we examined the impact of these two adjuvants on vaccine-induced T-cell responses.

IFN-γ-producing $CD4^+$ Th cells are important for optimal antibody responses and the induction

of cellular immunity to clear viruses (*63-65*). To assess the effect of STING and CpG on

vaccine-induced Th cells, we isolated splenocytes from mice 8 weeks after vaccination and

cultured them in the presence of BALB/c mouse DCs pulsed with the spike-presenting I3-01v9

SApNP. Compared with the non-adjuvanted control, STING and CpG (TLR9) induced 3.7 and

5.5-fold more IFN-γ-producing $CD4^+$ Th1 cells, and 5.5 and 16-fold more IL-4-producing $CD4^+$

Th2 cells, respectively (**Fig. 2E, fig. S2I**). A visible but nonsignificant trend toward a higher

frequency of both Th1 and Th2 cells was noted in mice immunized with the CpG-formulated I3-

01v9 SApNP than other formulations. Nonetheless, both adjuvants induced more IFN-γ-

producing $CD4^+$ Th1 cells than IL-4-producing $CD4^+$ Th2 cells, suggesting a dominant Th1

response in these mice. This is consistent with the results for the S-Trimer (SCB-2019), which,

when formulated with the AS03 adjuvant or mixed CpG 1018/alum adjuvants, induced Th1-

biased cellular responses in mice (*61*). STING and CpG also enhanced $CD8^+$ T-cell responses by

6- and 10-fold, respectively, compared with the PBS control. Notably, this effect was more

visible for CpG in terms of both the frequency and number of IFN-γ-producing $CD8^+$ effector T

cells (**Fig. 2F, fig. S2J**).

Our results demonstrate that the I3-01v9 SApNP itself is immunogenic, and adjuvants

can further enhance vaccine-induced NAb responses in plasma by up to 35-fold. The I3-01v9

SApNP, when formulated with the STING or TLR9 agonist, yielded the highest ID_{50} titers with

robust $CD4^+$ and $CD8^+$ T-cell responses, highlighting their potential as adjuvants in the

development of more effective SARS-CoV-2 vaccines.

Diverse variant-neutralizing mouse antibody lineages identified by single-cell analysis.

Although plasma neutralization confirmed the effectiveness of our newly designed SARS-CoV-2

vaccines (*41*) against variants, the nature of this response was unclear. It might result from

multiple NAb lineages that each target a specific strain (non-overlapping), a few bNAb lineages

that are each able to block multiple strains (overlapping), or a combination of both. Previously,

we used antigen-specific single-cell sorting to identify potent mouse NAbs elicited by an I3-01

SApNP that presents 20 stabilized HIV-1 Env trimers (*66*). Here, we applied a similar strategy to

decipher NAb responses induced by SARS-CoV-2 vaccines using mouse samples from our

previous study (*41*), for which potent plasma neutralization against four variants has been

verified (**Fig. 1C**).

Spleen samples from M4 in the spike group (S2GΔHR2-5GS-1TD0) and M2 in the spike-

SApNP group (S2GΔHR2-10GS-I3-01v9-L7P), along with a control sample from M2 in the

RBD (RBD-5GS-1TD0) group were analyzed. Two probes, RBD-5GS-foldon-Avi and

S2GΔHR2-5GS-foldon-Avi, were produced, biotinylated, and purified to facilitate antigen-

specific B-cell sorting (**fig. S3A**, **B**)**.** Following antibody cloning, reconstituted mouse mAbs

were tested for neutralizing activity against the wildtype strain, Wuhan-Hu-1, in SARS-CoV-2-

pp assays. A total of 20 mAbs, four from the RBD group (**fig. S3C**), six from the spike group

(**fig. S3D**), and 10 from the I3-01v9 SApNP group (**fig. S3E**), were found to be NAbs. The

genetic analysis of mAb sequences revealed some salient features of the vaccine-induced NAb

response in mice (**Fig. 3A**). Overall, these mAbs evolved from diverse germline origins. The

RBD-elicited mAbs appeared to use distinct germline variable (V) genes for both heavy chain

(HC) and κ-light chain (KC), or V_H and V_K, respectively, whereas the spike and I3-01v9

SApNP-elicited mAbs shared some common V_H genes, such as IGHV14-1/3 and IGHV1S81.

This result was not unexpected because the RBD vaccine presents a structurally distinct antigen

to the immune system compared with the spike and I3-01v9 vaccines, which both present the

S2GΔHR2 spike. These mAbs showed low levels of somatic hypermutation (SHM) with respect

to their germline genes. Heavy chain complementarity-determining region (HCDR3) loops

ranged from 4 to 12 aa in length, whereas most KCs contained 9 aa KCDR3 loops. Collectively,

diverse germline genes and HCDR3 loops, accompanied by low degrees of SHM, suggest that

many antibody lineages must have been generated upon vaccination, and some could achieve

neutralizing activity without an extensive maturation process.

We then examined the biological function of these mouse mAbs. Neutralizing activity

was assessed in SARS-CoV-2-pp assays against the wildtype strain and four variants (**Fig. 3B**,

fig. S3F). Overall, diverse yet consistent patterns were observed for the three sets of mAbs. Both

the RBD vaccine (an RBD scaffold) and the two spike vaccines, albeit in different forms,

appeared to elicit potent NAbs against the wildtype strain. MAbs TRBD-R-4G5, S2GD-S-2C10,

and I3V9-R-1G9 showed similar IC_{50} values (0.02-0.03 μg/ml) against Wuhan-Hu-1, on par with

the human NAbs CB6 (*67*) and CC12.1/3 (*68*) (**Fig. 1F**). All three vaccines elicited bNAb

responses, despite variation in potency for different mAbs against different strains. Notably,

I3V9-R-1G9, which was isolated from an I3-01v9 SApNP-immunized mouse, demonstrated high

potency across all four variants (IC_{50}: 0.01-0.02 μg/ml). This bNAb provided evidence that

individual bNAb lineages may critically contribute to the plasma neutralization of diverse

variants (**Fig. 1C**). All three vaccines generated NAbs that preferentially neutralize specific

SARS-CoV-2 strains. For example, TRBD-R-4B6 was more effective against the wildtype strain

and an early VOC, B.1.1.7, whereas S2GD-R-2E4 neutralized B.1.351, P.1, and $B.1.617_{Rec}$ with

greater potency. Notably, more than 60% (13) of the mAbs exhibited different patterns in the

neutralization of $B.1.617_{Rec}$ *vs.* VOCs B.1.351 and P.1, as indicated by the fold change in IC_{50},

suggesting that B.1.617 may represent a distinct SARS-CoV-2 lineage. Although RBD-isolated

NAbs likely neutralized SARS-CoV-2 by blocking its receptor binding, those spike-isolated

NAbs could target the RBD, N-terminal domain (NTD), or epitopes in the S2 subunit. Thus, we

tested these mAbs in an enzyme-linked immunosorbent assay (ELISA) against the RBD

monomer and S2GΔHR2-5GS-1TD0 spike, both based on the wildtype Wuhan-Hu-1 backbone

(**Fig. 3C, fig. S3G, H**). Overall, all of the NAbs bound the RBD and spike with a half-maximal

concentration (EC_{50}) of 0.034 µg/ml or lower, except for I3V9-R-2F2 (1.973 µg/ml for the RBD

and 2.051 µg/ml for the spike). Most (15) NAbs showed greater binding affinity (or lower EC_{50}

values) for the spike, suggesting that the two arms of the immunoglobulin (Ig) can each interact

with one RBD of the spike, resulting in an avidity effect. Notably, diverse binding patterns were

observed for I3-01v9 SApNP-elicited NAbs. Although I3V9-S-1C9 and I3V9-S-1F5 bound to

the spike more favorably than the RBD, as indicated by a 4.7 to 4.9-fold reduction of their EC_{50}

values, three NAbs from this group (I3V9-R-1G3, I3V9-R-1G9, and I3V9-R-2F10) preferred the

RBD monomer over the spike. This preference might be explained by steric hindrance when

these NAbs approach the RBDs on a trimeric spike at specific angles.

Lastly, we characterized these mouse NAbs in antigen-specific B-cell repertoires by next-

generation sequencing (NGS), as previously demonstrated for NAbs isolated from HIV-1

SApNP-immunized mice and rabbits (*66*). Using the same RBD and spike probes (**fig. S3A**),

~1500 splenic B cells were bulk-sorted from each of the three mice that were analyzed by single-

cell sorting for mAb isolation (**fig. S4A**). Unbiased mouse antibody HC and KC libraries were

constructed and sequenced on an Ion S5 platform, which yielded up to 4 million raw reads (**fig.**

S4B). The antibody NGS data were then processed using a mouse antibodyomics pipeline (*69*) to

remove low-quality reads, resulting in 0.11-0.41 full-length HCs and KCs (**fig. S4B**).

Quantitative profiles of critical antibody properties, such as germline gene usage, the degree of

SHM, and CDR3 loop length, were determined for the RBD and spike-specific B-cell

populations (**fig. S4C**). All 20 single-cell-sorted mouse NAbs could well fall in the range of

these repertoire profiles, but some V_H/V_K genes that accounted for large portions of antigen-

specific B cells, such as IGHV9 and IGHV5, were not used by any NAbs, suggesting that they

might give rise to non-neutralizing binding antibodies. Two-dimensional (2D)

divergence/identity plots were generated to visualize these NAbs in the context of NGS-derived

B-cell repertoires (**Fig. 3D**, **fig. S4D-F**). Somatic variants were identified for each NAb by

searching for sequences of the same V_H/V_K gene with a CDR3 identity cutoff of 90% (or 85%

for evolutionarily more remote variants). For the most potent NAb, TRBD-R-4G5, from an

RBD-immunized mouse (M2), 34 HC variants were identified that overlapped with an "island"

of high sequence similarity to TRBD-R-4G5 on the plot, whereas more KC variants (1183) were

found, likely due to the lack of diversity in the KCDR3 region. A similar pattern was observed

for the potent bNAb, I3V9-R-1G9, from an SApNP-immunized mouse (M2). By comparison,

fewer putative somatic variants were identified for other NAbs in the antigen-specific B-cell

repertoires regardless of the sorting probe used (**fig. S4D-F**), suggesting that these NAbs either

were from less prevalent lineages or were generated in response to a previous injection (each

mouse received four doses) (*41*). Similar observations were reported for the vaccination of

nonhuman primates and humans in longitudinal repertoire analyses of single-cell-sorted NAbs

(*70, 71*).

Single-cell isolation identified a panel of mouse mAbs with different neutralization

breadth and potency against the wildtype SARS-CoV-2 strain and four major variants. The

ELISA analysis suggested that the I3-01v9 SApNP can elicit NAbs with more diverse angles of

approach to their epitopes than the RBD and soluble spike vaccine. Structural analysis by

crystallography and EM will provide a more detailed understanding of epitope recognition by

these mouse mAbs.

Distribution and trafficking of I3-01v9 SApNP in mouse lymph node.

After validating these vaccines against variants at both the plasma and mAb levels, we studied *in*

vivo behaviors of the S2GΔHR2 spike and two large 60-meric SApNPs to understand why

SApNPs outperform soluble spikes in bNAb elicitation. In principle, these SApNPs need to be

transported to lymph nodes, retained, and presented to various immune cell populations to induce

robust innate and adaptive immune responses. Here, we first examined the transport and

distribution of I3-01v9 SApNPs in mouse lymph nodes via footpad injections (10 μg/footpad).

The mice were sacrificed 12 h after single-dose (**Fig. 4A**) and prime-boost (**Fig. 4B**) regimens.

The axillary, brachial, and popliteal sentinel lymph nodes were isolated for histological analysis.

The lymph node tissues were stained with the human anti-spike antibody P2B-2F6 (*72*) to

characterize SARS-CoV-2 spikes presented on the I3-01v9 SApNPs. Consistent with our

previous study (*73*), SApNPs accumulated in lymph node follicles, regardless of the number of

doses. SApNPs were sequestrated in the center of lymph node follicles after a single dose (**Fig.**

4A, images on the left, schematics on the right) but were located along the outer layer of

expanded lymph node follicles after the second injection due to preexisting humoral immunity

(i.e., GC reactions) that was induced by the first dose (**Fig. 4B**, images on the left, schematics on

the right). Overall, the majority of SApNPs accumulated in lymph node follicles, but their

distribution differed slightly, depending on the doses.

In this context, we examined patterns of trafficking and lymph node follicle retention for

soluble S2GΔHR2 spike *vs*. the S2GΔHR2-presenting E2p and I3-01v9 SApNPs. To facilitate

this analysis, the mice were sacrificed 2 h to 8 weeks after a single dose (**Fig. 4C**) and 2 h to 5

weeks after the boost (**Fig. 4D**). The injection dose was normalized to the total amount of protein

(10 µg) per injection into each footpad (40 µg/mouse). As shown in **Fig. 4C**, the S2GΔHR2

spikes that trafficked into lymph node follicles at 2 h cleared within 48 h. In contrast, the two

large SApNPs accumulated in the subcapsular sinus at 2 h and then trafficked into follicles 12 h

after the single-dose injection. Remarkably, I3-01v9 SApNPs remained detectable in lymph node

follicles after 2 weeks, suggesting 6-fold longer retention than the S2GΔHR2 spike (**Fig. 4C**).

The results for these protein nanoparticles are thus consistent with the pattern of size dependency

that was observed for ovalbumin-conjugated gold nanoparticles in our previous study (*73*), in

which small (5-15 nm) nanoparticles cleared shortly after the injection, whereas large (50-100

nm) nanoparticles were retained in lymph node follicles for weeks. Similar patterns of antigen

retention were observed after the second injection, although the boost appeared to exert a more

positive effect on the soluble spike, which could be detected in lymph node follicles at 48 h (**Fig.**

4D). Nonetheless, prolonged retention was observed for both E2p and I3-01v9 SApNPs 2 weeks

after the boost injection. Overall, the multivalent display of S2GΔHR2 spikes on the I3-01v9

SApNP resulted in 325- and 4-fold greater accumulation in lymph node follicles compared with

the soluble spike 48 h after the single-dose **(Fig. 4E**) and prime-boost (**Fig. 4F**) injections,

respectively. These findings reveal the advantage of a leading vaccine candidate identified in our

previous study, S2GΔHR2-10GS-I3-01v9-L7P (*41*), in terms of spike antigen retention in lymph

node follicles.

Retention and presentation of I3-01v9 SApNP on follicular dendritic cell dendrites.

Antigen retention and presentation in lymph node follicles are prerequisites to the stimulation of

robust B cell responses and GC reactions (*34, 36*). Resident cells spatially rearrange antigens and

present them to B cells. Follicular dendritic cells (FDCs) are resident stromal cells in follicles

and retain soluble antigens, immune complexes, virus-like particles (VLPs), viruses, and bacteria

(*73-76*). FDCs are also key to GC initiation, maintenance, and B-cell affinity maturation (*37, 77,*

78). Here, we hypothesized that FDCs comprise the major cell population in lymph node follicles

that retain SARS-CoV-2 spikes and spike-presenting SApNPs. To test this hypothesis, we

administered vaccines via footpad injections and collected mouse lymph nodes at the peak of

accumulation (12 h) after single-dose (**Fig. 5A**) and prime-boost (**Fig. 5B**) injections. Lymph

node tissue samples were stained with the anti-spike antibody P2B-2F6 (*72*) for the S2GΔHR2

spike, as well as anti-CD21 and anti-CD169 antibodies for FDCs and subcapsular sinus

macrophages, respectively. The spike and SApNP (E2p or I3-01v9) signals colocalized with

FDC (CD21$^+$) networks in lymph node follicles (**Fig. 5A, B**). This result confirmed the critical

role of FDC networks in mediating vaccine retention in lymph node follicles.

The induction of potent bNAb responses by spike-presenting SApNPs in mice suggests

the effective activation of naïve B cells and subsequent recalls by crosslinking B cell receptors

(*76, 79, 80*). We visualized the interface between FDC networks and B cells to better understand

how FDC networks present SApNPs to engage B cells. Briefly, fresh lymph nodes were isolated

and directly immersed in fixative. The processed tissue samples were sectioned and stained on

copper grids for TEM analysis. We first determined whether SApNPs, such as the S2GΔHR2-

presenting I3-01v9 SApNP, remain intact *in vivo* (**fig. S5**). Mouse lymph nodes were isolated 2 h

after the injection of a high dose (50 μg) of the non-adjuvanted I3-01v9 SApNP. The TEM

images revealed that round-shape granules corresponding to intact SApNP aligned on the

macrophage surface or inside endolysosomes of the macrophage in a lymph node (**fig. S5**). We

next studied the relative location between FDCs and I3-01v9 SApNPs and how FDCs present

SApNPs to B cells. Mouse lymph nodes were collected 2, 12, and 48 h after a single-dose (50

µg) and 12 h after the boost of the I3-01v9 SApNP vaccine. The FDCs exhibited the

characteristic morphology of long dendrites that surrounded and interacted with B cells in lymph

node follicles (**Fig. 5C**, **fig. S6**). Few I3-01v9 SApNPs were observed on FDC dendrites at 2 h

(**fig. S6D**), whereas notably more nanoparticles migrated to and aligned on FDC dendrites at 12

and 48 h (**Fig. 5C**, **fig. S6A-C**, yellow arrows). The TEM images indicated that FDCs can

present many SApNPs to neighboring B cells in this "hugging mode", in which their long

dendrites brace B cells to maximize interactions between multivalently displayed spikes and B

cell receptors. These results demonstrated the intrinsic nature of FDCs as a reservoir for the

sequestration, retention, and presentation of virus-like particles, or SApNPs with similar

molecular traits, to initiate GC reactions.

Robust germinal center reactions induced by spike-presenting SApNPs.

Long-lived GC reactions induce immune stimulation for B-cell selection and affinity maturation,

as well as production of immune memory and bNAb responses (*34, 35, 40*). Here, we

investigated whether the prolonged retention of S2GΔHR2-presenting E2p and I3-01v9 SApNPs

induces more robust GCs in lymph node follicles than the soluble S2GΔHR2 spike.

Immunohistological analysis was performed to characterize GC B cells (GL7$^+$) and T follicular

helper (T$_{fh}$) cells (CD4$^+$Bcl6$^+$). For the I3-01v9 SApNP, 2 weeks after immunization, we

observed robust GCs in lymph node B cell follicles (B220$^+$) with well-formed dark zone (DZ)

and light zone (LZ) compartments, which contain GC B cells, FDCs, and T$_{fh}$ cells (*35, 81-83*)

(**Fig. 6A**). We then extended the analysis to the S2GΔHR2 spike and spike-presenting SApNPs

2, 5, and 8 weeks after the single-dose injection (**Fig. 6B, fig. S7A-C**) and 2 and 5 weeks after

the boost (**Fig. 6C, fig. S7D, E**). Two metrics, the GC/FDC ratio (i.e., whether GC formation is

associated with an FDC network, %) and GC size (i.e., occupied area), were used. Overall, the

soluble spike and both large SApNPs induced robust GCs 2 weeks after immunization (**Fig. 6B,**

fig. S7A). The E2p and I3-01v9 SApNPs that present 20 spikes induced robust, long-lived GCs,

whereas the spike alone failed to sustain robust GCs at week 8 with either the single-dose (**Fig.**

6B, D) or prime-boost (**Fig. 6C, E**) injections. The I3-01v9 SApNP generated larger GCs than

the soluble spike, 2.0-fold larger after the single dose (**Fig. 6B, D**) and 2.4-fold larger after the

boost (**Fig. 6C, E**), measured at week 8.

We further characterized GC reactions by flow cytometry. Fresh mouse lymph nodes

were disaggregated into a single cell suspension and stained with an antibody cocktail to quantify

GC B cells and T_{fh} cells (**fig. S8A**). The results were consistent with the immunohistological

analysis, in which all spike-based vaccine antigens, including the S2GΔHR2 spike and SApNPs,

showed robust GCs at week 2 after the injection that declined over time, as measured at weeks 5

and 8 (**Fig. 6F**). The E2p and I3-01v9 SApNPs generated a larger population of GC B cells than

both the S2P$_{ECTO}$ and S2GΔHR2 spikes at week 2 (**fig. S8B, C**). Although the boost dose had

little impact on the frequency of GC B cells and T_{fh} cells, it appeared to extend GC formation

within lymph nodes (**Fig. 6F, G**), which may promote B cell development toward bNAbs.

Notably, the GC B cell and T_{fh} cell populations elicited by the soluble S2GΔHR2 spike were

barely detectable 5 weeks after immunization (**Fig. 6F, G**). This result was reminiscent of a

recent study of an mRNA vaccine, in which GC reactions diminished to baseline levels at week 4

after a single-dose injection (*84*). The S2GΔHR2-presenting I3-01v9 SApNP generated 3.7/5.2-

fold more GC B cells and 3.7/4.4-fold more T_{fh} cells than the soluble S2GΔHR2 spike at week 8

after one/two-dose immunization (**Fig. 6F, G**). Therefore, SApNPs that were retained on FDC

dendrites could present NAb epitopes to enable more effective B cell recognition than the soluble

spike, and consequently induce more robust and long-lived GCs in lymph nodes. Patterns of

trafficking and retention may be specific to antigen size, as shown previously (*73*) and in the

present study (**Figs. 4** and **5**), but GC reactions are largely determined by vaccine adjuvants. This

effect was briefly demonstrated for the E2p and I3-01v9 SApNPs, which were previously

formulated with the AddaVax and AP adjuvants (*41*). At week 2 after a single-dose injection, the

adjuvanted SApNPs induced stronger GC reactions than the non-adjuvanted groups (**fig. S9**).

This result can also explain the differences in plasma neutralization between the adjuvanted and

non-adjuvanted I3-01v9 SApNPs **(Fig. 2)**.

NGS has been used to assess vaccine-draining lymph node B-cell responses (*85*). Here,

we characterized lymph node B cells at the repertoire level for three groups of mice immunized

with two doses (3.3 μg each) of the S2GΔHR2 spike, E2p, and I301v9 SApNPs via footpad

injections. At this dosage, the spike showed less effective plasma neutralization of variants than

the large SApNPs (**Fig. 1E**). Given their differences in retention, presentation, and GC reaction

(**Figs. 4-6**), they were expected to yield different lymph node B-cell profiles. Interestingly,

antigen-specific sorting identified more spike-targeting lymph node B cells from the I3-01v9

SApNP group than both the spike and E2p SApNP groups (**fig. S10A**). The antibody NGS data

were processed by the mouse antibodyomics pipeline (*69*) (**fig. S10B**) to derive quantitative B-

cell profiles (**fig. S10C-E**). Compared with the spike, the I3-01v9 SApNP appeared to activate

fewer V_H/V_K genes (**fig. S10F**, left two), while generating a larger population of spike-specific

lymph node B cells (**fig. S10A**). The three vaccine groups exhibited a similar degree of SHM for

V_H genes, with the I3-01v9 SApNP showing the highest SHM for V_K genes (**fig. S10F**, middle

two). A highly uniform HCDR3 loop length distribution (~10 aa) was observed for mice in the

I3-01v9 SApNP group with little variation, as measured by the root-mean-square fluctuation

(RMSF) (**fig. S10F**, right two). In our previous studies (*86, 87*), a similar approach was applied

to assess hepatitis C virus (HCV) and Ebola virus (EBOV) vaccine-induced B cell responses in

the spleen, a major lymphoid organ (*88*), after mice received four i.p. injections. We observed

distinct B-cell profiles associated with the viral antigen and NP platform (*86, 87*). Here, the

lymph node B-cell profiles appeared to be rather different, revealing the complex inner workings

of another primary site for vaccine-induced immunity. Notably, I3-01v9 SANP exhibited more

"focused" B-cell activation and development in vaccine-draining lymph nodes, as indicated by

fewer activated germline genes and a narrower HCDR3 length distribution. More in-depth

studies are needed to investigate the effect of injection route, adjuvant, and lymphoid organ, in

addition to viral antigen and NP platform, on the resulting B-cell profiles. Single-cell immune

profiling and antibody isolation (*89*) may provide further insights into the clonality of vaccine-

induced B-cell lineages within lymph nodes.

DISCUSSION

To end the COVID-19 pandemic, vaccines need to effectively block current and emerging

SARS-CoV-2 variants that evade NAb responses by mutating key epitopes on the viral spike

(*31*). To overcome this challenge, some suggested that COVID-19 vaccines need to be updated

on a regular basis (*30-32*), whereas others developed mosaic or cocktail vaccines for related

sarbecoviruses (*46, 90*). These vaccine strategies need to be evaluated for long-term protection,

because SARS-CoV-2 is evolving rapidly and may acquire new mutations to evade vaccine-

induced immunity (e.g., B.1.617) (*11*). In our previous study (*41*), the spike-presenting SApNPs

induced a potent NAb response to SARS-CoV-1, which is evolutionarily much more distant to

the wildtype SARS-CoV-2 strain, Wuhan-Hu-1, than all its circulating variants. Emerging data

from human serum analysis suggested that vaccines derived from early pandemic strains may

provide broad protection against current variants (*33*). Based on these findings, we hypothesized that SApNPs presenting stabilized ancestral Wuhan-Hu-1 spikes may provide an effective vaccine against SARS-CoV-2 variants. In the present study, we sought to confirm this hypothesis by testing four major variants and, if proven true, investigate the mechanism underlying such a broadly protective vaccine.

We explored several critical aspects related to the vaccine response, with a focus on the lead candidate identified in our previous study, S2GΔHR2-10GS-I3-01v9-L7P (*41*). We first tested vaccine-induced mouse plasma, which represents a polyclonal response, against four SARS-CoV-2 variants. Mouse plasma generated previously (*41*) and in new studies using different regimens (e.g., injection route, dosage, and adjuvant) potently neutralized the variants. Notably, SApNPs retained their high ID_{50} titers at a dosage as low as 3.3 μg, whereas formulations with the STING and TLR9 agonists further enhanced the I3-01v9 SApNP-induced neutralizing response. While plasma neutralization data may be interpreted with caution due to assay variation (*44*), single-cell-sorted mAbs provided unambiguous evidence of the vaccine-induced bNAb response. Our results revealed that a plethora of NAb lineages were generated upon vaccination, with I3-01v9 SApNP being the most effective at eliciting bNAbs. Additionally, our results confirmed the necessity of a prime-boost strategy for eliciting a potent NAb response, regardless of the regimen (e.g., injection route, dosage, and adjuvant). Such an NAb response, once generated, can persist for an extended period of time post vaccination. Although SARS-CoV-2 challenge in relevant animal models gives more accurate assessment of vaccine protection (*91*), NAb titers have been found to be highly predictive of immune protection from symptomatic infections in a large cohort study (*92*). Protein vaccines, despite the well-established records of safety and effectiveness, have yet to be deployed to mitigate the

COVID-19 pandemic (*93-95*). One protein vaccine, NVX-CoV2373 (micelle-attached spikes

formulated with the Matrix-M^{TM} adjuvant), showed ~90% efficacy in human trials (*19*). Our

study indicates that SApNPs displaying 20 stabilized spikes provide a promising protein vaccine

candidate that can be used either alone or as a booster for nucleic acid (e.g., mRNA and viral

vector) vaccines in the battle against emerging SARS-CoV-2 variants (*11*).

We explored the mechanism of SApNP *vs.* spike vaccines following the previously used

strategy to analyze the *in vivo* behaviors of antigen-attached gold nanoparticles (*73*). In principle,

SApNP vaccines must induce long-lasting GCs to facilitate the development of bNAbs. Effective

vaccine retention and presentation are critical for inducing and sustaining GC reactions, which in

turn promote the proliferation and affinity maturation of antigen-specific B cells. Indeed, we

found that the I3-01v9 SApNP, our leading vaccine candidate (*41*), elicited 6-fold longer

retention and 4-fold greater accumulation in lymph node follicles than the stabilized S2GΔHR2

spike alone with a prime-boost regimen. This can be attributed to the intrinsic physiological

properties of lymph nodes that mediate vaccine trafficking and retention in follicles in a size-

dependent manner, which would favor retaining large (> 50 nm) virus-like particles (*73-75, 80,*

96). Supporting this notion are the TEM images of retained SApNPs aligned on long FDC

dendrites, suggesting that such protein nanoparticles can present spike antigens to B cells for

rapid initiation and then sustain GC reactions in lymph node follicles for an extended period of

time. Specifically, the I3-01v9 SApNP generated 2.4-fold larger GCs and greater numbers of GC

B cells (5.2-fold) and T_{fh} cells (4.4-fold) than the soluble S2GΔHR2 spike with the prime-boost

regimen. These findings provide quantitative evidence that spike-presenting SApNPs are

uniquely suited for inducing long-lived robust GCs in lymph node follicles. Our analyses thus

shed light on the mechanism by which the I3-01v9 SApNP can elicit a more effective bNAb

response than the soluble spike.

Rational design of next-generation COVID-19 vaccines requires an in-depth

understanding of bNAb elicitation (*31*). Superior NAb (but not necessarily bNAb) responses

have been reported for several vaccine candidates that employ particulate display (*90, 97-104*).

The I3-01v9 SApNP elicited a potent bNAb response to four variants, overcoming a major

challenge facing the current COVID-19 vaccines. Mechanistic studies of vaccine trafficking,

retention, presentation, and GC reactions provided valuable insights into the spike and SApNP-

induced immunity (*95, 105, 106*). Such knowledge, if can be obtained for other vaccine

platforms (e.g., inactivated whole virions, mRNAs, and viral vectors) will facilitate rational

selection of the most effective vaccine candidates to mitigate the pandemic and ultimately stop

the spread of SARS-CoV-2.

MATERIALS AND METHODS

SARS-CoV-2 spike and SApNP vaccine antigens

The design, expression, and purification of a stabilized SARS-CoV-2 spike, S2GΔHR2, and

three SApNPs that present either 8 or 20 S2GΔHR2 spikes were described in our recent study

(*41*). Briefly, the spike gene of the SARS-CoV-2 isolate Wuhan-Hu-1 (GenBank accession no.

MN908947) was modified to include the mutations ^{682}GSAGSV^{687} and K986G/V987G, in

addition to truncation of the HR2 stalk (ΔE1150-Q1208). The viral capsid protein SHP (Protein

Data Bank: 1TD0) was added as a C-terminal trimerization motif to stabilize the S2GΔHR2

trimer, resulting in a soluble S2GΔHR2-5GS-1TD0 spike (*41*). The S2GΔHR2 spike was

genetically fused to ferritin (FR), multilayered E2p, and multilayered I3-01v9 with 5GS, 5GS,

and 10GS linkers, respectively, resulting in three S2GΔHR2-presenting SApNPs (*41*). An

S2P$_{ECTO}$-5GS-1TD0 spike construct that contained the mutations [682]GSAGSV[687] and

K986G/V987G but without HR2 deletion (*41*) was included for comparison. All vaccine

antigens were transiently expressed in ExpiCHO cells and purified by a CR3022 antibody

column and size-exclusion chromatography (SEC) as described previously (*41*). Briefly,

ExpiCHO cells were thawed and incubated with ExpiCHO^{TM} Expression Medium (Thermo

Fisher) in a shaker incubator at 37 °C at 135 rotations per minute (rpm) with 8% CO_2. When the

cells reached a density of 10×10^6 ml^{-1}, ExpiCHO^{TM} Expression Medium was added to reduce

cell density to 6×10^6 ml^{-1} for transfection. The ExpiFectamine^{TM} CHO/plasmid DNA complexes

were prepared for 100-ml transfection in ExpiCHO cells according to the manufacturer's

instructions. For a given construct, 100 μg of plasmid and 320 μl of ExpiFectamine^{TM} CHO

reagent were mixed in 7.7 ml of cold OptiPRO™ medium (Thermo Fisher). After the first feed

on day 1, ExpiCHO cells were cultured in a shaker incubator at 33 °C at 115 rpm with 8% CO_2

according to the Max Titer protocol with an additional feed on day 5 (Thermo Fisher). Culture

supernatants were harvested 13-14 days after transfection, clarified by centrifugation at 4000

rpm for 25 min, and filtered using a 0.45 μm filter (Thermo Fisher). The CR3022 antibody

column was used to extract SARS-CoV-2 antigens from the supernatants, followed by SEC on a

Superdex 200 10/300 GL column (for scaffolded RBD trimers), a Superose 6 16/600 GL column

(for the S2GΔHR2 spike, with and without Avi-tag), or a Superose 6 10/300 GL column (for

SApNPs). Protein concentration was determined using UV$_{280}$ absorbance with theoretical

extinction coefficients.

Animal immunization and sample collection

Similar immunization protocols were reported in our previous vaccine studies (*41, 86, 87*).

Briefly, Institutional Animal Care and Use Committee (IACUC) guidelines were followed for all

of the animal studies. BALB/c mice (6 weeks old) were purchased from the Jackson Laboratory

and kept in ventilated cages in environmentally controlled rooms at The Scripps Research

Institute. The mouse studies were conducted according to Association for the Assessment and

Accreditation of Laboratory Animal Care guidelines, and the protocols were approved by the

IACUC. For the immunogenicity study, the mice were intraperitoneally immunized at weeks 0

and 3 with 200 µl of antigen/adjuvant mix containing 5-50 µg of vaccine antigen and 100 µl of

adjuvant (*41*), or intradermally immunized at weeks 0 and 3 with 80 µl of antigen/adjuvant mix

containing 3.3 µg of vaccine antigen and 40 µl of adjuvant. The intradermal (i.d.) immunization

was done through injections into four footpads, each with 20 µl of antigen/adjuvant mix. For the

mechanistic study of vaccine trafficking, retention, and induced GCs, the mice were immunized

at weeks 0 and 3 with 80 µl of antigen/adjuvant mix containing 40 µg of vaccine antigen per

mouse. To visualize the I3-01v9 SApNPs in lymph node tissues using TEM, each mouse was

immunized at weeks 0 and 3 with 140 µl of antigen/adjuvant mix containing 100 µg of vaccine

antigen (40 µl of adjuvant) into the two hind footpads. Vaccines were intradermally administered

into mouse footpads using a 29-gauge insulin needle under 3% isoflurane anesthesia with

oxygen. Blood was drawn from the maxillary/facial vein into an ethylenediaminetetraacetic acid

(EDTA)-coated tube 2 weeks after each immunization. Plasma was isolated from blood after

centrifugation at 14000 rpm for 10 min. Plasma was heat-inactivated at 56°C for 30 min, with

the supernatant collected after centrifugation at 8000 rpm for 10 min. Plasma was used in

pseudovirus neutralization assays to determine the vaccine-induced NAb responses. The axillary,

brachial, and popliteal sentinel lymph nodes were collected at the end timepoint for further

analysis.

Experimental adjuvants and formulation

The adjuvants squalene-oil-in-water (AddaVax), aluminum hydroxide (AH), aluminum

phosphate (AP), 2'3'-c-di-AM(PS)2 (Rp,Rp) (STING ligand), monophosphoryl lipid A from S.

minnesota (MPLA-SM) R595 (TLR4 agonist), imidazoquinoline compound R848 (TLR7/8

agonist), and CpG ODN 1826, Class B (murine) (TLR9 agonist) were purchased from

InvivoGen. PIKA, a TLR3 agonist with enhanced T cell and antibody responses reported for a

Phase I rabies vaccine trial (*107*), was used as an adjuvant. PIKA was generously provided by

Yisheng Biopharma and included in this study as an adjuvant that activates the TLR3 pathway.

Macrophage inhibitors clodronate liposomes (Liposoma BV, catalog no. CP-005-005) were used

to eliminate subcapsular sinus macrophages in lymph nodes to promote more robust B-cell

activation. Mouse immunization was performed to examine the effects of 16 adjuvants or

adjuvant combinations on the I3-01v9 SApNP-induced immune response with respect to the non-

adjuvanted vaccine (PBS instead of an adjuvant). Vaccine antigen and adjuvants were mixed

thoroughly 10 min before immunization. Each mouse was intradermally immunized at weeks 0,

3, and 6 with 120-140 μl of antigen/adjuvant mix containing 20 μg of vaccine antigen (I3-01v9

SApNP) and 80-100 μl of adjuvant, which was evenly split and injected into four footpads.

Mouse blood was isolated at weeks 5 and 8 after two and three intradermal injections,

respectively. Spleens and lymph nodes were harvested at week 8 for immunological analyses.

Spleen samples were ground through a 70 μm cell strainer to release splenocytes into a cell

suspension. Splenocytes were spun down at $400 \times g$ for 10 min, washed with PBS, and treated

with the ammonium-chloride-potassium (ACK) lysing buffer (Lonza). Splenocytes were then

frozen with 3 ml of Bambanker freezing media.

SARS-CoV-2 pseudovirus neutralization assay

The SARS-CoV-2-pp neutralization assays were described in our previous study (*41*). Briefly,

SARS-CoV-2-pps were generated by the co-transfection of HEK293T cells with the HIV-1

pNL4-3.lucR-E- plasmid (obtained from the National Institutes of Health AIDS reagent program;

https://www.aidsreagent.org/) and the expression plasmid encoding the *S* gene of five SARS-

CoV-2 strains, including the wildtype Wuhan-Hu-1 strain (GenBank accession no. MN908947),

three VOCs (GISAID accession no. EPI_ISL_601443, EPI_ISL_678597, and EPI_ISL_792680

for B.1.1.7, B.1.351, and P.1, respectively), and B.1.617$_{Rec}$, a reconstituted strain based on an

early analysis of the B.1.617 lineage (*11*). The HEK293T-hACE2 cell line (catalog no. NR-

52511) and pcDNA3.1(-) vector containing the *S* gene of the wildtype Wuhan-Hu-1 strain

(catalog no. NR52420) were requested from the BEI Resources (https://www.beiresources.org/)

on September 23, 2020 and used in the pseudovirus neutralization assays (*43*). Based on

sequence alignment, spike mutations were incorporated into the *S* gene of the Wuhan-Hu-1 strain

(catalog no. NR52420) to create respective expression plasmids for B.1.1.7, B.1.351, P.1, and

B.1.617$_{Rec}$. For B.1.617$_{Rec}$, G142D, L452R, E484Q, D614G and P681R were included as

representative spike mutations in this SARS-CoV-2 lineage (*11*). SARS-CoV-2-pp neutralization

by immunized mouse plasma and human or mouse mAbs was performed according to our

previously described protocol (*41*). Using the same co-transfection expression system as

described above for the SARS-CoV-2-pps, we produced pseudoviruses carrying the murine

leukemia virus (MLV) Env, MLV-pps, for use as a negative control (*41*). Percent neutralization

data were analyzed using GraphPad Prism 9.1.2 software. ID$_{50}$/IC$_{50}$ values were calculated using

constraints for percent neutralization (0-100%), whereas unconstrained neutralization plots are

shown in **Fig. 1** and **figs. S1-S3**.

Enzyme-linked immunosorbent assay

Each well of a Costar™ 96-well assay plate (Corning) was first coated with 50 µl of PBS

containing 0.2 µg of the appropriate antigens. The plates were incubated overnight at 4 °C, and

then washed five times with wash buffer containing PBS and 0.05% (v/v) Tween 20. Each well

was then coated with 150 µl of blocking buffer consisting of PBS and 40 mg/ml blotting-grade

blocker (Bio-Rad). The plates were incubated with blocking buffer for 1 h at room temperature,

and then washed five times with wash buffer. Mouse mAbs, in the immunoglobulin G (IgG)

form, were diluted in blocking buffer to a maximum concentration of 10 µg/ml followed by a 10-

fold dilution series. For each dilution, a total volume of 50 µl was added to the appropriate wells.

Each plate was incubated for 1 h at room temperature and then washed five times with PBS

containing 0.05% Tween 20. A 1:5000 dilution of horseradish peroxidase (HRP)-conjugated goat

anti-human IgG antibody (Jackson ImmunoResearch Laboratories) was then made in wash

buffer (PBS containing 0.05% Tween 20), with 50 µl of this diluted secondary antibody added to

each well. The plates were incubated with the secondary antibody for 1 h at room temperature

and then washed six times with PBS containing 0.05% Tween 20. Finally, the wells were

developed with 50 µl of TMB (Life Sciences) for 3-5 min before stopping the reaction with 50 µl

of 2 N sulfuric acid. The resulting plate readouts were measured at a wavelength of 450 nm. The

ELISA data were analyzed to calculate EC_{50} values using GraphPad Prism 9.1.2 software.

Histology, immunostaining, and imaging

The mice were sacrificed 2 h to 8 weeks after a single-dose immunization and 2 h to 5 weeks

after the boost immunization. The axillary, brachial, and popliteal sentinel lymph nodes were

isolated for histological analysis. Fresh lymph nodes were rapidly merged into frozen section

compound (VWR International, catalog no. 95057-838) in a plastic cryomold (Tissue-Tek at

VWR, catalog no. 4565) using liquid nitrogen to preserve antigens on the cell membrane and

spike. Lymph node samples were stored at -80°C and sent to the Centre for Phenogenomics

(http://phenogenomics.ca) on dry ice for sample processing and imaging. Tissue sections (8 μm)

were cut on a cryostat (Cryostar NX70) and collected on charged slides. Sections were post-fixed

in 10% neutral buffered formalin and permeabilized in PBS containing 0.5% Triton X-100

before immunostaining. Protein Block (Agilent) was used to block nonspecific antibody binding

before incubating the sections with primary antibody overnight at 4°C. After washing in TBST,

the sections were incubated in fluorophore-conjugated secondary antibodies for 1 h at room

temperature. Lymph node tissue sections were stained with human anti-spike antibody P2B-2F6

(*72*) (1:50) and biotinylated goat anti-human secondary antibody (Abcam, catalog no. ab7152,

1:300), followed by streptavidin-HRP reagent (Vectastain Elite ABC-HRP Kit, Vector, catalog

no. PK-6100) and diaminobenzidine (DAB) (ImmPACT DAB, Vector, catalog no. SK-4105) to

study the distribution and retention of the soluble S2GΔHR2 spike alone and S2GΔHR2 spike-

presenting E2p and I3-01v9 SApNPs. For immunofluorescent staining, tissue sections were

stained for FDCs using anti-CD21 antibody (Abcam, catalog no. ab75985, 1:1800) followed by

anti-rabbit secondary antibody conjugated with Alexa Fluor 555 (Thermo Fisher, catalog no.

A21428, 1:200), stained for B cells using anti-B220 antibody (eBioscience, catalog no. 14-0452-

82, 1:100) followed by anti-rat secondary antibody conjugated with Alexa Fluor 674 (Thermo

Fisher, catalog no. A21247, 1:200), and stained for subcapsular sinus macrophages using anti-

sialoadhesin (CD169) antibody (Abcam, catalog no. ab53443, 1:600) followed by anti-rat

secondary antibody conjugated with Alexa Fluor 488 (Abcam, catalog no. ab150165, 1:200).

Germinal center B cells were labeled using rat anti-GL7 antibody (FITC; BioLegend, catalog no.

144604, 1:250). T_{fh} cells were labeled using anti-CD4 antibody (BioLegend, catalog no. 100402,

1:100) followed by anti-rat secondary antibody conjugated with Alexa Fluor 488 (Abcam,

catalog no. ab150165, 1:1000) and Bcl6 antibody (Abcam, catalog no. ab220092, 1:300)

followed by anti-rabbit secondary antibody conjugated with Alexa Fluor 555 (Thermo Fisher,

catalog no. A21428, 1:1000). Nuclei were then counterstained with 4′,6-diamidino-2-

phenylindole (DAPI) (Sigma-Aldrich, catalog no. D9542, 100 ng/ml). The stained tissue sections

were scanned using an Olympus VS-120 slide scanner and imaged using a Hamamatsu ORCA-

R2 C10600 digital camera for all bright-field and fluorescent images. Bright-field images of

stained S2GΔHR2 spike and S2GΔHR2 spike-presenting SApNPs in lymph node follicles and

fluorescent images of GCs were quantified using ImageJ software (*108*).

Electron microscopy analysis of protein nanoparticles and lymph node tissues

Electron microscopy (EM) analysis was performed by the Core Microscopy Facility at The

Scripps Research Institute. For the negative-staining EM analysis of protein nanoparticles, the

S2GΔHR2-10GS-I3-01v9-L7P SApNP samples were prepared at a concentration of 0.01 mg/ml.

Carbon-coated copper grids (400 mesh) were glow-discharged, and 10 µl of each sample was

adsorbed for 2 min. Excess sample was wicked away and grids were negatively stained with 2%

uranyl formate for 2 min. Excess stain was wicked away and the grids were allowed to dry. For

the EM analysis of mouse tissues, the lymph nodes were dissected from each animal and

immersed in oxygenated 2.5% glutaraldehyde and 4% paraformaldehyde in 0.1M Na cacodylate

buffer (pH 7.4) fixative overnight at 4ºC . After washing in 0.1 M sodium cacodylate buffer, the

tissue samples were post-fixed in buffered 1% osmium tetroxide and 1.5% potassium

ferrocyanide for 1-1.5 h at 4°C, rinsed in the same buffer, and then stained *en bloc* with 0.5%

uranyl acetate overnight at 4°C. The tissue samples were washed in double-distilled H_2O and

dehydrated through a graded series of ethanol followed by acetone, infiltrated with LX-112

(Ladd) epoxy resin, and polymerized at 60°C. Ultrathin lymph node sections (at 70-nm

thickness) were prepared for imaging. Samples were analyzed at 80 kV with a Talos L120C

transmission electron microscope (Thermo Fisher), and images were acquired with a CETA 16M

CMOS camera.

Lymph node disaggregation, cell staining, and flow cytometry

Germinal center reactions, including the percentage of GC B cells (GL7$^+$B220$^+$) and T$_{fh}$ cells

(CD3$^+$CD4$^+$CXCR5$^+$PD-1$^+$), and the number of GC B cells and T$_{fh}$ cells were studied by flow

cytometry (**fig. S5A**). The mice were sacrificed 2, 5, and 8 weeks after a single-dose

immunization and 2 and 5 weeks after the boost immunization. Fresh axillary, brachial, and

popliteal sentinel lymph nodes were collected and mechanically disaggregated. These lymph

node samples were merged in enzyme digestion solution containing 958 μl of Hanks' balanced

salt solution (HBSS) buffer (Thermo Fisher Scientific, catalog no. 14185052), 40 μl of 10 mg/ml

collagenase IV (Sigma-Aldrich, catalog no. C5138), and 2 μl of 10 mg/ml of DNase (Roche,

catalog no. 10104159001) in an Eppendorf tube. After incubation at 37°C for 30 min, lymph

node samples were filtered through a 70 μm cell strainer and spun down at $400 \times g$ for 10 min.

The supernatant was discarded, and the cell pellet was resuspended in HBSS blocking solution

containing 0.5% (w/v) bovine serum albumin and 2 mM EDTA. The nonspecific binding of Fc

receptors was blocked using anti-CD16/32 antibody (BioLegend, catalog no. 101302) on ice for

30 min. Cocktail antibodies, Zombie NIR live/dead stain (BioLegend, catalog no. 423106),

Brilliant Violet 510 anti-mouse/human CD45R/B220 antibody (BioLegend, catalog no. 103247),

FITC anti-mouse CD3 antibody (BioLegend, catalog no. 100204), Alexa Fluor 700 anti-mouse

CD4 antibody (BioLegend, catalog no. 100536), PE anti-mouse/human GL7 antibody

(BioLegend, catalog no. 144608), Brilliant Violet 605 anti-mouse CD95 (Fas) antibody

(BioLegend, catalog no. 152612), Brilliant Violet 421 anti-mouse CD185 (CXCR5) antibody

(BioLegend, catalog no. 145511), and PE/Cyanine7 anti-mouse CD279 (PD-1) antibody

(BioLegend, catalog no. 135216) were then mixed with the cells and placed on ice for 30 min.

After washing cells with HBSS blocking solution after antibody staining, the samples were fixed

using 1.6% paraformaldehyde (Thermo Fisher Scientific, catalog no. 28906) in HBSS on ice for

30 min. The cell samples were stored in HBSS blocking solution for the flow cytometry study.

Sample events were acquired by a 5-laser BD Biosciences LSR II analytical flow cytometer with

BD FACS Diva 6 software at the Core Facility of The Scripps Research Institute. The data were

further processed using FlowJo 10 software.

DC production, T cell culture, activation, and flow cytometry analysis

Mouse bone marrow (BM) was cultured in RPMI 1640 medium containing 10% fetal bovine

serum (FBS) and recombinant mouse Fms-like tyrosine kinase 3 ligand (Flt3L, 50 ng/ml) and

stem cell factor (SCF, 10 ng/ml) for 9 days as previously described (*109*). To induce DC

activation, immature DCs were incubated with lipopolysaccharide (LPS, 100 ng/ml) plus R848

(Resiquimod, 100 ng/ml) overnight, which activated TLR4 or TLR7/8 signaling, respectively.

Cells were harvested for the experiments. CD11c$^+$ DCs were sorted using magnetic beads

(Miltenyi-Biotech, CA). Splenic mononuclear cells from each group of immunized mice were

cultured in the presence of DCs pulsed with or without I3-01v9 SApNP (1 × 10^{-7} mM) in

complete IMDM medium containing IL-2 (5.0 ng/ml). Cells were collected 16 h later for

intracellular cytokine staining and flow cytometry. All antibodies used for immunofluorescence

staining were purchased from eBioscience (San Diego, CA), BioLegend (San Diego, CA) or BD

Biosciences (San Jose, CA). Magnetic microbead-conjugated streptavidin was purchased from

Miltenyi-Biotech (Auburn, CA). Recombinant human IL-2 protein was purchased from R&D

Systems (Minneapolis, MN). Recombinant mouse Flt3 ligand (Flt3L) and mouse SCF were

purchased from Shenandoah Biotech (Warwick, PA). Cells were stained with appropriate

concentrations of mAbs. Dead cells were excluded using Fixable Viability Dye (eBioscience,

CA). Flow cytometry was performed using LSRII (BD Bioscience, CA).

Bulk and single-cell sorting of SARS-CoV-2 antigen-specific mouse B cells.

Spleens or lymph nodes were harvested from mice 15 days after the last immunization, and the

cell suspension was prepared. Dead cells were excluded by staining with the Fixable Aqua Dead

Cell Stain kit (Thermo Fisher, catalog no. L34957). FcγIII (CD16) and FcγII (CD32) receptors

were blocked by adding 20 µl of 2.4G2 mAb (BD Pharmigen, catalog no. N553142). The cells

were then incubated with 10 µg of a biotinylated RBD-5GS-foldon-Avi trimer or biotinylated

S2GΔHR2-5GS-foldon-Avi spike. Briefly, the probes were generated by the biotinylation of

Avi-tagged SARS-CoV-2 antigens using biotin ligase BirA according to the manufacturer's

instructions (Avidity). Biotin excess was removed by SEC on either a Superdex 200 10/300

column (GE Healthcare) for the RBD probe or a HiLoad Superose 6 16/600 column (GE

Healthcare) for the spike probe. In the SEC profiles, the probe peak was well separated from the

peak of biotin ligase (**fig. S3A**). Cells and biotinylated proteins were incubated for 5 min at 4 °C,

followed by the addition of 2.5 µl of anti-mouse IgG fluorescently labeled with FITC (Jackson

ImmunoResearch catalog no. 115-095-071) and incubated for 15 min at 4 °C. Finally, 5 µl of

premium-grade allophycocyanin (APC)-labeled streptavidin was added to the cells and incubated

for 15 min at 4 °C. In each step, the cells were washed with 0.5 ml of PBS and the sorting buffer

(PBS with 2% FBS). FITC$^+$ APC$^+$ probe-specific B cells were sorted using MoFloAstrios EQ

(Beckman Coulter). For bulk sorting, positive cells were sorted into an Eppendorf microtube

with 20 µl of lysis buffer. For single B-cell sorting, individual positive cells were sorted into the

inner wells of a 96-well plate with 20 µl of pre-reverse transcription (RT) lysis mix containing

0.1 µl of NP40 (Sigma-Aldrich), 0.5 µl of RNAse Inhibitor (Thermo Fisher), 5 µl of 5× First

Strand Buffer, and 1.25 µl of DTT from the SuperScript IV kit (Invitrogen), with 13.15 µl of

H_2O per well.

Antibody cloning from Env-specific single B cells and antibody production.

The antibody cloning of SARS-CoV2-2 antigen-sorted single B cells was conducted as follows.

A mix containing 3 µl of Random Hexamers (GeneLink), 2 µl of dNTPs, and 1 µl of SuperScript

IV enzyme (Thermo Fisher) was added to each well of a single-cell-sorted 96-well plate that

underwent thermocycling according to the program outlined in the SuperScript IV protocol,

resulting in 25 µl of cDNA for each single cell. cDNA (5 µl) was then added to a polymerase

chain reaction (PCR) mix containing 12.5 µl of 2× Multiplex PCR mix (Qiagen), 9 µl of H_2O,

0.5 µl of forward primer mix, and 0.5 µl of reverse mouse primer mix (*110*) for heavy and κ-

light chains within each well. A second PCR reaction was then performed using 5 µl of the first

PCR as the template and respective mouse primers (*110*) according to the same recipe as the first

PCR. The PCR products were run on 1% Agarose gel and those with correct heavy and light

chain bands were then used for Gibson ligation (New England Biolabs), cloning into human IgG

expression vectors, and transformation into competent cells. Mouse mAbs were expressed by the

transient transfection of ExpiCHO cells (Thermo Fisher) with equal amounts of paired heavy and

κ-light chain plasmids. Antibody proteins were purified from the culture supernatant after 12-14

days using Protein A bead columns (Thermo Fisher).

NGS and bioinformatics analysis of mouse B cells.

Previously, a 5′-rapid amplification of cDNA ends (RACE)-PCR protocol was developed for the

deep sequencing analysis of mouse B-cell repertoires (*69*). In the present study, this protocol was

applied to analyze bulk-sorted, RBD/spike-specific mouse B cells. Briefly, 5´-RACE cDNA was

obtained from bulk-sorted B cells of each mouse with the SMART-Seq v4 Ultra Low Input RNA

Kit for Sequencing (TaKaRa). The IgG PCRs were set up with Platinum *Taq* High-Fidelity DNA

Polymerase (Life Technologies) in a total volume of 50 µl, with 5 µl of cDNA as the template, 1

µl of 5´-RACE primer, and 1 µl of 10 µM reverse primer. The 5'-RACE primer contained a

PGM/S5 P1 adaptor, and the reverse primer contained a PGM/S5 A adaptor. The mouse $3´-C_\gamma1-$

$3/3´-C_\mu$ inner primers and $3´-mC_\kappa$ outer primer (*110*) were adapted as reverse primers for the 5´-

RACE PCR processing of heavy and κ-light chains. A total of 25 cycles of PCR was performed

and the expected PCR products (500-600 bp) were gel purified (Qiagen). NGS was performed on

the Ion S5 GeneStudio system. Briefly, heavy and κ-light chain libraries from the same mouse

were quantitated using a Qubit 2.0 Fluorometer with the Qubit dsDNA HS Assay Kit and then

mixed at a 3:1 ratio before being pooled with antibody libraries of other mice at an equal ratio for

sequencing. Template preparation and (Ion 530) chip loading were performed on Ion Chef using

the Ion 520/530 Ext Kit, followed by sequencing on the Ion S5 system with default settings (*86*).

The mouse Antibodyomics pipeline (*69*) was used to process raw NGS data, derive quantitative

profiles for germline gene frequency, the degree of SHM, and CDR3 loop length distribution,

and generate 2D divergence/identity plots to visualize mAbs in their respective repertoires (*86*).

Statistical analysis

Data were collected from 4-7 mice per group. All of the statistical analyses were performed and

graphs were generated using GraphPad Prism 9.1.2 software. In the analysis of vaccine-induced

plasma neutralization, different vaccine groups were compared using one-way analysis of

variance (ANOVA), whereas for a given vaccine group ID_{50} titers of the same plasma sample

against different variants were compared using repeated measures one-way ANOVA. In both

cases, they were followed by Dunnett's multiple comparison *post hoc* test. For the vaccine

accumulation and GC study, different vaccine groups were compared using one-way ANOVA,

followed by Tukey's multiple comparison *post hoc* test. Statistical significance was indicated as

the following: ns (not significant), $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$.

SUPPLEMENTARY MATERIALS

Supplementary material for this article is available at http://xxx/xxx/xxx.

fig. S1. Spike and spike-presenting SApNP vaccine-induced neutralizing antibody responses

against the wildtype SARS-CoV-2 strain and four variants.

fig. S2. Immune responses against the wildtype SARS-CoV-2 strain and four variants induced by

the I3-01v9 SApNP formulated with different adjuvants.

fig. S3. Single-cell isolation and functional evaluation of monoclonal neutralizing antibodies

from mice immunized with the RBD, spike, and SApNP vaccines.

fig. S4. Unbiased repertoire analysis of bulk-sorted SARS-CoV-2 antigen-specific mouse splenic

B cells and tracing of mouse neutralizing antibodies in the NGS-derived repertoires.

fig. S5. SARS-CoV-2 spike-presenting SApNP interaction with macrophages in a lymph node.

fig. S6. TEM images of SARS-CoV-2 spike-presenting I3-01v9 SApNP interaction with FDCs in

a lymph node.

fig. S7. Immunohistological analysis of SARS-CoV-2 spike/spike-presenting SApNP vaccine-

induced GCs.

fig. S8. Flow cytometry analysis of SARS-CoV-2 spike/spike-presenting SApNP vaccine-

induced GCs.

fig. S9. Adjuvant effect on SARS-CoV-2 spike/spike-presenting SApNP vaccine-induced GCs.

fig. S10. NGS analysis of SARS-CoV-2 spike-specific lymph node (LN) B cells from mice

immunized with the spike and SApNP vaccines.

<div style="text-align: center;">References</div>

863
864
1. J. M. Dan *et al.*, Immunological memory to SARS-CoV-2 assessed for up to 8 months
after infection. *Science* **371**, eabf4063 (2021).
2. B. Isho *et al.*, Persistence of serum and saliva antibody responses to SARS-CoV-2 spike
antigens in COVID-19 patients. *Sci. Immunol.* **5**, eabe5511 (2020).
3. A. S. Iyer *et al.*, Persistence and decay of human antibody responses to the receptor
binding domain of SARS-CoV-2 spike protein in COVID-19 patients. *Sci. Immunol.* **5**,
eabe0367 (2020).
4. Y. Chen *et al.*, Quick COVID-19 healers sustain anti-SARS-CoV-2 antibody production.
Cell **183**, 1496-1507.e1416 (2020).
5. R. Burioni, E. J. Topol, Assessing the human immune response to SARS-CoV-2 variants.
Nat. Med., Published Ahead-of-Print (2021).
6. H. Tegally *et al.*, Emergence of a SARS-CoV-2 variant of concern with mutations in
spike glycoprotein. *Nature*, Published Ahead-of-Print (2021).
7. R. Challen *et al.*, Risk of mortality in patients infected with SARS-CoV-2 variant of
concern 202012/1: matched cohort study. *BMJ* **372**, n579 (2021).
8. N. G. Davies *et al.*, Estimated transmissibility and impact of SARS-CoV-2 lineage
B.1.1.7 in England. *Science*, eabg3055 (2021).
9. E. C. Sabino *et al.*, Resurgence of COVID-19 in Manaus, Brazil, despite high
seroprevalence. *Lancet* **397**, 452-455 (2021).
10. J. R. Mascola, B. S. Graham, A. S. Fauci, SARS-CoV-2 viral variants—tackling a
moving target. *JAMA*, Published Ahead-of-Print (2021).
11. S. Cherian *et al.*, Convergent evolution of SARS-CoV-2 spike mutations, L452R, E484Q
and P681R, in the second wave of COVID-19 in Maharashtra, India. *bioRxiv*,
2021.2004.2022.440932 (2021).
12. E. Callaway, Delta coronavirus variant: scientists brace for impact. *Nature*, Published
Ahead-of-Print (2021).
13. C. Zimmer, J. Corum, W. S.-L., in *Coronavirus vaccine tracker*.
(https://www.nytimes.com/interactive/2020/science/coronavirus-vaccine-tracker.html).
14. L. R. Baden *et al.*, Efficacy and safety of the mRNA-1273 SARS-CoV-2 vaccine. *N.
Engl. J. Med.* **384**, 403-416 (2020).
15. T. C. Williams, W. A. Burgers, SARS-CoV-2 evolution and vaccines: cause for concern?
Lancet Respir. Med., Published Ahead-of-Print (2021).
16. D. Y. Logunov *et al.*, Safety and efficacy of an rAd26 and rAd5 vector-based
heterologous prime-boost COVID-19 vaccine: an interim analysis of a randomised
controlled phase 3 trial in Russia. *Lancet* **397**, 671-681 (2021).
17. F. P. Polack *et al.*, Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine. *N.
Engl. J. Med.* **383**, 2603-2615 (2020).
18. M. G. Thompson *et al.*, Prevention and attenuation of Covid-19 with the BNT162b2 and
mRNA-1273 vaccines. *N. Engl. J. Med.*, Published Ahead-of-Print (2021).
19. P. T. Heath *et al.*, Safety and efficacy of NVX-CoV2373 Covid-19 vaccine. *N. Engl. J.
Med.*, Published Ahead-of-Print (2021).
20. Q. Li *et al.*, The impact of mutations in SARS-CoV-2 spike on viral infectivity and
antigenicity. *Cell* **182**, 1284-1294.e1289 (2020).

21. C. Rees-Spear *et al.*, The impact of spike mutations on SARS-CoV-2 neutralization.
bioRxiv, 2021.2001.2015.426849 (2021).
22. C. K. Wibmer *et al.*, SARS-CoV-2 501Y.V2 escapes neutralization by South African
COVID-19 donor plasma. *bioRxiv*, 2021.2001.2018.427166 (2021).
23. E. Andreano *et al.*, SARS-CoV-2 escape in vitro from a highly neutralizing COVID-19
convalescent plasma. *bioRxiv*, 2020.2012.2028.424451 (2020).
24. M. Hoffmann *et al.*, SARS-CoV-2 variants B.1.351 and P.1 escape from neutralizing
antibodies. *Cell*, Published Ahead-of-Print (2021).
25. W. F. Garcia-Beltran *et al.*, Multiple SARS-CoV-2 variants escape neutralization by
vaccine-induced humoral immunity. *Cell*, Published Ahead-of-Print (2021).
26. P. Wang *et al.*, Antibody resistance of SARS-CoV-2 variants B.1.351 and B.1.1.7.
bioRxiv, 2021.2001.2025.428137 (2021).
27. P. Supasa *et al.*, Reduced neutralization of SARS-CoV-2 B.1.1.7 variant by convalescent
and vaccine sera. *Cell*, Published Ahead-of-Print (2021).
28. D. A. Collier *et al.*, SARS-CoV-2 B.1.1.7 sensitivity to mRNA vaccine-elicited,
convalescent and monoclonal antibodies. *medRxiv*, 2021.2001.2019.21249840 (2021).
29. D. Planas *et al.*, Sensitivity of infectious SARS-CoV-2 B.1.1.7 and B.1.351 variants to
neutralizing antibodies. *Nat. Med.* **27**, 917-924 (2021).
30. Z. Wang *et al.*, mRNA vaccine-elicited antibodies to SARS-CoV-2 and circulating
variants. *Nature*, Published Ahead-of-Print (2021).
31. D. R. Burton, E. J. Topol, Variant-proof vaccines - invest now for the next pandemic.
Nature **590**, 386-388 (2021).
32. Q. Li *et al.*, No higher infectivity but immune escape of SARS-CoV-2 501Y.V2 variants.
Cell, Published Ahead-of-Print (2021).
33. C. Liu *et al.*, Reduced neutralization of SARS-CoV-2 B.1.617 by vaccine and
convalescent serum. *Cell*, (2021).
34. A. Singh, Eliciting B cell immunity against infectious diseases using nanovaccines. *Nat.*
Nanotechnol. **16**, 16-24 (2021).
35. G. D. Victora, M. C. Nussenzweig, Germinal centers. *Annu. Rev. Immunol.* **30**, 429-457
(2012).
36. R. Rappuoli, Glycoconjugate vaccines: principles and mechanisms. *Sci. Transl. Med.* **10**,
eaat4615 (2018).
37. M. Akkaya, K. Kwak, S. K. Pierce, B cell memory: building two walls of protection
against pathogens. *Nat. Rev. Immunol.* **20**, 229-238 (2020).
38. O. Bannard, J. G. Cyster, Germinal centers: programmed for affinity maturation and
antibody diversification. *Curr. Opin. Immunol.* **45**, 21-30 (2017).
39. P. A. Robert, A. L. J. Marschall, M. Meyer-Hermann, Induction of broadly neutralizing
antibodies in germinal centre simulations. *Curr. Opin. Biotechnol.* **51**, 137-145 (2018).
40. D. R. Burton, L. Hangartner, Broadly neutralizing antibodies to HIV and their role in
vaccine design. *Annu. Rev. Immunol.* **34**, 635-659 (2016).
41. L. He *et al.*, Single-component, self-assembling, protien nanoparticles presenting the
receptor binding domain and stabilized spike as SARS-CoV-2 vaccine candidates. *Sci.*
Adv. **7**, eabf1591 (2021).
42. D. Wrapp *et al.*, Cryo-EM structure of the 2019-nCoV spike in the prefusion
conformation. *Science* **367**, 1260-1263 (2020).

43. K. H. D. Crawford *et al.*, Protocol and reagents for pseudotyping lentiviral particles with
SARS-CoV-2 spike protein for neutralization assays. *Viruses* **12**, 513 (2020).
44. A. M. Sholukh *et al.*, Evaluation of SARS-CoV-2 neutralization assays for antibody
monitoring in natural infection and vaccine trials. *medRxiv*, 2020.2012.2007.20245431
(2020).
45. D. Pinto *et al.*, Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV
antibody. *Nature* **583**, 290-295 (2020).
46. A. C. Walls *et al.*, Elicitation of broadly protective sarbecovirus immunity by receptor-
binding domain nanoparticle vaccines. *bioRxiv*, 2021.2003.2015.435528 (2021).
47. B. Pulendran, R. Ahmed, Translating innate immunity into immunological memory:
implications for vaccine development. *Cell* **124**, 849-863 (2006).
48. S. G. Reed, M. T. Orr, C. B. Fox, Key roles of adjuvants in modern vaccines. *Nat. Med.*
19, 1597-1608 (2013).
49. B. Pulendran, R. Ahmed, Immunological mechanisms of vaccination. *Nat. Immunol.* **12**,
509-517 (2011).
50. K. A. Fitzgerald, J. C. Kagan, Toll-like receptors and the control of immunity. *Cell* **180**,
1044-1066 (2020).
51. T. Kawai, S. Akira, The role of pattern-recognition receptors in innate immunity: update
on Toll-like receptors. *Nat. Immunol.* **11**, 373-384 (2010).
52. S. P. Kasturi *et al.*, Programming the magnitude and persistence of antibody responses
with innate immunity. *Nature* **470**, 543-547 (2011).
53. T. L. Flach *et al.*, Alum interaction with dendritic cell membrane lipids is essential for its
adjuvanticity. *Nat. Med.* **17**, 479-487 (2011).
54. P. J. Hotez, D. B. Corry, U. Strych, M. E. Bottazzi, COVID-19 vaccines: neutralizing
antibodies and the alum advantage. *Nat. Rev. Immunol.* **20**, 399-400 (2020).
55. R. Cantisani *et al.*, Vaccine adjuvant MF59 promotes retention of unprocessed antigen in
lymph node macrophage compartments and follicular dendritic cells. *J. Immunol.* **194**,
1717-1725 (2015).
56. G. N. Barber, STING: infection, inflammation and cancer. *Nat. Rev. Immunol.* **15**, 760-
770 (2015).
57. B. Guy, The perfect mix: recent progress in adjuvant research. *Nat. Rev. Microbiol.* **5**,
396-397 (2007).
58. M. S. Duthie, H. P. Windish, C. B. Fox, S. G. Reed, Use of defined TLR ligands as
adjuvants within human vaccines. *Immunol. Rev.* **239**, 178-196 (2011).
59. M. P. Steinbuck *et al.*, A lymph node–targeted Amphiphile vaccine induces potent
cellular and humoral immunity to SARS-CoV-2. *Sci. Adv.* **7**, eabe5819 (2021).
60. Y.-N. Zhang, W. Poon, E. Sefton, W. C. W. Chan, Suppressing subcapsular sinus
macrophages enhances transport of nanovaccines to lymph node follicles for robust
humoral immunity. *ACS Nano* **14**, 9478-9490 (2020).
61. J. G. Liang *et al.*, S-Trimer, a COVID-19 subunit vaccine candidate, induces protective
immunity in nonhuman primates. *Nat. Commun.* **12**, 1346 (2021).
62. P. Richmond *et al.*, Safety and immunogenicity of S-Trimer (SCB-2019), a protein
subunit vaccine candidate for COVID-19 in healthy adults: a phase 1, randomised,
double-blind, placebo-controlled trial. *Lancet* **397**, 682-694 (2021).
63. E. S. Rosenberg *et al.*, Vigorous HIV-1-specific CD4+ T cell responses associated with
control of viremia. *Science* **278**, 1447-1450 (1997).

64. L. M. Snell *et al.*, Overcoming CD4 Th1 cell fate restrictions to sustain antiviral CD8 T
cells and control persistent virus infection. *Cell Rep.* **16**, 3286-3296 (2016).
65. J. Zhu, H. Yamane, W. E. Paul, Differentiation of effector CD4 T cell populations (*).
Annu. Rev. Immunol. **28**, 445-489 (2010).
66. S. Kumar *et al.*, Neutralizing antibodies induced by first-generation gp41-stabilized HIV-
1 envelope trimers and nanoparticles. *mBio* **12**, e00429-00421 (2021).
67. R. Shi *et al.*, A human neutralizing antibody targets the receptor-binding site of SARS-
CoV-2. *Nature* **584**, 120-124 (2020).
68. T. F. Rogers *et al.*, Isolation of potent SARS-CoV-2 neutralizing antibodies and
protection from disease in a small animal model. *Science* **369**, 956-963 (2020).
69. C. D. Morris *et al.*, Differential antibody responses to conserved HIV-1 neutralizing
epitopes in the context of multivalent scaffolds and native-like gp140 trimers. *mBio* **8**,
e00036-00017 (2017).
70. F. Chen *et al.*, Functional convergence of a germline-encoded neutralizing antibody
response in rhesus macaques immunized with HCV envelope glycoproteins. *Immunity* **54**,
781-796.e784 (2021).
71. G. P. Wen *et al.*, Quantitative evaluation of protective antibody response induced by
hepatitis E vaccine in humans. *Nat. Commun.* **11**, 3971 (2020).
72. B. Ju *et al.*, Human neutralizing antibodies elicited by SARS-CoV-2 infection. *Nature*
584, 115-119 (2020).
73. Y.-N. Zhang *et al.*, Nanoparticle size influences antigen retention and presentation in
lymph node follicles for humoral immunity. *Nano Lett.* **19**, 7226-7235 (2019).
74. B. A. Heesters, R. C. Myers, M. C. Carroll, Follicular dendritic cells: dynamic antigen
libraries. *Nat. Rev. Immunol.* **14**, 495-504 (2014).
75. J. G. Cyster, B cell follicles and antigen encounters of the third kind. *Nat. Immunol.* **11**,
989-996 (2010).
76. R. Rappuoli, D. Serruto, Self-assembling nanoparticles usher in a new era of vaccine
design. *Cell* **176**, 1245-1247 (2019).
77. C. D. C. Allen, T. Okada, J. G. Cyster, Germinal-center organization and cellular
dynamics. *Immunity* **27**, 190-202 (2007).
78. L. Mesin, J. Ersching, Gabriel D. Victora, Germinal center B cell dynamics. *Immunity* **45**,
471-482 (2016).
79. J. López-Sagaseta, E. Malito, R. Rappuoli, M. J. Bottomley, Self-assembling protein
nanoparticles in the design of vaccines. *Comput. Struct. Biotechnol. J.* **14**, 58-68 (2016).
80. Y. Kato *et al.*, Multifaceted effects of antigen valency on B cell response composition
and differentiation in vivo. *Immunity* **53**, 548-563.e548 (2020).
81. S. Crotty, Follicular helper CD4 T cells (TFH). *Annu. Rev. Immunol.* **29**, 621-663 (2011).
82. J. Merkenschlager *et al.*, Dynamic regulation of TFH selection during the germinal centre
reaction. *Nature* **591**, 458-463 (2021).
83. C. Viant *et al.*, Antibody affinity shapes the choice between memory and germinal center
B cell fates. *Cell* **183**, 1298-1311.e1211 (2020).
84. K. Lederer *et al.*, SARS-CoV-2 mRNA vaccines foster potent antigen-specific germinal
center responses associated with neutralizing antibody generation. *Immunity* **53**, 1281-
1295.e1285 (2020).

85. G. S. Shukla, Y. J. Sun, S. C. Pero, G. S. Sholler, D. N. Krag, Immunization with tumor neoantigens displayed on T7 phage nanoparticles elicits plasma antibody and vaccine-draining lymph node B cell responses. *J. Immunol. Methods* **460**, 51-62 (2018).

86. L. He *et al.*, Single-component multilayered self-assembling nanoparticles presenting rationally designed glycoprotein trimers as Ebola virus vaccines. *Nat. Commun.* **12**, 2633 (2021).

87. L. He *et al.*, Proof of concept for rational design of hepatitis C virus E2 core nanoparticle vaccines. *Sci. Adv.* **6**, eaaz6225 (2020).

88. S. M. Lewis, A. Williams, S. C. Eisenbarth, Structure and function of the immune system in the spleen. *Sci. Immunol.* **4**, (2019).

89. Y. L. Cao *et al.*, Potent neutralizing antibodies against SARS-CoV-2 identified by high-throughput single-cell sequencing of convalescent patients' B cells. *Cell* **182**, 73-84.e16 (2020).

90. A. A. Cohen *et al.*, Mosaic nanoparticles elicit cross-reactive immune responses to zoonotic coronaviruses in mice. *Science* **371**, 735-741 (2021).

91. C. Munoz-Fontela *et al.*, Animal models for COVID-19. *Nature* **586**, 509–515 (2020).

92. D. S. Khoury *et al.*, Neutralizing antibody levels are highly predictive of immune protection from symptomatic SARS-CoV-2 infection. *Nat. Med.*, (2021).

93. R. Rappuoli *et al.*, Vaccinology in the post–COVID-19 era. *Proc. Natl. Acad. Sci. U.S.A.* **118**, e2020368118 (2021).

94. L. Corey, J. R. Mascola, A. S. Fauci, F. S. Collins, A strategic approach to COVID-19 vaccine R&D. *Science* **368**, 948-950 (2020).

95. L. DeFrancesco, Whither COVID-19 vaccines? *Nat. Biotechnol.* **38**, 1132-1145 (2020).

96. T. Tokatlian *et al.*, Innate immune recognition of glycans targets HIV nanoparticle immunogens to germinal centers. *Science* **363**, 649-654 (2019).

97. B. Zhang *et al.*, A platform incorporating trimeric antigens into self-assembling nanoparticles reveals SARS-CoV-2-spike nanoparticles to elicit substantially higher neutralizing responses than spike alone. *Sci. Rep.* **10**, 18149 (2020).

98. A. C. Walls *et al.*, Elicitation of potent neutralizing antibody responses by designed protein nanoparticle vaccines for SARS-CoV-2. *Cell* **183**, 1367-1382.e1317 (2020).

99. A. E. Powell *et al.*, A single immunization with spike-functionalized ferritin vaccines elicits neutralizing antibody responses against SARS-CoV-2 in mice. *ACS Cent. Sci.* **7**, 183-199 (2021).

100. P. J. M. Brouwer *et al.*, Two-component spike nanoparticle vaccine protects macaques from SARS-CoV-2 infection. *Cell* **184**, 1188-1200.e1119 (2021).

101. Y.-F. Kang *et al.*, Rapid development of SARS-CoV-2 spike protein receptor-binding domain self-assembled nanoparticle vaccine candidates. *ACS Nano* **15**, 2738-2752 (2021).

102. T. K. Tan *et al.*, A COVID-19 vaccine candidate using SpyCatcher multimerization of the SARS-CoV-2 spike protein receptor-binding domain induces potent neutralising antibody responses. *Nat. Commun.* **12**, 542 (2021).

103. X. Ma *et al.*, Nanoparticle vaccines based on the receptor binding domain (RBD) and heptad repeat (HR) of SARS-CoV-2 elicit robust protective immune responses. *Immunity* **53**, 1315-1330.e1319 (2020).

104. B. Nguyen, N. H. Tolia, Protein-based antigen presentation platforms for nanoparticle vaccines. *npj Vaccines* **6**, 70 (2021).

105. I. Quast, D. Tarlinton, B cell memory: understanding COVID-19. *Immunity* **54**, 205-210
(2021).
106. M. D. Shin *et al.*, COVID-19 vaccine development and a potential nanomaterial path
forward. *Nat. Nanotechnol.* **15**, 646-655 (2020).
107. L. Wijaya *et al.*, An accelerated rabies vaccine schedule based on toll-like receptor 3
(TLR3) agonist PIKA adjuvant augments rabies virus specific antibody and T cell
response in healthy adult volunteers. *Vaccine* **35**, 1175-1183 (2017).
108. C. A. Schneider, W. S. Rasband, K. W. Eliceiri, NIH Image to ImageJ: 25 years of image
analysis. *Nat. Methods* **9**, 671-675 (2012).
109. K. Mochizuki *et al.*, Programming of donor T cells using allogeneic δ-like ligand 4-
positive dendritic cells to reduce GVHD in mice. *Blood* **127**, 3270-3280 (2016).
110. T. Tiller, C. E. Busse, H. Wardemann, Cloning and expression of murine Ig genes from
single B cells. *J. Immunol. Methods* **350**, 183-193 (2009).
1102

Acknowledgements

Funding: This work was funded by National Institutes of Health grants AI137472, AI139092 (to

J.Z.), Ufovax/SFP-2018-0416, Ufovax/SFP-2018-1013, and Ufovax/SFP-2020-0111 (to J.Z.).

Y.-N.Z. thanks the Natural Sciences and Engineering Research Council of Canada (NSERC) for

a postdoctoral fellowship. We thank V. Bradaschia, K. Duffin, and M. Ganguly at the Centre for

Phenogenomics for their expertise in histology and immunostaining. We acknowledge the expert

assistance of S. Henderson, K. Vanderpool, and T. Fassel at the Core Microscopy Facility at The

Scripps Research Institute. We thank A. Saluk, B. Seegers, and B. Monteverde at the Flow

Cytometry Core Facility of The Scripps Research Institute for their expertise in flow cytometry.

The authors thank M. Arends for proofreading the manuscript. **Author contributions:** Project

design by Y.-N.Z., L.H., and J.Z. SARS-CoV-2 variant plasmid design and processing by L.H.

and C.S. Antigen production, purification, and basic characterization by T.N., T.F., and L.H.

Antibody and mouse plasma neutralization by J.P., T.F., and L.H. Mouse immunization, plasma

collection, and lymph node isolation by Y.-N.Z. Vaccine-induced T-cell response analysis by

Y.W., C.A., and Y.Z. B cell sorting, antibody cloning, and synthesis by C.S. and L.H. Antibody

expression, purification, and ELISA by C.S. and T.N. NGS and bioinformatics by L.H. and J.Z.

Immunohistology, TEM, and flow cytometry by Y.-N.Z. Manuscript written by Y.-N.Z., Y.Z.,

L.H., and J.Z. All authors commented on the manuscript. This is manuscript number 30082 from

The Scripps Research Institute. **Competing interests:** The authors declare no competing

interests. **Data and material availability:** All data are available in the main text or in the

supplementary materials.

1124

Figure Legends

Fig. 1. SApNP vaccines induce broadly neutralizing plasma responses to four

representative SARS-CoV-2 variants. (**A**) Molecular surface representations of two spike

(S2P$_{ECTO}$-5GS-1TD0 and S2GΔHR2-5GS-1TD0) and three spike-SApNP (S2GΔHR2-5GS-

ferritin [FR], S2GΔHR2-5GS-E2p-LD4-PADRE [E2p-L4P], and S2GΔHR2-10GS-I3-01v9-

LD7-PADRE [I3-01v9-L7P]) vaccines. Representative EM image of S2GΔHR2-10GS- I3-01v9-

L7P SApNPs is shown on the right. (**B**) Neutralization of the wildtype Wuhan-Hu-1 strain by

mouse plasma induced by five different vaccines at week 5 after two intraperitoneal (i.p.)

injections ($n = 5$ mice/group). ID$_{50}$ titers derived from SARS-CoV-2-pp neutralization assays are

plotted, with average ID$_{50}$ values labeled on the plots. (**C**) Mouse plasma neutralization against

Wuhan-Hu-1 and the B.1.1.7, B.1.351, P.1, and B.1.617$_{Rec}$ variants at week 5 after two i.p.

injections of the adjuvanted S2GΔHR2-10GS-I3-01v9-L7P vaccine (Left panels 1-5: percent

neutralization plots; Right panel: ID$_{50}$ plot). In (B) and (C), the plasma samples were generated

in our previous study (*41*), in which mice were immunized with 50 μg of adjuvanted vaccine

antigen. (**D**) Neutralization of mouse plasma against the wildtype Wuhan-Hu-1 strain induced by

the S2GΔHR2 spike and two large SApNPs at week 5. Vaccines were administered via

intradermal (i.d.) footpad injections (0.8 μg/injection, for a total of 3.3 μg/mouse). (**E**) Mouse

plasma neutralization against Wuhan-Hu-1 strain and the B.1.1.7, B.1.351, P.1, and B.1.617$_{Rec}$

variants at week 5 after two i.d. footpad injections. (**F**) Neutralization of mouse plasma against

Wuhan-Hu-1 induced by the S2GΔHR2 spike and two large SApNPs at week 26. In (B)-(F), the

ID$_{50}$ values are plotted as mean ± SEM. The data were analyzed using one-way ANOVA for

comparison between different vaccine groups or repeated measures ANOVA for comparison of

ID$_{50}$ titers from the same plasma sample against different SARS-Cov-2 strains. Dunnett's

multiple comparison *post hoc* test was performed. ns (not significant), **$p < 0.01$, ****$p <$

0.0001. (**G**) Neutralization of five SARS-CoV-2 strains by eight human monoclonal antibodies.

The IC$_{50}$ values were calculated with the % neutralization range constrained within 0.0-100.0%

and color-coded (white: IC$_{50}$ > 10 µg/ml; green to red: low to high).

Fig. 2. Adjuvants enhance the I3-01v9 SApNP vaccine -induced plasma neuralization of

both the wildtype strain and four variants. (A) Schematic representation of mouse

immunization with the I3-01v9 SApNP with diverse adjuvant formulations and functional

assessment by SARS-CoV-2-pp neutralization assays and T-cell analysis. Conventional

adjuvants, STING/TLR agonists, macrophage inhibitors, and adjuvant combinations were

compared to non-adjuvanted control (PBS). (**B, C**) Mouse plasma neutralization against the

wildtype SARS-CoV-2 strain, Wuhan-Hu-1, at weeks 5 and 8 after two and three footpad

injections, respectively. ID$_{50}$ titers derived from SARS-CoV-2-pp neutralization assays are

plotted, with average ID$_{50}$ values labeled on the plots. (**D**) Neutralization against four variants by

mouse plasma from STING (top) and CpG (bottom)-formulated vaccine groups. ID$_{50}$ titers

derived from SARS-CoV-2-pp neutralization assays are plotted. Neutralization data were

analyzed using either one-way ANOVA (B-C) or repeated measures one-way ANOVA (D) to

compare ID$_{50}$ titers. Dunnett's multiple comparison *post hoc* test was performed. Splenic

mononuclear cells derived from mice in the STING and CpG groups ($n = 5$ mice/group) at week

8 were cultured in the presence of BALB/C DCs pulsed with I3-01v9 SApNP (1×10^{-7} mM).

Cells were harvested 16 h following re-activation. (**E**) Production of IFN-γ-producing Th1 CD4$^+$

T cells and IL-4-producing Th2 CD4$^+$ T cells. (**F**) IFN-γ-producing CD8$^+$ effector T cells. T-cell

responses were analyzed using one-way ANOVA followed by Tukey's multiple comparison *post*

hoc test. ns (not significant), *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

Fig. 3. Single-cell isolation identifies vaccine-elicited mouse neutralizing antibody lineages

with diverse breadth and potency. (**A**) Genetic analysis of 20 mouse antibodies identified from

M2 in the RBD-5GS-1TD0 trimer group (four), M4 in the S2GΔHR2-5GS-1TD0 spike group

(six), and M2 in the S2GΔHR2-10GS-I3-01v9-L7P SApNP group (ten). Antibodies isolated by

the RBD and spike probes are highlighted in light gray and orange shade, respectively. (**B**)

Neutralization of five SARS-CoV-2 strains by 10 RBD and spike-elicited mouse antibodies (left)

and 10 SApNP-elicited mouse antibodies (right). The IC$_{50}$ values were calculated with the

%neutralization range constrained within 0.0-100.0% and color-coded (white: IC$_{50} > 100$ μg/ml;

green to red: low to high). (**C**) EC$_{50}$ (μg/ml) values of 20 mouse antibodies binding to the two

SARS-CoV-2 antigens, the RBD monomer and S2GΔHR2-5GS-1TD0 spike, both with the

Wuhuan-Hu-1 backbone. Antigen binding was measured by ELISA in duplicate, with mean

value and standard deviation (SD) shown as black and red lines, respectively. (**D**) Divergence-

identity analysis of selected mouse NAbs in the context of RBD/spike-specific splenic B cells.

HCs and KCs are plotted as a function of sequence identity to the template and sequence

divergence from putative germline genes. Color coding denotes sequence density. The template

and sequences identified based on the *V* gene assignment and a CDR3 identity of 90%/85% or

greater to the template are shown as black and orange/magenta dots on the 2D plots, with the

number of related sequences labeled accordingly. The 2D plots for other NAbs are shown in **fig.**

S4D-F.

Fig. 4. SARS-CoV-2 SApNP vaccines induce long-term lymph node follicle retention. (**A**, **B**)

S2GΔHR2-presenting I3-01v9 SApNP vaccine distribution in a lymph node 12 h after (**A**) a

single-dose or (**B**) prime-boost footpad injections (10 μg/footpad, 40 μg/mouse). A schematic

illustration of SApNPs in lymph node follicles is shown. (**C**, **D**) Histological images of the

S2GΔHR2 spike and S2GΔHR2-presenting E2p and I3-01 SApNP vaccine trafficking and

retention in lymph node follicles 2 h to 8 weeks after (**C**) single-dose or (**D**) prime-boost

injections, with a scale bar of 50 μm shown for each image. (**E**, **F**) Quantification of vaccine

accumulation in lymph node follicles 48 h after (**E**) a single-dose or (**F**) prime-boost injections.

Data were collected from more than 10 lymph node follicles (n = 3-4 mice/group). The data

points are expressed as mean ± SD. The data were analyzed using one-way ANOVA followed by

Tukey's multiple comparison *post hoc* test. ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

Fig. 5. SARS-CoV-2 SApNP vaccines interact with follicular dendritic cells (FDCs) and are

presented on FDC dendrites to B cells. (**A**, **B**) S2GΔHR2 spike and S2GΔHR2-presenting E2p

and I3-01 SApNP vaccine interaction with FDC networks in lymph node follicles 12 h after (**A**)

a single-dose or (**B**) prime-boost injections (10 μg/footpad, 40 μg/mouse). Vaccine antigens (the

S2GΔHR2 spike and S2GΔHR2-presenting E2p and I3-01 SApNPs) colocalized with FDC

networks. Immunostaining is color-coded (Green: CD21; Red: CD169; White: anti-spike), with

scale bars of 500 μm and 100 μm shown for a complete lymph node and an enlarged image of a

follicle, respectively. (**C**) Representative TEM images of an FDC surrounded by multiple B cells.

S2GΔHR2-presenting I3-01 SApNPs (yellow arrows) presented on FDC dendrites.

Fig. 6. SARS-CoV-2 SApNP vaccines induce robust long-lived germinal centers. (**A**) Top: Representative immunohistological images of germinal centers at week 2 after a single-dose injection of the S2GΔHR2-presenting I3-01 SApNP vaccine (10 μg/injection, 40 μg/mouse). Bottom: Germinal center B cells (GL7$^+$, red) adjacent to FDCs (CD21$^+$, green) in lymph node follicles (left) and T$_{fh}$ cells in the light zone (LZ) of germinal centers (right). Scale bars of 500 μm and 50 μm are shown for a complete lymph node and an enlarged image of a follicle, respectively. (**B**, **C**) Quantification of germinal center reactions using immunofluorescent images: GC/FDC ratio and sizes of germinal centers 2, 5, and 8 weeks after (**B**) single-dose or (**C**) prime-boost injections ($n = 4$-7 mice/group). The GC/FDC ratio is defined as whether the germinal center formation is associated with an FDC network (%). (**D, E**) Representative immunohistological images of germinal centers in mice immunized using S2GΔHR2 spike or S2GΔHR2-presenting E2p and I3-01 SApNP vaccines at week 8 after (**D**) single-dose or (**E**) prime-boost injections, with a scale bar of 50 μm shown for each image. (**F**, **G**) Quantification of germinal center reactions using flow cytometry: percentage and number of germinal center B cells and T$_{fh}$ cells 2, 5, and 8 weeks after (**F**) single-dose or (**G**) prime-boost injections. The data points are shown as mean ± SD. The data were analyzed using one-way ANOVA followed by Tukey's multiple comparison *post hoc* test for each timepoint. ns (not significant), *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

Figure 2

A

Antibodies identified from SARS-CoV-2 RBD, spike, and nanoparticle-immunized mice by single-cell sorting [a]

Antibody	V_H family [b]	V_H identity [b]	HCDR3 (# aa) [c]	V_K family [b]	V_K identity [b]	KCDR3 (# aa) [c]
TRBD-R-4B6	IGHV2-2*02	97.6%	WGYWYFDV (8)	IGKV8-24*01	98.0%	QQHYSTPLT (9)
TRBD-R-4F3	IGHV1-12*01	95.6%	DDYYALDY (8)	IGKV8-30*01	97.0%	QQYYSYPWT (9)
TRBD-R-4G5	IGHV1-5*01	94.9%	DYYGYGNY (8)	IGKV12-44*01	99.6%	QHHYGTPLT (9)
TRBD-R-5B9	IGHV1S126*01	88.0%	RDYYGNNPFDF (11)	IGKV2-137*01	97.7%	MQHLEYPYT (9)
S2GD-S-1F3	IGHV1-7*01	95.9%	SEYGNLYYAMDY (12)	IGKV3-4*01	97.6%	QQSNEDPLT (9)
S2GD-S-2C10	IGHV1-14*01	98.6%	FTKVEGY (7)	IGKV3-2*01	97.3%	HQSKEVPWT (9)
S2GD-R-1F9	IGHV14-1*02	98.6%	WEGRAMDY (8)	IGKV10-94*01	95.8%	QQYSKLPYT (9)
S2GD-R-2D10	IGHV14-3*02	95.1%	WDNAAYYYGMDY (12)	IGKV14-111*01	95.7%	LQYDELYT (8)
S2GD-R-2E4	IGHV1S81*02	92.8%	YGRYFDC (7)	IGKV14-130*01	98.2%	LQFYEFPYT (9)
S2GD-R-2F4	IGHV1-37*01	95.8%	ERHY (4)	IGKV14-100*01	98.9%	VQYDQFPYT (9)
I3V9-S-1C9	IGHV14-1*02	98.6%	GDFDY (5)	IGKV4-55*01	96.1%	QQWGTYPPRT (10)
I3V9-S-2D8	IGHV14-3*02	95.6%	GLYDYDGSPFAY (12)	IGKV10-96*01	96.8%	QQGNTLPYT (9)
I3V9-S-2D10	IGHV14-1*02	95.5%	GDGNY (5)	IGKV4-57*01	97.8%	QQRSSYPIFT (10)
I3V9-R-1G3	IGHV1S81*02	94.2%	DGSSAY (6)	IGKV9-120*01	97.1%	LQYAGSPLT (9)
I3V9-R-1G9	IGHV1-18*01	93.2%	DGYPYYYAMDY (11)	IGKV6-15*01	97.9%	QQYNSYPWT (9)
I3V9-R-1F5	IGHV1-82*01	92.5%	SWDGLVFAY (9)	IGKV4-55*01	93.3%	QQWNNFPPT (9)
I3V9-R-2E11	IGHV1S81*02	92.2%	DGSIAY (6)	IGKV9-120*01	98.5%	LQYASSPYT (9)
I3V9-R-2F2	IGHV1-18*01	89.4%	DRYDRYFDV (9)	IGKV4-55*01	92.9%	QQWNNFPPT (9)
I3V9-R-2F5	IGHV1-20*02	96.2%	SHDYPFDY (8)	IGKV9-123*01	97.5%	LQRNAYPYT (9)
I3V9-R-2F10	IGHV1-18*01	92.9%	DGYPYYYALDY (11)	IGKV12-98*01	97.6%	QQLYSTPLT (9)

[a] Splenic B cells from three mice immunized with the RBD-5GS-1TD0 trimer, S2GΔHR2-5GS-1TD0 spike, and S2GΔHR2-10GS-I3-01v9-L7P NP were sorted to isolate monoclonal neutralizing antibodies. Antibodies are named as [vaccine]-[probe]-[sorting index]. [vaccine]: TRBD, S2GD, and I3V9 stand for RBD-5GS-1TD0, S2GΔHR2-5GS-1TD0, and S2GΔHR2-10GS-I3-01v9-L7P, respectively. [probe]: R and S stand for RBD-5GS-foldon-Avi-Biot and S2GΔHR2-5GS-foldon-Avi-Biot, respectively. Spike and RBD-sorted antibodies are highlighted in light gray and light brown shade, respectively.
[b] $V_{H/K}$ gene assignment and identity were determined by IGBLAST (https://www.ncbi.nlm.nih.gov/igblast/).
[c] H/KCDR3 sequence and length are determined following the Kabat numbering scheme.

B **RBD and spike-elicited mouse neutralizing antibodies against SARS-CoV-2 wildtype and variants (IC$_{50}$, µg/ml)**

	Antibody	Wuhan-Hu-1 (WT)	B.1.1.7 (Alpha)	B.1.351 (Beta)	P.1 (Gamma)	B.1.617$_{Rec}$
RBD	TRBD-R-4B6	0.11	0.05	>100	>100	>100
	TRBD-R-4F3	4.07	6.22	0.13	0.08	1.68
	TRBD-R-4G5	0.02	0.03	1.16	0.82	0.16
	TRBD-R-5B9	1.00	2.53	1.45	0.69	1.29
Spike	S2GD-S-1F3	0.27	24.13	>100	>100	10.51
	S2GD-S-2C10	0.03	0.02	0.91	1.18	0.16
	S2GD-R-1F9	1.20	0.94	4.02	2.72	>100
	S2GD-R-2D10	16.97	3.62	14.43	1.50	17.57
	S2GD-R-2E4	2.60	1.87	0.24	0.16	0.08
	S2GD-R-2F4	1.92	1.13	3.67	3.87	0.05

Spike nanoparticle-elicited mouse neutralizing antibodies against SARS-CoV-2 wildtype and variants (IC$_{50}$, µg/ml)

	Antibody	Wuhan-Hu-1 (WT)	B.1.1.7 (Alpha)	B.1.351 (Beta)	P.1 (Gamma)	B.1.617$_{Rec}$
I3-01v9 SApNP	I3V9-S-1C9	0.12	0.09	0.05	0.07	71.85
	I3V9-S-2D8	>100	2.63	11.82	6.47	>100
	I3V9-S-2D10	0.09	0.05	0.10	0.18	93.65
	I3V9-R-1F5	9.29	3.08	6.24	3.10	8.35
	I3V9-R-1G3	0.03	0.03	10.43	9.31	0.04
	I3V9-R-1G9	0.02	0.01	0.01	0.01	0.01
	I3V9-R-2E11	0.15	0.05	>100	>100	0.11
	I3V9-R-2F2	3.60	14.35	3.17	1.58	>100
	I3V9-R-2F5	2.58	3.66	3.76	8.76	2.65
	I3V9-R-2F10	54.52	>100	14.10	11.74	77.34

C

Mouse neutralizing antibodies binding to RBD and spike antigens

D

Figure 4

Figure 6