GenTB: A user-friendly genome-based predictor for tuberculosis resistance powered by machine learning

Authors:

Matthias I Gröschel[1], Martin Owens[1], Luca Freschi[1], Roger Vargas Jr[1,2], Maximilian G Marin[1,2], Jody Phelan[3], Zamin Iqbal[4], Avika Dixit[1,5] and Maha R Farhat[1,6]

Affiliations

[1] Department of Biomedical Informatics, Harvard Medical School, Boston, MA, USA

[2] Department of Systems Biology, Harvard Medical School, Boston, MA, USA

[3] Faculty of Infectious and Tropical Diseases, London School of Hygiene & Tropical Medicine, London WC1E 7HT, UK

[4] European Bioinformatics Institute, Hinxton, Cambridge CB10 ISD, UK

[5] Division of Infectious Diseases, Boston Children's Hospital, Boston, MA, USA

[6] Division of Pulmonary and Critical Care Medicine, Massachusetts General Hospital, Boston, MA, USA

Keywords

Tuberculosis, Drug resistance, Drug-susceptibility testing, Diagnostics, Whole Genome Sequencing, Machine Learning, MDR-TB, XDR-TB

Corresspondence

MRF (maha_farhat@hms.harvard.edu) or MIG (matthias_groeschel@hms.harvard.edu)

ABSTRACT

Introduction. Multidrug-resistant *Mycobacterium tuberculosis* (*Mtb*) is a significant global

public health threat. Genotypic resistance prediction from *Mtb* DNA sequences offers an

alternative to laboratory-based drug-susceptibility testing. User-friendly and accurate

resistance prediction tools are needed to enable public health and clinical practitioners to

rapidly diagnose resistance and inform treatment regimens.

Methods. We present Translational Genomics platform for Tuberculosis (GenTB), a web-

based application to predict antibiotic resistance from next-generation sequence data. The

user can choose between two potential predictors, a Random Forest (RF) classifier and a

Wide and Deep Neural Network (WDNN) to predict phenotypic resistance to 13 and 10 anti-

tuberculosis drugs, respectively. We benchmark GenTB's predictive performance along with

leading TB resistance prediction tools (Mykrobe and TB-Profiler) using a ground truth

dataset of 20,408 isolates with laboratory-based drug susceptibility data.

Results. All four tools reliably predicted resistance to first-line tuberculosis drugs but had

varying performance for second-line drugs. The mean sensitivities for GenTB-RF and

GenTB-WDNN across the nine shared drugs was 77.6% (95% CI 76.6 - 78.5%) and 75.4%

(95% CI 74.5 - 76.4%) respectively, and marginally higher than the sensitivities of TB-Profiler

at 74.4% (95% CI 73.4 - 75.3%) and Mykrobe at 71.9% (95% CI 70.9 - 72.9%). The higher

sensitivities were at an expense of ≤1.5% lower specificity: Mykrobe 97.6% (95% CI 97.5 -

97.7%), TB-Profiler 96.9% (95% CI 96.7 to 97.0%), GenTB-WDNN 96.2% (95% CI 96.0 to

96.4%), and GenTB-RF 96.1% (95% CI 96.0 to 96.3%). Genotypic resistance sensitivity was

11% and 9% lower for isoniazid and rifampicin respectively, on isolates sequenced at low

depth (<10x across 95% of the genome) emphasizing the need to quality control input

sequence data before prediction. We discuss differences between tools in reporting results

to the user including variants underlying the resistance calls and any novel or indeterminate

variants

Conclusion. GenTB is an easy-to-use online tool to rapidly and accurately predict

resistance to anti-tuberculosis drugs. GenTB can be accessed online at

https://gentb.hms.harvard.edu, and the source code is available at https://github.com/farhat-

lab/gentb-site.

INTRODUCTION

Human tuberculosis, a chronic infectious disease caused by members of the *Mycobacterium tuberculosis* complex, is a leading cause of death from a bacterial infectious agent [1]. The proliferation of multidrug-resistant tuberculosis (MDR-TB) is threatening TB prevention and control activities worldwide [1]. Timely detection of antimicrobial resistance is vital to guide therapeutic options and contain transmission. Antimicrobial resistance is conventionally determined by *in vitro* drug susceptibility tests (DST) on solid or liquid antibiotic-containing culture, which uses drug-specific testing breakpoints ('critical concentration') to classify the infecting strain into drug-susceptible or drug-resistant [2]. Being contingent on mycobacteria's slow growth rate, these phenotypic tests require days to weeks and often deliver unreliable and poorly reproducible results for some drugs, such as ethambutol and pyrazinamide [3,4]. In contrast, molecular methods have emerged as rapid resistance prediction alternatives to complement and speed up traditional DST, leveraging known and reliable genotype-phenotype relationships between variants in the *M. tuberculosis* genome and *in vitro* drug resistance [5].

Over recent years, whole-genome sequencing (WGS) of *M. tuberculosis* has become an affordable tool to provide genetic information for genotypic resistance prediction and high-resolution outbreak reconstruction [6]. Large scale genotype-phenotype assessments have demonstrated high diagnostic accuracy for clinical use to predict susceptibility to first-line drugs based on WGS [7]. Following these results, public health authorities have begun to discontinue phenotypic testing when pan susceptibility is predicted from the genotype, a step with considerable cost- and time benefits [8]. Start-to-end applications which analyze sequencing data to predict resistance phenotypes and are accessible to non-bioinformatic experts are required as WGS based analyses become part of the standardized diagnostic process in clinical laboratories. A range of published tools available for command-line [9,10] or web-based/desktop use [11–13] or both [14,15] exists. These applications vary in quality control and sequence preprocessing steps and rely on detecting pre-defined resistance-

conferring mutations such as single nucleotide polymorphisms (SNPs) or small

insertions/deletions (indels) in the WGS data to predict the resistance phenotype. They also

vary in the type of information fed back to the user including error rates and specific variants

detected.

Here, we present GenTB (https://gentb.hms.harvard.edu), an open user-friendly start-to-end

application to predict drug resistance phenotypes to 13 drugs from WGS data. The GenTB

analysis pipeline is also available for command-line use wrapped in *Snakemake* [16]. The

online user interface allows users to interactively explore the sequencing data, prediction

results and geographic distributions. Resistance prediction is made based on a previously

observed set of variant positions spanning 18 resistance-associated genetic loci and a

validated random forest (RF) classifier [17] as well as a wide and deep neural network

(WDNN) combining a logistic regression model with a multilayer perceptron to predict the

resistance phenotype [18]. In this study, we benchmark these two classification models

implemented in GenTB along with two other tools with a command-line interface, *TB-profiler,*

and *Mykrobe*, on a large dataset of >20k clinical *M. tuberculosis* isolates starting from raw

Illumina sequence data.

METHODS

Backend and website build

GenTB is a bespoke Django website hosted by the Harvard Medical School O2 high

performance computing environment and collaboratively developed on GitHub

(https://github.com/farhat-lab/gentb-site). The website uses off-the-shelf frontend

components; Bootstrap for styling and mobile-friendly delivery, nvd3 for plots and graphs,

resumable.js for robust uploading and supplements these with custom Javascript

functionality for integration. The backend is a Python-Django web service using a

PostgreSQL database which integrates with Dropbox for file uploading, and python-chore for

slurm cluster job submission and management. GenTB predict jobs are run by modular

programs organized into pipelines. The modularity allows for easy maintenance and

management of dependencies and outputs. Administration screens allow a non-expert

developer design new program calls and construct new pipelines and integrate them without

redeployment of the website. Further tools provide error tracking. GenTB predict results are

integrated into the PostgreSQL database allowing website generated plots to be populated

quickly. All generated files for the intermediary pipeline steps are provided for download by

the user. GenTB Map uses a PostGIS database to rapidly link strain mutation and lineage

information with geo-spatial objects; these are fed into the leaflet.js display to render strain

information to the user. Map allows users to display strain data groupings by country,

lineage, drug resistance phenotype or specific genetic mutation through tabs that can nest

the groups in any order.

Raw read processing

Upon uploading single-end or paired-end FastQ files, GenTB first validates the input using

fastQValidator (Fig. 1). Low-quality reads and sequencing adapters are then trimmed with

fastp [19]. Read mapping taxonomy is assessed with a custom-built *Kraken* database

comprising *M. tuberculosis* complex reference sequences [20] followed by *minimap2*

alignment (parameters: default) of reads to the H37Rv reference genome (AL123456) [21].

Samtools is used for sorting the aligned reads, removing duplicates, and indexing [22].

Sequence read datasets with a coverage of <95% at 10x or less across the genome or that

had a mapping percentage of <90% to *M. tuberculosis* complex strains will not be further

processed, and an error message is displayed to the user. Variants are called with *pilon*

(parameters: default) [23] to obtain SNPs and indels in the variant calling format (VCF)

requiring that they have a PASS or Amb filter tags with read allele frequency >0.40. *Fast-*

Lineage-Caller then detects the *M. tuberculosis* lineage based on five lineage typing

schemes as implemented by Freschi *et al.* [24]. Subsequently, invariant sites in the VCF file

are removed, and a custom Perl script annotates each variant as frameshift, synonymous or

non-synonymous, stop codon, indel along with the H37Rv locus tag for each respective

gene. A custom python script generates a matrix file with all model features/variables in the

columns used as input to the two prediction steps specified below. These scripts are

available from Github (https://github.com/farhat-lab/gentb-site) and are open source

(AGPLv3 license). All intermediate sequence files are accessible to the user for download

and verification.

Operation

Users must create an account to run predictions and track uploaded datasets, intermediary

files and results. Users with low internet bandwidth can use the *Dropbox* integration to

upload files. Both raw sequence reads and variants in variant call format (VCF) can be

uploaded for resistance prediction. The user can select an option to delete uploaded source

data after prediction or otherwise to save it for their future access through GenTB. Files are

user specific and not shared or accessible by others.

GenTB online interface has been tested with batches of up to 300 isolates. For batch

processing of larger numbers of raw sequence data, we provide a command-line GenTB

workflow based on *Snakemake* v5.20.1 [16] where dependent software will be sourced via

conda [25]. The *Snakemake* workflow can be accessed via Github (https://github.com/farhat-

lab/gentb-snakemake). The README file details how resistance prediction results on a

paired-end sample can be obtained.

Genotypic resistance prediction using two statistical models

Two multivariate models are used to predict the resistance phenotype, an RF model

(GenTB-RF) and a WDNN (GenTB-WDNN). GenTB-RF was trained on isolates with

available resistance phenotype data and was validated as previously described [17]. Briefly,

1,397 clinical isolates sampled as detailed in reference [17] underwent targeted sequencing

at 18 drug resistance loci using molecular inversion probes and in parallel underwent binary

drug culture-based DST to 13 drugs. One RF was built for each drug using the

randomForest R package (v. 4.6.7) with a subset of the total 992 SNPs/indels observed.

Variants of highest importance for resistance prediction to each drug were selected by

iteratively paring down the model and measuring loss of performance. Important variants are

shown in Suppl. Figure S1 for isoniazid and rifampicin.

Pyrazinamide resistance is known to rely on a large number of individually rare variants.

Given the large increase in published *M. tuberculosis* WGS and linked DST data as well as

the recent implication of novel resistance loci we retrained the pyrazinamide RF here using a

newer version of randomForest R package (v. 4.6.-14) on variants in the genes *pncA*, *panD*,

clpC1, *clpP* [26]. We used 75% (15,267 isolates) of the dataset to train the model and 25%

(5,098 isolates) to validate its performance. During retraining, we excluded silent variants,

those that occurred only in phenotypically susceptible isolates, or known phylogenetic

variants, and the final model was trained on 393 variants occurring in 3,262 phenotypically

pyrazinamide resistant isolates [24]. We chose the randomForest *mtry* variable that yielded

the smallest out-of-bag error and varied the *classwt* variable to maximize the sum of

sensitivity and specificity.

GenTB-WDNN is a multitask logistic regression model combined with a multilayer

perceptron. It has been previously shown to have equal or higher performance than the RF

architecture when both are trained on the same data [18]. GenTB-WDNN was trained on

3,601 isolates (sampled as detailed in reference [18]) for 11 drugs using the Keras 2.2.4

library in Python 3.6 with a TensorFlow 1.8.0 backend. The model uses 222 features (i.e.,

SNPs or small insertions/deletions) along with derived variables (i.e., the number of non-

synonymous SNPs across all resistance-conferring genes) to predict the resistance

phenotype.

Validation sequencing and phenotype data

We collated a database of 20,408 Illumina raw sequence read datasets for which laboratory-

based phenotypic DST data was available from public sources (Suppl. Table S1). Sequence

data was downloaded from NCBI nucleotide databases. Custom scripts were used to pool

the phenotype data from NCBI, Patric, ReseqTB, and the supplementary information from

published literature (detailed methods in https://github.com/farhat-lab/resdata-ng). Sequence

data was merged in case of multiple sequencing runs per isolate for downstream processing

and resistance prediction. In isolates where >10% of reads did not classify as *M.*

tuberculosis complex, we removed unclassified reads using seqtk

(https://github.com/lh3/seqtk).

Performance of GenTB and comparison with other tools

To assess the performance of GenTB for predicting resistance, all isolates were processed

through the GenTB pipeline. We compared the diagnostic accuracy with two leading

resistance prediction tools, *TB-profiler* 2.8.12 [14] and *Mykrobe* v0.9.0 [15], that were run

with default parameters. The two tools and two GenTB prediction models' predictive ability

were obtained by comparing the genotypic prediction to the phenotype data that was

considered the ground truth. We calculated the true positive rate (sensitivity), the true

negative rate (specificity), and area under the receiver operating curve (AUC for short) to

measure test accuracy for each drug and tool. We evaluated 1,000 probability thresholds per

drug to call resistance or susceptibility for GenTB-RF while using the GenTB-WDNN

thresholds previously described [18] (Suppl. Fig S2 and S3).

Statistical Analyses and data visualization

Prediction files from all tools were parsed and analyzed in Jupyter Notebooks running

Python 3.7 using the Pandas [27] and JSON libraries. Receiver operating characteristic

curves were plotted using the Seaborn library [28]. The Vioplot package was used for violin

plots [29]. Summary tables were created in R version 3.6.3 [30] using the packages from the

tidyverse [31] and kable (https://cran.r-project.org/web/packages/kableExtra/index.html).

Sequencing depth in resistance loci was calculated and plotted using *Mosdepth* version

0.2.9 [32]. Confidence intervals were obtained by bootstrapping, comparing 5000 predictions

per tool and drug on a resampled dataset.

Code and Data Availability

Code is available here: https://github.com/farhat-lab-gentb-site. The *snakemake*

implementation is available here: https://github.com/farhat-lab/gentb-snakemake.

Comparison of output between tools

We collated the output files and information produced by the GenTB online application, the

webserver of TB-Profiler (https://tbdr.lshtm.ac.uk, version 3.0.0), and the Desktop version of

Mykrobe (MacOS app v0.90) using one example raw sequence dataset (accession

ERR1664619). The tools' output was compared based on the following criteria: 1) Type and

accessibility of output data formats; 2) Communication of genotypic prediction results, i.e.

binary classification versus probability; 3) Disclosure of the prediction model's error rate; 4)

Description of known resistance conferring variants identified, 5) Reporting any novel

mutation not listed in the resistance variant database, 6) Detailed account of detected

lineage variants and what lineage typing scheme was used, 7) Report quality metrics on the

input sequence data.

RESULTS

A user-friendly application to analyze *M. tuberculosis* sequencing data

GenTB was developed as a free and benchmarked online application to help public health

and clinical practitioners deconvolute the complexity of *M. tuberculosis* WGS data. *GenTB*

Predict allows users to predict resistance to 13 anti-TB drugs from a clinical isolate's raw

Illumina sequence data (FASTQ). Two validated machine learning models are used to make

predictions: GenTB-RF and GenTB-WDNN (**Methods** and [17,18]). GenTB-RF is the default

prediction model. In addition to the *GenTB Predict* function that we focus on here, the web-

application has additional features for sharing, mapping, and exploring *M. tuberculosis*

genetic and phenotypic data (Fig. 2). *GenTB Data* enables researchers to store, version, and

share *M. tuberculosis* sequence and phenotype data and is powered by the Dataverse

research data repository [33]. Users can select an option to delete source files upon

processing the prediction. *GenTB Map* enables users to geographically visualize genetic and

phenotype data. Users can explore the subset of 20,408 isolates with geographic tags (n=

12,547 isolates) used for GenTB predict validation (**Methods**), or can upload and explore

their own data in enriched-VCF format (https://gitlab.com/doctormo/evcf/-

/blob/master/docs/Enriched_VCF_Format.md). Raw data and results can be exported to a

tabular data format.

Dataset description

We curated a dataset of 20,408 *M. tuberculosis* isolates with known phenotypic resistance

status to benchmark *GenTB Predict* performance (**Methods** and Suppl. Table S1). We

excluded 29 isolates as they failed FastQ validation. Of the remaining, 1,339 isolates did not

pass our taxonomy filter criterion, and their non-*M. tuberculosis complex* reads were

removed. The GenTB pipeline identified an additional 499 isolates where more than 5% of

the genome was covered at depth <10x and these isolates were excluded from further

analysis. These isolates had a median depth of 21x (IQR 17 to 26). The remaining 19,880

isolates with high quality sequencing data were majority lineage 4 (52%), with lesser

representation of lineage 2 (21%), lineage 3 (15%), lineage 1 (10%), *M. bovis* (0.6%),

lineage 6 (0.3%), and lineage 5 (0.2%). Completeness of phenotypic DST data varied by

drug and was highest for the first-line drugs rifampicin (98.3%), isoniazid (96.4%),

ethambutol (77.5%), and pyrazinamide (71.5%) (Suppl. Table S2). The second and third-line

drug phenotype data ranged from 35.1% completeness for streptomycin to 7.8% for

ethionamide. Of the 20,408 isolates, 13,817 were phenotypically susceptible to first line

drugs, 4,743 (23.3%) were phenotypically MDR (i.e., resistant to isoniazid and rifampicin)

and 396 (1.9%) were phenotypically XDR (MDR and resistant to fluoroquinolones and the

second-line injectables – amikacin, kanamycin or capreomycin). We ran GenTB-RF and

GenTB-WDNN to predict resistance on 19,880 isolates and compared the predictions to

phenotypic data.

Predictive performance of the GenTB-Random Forest

We assessed each tools' predictive performance by comparison with phenotypic culture-

based DST results. Overall, the four tools had comparable performance characterized by

varying sensitivities and high specificities (Tables 1 & 2, Fig 3A). Diagnostic performance

was better for first-line than second-line drugs. As sensitivity varied most widely, we discuss

it by drug class below. Specificities varied less by tool or by drug. GenTB-RF's diagnostic

specificity was >92% for all drugs including the second-line injectables and fluoroquinolones

with the exception ethionamide (specificity = 78% [95% CI 75-80]) and streptomycin

(specificity = 89% [95% CI 88-90]). GenTB-RF's specificities were similar or higher than the

other three tools with the exception of pyrazinamide (94% [95% CI 93-95]) and streptomycin

(89% [95% CI = 88-90]) compared to TB-Profiler (96% and 95%, respectively) as well as

Mykrobe (98% and 95%, respectively).

First-line drugs: Rifampicin resistance prediction by GenTB-RF was most accurate

compared to other tools: AUC 0.96 (95% CI = 0.95-0.96), sensitivity 93% (95% CI = 93-94),

second highest sensitivity was for TB-Profiler at 92% (95% CI = 91-93) (Tables 1 & 2, Figure

4). The accuracy of isoniazid resistance prediction was high and comparable across three of

the four tools including GenTB-RF (sensitivity 91% [95% CI =91-92]). For ethambutol,

GenTB-RF and TB-Profiler had the best and comparable performance with sensitivity 86%

(95% CI =85-87).

GenTB-RF predictions for pyrazinamide using the original model (v1.0) had low sensitivity at

56% (95% CI 54-58) with adequate specificity (98% [95% CI = 98-99]) compared to the other

tools when evaluated on the 19,880 isolates (2,336 phenotypically resistant and 11,932

susceptible) [17]. Pyrazinamide resistance is known to be caused by a large number of

individually rare variants in the gene *pncA* [34]. Given the large interval increase in available

WGS data and recent implication of novel resistance loci (*panD, clpC1, clpP*) [26] since

GenTB-RF was last trained, we assessed the number of rare variants in the four

aforementioned genes linked to pyrazinamide resistance. In a random 75% subset of the

20,379 isolates, we detected a total of 393 different variants in *pncA, panD, clpC1 and clpP*

with 40% (158/393) occurring only once. The majority of these variants, i.e., 73% (285/393)

were not previously seen by the original model. As a result of these observations, we

retrained a GenTB-RFv2.0, on 75% of the data using all 393 non-synonymous variants

including singletons and insertion/deletion variants from *pncA, panD, clpC1* and *clpP*. The

retrained model, when benchmarked on an independent validation dataset of 5,098 isolates,

offered a sensitivity similar to that of the other tools (79%, 76 to 83) (Table 1).

Second-line drugs: For second-line drugs, larger discrepancies between genotype and

resistance phenotype have been previously described compared with first-line drugs [14,15].

Resistance to the second-line injectable drugs amikacin and kanamycin ranged between 63-

68% across the four tools, with the exception of a sensitivity of 55% by TB-Profiler for

amikacin (Table 1). For the fluoroquinolone ofloxacin, sensitivity ranged from 62%-68%

across the four tools. Three drugs had too few isolates with known phenotypic resistance

(ciprofloxacin [n = 63], levofloxacin [n =111], and para-aminosalicylic acid [n = 46]), and

hence the tool's predictions had wide confidence intervals for these drugs (Supplemental

Tables S3 and S4).

Predictive performance of GenTB-WDNN.

Similar to GenTB-RF, the overall GenTB-WDNN performance was marked by high prediction

accuracy of first-line drug resistance and lower accuracy of second-line resistance (Table 1).

AUC 95% CI overlapped for all drugs between the two models except for ofloxacin and

rifampicin for which the GenTB-RF AUC was higher (Table 2). For streptomycin the GenTB-

WDNN offered the best sensitivity and specificity of all four models (sensitivity 87%, 95% CI

85-88%, specificity 87% (95%CI 86-88%). Specificities were >95% for all drugs except for

streptomycin (87%, 95% CI 85 to 88) and ethambutol (93%, 95% CI 93 to 94).

Predictive performance depends on sequencing depth

We evaluated the need for quality control on sequencing depth as several tools do not

currently implement this prior to resistance prediction [9,14,15]. We observed predictive

performance to be highly dependent on sequencing depth as indicated by lower sensitivity to

predict rifampicin or isoniazid resistance by all four tools for the 499 isolates that did not

meet the threshold of ≥10x depth across >95% of the genome (median depth of 21x, IQR 17

to 26, Figures 3E,3F). Using GenTB-RF, the mean sensitivity of isoniazid and rifampicin

prediction was 84.6% (SD 3.6) and 87.3% (SD 3.6) respectively among low-depth isolates,

compared with 91% and 93%, respectively, on high-depth isolates (Suppl. Table S5, Figures

3E, 3F). Loss of sensitivity due to low sequencing depth was comparable across the four

tools.

Discordant resistance predictions

To gain insight into model performance, we probed discrepancies between GenTB-RF's

genotype-based prediction and the resistance phenotype. We focused on this model as it

had the highest overall sensitivity. We examined specifically rifampicin and isoniazid as

resistance to these two drugs defines MDR-TB, and their genetic resistance mechanisms

are well understood. We investigated isolates for which GenTB-RF predicted resistance

while the phenotype was reported as susceptible (false positives) and isolates for which

GenTB-RF predicted susceptibility with a resistant phenotype (false negatives). We

confirmed that false negative predictions were not due to low sequencing depth in relevant

drug resistance loci (*i.e.* that depth was ≥10x across all bases, Suppl. Figures S4 and S5).

Rifampicin false positives: Variants causative of rifampicin resistance are concentrated in a

81bp window in the *rpoB* gene *a.k.a* the rifampicin resistance determining region (RRDR,

H37Rv coordinates 761081 to 761162, accession AL123456) [35]. For rifampicin, we

observed 254 false positive predictions (phenotypically susceptible isolates predicted

resistant). GenTB-RF detected one or more non-silent RRDR variants in 198 of these 254

isolates (78%). The most common RRDR variants were S450L (occurred in 49/254 isolates),

L430P (in 33/254), and H445N (in 31/254) (Suppl. Table S6). The remaining 56 of 254

isolates, harbored non-RRDR variants, the two most common were *rpoB* I491F (occurred in

29/56) and *rpoB* V695L (occurred in 24/56). Twenty eight of the 56 isolates (50%) were

phenotypically resistant to isoniazid and a further 16 (29%) were resistant to ethambutol.

Rifampicin false negatives: Among the 333 false negative rifampicin predictions

(phenotypically resistant isolates predicted susceptible), 96 (29%) isolates harbored a

variant in *rpoB* and of these 75 (23% of the 333) were in the RRDR (Suppl. Table S6). These

included most commonly three base pair insertion in *rpoB* codon 433 (occurred in 14/333

isolates) and *rpoB codon* 443 (occurred in 9/333 isolates) and *rpoB* substitution Q432L (in

9/333) [36]. These *rpoB* variants were not previously seen by the GenTB-RF model when

initially trained. For the remaining 237 of 333 isolates (71%) phenotypic resistance remained

unexplained.

<u>Isoniazid false positives:</u> For isoniazid, we observed 315 false positive predictions

(phenotypically susceptible isolates predicted resistant by GenTB-RF). Among these

isolates, 119/315 (38%) had a total of 40 unique non-silent non-lineage variants in genes

linked to isoniazid resistance (*inhA*, *katG*, *ahpC*, *fabG1*) (Suppl. Table S7). Most variants,

36/40, were rare, occurring in only 2 or fewer isolates. Five out of the 40 unique mutations

detected in 75/315 (24%) isolates are considered important for isoniazid resistance

prediction by GenTB-RF [17]. The most frequent INH resistance variants were the canonical

isoniazid resistance mutation *katG* S315T [37] (occurred in 56/315 isolates) and non-silent

variants at *inhA* codon 94 (occurred in 14/315 isolates). Seventy-six of the 315 (24%)

apparent false positive isolates were phenotypically resistant to rifampicin and 189 (60%)

isolates had a phenotypic resistance to at least one other drug.

<u>Isoniazid false negatives:</u> Among the 518 false negative isoniazid predictions (phenotypically

resistant isolates predicted susceptible by GenTB-RF), 194/518 (37%) harbored non-silent

variants in isoniazid resistance associated genes (Suppl. Table S7). Only 13 of the 139

unique variants observed in the 518 isolates were seen before by GenTB-RF and none of

these were considered important isoniazid resistance mutations. *KatG* W328L was the

variant detected most frequently (occurred in 10/518 isolates predicted false negative) and

although not previously seen by GenTB-RF was described to occur in 0.2% of isoniazid

resistance in one study [38]. Most variants linked to isoniazid resistance observed in these

isolates were rare, i.e., 134/139 (96%) occurred in ≤ 3 isolates.

Output comparison across the three tools

All four tools are accessible to the non-experienced user via either an online interface

(GenTB, TB-Profiler) or via a Desktop application. We compared each tool's output using the

criteria specified in **Methods** (Table 3). GenTB-RF provides a heatmap indicating the

probability of resistance including the models' error rate with all prediction and intermediary

files available for download. TB-Profiler and Mykrobe present binary (resistant or

susceptible) predictions in overview tables with download options in CSV or JSON formats,

respectively. TB-Profiler and GenTB present resistance causing variants and variants not

associated with resistance. All tools provide the lineage call made but GenTB also specifies

the lineage typing schemes used.

392   DISCUSSION

The increasing affordability of WGS and our improving comprehension of mycobacterial drug

resistance mechanisms has placed sequencing at the forefront of *M. tuberculosis* resistance

diagnosis in clinical and public health laboratories (*e.g.* Public Health England in the United

Kingdom and the Centers for Disease Control and Prevention in the United States) [7,39].

Yet, the complexity of resistance biology is such that large and diverse bacterial isolate

datasets are needed to confirm the accuracy of genotype-based resistance prediction and its

generalizability. Further, the required computational resources and knowledge to conduct

sequencing analysis prohibit both the access to and confidence in WGS based resistance

prediction in clinics in both low- and high-incidence settings. High confidence automated

tools that are systematically benchmarked on diverse datasets are needed to facilitate

adoption, and to act as the standard for future tool development and regulation by oversight

agencies such as the World Health Organization (WHO).

GenTB is an automated open tool for resistance prediction from WGS. Here we

benchmarked its two prediction models against two other leading TB prediction tools. Both

GenTB models predicted resistance and susceptibility against first-line drugs with high

accuracy. Predictive performance for second line drugs showed lower sensitivity, although

with high specificity for some of those drugs, i.e., capreomycin, kanamycin, and ofloxacin.

This high specificity may be used to rule out resistance when no resistance conferring

variant for these drugs was found. A detailed analysis of discrepant predictions made by

GenTB-RF illustrated that a number of false positive predictions were supported by

canonical resistance variants, e.g., non-silent mutation in the *rpoB* RRDR in case of

rifampicin, suggesting that their phenotypes were erroneously labeled as susceptible.

Similarly, nearly half (48%) of the variants found in isoniazid false positive predictions are

canonical resistance variants. These isoniazid resistance variants, the large proportion

(60%) of phenotypic resistance to another drug among these isolates, and the knowledge

that isoniazid is usually a gateway drug resistance, suggest that some phenotypes were

erroneously characterized as susceptible [40]. Accordingly, specificity of genotype-based

prediction in practice maybe even higher than reported here (Table 1).

For isolates with a resistant rifampicin phenotype that were predicted susceptible by GenTB-

RF, we found a mutation in the *rpoB* RRDR in a nearly a quarter (23%) of isolates that

reasonably accounts for the resistance phenotype, but had not been seen by the model

previously. For the remaining majority of false negatives (71% for rifampicin) no relevant

resistance variant was found. In these cases, phenotypic resistance remained unexplained

and could be due to erroneous phenotypes or yet unknown resistance mechanisms. For

isolates with a resistant isoniazid phenotype predicted susceptible, no important resistance

conferring mutations were found. In these cases, phenotypic resistance could be due to rare

and yet undescribed resistance variants. A substantial proportion of false negative

predictions to isoniazid or rifampicin had genotypic resistance to at least another drug (48%

of rifampicin false negatives and 40% of isoniazid false negatives). These observations

overall suggest that a viable option to reduce false negative predictions by current models

would be to leverage genotypic predictions to other drugs and flag such isolates for

complementary phenotypic DST. In the future as new larger datasets of paired genotype and

resistance phenotype are curated, e.g. by efforts sponsored by the WHO [41], retraining

existing resistance prediction models will improve diagnostic sensitivity.

The final output produced by the four tools varies in terms of detail and type of variants

reported with GenTB providing the most detail. GenTB's output reports novel variants not

linked to resistance in addition to those that are resistance associated. The phylogenetic

lineage calling procedure implemented in GenTB [24] uses currently available typing

schemes, including the spoligotype nomenclature, to facilitate comparisons across lineage

schemes.

Unlike other published resistance prediction tools that rely on a curated list of resistance

conferring mutations that call resistance when a specific variant is present, GenTB-RF and

GenTB-WDNN use multivariable statistical models to predict resistance phenotype. These

models are better suited to account for the complex relationships between resistance

genotype and phenotype. Among the advantages of multivariate prediction models is that

relationships between variables are taken into account as both individual variants and gene-

gene interactions cause phenotypic drug resistance. As such, the two models provide a

probability value that a given isolate is resistant or susceptible rather than a binary

classification. This is relevant in case of variants that, if present alone, confer only weak to

no resistance, but may confer complete resistance if present in combination. Also, each

variable in a multivariable model has different weights depending on the strength of

association with resistance in the training data, reflecting the biological reality where variants

cause differing levels of resistance. The benchmarking data presented here confirm that

these multivariate models offer gains in sensitivity over the other two tools that use curated

mutation lists, however this comes at a small decrease in specificity overall. Seen its higher

overall performance GenTB-RF is currently implemented as the default prediction model. As

larger and more diverse data will become available for model training, especially for

prediction of resistance more quantitatively, i.e., to predict minimum inhibitory concentrations

or MICs, we anticipate multivariate models including the more complex GenTB-WDNN

architecture to have an even bigger advantage over direct association of mutation lists.

This study was not without limitations. An important prerequisite for reliable genotypic

resistance prediction is the quality of the raw sequencing data. Variants and small indels in

resistance conferring genes can be accurately and confidently called from Illumina raw

sequence data if the genes are adequately covered at an acceptable sequencing depth

[Marin *et al.*, in preparation]. However, short-read sequencing data is recognized to have

lower sensitivity for detecting more complex genomic variants including long indels or

structural variation and these may have been missed in this study. But these latter types of

variants are expected to be rare. Our finding of 'apparent' false positive predictions (i.e.,

resistance call by GenTB-RF while susceptible phenotype) in isolates harboring canonical

resistance variants portends some erroneous phenotypes in our ground truth dataset. Due to

the scale and public nature of the dataset used for benchmarking in this study, we were

unable to retest the laboratory-based drug susceptibility profiles of isolates with discordant

predictions, but hope that it provides a test closer to a 'real-world' scenario for these tool's

application.

CONCLUSION

The rapid emergence and affordability of sequencing of *M. tuberculosis* along with the herein

confirmed high accuracy of several genotypic resistance prediction tools supports the use of

informatically assisted treatment design in the clinical setting. Independent benchmarking

efforts will facilitate regulatory reviews and assessments and build confidence in the tools'

performances. As genotypic resistance predictions will accompany and increasingly replace

laboratory-based resistance phenotyping performance criteria will need to be defined to

guide clinical and public health laboratories in their use. Lastly, it will be important to

communicate the confidence and uncertainty that is inherent to all genotypic predictions to

clinicians, and provide clear diagnostic algorithms in case of genotype-phenotype

discordances.

REFERENCES

1. World Health Organization. Global Tuberculosis Report 2020 [Internet]. World Health Organization; 2020. Available from: https://www.who.int/publications/i/item/9789240013131

2. World Health Organization. Guidelines for surveillance of drug resistance in tuberculosis 5th Edition [Internet]. WHO; 2015. Available from: https://apps.who.int/iris/bitstream/handle/10665/174897/;jsessionid=52537DA4A0AB0E19A1 0382076AC23874?sequence=1

3. World Health Organization. Implementing tuberculosis diagnostics: A policy framework [Internet]. WHO; 2015. Available from: http://www.who.int/tb/publications/implementing_TB_diagnostics/en/

4. World Health Organization. Technical manual for drug susceptibility testing of medicines used in the treatment of tuberculosis [Internet]. WHO; 2018. Available from: https://apps.who.int/iris/bitstream/handle/10665/275469/9789241514842-eng.pdf?ua=1

5. Cirillo DM, Miotto P, Tortoli E. Evolution of Phenotypic and Molecular Drug Susceptibility Testing. Adv Exp Med Biol. 2017;1019:221–46.

6. Meehan CJ, Goig GA, Kohl TA, Verboven L, Dippenaar A, Ezewudo M, et al. Whole genome sequencing of Mycobacterium tuberculosis: current standards and open issues. Nat Rev Microbiol [Internet]. 2019; Available from: https://doi.org/10.1038/s41579-019-0214-5

7. CRyPTIC Consortium and the 100,000 Genomes Project, Allix-Béguec C, Arandjelovic I, Bi L, Beckert P, Bonnet M, et al. Prediction of Susceptibility to First-Line Tuberculosis Drugs by DNA Sequencing. N Engl J Med. 2018;379:1403–15.

8. McNerney R, Zignol M, Clark TG. Use of whole genome sequencing in surveillance of drug resistant tuberculosis. Expert Rev Anti Infect Ther. 2018;16:433–42.

9. Kohl TA, Utpatel C, Schleusener V, De Filippo MR, Beckert P, Cirillo DM, et al. MTBseq: a comprehensive pipeline for whole genome sequence analysis of Mycobacterium tuberculosis complex isolates. PeerJ. 2018;6:e5895.

10. Steiner A, Stucki D, Coscolla M, Borrell S, Gagneux S. KvarQ: targeted and direct variant calling from fastq reads of bacterial genomes. BMC Genomics. 2014;15:881.

11. Feuerriegel S, Schleusener V, Beckert P, Kohl TA, Miotto P, Cirillo DM, et al. PhyResSE: a Web Tool Delineating Mycobacterium tuberculosis Antibiotic Resistance and Lineage from Whole-Genome Sequencing Data. J Clin Microbiol. 2015;53:1908–14.

12. Iwai H, Kato-Miyazawa M, Kirikae T, Miyoshi-Akiyama T. CASTB (the comprehensive analysis server for the Mycobacterium tuberculosis complex): A publicly accessible web server for epidemiological analyses, drug-resistance prediction and phylogenetic comparison of clinical isolates. Tuberculosis . 2015;95:843–4.

13. Sekizuka T, Yamashita A, Murase Y, Iwamoto T, Mitarai S, Kato S, et al. TGS-TB: Total Genotyping Solution for Mycobacterium tuberculosis Using Short-Read Whole-Genome Sequencing. PLoS One. 2015;10:e0142951.

14. Phelan JE, O'Sullivan DM, Machado D, Ramos J, Oppong YEA, Campino S, et al. Integrating informatics tools and portable sequencing technology for rapid detection of resistance to anti-tuberculous drugs. Genome Med. 2019;11:41.

15. Hunt M, Bradley P, Lapierre SG, Heys S, Thomsit M, Hall MB, et al. Antibiotic resistance
prediction for *Mycobacterium tuberculosis* from genome sequence data with Mykrobe.
Wellcome Open Res. 2019;4:191.

16. Köster J, Rahmann S. Snakemake—a scalable bioinformatics workflow engine.
Bioinformatics. Oxford Academic; 2012;28:2520–2.

17. Farhat MR, Sultana R, Iartchouk O, Bozeman S, Galagan J, Sisk P, et al. Genetic
Determinants of Drug Resistance in Mycobacterium tuberculosis and Their Diagnostic Value.
Am J Respir Crit Care Med. 2016;194:621–30.

18. Chen ML, Doddi A, Royer J, Freschi L, Schito M, Ezewudo M, et al. Beyond multidrug
resistance: Leveraging rare variants with machine and statistical learning models in
Mycobacterium tuberculosis resistance prediction. EBioMedicine. 2019;43:356–69.

19. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics. 2018;34:i884–90.

20. Wood DE, Salzberg SL. Kraken: ultrafast metagenomic sequence classification using
exact alignments. Genome Biol. 2014;15:R46.

21. Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics.
2018;34:3094–100.

22. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25:2078–9.

23. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, et al. Pilon: an
integrated tool for comprehensive microbial variant detection and genome assembly
improvement. PLoS One. 2014;9:e112963.

24. Freschi L, Vargas R Jr, Hussain A, Kamal SMM, Skrahina A, Tahseen S, et al.
Population structure, biogeography and transmissibility of Mycobacterium tuberculosis
[Internet]. bioRxiv. bioRxiv; 2020. Available from:
http://dx.doi.org/10.1101/2020.09.29.293274

25. Grüning B, Dale R, Sjödin A, Chapman BA, Rowe J, Tomkins-Tinch CH, et al. Bioconda:
sustainable and comprehensive software distribution for the life sciences. Nat Methods.
Springer Science and Business Media LLC; 2018;15:475–6.

26. Gopal P, Sarathy JP, Yee M, Ragunathan P, Shin J, Bhushan S, et al. Pyrazinamide
triggers degradation of its target aspartate decarboxylase. Nat Commun. Springer Science
and Business Media LLC; 2020;11:1661.

27. McKinney W. Data Structures for Statistical Computing in Python. Proceedings of the 9th
Python in Science Conference [Internet]. SciPy; 2010. Available from:
http://dx.doi.org/10.25080/majora-92bf1922-00a

28. Waskom M, Botvinnik O, O'Kane D, Hobson P, Lukauskas S, Gemperline DC, et al.
Mwaskom/Seaborn: V0.8.1 (September 2017) [Internet]. Zenodo; 2017. Available from:
http://dx.doi.org/10.5281/zenodo.883859

29. Adler D, Kelly ST. vioplot: violin plot [Internet]. 2020. Available from:
https://github.com/TomKellyGenetics/vioplot

30. Team RC, Others. R: A language and environment for statistical computing [Internet].
Vienna, Austria; 2013. Available from: http://cran.univ-

paris1.fr/web/packages/dplR/vignettes/intro-dplR.pdf

31. Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, et al. Welcome to
the Tidyverse. JOSS. 2019;4:1686.

32. Pedersen BS, Quinlan AR. Mosdepth: quick coverage calculation for genomes and
exomes. Bioinformatics. Oxford University Press (OUP); 2018;34:867–8.

33. King G. An introduction to the dataverse network as an infrastructure for data sharing.
Sociol Methods Res. SAGE Publications; 2007;36:173–99.

34. Yadon AN, Maharaj K, Adamson JH, Lai Y-P, Sacchettini JC, Ioerger TR, et al. A
comprehensive characterization of PncA polymorphisms that confer resistance to
pyrazinamide. Nat Commun. Springer Science and Business Media LLC; 2017;8:588.

35. Donnabella V, Martiniuk F, Kinney D, Bacerdo M, Bonk S, Hanna B, et al. Isolation of the
gene for the beta subunit of RNA polymerase from rifampicin-resistant Mycobacterium
tuberculosis and identification of new mutations. Am J Respir Cell Mol Biol. 1994;11:639–43.

36. Miotto P, Cabibbe AM, Borroni E, Degano M, Cirillo DM. Role of disputed mutations in
the rpoB gene in interpretation of automated liquid MGIT culture results for rifampin
susceptibility testing of Mycobacterium tuberculosis. J Clin Microbiol [Internet]. American
Society for Microbiology; 2018;56. Available from:
http://jcm.asm.org/cgi/pmidlookup?view=long&pmid=29540456

37. Heym B, Alzari PM, Honoré N, Cole ST. Missense mutations in the catalase-peroxidase
gene, katG, are associated with isoniazid resistance in Mycobacterium tuberculosis. Mol
Microbiol. Wiley; 1995;15:235–45.

38. Seifert M, Catanzaro D, Catanzaro A, Rodwell TC. Genetic mutations associated with
isoniazid resistance in Mycobacterium tuberculosis: a systematic review. PLoS One. Public
Library of Science (PLoS); 2015;10:e0119628.

39. Miotto P, Tessema B, Tagliani E, Chindelevitch L, Starks AM, Emerson C, et al. A
standardised method for interpreting the association between mutations and phenotypic drug
resistance in Mycobacterium tuberculosis. Eur Respir J [Internet]. 2017;50. Available from:
http://dx.doi.org/10.1183/13993003.01354-2017

40. Ektefaie Y, Dixit A, Freschi L, Farhat MR. Globally diverse Mycobacterium tuberculosis
resistance acquisition: a retrospective geographical and temporal analysis of whole genome
sequences. The Lancet Microbe. Elsevier BV; 2021;2:e96–104.

41. Ezewudo M, Borens A, Chiner-Oms Á, Miotto P, Chindelevitch L, Starks AM, et al.
Integrating standardized whole genome sequence analysis with a global Mycobacterium
tuberculosis antibiotic resistance knowledgebase. Sci Rep. Springer Science and Business
Media LLC; 2018;8:15382.

GenTB: A user-friendly genome-based predictor for

tuberculosis resistance powered by machine learning

TABLES AND FIGURES

Authors:

Matthias I Gröschel[1], Martin Owens[1], Luca Freschi[1], Roger Vargas Jr[1,2], Maximilian G Marin[1,2], Jody Phelan[3], Zamin Iqbal[4], Avika Dixit[1,5] and Maha R Farhat[1,6]

Affiliations

[1] Department of Biomedical Informatics, Harvard Medical School, Boston, MA, USA

[2] Department of Systems Biology, Harvard Medical School, Boston, MA, USA

[3] Faculty of Infectious and Tropical Diseases, London School of Hygiene & Tropical Medicine, London WC1E 7HT, UK

[4] European Bioinformatics Institute, Hinxton, Cambridge CB10 ISD, UK

[5] Division of Infectious Diseases, Boston Children's Hospital, Boston, MA, USA

[6] Division of Pulmonary and Critical Care Medicine, Massachusetts General Hospital, Boston, MA, USA

Figure 1. Schematic overview of the GenTB pipeline. Raw sequence data is quality checked and adapter trimmed before alignment to the H37Rv reference strain (accession AL123456). Variants are called with Pilon, and a variant matrix used by the prediction models are prepared using custom scripts available on Github. The analysis will fail if quality criteria are not met (blunt end arrows). Numbers represent the three moments in the pipeline where users can upload their data to predict resistance for their isolate.

Figure 2. GenTB online user interface. A) The user is presented with the three main features offered by GenTB, i.e., to run predictions from user input data, to upload, share, and cite their data with the GenTB project, and to geographically map resistance frequencies or phenotype data. **B)** Example of a resistance prediction output where boxes are colored in the function of the prediction model's output probability. **C)** Mutation plot that appears when clicked on one of the drugs heatmaps in **(B)**. Mutations will be shown when hovering the mouse over the genetic loci. INH = isoniazid, RIF = rifampicin, PZA = pyrazinamide, STR = streptomycin, EMB = ethambutol, ETH = ethionamide, KAN = kanamycin, CAP = capreomycin, AMK = amikacin, LEVO = levofloxacin, OFL = ofloxacin, PAS = Para-aminosalicylic acid.

Figure 3: Diagnostic performance of the four prediction tools across antituberculosis drugs. Paired violin plots displaying sensitivity and specificity to predict drug resistance for **A)** GenTB-Random Forest, **B)** GenTB-Wide and Deep Neural Network, **C)** TB-Profiler and **D)** Mykrobe. **E)** Violinplot of diagnostic performance to predict rifampicin resistance comparing isolates passing depth filters (in black) to isolates that failed the depth-filters (in grey) arranged by prediction tool. **F)** Violinplot of diagnostic performance to predict isoniazid resistance comparing isolates passing depth filters (in black) to isolates that failed the depth-filters (in grey) arranged by prediction tool.

AMK = amikacin, CAP = capreomycin, EMB = ethambutol, ETH = ethionamide, INH = isoniazid, KAN = kanamycin, OFL = ofloxacin, PZA = pyrazinamide, RIF = rifampicin, STR = streptomycin.

Figure 4: ROC performance curve of the GenTB-RF and GenTB-WDNN prediction models. A ROC plot of the GenTB-Random Forest (top) and GenTB-WDNN (bottom) predictive performance on the study dataset for first line (**A)** and **C)**) and second line drugs (**B)** and **D)**).

Table 1: Diagnostic accuracy of GenTB RandomForest and GenTB Wide and Deep Neural Network compared with two other leading prediction tools on a depth filtered dataset.

DrugName	Phenotype		GenTB - RF		GenTB - WDNN		Mykrobe		TB-Profiler	
			Isolates sequenced with high depth (n = 19.880)							
	R (n)	S (n)	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)
isoniazid	6,043	13,112	91% (91 to 92)	98% (97 to 98)	90% (89 to 91)	99% (99 to 99)	87% (86 to 88)	98% (98 to 98)	91% (90 to 92)	98% (97 to 98)
rifampicin	5,068	14,474	93% (93 to 94)	98% (98 to 98)	88% (88 to 89)	99% (99 to 99)	90% (89 to 91)	98% (98 to 99)	92% (91 to 93)	98% (98 to 99)
ethambutol	2,936	12,362	86% (85 to 87)	92% (92 to 93)	82% (80 to 83)	93% (93 to 94)	79% (77 to 80)	93% (93 to 94)	86% (85 to 88)	92% (92 to 93)
pyrazinamide	508	1,544	79% (76 to 83)	94% (93 to 95)	80% (79 to 82)	95% (94 to 95)	72% (71 to 74)	98% (97 to 98)	83% (80 to 86)	96% (96 to 97)
amikacin	618	3,458	67% (63 to 71)	99% (99 to 100)	66% (62 to 70)	99% (99 to 100)	63% (60 to 67)	99% (99 to 100)	55% (51 - 59)	99% (99 to 100)
capreomycin	648	3,733	63% (59 to 67)	97% (97 to 98)	57% (53 to 61)	98% (98 to 99)	60% (56 to 64)	98% (98 to 99)	56% (52 to 60)	96% (95 to 96)
ethionamide	502	1,094	67% (63 to 72)	78% (75 to 80)	-	-	-	-	70% (66 to 74)	73% (70 to 76)
kanamycin	576	3,707	68% (64 to 72)	99% (98 to 99)	66% (62 to 70)	100% (99 to 100)	66% (63 to 70)	99 (99 to 100)	68% (64 to 71)	98% (98 to 99)
streptomycin	2,126	4,968	82% (80 to 83)	89% (88 to 90)	87% (85 to 88)	87% (86 to 88)	68% (66 to 70)	95% (95 to 96)	71% (70 to 73)	95% (95 to 96%)
ofloxacin	743	4,038	68% (65 to 72)	99% (98 to 99)	62% (58 to 66)	96% (95 to 96)	62% (58 to 65)	99% (98 to 99)	67% (63 to 70)	98 (98 to 99)

Table 2: Area under the Receiver Operating Characteristic curve for GenTB-RF and GenTB-WDNN

Drug	GenTB-RF	GenTB-WDNN
	Area under the ROC curve (95% CI)	
isoniazid	0.94 (0.94 to 0.95)	0.94 (0.94 to 0.95)
rifampicin	0.96 (0.95 to 0.96)	0.94 (0.93 to 0.94)
ethambutol	0.89 (0.88 to 0.9)	0.87 (0.87 to 0.87)
pyrazinamide	0.90 (0.88 to 91)	0.88 (0.87 to 0.88)
amikacin	0.83 (0.81 to 0.85)	0.83 (0.81 to 0.84)
capreomycin	0.80 (0.78 to 0.82)	0.78 (0.76 to 0.80)
ethionamide	0.73 (0.7 to 0.75)	-
kanamycin	0.83 (0.81 to 0.85)	0.83 (0.81 to 0.85)
streptomycin	0.85 (0.84 to 0.86)	0.87 (0.86 to 0.88)
ofloxacin	0.83 (0.82 to 0.85)	0.79 (0.77 to 0.81)

RF = Random Forest, WDNN = Wide and Deep Neural Network

Table 3: Output comparison across tools

Criteria	GenTB	TB-Profiler	Mykrobe
1) Output			
Type	Heatmap and barplot	Overview tables	Overview table
Download	all intermediate and output files (JSON)	yes (CSV)	Yes (JSON)
2) Genotypic predictions	Probability	Binary	Binary
3) Error rate	Yes	N.A.	N.A.
4) Resistance variants	Variant by drug	Variant by drug incl. fraction of mutant / wild-type allele	Variant by drug incl. depth of mutant and wild-type alleles
5) Unknown variants	Yes, in all genes	Yes, in candidate resistance genes	No
6) *M. tuberculosis* Lineage			
Lineage	Yes	Yes	Yes
Typing scheme	Yes	No	No
7) Quality metrics	Trimming and Kraken report downloadable	No. of reads, Percentage of reads mapped	No