mTORC2 contributes to murine systemic autoimmunity

Xian Zhou[1], Haiyu Qi[1,2], Meilu Li[1,3], Yanfeng Li[1], Xingxing Zhu[1], Shreyasee Amin[1], Mariam Alexander[4], Catherine Diadhiou[5,6], Anne Davidson[5,6], Hu Zeng[1,7]

[1]Division of Rheumatology, Department of Medicine, Mayo Clinic Rochester, MN 55905, USA

[2]Department of Rheumatology, Beijing Friendship Hospital, Capital Medical University, Beijing, 100050, P. R. China

[3]Department of Dermatology, the Second Hospital of Harbin Medical University, Harbin Medical University, Harbin, 150001, P. R. China

[4]Division of Laboratory Medicine and Pathology, Mayo Clinic Rochester, MN 55905, USA

[5]Institute of Molecular Medicine, Feinstein Institutes for Medical Research, Manhasset, NY 11030, USA

[6]Donald and Barbara Zucker School of Medicine at Hofstra/Northwell, Manhasset, NY 11030, USA

[7]Department of Immunology, Mayo Clinic Rochester, MN 55905, USA

Address correspondence and reprint request to Dr. Hu Zeng, Division of Rheumatology, Department of Medicine, Mayo Clinic Rochester, Guggenheim Building 3-42, 200 First ST SW, Rochester, MN, 55902, USA. E-mail address: zeng.hu1@mayo.edu.

Abbreviations used in this article: Tfh, follicular helper T cells; Treg, regulatory T cells; SLE, systemic lupus erythematosus; mTOR, mechanistic target of rapamycin; IFN, interferon; GC, germinal center; EF, extrafollicular; Tfr, follicular regulatory T cells; ICOS, inducible T-cell costimulator.

ABSTRACT

The development of many systemic autoimmune diseases, including systemic lupus erythematosus, is associated with overactivation of the type I interferon (IFN) pathway, lymphopenia, and increased follicular helper T (Tfh) cell differentiation. However, the cellular and molecular mechanisms underlying these immunological perturbations remain incompletely understood. Here we show that the mechanistic target of rapamycin complex 2 (mTORC2) promotes Tfh differentiation and disrupts Treg homeostasis. Inactivation of mTORC2 in total T cells, but not in Tregs, greatly ameliorated the immunopathology in a systemic autoimmunity mouse model. This was associated with reduced Tfh differentiation, B cell activation, and reduced T cell glucose metabolism. Finally, we show that type I IFN can synergize with TCR ligation to activate mTORC2 in T cells, which partially contributes to T cell lymphopenia. These data indicate that mTORC2 may act as downstream of type I IFN, TCR, and costimulatory receptor ICOS, to promote glucose metabolism, Tfh differentiation, and T cell lymphopenia, but not to suppress Treg function in systemic autoimmunity. Our results suggest that mTORC2 might be a rational target for systemic autoimmunity treatment.

INTRODUCTION

The cellular and molecular mechanisms leading to systemic autoimmunity remain poorly understood. A prototype of systemic autoimmunity is systemic lupus erythematosus (SLE). Two major immunological alterations have been identified in lupus, overactivation of the type I IFN signaling pathway and a loss of balance between effector T cells, particularly follicular helper T (Tfh) cells, and regulatory CD4[+] T cells (Tregs)[1-3]. Therapeutical options like blocking type I IFN, or its receptor have shown clinical efficacy[4]. Yet, the specific cellular and molecular mechanisms by which type I IFN promotes SLE pathology remain incompletely understood. Recent studies using murine lupus models have demonstrated that type I IFN signaling in CD4[+] T cells are required for development of lupus pathology[5,6]. In addition, it is well established that type I IFN can block T cell egress from lymph nodes and induce lymphopenia partly by upregulating the expression of CD69[7,8], a marker that is also associated with lupus pathology[9,10]. Clinical studies[11] and large-scale single cell RNA sequencing analysis[12] confirmed the inverse association between type I IFN activity and the abundance of circulating lymphocytes or naïve CD4[+] T cells in SLE patients. However, the type I IFN signaling pathway in CD4[+] T cells remains incompletely understood and it is unclear how type I IFN overactivation may be linked to T cell lymphopenia in lupus.

Increased differentiation of Tfh cells has been associated with SLE development and severity in mouse models and human patients[13,14]. Germinal center (GC) Tfh cells, distinguished by expression of the transcription factor BCL6, chemokine receptor CXCR5, and immune checkpoint receptor PD-1, are a specialized effector T cell (Teff) lineage that stimulates B cells in GC to undergo affinity maturation and clonal expansion. The costimulatory receptor ICOS

delivers critical signals for Tfh differentiation[15]. In addition, a special subset of Tfh cells with

low expression of CXCR5 and P-selectin ligand, PSGL-1, are localized in extrafollicular (EF) to

promote generation of plasmablasts, which produce antibodies with modest affinity[16-18]. Both

GC and EF responses may contribute to SLE development[19]. In contrast, impaired Treg function

is observed in some SLE patients[20]. However, what drives the imbalance between Tfh and Tregs

in lupus is not well understood.

Mechanistic target of rapamycin (mTOR) is a central metabolic pathway, which consists of two

complexes, mTORC1 and mTORC2, with scaffolding molecules RAPTOR (encoded by *RPTOR*)

and RICTOR (encoded by *RICTOR*) as their defining components, respectively[21]. mTORC1 is

critical for naïve T cell quiescence exit and proliferation, and all effector T cells differentiation[22].

In contrast, mTORC2 is essential for Tfh differentiation, but dispensable for T cell activation and

modestly contributes to other effector T cell lineages[22]. Ligation of CD3 and ICOS activates

mTORC2, which in turn promotes glucose metabolism in Tfh cells[23]. In terms of Tregs,

mTORC1 maintains natural or thymic derived Treg (nTreg, or tTreg) functional competency[24-27].

Overactivation mTORC2 impairs Treg suppressive function and stability, especially their ability

to suppress Tfh cells, which leads to systemic autoimmune disease with lupus characteristics[28,29].

Inhibition of mTORC2 in Tregs can partly restore immune tolerance in the absence of

mTORC1[24], or FOXP3[30]. Therefore, mTORC2 signaling controls the balance between Tfh and

Treg by promoting the former and suppressing the latter. These fundamental discoveries suggest

that targeting mTORC2 in T cells might benefit lupus disease.

Several studies have examined the contribution of mTORC1 to SLE development[31]. How

mTORC2 contributes to systemic autoimmunity has been much less understood. A significant

increase of phosphorylation of AKT at serine 473, an established direct target of mTORC2, was

observed in T cells from lupus patients and it was associated with disease severity[32,33]. These

studies suggest that mTORC2 overactivation might be a potential contributing factor for lupus.

The underlying mechanisms, however, remain obscure. Here, we tested the hypothesis that

genetic targeting mTORC2 in T cells may improve systemic autoimmunity in C57BL/6-Fas^{Lpr}

(Lpr) mice. We show that Lpr T cells have an elevated baseline activity of mTORC2. Loss of

Rictor specifically in T cells significantly improves immunopathology in Lpr mice, including T

cell lymphopenia and generation of autoantibodies, which is associated with normalization of the

balance between Tfh cells and Tregs. Surprisingly, mTORC2 contributes to systemic

autoimmunity primarily through promoting Tfh cells, but not suppressing Tregs, because specific

deletion of *Rictor* in Tregs largely fails to ameliorate the immune activation in Lpr mice. Lastly,

we demonstrate that type I IFN can synergize with TCR signaling to sustain mTORC2 activation

and promote CD69 expression, which suppress T cell egress and lead to lymphopenia, partly in

an mTORC2 dependent manner. Our data supports the notion that mTORC2 dependent Tfh

expansion, but not Treg impairment, may contribute to the lupus-associated systemic

immunopathology.

MATERIAL and METHODS
Mice

Cd4[Cre]*Rictor*[fl/fl] mice have been described before[23]. C57BL/6-*Fas*[Lpr] and Foxp3[Cre] mice were

purchased from The Jackson Laboratory. Mice were bred and maintained under specific

pathogen-free conditions in the Department of Comparative Medicine of Mayo Clinic with

IACUC approval.

ELISA

For autoantibodies detection in sera, 96-well plates (2596; Costar) were coated with ssDNA and

chromatin (Sigma-Aldrich) in PBS overnight at 4°C used at a concentration of 1.6 μg/ml and 5

μg/ml, respectively. Plates were washed twice (0.05% Tween 20 in PBS), blocked with 5%

blocking protein (Bio-Rad) for 1 h, and washed twice, and serially diluted sera samples were

added for 1.5 h at 37 °C. Plates were washed four times and horseradish peroxidase (HRP)-

conjugated detection Abs for IgG (Bethyl Laboratories) were added for 1 h at RT, washed four

times, and tetramethylbenzidine (TMB) substrate was added. Reaction was stopped using 2N

H_2SO_4 and read at 450 nm. The titers of mouse IgM, IgG, IgG2b and IgG3 (Fig. 1C) in sera were

measured with the ELISA kits from Invitrogen.

Mouse immunoglobulin isotyping panel detection

The concentration of mouse immunoglobulin isotypes IgG1, IgG2a, IgG2b and IgA (Fig. 4G) in

sera were measured with the LEGENDplex mouse immunoglobulin isotyping panel, according to

manufacturer's instructions (Biolegend; cat# 740493).

Immunofluorescence staining

Kidneys were obtained from 6-month-old B6, Lpr and Lpr. *Rictor*[fl/fl] mice, and fixed in 4%

paraformaldehyde (PFA) overnight, osmotically dehydrated in 20 % (w/v) sucrose, embedded in

OCT, and cryosectioned to 6 μM. Sections were fixed in −20°C acetone for 2 min and blocked in

PBS containing 8% goat serum, 1% bovine serum albumin (BSA) and 0.1% Tween 20 for 45

min. Deposition of immunoglobulin (IgG) and complement component 3 (C3) in the kidney

tissue were stained with PE labeled goat anti-mouse IgG (Invitrogen) and FITC labeled rat anti-

mouse C3 (Cedarlane) antibodies. Similarly, spleens were obtained from 6 months old mice, and

processed as above. Frozen sections were fixed and blocked as described above, and incubated

with primary antibodies Biotin Peanut Agglutinin (PNA) (Vectorlabs), purified rat anti-mouse

CD3 (Biolegend) and PE labeled anti-mouse IgD (BD), or stained with Biotin anti-mouse CD138

(Biolegend) and PE labeled anti-mouse IgD at 4°C for 4 hours. Slides were then washed before

streptavidin-Alexa Fluor 488 conjugate (ThermoFisher) and Alexa Fluor 647 labeled goat anti-

rat secondary antibody (ThermoFisher) were added for 1 h at room temperature. Sections were

covered with coverslip and visualized under an Olympus BX51 fluorescence microscope.

Flow cytometry

Single-cell suspension was prepared by passing spleen and peripheral lymph nodes through a 70-

μm nylon mesh. For analysis of surface markers, cells were stained in PBS containing 1% (w/v)

BSA on ice for 30 min, with APC-labeled anti-ICOS (clone: 7E.17G9; BD), PE-Cy7-labeled anti

PD-1 (clone: RMPI-30; BioLegend), Super Bright 600–labeled anti-CD4 (Clone: SK-3;

eBioscience), BV510-labeld anti-CD8a (clone:53-6.7; BioLegend); APC-Cy7-labeled TCRβ

(clone: H57-597; BioLegend), APC anti-CD162 (PSGL1; clone: 2PH1; BD), BV605-labeled

anti-CD25 (clone: PC61; BioLegend), PE-Cy7–labeled anti-CD19 (clone: 6D5; BioLegend),

BV605-labeled B220 (clone: RA3-6B2; BioLegend), PE-labeled anti-Fas (clone: Jo2; BD),

BV785-labeled anti-CD138 (clone: 281-2; BioLegend), APC-labeled anti-IgD (clone: IA6-2;

BioLegend), PerCP-Cyanine5.5-labeled anti-CD38 (clone: 90; BioLegend), FITC-labeled anti-

IgG1 (clone: RMG1-1; BioLegend), APC-labeled anti-IL7Rα (clone: SB/199; BioLegend),

FITC-labeled anti-CD69 (clone: H1.2F3; Tonbo Bio), PE-labeled anti-CD153 (clone: RM153;

eBioscience), PE-labeled anti-CD73 (clone: TY/11.8; BioLegend) and APC-labeled anti-FR4

(clone: 12A5; BD). Cell viability was assessed using the Fixable Dye Ghost 540 (Tonbo

Bioscience), or 7AAD. CXCR5 and PNA were stained with biotinylated anti-CXCR5 (clone:

2G8, BD Biosciences) or biotinylated peanut agglutinin (FL10-71; Vectorlabs), followed by

staining with streptavidin-conjugated PE (BD Biosciences). Splenocytes and peripheral lymph

node cells were cultured for 4 h at 37°C in complete medium (1640 RPMI + 10% FBS)

containing ionomycin (1 µg/ml), PMA (50 ng/ml) and Monensin solution (1,000×; BioLegend)

for cytokines detection. Intracellular cytokine staining of IFN-γ (clone: XMG1.2; eBioscience),

IL-2 (clone: JES6-5H4; BioLegend) and IL-17A (clone: TC11-18H10.1; BioLegend) were

performed using the BD Cytofix/Cytoperm and Perm/Wash buffers or, for transcriptional factors

FoxP3 (clone: FJK-16s; eBioscience), Bcl6 (clone: K112-91, BD) and Ki67 (clone: SolA15;

eBioscience) staining using the eBioscience Transcription Factor Staining Buffer Set. Phosflow

staining for AF647 conjugated phosphor-AKT (S473) (D9E; Cell signaling) was performed

using Phosflow Fix/Perm kit (BD Biosciences). Flow cytometry was performed on the Attune

NxT (ThermoFisher) cytometer, and analysis was performed using FlowJo software (Tree Star).

Immunoblotting

For immunoblotting, murine CD4$^+$ T cells were isolated from mouse spleen and peripheral

lymph node single cell suspension, and human CD4$^+$ T cells were enriched from healthy donor

PBMC with STEMCELL mouse or human CD4$^+$ T cell isolation kits, respectively. Cells were

lysed in lysis buffer with protease and phosphatase inhibitors (Sigma-Aldrich). Protein

concentration in samples was quantified by BCA assay (Thermo Fisher Scientific) before loading

the samples for electrophoresis and membrane transfer. The transferred membrane was blocked

with TBST (0.1% Tween 20) containing 5% BSA for 1 h at room temperature. The membrane

was incubated with primary antibodies overnight including anti-phospho-AKT (Ser473) (clone:

D9E; Cell Signaling), AKT (pan) (Clone: 40D4; Cell Signaling), phospho-STAT1 (Tyr701)

(clone: D4A7; Cell Signaling), phospho-STAT2 (Tyr690) (Cell Signaling), phospho-S6

(Ser235/236) (clone: D57.2.2E; Cell Signaling) and anti-β-actin (clone: 13E5; Sigma-Aldrich).

Then, the membrane was washed and incubated with the corresponding secondary antibody for

subsequent enhanced chemiluminescence (ECL; Thermo Fisher) exposure. The band intensity of

all the immunoblot was analyzed by ImageJ software.

Inflammatory cytokines detection

Murine sera were collected from 6-month-old B6, Lpr and Lpr.*Rictor* T-KO mice. The levels of

inflammatory cytokines in serum were measured with the LEGENDplex Multi-Analyte Flow

Assay Kit, according to manufacturer's instructions (Biolegend, Mouse Inflammation Panel, cat#

740446).

ICOS stimulation

Cell isolation and flow cytometry of lymphocytes were as described. One million CD4$^+$ T cells

were activated with plate coated anti-CD3 and anti-CD28 (Bio X Cell). After 2 days activation,

T cells were removed from stimulation and rested overnight. Live cells were purified using

lymphocyte isolation buffer (MP Biomedicals), followed by stimulation with plate bound anti-

CD3 (2 μg/ml) and anti-ICOS (5 μg/ml; C398.4A; Biolegend) for 24 hours.

Metabolic assay

The bioenergetic activities of the OCR and ECAR were measured using a Seahorse XFe96

Extracellular Flux Analyzed following established protocols (Agilent). Briefly, equal number of

live CD4$^+$ T cells were seeded at 200,000 cells/well on Cell-Tak (Corning) coated XFe96 plate

with fresh XF media (Seahorse XF RPMI medium containing 10 mM glucose, 2 mM L-

glutamine, and 1 mM sodium pyruvate, PH 7.4; all reagents from Agilent). Basal OCR and

ECAR were measured in the presence of Oligomycin (1.5 μM; Sigma-Aldrich), FCCP (1.5 μM;

Sigma-Aldrich), and Rotenone (1 μM; Sigma-Aldrich)/ Antimycin A (1 μM; Sigma-Aldrich) in

Mito Stress assay. For ECAR measurement in ICOS secondary stimulated CD4$^+$ T cells,

glycolytic rate assay was performed according to the manufacturer's manual (Agilent) in

presence of Rotenone/Antimycin A and 2-DG (50 mM, Sigma-Aldrich)

Poly(I:C) treatment and cell counting in peripheral blood

Cd4^{Cre}*Rictor*$^{fl/fl}$ and wild-type mice were given intraperitoneal injection of 200 μg poly(I:C)

(Sigma-Aldrich) in 200 μL PBS, and blood lymphocyte counts were assessed 16 and 40 hours

later. Briefly, to determine absolute blood cell counts, 50 μL peripheral blood with 5 μL EDTA

was collected, lysed with 1 mL ACK lysing buffer (ThermoFisher), and washed with PBS.

Samples were stained with anti-mouse CD19, CD4, CD8 and CD69 antibodies, washed with

PBS once, suspended with 500 μL PBS and loaded to Attune NxT cytometer with the accurate

acquisition of 250 μL samples. Cell counts were calculated per 5 μL blood.

Statistical Analysis

Statistical analysis was performed using GraphPad Prism (version 8). P values were calculated

with Student's *t* test, or one-way ANOVA. *P* < 0.05 was considered significant.

RESULTS
Loss of mTORC2 in T cells improves immunopathology in murine systemic autoimmunity

Lpr mice recapitulate many of the systemic immunoproliferative phenotypes in human lupus

disease without developing overt nephritis phenotypes[34]. We tested whether mTORC2 activity is

altered in T cells from Lpr mice. Immunoblot analysis showed that $CD4^+$ T cells from Lpr mice

had elevated phosphorylation of AKT at Serine 473 (p-AKT_{473}), a direct target of mTORC2,

indicating overactivation of mTORC2 in Lpr T cells at baseline (Fig. 1A). To test the hypothesis

that targeting mTORC2 may improve systemic autoimmunity, we crossed $Cd4^{Cre}Rictor^{fl/fl}$ mice,

in which $Rictor$ is specifically deleted and hence abrogates mTORC2 activity in T cells[23,24], with

Lpr mice to generate C57BL/6.$Fas^{Lpr}Cd4^{Cre}Rictor^{fl/fl}$ (Lpr.$Rictor$ T-KO) mice. RICTOR

deficiency in T cells alone does not significantly impact immune cell homeostasis and overall

mouse health[24,35,36]. Therefore, we focused on the comparison among B6, Lpr and Lpr.$Rictor$ T-

KO mice. Fas^{Lpr} mutation drives dramatic lymphoid hyperplasia, including lymphadenopathy.

We observed that deletion of $Rictor$ in T cells significantly rescued lymphadenopathy

phenotypes (Fig. 1B). Consistent with these findings, Lpr mice had increased total IgM and IgG,

and isotypes IgG2b and IgG3 immunoglobulin levels (Fig. 1C) and increased anti-ssDNA and

anti-chromatin autoantibodies (Fig. 1D), most of which were significantly reduced in Lpr.$Rictor$

T-KO mice. Furthermore, we examined IgG and complement deposition in kidneys, which

reflects renal involvement in autoimmunity. Lpr mice on B6 background have very mild kidney

pathology compared to those on MRL/MpJ background[37]. Consistent with this observation, we

observed a mild increase of IgG and complement C3 deposition in kidneys from Lpr mice.

Deletion of $Rictor$ reduced IgG and C3 deposition in Lpr.$Rictor$ T-KO kidney, although the latter

did not reach statistical significance (Supplemental Fig. 1A). Thus, inactivation of mTORC2 in T

cells substantially attenuates lymphoid hyperplasia, antibody secretion, and renal immune

complex deposition in a mouse model of systemic autoimmunity.

Consistent with the attenuated lymphoid hyperplasia in Lpr.*Rictor* T-KO mice, the accumulation

of the aberrant B220$^+$TCRβ$^+$ cells in Lpr mice was nearly completely reversed in Lpr.*Rictor* T-

KO mice (Fig. 1E). Although Lpr.*Rictor* T-KO mice had a significant increase of B220$^+$TCRβ$^-$

B cell percentage, the absolute number of this population modestly, but significantly, reduced in

Lpr.*Rictor* T-KO mice compared to Lpr mice (Fig. 1E). B220$^-$TCRβ$^+$ T cell percentages were

highly reduced in both Lpr and Lpr.*Rictor* T-KO mice, but Lpr mice had an increase of B220$^-$

TCRβ$^+$ T cell number likely due to the lymphoid hyperplasia, which was also reversed in

Lpr.*Rictor* T-KO mice (Fig. 1E). Moreover, the accumulation of TCRβ$^+$CD4$^-$CD8$^-$ DN T cells

was substantially restored by *Rictor* deletion (Fig. 1F**)**. Finally, there were elevated, albeit to a

widely variable degree, levels of multiple inflammatory cytokines in Lpr mice, such as IFNβ, IL-

1β, TNFα and IL-17A, many of which were rectified in Lpr.*Rictor* T-KO mice (Fig. 1G).

Therefore, the inactivation of mTORC2 in T cells significantly reverses the abnormal

accumulation of T and B cells, production of immunoglobulin and inflammatory cytokine, and

restores immune homeostasis in peripheral lymphoid organs of Lpr mice.

mTORC2 in T cells promotes B cell hyper-activation

Reflecting the enhanced antibody levels, immunofluorescence staining revealed a dramatic

increase of peanut agglutinin (PNA) staining in Lpr mice, suggesting increased GC reaction. The

pattern of the PNA staining was variable. Some are dense aggregations like defined GC

formation, but others are more diffused without defined boundaries (Fig. 2A). T cell distribution

was also disorganized in Lpr lymphoid tissues. Both the increased PNA staining and

disorganized T cell distribution were greatly restored by RICTOR deficiency (Fig. 2A). Flow

cytometry analysis confirmed these observations. B cells from Lpr mice had highly increased

expression of GC B cell marker GL-7, and significant loss of IgD expression, indicating

increased B cell activation and class switch (Fig. 2B). Consistent with this notion, Lpr B cells

had greatly increased IgG1 expression (Supplemental Fig. 2A). A large proportion of Lpr B cells

lost expression of CD38, a NADase with strong immunosuppressive capacity[38], and the loss of

which is another characteristic of germinal center B cells (Supplemental Fig. 2A). Because

plasmablasts contribute to extrafollicular response, we stained CD138, a marker of plasmablasts.

The results showed a markedly increase of CD138 expression in extrafollicular region of Lpr

spleens, which was reduced in Lpr.*Rictor* T-KO mice (Fig. 2C). The RICTOR-dependent

increase of extrafollicular response in Lpr mice was confirmed with flow cytometry analysis. Lpr

mice accumulated large number of cells with modest expression of CD138, and $B220^{lo}CD138^{hi}$

plasmablast/plasma cells, while deletion of *Rictor* in T cells dramatically suppressed the

generation of these populations (Fig. 2D). Therefore, mTORC2 activity in T cells is a major

contributor to B cell abnormality in Lpr mice. Taken together, our data indicate that inhibition of

mTORC2 in T cells attenuates GC and plasmablast formation in Lpr mice.

Increased Tfh differentiation in Lpr mice is dependent on mTORC2

Tfh is key for GC formation. Indeed, we observed many T cells in GC of Lpr mice from

immunofluorescence staining, presumably Tfh cells (Fig. 2A). This was confirmed by flow

cytometry data, which showed a strong increase of $BCL6^+CXCR5^{hi}$ GC Tfh and $BCL6^-CXCR5^+$

pre-Tfh cells in Lpr mice, that was significantly dependent on RICTOR expression (Fig. 3A and

Supplemental Fig. 3A for gating strategy). A RICTOR-dependent CXCR5 expression was also

found on TCRβ⁺CD4⁻CD8⁻ DN T cells (data not shown). Similar observations were found when

using other conventional Tfh markers, PD-1 and CXCR5 (Supplemental Fig. 3B). RICTOR

deficiency also reduced PSGL-1⁻CXCR5⁻ EF Tfh cell frequency (Fig. 3B), consistent with the

reduction of plasmablast/plasma cells. These results indicated a critical requirement of mTORC2

signaling for autoreactive GC and EF Tfh differentiation in Lpr lupus-prone model. Similar

rescue effects were observed at different ages (at 3 and 9 months, data not shown). Like SLE

patients[14], Lpr mice had highly increased ICOS expression, both on CXCR5⁺ and CXCR5⁻ T

cells. Interestingly, deletion of *Rictor* failed to restore ICOS expression (measured by both

positive percentages and mean fluorescence intensity), indicating that the reduced Tfh

differentiation in Lpr.*Rictor* T-KO mice is not a consequence of reduced ICOS expression (Fig.

3C).

Treg cells are key for maintaining immune tolerance and preventing autoimmunity.

Counterintuitively, in an inflammatory condition, Treg percentage sometimes paradoxically

increases. We found that Lpr mice had variable Treg perturbations dependent on anatomic

location: while Treg percentages increased the most in spleens and modestly in peripheral lymph

nodes (pLN), they trended lower in the gut mucosal site relative to WT B6 mice (Fig. 3D), in a

manner consistent with the graded mTORC2 activity, gradually increasing from spleen to pLN,

and to gut mucosal site[23]. In all cases, deletion of *Rictor* significantly restored the Treg

homeostasis in Lpr mice (Fig. 3D). The magnitude of changes of Treg frequencies in Lpr mice

were much smaller than that of Tfh cells, thus the ratio between Tfh and Treg was drastically

increased in Lpr mice, suggesting an imbalance between Tfh and Tregs in Lpr mice. Such

imbalance was significantly restored by mTORC2 deficiency (Fig. 3E). FOXP3⁺CXCR5⁺

follicular regulatory T (Tfr) cells are specialized to suppress Tfh cells and GC formation[39].

RICTOR deficiency alone does not impact Tfr differentiation[23]. Lpr mice had a highly increased

ratio between Tfh and Tfr cells, which was partly restored in Lpr.*Rictor* T-KO mice (Fig. 3E).

However, Lpr mice also had significantly increased frequency of FOXP3$^+$CXCR5$^+$ Tfr cells

compared to B6 control mice, which was also reversed by *Rictor* deletion (Supplemental Fig.

3C). Therefore, the data indicate that RICTOR deficiency impacts Tfh/Tfr ratio primarily by

reducing Tfh, not increasing Tfr.

Furthermore, we found that *Rictor* deficiency did not restore the increased IFNγ expression (Fig.

3F) or alter IL-2 and IL-17 expression in CD4$^+$ T cells from Lpr mice (Fig. 3F, 3G), or change

the expression of lineage transcription factor Tbet and Rorγt (Supplemental Fig. 3D), consistent

with our hypothesis that mTORC2 inactivation preferentially affects Tfh differentiation, but not

other effector T cell lineages. These data also suggested that Tregs from Lpr.*Rictor* T-KO mice

were not able to control the excessive cytokine production in Lpr CD4$^+$ T cells, despite

normalization of their homeostasis. Finally, the restoration of T cell phenotypes in Lpr.*Rictor* T-

KO mice was likely independent of cell proliferation, because CD4$^+$ T cells from Lpr.*Rictor* T-

KO mice had similarly increased Ki-67$^+$ percentage at steady state as those from Lpr mice (Fig.

3H). Therefore, our data showed that mTORC2 contributes to the increased GC and EF Tfh cells

in Lpr mice, independent of ICOS receptor expression and T cell proliferation.

mTORC2 inactivation in Tregs fails to restore immune homeostasis in Lpr mice

The two best described functions of mTORC2 in T cells are promoting Tfh differentiation[23,40]

and suppression of Treg function[24,28]. Thus, it is possible the above-described phenotypic

improvements in Lpr.*Rictor* T-KO mice (deletion in all T cells) can be partly attributed to

enhanced Treg function. To dissect whether mTORC2 contributes to systemic autoimmunity

mainly through inhibiting Tregs, promoting Tfh, or both, we generated

C57BL/6.*Fas[Lpr]Foxp3*[Cre]*Rictor*[fl/fl] (Lpr.*Rictor* Treg-KO) mice, in which *Rictor* was specifically

deleted in Treg cells (Supplemental Fig. 4A). Although *Rictor* deficiency in Tregs can boost

Treg suppressive function in the absence of mTORC1[24], *Pten*[28] or *Foxp3*[30], it also reduces

peripheral Treg frequency compared to WT controls[24]. Lpr.*Rictor* Treg-KO mice also had

reduced Treg percentage compared to Lpr mice, although their Treg percentage was similar to

Lpr.*Rictor* T-KO mice (Fig. 4A). We compared the immunological phenotypes between Lpr,

Lpr.*Rictor* T-KO, and Lpr.Rictor Treg-KO mice. To our surprise, Lpr.*Rictor* Treg-KO mice had

equivalent lymphoplasia found in Lpr mice, as illustrated by the lymphadenopathy (Fig. 4B). The

expansion of $B220^+TCR\beta^+$ cells was not affected by Treg specific deletion of *Rictor* (Fig. 4C).

Similarly, the elevated Tfh, pre-Tfh and GC B cells in Lpr.*Rictor* Treg-KO mice were all at a

comparable level as those found in Lpr mice, in sharp contrast to Lpr.*Rictor* T-KO mice (Fig.

4D, E). Because of the elevated Tfh frequency, Lpr.*Rictor* Treg-KO mice retained a high Tfh vs

Treg ratio similar to that of Lpr mice (Fig. 4F). Quantification of immunoglobulin concentrations

revealed that Lpr.*Rictor* Treg-KO mice had highly elevated antibody levels, some of which

(including IgG1 and IgG2a) even surpassed those in Lpr mice, whereas Lpr.*Rictor* T-KO mice

had antibody levels close to those in B6 control mice (Fig. 4G). These data support the idea that

mTORC2-dependent overactivation of Tfh, but not mTORC2-mediated impairment of Treg

cells, contributes to the systemic lymphoproliferation in Lpr mice.

Inactivation of mTORC2 restores T cell glucose metabolism

Recent studies indicate that lupus development is associated with dysregulation of T cell

immunometabolism[41]. mTORC2 is known as one of the key regulators for cell metabolism. We

compared the basal respiration and glycolysis of T cells among B6, Lpr and Lpr.*Rictor* T-KO

mice at different ages under TCR signaling stimulation. Consistent with recent publications[42,43],

T cells from 2 months old Lpr mice had modestly increased basal respiration and glycolysis (Fig.

5A, 5B). Strikingly, T cells from 3 and 9 months old Lpr mice had dramatically decreased basal

respiration and glycolysis, relative to B6 mice (Fig. 5A, 5B). The opposite metabolic activities

between 2 and 3 months old Lpr mice T cells were associated with drastic changes of naive and

effector T cell composition at different ages of Lpr mice during lupus development

(Supplemental Fig. 5A). Lpr.*Rictor* T-KO mice had much lower levels of basal respiration and

glycolysis compared to B6 mice at all ages (Fig. 5A, 5B). The reduced metabolic activity in

primary culture of T cells from older Lpr and Lpr.*Rictor* T-KO mice (compared to 2 months old

mice) was consistent with reduced T cell proliferation during primary culture (Supplemental Fig.

5B). Because anergic T cells are known to have stunted metabolic activity and reduced

glycolysis[44,45], we tested whether the reduced metabolism in 6 months old Lpr T cells was

associated with increased anergic T cells. Indeed, we found drastically increased anergy

(FR4$^+$CD73$^+$)[48] markers on Lpr CD4$^+$ T cells at steady state, which was mildly reduced in

Lpr.*Rictor* T-KO mice (Supplemental Fig. 5C). Costimulatory signaling from ICOS is important

to limit T cell anergy[49]. Furthermore, we have previously shown that mTORC2 functioned as

downstream of ICOS receptor to promote Tfh differentiation, which could be modeled by

restimulating previously activated T cells with anti-CD3/anti-ICOS[23]. Interestingly, Lpr T cells

had enhanced proliferation during secondary activation with anti-CD3/anti-ICOS regardless of

the ages (Supplemental Fig. 5D, and data not shown), suggesting that re-stimulation from TCR

and ICOS signaling not only reversed the proliferation defect, and possibly broke the anergic

status, in Lpr T cells, but also promote more accumulation of Lpr T cells relative to WT control.

All these defects were not markedly affected by RICTOR deficiency. mTORC2 promotes

TCR/ICOS mediated glucose metabolism during Tfh differentiation[23]. Therefore, we examined

the T cell glycolysis after anti-CD3/anti-ICOS secondary stimulation. Lpr CD4$^+$ T cells exhibited

a significant increase of compensatory glycolytic rate relative to B6 controls, regardless of the

ages. *Rictor* deletion substantially reduced both basal and compensatory glycolytic rate in Lpr

CD4$^+$ T cells (Fig. 5C). Therefore, mTORC2 contributes to the TCR/ICOS mediated enhanced

glycolysis in Lpr T cells without affecting T cell proliferation.

mTORC2 functions downstream of type I IFN to promote CD69 expression and suppress T

cell egress

Previous studies have identified that type I IFN contributes to the lupus phenotypes in Lpr

mice[50,51]. We also observed increased type I IFN in Lpr mouse serum (Fig. 1G). Interestingly, we

found that addition of IFNα significantly enhanced anti-CD3/anti-CD28 induced mTORC2

target p-AKT$_{473}$, but not mTORC1 target p-S6, at 6 h stimulation, which was completely

dependent on RICTOR (Fig. 6A). IFNα alone could also induced weak p-AKT$_{473}$, but the

combination of anti-CD3/anti-CD28 and IFNα appeared to be the most potent activators of

RICTOR dependent p-AKT$_{473}$ (Fig. 6A, 6B). RICTOR deficiency did not markedly affect

canonical IFNα−STAT1 signaling or mTORC1 activation (Fig. 6A, Supplementary Fig. 6A).

Similar results were observed using IFNβ (Supplementary Fig. 6B). Thus, type I IFN, in synergy

with anti-CD3/anti-CD28, promoted p-AKT$_{473}$ in a RICTOR dependent manner. We also

observed similar synergy between TCR and type I IFN on mTORC2 activation in human CD4$^+$ T

cells (Fig. 6C). Type I IFN is known to suppress lymphocyte egress from lymph nodes to blood,

partly through induction of CD69 expression[7,8]. Poly(I:C) is a potent inducer of type I IFN. It

induces CD4$^+$ T cell lymphopenia in an IFNAR dependent manner[8]. Indeed, administering

poly(I:C) induced potent CD69 expression on blood CD4$^+$ T cell (Fig. 6D), and induced CD4$^+$ T

cell lymphopenia (Fig. 6E). Both phenotypes were significantly reversed in CD4$^+$ T cells from

$Cd4^{Cre}Rictor^{fl/fl}$ mice. Such phenotype was not observed in CD19$^+$ B cells, in which RICTOR

was intact (Supplemental Fig. 6C), indicating that type I IFN promotes CD4$^+$ T cell lymphopenia

partly through T cell intrinsic mTORC2. Consistent with these observations, Lpr CD4$^+$ T cells

had increased CD69 level, which was restored in Lpr.$Rictor$ T-KO mice in both lymph nodes and

blood (Fig. 6F, Supplemental Fig. 6D). Importantly, the severe CD4$^+$ T cell lymphopenia

phenotype in Lpr mice was also modestly, but significantly, reversed by RICTOR deficiency

(Fig. 6G, Supplemental Fig. 6E). The rescue effect on CD69 expression was consistently more

pronounced than that on CD4$^+$ T cell lymphopenia (Fig. 6D-G), suggesting that a large part of

lymphopenia phenotype in Lpr mice may be CD69 independent. Altogether, CD4$^+$ T cells from

Lpr mice exhibited increased mTORC2 activity. Type I IFN, together with TCR signaling,

activates mTORC2 to induce CD69 expression on CD4$^+$ T cells and blocks CD4$^+$ T cell egress

into blood. RICTOR deficiency partially restores CD4$^+$ T cell lymphopenia in Lpr mice.

DISCUSSION

In this study, we experimentally tested the hypothesis that genetic targeting mTORC2 in T cells

may benefit systemic autoimmunity using a classic systemic lymphoproliferative mouse model,

Lpr mice. We made several key observations: 1) T cells in Lpr mice have elevated mTORC2

activity and mTORC2 inhibition through genetic deletion of RICTOR in T cells can significantly

ameliorate immunopathology of Lpr mice, including increased GC B cells and extrafollicular B

cells; 2) RICTOR deficiency restores Tfh and Treg balance in Lpr mice, without significant

alteration of T cell proliferation and Th1 and Th17 lineages; 3) RICTOR deficiency in Tregs

fails to rectify most of the lymphoproliferative phenotypes in Lpr mice; 4) RICTOR deficiency

reduces ICOS-mediated glycolysis in Lpr T cells. 5) Type I IFN promotes mTORC2 activation

in CD4$^+$ T cells partly to drive CD69 expression and T cell lymphopenia. Thus, our study

suggests that mTORC2 might be a potential therapeutic target for systemic autoimmunity.

Our genetic studies demonstrated that mTORC2-dependent overactivation of Tfh cells, but not

impairment of Tregs, contributes to systemic autoimmunity in Lpr mice. Although previous

study showed deletion of *Rictor* in Tregs restores Treg suppressive activity, including the Tfh

suppressing ability, our data demonstrated that deletion of mTORC2 in Tregs does not improve

their ability to suppress Tfh cells in Lpr mice. These observations suggest that the ability to

promote Tfh differentiation is likely a more dominant function of mTORC2 in systemic

autoimmunity, at least in Lpr background. Furthermore, it is worth noting that while RICTOR

deficiency greatly reduced Tfh differentiation in Lpr mice, it did not significantly affect Th1 and

Th17 subsets, T cell IL-2 production and proliferation. These observations are consistent with

our hypothesis that mTORC2 specifically promotes Tfh differentiation without substantial

effects on T cell activation, proliferation, and other effector T cell lineages. This contrasts with

mTORC1, which is required for T cell quiescence exit and all effector T cell lineages[22]. Thus,

our data suggest that targeting mTORC2 may specifically suppress Tfh lineage, which may have

relatively smaller immunosuppressive side effects, compared to inhibition of mTORC1.

It has long been established that exuberant type I IFN activity contributes to lupus development.

Yet, the molecular mechanism through which type I IFN impact T cells in lupus remains largely

unknown. We show that type I IFN may be one of the upstream signals that activate mTORC2 in

T cells. One of the downstream effects of type I IFN-mTORC2 activation is to promote CD69

expression and T cell lymphopenia. However, our data also showed that mTORC2 is only part of

type I IFN program that controls T cell egress, because RICTOR deficiency had a relatively

small rescue effect on poly(I:C) induced lymphopenia or lymphopenia in Lpr mice. The

prevailing theory on the pathogenesis of the lymphopenia in lupus has been centered on

antilymphocyte antibodies[52]. But a large-scale multiplex single cell RNA-seq analysis

demonstrated a clear inverse correlation between type I IFN activity and circulating naïve CD4$^+$

T cells in SLE patients[12]. Our study corroborates with such observation indicating that elevated

type I IFN production may be another contributing factor for the lymphopenia phenotype in

lupus partly by potentiating mTORC2 activation. Consistent with this idea, a recent clinical trial

showed that type I IFN receptor blockade can correct the T cell lymphopenia phenotype in

patients with SLE[53]. One major limitation of our study is that our conclusions are based on one

genetic mouse model, Lpr mice. Lpr mice carry spontaneous mutation in *Fas* gene. On C57BL/6

background, *Fas* mutation does not lead to full-blown clinical symptoms characteristic of human

SLE, although many of the immune cell hyperactivation phenotypes resemble SLE. Further

investigations using other mouse models and patient samples are needed. Altogether, our results

support the notion that mTORC2 could be a novel target for systemic autoimmunity.

ACKNOWLEDGEMENTS

We thank Drs. Virginia Shapiro and Jie Sun for thoughtful discussions. This work was partly

supported by NIH R01AR077518 and R01Al162678, Mayo Foundation for Medical Education

and Research, and Center for Biomedical Discovery at Mayo Clinic, and Lupus Research

Alliance (696599, to H.Z.).

REFERENCE

1. Muskardin, T.L.W. & Niewold, T.B. Type I interferon in rheumatic diseases. *Nat Rev Rheumatol* **14**, 214-228 (2018).
2. Banchereau, J. & Pascual, V. Type I interferon in systemic lupus erythematosus and other autoimmune diseases. *Immunity* **25**, 383-392 (2006).
3. Liu, Z. & Davidson, A. Taming lupus-a new understanding of pathogenesis is leading to clinical advances. *Nature medicine* **18**, 871-882 (2012).
4. Morand, E.F.*, et al.* Trial of Anifrolumab in Active Systemic Lupus Erythematosus. *The New England journal of medicine* **382**, 211-221 (2020).
5. Klarquist, J.*, et al.* Type I IFN Drives Experimental Systemic Lupus Erythematosus by Distinct Mechanisms in CD4 T Cells and B Cells. *Immunohorizons* **4**, 140-152 (2020).
6. Liu, Z.*, et al.* Interferon-alpha accelerates murine systemic lupus erythematosus in a T cell-dependent manner. *Arthritis Rheum* **63**, 219-229 (2011).
7. Shiow, L.R.*, et al.* CD69 acts downstream of interferon-alpha/beta to inhibit S1P1 and lymphocyte egress from lymphoid organs. *Nature* **440**, 540-544 (2006).
8. Kamphuis, E., Junt, T., Waibler, Z., Forster, R. & Kalinke, U. Type I interferons directly regulate lymphocyte recirculation and cause transient blood lymphopenia. *Blood* **108**, 3253-3261 (2006).
9. Su, C.C., Shau, W.Y., Wang, C.R., Chuang, C.Y. & Chen, C.Y. CD69 to CD3 ratio of peripheral blood mononuclear cells as a marker to monitor systemic lupus erythematosus disease activity. *Lupus* **6**, 449-454 (1997).
10. Tsujimura, S., Adachi, T., Saito, K. & Tanaka, Y. Role of P-glycoprotein on CD69(+)CD4(+) cells in the pathogenesis of proliferative lupus nephritis and non-responsiveness to immunosuppressive therapy. *RMD Open* **3**, e000423 (2017).
11. Oke, V.*, et al.* High levels of circulating interferons type I, type II and type III associate with distinct clinical features of active systemic lupus erythematosus. *Arthritis Res Ther* **21**, 107 (2019).
12. Perez, R.K.*, et al.* Single-cell RNA-seq reveals cell type-specific molecular and genetic associations to lupus. *Science* **376**, eabf1970 (2022).
13. Mountz, J.D., Hsu, H.C. & Ballesteros-Tato, A. Dysregulation of T Follicular Helper Cells in Lupus. *J Immunol* **202**, 1649-1658 (2019).
14. Choi, J.Y.*, et al.* Circulating follicular helper-like T cells in systemic lupus erythematosus: association with disease activity. *Arthritis Rheumatol* **67**, 988-999 (2015).
15. Crotty, S. T Follicular Helper Cell Biology: A Decade of Discovery and Diseases. *Immunity* **50**, 1132-1148 (2019).
16. Odegard, J.M.*, et al.* ICOS-dependent extrafollicular helper T cells elicit IgG production via IL-21 in systemic autoimmunity. *J Exp Med* **205**, 2873-2886 (2008).
17. Poholek, A.C.*, et al.* In vivo regulation of Bcl6 and T follicular helper cell development. *J Immunol* **185**, 313-326 (2010).
18. Lee, S.K.*, et al.* B cell priming for extrafollicular antibody responses requires Bcl-6 expression by T cells. *J Exp Med* **208**, 1377-1388 (2011).
19. Jenks, S.A., Cashman, K.S., Woodruff, M.C., Lee, F.E. & Sanz, I. Extrafollicular responses in humans and SLE. *Immunol Rev* **288**, 136-148 (2019).
20. Mizui, M. & Tsokos, G.C. Targeting Regulatory T Cells to Treat Patients With Systemic Lupus Erythematosus. *Front Immunol* **9**, 786 (2018).

21.   Saxton, R.A. & Sabatini, D.M. mTOR Signaling in Growth, Metabolism, and Disease.
Cell **169**, 361-371 (2017).
22.   Zeng, H. & Chi, H. mTOR signaling in the differentiation and function of regulatory and
effector T cells. *Curr Opin Immunol* **46**, 103-111 (2017).
23.   Zeng, H.*, et al.* mTORC1 and mTORC2 Kinase Signaling and Glucose Metabolism Drive
Follicular Helper T Cell Differentiation. *Immunity* **45**, 540-554 (2016).
24.   Zeng, H.*, et al.* mTORC1 couples immune signals and metabolic programming to
establish T(reg)-cell function. *Nature* **499**, 485-490 (2013).
25.   Shi, H.*, et al.* Amino Acids License Kinase mTORC1 Activity and Treg Cell Function
via Small G Proteins Rag and Rheb. *Immunity* **51**, 1012-1027 e1017 (2019).
26.   Chapman, N.M.*, et al.* mTOR coordinates transcriptional programs and mitochondrial
metabolism of activated Treg subsets to protect tissue homeostasis. *Nat Commun* **9**, 2095 (2018).
27.   Hosokawa, T.*, et al.* Lamtor1 Is Critically Required for CD4(+) T Cell Proliferation and
Regulatory T Cell Suppressive Function. *J Immunol* **199**, 2008-2019 (2017).
28.   Shrestha, S.*, et al.* Treg cells require the phosphatase PTEN to restrain TH1 and TFH cell
responses. *Nat Immunol* **16**, 178-187 (2015).
29.   Huynh, A.*, et al.* Control of PI(3) kinase in Treg cells maintains homeostasis and lineage
stability. *Nat Immunol* **16**, 188-196 (2015).
30.   Charbonnier, L.M.*, et al.* Functional reprogramming of regulatory T cells in the absence
of Foxp3. *Nat Immunol* **20**, 1208-1219 (2019).
31.   Wyman, B. & Perl, A. Metabolic pathways mediate pathogenesis and offer targets for
treatment in rheumatic diseases. *Curr Opin Rheumatol* **32**, 184-191 (2020).
32.   Suarez-Fueyo, A., Barber, D.F., Martinez-Ara, J., Zea-Mendoza, A.C. & Carrera, A.C.
Enhanced phosphoinositide 3-kinase delta activity is a frequent event in systemic lupus
erythematosus that confers resistance to activation-induced T cell death. *J Immunol* **187**, 2376-
2385 (2011).
33.   Kato, H. & Perl, A. Blockade of Treg Cell Differentiation and Function by the
Interleukin-21-Mechanistic Target of Rapamycin Axis Via Suppression of Autophagy in Patients
With Systemic Lupus Erythematosus. *Arthritis Rheumatol* **70**, 427-438 (2018).
34.   Li, W., Titov, A.A. & Morel, L. An update on lupus animal models. *Curr Opin
Rheumatol* **29**, 434-441 (2017).
35.   Lee, K.*, et al.* Mammalian target of rapamycin protein complex 2 regulates differentiation
of Th1 and Th2 cell subsets via distinct signaling pathways. *Immunity* **32**, 743-753 (2010).
36.   Delgoffe, G.M.*, et al.* The kinase mTOR regulates the differentiation of helper T cells
through the selective activation of signaling by mTORC1 and mTORC2. *Nat Immunol* **12**, 295-
303 (2011).
37.   Kelley, V.E. & Roths, J.B. Interaction of mutant lpr gene with background strain
influences renal disease. *Clin Immunol Immunopathol* **37**, 220-229 (1985).
38.   Chen, L.*, et al.* CD38-Mediated Immunosuppression as a Mechanism of Tumor Cell
Escape from PD-1/PD-L1 Blockade. *Cancer Discov* **8**, 1156-1175 (2018).
39.   Clement, R.L.*, et al.* Follicular regulatory T cells control humoral and allergic immunity
by restraining early B cell responses. *Nat. Immunol.* **20**, 1360-1371 (2019).
40.   Yang, J.*, et al.* Critical roles of mTOR Complex 1 and 2 for T follicular helper cell
differentiation and germinal center responses. *Elife* **5**(2016).
41.   Morel, L. Immunometabolism in systemic lupus erythematosus. *Nat Rev Rheumatol* **13**,
280-290 (2017).

42.   Yin, Y.*, et al.* Glucose Oxidation Is Critical for CD4+ T Cell Activation in a Mouse
Model of Systemic Lupus Erythematosus. *J Immunol* **196**, 80-90 (2016).
43.   Yin, Y.*, et al.* Normalization of CD4+ T cell metabolism reverses lupus. *Sci Transl Med*
7, 274ra218 (2015).
44.   Frauwirth, K.A.*, et al.* The CD28 signaling pathway regulates glucose metabolism.
Immunity **16**, 769-777 (2002).
45.   Zheng, Y., Delgoffe, G.M., Meyer, C.F., Chan, W. & Powell, J.D. Anergic T cells are
metabolically anergic. *J. Immunol.* **183**, 6095-6101 (2009).
46.   Shirakawa, K.*, et al.* Obesity accelerates T cell senescence in murine visceral adipose
tissue. *J Clin Invest* **126**, 4626-4639 (2016).
47.   Tahir, S.*, et al.* A CD153+CD4+ T follicular cell population with cell-senescence
features plays a crucial role in lupus pathogenesis via osteopontin production. *J Immunol* **194**,
5725-5735 (2015).
48.   Kalekar, L.A.*, et al.* CD4(+) T cell anergy prevents autoimmunity and generates
regulatory T cell precursors. *Nat Immunol* **17**, 304-314 (2016).
49.   Xie, D.L., Wu, J., Lou, Y.L. & Zhong, X.P. Tumor suppressor TSC1 is critical for T-cell
anergy. *Proc Natl Acad Sci U S A* **109**, 14152-14157 (2012).
50.   Braun, D., Geraldes, P. & Demengeot, J. Type I Interferon controls the onset and severity
of autoimmune manifestations in lpr mice. *Journal of autoimmunity* **20**, 15-25 (2003).
51.   Nickerson, K.M., Cullen, J.L., Kashgarian, M. & Shlomchik, M.J. Exacerbated
autoimmunity in the absence of TLR9 in MRL.Fas(lpr) mice depends on Ifnar1. *J Immunol* **190**,
3889-3894 (2013).
52.   Fayyaz, A.*, et al.* Haematological manifestations of lupus. *Lupus Sci Med* **2**, e000078
(2015).
53.   Casey, K.A.*, et al.* Type I interferon receptor blockade with anifrolumab corrects innate
and adaptive immune perturbations of SLE. *Lupus Sci Med* **5**, e000286 (2018).
597

    FIGURE LEGENDS

Figure 1. *Rictor* deletion in T cells rectifies immunopathology in Lpr mice. (A) Immunoblot

analysis of p-AKT473 in B6 and Lpr CD4$^+$ T cells isolated from peripheral lymph nodes (pLN).

Right, summary of the relative p-AKT473 expression (normalized to that in B6 CD4$^+$ T cells).

(B) Image of peripheral lymph nodes taken from 6 months old B6, Lpr and Lpr.*Rictor* T-KO

mice. Right, summary of total cellularity of lymph nodes. (C) Titers of total immunoglobulin (Ig)

M, IgG, IgG2b and IgG3 were measured by ELISA. (D) Titers of anti-ssDNA (left) and anti-

chromatin (right) were measured by ELISA. Samples were from 9 months old B6, Lpr and

Lpr.*Rictor* T-KO mice. (E) Expression of B220 and TCRβ on lymphocytes. Right, the

frequencies (upper panels) and absolute numbers (lower panels) of B220$^+$TCRβ$^+$, B220$^-$TCRβ$^+$,

B220$^+$TCRβ$^-$ cells. (F) Flow cytometry analysis of CD4 and CD8 T cells among B220$^-$TCRβ$^+$

cells. Right, the frequency (left) and absolute number (right) of CD4$^-$CD8$^-$ T cell population. (E)

and (F) Cells were from peripheral lymph nodes (pLN) of 6 months old B6, Lpr and Lpr.*Rictor*

T-KO mice. (G) The inflammatory cytokine levels in mouse sera samples collected from 5-6

months old B6, Lpr and Lpr.*Rictor* T-KO mice. NS, not significant; * $P < 0.05$, ** $P < 0.01$, ***

P < 0.001, **** *P* < 0.0001. *p* values were calculated with one-way ANOVA. Results were

presentative of 3 (A) independent experiment and pooled from at least 3 (A-G) independent

experiments. Error bars represent SEM.

Figure 2. mTORC2 in T cells is required for B cell hyper-activation in Lpr mice. (A)

Immunofluorescence staining of peanut agglutinin (PNA), CD3 and IgD on spleen sections from

6 months old B6, Lpr and Lpr.*Rictor* T-KO mice. (B) Flow cytometry analysis of GL-7 and IgD

expression among B220$^+$TCRβ$^-$ cells. Right, the frequency of IgD$^-$GL-7$^+$ germinal center (GC)

B cells. (C) Immunofluorescence staining of CD138 and IgD on spleen sections from 6 months

old B6, Lpr and Lpr.*Rictor* T-KO mice. (D) Flow cytometry analysis of B220$^+$CD138^{int} and

B220^{lo}CD138^{hi} two populations. Right, the frequencies of B220$^+$CD138^{int} (left) and

B220^{lo}CD138^{hi} (right) among total lymphocytes. (B) and (D) Cells were from pLNs of 6 months

old B6, Lpr and Lpr.*Rictor* T-KO mice. NS, not significant; *** $P < 0.001$, **** $P < 0.0001$

(one-way ANOVA). Results were pooled from 3 (B and D) independent experiments. Error bars

represent SEM.

Figure 3. *Rictor* deletion reduces Tfh differentiation, normalizes Treg frequencies, but does

not affect Th1 and Th17 differentiation or T cell proliferation. (A) Expression of CXCR5

and BCL6 on B220$^-$CD4$^+$ T cells. Right, the percentages (left) and absolute numbers (right) of

BCL6$^+$CXCR5$^+$ and BCL6$^-$CXCR5$^+$ cells. (B) Expression of PSGL1 and CXCR5 on B220$^-$

CD4$^+$ T cells. Right, the summary of PSGL1^{lo}CXCR5^{lo} extrafollicular (EF) Tfh cell percentages.

(C) Expression of ICOS on B220$^-$CD4$^+$ T cells. Right, the summary of ICOS$^+$ cell frequency

(left) and normalized ICOS median fluorescence intensity (MFI) (right). (D) Flow cytometry

analysis of Treg cells (B220$^-$CD4$^+$FOXP3^{++}). Frequencies of FOXP3$^+$ cells in CD4$^+$ T cells from

spleen (left), pLN (middle) and Peyer's patches (PP) (right). (E) The ratio between Tfh and Treg

frequency (left) and the ratio between Tfh and Tfr frequency (right) in CD4$^+$ T cells from B6,

Lpr and Lpr.*Rictor* T-KO mice. (F) Expression of IFNγ and IL-17 in CD4$^+$ T cells activated by

PMA and ionomycin. Right, frequencies of IFNγ$^+$ cells and IL-17$^+$ cells. (G) Expression of IL-2

in CD4$^+$ T cells activated by PMA and ionomycin. Right, the frequency of IL-2$^+$ cell. (H)

Expression of Ki-67 in CD4$^+$ T cells from B6, Lpr and Lpr.*Rictor* T-KO mice. Right, the

summary of Ki-67$^+$ cell frequency. (A-C, and E-H) Cells were from pLNs of 6 months old B6,

Lpr and Lpr.*Rictor* T-KO mice. NS, not significant; * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, ****

P < 0.0001 (one-way ANOVA). Results were pooled from at least 3 (A-H) independent

experiments. Error bars represent SEM.

Figure 4. *Rictor* deletion in Treg is unable to restore immune dysregulation in Lpr mice.

(A) Frequency of FOXP3$^+$ Treg cells in splenic CD4$^+$ T cells. (B) Image of peripheral lymph

nodes taken from 6 months old B6, Lpr, Lpr.*Rictor* T-KO mice and Lpr.*Rictor* Treg-KO. Right,

summary of total cellularity of lymph nodes. (C) Frequency of B220$^+$TCRβ$^+$ cells in pLN among

B6, Lpr, Lpr.*Rictor* T-KO mice and Lpr.*Rictor* Treg-KO. (D) Frequencies of pre-Tfh (BCL6$^-$

CXCR5$^+$) and Tfh (BCL6$^+$CXCR5$^+$) cells in pLN. (E) Frequency of GC B cells. (F) The ratio

between Tfh and Treg frequency in CD4$^+$ T cells from B6, Lpr, Lpr.R*ictor* T-KO and Lpr.*Rictor*

Treg-KO mice. (G) Serum concentrations of immunoglobulin IgG1, IgG2a, IgG2b and IgA were

measured by LEGENDplex. NS, not significant; * P < 0.05, ** P < 0.01, *** P < 0.001, **** P

< 0.0001 (one-way ANOVA). Results were pooled from at least 3 (A-G) independent

experiments. Error bars represent SEM.

Figure 5. mTORC2 supports TCR/ICOS mediated glucose metabolism in Lpr T cells. (A)

The basal respiration and (B) basal extracellular acidification rate (ECAR) of CD4$^+$ T from 2-,

3-, and 9-months old mice cells under anti-CD3/anti-CD28 overnight activation, respectively.

(C) The ECAR of CD4$^+$ T cells followed by sequential anti-CD3/anti-CD28, and anti-CD3/anti-

ICOS stimulation measured by glycolytic rate assay. Cells were from pLNs of 6 months old B6,

Lpr and Lpr.*Rictor* T-KO mice. Right, summaries of basal glycolysis and compensatory

glycolysis. Rot, Rotenone; AA, Antimycin A; 2-DG, 2-deoxyglucose. NS, not significant; * *P* <

0.05, ** *P* < 0.01, *** *P* < 0.001, **** *P* < 0.0001 (one-way ANOVA). Results were

presentative of 2 (A-B) and 3 (C) independent experiments. Error bars represent SEM.

Figure 6. Type I IFN synergizes with TCR to promote CD69 expression and suppress T cell egress. (A) Immunoblot analysis of p-AKT$_{473}$, AKT (pan), p-STAT1 and p-S6 in CD4$^+$ T cells from B6 and $Cd4^{cre}Rictor^{fl/fl}$ mice stimulated with or without anti-CD3/anti-CD28 in the presence or absence of IFNα for 1 and 6 hours. Right, summaries of the relative p-AKT$_{473}$ expression (normalized to total AKT and then compared with baseline B6 CD4$^+$ T cells) stimulated for 6 hours. (B) Flow cytometry analysis of p-AKT$_{473}$ expression in CD4$^+$ T cells treated with IFNα alone, or in combination with anti-CD3/anti-CD28 overnight. Right, summary of the relative pAKT$_{473}$ expression (normalized to that in B6 CD4$^+$ T cells without any stimulation). (C) Immunoblot analysis of p-AKT$_{473}$, AKT (pan), and p-STAT1 in human CD4$^+$ T cells with or without anti-CD3/anti-CD28 activation in presence or absence of human IFNα for 3 hours. Right, summary of the relative p-AKT$_{473}$ expression normalized to total AKT and then compared with baseline. (D) and (E) B6 and $Cd4^{cre}Rictor^{fl/fl}$ mice were administered with poly(I:C) intraperitoneally. (D) Expression of CD69 in blood CD4$^+$ T cells from B6 and $Cd4^{cre}Rictor^{fl/fl}$ mice after poly(I:C) administration. Numbers indicate the percentages of CD69$^+$CD4$^+$ T cells. Right, summary of CD69$^+$ percentage in CD4$^+$ T cells at baseline or treated with poly(I:C) for 16 h and 40 h. (E) Blood CD4$^+$ T cell counts were determined before and after poly(I:C) treatment. (F) Expression of CD69 in CD4$^+$ T cells from B6, Lpr and Lpr.$Rictor$ T-KO mice. Right, summary of CD69$^+$ percentages among pLN CD4$^+$ T cells. (G) Blood CD4$^+$ T cell counts were determined in 4-6 months old B6, Lpr and Lpr.$Rictor$ T-KO mice. NS, not significant; * $P <$ 0.05, ** $P <$ 0.01, *** $P <$ 0.001, **** $P <$ 0.0001 (A, B, D and E, unpaired Student's t test, C, F and G, one-way ANOVA). Results were presentative of 4 (A, C), or pooled from at least 3 (A-G) independent experiments. Error bars represent SEM.

Figure 1

Figure 2

Figure 3

Figure 4

A spleen

FOXP3+%

● B6
○ Lpr
▲ Lpr.*Rictor* T-KO
□ Lpr.*Rictor* Treg-KO

B

B6   Lpr   Lpr.*Rictor* T-KO   Lpr.*Rictor* Treg-KO

Cell number (× 10⁷)

● B6
○ Lpr
▲ Lpr.*Rictor* T-KO
□ Lpr.*Rictor* Treg-KO

C TCRβ⁺B220⁺ %

● B6
○ Lpr
▲ Lpr.*Rictor* T-KO
□ Lpr.*Rictor* Treg-KO

D BCL6⁻CXCR5⁺%   BCL6⁺CXCR5⁺%

● B6
○ Lpr
▲ Lpr.*Rictor* T-KO
□ Lpr.*Rictor* Treg-KO

E IgD⁻GL7⁺%

● B6
○ Lpr
▲ Lpr.*Rictor* T-KO
□ Lpr.*Rictor* Treg-KO

F Tfh (%) vs Treg (%)

● B6
○ Lpr
▲ Lpr.*Rictor* T-KO
□ Lpr.*Rictor* Treg-KO

G IgG1 (mg/mL)   IgG2a (mg/mL)   IgG2b (mg/mL)   IgA (mg/mL)

● B6
○ Lpr
▲ Lpr.*Rictor* T-KO
□ Lpr.*Rictor* Treg-KO

Figure 5

A

2 months | 3 months | 9 months

Basal Respiration (pmol/min) — B6, Lpr, Lpr.*Rictor* T-KO

B

2 months | 3 months | 9 months

Basal ECAR — B6, Lpr, Lpr.*Rictor* T-KO

C

ECAR (mpH/min) — Rot/AA, 2-DG

Basal glycolysis (pMoles/min)

Compensatory Glycolysis (pmol/min)

B6
Lpr
Lpr.*Rictor* T-KO

Figure 6

Red, IgG
Green, C3

Supplemental Figure 1
(A) Immunofluorescence staining of IgG (Red) and C3 (Green) on mice kidney sections. Right, the scores of IgG and C3 deposition. * $P < 0.05$ (one-way ANOVA). Results were pooled from 2 independent experiments. Error bars represent SEM.

Supplemental Figure 2

(A) Expression of CD38 and IgG1 expression on B220$^+$TCRb$^-$ B cells. Right, the frequencies of CD38$^+$IgG1$^+$ B cells (left) and CD38$^+$ percentages (right). (NS, not significant; * P < 0.05, ** P < 0.01, *** P < 0.001, **** P < 0.0001, one-way ANOVA, Results were pooled from 3 independent experiments. Error bars represent SEM.

Supplemental Figure 3

(A) Gating strategy for analysis of Tfh cells.(B) Flow cytometry analysis of CXCR5 and PD-1 on B220⁻CD4⁺ T cells. Right, the percentages of PD-1⁺CXCR5⁺ cells. (C) Expression of CXCR5 in Tregs. Right, summary of CXCR5⁺ Tfr cell percentages. (D) Expression of CD25 on FOXP3⁺ Tregs. Numbers indicate the percentages of CD25⁻ Tregs. Right, summary of the percentages of CD25⁻ Tregs. (F) Expression of Tbet and Rorgt on on B220⁻CD4⁺ T cells. Right, summary of Tbet⁺ and Rorgt⁺ cell percentages. Cells were from peripheral lymph nodes (pLNs) of 6 months old B6, Lpr and Lpr.*Rictor* T-KO mice. NS, not significant; ** *P* < 0.01, *** *P* < 0.001, **** *P* < 0.0001 (one-way ANOVA). Results were pooled from at least 3 independent experiments. Error bars represent SEM.

A

Supplemental Figure 4

(A) Expression of *Rictor* in sorted CD25$^+$ CD4$^+$ T cells from B6, Lpr, and Lpr.*Rictor* Treg-KO mice. Data represent 2 independent experiments.

Supplemental Figure 5

(A) Flow cytometry analysis of CD44 vs. CD62L in B220⁻CD4⁺ T cells among 2 and 3 months old B6, Lpr and Lpr.*Rictor* T-KO mice, respectively. (B) Summary of cell numbers for B220⁻ CD4⁺ T cells with anti-CD3/anti-CD28 activation for 72 hours. Numbers were normalized to B6 control. (C) Flow cytometry analysis of CD73 vs. FR4 expression in B220⁻CD4⁺ T cells among 6 months old B6, Lpr and Lpr.*Rictor* T-KO mice. Right, summaries of frequencies of CD73^{hi}FR4^{hi} populations. (D) Summary of cell numbers for B220⁻CD4⁺ T cells activated with anti-CD3/anti-CD28, and then with secondary anti-CD3/ICOS stimulation. Numbers were normalized to B6 control. NS, not significant; * P < 0.05, ** P < 0.01, *** *P* < 0.001, **** P < 0.0001 (one-way ANOVA). Results were pooled from at least 3 independent experiments. Error bars represent SEM.

Supplemental Figure 6

(A) Immunoblot analysis of p-STAT1 and p-S6 expression in CD4$^+$ T cells from B6 and *Cd4^{cre}Rictor$^{fl/fl}$* mice treated with or without IFNα in presence of anti-CD3/anti-CD28 for 6 hours. The relative expressions were normalized to them in B6 CD4$^+$ T cells without any stimulation.

(B) Immunoblot analysis of p-AKT473, p-STAT1 and Actin in CD4 T cells from B6 and *Cd4^{cre}Rictor$^{fl/fl}$* mice activated with anti-CD3/anti-CD28 in the presence or absence of IFNb. (C) Blood CD19$^+$ B cell counts were determined before and after poly(I:C) treatment in B6 and *Cd4^{cre}Rictor$^{fl/fl}$* mice. (D) Expression of CD69 was measured by flow cytometry in blood CD4$^+$ T cells from 4-6 months old B6, Lpr and Lpr.*Rictor* T-KO mice. Right, summary of CD69$^+$ percentages among blood CD4$^+$ T cells. (E) Percentages of CD19$^-$CD4$^+$ T cells in blood from B6, Lpr and Lpr.*Rictor* T-KO mice among total lymphocytes. NS, not significant; * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$ (A, unpaired Student's *t* test, C-E, one-way ANOVA). Results were pooled from at least 3 independent experiments. Error bars represent SEM.