Single cell trajectory modeling identifies a primitive trophoblast state defined by BCAM enrichment

Matthew Shannon[1,2], Jennet Baltayeva[,1,2], Barbara Castellana[1,2], Jasmin Wächter[1,2], Samantha Yoon[1,3], Jenna Treissman[1,2], Hoa Le[1,3], Alexander G. Beristain[1,2]

[1] The British Columbia Children's Hospital Research Institute, Vancouver, Canada.
[2] Department of Obstetrics & Gynecology, The University of British Columbia, Vancouver, Canada.
[3] Department of Surgery, The University of British Columbia, Vancouver, Canada.

To whom correspondence should be addressed: Alexander G. Beristain, The British Columbia Children's Hospital Research Institute, The University of British Columbia, Vancouver, British Columbia, Canada. V5Z 4H4. Tel: (604) 875-3573; E-mail: aberista@mail.ubc.ca

Keywords: Placenta, trophoblast, single cell RNA-sequencing, extravillous trophoblast, villous cytotrophoblast, cell differentiation,

Summary Statement: A trophoblast progenitor niche enriched in BCAM shows complex cell trajectory dynamics and gene signaling pathways

Abbreviations
CTB: Cytotrophoblast
DCT: Distal column trophoblast
DGE: Differential gene expression
EVT: Extravillous trophoblast
FDR: False discovery rate
GO: Gene ontology
HLA-G: Human leukocyte antigen G
Hr: Hour
RT: Room Temperature
IF: Immunofluorescence
scRNA-seq: single cell RNA sequencing
SCT: Syncytiotrophoblast
UMAP: Uniform manifold approximation and projection
47

ABSTRACT

The establishment and function of the human placenta is dependent on specialized cells called trophoblasts. Progenitor cytotrophoblasts (CTBs) differentiate along one of two cellular trajectories: the villous or extravillous pathways. CTBs committed to the villous pathway fuse with neighboring CTBs forming the outer multinucleated syncytiotrophoblast layer (SCT), while CTBs committed to the extravillous pathway develop into multi-layered cell columns that anchor the placenta to uterine tissue. At distal column sites, column CTBs differentiate into invasive extravillous trophoblasts (EVT) that facilitate uterine remodeling of spiral arteries and modulate the maternal immune response to fetal antigen. Unfortunately, little is known about the cellular and molecular processes controlling human trophoblast stem cell maintenance and differentiation into these critical trophoblast sub-lineages. To address this, our laboratory established a single cell RNA sequencing (scRNA-seq) dataset from first trimester placentas to identify molecular programs and cell states important in trophoblast progenitor establishment, renewal, and differentiation. Using this dataset, eight distinct trophoblast states were identified, representing progenitor cytotrophoblasts, intermediate column CTBs, syncytiotrophoblast progenitors, and terminally differentiated EVT. Lineage trajectory analysis identified a CTB origin that was reproduced in human trophoblast stem cell organoids. Heightened expression of basal cell adhesion molecule (*BCAM*) defined this primitive state, where CTBs selected for high levels of surface BCAM expression generated larger clonally-derived organoids than did CTB counterparts expressing lower levels of BCAM. Together, this work resolves complex gene networks within human trophoblast progenitor cells, and identifies BCAM as marker that defines an up-stream progenitor origin.

INTRODUCTION

The placenta is a blastocyst-derived tissue that shares the genetic identity of the developing fetus. Derived from cells of the trophectoderm and extraembryonic mesoderm, the human placenta matures into a functional organ by 10-12 weeks' gestation (GA)[1-3]. Through its direct interaction with maternal uterine epithelial, stromal, and immune cells, the placenta coordinates the establishment of the maternal-fetal-interface and serves as a critical barrier important for nutrient-waste exchange between maternal and fetal circulations. Trophectoderm-derived trophoblasts perform many functions of the placenta. These include, but are not limited to the facilitation of nutrient and oxygen transfer between mother and fetus[4], the modulation of the maternal immune response towards the semi-allogeneic fetus and placenta[5,6], and the production of hormones (i.e. chorionic gonadotropin, progesterone, placental lactogen) and other factors required for pregnancy[7].

Of all eutherian mammals, the human placenta is the most invasive[3]. Though humans and rodents both share a haemochorial arrangement of trophoblast and uterine tissue, control of uterine surface epithelial and stromal erosion, as well as artery remodeling by trophoblasts during blastocyst implantation and early placental establishment is far more extensive in humans than in rodents[8]. Importantly, key regulators of trophoblast lineage specification in rodents (i.e. *Cdx2*, *Eomes*, *Esrrb*, and *Sox2*) do not appear to play essential (or identical in the case of *Cdx2*) roles in human trophoblasts[9]. These important differences in human trophoblast biology underlie the need to use human placentas and cell systems for generating fundamental knowledge central to human trophoblast and placental development. While recent advances in regenerative trophoblast culture systems have created the necessary tools required for in-depth cellular and molecular understandings central to trophoblast differentiation[10-12], to date, few studies have used these platforms to examine at high resolution trophoblast progenitor and/or stem cell dynamics.

In humans, two major trophoblast differentiation pathways – villous and extravillous – give rise

to all trophoblasts in the fetal-maternal interface (Figure 1A). In the villous pathway, progenitor

cytotrophoblasts (CTBs) fuse with neighboring CTBs to replenish or generate new syncytiotrophoblast

(SCT), a multinucleated outer layer of the placenta that is the site of transport of most substances across

the placenta, and is also the placenta's major metabolic and endocrine engine. Cells committed to

differentiate along the extravillous pathway are predicted to originate from CTB progenitors proximal

to anchoring villi[9], though populations of villous CTB may already be programmed towards

extravillous differentiation or may alternatively harbor a level of bi-potency[2,13]. Nonetheless,

progenitors devoted to this pathway give rise to extravillous trophoblasts (EVTs) that anchor the

placenta to the uterine wall (i.e. proximal and distal column CTB) and these cells adopt invasive

characteristics hallmarked by interstitial and endovascular EVT subsets that facilitate uterine tissue and

artery remodeling, and interact with and modulate maternal immune cell activity[14-16]. Defects in

trophoblast differentiation along either pathway associate with impaired placental function that may

contribute to, or drive, the development of aberrant conditions of pregnancy[17].

In this study, we integrate an in-house first trimester placenta single cell RNA-sequencing

(scRNA-seq) dataset (7 unique placentas) with a publicly available first trimester placenta/decidual

scRNA-seq dataset consisting of 4 additional samples[18]. Using this data, we investigate trophoblast

heterogeneity across the first trimester and identify a CTB origin state that likely represents a

trophoblast stem cell/progenitor population. Within this upstream progenitor state, we show elevated

expression of the gene *Basal Cell Adhesion Molecule* (*BCAM*), where isolation and culture of BCAM[hi]

CTBs result in robust organoid establishment and growth. Together, this work reaffirms prior findings

related to CTB biology, maintenance, and renewal, and further identifies new gene pathways

underlying distinct CTB progenitor/stem states upstream of SCT and EVT sub-lineage commitment

that are central to placental development.

RESULTS

Single cell transcriptomics resolves trophoblast heterogeneity and cell state dynamics

Single-cell transcriptomic analysis enables the identification of novel cell types and cell states in a systematic and quantitative manner. Here, we applied the 10X Genomics Chromium platform to sequence the healthy human placental transcriptome at single cell resolution (n=7 placentas), and merged this data with publicly available single-cell transcriptomic profiles of healthy human maternal-fetal interface tissues (n=4 placentas, n=4 deciduas) (Figure 1B). To examine developmental differences, we grouped individual samples into early (6-7 weeks') and late (9-12 weeks') gestational age (GA) ranges. Clinical characteristics and sequencing metrics from each patient and sample are summarized in Supplemental Tables 1 and 2. 50,790 cells remained after quality control and data pre-processing (Supplemental Figure 1A-C). Principal component (PC) analysis validated data integration between our placental data and the maternal-placental interface data collected from ArrayExpress[18] with separation of cells explained by tissue type, not data source (Supplemental Figure 1D). These cells were clustered into 14 cell types and 31 distinct cell states by transcriptomic similarity (Supplemental Figure 1E). The influence of fetal gestational age on placental and decidual cell composition was compared, where late first trimester samples showed greater uterine cell coverage than early samples (Supplemental Figure 1F). All clusters were identified by comparing gene expression profiles (Supplemental Figure 1G), and cell cycle phase proportions per cluster informed the level of proliferation in each cell state within the first trimester (Supplemental Figure 1H). Decidual cells were identified by vimentin expression and decidual immune cells were specifically identified by *PTPRC* transcripts that encode the leukocyte CD45 antigen (Supplemental Figure 1I). Additional markers were used to identify endothelial, stromal, and fibroblastic cell compartments of the maternal fetal interface, defining the heterogeneous landscape and dynamic nature of these cells (Supplemental Figure 1I).

For trophoblast identification, cells expressing combinations of trophoblast-related transcripts

(i.e. *KRT7*, *EGFR*, *TEAD4*, *TP63*, *TFAP2A*, *TFAP2C*, *GATA2*, *GATA3*, *HLA-G*, *ERVFRD-1*), and

lacking mesenchymal- and immune-related transcripts (i.e. *VIM*, *WARS*, *PTPRC*, *DCN*, *CD34*, *CD14*,

CD86, *CD163*, *NKG7*, *KLRD1*, *HLA-DPA1*, *HLA-DPB1*, *HLA-DRA*, *HLA-DRB1*, *HLA-DRB5*, and

HLA-DQA1), were subset for further analysis (Supplemental Figure 1G, 1J). Eight distinct trophoblast

clusters from 7,798 cells were resolved (Figure 1C). Within these 8 clusters, we identified four CTB

states (CTB1-4; *EGFR*$^+$, *ELF5*$^+$), a CTB state expressing the SCT-related genes that we termed

progenitor syncytiotrophoblast (SCTp; *ERVFRD-1*$^+$, *CGB*$^+$), two column trophoblast states (cCTB1-2;

NOTCH1$^+$, ITGA5$^+$), and a state specific to EVT (EVT; *HLA-G*$^+$, *ITGA1*$^+$) (Figure 1C, 1D;

Supplemental Table 3). Moreover, CTB1-4 showed varying degrees of enrichment for genes associated

with trophoblast progenitor/stem states (i.e. *CDX2, TP63, ELF5, TEAD4*), where CTB2 and CTB3

showed the greatest enrichment for progenitor genes (Figure 1D).

Column trophoblasts are precursors to EVT, where cells situated within proximal and distal

sites of anchoring columns can be molecularly distinguished by expression of distinct NOTCH and

integrin genes[19,20]. To this end, cCTB2 expresses high levels of *NOTCH1*, *SOX9,* and *ITGA2*, the latter

of which defines a distinct EVT progenitor population[13] and therefore likely represents a proximal

column EVT progenitor (Figure 1D). Compared to cCTB2, cells within the cCTB1 state showed higher

levels of *ITGA5*, *NOTCH2*, and *HLA-G*, indicating that this state is developmentally downstream of

cCTB2 and cells within this state likely reside within central and distal regions of anchoring columns[21]

(Figure 1D). Stratifying cells by phases of the cell cycle showed that cCTB1 column cells are

predominately in G2, M, and S phases, while cCTB2 column cells are not actively cycling (Figure 1E).

As expected, terminally differentiated EVT and SCTp states are largely non-cycling G1 cells, as are

progenitor CTB1-3 states (Figure 1E). This was in contrast to CTB4, where the vast majority of cells

are actively cycling and express hallmark genes linked to proliferation (i.e. *MKI67*, *CCNA2*) (Figure

1D-F). Importantly, these distinct trophoblast states largely align with recent single cell transcriptomic

characterizations of cells from first trimester pregnancies[18,22,23].

Having defined eight trophoblast types/states within first trimester placental villi, we next

examined their stability across the first trimester developmental period. Relative steady-state

frequencies in CTB1 and CTB3 states were observed in placentas from 6-7 weeks' and 9-12 weeks'

GA, whereas more dynamic changes were seen in other states (Figure 1G). The proportion of cells

comprising the proliferative CTB4 state decreased slightly from the early (18%) to late (11%) first

trimester (Figure 1G). Moderate decreases in frequency were also seen in cells within column (cCTB1,

cCTB2), EVT, and SCTp states (Figure 1G). This was in mark contrast to CTB2, which showed a ~3-

fold increase from 8% to 29% (Figure 1G). Together, these findings define unique states of trophoblast

heterogeneity in the early and late first trimester developmental window, a developmental period

correlating with an onset in maternal blood perfusion within the intervillous space around 10 weeks'

GA, and a significant expansion/branching of floating placental villi.

Defining known and novel gene programs in CTB progenitor maintenance and differentiation

With the recent development of regenerative human trophoblast models that better facilitate the

study of progenitor CTB differentiation into SCT and EVT, significant strides have been made in our

understanding of key pathways underlying CTB establishment, maintenance, and the initial molecular

programs driving CTB differentiation[10-12,24,25]. For example, WNT, HIPPO, and EGFR signaling are

essential in human trophoblast stem cell establishment[10-12], and impairments in trophoblast stem cell

regeneration in the context of YAP-TEAD4 complex perturbation[24,25] underscore the necessity of these

pathways in placental development. To examine how previously described pathways critical to CTB

development are altered during differentiation, we applied a simple branched pseudotime trajectory

analysis (using Monocle2) to order cells and states in pseudotime (Supplemental Figure 2A, 2B). We

focused on the transitions occurring between trophoblast cell type, and more specifically cell state, with

increased pseudotime maturation along either of the defined villous or extravillous trajectories

(Supplemental Figure 2B). While simple branched pseudotime ordering did not resolve a

developmental lineage within the four CTB states, a predicted cell state/type origin was identified

comprising cells within CTB1-4, and pseudotime ordering did delineate a transition from cCTB2 into

cCTB1 which further progress towards EVT (Supplemental Figure 2B). Monocle2 ordering also

highlighted the transition from CTB to SCTp; these findings are in line with recent pseudotime

ordering characterization of trophoblasts showing a defined origin and two differentiation trajectory

branches[18].

Aligning WNT, HIPPO, and EGFR pathway genes to the Monocle2 pseudotime trajectory

showed distinct patterns of gene expression in cells with proximity to the predicted origin, within

intermediate states along both the SCT and EVT differentiation pathways, as well as in terminally

differentiated SCTp and EVT states (Supplemental Figure 2C, 2D). In agreement with previous studies,

Dickkopft-1 (*DKK1*; an inhibitor of Wnt signaling), leucine-rich repeat-containing G-protein coupled

receptor 6 (*LGR6*; a Wnt receptor), the Wnt ligand genes *WNT5B* and *WNT5A*, the HIPPO signaling

genes transcriptional enhancer factor TEF-3 (*TEAD4*) and protein salvador homolog 1 (*SAV1*), and the

EGFR ligand betacellulin (*BTC*) are enriched within CTB cell states aligning to or proximal to the

predicated origin (Supplemental Figure 2C, 2D). As trophoblasts progress along the EVT pathway,

expression levels of specific Wnt-related genes (i.e. *AXIN2*, *WNT11*, *WNT9A*, *FZD4*) and the EGFR

ligand *AREG* transiently increase within intermediate cCTB states (Supplemental Figure 2C, 2D).

Nearing terminal EVT differentiation, the Wnt-ligand *WNT10B* markedly increases, as do specific

HIPPO-related transcription factors (*TEAD2*, *TEAD1*), and modulators (*FAT1*, *FAT2*, *WWTR1*), and the

EGFR receptor *ERBB2* (Supplemental Figure 2C, 2D). Along the villous trajectory, SCT progenitors

up-regulate the Wnt receptor *FZD1* and multiple Wnt ligands (*WNT2*, *WNT2B*, *WNT3A*, *WNT7A*), as

well as the HIPPO pathway transcription factor TEAD3 and the EGFR's *ERBB3* and *ERBB4*

(Supplemental Figure 2C, 2D). Together, lineage trajectory modeling and alignment of key

genes/pathways known to be important in CTB regeneration and differentiation provides a

comprehensive landscape of distinct effectors and regulators putatively controlling trophoblast

development in early pregnancy.

To resolve the genetic similarities and differences between distinct CTB states, all four CTB

states were subset for targeted transcriptomic analyses (Figure 2A). Next, we compared CTB1-4 using

the top 10 unique genes expressed in each state (Figure 2B). This approach characterized CTB1 as a

translationally active cell state, CTB2 as a cell state made up of more classically defined CTBs (i.e.

expressing *SPINT1*), CTB3 as a metabolically active state defined by mitochondrial gene enrichment,

and CTB4 as a proliferative state (Figure 2B). The similarities in gene expression across all four CTB

states revealed 944 genes upregulated within all CTBs (compared to more differentiated trophoblast

states), including the known CTB markers *TP63*, *EGFR*, and *TEAD4* (Figure 2D; Supplemental Table

4). Specific to each CTB state, we found upregulation of *PGF* in CTB1, upregulation of the

extracellular laminin gene *LAMC2* and the transcription factor *HOXB2* in CTB2, upregulation of

GATA2, *FZD3*, and *AXIN2* in CTB3, and upregulation of *TFAP4* and *TFAP2B* in CTB4 (Supplemental

Table 4). Gene ontology terms enriched from those genes uniquely upregulated in each state revealed

that CTB1 is involved in general metabolic processes (Figure 2E), CTB2 is involved in cell stemness

and general responses to hypoxia (Figure 2E), CTB3 has terms describing the suppression of

mitochondrial activity and promoting cell anabolism (Figure 2E), and CTB4, is involved in cell cycle

progression and chromosome organization/segregation (Figure 2E). Finally, *cis*-regulatory sequence

analysis of the enriched genes per CTB state revealed a list of 62 transcription factors (TFs) that may

act as CTB-specific regulators (Figure 2F). Included in this list are TFs known to be important in CTB

biology like *TEAD4* and *TFAP2A*[25]. Interestingly, TFs such as transcriptional regulator KAISO

(*ZBTB33*) involved the negative regulation of the WNT pathway[26] shows enrichment in CTB3. These

findings highlight the cellular and molecular complexity within the CTB compartment of the placenta,

and identify possible transcriptional regulators important in controlling the maintenance of progenitor

CTBs as well as regulating the possible transition between unique CTB states.

Single cell trajectory modeling identifies a CTB origin

RNA velocity, the time derivative of the gene expression state, can be used to predict the future

transcriptional state of a cell. RNA velocity analysis in combination with single-cell trajectory analyses

provides us with insights into the transcriptional dynamics of developing cells and enables high-

resolution lineage pathway ordering. By applying scVelo to our dataset to further refine progenitor and

cell end-point relationships, we identified CTB2 as a first trimester placental trophoblast progenitor

origin from which all other trophoblasts derive (Figure 3A). We identified two paths extending from

this origin towards the EVT state and one path developing from the origin towards the SCTp state. As

well, we found an additional differentiation trajectory, extending from CTB2 into CTB3, representing

possible CTB progenitor reserves or progenitor states in transition.

To more completely understand the developmental steps involved in CTB differentiation, we

applied complex pseudotime ordering using Monocle3. Similar to RNA velocity, CTB2 was identified

as the CTB origin showing increased complexity as CTBs progress along one of two trajectories that

lead to either EVT or SCT end-point states (Supplemental Figure 3A, 3B). Specifically, complex

pseudotime ordering revealed a looped trajectory within CTB2 that exits towards CTB3, SCTp, and

EVT states (Supplemental Figure 3C). We compared these refined trajectories with expression patterns

of known CTB progenitor/stem cell genes. Predominant expression of *ELF5*, *YAP1*, and *ITGA6* was

observed in CTB3, while expression of CTB stem cell transcription factor *CDX2* showed predominant

expression within the CTB2 state (Figure 3B). Because CTB2 and CTB3 shared expression of multiple

trophoblast progenitor genes, we examined genes both shared and unique to these states (Figure 3C).

Twenty-five genes were shown to overlap between CTB2 and CTB3, including the genes *EGFR, IGF2,*

and *FBLN1* (Figure 3C). The CTB3 progenitor state expressed 274 unique genes, and of these genes,

many associate with trophoblast stem cell and regenerative properties (i.e. *YAP1, TP63*), Wnt signaling

(*LRP5*), and cell-matrix interactions (*MMP14, MMP15, ITGA6, LAMA5, LAMB1*) (Figure 3C). The

predicted CTB2 origin expressed 269 unique genes, including the Wnt ligand, *WNT7A*, the laminin

receptor basal cell adhesion molecule (*BCAM*), and the transcriptional modulators *TFAP2A* and

CITED2 (Figure 3C). The expression patterns of select genes in these CTB states over pseudotime

support scVelo and Monocle3 trajectory ordering suggesting that CTB2 is upstream of CTB3 (Figure

3D; Supplemental Figure 3C). Focusing specifically on the CTB2 origin state, we identified the

enrichment of *BCAM*, a gene encoding a laminin cell surface receptor (Figure 3E). Also enriched

within this primitive CTB2 state were the genes *SOD3*, *GPC3*, *EFEMP1*, *EPN3*, and *C10orf54* (Figure

3E). Cumulatively, we have identified a primitive CTB state within the first trimester placenta that

shows molecular features of a putative CTB stem cell population possessing a unique gene expression

profile that includes the early stem marker *CDX2* and the cell surface glycoprotein *BCAM*.

Trophoblast stem cells recapitulate scRNA-seq-informed CTB heterogeneity and hierarchy

Recently described regenerative human trophoblast models provide an improved method for

modeling trophoblast differentiation *in vitro*[10]. Three-dimensional organoid-like cultures of human

trophoblast stem cell lines (hTSCs) have also been shown to reproduce molecular features underlying

CTB maintenance and differentiation[25]. To examine the reproducibility of trophoblast-intrinsic

differentiation programs defined by scRNA-seq analysis of *in vivo* placental specimens, we subjected

three hTSC lines to scRNA-seq analysis prior to and following EVT-directed differentiation. Regenerative and EVT differentiating hTSC organoids (i.e. CT27, CT29, and CT30 CTB lines) were cultured for 7 days in regenerative Wnt$^+$ (containing the Wnt potentiators R-spondin and CHIR99021) and EVT-differentiating Wnt$^-$ (lacking R-spondin and CHIR99021) media, respectively, prior to single cell workflow. To confirm the intactness of our model system, a separate cohort of hTSC organoids were cultured over 21 days in these media and differentiation was assessed by light microscopy, flow cytometry, and qPCR analysis. hTSC-derived organoids cultured in regenerative media established large spherical colonies characterized in part by elevated expression of the CTB progenitor genes *TEAD4* and *ITGA6* (Figure 4A, 4C). Moreover, expression of surface MHC-I (HLA-A, -B, -C) by flow cytometry did not detect these MHC-I ligands in undifferentiated cultures, (Figure 4B). Comparatively, hTSC organoids cultured in EVT differentiating Wnt$^-$ media showed characteristic cellular elongations and extensions from the organoid body (Figure 4A), as well as a marked decrease in levels of *TEAD4* mRNA (Figure 4C). As expected, both immature (*ITGA5*, *NOTCH1*) and mature (*NOTCH2*, *HLA-G*) EVT marker genes, as well as HLA-C and HLA-G surface expression increased along the EVT differentiation time-line (Figure 4B, 4C), indicating that the 3D organoid model to study CTB regeneration and EVT differentiation is intact.

13,395 regenerative and 40,572 EVT-differentiated hTSC organoid cells were then used to generate single cell cDNA libraries following the 10X Genomics Chromium workflow. Combining single cell transcriptomic data from CT27, CT29, and CT30 hTSC organoids resolved 7 distinct cell states based on global gene expression signatures; three CTB (CTB1-3), a pre-syncytiotrophoblast (SCTp), a putative column/EVT progenitor (cCTB), and two EVT-like states (EVT1, EVT2) were identified (Figure 4D). Hallmark gene markers of trophoblast-lineage (*TFAP2A, TFAP2C*), CTB progenitors (*TEAD4, YAP1, TP63, ITGA6, ELF5*), syncytiotrophoblast (*ERVFRD-1, ERVV-1, SDC1*), immature (*ITGA5, ITGA2, NOTCH1*) and mature EVT (*ITGA1, NOTCH2, HLA-G*), as well as gene

markers of proliferation (*MKI67, CCNA2*) are shown corresponding to individual cell states (Figure

4D). CTB1-3 shared great overlap in gene expression, though CTB1 distinctly expressed proliferative

genes that also co-expressed at high levels the progenitor genes *TEAD4*, *TP63*, and *ELF5* (Figure 4D).

While CTB2 and CTB3 also expressed progenitor genes, their levels were comparatively lower than

levels in CTB1 (Figure 4D). The putative column/EVT progenitor state cCTB uniquely expressed

ITGA2, *SOX9*, and the immature EVT marker *NOTCH1*; at the gene level, this unique state resembles

the EVT progenitor population described in Lee *et al*[13], though unlike that progenitor population, this

state did not express high levels of proliferative genes (Figure 4D). States resembling more terminally

differentiated cell types showed strong enrichment for syncytial-like genes (SCTp) and EVT-associated

genes, with EVT1 showing the highest expression of *HLA-G* and *NOTCH2* (Figure 4D).

When scRNA-seq data for hTSC-derived organoids are plotted individually, overall cell state

complexity diminishes from 7 unique states to 5 states (Figure 4E). When comparing the three hTSCs

with each other, notable variances in cell states are observed (Figure 4E). While clearly defined

endpoint states are evident, variances defined by gene expression in CTB and cCTB states suggest that

the hTSC lines harbor cell differentiation kinetics that are different from each other (Figure 4E).

Nonetheless, within each hTSC-derived organoid, a CTB origin was revealed by estimation of RNA

velocity that overlapped with CTB1 (Figure 4E). When this origin was compared across each hTSC

line by relative gene expression, we found consistent up-regulation of the trophoblast stem cell genes

TP63, *YAP1*, and *PAGE4* compared to other states. However, we did see differences in the expression

of the *ELF5* stem gene between hTSCs and individual states. Expression of *ELF5* localized to the

origin in the CT27 hTSC line, but not within the origins of the CT29 or CT30 hTSC lines (Figure 4E).

Similar differences were observed for the proliferative gene markers *MKI67* and *CCNA2*, with positive

expression occurring in the origins of CT27 and CT30, but not in the origins of CT29 (Figure 4E).

When averaged across all hTSC lines, however, we observed consistent expression of *ELF5*, *BCAM*,

MKI67, and *CNNA2* within the CTB origin (Figure 4D). Comparison of transcriptional signatures

between the scVelo-determined origins in *in vivo* trophoblasts from chorionic villi and hTSC-derived

organoids highlights that there is considerable overlap between the CTB2 & 3 (*in vivo*) and CTB1& 2

(*in vitro*) states (Figure 4F). Notably, we show *BCAM* to be one of the most highly expressed genes

within primitive states identified *in vivo* and in hTSC (Figure 4G, 4H). This suggests that BCAM may

be a novel CTB stem cell or progenitor marker.

BCAM-enrichment confers increased CTB regenerative potential

To our knowledge, BCAM localization within the human placenta has never been reported. To

this end, we probed early (6.3 week) and late (11.2 week) first trimester placental villi specimens with

BCAM antibody (Figure 5A). A specific and strong BCAM signal was observed in CTB of both

developmental time-points, and serial sections stained with keratin-7 (KRT7) and HLA-G showed that

BCAM was present within proximal column cells, though the intensity of the signal was less than in

CTB (Figure 5A). Serial sections of villi stained with KRT7 and hCG clearly showed that BCAM is

absent from the syncytiotrophoblast outer layer (Figure 5A). In hTSC-derived organoids cultured in

regenerative Wnt$^+$ media, BCAM signal was specific to the outer CTB layer and absent within the

hCG$^+$ syncytium (Figure 5B). Dual-labeling with BCAM and Ki67 antibody in CT29 hTSC-derived

organoids showed that BCAM$^+$ CTB and proximal column trophoblasts also stained positive for the

proliferation marker (Figure 5B). Flow cytometry, which enables a quantitative measure of BCAM cell

frequency, showed that in dissociated EGFR$^+$ CTB from CT29-derived organoids, ~90% of CTBs are

BCAM positive (Figure 5C).

To test the regenerative potential of BCAM-expressing cells, BCAM^{hi} and BCAM^{lo} CTBs (i.e.

EGFR$^+$ trophoblasts) were sorted from CT29 hTSC organoids and seeded as single cells at a density of

125 cells/µl within Matrigel domes and cultured over 11 days (Figure 5D). While the colony take-rate

did not differ between BCAM^{hi} and BCAM^{lo} CTB organoids (Supplemental Figure 4), BCAM^{hi}

organoid colony size appeared to be larger by day 7 of culture, with BCAM^{hi} colonies being

significantly larger than BCAM^{lo} colonies on days 9-11 in culture (Figure 5E). Notably, flow

cytometry assessment of endpoint day 11 BCAM^{hi} and BCAM^{lo} organoids showed that while BCAM^{hi}-

sorted cells retained frequencies of >90% BCAM high-expressing cells (Figure 5F), the frequency of

CTBs expressing BCAM in BCAM^{lo}-sorted cells was only marginally less, suggesting that the BCAM

phenotype is dynamic. Comparing differentially expressed genes between BCAM^{hi} and BCAM^{lo}

EGFR+ CTBs showed that BCAM^{hi} cells expressed higher levels of the paternally imprinted

transcription factors *PEG3* and *PEG10* involved in potentiating progression through the cell cycle and

stem cell renewal genes *EPCAM* and *HMGB3*[27] (Figure 5G). In summary, these data show that

enrichment of BCAM high-expressing CTB confers organoid growth and regenerative potential.

DISCUSSION

Here we applied single cell RNA sequencing of human placentas at two gestational age time-points within the first trimester to further our understanding of human trophoblast development. In doing this, we provide here novel transcriptomic data from 7,798 human trophoblasts, resolve the first trimester trophoblast landscape at increased resolution, and clarify how gestational age informs trophoblast heterogeneity, composition, and gene expression. To our knowledge, this is the first trophoblast-focused study that accounts for multiple gestational age time-points within the first trimester. Using state-of-the-art single cell lineage trajectory tools, we trace back the lineage commitments made by developing CTBs to identify a novel trophoblast progenitor cell niche marked by increased expression of the cell surface glycoprotein BCAM.

Previous studies have explored the maternal-placental interface transcriptome[18] as well as the placental transcriptome across gestation at single cell resolution[22,23]. While these studies have contributed important insight into placental cell heterogeneity, complexity, and have defined detailed maternal-trophoblast interactions, a strict focus on understanding CTB hierarchy, the CTB progenitor transcription factor regulatory landscape, and gene pathways underlying CTB commitment to EVT or SCT has not been specifically addressed. Our study defines four CTB states. The first CTB state is identified by increased expression of large (RPL) and small (RPS) ribosomal subunit genes. Upregulation of these ribosomal subunit genes suggests increased ribosome content, and therefore increased protein translation occurring within these cells[28]. Some studies have reported a link between ribosomal subunit protein activity and regulation the cell cycle through P53-mediated arrest or the promotion of proliferation[29]. The mixed expression of both activators (*RPS15A*[29]) and inhibitors (*RPS14*[29]) of proliferation within this CTB state suggests that this cell state may represent a transient CTB population increasing expression of ribosome genes to regulate entry into the cell cycle[28,29]. Further, previous work by Khajuria et al[30] has described a link between ribosome protein level and cell

differentiation, suggesting an alternative explanation that these cells are increasing expression of

ribosome subunits to facilitate differentiation out of the trophoblast stem cell state.

The second CTB state we define is a population of proliferative CTBs. Previous studies have

proposed that proliferating CTBs will generally contain trophoblast stem cells[23], allowing these

dividing cells to supply and replenish the placenta across gestation. Expression patterns of known

trophoblast stem cell markers, however, shows little overlap with these cells in our data, the exception

being *TEAD4*, which is identified as a potential key regulator of this state. Previous work has

demonstrated multiple roles for the TEA domain (TEAD) proteins in cell proliferation, differentiation,

and apoptosis[31-34]. Specifically, TEAD4 co-activation with VGLL1/2 proteins is shown to promote

proliferation, while TEAD4 co-activation with the Hippo pathway effector protein YAP has been

demonstrated to promote trophoblast stemness[24]. Comparison of the expression patterns of *TEAD4* and

YAP1 (encoding the YAP protein), in our proliferative CTBs and in CTB2 suggests a dual role of

TEAD4 expression in the CTB, dependent on the expression of these co-activators. Within our

proliferative CTBs, *TEAD4* is co-expressed with *VGLL1* without *YAP1*, demonstrating that here

TEAD4 is driving a proliferative and not stem phenotype.

The third CTB state identified is a metabolically active state, identified by increased expression

of mitochondria genes. Previous sequencing reports have noted that increased percent expression of

mitochondrial RNA may be indicative of cell stress or damage[35-37]. However, our quality control

pipeline filters on the basis of percent mitochondrial RNA content. As well, comparison of cell specific

feature detection and the detection of unique molecular identifiers indicates no observable difference

that would support the presence of damaged or dying cells. As a result, we theorize that these cells are

expressing mitochondrial RNA transcripts in response to increased metabolic demand. Unique to this

CTB state is the upregulation of many genes shown to be involved in trophoblast stemness, suggesting

a role for mitochondrial activity in CTB stemness. Fitting with this, recent work has shown a direct role

for WNT signaling in the promotion of CTB stemness[10-12], suggesting that these cells, which express a

CTB progenitor-like transcriptome but are not the CTB progenitors, may represent a transient CTB

state responding to WNT activation. To further support this hypothesis, recent work has demonstrated a

correlation between WNT activity and mitochondrial biogenesis/activation[38,39], however further work

is needed to test this in the placenta. Interestingly, three transcription factors were identified as master

regulators of this CTB state, ZBTB33, TFCP2L1, and FOXK1, all of which show involvement in the

regulation of WNT signaling.

We then sought to identify and characterize a trophoblast stem cell niche. Our initial approaches

involved branched pseudotime trajectory inference, which captured transitions between cell type as

CTB differentiate into SCT or EVT, however, this approach proved insufficient for delineating the

CTB specific developmental hierarchy. To circumvent this and identify the most primitive trophoblast

population, we applied a sophisticated RNA velocity approach for quantifying the relative amounts of

spliced and unspliced gene abundances. RNA velocity was able to identify CTB2, the fourth state, as

an origin population from which all trophoblasts arise. Within these cells, we observe increased

expression of the most primitive trophoblast stem cell gene *CDX2*, further indicating that these cells

represent a primitive CTB state. Gene ontology enrichment analysis of genes in CTB2 identified

biological process terms relating to hematopoietic progenitor cell differentiation and cellular response

to hypoxia. The first 9 weeks of gestation are characterized by a hypoxic oxygen concentration < 20

mm Hg that is maintained until maternal trophoblastic plugs are eroded at 10 weeks GA, bringing the

oxygen concentration of the maternal fetal interface to 40-80 mmHg[40-43]. Enrichment of hypoxia cell-

related terms within this origin indicates that these cells are highly sensitive to their environment, but

also that hypoxia-driven programs may play roles in coordinating progenitor functions in CTB. Indeed,

exposure to low levels of oxygen drive CTBs to differentiate into immature EVT[44-46], but the

contribution of hypoxia on progenitor biology is not as well developed. Future work aimed at

understanding the relationship between hypoxia and stemness in trophoblast development, including

the importance of HIF transcription factors in stem cell maintenance will be important.

Comparisons between our placental tissue sequencing data and our trophoblast organoid

sequencing data demonstrate similarities in general CTB, SCT and EVT function, but also subtle

differences between the different states of CTB found in each stem cell derived organoid condition and

our tissue CTBs. Comparison of the CTB progenitor niche identified in both the organoid and tissue

datasets identifies the common transcription factors *YAP1*, *SPINT2*, *PEG10*, and *MSI1* as well as

consistent upregulation of *BCAM*, supporting identification of BCAM as a CTB progenitor marker.

Interestingly, quantification of mRNA levels of known CTB related genes (*TEAD4* and *ITGA6*) as well

as SCT (*CGB* and *CSH1*) and EVT related genes (*ITGA5*, *NOTCH1*, *NOTCH2*, and *HLA-G*) on days 0,

11, and 21 of differentiation indicate that hTSC specific organoid colonies show different rates of

differentiation or capacities to differentiate. Our EVT-differentiated organoid transcriptomes were

sequenced on day 7 of differentiation, indicating that the differences we are observing may in part be

due to groups of organoid colonies being at a different stage of differentiation, with CT29 derived

organoids showing the most rapid capacity for EVT differentiation, followed by CT30 and CT27.

Consistent between our *in vivo* and *in vitro* datasets is the identification of two CTB progenitor

cell niches in the first trimester placenta, one representing a CTB origin state marked by increased

BCAM gene expression, and one representing a more specified bipotent EVT progenitor cell state,

marked by *ITGA2* and *NOTCH1* gene expression. Immunofluorescence microscopy and

immunohistochemistry of 8-9 week GA placental villi shows that ITGA2 is specific to proximal cells

within anchoring placental villi near the basement membrane[13]. It is thought that as chorionic villi

develop, trophoblast progenitors within close proximity to uterine stroma adopt a unique gene

expression profile that predisposes their differentiation trajectory to the extravillous pathway. We show

that the transcriptome of *ITGA2*[+] putative column progenitors express EVT-associated genes such as

NOTCH1 and *ITGA5,* as well as the CTB/progenitor-associated genes *TFAP2C*, *GATA3*, and *ELF5*.

Lee et al[13] further describe ITGA2$^+$ cCTBs as having a mixed endothelial- and mesenchymal-like

identity. This unique gene signature, harboring stem-like and EVT-like genes, supports the possibility

that *ITGA2$^+$* CTB retain a level of bipotency. Analyses using scVelo trajectory inference suggests

*ITGA2$^+$*column/EVT progenitors descend from the *BCAM$^+$* CTB progenitor origin and are maintained

as a reserve progenitor population within anchoring columns.

We identify *BCAM* as a gene expressed by CTB progenitors, where its expression is particularly

elevated in the most upstream CTB states. BCAM interacts predominantly with its ligand *LAMA5*, a

component of the laminin-511 heterotrimers found within the placental extracellular matrix[47,48]. BCAM

is an antigen upregulated in ovarian carcinoma, and facilitates tumor migration by competing for

laminin binding with integrins [49,50,51]. Here, we show that flow-cytometry enrichment of BCAM in

CTBs significantly potentiates CTB organoid growth. Interestingly, no differences were observed in the

capacity of BCAM^{hi} or BCAM^{lo} CTBs to form organoid colonies from single cells, suggesting that

BCAM^{hi} CTB may not have intrinsically greater stem-like properties over BCAM^{lo} CTB. Rather,

BCAM may enhance organoid growth by potentiating survival signals induced by cell-ECM

engagement, or by initiating/driving CTB proliferation pathways. However, we show that genes

associated with stem cell regeneration are elevated in BCAM^{hi} hTSC-derived organoids as compared to

organoids derived from BCAM^{lo} cells. Specifically, we show increased expression of *PEG10*, *EPCAM*,

and *HMGB3*, indicating that BCAM may confer increased organoid growth by controlling stem cell

maintenance or regeneration. In our experimental organoid system, even though BCAM^{lo} organoids

form smaller colonies by day 11, assessment of BCAM expression in BCAM^{lo} organoids at end-point

shows that BCAM levels and the frequency of BCAM-expressing cells is comparable to BCAM^{hi}

CTBs. This suggests that BCAM expression in progenitor CTBs is dynamic, and that requirements for

growth, stemness, or survival result in the induction of BCAM expression. Future work designed to

examine how BCAM drives trophoblast organoid growth will be necessary.

<u>SIGNIFICANCE</u>

The identification of a novel cell surface human CTB progenitor cell marker provides improved

methods for isolating and studying the mechanisms regulating human CTB progenitor cells, trophoblast

differentiation, and human placental development.

<u>CONCLUSIONS</u>

In conclusion, we have used single cell RNA sequencing to define trophoblast differentiation

trajectories in early human placental development. Importantly, we identify BCAM as a new CTB

progenitor marker that confers enhanced clonal growth. Future work is needed to clarify how increased

BCAM expression facilitates improved CTB growth, and the exact molecular paths through which this

occurs. We hope that future experiments will be able to capitalize on the identification of BCAM as a

cell surface CTB progenitor marker through which these cells can be isolated and further studied.

MATERIALS AND METHODS

Patient recruitment and tissue collection

Decidual and placental tissues were obtained with approval from the Research Ethics Board on the use of human subjects, University of British Columbia (H13-00640). All samples were collected from women (19 to 39 years of age) providing written informed consent undergoing elective terminations of pregnancy at British Columbia's Women's Hospital, Vancouver, Canada. First trimester placental tissues (*N*=7) were collected from participating women (gestational ages ranging from 5–12 weeks) having confirmed viable pregnancies by ultrasound-measured fetal heartbeat. Patient clinical characteristics i.e. height and weight were additionally obtained to calculate body mass index (BMI: kg/m^2) and all consenting women provided self-reported information via questionnaire to having first hand exposure to cigarette smoke or taken anti-inflammation or anti-hypertensive medications during pregnancy. Patient and sample phenotype metadata are summarized in Supplemental Table 1.

Human trophoblast organoids

Establishment and culture: Human trophoblast stem cells (hTSC) CT27 (RCB4936), CT29 (RCB4937), and CT30 (RCB4938) from Riken BRC Cell Bank, were established from first–trimester CTBs by Okae et al[10] and used here to generate organoid cultures using a modification of previously described protocols[11,12]. In summary, these cells were re-suspended in ice-cold growth factor-reduced Matrigel (GFR-M, Corning) to a final concentration of 100%. Cell suspensions (40µl) containing 5-20 x 10^3 cells were carefully seeded in the center of each well of a 24-well plate. Plates were placed at 37°C for 2 minutes. Following this, the plates were turned upside down to promote equal spreading of the cells throughout the Matrigel domes. After 15 minutes, the plates were flipped back to normal and the Matrigel domes were overlaid with 0.5mL pre-warmed trophoblast organoid media containing advanced DMEM/F12 (Gibco), 10mM HEPES, 1 x B27 (Gibco), 1 x N2 (Gibco), 2mM L-glutamine

(Gibco), 100μg/ml Primocin (Invivogen), 100ng/mL R-spondin (PeproTech), 1μM A83-01 (Tocris),

100ng/mL rhEGF (PeproTech), 3μM CHIR99021 (Tocris), and 2μM Y-27632 (EMD Millipore).

Organoids were cultured for 7-21 days.

EVT Differentiation: EVT differentiation of trophoblast organoids was performed using an

adapted of previously published protocols[10]. In summary, organoids were cultured until > 50% colonies

reached at least 100μm in diameter (~3-5 days). Following this, organoids were maintained in EVT

differentiation media containing advanced DMEM/F12, 0.1mM 2-mercaptoethanol (Gibco), 0.5%

penicillin/streptomycin (Gibco), 0.3% BSA (Sigma Aldrich), 1% ITS-X supplement (Gibco),

100ng/mL NRG1 (Cell Signaling Technology), 7.5μM A83-01 (Tocris), 4% Knockout Serum

Replacement (Gibco). Media exchanged for EVT differentiation media lacking NRG1 around day 7-10

of differentiation following the visible outgrowth of cells. Differentiation was stopped at day 7 (D7) for

10X single cell RNA sequencing and flow cytometry analysis. Differentiation was stopped at day 21

(D21) for qPCR analysis.

Placental tissue and trophoblast organoid single cell RNA-seq library construction

Placental tissue: Healthy placental villi (BMI < 25.00 kg/m², MA: 20-32 years, and no

exposure to first hand cigarette smoke or anti-inflammation and anti-hypertensive medications during

pregnancy) were physically scraped form chorionic membrane by scalpel and enzymatically digested in

0.5ml collagenase/hyaluronidase and 1.5ml DMEM/F12 with 500μl penicillin/streptomycin and 500μl

100X antibiotic/antimycotic at 37˚C for 45 minutes. Placental villi cell suspensions were diluted in 1X

Hank's Balanced Salt Solution (HBSS) containing 2% FBS, pelleted at 1200rpm for 5 minutes, and

resuspended in 0.25% trypsin-EDTA. The solution was diluted again in 1X HBSS containing 2% FBS,

pelleted at 1200rpm for 5 minutes, and resuspended in dispase and 1mg/ml DNase I. 1X Hank's

Balanced Salt Solution (HBSS) containing 2% FBS was added once more and the solution was sieved

through a 40μm filter.

Trophoblast organoids: For all organoid cultures, media was aspirated and organoids were

diluted in Cell Recovery Solution at 4°C for 40 minutes on either culture day 0 in complete media (for

non-differentiated organoids) or culture day 7 in differentiation media (for EVT-differentiated

organoids). Cells were then washed with cold PBS and centrifuged at 1000rpm for 5 minutes. Cells

were disaggregated in TrypLE containing 5% DNase I at 37°C for 7 minutes before the addition of

100μl complete media. The resulting cell suspension was then sieved through a 40μm filter.

Library construction: Single cell suspensions (tissue- and organoid-derived) were stained with

7-Aminoactinomycin D (7-AAD) viability marker and live/dead sorted using a Becton Dickinson (BD)

FACS Aria. Cell suspensions were loaded onto the 10X Genomics single cell controller for library

preparation. Placental villi-derived single cell suspensions were prepared using the chromium single

cell 3' reagent v2 chemistry kit, following standard manufacturer protocol for all steps. Non-

differentiated and differentiated trophoblast organoid-derived single cell suspensions were prepared

using the chromium single cell 3' reagent v3 chemistry kit, following standard manufacturer protocol

for all steps. Single cell libraries were sequenced on an Illumina Nextseq 500 instrument at a

sequencing depth of 50,000 reads/cell for tissue-derived cells and 150,000 reads/cell for organoid-

derived cells. For placental villus samples, the 10X Genomics Cell Ranger version 2.0 software was

used for STAR read alignment and counting against the provided GRCh38 reference while the

trophoblast organoid samples used the 10X Genomics Cell Ranger version 3.0 software for STAR read

alignment and counting against the GRCh38 reference.

Data Repository Integration: Droplet-based first trimester scRNA-seq decidual and placental

tissue data was obtained from the public repository ArrayExpress (available at: E-MTAB-6701[18]). Raw

BAM files were downloaded and pre-processed using 10X Genomics Cell Ranger version 2.0 under

identical parameters as previously described for our placental villi and trophoblast organoid samples

using the GRCh38 reference. Sample specific metrics are summarized in Supplemental Table 2.

Single cell RNA-seq data analysis

Data pre-processing and quality control: A total of 50,790 single cells were sequenced and pre-

processed using the Seurat R package[52,53]. To ensure quality cells were used in downstream analyses,

cells containing fewer than 1000 detected genes and greater than 20% mitochondrial DNA content

were removed. Individual samples were scored based on expression of G2/M and S phase cell cycle

gene markers, scaled, normalized, and contaminating cell doublets were removed using the

DoubletFinder package[54] with default parameters and a doublet formation rate estimated from the

number of captured cells. Following pre-processing, all samples were merged and integrated using cell

pairwise correspondences between single cells across sequencing batches. During integration, the

Pearson residuals obtained from the default regularized negative binomial model were re-computed and

scaled to remove latent variation in percent mitochondrial DNA content as well as latent variation

resulting from the difference between the G2M and S phase cell cycle scores.

Cell clustering, identification, and trophoblast sub-setting: Cell clustering was accomplished in

Seurat at a resolution of 0.90 using 50 principal components. Cell identity was determined through

observation of known gene markers found within the maternal fetal interface (vimentin, CD45, neural

cell adhesion molecule, CD14, etc.) and identifying cluster marker genes via the "FindAllMarkers"

function using a model-based analysis of single-cell transcriptomics (MAST[55]) GLM-framework

implemented in Seurat. Gene markers were selected as genes with a minimum log fold change value >

0.20 and an adjusted p-value < 0.05 within a specific cluster. Trophoblasts were identified as

expressing *KRT7, EGFR, TEAD4, TP63, TFAP2A, TFAP2C, GATA2, GATA3, HLA-G, ERVFRD-1*

transcripts at a level greater than zero, and *VIM, WARS, PTPRC, DCN, CD34, CD14, CD86, CD163,*

NKG7, KLRD1, HLA-DPA1, HLA-DPB1, HLA-DRA, HLA-DRB1, HLA-DRB5, HLA-DQA1 transcripts

at a level equal to zero. Trophoblasts were subset and re-clustered in Seurat at a resolution of 0.375

using 30 principal components. As mentioned previously, trophoblast cluster identity was determined

through observation of known trophoblast gene marker expression, proliferative gene marker

expression, and from unique cluster genes identified by differential expression analysis in Seurat.

Trophoblast projections were visualized using Uniform Manifold Approximation and Projections

(UMAPs).

Differential gene expression: Differential expression (DEG) analyses were performed in Seurat

using the "FindMarkers" function using a MAST framework on the raw count data. Parameters were

limited to genes with a log fold-change difference between groups of 0.20 and genes detected in a

minimum 10% cells in each group. Results were filtered for positive gene markers and significant

DEGs were taken as having a fold change difference > 0.25 and an adjusted p value < 0.05.

Gene set enrichment analysis: GSEA analyses were performed in Seurat using the

"FindMarkers" function using a MAST framework on the raw count data. Parameters included a

minimum log fold-change difference between groups of "−INF", a minimum gene detection of "−INF"

in cells in each group, and a minimum difference in the fraction of detected genes within each group of

"−INF", allowing all genes to be compared. Results were visualized using the R package

EnhancedVolcano (version 1.8.0), with a fold-change cut-off of 0.25 and a p value cut off of 0.00005

taken as the thresholds for significance.

Gene ontology: GO analysis of significant differentially expressed genes was performed using

the clusterProfiler R package[56] and the top enriched terms (Benjamini-Hochberg corrected p value

<0.05) were visualized as dot plots using the R package enrich plot (version 1.10.2).

CTB gene regulatory network establishment: Placental CTBs were subset and state-specific CTB gene

markers were identified as described in the *Differential gene expression* section, above. Results were

exported as an excel file for import into Cytoscape (version 3.7.1). State-specific, as well as aggregated

general CTB gene marker lists were imported as networks and the iRegulon package[59] was applied

using default parameters to define CTB gene regulatory networks.

Pseudotime: The Monocle2 R package[58-60] was used to explore the differentiation of progenitor

CTBs into specialized SCT and EVT. The trophoblast cell raw counts were loaded into Monocle2 for

semi-supervised single cell ordering in pseudotime, taking increased *BCAM* expression to represent

cells at the origin and *HLA-G* and *ERVFRD-1* gene expression to indicate progress towards EVT and

SCT, respectively. Branched expression analysis modelling (BEAM) was applied to observe the

expression of identified trophoblast cluster gene markers and genes related to the WNT, HIPPO,

NOTCH, and EGFR signaling pathways over pseudotime, progressing towards either EVT or SCT.

Results were visualized using the "plot_cell_trajectory" function. To resolve cytotrophoblast

differentiation trajectories at higher resolution, the Monocle3 package[61] was utilized. Trophoblast raw

counts were input, clustered, and the principal graph was created using the "learn_graph" function. The

"order_cells" function was then used to order cells in pseudotime, designating the trophoblast root

within the CTB 2 state.

Lineage Trajectory: The Velocyto package[62] was run in terminal using the command line

interface to generate sample-specific files containing count matrices of unspliced, spliced, and

ambiguous RNA transcript abundances. The velocyto.R implementation of the package was then used

to merge each file with its corresponding 10X sample during data pre-processing in Seurat. The final,

processed trophoblast object and UMAP embeddings were exported to a Jupyter Notebook and the

scVelo Python package[63] was used to read, prepare, and recover holistic trophoblast gene splicing

kinetics. Velocities were projected on top of the extracted UMAP embeddings generated in Seurat.

Signaling Entropy: The SCENT R package[64] was applied using default parameters and the

CCAT method for potency estimation to quantify the degree of differentiation potency and gene

expression promiscuity across single cells.

Flow cytometry analysis and cell sorting

Organoid single cell work-flow: Single cells from CT29 organoids were isolated by incubation with Cell Recovery Solution (Corning) for 40 minutes at 4°C. Then, cells were washed in ice-cold PBS and dissociated into single cells using TrypLE™ supplemented with 5% DNAse I for 7 minutes at 37°C.

Phenotyping EVT differentiated CT27, CT29, and CT30 organoids: Following single cell dissociation, organoid cells were stained with EGFR BV605 (clone AY13; 1:50; Biolegend), HLA-G APC (clone MEMG-9; 1:50; Abcam), HLA-ABC Pacific Blue (clone W6/32; 1:100; Biolegend), CD49e/ITGA5 PeCy7 (clone eBioGoH3; 1:200; Thermo Fisher Scientific), CD49f/ITGA6 Alexa 488 (clone P1D6; 1:200; Thermo Fisher Scientific), and BCAM BV711 (clone B64; 1:50; BD Biosciences).

Phenotyping BCAM^{hi} and BCAM^{lo} organoids: Single cells were incubated with anti-human BCAM BV421 (clone B64; 1:50; BD Biosciences), EGFR BV605, and CD49f/ITGA6 PeCy7 for 30 minutes in the dark on ice, followed by viability staining with 7-AAD. Samples were analyzed using LSRII (Becton Dickinson) and a minimum of 20,000 events were acquired.

hTSC BCAM$^{hi/lo}$ sorting: CT29 hTSCs were dissociated using TrypLE™ (Thermo Fisher Scientific) with 5% DNAse I (Sigma Aldrich) for 7 minutes at 37°C. Following resuspension in PBS supplemented with Antibiotic-Antimycotic (Gibco) and 1% P/S (Sigma Aldrich), cells were stained with BCAM BV421 (clone B64; 1:50; BD Biosciences) and EGFR Pecy7 (clone AY13; 1:50; Biolegend) for 30 minutes at 4°C in the dark, followed by incubation with 7-AAD (Thermo Fisher) for viability assessment. CT29 BCAM^{hi} and BCAM^{lo} populations were sorted using a FACS Aria (BD Biosciences).

Trophoblast organoid growth assay

CT29 sorted BCAM^{hi} and BCAM^{lo} cells were seeded at a density of 5,000 cells in 40μL of Matrigel in the presence of trophoblast organoid media. Phase contrast images were taken with AxioObserver inverted microscope (Carl Zeiss) using 20X objective, and organoids were further analyzed by qPCR, immunofluorescence, and flow cytometry. Organoid formation efficiency was assessed by measuring organoid number and size (area in μm^2). Images were taken daily and analyzed using imageJ (NIH). The scale was set by measuring the scale bar in pixels (285.5 pixels = 200μm).

RNA isolation

Prior to RNA extraction, organoids were dissociated into single cells as described above. Total RNA was prepared from these cells using RiboZol reagent (VWR Life Science) followed by RNeasy Mini kit (Qiagen) according to manufacturer's protocol. RNA purity was confirmed using a NanoDrop Spectrophotometer (Thermo Fisher Scientific).

cDNA synthesis and qPCR

RNA was reverse transcribed using qScript cDNA SuperMix synthesis kit (Quanta Biosciences) and subjected to qPCR (ΔΔCT) analysis, using PowerUP SYBR Green Master Mix (Thermo Fisher Scientific) on a StepOnePlus Real-Time PCR System (Thermo Fisher Scientific) and forward and reverse primer sets for *TEAD4* (F: 5'-CGGCACCATTACCTCCAACG-3'; R: 5'-CTGCGTCATTGTCGATGGGC-3'), *ITGA6* (F: 5'-CACTCGGGAGGACAACGTGA-3'; R: 5'-ACAGCCCTCCCGTTCTGTTG-3'), *CGB* (F: 5'-GCCTCATCCTTGGCGCTAGA-3'; R: 5'-TATACCTCGGGGTTGTGGGG-3'), *CSH1* (F: 5'-GACTGGGCAGATCCTCAAGCA-3'; R: 5'-CCATGCGCAGGAATGTCTCG-3'), *ITGA5* (F: 5'-GGGTCGGGGGCTTCAACTTA-3'; R: 5'-ACACTGACCCCGTCTGTTCC-3'), *HLA-G* (F: 5'-CCCACGCACAGACTGACAGA-3'; R: 5'-

AGGTCGCAGCCAATCATCCA-3'), *NOTCH1* (F: 5'-GCCTGAATGGCGGGAAGTGT-3'; R: 5'-

TAGTCTGCCACGCCTCTGC-3'), *NOTCH2* (F: 5'-ACTCGGGGCCTACTCTGTGA-3'; R: 5'-

TGTCTCCCTCACAACGCTCC-3'), *GAPDH* (F: 5'-AGGGCTGCTTTTAACTCTGGT-3' R: 5'-

CCCCACTTGATTTTGGAGGGA-3'). All raw data were analyzed using StepOne Software v2.3

(Thermo Fisher Scientific). The threshold cycle values were used to calculate relative gene expression

levels. Values were normalized to endogenous *GAPDH* transcripts.

Immunofluorescence microscopy

Placental villi (6-12 weeks' gestation; n=2) were fixed in 2% paraformaldehyde overnight at 4

°C. Tissues were paraffin embedded and sectioned at 6 μm onto glass slides. Immunofluorescence was

performed as previously described (Aghababaei et al., 2015). Briefly, organoids or placental tissues

underwent antigen retrieval by heating slides in a microwave for 5 X 2 minute intervals in citrate buffer

(pH 6.0). Sections were incubated with sodium borohydride for 5 minutes at room temperature (RT),

followed by Triton X-100 permeabilization for 5 minutes, RT. Slides were blocked in 5% normal goat

serum/0.1% saponin for 1hr, RT, and incubated with combinations of the indicated antibodies

overnight at 4 °C: mouse monoclonal anti-HLA-G (clone 4H84; 1:100; Exbio, Vestec, Czech

Republic); rabbit monoclonal anti-cytokeratin 7 (clone SP52; 1:50; Ventana Medical Systems), rabbit

polyclonal anti-BCAM (1:100; Abcam); mouse monoclonal anti-hCG beta (clone 5H4-E2; 1:100;

Abcam). Following overnight incubation, sections and coverslips were washed with PBS and incubated

with Alexa Fluor goat anti-rabbit 568 and goat anti-mouse 488 conjugated secondary antibodies (Life

Technologies, Carlsbad, CA) for 1h at RT, washed in PBS and mounted using ProLong Gold mounting

media containing DAPI (Life Technologies).

Slides were imaged with an AxioObserver inverted microscope (Car Zeiss, Jena, Germany)

using 20X Plan-Apochromat/0.80NA or 40X Plan-Apochromat oil/1.4NA objectives (Carl Zeiss). An

ApoTome .2 structured illumination device (Carl Zeiss) set at full Z-stack mode and 5 phase images

was used for image acquisition.

Statistical analysis

Data are reported as median values with standard deviations. All calculations were carried out

using GraphPad Prism software (San Diego, CA). For single comparisons, Mann-Whitney non-

parametric unpaired t-tests were performed. For multiple comparisons, one-way Kruskal-Wallis

ANOVA followed by Dunn's multiple comparison test was performed. One-way ANOVA followed by

Tukey post-test were performed for all other multiple comparisons. The differences were accepted as

significant at $P < 0.05$. For scRNA-seq statistical analyses, please refer to scRNA-seq analysis sections

in methods.

DATA AVAILABILITY/ACCESSION NUMBER

The ArrayExpress accession number for the publicly available data reported in this paper is: E-MTAB-6701.

AUTHOR CONTRIBUTIONS

AGB designed the research. AGB, MS, JB, BC, JW, SY, JT, and HTL performed experiments and analysed data. AGB, MS, JB, and BC wrote the paper. All authors read and approved the manuscript.

FUNDING

This work was supported by a Natural Sciences and Engineering Research Council of Canada (NSERC; RGPIN-2014-04466) Discovery Grant (to AGB), Canadian Institutes of Health Research 201809PJT-407571-CIA-CAAA) operating grants to AGB, and a CIHR Master's graduate studentship (to MS).

ACKNOWLEDGEMENTS

The authors extend their sincere gratitude to the hard work of staff at British Columbia's Women's Hospital's CARE Program for recruiting participants to our study and the staff at the University of British Columbia Biomedical Research Centre for assistance with all drop-seq library preparation and sequencing.

COMPETING INTERESTS

The authors declare that no competing interests exist.

REFERENCES

1. Lee CQE, et al. What is a trophoblast? A combination of criteria define human first-trimester
trophoblast. *Stem Cell Reports*. 2016. 6(2): 257-272. Doi: 10.1016/j.stemcr.2016.01.006.

2. Chang CW, et al. Trophoblast lineage specification, differentiation, and their regulation by oxygen
tension. *J Endocrinol*. 2018. 236(1): R43-56. Doi: https://doi.org/10.1530/JOE-17-0402.

3. Pjinenborg R, et al. Deep placentation. *Best Pract Res Clin Obstet Gynaecol*. 2011. 25(3): 273-285.
Doi: 10.1016/j.bpobgyn.2010.10.009.

4. Burton GJ, et al. Review Histotrophic nutrition and the placental-endometrial dialogue during
human early pregnancy. *Placenta*. 2020 102: 21-26. Doi: 10.1016/j.placenta.2020.02.008.

5. Tilburgs T, et al. Human HLA-G+ extravillous trophoblasts: Immune-activating cells that interact
with decidual leukocytes. *Proc Natl Acad Sci USA*. 2015. 112(23): 7219-7224. Doi:
10.1073/pnas.1507977112.

6. Turco MY and Moffett A. Development of the human placenta. *Development*. 2019. 146(22):
dev163428. Doi: 10.1242/dev.163428.

7. Fowden AL, et al. Review: Endocrine regulation of placental phenotype. *Placenta*. 2015. 36 Suppl
1: S50-59. Doi: 10.1016/j.placenta.2014.11.018.

8. Geusens N, et al. Endovascular trophoblast invasion, spiral artery remodelling and uteroplacental
haemodynamics in a transgenic rat model of pre-eclampsia. *Placenta*. 2008. 29(7): 614-623. Doi:
10.1016/j.placenta.2008.04.005.

9. Knöfler M, et al. Human placenta and trophoblast development: key molecular mechanisms and
model systems. *Cell Mol Life Sci*. 2019. 76(18): 3479-3496. Doi: 10.1007/s00018-019-03104-6.

10. Okae, et al. Derivation of Human Trophoblast Stem Cells. *Cell Stem Cell*. 2018. 22(1): 50-63.e6.
Doi: 10.1016/j.stem.2017.11.004.

11. Haider S, et al. Self-Renewing Trophoblast Organoids Recapitulate the Developmental Program of
the Early Human Placenta. *Stem Cell Reports*. 2018. 11(2): 537-551. Doi:
10.1016/j.stemcr.2018.07.004.

12. Turco MY, et al. Trophoblast organoids as a model for maternal-fetal interactions during human
placentation. *Nature*. 2018. 564(7735): 263-267. Doi: 10.1038/s41586-018-0753-3.

13. Lee CQE, et al. Integrin α2 marks a niche of trophoblast progenitor cells in first trimester human
placenta. *Development*. 2018. 145(16): dev162305. Doi: 10.1242/dev.162305.

14. Pollheimer J, et al. Regulation of Placental Extravillous Trophoblasts by the Maternal Uterine
Environment. *Front Immunol*. 2018. 9: 2597. Doi: 10.3389/fimmu.2018.02597.

15. Papuchova H, et al. Three types of HLA-G+ extravillous trophoblasts that have distinct immune

regulatory properties. *Proc Natl Acad Sci USA*. 2020. 117(27): 15772-15777. Doi:

10.1073/pnas.2000484117.

16. Moffett A, et al. Maternal allo-recognition of the fetus. *Fertil Steril*. 2017 107(6): 1269-1272. Doi:

10.1016/j.fertnstert.2017.05.001.

17. Jauniaux E, et al. Placental Implantation Disorders. *Obstet Gynecol Clin North Am*. 2020. 47(1):

117-132. Doi: 10.1016/j.ogc.2019.10.002.

18. Vento-Tormo R, et al. Single-cell reconstruction of the early maternal-fetal interface in humans.

Nature. 2018. 563 (7731): 347-353. Doi: 10.1038/s41586-018-0698-6.

19. Haider S, et al. Notch1 controls development of the extravillous trophoblast lineage in the human

placenta. *Proc Natl Acad Sci USA*. 2016. 113(48): E7710-E7719. Doi: 10.1073/pnas.1612335113.

20. Zhou Y, et al. Human cytotrophoblasts adopt a vascular phenotype as they differentiate. A strategy

for successful endovascular invasion? *J Clin Invest*. 1997. 99(9): 2139-2151. Doi:

10.1172/JCI119387.

21. Haider S, et al. Notch signaling plays a critical role in motility and differentiation of human first-

trimester cytotrophoblasts. *Endocrinology*. 2014. 155(1): 263-74. Doi: 10.1210/en.2013-1455.

22. Suryawanshi H, et al. A single-cell survey of the human first-trimester placenta and decidua. *Sci*

Adv. 2018. 4(10):eaau4788. Doi: 10.1126/sciadv.aau4788.

23. Liu Y, et al. Single-cell RNA-seq reveals the diversity of trophoblast subtypes and patterns of

differentiation in the human placenta. *Cell Res*. 2018. 28(8):8 19-832. Doi: 10.1038/s41422-018-

0066-y.

24. Meinhardt G, et al. Pivotal role of the transcriptional co-activator YAP in trophoblast stemness of

the developing human placenta. *Proc Natl Acad Sci USA*. 2020. 117(24): 13562-13570. Doi:

10.1073/pnas.2002630117.

25. Saha B, et al. TEAD4 ensures postimplantation development by promoting trophoblast self-

renewal: An implication in early human pregnancy loss. *Proc Natl Acad Sci USA*. 2020. 117(30):

17864-17875. Doi: 10.1073/pnas.2002449117.

26. Del Valle-Pérez B, et al. Wnt controls the transcriptional activity of Kaiso through CK1ε-dependent

phosphorylation of p120-catenin. *J Cell Sci*. 2016. 129(4): 873. Doi: 10.1242/jcs.186288.

27. Nemeth MJ, et al. Hmgb3 regulates the balance between hematopoietic stem cell self-renewal and

differentiation. *Proc Natl Acad Sci USA*. 2006. 103(37): 13783-8. Doi: 10.1073/pnas.0604006103.

28. Thomas G. An encore for ribosome biogenesis in the control of cell proliferation. *Nature*. 2000. 2:

E71-72.

29. Xu X, et al. The role of ribosomal proteins in the regulation of cell proliferation, tumorigenesis, and genomic integrity. *Sci China Life*. 2016. 59(7): 656-672.

30. Khajuria RK, et al. Ribosome levels selectively regulate translation and lineage commitment in human hematopoiesis. *Cell*. 2018. 173(1): Doi: 10.1016/j.cell.2018.02.036.

31. Shi Z, et al. DNA-binding mechanism of the Hippo pathway transcription factor TEAD4. *Oncogene*. 2017. 36: 4362-4369. Doi: https://doi.org/10.1038/onc.2017.24.

32. Saha B, et al. TEAD4 ensures postimplantation development by promoting trophoblast self-renewal: an implication in early human pregnancy loss. PNAS. 2020. 117(30): 17864-17875.

33. Xiao L, et al. Deciphering a distinct regulatory network of TEAD4, CDX2 and GATA3 in humans for trophoblast transition from embryonic stem cells. *Biochim Biophys Acta Mol Cell Res*. 2020. 1867(9): 118736.

34. Yagi R, et al. Transcription factor TEAD4 specifies the trophectoderm lineage at the beginning of mammalian development. *Development*. 2007. 134: 3827-3836. doi: 10.1242/dev.010223.

35. Ilicic T, et al. Classification of low quality cells from single-cell RNA-seq data. *Genome Biol*. 2016. 17(29). Doi: https://doi.org/10.1186/s13059-016-0888-1.

36. Galluzzi L, et al. Mitochondria: master regulators of danger signalling. *Nat Rev Mol Cell Biol*. 2012. 13: 780-788. Doi: https://doi.org/10.1038/nrm3479.

37. Detmer SA and Chan DC. Functions and dysfunctions of mitochondrial dynamics. *Nat Rev Mol Cell Biol*. 2007. 8: 870-879. Doi: https://doi.org/10.1038/nrm2275.

38. Yoon JC, et al. Wnt signaling regulates mitochondrial physiology and insulin sensitivity. *Genes & Dev*. 2010. 24: 1507-1518. Doi: 10.1101/gad.1924910.

39. Brown K, et al. WNT/β-catenin signaling regulates mitochondrial activity to alter the oncogenic potential of melanoma in a PTEN-dependent manner. *Oncogene*. 2017. 36(22): 3119-3136. Doi: 10.1038/onc.2016.450.

40. Jauniaux E, et al. Onset of maternal arterial blood flow and placental oxidative stress. A possible factor in human early pregnancy failure. *Am J Pathol*. 2000. 157(6): 2111-2122. Doi: 10.1016/S0002-9440(10)64849-3.

41. Jauniaux E, et al. Evaluation of respiratory gases and acid-base gradients in human fetal fluids and uteroplacental tissue between 7 and 16 weeks' gestation. *Am J ObGyn*. 2001. 184(5): 998-1003.

42. Weiss G, et al. The trophoblast plug during early pregnancy: a deeper insight. *Histochem Cell Biol*. 2016. 146: 749-756. Doi: https://doi.org/10.1007/s00418-016-1474-z.

43. Tuuli MG, et al. Review: oxygen and trophoblast biology—a source of controversy. Placenta. 2011.
32(2): S109-S118. doi: 10.1016/j.placenta.2010.12.013.

44. Chang CW, et al. Trophoblast lineage specification, differentiation, and their regulation by oxygen
tension. J Endocrinol. 2018. 236(1): R43-56. Doi: https://doi.org/10.1530/JOE-17-0402.

45. Wakeland AK, et al. Hypoxia directs human extravillous trophoblast differentiation in a hypoxia-
inducible factor-dependent manner. Am J Pathol. 2017. 187(4): 767-780. Doi:
https://doi.org/10.1016/j.ajpath.2016.11.018.

46. Treissman J, et al. Low oxygen enhances trophoblast column growth by potentiating differentiation
of the extravillous lineage and promoting LOX activity. Development. 2020. 147(2). DOI:
10.1242/dev.181263.

47. Zen Q, et al. Critical factors in basal cell adhesion molecule/Lutheran-mediated adhesion to
laminin. J Biol Chem. 1999. 274(2): 728-734. Doi: https://doi.org/10.1074/jbc.274.2.728.

48. Kikkawa Y, et al. The Lutheran/basal cell adhesion molecule promotes tumor cell migration by
modulating integrin-mediated cell attachment to laminin-511 protein. J Biol Chem. 288(43): 30990-
31001. Doi: https://doi.org/10.1074/jbc.M113.486456.

49. Kannan K et al. Recurrent BCAM-AKT2 fusion gene leads to a constitutively activated AKT2
fusion kinase in high-grade serous ovarian carcinoma. PNAS. 2015. 112(11): E1272-1277. Doi:
10.1073/pnas.1501735112.

50. Jin J, et al. Upregulation of BCAM and its sense lncRNA BAN are associated with gastric cancer
metastasis and poor prognosis. Mol Oncology. 2020. 14(4): 829-845. Doi:
https://doi.org/10.1002/1878-0261.12638.

51. Akiyama H, et al. The FBI/Akirin2 target gene, BCAM, acts as a suppressive oncogene. PLOS
One. 2013. 8(11): e78716. Doi: https://doi.org/10.1371/journal.pone.0078716.

52. Butler A, et al. Integrating single-cell transcriptomic data across different conditions, technologies,
and species. *Nature Biotechnol*. 2018. 36(5): 411-420.

53. Stuart T, et al. Comprehensive integration of single-cell data. *Cell*. 2019. 177(7): 1888-1902.

54. McGinnis CS, et al. DoubletFinder: Doublet detection in single-cell RNA sequencing data using
artificial nearest neighbours. Cell Systems. 2019. 8(4): 329-337.

55. Finak G, et al. MAST: a flexible statistical framework for assessing transcriptional changes and
characterizing heterogeneity in single-cell RNA sequencing data. *Genome Biol*. 2015. 16: 278.

56. Yu G, et al. clusterProfiler: an R package for comparing biological themes among gene clusters.
OMICS. 2012. 16(5): 284-287.

57. Janky R, et al. iRegulon: from a gene list to a gene regulatory network using large motif and track

collections. *PLoS Comput Biol*. 2014. 10(7): e1003731.

58. Trapnell C, Cacchiarella D, et al. The dynamics and regulators of cell fate decisions are revealed by

pseudo-temporal ordering of single cells. *Nature Biotechnol*. 2014. 32(4): 381.

59. Qiu X, Hill A, Trapnell C, et al. Single-cell mRNA quantification and differential analysis with

Census. *Nat Methods*. 2017. 14(3): 309-315.

60. Qiu X, Trapnell C, et al. Reverse graph embedding resolves complex single-cell developmental

trajectories. *Nat Methods*. 2017. 14(10): 979-982.

61. Cao J, et al. The single-cell transcriptional landscape of mammalian organogenesis. *Nature*. 2019.

566(7745): 496-502. Doi: 10.1038/s41586-019-0969-x.

62. La Manno G, et al. RNA velocity of single cells. *Nature*. 2018. 560: 494-498.

63. Bergen V, et al. Generalizing RNA velocity to transient cell states through dynamical modelling.

Nature Biotechnol. 2020: https://doi.org/10.1038/s41587-020-0591-3

64. Teschendorff A, Enver T. Single-cell entropy for accurate estimation of differentiation potency

from a cell's transcriptome. *Nat Commun*. 2017. 8(1): 1-15.

TITLES AND LEGENDS TO FIGURES

Figure 1: Single cell characterization of first trimester trophoblasts

(A) Schematic of a first trimester chorionic villous including anatomical representations of both an anchoring villous and floating villous. Highlighted are trophoblast sub-populations and the villous and extravillous cell differentiation trajectories: cytotrophoblast (CTB), column cytotrophoblast (cCTB), syncytiotrophoblast (SCT), and invasive extravillous trophoblast (EVT).

(B) Summary of workflow used for collection, processing, and single cell analysis of placental villous cells. MFI, Maternal-Fetal Interface, 7AAD: 7-Aminoactinomycin D, FSC-A: Forward Scatter Area.

(C) Uniform Manifold Approximation and Projection (UMAP) plot of 7798 captured trophoblasts, demonstrating 8 distinct trophoblast clusters: CTB1-4, cCTB1-2, SCTp, and EVT.

(D) Heatmap showing the average expression of known trophoblast and proliferation gene markers in each trophoblast cluster. Cell cluster names and colours correspond to those in Figure 1C.

(E) Stacked bar plot showing cell state distribution in distinct phases of the cell cycle (G1, G2M, S).

(F) UMAP plots showing expression (red) of CTB progenitor (*EGFR*, *TP63*), proliferation (*MKI67*), EVT (*HLA-G*), SCT (*ERVFRD-1*), and an EVT progenitor (*ITGA2*) gene(s).

(G) Proportion of 4687 early first trimester trophoblasts in each scRNA-seq-defined state in early (6-7 weeks) and late (9-12 weeks) gestational time-points.

Figure 1

918   **Figure 2: Single cell transcriptomic resolution of distinct CTB states**
919

(A) UMAP plot of 5477 captured CTBs highlighting 4 distinct cell states (CTB1-4).
(B) Heatmap showing the average expression of the top-10 most highly-expressed (FDR-ranked) genes
in CTB states 1-4.
(C) Venn diagram comparing the intersection of genes upregulated within each CTB1-4 states.
(D) Dot plots of the top significant terms enriched by gene ontology analysis informed by gene
signatures enriched within distinct CTB (CTB1-4) states.
(E) Heatmap showing the average expression of transcription factors identified as putative
regulators/drivers enriched within CTB1-4 states.
928

Figure 2

Figure 3: scVelo-informed cell differentiation identifies a primitive CTB origin.

(A) UMAP plot overlain with projected RNA velocity (scVelo) vector field, indicating trophoblast directionality (represented by the velocity arrows) away from a putative origin towards endpoint SCTp and EVT states. Cell cluster names and colours correspond to those in Figure 1C.

(B) UMAP plots showing gene expression (red) of progenitor- and stem cell-associated genes in all scRNA-seq-informed trophoblast states.

(C) Venn diagram comparing the genes upregulated within CTB states 2 and 3. Highlighted are genes related to WNT and EGFR signaling pathways, genes involved in extracellular matrix (ECM) interactions, and miscellaneous (Misc) genes of interest.

(D) Scatter plots showing gene expression levels of EGFR related genes (*EFEMP1*), transcription factors (*EGR1* and *ELF5*), and genes involved in ECM-cell interactions (*ITGA6*, *BCAM*, and *LAMA5*) across pseudotime as the CTB2 state transitions to the CTB3 state.

(E) UMAP plots showing gene expression (red) of genes upregulated within the CTB2 state origin.

Figure 3

Figure 4: Single cell transcriptomic analyses in hTSC organoids recapitulate a primitive CTB origin

(A) Bright field microscopy images of regenerative (Wnt+) and EVT-differentiated (Wnt-) hTSC-derived organoids following 10 days of culture. Bar = 400 μm.

(B) Flow cytometry plots showing cell frequencies of HLA-G, CD49e, and HLA-ABC positive cells in regenerative (Wnt+) and EVT-differentiated hTSC-derived (CT27) organoids.

(C) Relative measurement of mRNA levels by qPCR in EVT-differentiated hTSC-derived organoids (CT27, CT29, CT30) over 21 days (D) of culture. Shown are CTB/progenitor (*TEAD4, ITGA6*), syncytial (*CGB, CSH1*), immature EVT (*ITGA5, NOTCH1*), and mature EVT (*NOTCH2, HLA-G*) genes.

(D) UMAP and gene heatmap plots of combined CT27, CT29, and CT30 single cell hTSCs following culture in regenerative (Wnt+) and EVT-differentiated (Wnt-) conditions after 7 days in culture. UMAP plots are overlain with projected RNA velocity (scVelo) vector field, demonstrating 7 distinct trophoblast clusters representing 3 states (CTB1, CTB2, CTB3, bipotent cCTB, EVT1, EVT2, SCTp). Heatmap shows the average expression of trophoblast- and proliferation-associated genes in each cell state.

(E) UMAP and gene heatmap plots derived from individual hTSC lines (CT27, 2259 cells; CT29, 1446 cells; CT30, 2604 cells) subjected to scRNA-seq analyses. UMAP plots are overlain with projected RNA velocity vector field, demonstrating 5 distinct trophoblast clusters in each hTSC line.

(F) Venn diagram comparing genes shared and distinct between CTB2/CTB3 in chorionic villi and CTB1/CTB2 states in hTSC organoids.

UMAP plots showing expression of BCAM (red) in **(G)** combined (CT27, CT29, CT30) and **(H)** segregated hTSC-derived organoid cells/states.

(G) UMAP plot of all 7 cell states in hTSC-derived organoids (CT27, CT29, CT30) showing *BCAM* expression (red signal).

(H) UMAP plots of cell states of individual hTSC-derived organoids (CT27, CT29, CT30) showing *BCAM* expression (red signal).

Figure 4

A CT29 (10 days)
Wnt+
Wnt−
400μm

B CT27 (10 days)

Wnt+
HLA-G+ 2.09
HLA-ABC+ 0.75

Wnt−
HLA-G+ 28.2
HLA-ABC+ 17.8

HLA-G / CD49e
HLA-ABC / HLA-G

C
CTB/progenitor genes — TEAD4, ITGA6
Syncytial genes — CGB, CSH1
Immature EVT genes — ITGA5, NOTCH1
Mature EVT genes — NOTCH2, HLA-G

D0, D10, D21

CT27 CT29 CT30

D hTSC-Derived Organoids
CT27, CT29, CT30

E CT27 | CT29 | CT30

Genes: ERVFRD−1, ERVV−1, SDC1, TFAP2C, TFAP2A, GATA3, EGFR, GATA2, KRT7, HLA−G, ITGA5, NOTCH2, ITGA1, BMP4, NOTCH1, SOX9, ITGA2, YAP1, TEAD4, TP63, ITGA6, BCAM, PAGE4, ELF5, NANOG, MKI67, CCNA2

UMAP 2 / UMAP 1

● CTB 1 ● CTB 2 ● CTB 3 ● Bipotent cCTB ● EVT 1 ● EVT 2 ● SCTp

F
Chorionic villi CTB2, CTB3 | Organoid CTB1, CTB2

YAP1 PEG10 SPINT1 PEG3 MSI2 EFEMP1

ITGB5 | LAMA1

EGFR TFAP2C SOD3 ESRRA

1020 | 398 | 1014

MKI67 EPCAM TEAD4

BCAM LAMA5 FBN2 LAMB1 LAMC1 ITGB4 ITGA6

G BCAM
hTSC (CT27, CT29, CT30)

H BCAM
CT27
CT29
CT30

Figure 5: BCAM enrichment establishes a regenerative/growth advantage in trophoblast organoids

(A) Immunofluorescence (IF) images of BCAM and keratin-7 (KRT7), human chorionic gonadotrophin (hCG), and HLA-G in early (6.3 weeks) and late (11.2 weeks) first trimester placental villi. Nuclei are stained with DAPI (white). Bars = 100μm.

(B) IF images of BCAM, hCG and Ki67 in hTSC-derived organoid (CT29) cultured in regenerative (Wnt+) media for 7 days.

(C) Cell sorting workflow for isolating BCAM^{hi} and BCAM^{lo} EGFR+ CTBs in hTSC (CT29)-derived organoids.

(D) Bright field images of hTSC (CT29) BCAM^{hi} and BCAM^{lo} –derived organoids cultured over 11 days in regenerative Wnt+ media. Bars = 200μm.

(E) Colony area (μm^2) of BCAM^{hi} and BCAM^{lo} hTSC (CT29)-derived organoids cultured over 11 days (D).

(F) Flow cytometric panels showing expression of BCAM and EGFR in BCAM^{hi} and BCAM^{lo}, CT29-derived organoids following 11 days of culture.

(G) Heatmap showing levels of select differentially expressed genes from CTB expressing high (hi) or low (lo) levels of *BCAM* in the CTB1 state of CT29-derived organoids following 7 days of culture.

Figure 5

Supplemental Figure 1: Sample quality control and single cell characterization of the first
trimester human maternal-fetal interface

(A – C) Violin plots showing the distribution of the raw number of features **(A)**, the raw number of counts **(B)**, and the percent mitochondrial DNA content remaining post quality control and data pre-processing **(C)** captured per cell during sequencing for each sample. Samples 1-6 (teal) represent the placental tissue samples sequenced by the Beristain lab, samples 7-11 (red) represent the placental tissue samples acquired from ArrayExpress, and samples 12-15 (grey) represent the decidual tissue samples acquired from ArrayExpress.

(D) Scatter plot showing the distribution of captured cells across principal components 1 and 2 following dimensional reduction in principal component space. Cell positions are calculated based on cell embeddings calculated during dimensional reduction by principal component analysis. Cells are colour coordinated by source with teal representing those cells originating from placental tissue samples sequenced by the Beristain lab and red representing those cells originating from placental and decidual tissue samples acquired from ArrayExpress.

(E) UMAP plot of 50,790 captured cells, demonstrating 31 distinct clusters identified within the first trimester human maternal fetal interface. Dashed lines separate maternal decidual immune cells from non-immune cells and fetal placental trophoblasts. Cells are colour coordinated by identity and abbreviations are identical to those in Figure 1A. Epigland: epiglandular cell, fetal Endo: fetal endothelial cell, dEndo: decidual endothelial cell, dSC: decidual stromal cell PV: perivascular cell, pFib: placental fibroblast, Hb: fetal Hofbauer cell, M1: macrophage cell 1, M2: macrophage cell 2, uNK: uterine natural killer cell.

(F) UMAP projection of the captured maternal fetal interface cells, stratified by gestational age. Early (left panel): 9803 early first trimester cells from each identified cluster captured between 6-7 weeks gestational age, before the onset of maternal placental vascularization. Late (right panel): 40,987 late first trimester cells from each identified cluster captured between 9-12 weeks gestational age, after the onset of maternal placental vascularization. Cell cluster names and colours correspond to those in Supplemental Figure 1E.

(G) Heatmap showing the average expression of known immune, non-immune, and trophoblast gene markers, as well as proliferation gene markers in each identified cluster. Cell cluster names and colours correspond to those in Supplemental Figure 1E.

(H) Stacked bar plot showing the percent distribution of cells in each cluster with gene signatures representing a cell in the gap 1 phase of proliferation, the gap 2 or mitotic phases of proliferation, or the synthesis phase of proliferation. Cell cluster names and colours correspond to those in Supplemental Figure 1E and abbreviations are identical to those in Figure 1F.

(I) UMAP plots showing gene expression in red of a mesenchymal cell gene marker (*VIM*), an immune cell gene marker (*PTPRC*, encoding the CD45 protein), endothelial cell gene markers (*CD34* and *VCAM1*), and a fibroblast gene marker (*ACTA2*).

(J) UMAP plots showing gene expression in red of known trophoblast gene markers used to subset and purify first trimester human trophoblasts from other cells within the maternal fetal interface.

1033 **Supplemental Figure 2: Branched pseudotime ordering aligns key gene pathways to the villous**
1034 **and extravillous differentiation trajectories**
1035
(A) Pseudotime trajectory plot showing trophoblasts ordered along the branched trajectory. Cells are
colour coordinated by pseudotime value, with pseudotime increasing as cells move away from the root
along the trajectory.
(B) Cells are colour coordinated by cluster identity, with progress towards SCT indicated with a dashed
arrow and progress towards EVT indicated with a solid arrow, moving away from the indicated
trajectory root. Cell cluster names and colours correspond to those in Figure 1C.
(C-D) Branched pseudotime heatmap showing relative gene expression changes and average cluster
gene expression levels, respectively, of genes involved in the WNT signaling pathway, the HIPPO
signaling pathway, the NOTCH signaling pathway, and the EGFR signaling pathway.
1045
1046
1047 **Supplemental Figure 3: Monocle3 pseudotime quantification supports identification of CTB2 as**
1048 **the progenitor origin**
1049
(A) UMAP plot with constructed trajectory graph overlain. Cells are colour coordinated by relative
pseudotime value.
(B) UMAP plot with constructed trajectory graph overlain. Cells are colour coordinated by cluster
identity. Cell cluster names and colours correspond to those in Figure 1C.
(C) UMAP plot highlighting the constructed trajectory graph which describes the path taken as CTB
state 2 progresses into CTB state 3. Cells are colour coordinated by cluster. Cluster names and colours
correspond to those in Figure 1C.
1057
1058 **Supplemental Figure 4: Organoid colonies in BCAM^{hi} and BCAM^{lo} sorted organoids**
1059 Bar graph shows the average of numbers of organoid colonies from BCAM^{hi} and BCAM^{lo} –sorted
1060 CT29 organoids cultured over 11 days. Grey bars = BCAM^{lo}; Green bars = BCAM^{hi}.

Supplemental Figure 1

Supplemental Figure 2

Supplemental Figure 3

Supplemental Figure 4