1 *Major subject area:* Biochemistry and Chemical Biology, Cell Biology

3 Edge-strand of BepA interacts with immature LptD on the β-barrel assembly

4 machine to direct it to on- and off-pathways

6 Ryoji Miyazaki[1, 2], Tetsuro Watanabe[1], Kohei Yoshitani[1], Yoshinori Akiyama[1, *]

7 [1]Institute for Frontier Life and Medical Sciences, Kyoto University, Kyoto 606-8507, Japan

8 [2]Present address:

9 Division of Biological Science, Nara Institute of Science and Technology, Ikoma, Nara 630-0192, Japan

10 *Correspondence: yakiyama@infront.kyoto-u.ac.jp (Y. A.)

13 Abstract

The outer membrane (OM) of gram-negative bacteria functions as a selective permeability barrier. *Escherichia*

coli periplasmic Zn-metallopeptidase BepA contributes to the maintenance of OM integrity through its

involvement in the biogenesis and degradation of LptD, a β-barrel protein component of the lipopolysaccharide

translocon. BepA either promotes the maturation of LptD when it is on the normal assembly pathway (on-pathway)

or degrades it when its assembly is compromised (off-pathway). BepA performs these functions probably on the

β-barrel assembly machinery (BAM) complex. However, how BepA recognizes and directs an immature LptD to

different pathways remains unclear. Here, we explored the interactions among BepA, LptD, and the BAM complex.

We found that the interaction of the BepA edge-strand located adjacent to the active site with LptD was crucial

not only for proteolysis but also, unexpectedly, for assembly promotion of LptD. Site-directed crosslinking

analyses indicated that the unstructured N-terminal half of the β-barrel-forming domain of an immature LptD

contacts with the BepA edge-strand. Furthermore, the C-terminal region of the β-barrel-forming domain of the

BepA-bound LptD intermediate interacted with a "seam" strand of BamA, suggesting that BepA recognized LptD

assembling on the BAM complex. Our findings provide important insights into the functional mechanism of BepA.

Introduction

The cell envelope of diderm bacteria is composed of two membranes, namely the inner (cytoplasmic) membrane (IM) and the outer membrane (OM). The intermembrane space, known as periplasmic space, contains a peptidoglycan layer. The OM is the outermost layer of a cell directly facing the external milieu and acts as a selective permeability barrier that prevents the penetration of toxic compounds including antibiotics (Nikaido, 2003). The cell surface localization as well as the functional impotence of the OM make its components suitable drug targets.

Outer membrane proteins (OMPs), generally exhibiting a β-barrel fold formed by more than 8 β-strands, play important roles in maintaining the structural and functional integrity of the OM (Konovalova et al., 2017). Therefore, irregularities in OMP biogenesis result in elevated cellular sensitivity to toxic compounds (Hart et al., 2019; Oh et al., 2011). After synthesis in the cytoplasm and following translocation across the IM to the periplasm through the SecYEG translocon, OMPs are delivered to the OM by periplasmic chaperones such as DegP, Skp, and SurA, and are finally integrated into the OM (Konovalova et al., 2017; Plummer and Fleming, 2016; Ricci and Silhavy, 2019). The OM assembly of OMPs is mediated by the β-barrel assembly machinery (BAM) complex consisting of a β-barrel OMP, BamA, and four lipoproteins, namely BamB, BamC, BamD, and BamE (Plummer and Fleming, 2016; Ricci and Silhavy, 2019; Tomasek and Kahne, 2021). Among the BAM complex subunits, BamA and BamD are the only essential components, although recent studies have shown that certain BamA mutations render all other BAM subunits dispensable (Hart et al., 2020). The BAM complex has a silk-hat-like structure (Bakelar et al., 2016; Gu et al., 2016; Han et al., 2016; Iadanza et al., 2016; Tomasek et al., 2020); the OM-embedded C-terminal β-barrel domain of BamA forms the "crown" and the N-terminal periplasmic polypeptide transport-associated (POTRA) domains of BamA form the "brim" together with the BamB/C/D/E lipoproteins.

Lipopolysaccharide (LPS), another major OM constituent localized in the outer leaflet of the OM, is also important for the maintenance of the structure and function of the OM (Sperandeo et al., 2017). LPS is synthesized on the cytoplasmic side of the IM and flipped across the IM to the periplasm. After maturation, it is transported to the OM by the LPS transport (Lpt) proteins (Sperandeo et al., 2017). A heterodimer of LptD, a β-barrel OMP, and LptE, a lipoprotein, plays roles in the final step to insert LPS into the OM (Dong et al., 2014; Qiao et al., 2014; Wu et al., 2006). The OM assembly process of LptD is unique in that it is accompanied by

rearrangement of intramolecular disulfide bonds. The mature form of LptD possesses two disulfide bonds formed by non-consecutive pairs of Cys residues (C31-C724 and C173-C725) (Ruiz et al., 2010). It has been shown, however, that an assembly intermediate of LptD having disulfide bonds formed by consecutive pairs of the Cys residues (C31-C173 and C724-C725) (LptD^C; LptD with **C**onsecutive disulfide bods) is first generated and isomerized to LptD^{NC} (LptD with **N**on-**C**onsecutive disulfide bods) during its assembly/maturation, which is triggered by the association of LptD with LptE (Chng et al., 2012; Narita et al., 2013). The LptD^C to LptD^{NC} conversion should occur at a later step in the OM assembly because LptD presumably associates with LptE on the BAM complex (Chimalakonda et al., 2011; Chng et al., 2012; Lee et al., 2016; Narita et al., 2013).

BepA (formally called YfgC), a bi-functional periplasmic protein that plays an important role in maintaining OM integrity (Narita et al., 2013), belongs to the M48 family zinc metallopeptidases that include prokaryotic and eukaryotic proteases (such as Ste24, Oma1, and HptX) involved in membrane quality control (Rawlings et al., 2018).We have previously shown that BepA is involved in the biogenesis and quality control of LptD. While BepA promotes the LptD^C to LptD^{NC} conversion (chaperone-like function) (Narita et al., 2013), it also degrades the stalled or misassembled LptD^C molecules that are generated due to an *lptD* mutation (*lptD4213*) or decreased availability of or weakened interaction with LptE (protease function) (Narita et al., 2013; Soltes et al., 2017). The BepA protein consists of an N-terminal M48 metallopeptidase domain and a C-terminal tetratricopeptide repeat (TPR) domain that are associated closely to form a compact structure (**Figure 1A**) (Bryant et al., 2021; Shahrizal et al., 2019). Our previous study suggested that the TPR domain of BepA directly interacts with LptD and with the BAM complex with its TPR domain inserted into the interior of the periplasmic part (brim) of the BAM complex (Daimon et al., 2017; Narita et al., 2013). A mutational study has suggested that these interactions are important for BepA functions (Shahrizal et al., 2019). Recent studies have also shown that the His-246 residue of BepA that coordinates the zinc ion at the proteolytic active site acts as an ON/OFF switch (His switch) for the proteolytic activity of BepA (Bryant et al., 2021; Daimon et al., 2020). The dual functions of BepA should be appropriately regulated because the unregulated expression of the proteolytic activity of BepA caused by an H246 mutation leads to the degradation of the normally assembling LptD intermediate (Daimon et al., 2020). However, information on the molecular mechanism of this regulation and the modes of the BepA–LptD interaction in each BepA function remains elusive.

Here, we investigated the mechanism by which BepA established interaction with LptD in promotion

of its assembly and degradation. Our results showed that a conserved β-strand (edge-strand) located adjacent to

the BepA active site directly contacts with LptD and plays important roles in substrate proteolysis, like many other

proteases. In addition, we unexpectedly found that the edge-strand-mediated interaction with a substrate is also

required for the chaperone-like function of BepA, which should be enabled by the His switch-mediated repression

of the proteolytic activity. Crosslinking experiments demonstrated that BepA could interact with an LptD molecule

assembling on the BAM complex. Based on these observations, we propose a model explaining the edge-

strand/His switch-mediated functional regulation of BepA in LptD assembly/degradation.

Results

Interaction of the BepA edge-strand with LptD is crucial not only for proteolysis but also for assembly promotion of LptD by BepA

Zinc-metallopeptidases usually possess a β-strand, called edge-strand, located close to their proteolytic active sites (Akiyama et al., 2015; López-Pelegrín et al., 2013; Stöcker and Bode, 1995). While the edge-strand is known to play a critical role in substrate proteolysis by directly interacting with a substrate polypeptide by the strand addition mechanism and converts it into an extended conformation for its presentation to the active site and proteolysis, it has not been well-characterized in M48 proteases. The solved structures of *Escherichia coli* BepA (Bryant et al., 2021; Shahrizal et al., 2019) show that it has a β-strand (β2) that is conserved among the M48-peptidases and is located adjacent to the active site (**Figure1A** and **Figure 1-figure supplement 1**), suggesting that this strand presumably acts as an edge-strand. To examine the role of the β2 strand in BepA functions, we constructed BepA mutants by introducing Pro at each position in β2 (from Asn-105 to Phe-110; **Figure 1A**). We then investigated the effects of β2 mutations on the proteolytic activity of BepA against overproduced LptD. When LptD is overproduced from a multi-copy plasmid, it mainly accumulates in the form of LptD^C possibly due to the limited availability of its partner protein, LptE (Daimon et al., 2020, 2017). This species probably represents a "normal" assembly intermediate as it is associated with the BAM complex (See below) and can be converted to the mature form (LptD^{NC}) when LptE is co-expressed (Miyazaki et al., 2018). As reported previously (Daimon et al., 2020, 2017), overproduced LptD was degraded by co-expressed wild-type BepA to generate discrete degradation products (**Figure 1B**). To examine the possible roles of the β2 strand in the function of BepA, we introduced a Pro substitution into the β2 strand, as a Pro residue would affect the secondary structure, and thus the function, of this strand. We found that the expression of a few BepA mutants (N105P, A106P, F107P, and A108P) led to a significantly decreased generation of the LptD degradation products (**Figure 1B**). These mutants accumulated at a level comparable to that of wild-type BepA, suggesting that they did not exert drastic effects on the overall structure of BepA. Furthermore, some of these mutations compromised the degradation of BamA in a Δ*surA* strain (Daimon et al., 2017) and the self-cleavage of BepA-His$_{10}$ (BepA possessing a C-terminal His$_{10}$-tag) within the His$_{10}$-tag (Narita et al., 2013) (**Figure 1-figure supplement 2A** and **2B**). These results strongly suggest that the β2 strand is important for the proteolytic activity of BepA.

We next analyzed whether the β2 strand directly interacts with a substrate by using a site-directed *in*

vivo photo-crosslinking approach (Chin and Schultz, 2002; Miyazaki et al., 2020a). We expressed derivatives of

BepA-His$_{10}$ harboring a photoreactive amino acid analog, *p*-benzoyl-L-phenylalanine (*p*BPA), at each position in

the β2 strand in a Δ*bepA* strain and examined their complementation activity regarding erythromycin (EM)

sensitivity of the cell and self-cleavage of BepA-His$_{10}$. The results showed that certain mutants (F107*p*BPA,

F109*p*BPA, and F110*p*BPA) were as functional as the wild-type, but others exhibited neither significant

complementation activity nor self-cleavage (**Figure 1-figure supplement 3A** and **3B**). For the photo-crosslinking

experiments conducted in this study, we used BepA variants harboring the E137Q mutation in the H^{136}ExxH motif

for two reasons. First, this mutation would repress the possible degradation of LptD by the BepA derivatives.

Second, our previous study has shown that this mutation stabilizes the interaction of BepA with LptD (Daimon et

al., 2017). Following UV irradiation of cells expressing each of the BepA mutants, BepA-His$_{10}$ and its crosslinked

products were purified from the membrane fractions by affinity isolation using Ni-NTA agarose and subjected to

immunoblotting analysis (**Figure 1C**). The N105*p*BPA, A106*p*BPA, F107*p*BPA, and A108*p*BPA derivatives of

BepA generated, in a UV-dependent manner, evident crosslinked products that reacted with both anti-BepA and

anti-LptD antibodies. Taken together, these results imply that the β2 strand of BepA directly interacts with LptD

and plays an important role in their degradation. The structure, intramolecular disposition, as well as involvement

in substrate interaction and proteolysis of the β2 strand strongly supports the hypothesis that it indeed acts as the

edge-strand to recognize and present a substrate to the active site. We have thus referred to the β2 strand as "the

edge-strand" hereafter. BepA N105*p*BPA and F107*p*BPA derivatives also generated the unidentified crosslinked

products (**Figure 1C, open arrowheads**), suggesting that the edge strand would interact with some other substrates.

The differential effects of the mutations on the degradation of LptD and BamA and the self-cleavage of the C-

terminal tag (**Figure 1B** and **Figure 1-figure supplement 2A** and **2B**) might reflect their different interaction

properties and/or affinities for the BepA edge-strand. Our previous study also suggested substrate-specific

interaction of an edge-strand for another metallopeptidase, RseP (Akiyama et al., 2015).

BepA not only degrades the stalled or misassembled LptD (when LptD is on the normal (on-) pathway)

but also promotes the maturation of LptD through facilitating its assembly with the partner protein LptE

(chaperone-like activity) (on the off-pathway) (Narita et al., 2013). Therefore, we examined the effects of the edge-

strand Pro mutations on the chaperone-like activity of BepA. LptD possesses two intramolecular disulfide bonds.

While the mature (fully assembled) LptD has non-consecutive disulfide bonds, an assembly intermediate form of

LptD (LptD^C) that accumulates in the absence of functional BepA possesses consecutive disulfide bonds.

Consistent with our previous results (Narita et al., 2013), expression of the wild-type BepA in a $\Delta bepA$ strain

markedly decreased the accumulation of LptD^C. On the contrary, only a partial decrease in accumulation was

observed with the expression of the E137Q mutant. We found that the A106P mutant was defective in reducing

LptD^C accumulation, suggesting that a mutation in the edge-strand could affect the chaperone-like function

(**Figure 1-figure supplement 4A**). Subsequently, we examined the effects of two edge-strand mutations—A106P

and F1107P—that markedly compromised LptD degradation on the chaperone-like function of BepA using pulse-

chase experiments. Expression of the wild-type BepA in $\Delta bepA$ cells significantly accelerated the conversion of

LptD^C to LptD^{NC}, whereas expression of the protease-dead E137Q mutant demonstrated weakened conversion

(**Figure 1D**). The acceleration of the LptD^C to LptD^{NC} conversion by the F107P mutant was weaker than that by

the wild-type BepA (almost the same or slightly stronger as compared with the E137Q mutant), and no detectable

acceleration was observed with the A106P mutant (**Figure 1D**). These results unexpectedly revealed that the edge-

strand was also important for the chaperone-like activity of BepA.

161   BepA derivatives with a Cys substitution (note that BepA intrinsically does not possess a Cys residue),

instead of a Pro substitution, at the positions of Asn-105, Ala-106, or Phe-107 in the β2 strand exhibited almost

normal chaperone-like and proteolytic functions (**Figure 1-figure supplement 4B** and **4C**). This supports the idea

that the secondary structure of the edge-strand, but not the individual amino acid residues, is important for its

function. Note that, although it is possible that a Pro mutation also affects a local structure around the edge-strand,

the A106P mutant, which was defective in LptD and BamA degradation, still exhibited significant self-cleavage

of the probably unstructured C-terminal tag (**Figure 1-figure supplement 2B**), suggesting that this mutation, and

possibly the other Pro mutations, specifically affected the edge-strand structure, but not the active site or the

protein's overall structures.

BepA interacts with the N-terminal half of the β-barrel-forming domain of the LptD assembly intermediate

172   While BepA interacts with LptD to promote either its proper OM assembly or proteolytic elimination

depending on the situation (Narita et al., 2013), the details of the BepA–LptD interaction, including the region(s)

in LptD to which BepA binds, remain largely unknown. Thus, we performed a systematic photo-crosslinking

analysis to identify the BepA-contact region in the LptD assembly intermediate LptD^C. We performed photo-

crosslinking experiments in cells ectopically co-expressing an LptD derivative containing a photo-reactive amino

acid analog pBPA [LptD(pBPA)] and a protease-dead variant of BepA, BepA(E137Q). We first introduced pBPA

at each of the 50 positions (approximately every 15 residues) in the mature part of LptD and performed photo-

crosslinking analysis. Cells expressing LptD(pBPA)-His$_{10}$ and BepA(E137Q) were grown and UV-irradiated, and

the total cellular proteins were analyzed by immunoblotting with anti-BepA and anti-His antibodies. Under this

condition, expressed LptD-His$_{10}$ was considerably accumulated as LptD^C (**Figure 2-figure supplement 1**)

irrespective of co-expression of BepA(E137Q). We detected clear crosslinking with BepA mainly in the N-

terminal half of the LptD β-barrel-forming domain (**Figure 2A**). We then performed a detailed photo-crosslinking

analysis for the 20 additional sites in the N-terminal half of the LptD β-barrel-forming domain (**Figure 2B**) and

found that BepA was crosslinked at several of these sites. The residues at the BepA-cross-linkable sites were

oriented both inward and outward in the mature LptD β-barrel (**Figure 2C** and **2D**). Moreover, the residue Gln-

393 at which the strongest crosslinking was observed was oriented inward. These results suggest that LptD while

interacting with BepA would not assume a higher-order structure like closed β-barrel (See Discussion). We

selected a few LptD(pBPA) derivatives that had been crosslinked with BepA as representatives and examined their

functionality. They supported the growth of LptD-depleted cells when expressed from a plasmid, indicating that

they were functional (**Figure 2-figure supplement 2**). The above-mentioned crosslinking results thus likely reflect

a functional interaction of LptD with BepA in the normal assembly pathway.

BepA edge-strand directly interacts with the Tyr-331 residue in the β7 strand of the LptD β-barrel domain

195   We investigated further to identify the region of LptD that interacts with the BepA edge-strand. First,

we examined the effects of the BepA edge-strand Pro mutations (F107P and A106P) on the LptD(pBPA)–BepA

crosslinking. The F107P mutation significantly decreased the efficiency of the crosslinking of BepA(N105pBPA)

and BepA(A106pBPA) with LptD (**Figure 3-figure supplement 1**). Additionally, the A106P mutation exhibited

a similar effect on the crosslinking of BepA(N105pBPA) with LptD. Based on these effects on crosslinking, we

inferred that these mutations affected the interaction of the edge-strand with LptD. Subsequently, we selected

several LptD(pBPA) derivatives that showed relatively strong crosslinking with BepA and examined the effect of

F107P and A106P mutations in the BepA edge-strand on the crosslinking of LptD(pBPA) with BepA. We found

that these mutations altered LptD–BepA crosslinking in a site-specific manner. Further, the amount of crosslinked

products markedly decreased for LptD(Y331pBPA), but not for other mutants (**Figure 3A**). These results strongly

suggest that the region around Tyr-331 in the β7 strand of the LptD β-barrel domain (**Figure 2D**) is crosslinked

with the edge-strand of BepA.

To further confirm the direct interaction of the BepA edge-strand with the LptD β7 strand, we

conducted site-specific disulfide crosslinking experiments. For this analysis, we used the above-described single

Cys derivatives of BepA harboring a Cys residue at the position of Asn-105, Ala-106, or Phe-107, and derivatives

of LptD having a Cys substitution at either of the six positions, including Tyr-331 at which introduction of pBPA

showed clear crosslinking with BepA (**Figure 3A**). The wild-type LptD protein harbors intrinsic four Cys residues

that form two disulfide bonds essential for the LptD function; therefore, each of these Cys-substituted LptD

mutants possessed five Cys residues in total. We confirmed that these Cys-substituted LptD derivatives

accumulated normally and retained their function (**Figure 3-figure supplement 2**). Cells expressing a combination

of BepA Cys mutants and LptD Cys mutants were grown, and total proteins were acid-denatured and dissolved in

SDS containing N-ethylmaleimide (NEM; NEM was included to block free Cys residues). Then, LptD-His$_{10}$ and

its crosslinked products were affinity-isolated using the C-terminal His$_{10}$-tag, treated with or without 2-

mercaptoethanol (ME), and analyzed by SDS-PAGE and anti-BepA immunoblotting. We observed that certain

combinations of BepA and LptD derivatives showed a high-molecular-mass band in electrophoresis results.

Among them, the combination of BepA(N105C) and LptD(Y331C) showed the most intense band that exhibited

reaction with the anti-BepA antibody (**Figure 3B** and **Figure 3-figure supplement 2B**, no ME). These high-

molecular-mass bands were not observed with the wild-type LptD (no additional Cys) and disappeared upon

treatment with ME, suggesting that they were disulfide-crosslinked products (**Figure 3B** and **Figure 3-figure**

supplement 2B, + ME). These results are consistent with the photo-cross-linking experiments (**Figure 3A**) and

indicate that the edge-strand of BepA can directly bind to several regions in the N-terminal half of the LptD β-

barrel-forming domain, which includes the β7 strand containing Tyr-331.

BepA interacts with an LptD intermediate associating with the seam region of BamA on the BAM complex

We further investigated the mode of the interaction of the BepA-associated LptD with the BAM

complex. It has been recently shown that an assembly intermediate of LptD (LptD4213) interacts with the seam

region formed by the N- and C-terminal β-strands (β1 and β16, respectively) in the BamA barrel domain and forms

a hetero-complex, in which the C-terminal β-signal of the LptD4213was associated with the β1 strand of the BamA

seam (Lee et al., 2019). It has ben suggested that the interaction of the β-signal with the BamA β1 strand generally

facilitates the folding of the β-barrel domain of a substrate OMP (See **Figure 4D** and **5B**) (Tomasek and Kahne,

2021). We first examined the interaction of the BepA-associated LptD intermediate with BamA and BamD by

conducting *in vivo* photo-crosslinking experiments using the LptD derivatives with *p*BPA at the position of Glu-

749 or Tyr-726 in addition to the position of Tyr-331. *p*BPA at Glu-749 and Tyr-726 residue, both of which are

located near the β-signal of LptD, have been reported to be crosslinked with BamA and BamD, respectively, during

the LptD assembly (**Figure 4D**) (Lee et al., 2019, 2018). Complementation assay results showed that either of the

LptD derivatives containing one or two *p*BPA at Glu-749 and Tyr-726 were functional (**Figure 4-figure**

supplement 1). After UV-irradiation of the cells expressing LptD-His$_{10}$ and the *p*BPA derivatives, LptD-His$_{10}$ and

its crosslinked products were affinity-isolated from the membrane fractions and analyzed by immunoblotting. The

single *p*BPA derivatives indeed generated crosslinked products with the expected factors as: LptD(Y331*p*BPA)

with BepA, LptD(D749*p*BPA) with BamA, and LptD(Y726*p*BPA) with BamD (**Figure 4A** and **4B**). With the

double *p*BPA derivatives LptD(Y331/D749*p*BPA) and LptD(Y331/Y726*p*BPA), new crosslinked products with

higher molecular sizes were observed in addition to the ones observed with single *p*BPA mutants (**Figure 4A** and

4B). These results showed that the higher molecular-sized products represented BepA–LptD(Y331/D749*p*BPA)–

BamA and BepA–LptD(Y331/Y726*p*BPA)–BamD crosslinked products and that BepA interacts with an assembly

intermediate of LptD on the BAM complex.

250   We further conducted photo-crosslinking experiments using LptD locked on the BAM complex by

using an SH-crosslinker, 1,4-bismaleimidobutane (BMB). It has been previously shown (Lee et al., 2019) that a

cysteine placed near the N-terminal β-strand of the BamA β-barrel (S439C) was crosslinked with a cysteine

introduced near the β-signal of LptD (E733C) via BMB treatment (**Figure 4D**; **Figure 4-figure supplement 2A**).

We introduced a Y331*p*BPA mutation into LptD(E733C) and confirmed that the resultant mutant was functional

(Fig. S10*B*). When the LptD(Y331*p*BPA/E733C) mutant was expressed in a strain having a chromosomal

bamA(S439C) mutant gene, LptD–BepA crosslinked products were detected upon UV irradiation, whereas LptD–

BamA crosslinked products was detected upon BMB treatment. When cells were first treated with BMB and then

UV-irradiated, a higher mass product that reacted with both anti-BepA and anti-BamA antibodies was generated.

The generation of this product depended on both BMB-treatment and UV-irradiation (**Figure 4C**). These results

are fully consistent with the above photo-crosslinking results and further demonstrated that BepA could interact

with an LptD assembly intermediate associating with the seam site of BamA on the BAM complex.

Discussion

The involvement of BepA in the maintenance of structural and functional integrity of the OM was first suggested on the basis of increased sensitivities of the Δ*bepA*(*yfgC*) strain to several antibiotics and chemicals, which was similar to the characteristics of strains with a disruption of genes encoding proteins engaged in outer membrane biogenesis (Narita et al., 2013; Ruiz et al., 2005; Tamae et al., 2008). We have previously shown that BepA is involved in the biogenesis and quality control of LptD, probably on the BAM complex (Daimon et al., 2020, 2017; Narita et al., 2013). However, the mechanism by which BepA recognizes and interacts with LptD remains elusive. This information is important to understand the mechanism of BepA to distinguish between the normal (on-) and off-pathway intermediates of LptD that are either assembled into the OM or degraded, respectively. To gain insight into the BepA–LptD interaction, we examined the role of the conserved edge-strand of BepA in its function. Our results showed that the BepA edge-strand participates not only in the proteolytic activity but also, unexpectedly, in the chaperone-like activity through its direct interaction with LptD (**Figure 1**). The results of the photo- and disulfide-crosslinking analyses indicated that the N-terminal half of the LptD β-barrel-forming domain interacts with the BepA edge-strand (**Figure 2** and **Figure 3**). Moreover, we showed that BepA demonstrated interaction with an LptD assembly intermediate whose C-terminal region was associated with the seam strand of BamA on the BAM complex (**Figure 4**). These observations provide useful insights into the BepA functions involved in the biogenesis and quality control of LptD.

Our previous results have shown that BepA interacts with the BAM complex via its C-terminal TPR domain partly inserted into the periplasmic ring-like structure of the BAM complex (Daimon et al., 2017; Shahrizal et al., 2019). In this study, we observed that BepA showed crosslinking with the N-terminal half of the LptD β-barrel-forming domain but not with the C-terminal half. This observation was consistent with the localization of LptD on the BAM complex; the N-terminal region was localized at/near the periplasmic surface of the BAM complex, and the C-terminal region was inserted deep into the BAM complex's interior. The results of the Cys crosslinking experiments showed that several positions, including Tyr-331, in the N-terminal half of the LptD β-barrel-forming domain interacted with the edge-strand. The recently solved structures of BepA (Bryant et al., 2021; Shahrizal et al., 2019) showed that the active site region including the edge-strand was located inside the BepA molecule, leading to the suggestion that structural changes in BepA including the movement of the α6 and α9 loops covering the active site/edge-strand were necessary to enable access of a substrate to the active site/edge-

strand region (**Figure 5-figure supplement 1**). However, even after such structural changes, the edge-strand

should be located at a recessed position. This suggests that the regions around the BepA-crosslinked positions in

the LptD assembly intermediate do not form an extensive β-sheet structure as found in the mature LptD to gain

access to the edge-strand of BepA (**Figure 5-figure supplement 1**). Furthermore, *p*BPA at positions of both the

inward-pointing and outward-pointing residues in the mature β-barrel domain of LptD was crosslinked with BepA.

Collectively, these results support the hypothesis that the BepA-interacting region of the LptD intermediate is

largely unstructured. The unstructured nature of these proteins is in fact helpful to accommodate in or around the

narrow space inside the BAM complex. The TPR domain of BepA has also been shown to contact with LptD to

promote its biogeneen and degradation (Daimon et al., 2017). Currently, we have no information available on the

part of LptD that interacts with the TPR domain. The TPR domain may act either together with the edge-strand at

the same step or independently at other steps during LptD assembly and degradation.

Further, our results suggest that the proper interaction of LptD with the edge-strand of BepA is

important for the promotion of its assembly as well as degradation by BepA (**Figure 1** and **Figure 5A**). This

finding was unexpected as it raised a question of how the degradation of the normally assembling LptD

intermediate can be avoided despite its interaction with the edge-strand near the protease active site. In the BepA

structures, the conserved His-246 residue is coordinated to a zinc ion in the active site to block the activation of a

water molecule necessary for the catalysis of the proteolytic reaction (Bryant et al., 2021; Shahrizal et al., 2019).

We have recently reported that His-246 acts as a switch to regulate the proteolytic activity of BepA (Daimon et

al., 2020). This His-Switch-mediated repression of the protease activity would enable the interaction of LptD

intermediates at the protease active site of BepA without degradation. Indeed, the derepressed BepA(H246A)

mutant degrades an LptD intermediate on the normal assembly pathway, instead of promoting its biogenesis

(Daimon et al., 2020). One possible model for the discrimination of LptD intermediates by BepA for assembly or

degradation would be as follows (**Figure 5A**). In both assembly and degradation pathways of LptD, an unknown

signal(s), such as a specific interaction of BepA with the substrate and/or the BAM complex, induces a structural

change in BepA (dislocation of the α6 loop covering the active site) to expose its active site and to enable the

interaction between the LptD polypeptide and the edge-strand, whereas His-246 continues to repress the

degradation of LptD by BepA. This interaction is necessary for both proper assembly and degradation of LptD.

The transient interaction of normally assembling LptD with the "protease activity-repressed" state of BepA would

provide sufficient time for the progression of the maturation processes of LptD (including association with LptE),

resulting in the final release of mature LptD from BamA. In contrast, during the prolonged stay of stalled or

misassembled LptD molecules, BepA undergoes a further structural change including the movement of the α9

loop, stochastically or induced by a specific signal(s), to remove His-246 from the active site zinc, leading to the

degradation of the molecules that cannot be subjected to maturation successfully. Our previous results of pulse-

chase experiments showed that, although the conversion of LptD^C to LptD^{NC} was relatively slow, it started

immediately after the synthesis of LptD and proceeded rather constantly (Narita et al., 2013). In contrast,

degradation of stalled LptD in an LptE-depleted strain apparently occurred in a biphasic manner; little degradation

of stalled LptD was observed for several minutes after synthesis ((Narita et al., 2013), indicating the presence of

a lag period for degradation of stalled LptD to occur. These observations are consistent with the above model.

Notably, it has been reported that BepA expression can result in the degradation of a mutant form of LptD

(LptD4213) that probably mimics a late-state assembly intermediate of LptD (Lee et al., 2016) and possesses a

substantial degree of a higher order (β-barrel-like) structure. The mechanism by which LptD4213 is degraded by

BepA remains unclear, but its N-terminal region may interact with BepA.

BepA possibly facilitates the association of LptD with LptE on the BAM complex as the phenotypes

caused by the disruption of BepA, including drug sensitivity and retarded disulfide rearrangement in LptD, can be

suppressed by the overproduction of LptE (Narita et al., 2013). We have previously detected two species of LptD^C;

one is not associated with LptE and observed only in an extremely early phase of its membrane assembly, and the

other is associated with LptE and formed at a later phase (Miyazaki et al., 2018). It is likely that the LptD^C molecule

that was simultaneously crosslinked with BepA and BamA was not associated with LptE, as it was accumulated

in the LptE-limiting condition; therefore, it would represent the former species of LptD^C mentioned above. We

assume that this LptD intermediate may possess a partially folded structure in which the BamA-associating C-

terminal region of the barrel domain has a certain degree of a higher order (β-sheet) structure, but the BepA-

associating N-terminal region is largely unstructured. Such a state of LptD may be favorable for the association of

LptE with LptD (**Figure 4D**). Interestingly, Tyr-331 of LptD that contacts with the edge-strand of BepA is located

at the end of the LptE-surrounding region of the LptD β-barrel in the mature LptD/E complex (**Figure 2D**). The

interaction with the edge-strand of BepA may pin the partially folded structure of LptD transiently and facilitate

the association between LptD and LptE. It might also assist the formation of the β-sheet structure in the β barrel

forming domain of LptD. Although it is possible that BepA actively promotes the LptD–LptE association by

interacting with both LptD and LptE, it is also possible that BepA plays a passive role by maintaining an

appropriate structure of LptD for association with LptE without showing direct interaction with LptE.

The biogenesis of LptD has been well studied. LptD has been a focus of OM protein research because

it provides important information on its essential cellular function and it can be used as a model for OM protein

insertion into the OM by the BAM complex. Based on the results obtained in the previous and current studies, we

propose a model of the BepA-assisted biogenesis of LptD (**Figure 5B**): (i) After synthesis in the cytoplasm, LptD

is translocated to the periplasm through the SecYEG translocon, during or just after which the Cys residues are

oxidized by DsbA to form $LptD^C$ (Chng et al., 2012; Narita et al., 2013). It is then targeted to the BAM complex

with the aid of periplasmic chaperones including SurA (Schwalm et al., 2013; Vertommen et al., 2009); (ii) The

C-terminal β-signal region of LptD is inserted into the BAM complex possibly through the periplasmic ring-like

structure formed by the BamA POTRA domains and the lipoprotein subunits of the BAM complex, and it

establishes interaction with BamD and BamA (Lee et al., 2019); (iii) Although details of the exact juncture at

which interaction occurs are unknown, BepA interacts with the largely unstructured N-terminal half of the LptD

β-barrel-forming domain whose C-terminal β-signal region is associated with the seam region of BamA. This

helps the association of LptE with LptD; (iv) Then, the folding (β-sheet formation) of the unstructured N-terminal

region of the LptD β-barrel domain occurs to form a premature form of LptD with a substantially folded β-barrel

domain and LptE within it, like LptD4213 (Lee et al., 2019, 2016); (v) The β-barrel domain of LptD β-barrel is

finally released from BamA and closed to form the mature LptD/E complex. The isomerization of two disulfide

bonds in LptD ($LptD^C$ to $LptD^{NC}$) should occur at a later step after the association of LptD and LptE (Lee et al.,

2016; Miyazaki et al., 2018). The LptD intermediates that are stalled at certain steps in the above-mentioned

processes as a result of misfolding are eliminated by the action of several peptidases including DegP (in the

periplasm) and BepA /YcaL (on the BAM complex) (Soltes et al., 2017).

Our study reports several new findings on the interaction of BepA with LptD and BamA on the BAM

complex where BepA plays a crucial role in the biogenesis and degradation of LptD. Further, we proposed a model

explaining the role of BepA in these processes. Nonetheless, there are many questions that warrant further

investigation. It would be especially important to elucidate the mechanism by which the substrate gains access to

the active site buried inside the BepA molecule and the manner in which the switching of BepA from the state

with chaperone-like function to that with protease function occurs. The possible movement of the α6 and α9 loops

should be directly examined. It is possible that there are signals that arise from the BAM complex and/or from

(stalled) LptD to induce this structural/functional conversion of BepA. Further study, including structural and

biochemical analysis of the BepA−LptD−BAM complex, is necessary to substantiate our model and to elucidate

the molecular details of BepA functions. Cell surface proteins are suitable drug targets as they are more easily

accessible from the external milieu than cytoplasmic proteins. These studies can provide a basis for the

development of new drugs targeted to BepA/LptD/BAM.

Materials and methods

Key resources table

Reagent type or resource	Designation	Source reference	Identifiers	Additional information
Strains	*E. coli*	This study	N/A	Supplementary file 1
Virus	P1 bacteriophage	Laboratory stock	CGSC12133	
Recombinant DNA	Plasmids	This study	N/A	Supplementary file 2
Oligonucleotides		This study	PCR primers	described in the below
Antibodies	1st antibodies	This study	N/A	listed in the below
Antibody	Goat Anti-Rabbit IgG (H + L)-HRP Conjugate	Bio-Rad Laboratories	1706515	
Chemical compound	H-p-Bz-Phe-OH	Bachem	F2800	
Chemical compound	Methionine, L-[^{35}S] Translation Grade	American Radiolabeled Chemicals	ARS 01014	
Chemical compound	nProtein A Sepharose^{TM} 4 Fast Flow	GE Healthcare	17528004	
Chemical compound	Ni-NTA Agarose	QIAGEN	30250	
Critical Commercial Assays	ECL^{TM} Western Blotting Detection Reagents	GE Healthcare	RPN2106	
Critical Commercial Assays	ECL^{TM} Prime Western Blotting Detection Reagents	GE Healthcare	RPN2232	
Software, Algorithm	Microsoft Excel	Microsoft	N/A	
Software, Algorithm	Bio-imaging Analyzer BAS-1800, BAS-5000	Fujifilm/GE Healthcare	N/A	
Software, Algorithm	Image Qaunt LAS 4000 mini	Fujifilm/GE Healthcare	N/A	
Software, Algorithm	Multi Gauge	Fujifilm/GE Healthcare	N/A	

Bacterial strains, Plasmids and Primers

Escherichia coli K12 strains and plasmids used in this study are listed in Tables S1 and S2, respectively. Details

of the strain and plasmid construction and media are described in Construction of Mutant Strains and Plasmids

Construction, respectively.

Media and Bacterial Cultures

E. coli cells were grown in L rich medium (10 g/L bacto-tryptone, 5 g/L bacto-yeast extract, 5 g/L NaCl; pH

adjusted to 7.2 with NaOH) or M9 synthetic medium (without $CaCl_2$; (Miller, 1972)) supplemented with maltose

(final 0.2%), glycerol (final 0.4%), all amino acids (except Met and Cys; final concentration of 20 μg/mL each).

50 μg/mL ampicillin (Amp), 20 μg/mL chloramphenicol (Cm), 25 μg/mL kanamycin (Km), 25 μg/mL tetracycline

(Tet) and 50 μg/mL spectinomycin (Spc) were added as appropriate for growing plasmid-bearing cells and

selection of transformants and transductants. Bacterial growth was monitored with Mini photo 518R (660 nm;

TAITEC Co., Saitama, Japan).

Antibodies

Penta-His HRP conjugate was purchased from QIAGEN (Hilden, Germany). Anti-BepA (Narita et al., 2013), anti-

LptD (Narita et al., 2013) and anti-BamA (Gunasinghe et al., 2018) antibodies were described previously.

Construction of Mutant Strains

RM2091 (JE6631, *purC80*::Tn*10*) was constructed by transferring the *purC80*::Tn*10* marker, which is located

near the *bepA* gene, from CAG18470 (Nichols et al., 1998) into JE6631 (Miyazaki et al., 2018), respectively, by

P1 transduction. RM2243 (AD16, *bepA(E137Q) purC80*::Tn*10*) was constructed as follows. pRM330 (a plasmid

carrying *bepA(E137Q)*, see below) was introduced into RM2091 to yield cells with pRM330 integrated into the

chromosome by homologous recombination in the *bepA* region. They were then grown on an L-agar plate

containing 5% sucrose to select cells that had lost the integrated plasmid. The plasmid-cured cells were screened

for those having the chromosomal *bepA(E137Q)* allele at the *bepA* locus. The *bepA(E137Q)* allele was finally

transferred to AD16 (Kihara et al., 1995) by joint P1 transduction with the *purC80*::Tn*10* marker. One of such

strains was named RM2243. RM3654 (AD16, Δ*bepA*, *bamA*[+] *zae502*::Tn*10*) and RM3655 (AD16, *bamA(S439C)*

zae502::Tn*10*) were constructed as follows, pRM845 (a plasmid carrying *bamA(S439C)*, see below) was

introduced into YH188 (JE6631, *zae502*::Tn*10*) (Hizukuri and Akiyama, 2012) to yield cells with pRM845

integrated into the chromosome by homologous recombination in the *bamA* region. They were then grown on an

L-agar plate containing 5% sucrose to select cells that had lost the integrated plasmid. The plasmid-cured cells

were screened for those having the chromosomal *bamA(S439C)* allele at the *bamA* locus. The *bamA(S439C)* allele

was transferred to SN56 (Narita et al., 2013) by joint P1 transduction with the *zae502*::Tn*10* marker. Strains having

the *bamA+* allele and the *bamA(S439C)* allele were picked up and named RM3654 and RM3655, respectively.

RM2831 (HM1742, *kan araC*-P*araBAD*-*lptD*) were constructed by essentially the same procedure as the construction

of strains with a chromosomal C-terminal his$_{10}$-tagged gene (Miyazaki et al., 2020b). First, a *kan araC*-P*araBAD*-

lptD fragment having a sequence identical to the upstream or downstream region of the *lptD* start codon at the

respective ends of the fragment, was PCR-amplified from pRM741 (a plasmid carrying a *kan* cassette at the

upstream of an *araC*-P*araBAD*) using a pair of primers, ara-lptD-f (5'-

TTGTCACGCGCAACGTTACCGATGATGGAACAATAAAATCAACGTCATATGAATATCCTCCTTAG-

3') and ara-lptD-r (5'-

GGTGGCAATCATGGTGGCCAGGAGAGTGGGGATACGTTTTTTCATGGTGAATTCCTCCTGCTAG-3').

Then, the chromosomal *lptD* locus of the *E. coli* DY330 strain was replaced by this fragment using the λ-Red

recombination system (Yu et al., 2000). The *kan araC*-P*araBAD*-*lptD* was finally transferred to HM1742 by P1

transduction.

Plasmids Construction

pSTD689-derived plasmids carrying a *bepA* mutants were constructed from pRM290 (pSTD689-*bepA*) (Daimon

et al., 2017) by site-directed mutagenesis. Derivatives of pRM291 (pSTD689-*bepA(E137Q)*) (Daimon et al., 2017)

carrying an additional Cys mutation and derivatives of pUC-*bepA(E137Q)-his$_{10}$* (pUC18-*bepA(E137Q)-his$_{10}$*)

(Narita et al., 2013) carrying an *amber* mutation were constructed by site-directed mutagenesis. pUC18-

bepA(Pro)-his$_{10}$ and pUC18-*bepA(amb)-his$_{10}$* plasmids were constructed from pUC-*bepA-his$_{10}$* (pUC18-*bepA-*

his$_{10}$) (Narita et al., 2013) by site-directed mutagenesis. To construct pUC18-*bepA(Pro, amb, E137Q)-his$_{10}$*

plasmids, a mutation for the individual Pro substitutions was introduced into each of the pUC18-*bepA(amb,*

E137Q)-his$_{10}$ plasmids by site-directed mutagenesis. pNB91 (pMW118-*bepA(E137Q)*) was constructed by

subcloning an EcoRI-HindIII *bepA(E137Q)* fragment prepared from pUC-*bepA(E137Q)* (pUC18-*bepA(E137Q)*)

(Narita et al., 2013) into the same sites of pMW118. pRM807 (pMW118-*bepA(A106P, E137Q)*) and pRM808

(pMW118-*bepA(F107P, E137Q)*) were also constructed by subcloning an EcoRI-HindIII *bepA* fragment from

each of pSTD639-*bepA* plasmids into the same sites of pMW118.

Plasmids carrying *lptD(amb)-his$_{10}$* were constructed from pRM309 (pRM294-*lptD-his$_{10}$*) (Miyazaki

et al., 2018) by site-directed mutagenesis. pRM821 (pRM294-*lptD(Y331amb, D749amb)-his$_{10}$*) and pRM822

(pRM294-*lptD(Y331amb, Y726amb)-his₁₀*) were constructed from pRM626 (pRM294-*lptD(Y331amb)-his₁₀*) by

site-directed mutagenesis. pRM829 (pRM294-*lptD(E733C)-his₁₀*) and pRM831 (pRM294-*lptD(Y331, E733C)-*

his₁₀) were constructed from pRM309 and pRM626, respectively, by site-directed mutagenesis. Plasmid carrying

lptD(Cys)-his₁₀ were constructed as follows. Each mutation for the Cys substitutions was introduced to pRM309

by site-directed mutagenesis. The BamHI-HindIII *lptD(Cys)-his₁₀* fragment of the resulting plasmids was

subcloned into the same site of pTWV228 (Takara Bio inc., Shiga, Japan) to generate pTWV228- *lptD(Cys)-his₁₀*.

pRM320 (pUC118-*bepA-yfgD*) was constructed by PCR amplification of the *bepA-yfgD* fragment

from the genome of MC4100 using a pair of primers, bepA-f (5'-

GCGCGCGGATCCATTTGAGTGGGCTAATCTTCG-3') and yfgD-r (5'-

GCGCGCGTCGACCGAACTACGCGAAGTTAATCC-3'), and, cloning it into the BamHI-SalI site of pUC118

(Takara Bio) after digestion with these enzymes. For the construction of pRM324, the *bepA(E137Q)* mutation was

introduced into pRM320 by site-directed mutagenesis. pRM330 (pK18mobsacB-*bepA(E137Q)-yfgD*) was

constructed by subcloning the BamHI-SalI *bepA(E137Q)-yfgD* fragment from pRM324 into the same sites of

pK18mobsacB (Schäfer et al., 1994).

pRM823 (pUC118-*bamA*) was constructed by *in vitro* recombination using In-Fusion HD cloning kit

(Takara Bio) of a EcoRI-BamHI fragment from pUC118 and a *bamA* fragment prepared by PCR amplification

from the genome of MC4100 using a pair of primers, bamA-f (5'-

GCGCGAATTCAGGAAGAACGCATAATAACG-3') and bamA-r (5'-

GCGCGGATCCTTACCAGGTTTTACCGATG-3'). For the construction of pRM836, the *bamA(S439C)*

mutation was introduced into pRM823 by site-directed mutagenesis. pRM845 (pK18mobsacB-*bamA(S439C)*) was

constructed by subcloning the EcoRI-BamHI *bamA(S439C)* fragment from pRM823 into the same sites of

pK18mobsacB.

Immunoblotting analysis

Acid-denatured proteins were solubilized in SDS-sample buffer (62.5 mM Tris-HCl (pH 6.8), 2% SDS, 10%

glycerol, 5 mg/mL bromophenol blue) with or without 10% β-ME, separated by SDS-PAGE and electro-blotted

onto a PVDF membrane (Merck Millipore; Billerica, MA). The membrane was first blocked with 5% skim milk

in PBST (Phosphate Buffered Saline with Tween 20), and then incubated with Penta-His HRP conjugate (1/2,000

or 1/3,000 dilution)), anti-BepA (1/10,000), anti-LptD (1/50,000) or anti-BamA (1/20,000). After washing with

PBST, the membrane was incubated with a horseradish peroxidase (HRP)-conjugated secondary antibody

(1/5,000) (Goat Anti-Rabbit IgG (H + L)-HRP Conjugate; Bio-Rad Laboratories, Inc., Hercules, CA) in PBST

(This step was omitted for the detection using Penta-His HRP conjugate). Proteins were visualized with ECL^{TM}

Western Blotting Detection Reagents (GE Healthcare UK Ltd, *Amersham Place* Little Chalfont, England) or

ECL^{TM} Prime Western Blotting Detection Reagents (GE Healthcare) and LAS4000 mini lumino-image analyzer

(GE Healthcare).

Pulse-chase analysis for assay of the LptD disulfide-isomerization

Cells were first grown at 30°C in M9-medium supplemented with 2 µg/mL thiamine, 0.4% glycerol, 0.2% maltose,

all amino acids (except Met and Cys) with or without 0.05% arabinose until early log phase. After induction with

1 mM IPTG for 15 min, cells were pulse-labeled with 370 kBq/mL [^{35}S]Methionine for 1 min. At appropriate time

points after addition of excess nonradioactive Met (final conc. 250 µg/mL), total cellular proteins were precipitated

with 5% TCA, washed with acetone, solubilized in SDS-buffer (50 mM Tris-HCl (pH 8.1), 1% SDS, 1 mM EDTA)

and diluted 33-fold with Triton-buffer (50 mM Tris-HCl (pH 8.1), 150 mM NaCl, 2% Triton X-100, 0.1 mM

EDTA). After clarification, samples were incubated with anti-LptD antibodies and nProtein A Sepharose^{TM} 4 Fast

Flow (GE healthcare) at 4ºC over-night with slow rotation. Proteins bound to the antibody/ProteinA-Sepharose

were recovered by centrifugation, washed with Triton buffer and then with 10 mM Tris-HCl (pH 8.1) and eluted

by incubation at 98°C for more than 5 min in SDS-sample buffer. The samples were divided into two portions and

one was treated with 10% β-ME. The proteins were separated by SDS-PAGE, and visualized with BAS1800

phosphoimager. Relative LptD^{NC} were calculated by the following equation: Relative LptD^{NC} = [LptD$^{NC}_{(X}$

$_{min)}$]/[LptD$^{C}_{(X min)}$ + LptD$^{NC}_{(X min)}$], where LptD^{NC} and LptD^{C} are the intensities of the respective bands.

In vivo **photo-crosslinking analysis**

For the experiments in Figure 1D and Figure 3-figure supplement 1, cells were grown at 30°C in L medium

containing 0.5 mM *p*BPA and 0.02% arabinose until early log phase and induced with 1 mM IPTG for 1 h. A half

volume of the cell cultures was put on a petri dish and UV-irradiated at 4ºC for 10 min using B-100AP UV lamp

(365 nm; UVP, LLC., Upland, CA), at the distance of 4 cm. The other half was kept on ice as non-UV-irradiated

samples. Membrane fractions were prepared by sonical cell disruption and the following ultracentrifugation, and

solubilized with SDS-buffer (50 mM Tris-HCl (pH 8.1), 1% SDS, 1 mM EDTA). After 33-fold diluted with Triton-

buffer (50 mM Tris-HCl (pH 8.1), 150 mM NaCl, 2% Triton X-100, 0.1 mM EDTA) and clarification, samples

were subjected to pull-down with Ni-NTA Agarose (QIAGEN). The isolated proteins were solubilized in SDS-

sample buffer (62.5 mM Tris-HCl (pH 6.8), 2% SDS, 10% glycerol, 5 mg/mL bromophenol blue) containing ME

and analyzed by SDS-PAGE and immunoblotting analysis.

For the experiments in Figure 2A, 2B and 3A, cells were grown at 30°C in L medium containing 0.5

mM pBPA and 0.02% arabinose until early log phase and induced with 1 mM IPTG for 3 h. While a half volume

of the cell cultures was UV-irradiated for 10 min, the other half was kept on ice as non-UV-irradiated samples, as

above. Total cellular proteins were precipitated with 5% TCA, washed with acetone, and solubilized in SDS-

sample buffer containing ME. The samples were subjected to SDS-PAGE and immunoblotting analysis.

For experiments in Figure 4A and 4B, cells were grown at 30°C in L medium containing 0.5 mM

pBPA and 0.02% arabinose until early log phase and induced with 1 mM IPTG for 3 h. While a half volume of

the cell cultures was UV-irradiated for 10 min, the other half was kept on ice as non-UV-irradiated samples, as

above. Membrane fractions were prepared as above and solubilized in SDS-buffer. After dilution with Triton-

buffer and clarification, samples were subjected to pull-down with Ni-NTA Agarose. The isolated proteins were

solubilized in SDS-sample buffer containing ME and analyzed by SDS-PAGE and immunoblotting analysis.

BMB crosslinking combined with photo-crosslinking

Cells were grown at 30°C in L medium containing 0.5 mM pBPA and 0.02% arabinose until early log phase and

induced with 1 mM IPTG for 3 h. BMB crosslinking was performed essentially according to the previously

described procedures (Lee et al., 2019). After quenching the BMB crosslinking, cells were treated with or without

UV-irradiation for 30 min at 4°C. Total cellular proteins were precipitated with 5% TCA, washed with acetone,

and suspended in SDS-sample buffer containing 12.5 mM NEM for blocking the free Cys residues. After dilution

with Triton-buffer and clarification, samples were subjected to pull-down with Ni-NTA Agarose. The isolated

proteins were suspended in SDS-sample buffer containing ME and analyzed by SDS-PAGE and immunoblotting

analysis.

Article and author information

Author details

Ryoji Miyazaki

Institute for Frontier Life and Medical Sciences, Kyoto University, Kyoto, Japan

Contribution: Conceptualization, Methodology, Investigation, Resources, Writing–Original Draft, Writing–Review & Editing and Funding Acquisition

Competing interests: No competing interests declared

ORCID: 0000-0003-1626-2569

Tetsuro Watanabe

Institute for Frontier Life and Medical Sciences, Kyoto University, Kyoto, Japan

Contribution: Methodology, Investigation, Resources

Competing interests: No competing interests declared

ORCID:

Kohei Yoshitani

Institute for Frontier Life and Medical Sciences, Kyoto University, Kyoto, Japan

Contribution: Resources

Competing interests: No competing interests declared

ORCID:

Yoshinori Akiyama

Institute for Frontier Life and Medical Sciences, Kyoto University, Kyoto, Japan

Contribution: Writing–Original Draft, Writing–Review & Editing, Supervision and Funding Acquisition

Competing interests: No competing interests declared

ORCID: 0000-0003-4483-5408

Funding

This work was supported by the Japan Society for the Promotion of Science KAKENHI Grants 18H06047,

19K21179 and 20K15715 (to R.M.), and 15H01532 and 18H023404 (to Y.A.), and research grants from the

Institute for Fermentation Y-2020-02-027 (to R.M.). and from the Nagase Science and Technology Foundation (to

Y.A.).

Acknowledgments

We thank our laboratory members and S. Narita for discussion and helpful advices.

References

References

Akiyama K, Mizuno S, Hizukuri Y, Mori H, Nogi T, Akiyama Y. 2015. Roles of the membrane-reentrant β-

hairpin-like loop of RseP protease in selective substrate cleavage. *eLife* **4**:e08928. doi:10.7554/eLife.08928

Bakelar J, Buchanan SK, Noinaj N. 2016. The structure of the β-barrel assembly machinery complex. *Science*

351:180–186. doi:10.1126/science.aad3460

Bryant JA, Cadby IT, Chong ZS, Boelter G, Sevastsyanovich YR, Morris FC, Cunningham AF, Kritikos G, Meek

RW, Banzhaf M, Chng SS, Lovering AL, Henderson IR. 2021. Structure-function characterization of the

conserved regulatory mechanism of the *Escherichia coli* M48 metalloprotease BepA. *J Bacteriol*

203:e00434-20. doi:10.1128/JB.00434-20

Chimalakonda G, Ruiz N, Chng S-S, Garner RA, Kahne D, Silhavy TJ. 2011. Lipoprotein LptE is required for the

assembly of LptD by the β-barrel assembly machine in the outer membrane of *Escherichia coli*. *Proc Natl*

Acad Sci U S A **108**:2492–2497. doi:10.1073/pnas.1019089108

Chin JW, Schultz PG. 2002. In vivo photocrosslinking with unnatural amino acid mutagenesis. *ChemBioChem*

3:1135–1137. doi:10.1002/1439-7633(20021104)3:11<1135::AID-CBIC1135>3.0.CO;2-M

Chng SS, Xue M, Garner RA, Kadokura H, Boyd D, Beckwith J, Kahne D. 2012. Disulfide rearrangement

triggered by translocon assembly controls lipopolysaccharide export. *Science* **337**:1665–1668.

doi:10.1126/science.1227215

Daimon Y, Iwama-Masui C, Tanaka Y, Shiota T, Suzuki T, Miyazaki R, Sakurada H, Lithgow T, Dohmae N,

Mori H, Tsukazaki T, Narita S-I, Akiyama Y. 2017. The TPR domain of BepA is required for productive

interaction with substrate   proteins and the β-barrel assembly machinery complex. *Mol Microbiol* **106**:760–

776. doi:10.1111/mmi.13844

Daimon Y, Narita S-I, Miyazaki R, Hizukuri Y, Mori H, Tanaka Y, Tsukazaki T, Akiyama Y. 2020. Reversible

autoinhibitory regulation of *Escherichia coli* metallopeptidase BepA for selective β-barrel protein

degradation. *Proc Natl Acad Sci U S A* **117**:27989–27996. doi:10.1073/pnas.2010301117

Dong H, Xiang Q, Gu Y, Wang Z, Paterson NG, Stansfeld PJ, He C, Zhang Y, Wang W, Dong C. 2014. Structural

basis for outer membrane lipopolysaccharide insertion. *Nature* **511**:52–56. doi:10.1038/nature13464

Gu Y, Li H, Dong H, Zeng Y, Zhang Z, Paterson NG, Stansfeld PJ, Wang Z, Zhang Y, Wang W, Dong C. 2016.

Structural basis of outer membrane protein insertion by the BAM complex. *Nature* **531**:64–69.

doi:10.1038/nature17199

Gunasinghe SD, Shiota T, Stubenrauch CJ, Schulze KE, Webb CT, Fulcher AJ, Dunstan RA, Hay ID, Naderer T,

Whelan DR, Bell TDM, Elgass KD, Strugnell RA, Lithgow T. 2018. The WD40 Protein BamB Mediates

Coupling of BAM Complexes into Assembly Precincts in the Bacterial Outer Membrane. *Cell Rep* **23**:2782–

2794. doi:10.1016/j.celrep.2018.04.093

Han L, Zheng J, Wang Y, Yang X, Liu Y, Sun C, Cao B, Zhou H, Ni D, Lou J, Zhao Y, Huang Y. 2016. Structure

of the BAM complex and its implications for biogenesis of outer-membrane   proteins. *Nat Struct Mol Biol*

23:192–196. doi:10.1038/nsmb.3181

Hart EM, Gupta M, Wühr M, Silhavy TJ. 2020. The gain-of-function allele *bamA (E470K)* bypasses the essential

requirement for BamD in β-barrel outer membrane protein assembly. *Proc Natl Acad Sci U S A* **117**:18737–

18743. doi:10.1073/pnas.2007696117

Hart EM, Mitchell AM, Konovalova A, Grabowicz M, Sheng J, Han X, Rodriguez-Rivera FP, Schwaid AG,

Malinverni JC, Balibar CJ, Bodea S, Si Q, Wang H, Homsher MF, Painter RE, Ogawa AK, Sutterlin H,

Roemer T, Black TA, Rothman DM, Walker SS, Silhavy TJ. 2019. A small-molecule inhibitor of BamA

impervious to efflux and the outer membrane permeability barrier. *Proc Natl Acad Sci U S A* **116**:21748–

21757. doi:10.1073/pnas.1912345116

Hizukuri Y, Akiyama Y. 2012. PDZ domains of RseP are not essential for sequential cleavage of RseA or stress-

induced σE activation *in vivo*. *Mol Microbiol* **86**:1232–1245. doi:10.1111/mmi.12053

Iadanza MG, Higgins AJ, Schiffrin B, Calabrese AN, Brockwell DJ, Ashcroft AE, Radford SE, Ranson NA. 2016.

Lateral opening in the intact β-barrel assembly machinery captured by cryo-EM. *Nat Commun* **7**:12865.

doi:10.1038/ncomms12865

Kihara A, Akiyama Y, Ito K. 1995. FtsH is required for proteolytic elimination of uncomplexed forms of SecY,

an   essential   protein   translocase   subunit.   *Proc   Natl   Acad   Sci   U   S   A*   **92**:4532–4536.

doi:10.1073/pnas.92.10.4532

Konovalova A, Kahne DE, Silhavy TJ. 2017. Outer membrane biogenesis. *Annu Rev Microbiol* **71**:539–556.

doi:10.1146/annurev-micro-090816-093754

Lee J, Sutterlin HA, Wzorek JS, Mandler MD, Hagan CL, Grabowicz M, Tomasek D, May MD, Hart EM, Silhavy

TJ, Kahne D. 2018. Substrate binding to BamD triggers a conformational change in BamA to control

membrane insertion. *Proc Natl Acad Sci U S A* **115**:2359–2364. doi:10.1073/pnas.1711727115

Lee J, Tomasek D, Santos TM, May MD, Meuskens I, Kahne D. 2019. Formation of a β-barrel membrane protein

is catalyzed by the interior surface of the assembly machine protein BamA. *eLife* **8**:e49787.

doi:10.7554/eLife.49787

Lee J, Xue M, Wzorek JS, Wu T, Grabowicz M, Gronenberg LS, Sutterlin HA, Davis RM, Ruiz N, Silhavy TJ,

Kahne DE. 2016. Characterization of a stalled complex on the β-barrel assembly machine. *Proc Natl Acad*

Sci U S A **113**:8717–8722. doi:10.1073/pnas.1604100113

López-Pelegrín M, Cerdà-Costa N, Martínez-Jiménez F, Cintas-Pedrola A, Canals A, Peinado JR, Marti-Renom

MA, López-Otín C, Arolas JL, Gomis-Rüth FX. 2013. A novel family of soluble minimal scaffolds provides

structural insight into the   catalytic domains of integral membrane metallopeptidases. *J Biol Chem*

288:21279–21294. doi:10.1074/jbc.M113.476580

Miller JH. 1972. Experiments in Molecular Genetics. Cold Spring Harbor Laboratory Press.

Miyazaki R, Akiyama Y, Mori H. 2020a. A photo-cross-linking approach to monitor protein dynamics in living

cells. *Biochim Biophys acta Gen Subj* **1864**:129317. doi:10.1016/j.bbagen.2019.03.003

Miyazaki R, Akiyama Y, Mori H. 2020b. Fine interaction profiling of VemP and mechanisms responsible for its

translocation-coupled arrest-cancelation. *eLife* **9**:e62623. doi:10.7554/eLife.62623

Miyazaki R, Myougo N, Mori H, Akiyama Y. 2018. A photo-cross-linking approach to monitor folding and

assembly of newly synthesized proteins in a living cell. *J Biol Chem* **293**:677–686.

doi:10.1074/jbc.M117.817270

Narita S, Masui C, Suzuki T, Dohmae N, Akiyama Y. 2013. Protease homolog BepA (YfgC) promotes assembly

and degradation of β-barrel membrane proteins in *Escherichia coli*. *Proc Natl Acad Sci U S A* **110**:E3612–

E3621. doi:10.1073/pnas.1312012110

Nichols BP, Shafiq O, Meiners V. 1998. Sequence analysis of Tn*10* insertion sites in a collection of *Escherichia*

coli strains used for genetic mapping and strain construction. *J Bacteriol* **180**:6408–6411.

Nikaido H. 2003. Molecular basis of bacterial outer membrane permeability revisited. *Microbiol Mol Biol Rev*

67:593–656. doi:10.1128/mmbr.67.4.593-656.2003

Oh E, Becker AH, Sandikci A, Huber D, Chaba R, Gloge F, Nichols RJ, Typas A, Gross CA, Kramer G, Weissman

JS, Bukau B. 2011. Selective ribosome profiling reveals the cotranslational chaperone action of trigger factor

in vivo. *Cell* **147**:1295–1308. doi:10.1016/j.cell.2011.10.044

Plummer AM, Fleming KG. 2016. From Chaperones to the Membrane with a BAM! *Trends Biochem Sci* **41**:872–

882. doi:10.1016/j.tibs.2016.06.005

Qiao S, Luo Q, Zhao Y, Zhang XC, Huang Y. 2014. Structural basis for lipopolysaccharide insertion in the

bacterial outer membrane. *Nature* **511**:108–111. doi:10.1038/nature13484

Rawlings ND, Barrett AJ, Thomas PD, Huang X, Bateman A, Finn RD. 2018. The MEROPS database of

proteolytic enzymes, their substrates and inhibitors in 2017 and a comparison with peptidases in the

PANTHER database. *Nucleic Acids Res* **46**:D624–D632. doi:10.1093/nar/gkx1134

Ricci DP, Silhavy TJ. 2019. Outer membrane protein insertion by the β-barrel assembly machine. *EcoSal Plus*

8:1–9. doi:10.1128/ecosalplus.ESP-0035-2018

Ruiz N, Chng S-S, Hiniker A, Kahne D, Silhavy TJ. 2010. Nonconsecutive disulfide bond formation in an essential

integral outer membrane protein. *Proc Natl Acad Sci U S A* **107**:12245–12250.

doi:10.1073/pnas.1007319107

Ruiz N, Falcone B, Kahne D, Silhavy TJ. 2005. Chemical conditionality: a genetic strategy to probe organelle

assembly. *Cell* **121**:307–317. doi:10.1016/j.cell.2005.02.014

Schäfer A, Tauch A, Jäger W, Kalinowski J, Thierbach G, Pühler A. 1994. Small mobilizable multi-purpose

cloning vectors derived from the *Escherichia coli* plasmids pK18 and pK19: selection of defined deletions

in the chromosome of *Corynebacterium glutamicum*. *Gene* **145**:69–73. doi:10.1016/0378-1119(94)90324-7

Schwalm J, Mahoney TF, Soltes GR, Silhavy TJ. 2013. Role for Skp in LptD assembly in *Escherichia coli*. *J*

Bacteriol **195**:3734–3742. doi:10.1128/JB.00431-13

Shahrizal M, Daimon Y, Tanaka Y, Hayashi Y, Nakayama S, Iwaki S, Narita S-I, Kamikubo H, Akiyama Y,

Tsukazaki T. 2019. Structural basis for the function of the β-barrel assembly-enhancing protease BepA. *J*

Mol Biol **431**:625–635. doi:10.1016/j.jmb.2018.11.024

Soltes GR, Martin NR, Park E, Sutterlin HA, Silhavy TJ. 2017. Distinctive roles for periplasmic proteases in the

maintenance of essential outer membrane protein assembly. *J Bacteriol* **199**:e00418-17.

doi:10.1128/JB.00418-17

Sperandeo P, Martorana AM, Polissi A. 2017. The lipopolysaccharide transport (Lpt) machinery: A

675   nonconventional transporter for lipopolysaccharide assembly at the outer membrane of Gram-negative

676   bacteria. *J Biol Chem* **292**:17981–17990. doi:10.1074/jbc.R117.802512

677   Stöcker W, Bode W. 1995. Structural features of a superfamily of zinc-endopeptidases: the metzincins. *Curr Opin*

678   *Struct Biol* **5**:383–390. doi:10.1016/0959-440x(95)80101-4

679   Tamae C, Liu A, Kim K, Sitz D, Hong J, Becket E, Bui A, Solaimani P, Tran KP, Yang H, Miller JH. 2008.

680   Determination of antibiotic hypersensitivity among 4,000 single-gene-knockout mutants of *Escherichia coli*.

681   *J Bacteriol* **190**:5981–5988. doi:10.1128/JB.01982-07

682   Tomasek D, Kahne D. 2021. The assembly of β-barrel outer membrane proteins. *Curr Opin Microbiol* **60**:16–23.

683   doi:10.1016/j.mib.2021.01.009

684   Tomasek D, Rawson S, Lee J, Wzorek JS, Harrison SC, Li Z, Kahne D. 2020. Structure of a nascent membrane

685   protein as it folds on the BAM complex. *Nature* **583**:473–478. doi:10.1038/s41586-020-2370-1

686   Vertommen D, Ruiz N, Leverrier P, Silhavy TJ, Collet J-F. 2009. Characterization of the role of the *Escherichia*

687   *coli* periplasmic chaperone SurA using differential proteomics. *Proteomics* **9**:2432–2443.

688   doi:10.1002/pmic.200800794

689   Wu T, McCandlish AC, Gronenberg LS, Chng S-S, Silhavy TJ, Kahne D. 2006. Identification of a protein complex

690   that assembles lipopolysaccharide in the outer membrane of *Escherichia coli*. *Proc Natl Acad Sci U S A*

691   **103**:11754–11759. doi:10.1073/pnas.0604744103

692   Yu D, Ellis HM, Lee EC, Jenkins NA, Copeland NG, Court DL. 2000. An efficient recombination system for

693   chromosome engineering in *Escherichia coli*. *Proc Natl Acad Sci U S A* **97**:5978–5983.

694   doi:10.1073/pnas.100127597

Figure 1

Figure 1. The edge-strand of BepA is crucial for functional interaction with LptD

(**A**) Crystal structure of BepA (PDB code: 6AIT). The peptidase and the TPR domains of BepA are shown in gray and orange, respectively. The edge-strand, the proteolytic active site (the HExxH motif and the third zinc ligand, Glu-201) and the regulatory His-246 residue (His switch) in the peptidase domain are shown in red, blue, and green, respectively, and the coordinated zinc atom is shown in yellow. An enlarged view of the active site region is shown in *right*. (**B**) Protease activities of the BepA edge-strand mutants. Cells of SN56 (Δ*bepA*) carrying pTWV228-*lptD-his₁₀* and either pSTD689 or pSTD689-*bepA* plasmids were grown at 30°C in L-medium until early log phase and induced with 1 mM IPTG for 1 h. Total cellular proteins were acid-precipitated and analyzed by 7.5 or 10% Laemmli SDS-PAGE and immunoblotting with the indicated antibodies. (**C**) *In vivo* photo-crosslinking analysis of the BepA edge-strand. Cells of SN56 carrying pEVOL-pBpF and pUC18-*bepA(E137Q, amb)-his₁₀* plasmids were grown at 30°C in L medium containing 0.02% arabinose and 0.5 mM *p*BPA until early log phase, and induced with 1 mM IPTG for 1 h to express the indicated BepA(*p*BPA) variants. The cultures were divided into two portions, each of which was treated with or without UV-irradiation for 10 min at 4°C. Proteins of the total membrane fractions were subjected to pull-down with Ni-NTA agarose. Purified proteins were analyzed by 7.5% Laemmli SDS-PAGE and immunoblotting with the indicated antibodies. Open triangles indicate unknown crosslinked products. (**D**) Chaperone-like activities of the BepA edge-strand mutants. Cells of SN56 carrying pSTD689 or a pSTD689-*bepA* plasmid were grown at 30°C in M9-based medium until early log phase, induced with 1 mM IPTG for 15 min, pulse-labeled with ³⁵S-Met for 1 min and chased for the indicated periods. At each time point, total cellular proteins were acid-precipitated, subjected to IP with an anti-LptD antibody, and analyzed by 7.5% Laemmli SDS-PAGE followed by phosphorimaging. The ratio of the band intensities of LptD^NC to that of total LptD (LptD^C + LptD^NC) at the each time point was quantitated and the mean values were plotted with S.D. (n = 2 technical replicates). The result shown is a representative of two independent experiments that were conducted using the same transformants (i.e. two technical replicates).

Figure 2

Figure 2. Photo-crosslinking of the the β-barrel forming domain of LptD with BepA

(**A, B**) *In vivo* photo-crosslinking between LptD and BepA. Cells of RM2243 (*bepA(E137Q)*) carrying pEVOL-pBpF, pMW118-*bepA (E137Q, mut)* and pRM294-*lptD(amb)-his₁₀* plasmids were grown at 30°C in L medium containing 0.02% and 0.5 mM *p*BPA until early log phase and induced with 1 mM IPTG for 3 h to express the indicated LptD(*p*BPA) variants. The cultures were then divided into two portions, each of which was UV-irradiated for 10 min at 4°C. Total cellular proteins were acid-precipitated and analyzed by 7.5% Laemmli SDS-PAGE and immunoblotting with the indicated antibodies. Most of the LptD mutants were accumulated in comparable amounts. LptD-His₁₀xBepA crosslinked products were not detectable with an anti-His antibody due to its low reactivity to LptD-His₁₀ in this and the following experiments. Amino acid residues shown in red and blue indicate the ones whose side chain is pointing inward and outward, respectively. Amino acid residues shown in green indicate the ones located in the loop regions. The result shown is a representative of two technical replicates. (**C**) Summary of the BepA crosslinked positions in LptD. Positions where the crosslinking with BepA was clearly and reproducibly detected are indicated by colored dots. (**D**) Mapping of the BepA crosslinked positions on the barrel domain of LptD in the *E. coli* LptD–LptE structure (PDB code: 4RHB). LptD and LptE are shown in gray and light green, respectively. The N-terminal strand and the β-signal (C-terminal region) in the LptD β-barrel domain are shown in magenta and light blue, respectively. The top view of the LptD/E structure from extracellular space (*upper*) and the side view of the N-terminal region of LptD β-domain (*lower*) are shown. The positions where the crosslinking with BepA was observed were indicated by spheres colored as above.

Figure 3

Figure 3. BepA edge-strand directly contacts with the Tyr-331 residue in the N-terminal half region of the LptD β-barrel forming domain

(**A**) Effect of the BepA edge-strand mutations on the crosslinking between BepA and the LptD derivatives having *p*BPA in the N-terminal half region of the LptD b-barrel-forming domain. Cells of SN56 (Δ*bepA*) carrying pEVOL-pBpF, pMW118-*bepA(E137Q, mut)* and pRM294-*lptD(amb)-his$_{10}$* were grown, induced to express a BepA and a LptD*p*BPA derivative and subjected to photo-crosslinking analysis as described in Figure 2. (**B**) Disulfide crosslinking between the Cys residues in the edge-strand of BepA and the N-terminal half region of the LptD β-barrel-forming domain. Cells of SN56 (Δ*bepA*) carrying a combination of plasmids encoding WT or a Cys-introduced mutant of BepA and LptD-His$_{10}$ as indicated were grown in L-medium and induced with 1 mM IPTG for 3 h to express BepA (Cys) and LptD(Cys)-His$_{10}$. Total cellular proteins were acid-precipitated, solubilized with SDS-buffer containing NEM (for blocking free thiol groups) and subjected to pull-down with Ni-NTA agarose. The purified proteins were treated with or without 2-mercaptoethanol (ME) and analyzed by 7.5% Laemmli SDS-PAGE and immunoblotting with the indicated antibodies. The result shown is a representative of two technical replicates.

Figure 4

Figure 4. BepA interacts with a LptD intermediate assembling on the BAM complex

(**A, B**) *In vivo* photo-crosslinking of an LptD mutant having *p*BPA at two positions with BepA. Cells of RM2243 (*bepA(E137Q)*) carrying pEVOL-pBpF, pMW118‑*bepA(E137Q)* and pRM294-*lptD(amb)-his*$_{10}$ plasmids were grown at 30˚C in L medium containing 0.5 mM *p*BPA until early log phase and induced with 1 mM IPTG for 3 h to express the indicated LptD(*p*BPA) variants. The cultures were divided into two portions, each of which was treated with or without UV-irradiation for 30 min at 4˚C. Proteins of the total membrane fractions were subjected to pull-down with Ni-NTA agarose. Purified proteins were analyzed by 7.5% Laemmli SDS‑PAGE by immunoblotting with the indicated antibodies. (**C**) Simultaneous crosslinking of LptD having Y331*p*BPA and E733C with the BepA edge-strand and the seam region of BamA(S439C). Cells of RM3655 (*bamA(S439C)*, *ΔbepA*)/pEVOL-pBpF/pMW118‑*bepA(E137Q)* carrying pRM294-*lptD(E733C)-his*$_{10}$ or pRM294-*lptD(Y331amb, E733C)-his*$_{10}$ were grown and induced as in A. After treatment with or without BMB and the following quenching of BMB by addition of excess cysteine, the cultures were divided into two portions, each of which was treated with or without UV-irradiation for 10 min at 4°C. Total cellular proteins were acid-precipitated, solubilized with SDS-buffer containing NEM and subjected to pull-down with Ni-NTA agarose. Purified proteins were analyzed by 7.5% Laemmli SDS‑PAGE and immunoblotting with the indicated antibodies. The anti-BamA immunoblotting showed that the amount of the BepAxLptDxBamA crosslinked product was much lower than that of the LptDxBamA crosslinked product. Although the anti-BepA antibodies apparently cross-reacted weakly with the LptDxBamA crosslinked products (asterisks), the higher signal intensity of the BepAxLptDxBamA crosslinked product band as compared with the intensity of the LptDxBamA band indicate that the detection of the former band with the anti-BepA antibodies cannot be ascribed to this cross-reactivity. In **A–C**, we confirmed that the amounts of the isolated non-crosslinked LptD-His$_{10}$ derivatives was roughly equal by CBB staining or anti-His immunoblotting (Figure 4-figure supplement 3). The result shown is a representative of two technical replicates. (**D**) A schematic cartoon of the interaction of the LptD assembly intermediate with BepA and BamA/D on the BAM complex.

Figure 5

Figure 5. Model for the substrate recognition and discrimination by BepA
(**A**) A schematic cartoon of the substrate recognition by BepA at its active site. See the text for details. (**B**) An overview of the proposed LptD assembly process and BepA-mediated discrimination of the assembling and stalled LptD species. See the text for details.

List of Figure supplements.

Figure 1-figure supplement 1. Sequence alignment of BepA homologs and *E. coli* M48 family peptidases

Figure 1-figure supplement 2. Effects of the BepA edge strand Pro mutations on the BamA degradation and the BepA self-cleavage

Figure 1-figure supplement 3. Functionality of the BepA derivatives having *p*BPA in the edge strand

Figure 1-figure supplement 4. The chaperon-like and proteolytic activities of the BepA edge strand mutants

Figure 2-figure supplement 1. Over-expressed LptD molecules mainly accumulate as LptD^C

Figure 2-figure supplement 2. Complementation activity of the LptD(*p*BPA) derivatives

Figure 3-figure supplement 1. Effects of the BepA edge strand mutations on the crosslinking of the BepA edge strand to LptD

Figure 3-figure supplement 2. Disulfide crosslinking between BepA and LptD

Figure 4-figure supplement 1. Complementation activity of the LptD derivatives having *p*BPA at one or two positions

Figure 4-figure supplement 3. The amount of affinity purified LptD-His$_{10}$ derivatives

Figure 5-figure supplement 1. α6 and α9-loop regions shield the intramolecular active stie and edge strand of BepA

List of Supplementary files

Supplementary file 1: Table S1. Strains used in this study

Supplementary file 2: Table S2. Plasmids used in this study

A

Edge strand

```
E. coli          90 VKTPFHFFLINNDEINAFAFFGGNVVLH 117
K. pneumoniae    90 VRTPFHFYLINNDQINAFAFFGGNVVLH 117
X. nematophila   90 VKTPFHFYLINNPNINAYAFFGGNVMLH 117
Y. pestis        88 VRTPFHFYLVNNDQINAFAFFGGNVVLH 115
V. cholerae      87 VKTPFTFFMIRDRNINAFAFFGGYIALH 114
A. salmonicida   87 VRFPFTFFLVNDPSINAAAFLGGRVKVH 114
P. aeruginosa    81 QDHRLAFILIRDAQINAFAAPGGVIGVN 108
L. parisiensis   76 IKKP-DFFIVKSNEINAFAGPGGHIGIN 102

                    * ::.. .*** *  ** : :
```

B

Edge strand

```
BepA: 90 VKTPFHFFLINNDEINAFAFFG----GNVVLH 117
HtpX: 90 GIAMPQVAIYHAPDINAFATGARRDASLVAVS 121
LoiP: 85 NGQPVNYKVYMAKDVNAFAMAN----GCIRVY 112
YcaL: 89 NGTPVNYKVYMTSDVNAWAMAN----GCVRVY 116
            : :    ::**:*       . : :
```

Figure 1-figure supplement 1. Sequence alignment of BepA homologs and *E. coli* M48 family peptidases
(**A**) Alignment of the edge strand amino acid sequences of BepA homologs of *E. coli*, *Klebsiella pneumoniae*, *Xenorhabdus nematophila*, *Yersinia pestis*, *Vibrio cholerae*, *Aeromonas salmonicida*, *Pseudomonas aeruginosa* and *Legionella parisiensis*. (**B**) Alignment of the edge strand amino acid sequences of BepA and other *E. coli* M48 family peptidase (HtpX, LoiP, and YcaL). The alignments were conducted by using the Clustal Omega program. Conserved residues are marked in red.

Figure 1-figure supplement 2. Effects of the BepA edge strand Pro mutations on the BamA degradation and the BepA self-cleavage

(**A**) Degradation of BamA in a Δ*surA* strain expressing the BepA edge strand Pro mutants. Cells of SN259 (Δ*bepA*, Δ*surA*) carrying either pSTD689 or pSTD689-*bepA* plasmids were grown at 30°C in M9-based medium supplemented with 1 mM IPTG for 4 h. Total cellular proteins were acid-precipitated and analyzed by 7.5 or 10% Laemmli SDS-PAGE and immunoblotting with the indicated antibodies. (**B**) C-terminal self-cleavages of the BepA edge strand mutants. Cells of SN56 (Δ*bepA*) carrying either pUC18 or pUC18-*bepA*-*his*$_{10}$ plasmids were grown at 30°C in L medium until early log phase and induced with 1 mM IPTG for 1 h. Total cellular proteins were acid-precipitated and analyzed by 7.5% Laemmli SDS-PAGE and immunoblotting with an anti-BepA antibody. The result shown is a representative of two independent experiments that were conducted using the same transformants (i.e. two technical replicates).

Figure 1-figure supplement 3

A

B

Figure 1-figure supplement 3. Functionality of the BepA derivatives having *p*BPA in the edge strand

(**A**) Complementation activity of the BepA(*p*BPA) derivatives. Cells from 30 μL of overnight cultures of SN56 (Δ*bepA*) carrying pEVOL-pBpF and either pUC18 or pUC18-*bepA(amb)-his₁₀* plasmids were washed, suspended in saline, and serially diluted with saline (to about 10^9 cells/mL). 2.5 μL each of the diluted cells were spotted on L-agar plate or L-agar plates supplemented with 20 μg/mL erythromycin (EM) and 1 mM IPTG with or without 0.5 mM *p*BPA. Plates were incubated at 30°C for 22 h. (**B**) Self-cleavage of the BepA*p*BPA derivatives. Cells of the same strains in *A* were grown at 30°C in L medium supplemented with 0.02% arabinose and 0.5 mM *p*BPA until early log phase and induced with 1 mM IPTG for 1 h. Total cellular proteins were acid-precipitated and analyzed by 7.5% Laemmli SDS-PAGE and immunoblotting with an anti-BepA antibody. The result shown is a representative of two technical replicates.

Figure 1-figure supplement 4

Figure 1-figure supplement 4. The chaperone-like and proteolytic activities of the BepA edge strand mutants
(**A**, **B**) The accumulation of the BepA mutants and LptD^C in the Δ*bepA* cells expressing the BepA mutants. Cells of AD16 (*bepA*$^+$) carrying pSTD689 (vector) or SN56 (Δ*bepA*) carrying either pSTD689 or pSTD689-*bepA* plasmids were grown at 30°C in L medium until early log phase and induced with 1 mM IPTG for 1 h. Total cellular proteins were acid-precipitated, and analyzed by 10 or 7.5% Laemmli SDS-PAGE under a reducing (+ME) or non-reducing (no ME) condition and immunoblotting with anti-BepA (*upper* panels) or anti-LptD (*lower* panel) antibodies. (**C**) Protease activities of the BepA edge strand Cys mutants against overproduced LptD. Cells of SN56 (Δ*bepA*) carrying pTWV228-*lptD-his*$_{10}$ and either pSTD689 or pSTD689-*bepA* plasmids were grown at 30°C in L-medium until early log phase and induced with 1 mM IPTG for 1 h. Total cellular proteins were analyzed as in Figure 1B. The result shown is a representative of two technical replicates.

Figure 2-figure supplement 1. Over-expressed LptD molecules mainly accumulate as LptD^C

Cells of RM2243(*bepA(E137Q)*)/pEVOL-pBpF carrying either pMW118 or pMW118-*bepA(E137Q)* and either pRM294 or pRM294-*lptD-his$_{10}$* were grown n L medium containing 0.02% arabinose and 0.5 mM *p*BPA until early log phase at 30°C and induced with 1 mM IPTG for 3 h. Total cellular proteins were acid-precipitated and analyzed by 7.5 or 10% Laemmli SDS-PAGE under a reducing (+ME) or non-reducing (no ME) condition and immunoblotting with the indicated antibodies. The result shown is a representative of two biological replicates.

Figure 2-figure supplement 2

Figure 2-figure supplement 2. Complementation activity of the LptD(*p*BPA) derivatives
Cells of RM3588 (P*ara*-*lptD*) carrying pEVOL-pBpF and either pRM294 or pRM294-*lptD(amb)-his₁₀* plasmids were grown at 30˚C in L-medium supplemented with 0.1% arabinose for 2.5 h. Cells were washed, suspended in saline, and serially diluted with saline (to about 10^9 cells/mL). 2.5 μL each of the diluted cells were spotted on L-agar plate containing 0.1% arabinose or L medium-based agar plates supplemented with 1 mM IPTG with or without 0.5 mM *p*BPA. Plates were incubated at 30˚C for 22 h. The result shown is a representative of two technical replicates.

Figure 3-figure supplement 1

Figure 3-figure supplement 1. Effects of the BepA edge strand mutations on the crosslinking of the BepA edge strand to LptD

Cells SN56 (Δ*bepA*) carrying pEVOL-pBpF and pUC18-*bepA(E137Q, mut., amb)-his₁₀* plasmids were grown in L medium containing 0.02% arabinose and 0.5 mM *p*BPA until early log phase at 30°C and induced with 1 mM IPTG for 1 h to express the indicated BepA (*p*BPA) variants. The cultures were divided into two portions, each of which was treated with or without UV-irradiation for 10 min at 4°C. Proteins of the total membrane fractions were subjected to pull-down with Ni-NTA agarose. Purified proteins were analyzed by 7.5% Laemmli SDS-PAGE and immunoblotting with the indicated antibodies. For analysis of the cellular accumulation of each BepA derivative, total cellular proteins of cells of the same cultures were acid-precipitated, and analyzed by 10% Laemmli SDS-PAGE and immunoblotting with the indicated antibodies. Open triangles indicate unknown crosslinked products. The result shown is a representative of two technical replicates.

Figure 3-figure supplement 2

A

B

Figure 3-figure supplement 2. Disulfide crosslinking between BepA and LptD

(**A**) Complementation activity of the LptD derivatives with an engineered Cys residue. Cells of RM3588 (P_{ara}-*lptD*) carrying pRM294 or pRM294-*lptD(Cys)-his$_{10}$* plasmids were grown at 30°C in L medium supplemented with 0.1% arabinose for 2.5 h. Cells were washed, suspended in saline, and serially diluted with saline (to about 10^9 cells/mL). 2.5 μL each of the diluted cells were spotted on L-agar plate containing 0.1% arabinose or 1 mM IPTG. Plates were incubated at 30°C for 22 h. (**B**) BepA–LptD disulfide crosslinking. Cells of SN56 (Δ*bepA*) carrying a combination of plasmids encoding WT or a Cys-mutant form of BepA and LptD-His$_{10}$ as indicated were grown in L-medium and induced with 1 mM IPTG for 3 h to express BepA(Cys) and LptD(Cys)-His$_{10}$. Total cellular proteins were acid-precipitated, solubilized with SDS buffer containing NEM and subjected to pull-down with Ni-NTA agarose. The purified proteins were treated with or without 2-mercaptoethanol (ME) and analyzed by 7.5% Laemmli SDS-PAGE and immunoblotting with the indicated antibodies. The result shown is a representative of two technical replicates.

Figure 4-figure supplement 1

Figure 4-figure supplement 1. Complementation activity of the LptD derivatives having *p*BPA at one or two positions

Cells of RM3588 (P*ara*-*lptD*) carrying pEVOL-pBpF and either pRM294 or pRM294-*lptD(amb)-his10* plasmids were grown and analyzed as in Figure 2-figure supplement 2. The result shown is a representative of two technical replicates.

Figure 4-figure supplement 2. BMB crosslinking between LptD(Y331*p*BPA/E733C) and BamA(S439C)
(**A**) BMB crosslinking. Cells of RM3654 (*bamA⁺*, Δ*bepA*)/pEVOL-pBpF/pMW118-*bepA(E137Q)* or RM3655 (*bamA(S439C)*, Δ*bepA*)/ pEVOL-pBpF/pMW118-*bepA(E137Q)* carrying each pRM294-*lptD-his₁₀* derivative were grown at 30˚C in L medium containing 0.5 mM *p*BPA until early log phase and induced with 1 mM IPTG for 3 h to express the indicated LptD variants. Cell cultures were treated with TCEP to reduce possible artificially-formed disulfide bonds for E733C in LptD and S439C in BamA. The cultures were divided into two portions, each of which was treated with or without BMB. After quenching of BMB by addition of excess cysteine, total cellular proteins were acid-precipitated, solubilized with SDS-buffer containing NEM and subjected to pull-down with Ni-NTA agarose. The purified proteins were analyzed by 7.5% Laemmli SDS-PAGE and immunoblotting with the indicated antibodies. (**B**) Complementation activity of LptD derivatives. Cells of RM3588 (P*ₐᵣₐ*-*lptD*) carrying pEVOL-pBpF and pRM294 or each pRM294-*lptD-his₁₀* derivative were grown are analyzed as in Figure 2-figure supplement 2. The result shown is a representative of two technical replicates.

Figure 4-figure supplement 3. The amount of affinity purified LptD-His$_{10}$ derivatives
Affinity isolated proteins in Figure 4A-C were separated by 7.5% Laemmli SDS-PAGE and subjected to CBB staining (**A** and **B** for Figure 4A and B, respectively) or anti-His immunoblotting (**C** for Figure 4C). The result shown is a representative of two technical replicates.

Figure 5-figure supplement 1

Figure 5-figure supplement 1. α6 and α9-loop regions shield the intramolecular active stie and edge strand of BepA
The space-filling models of wild type BepA (*left*), BepA without the α6-loop (*middle*) and BepA without the α6- and α9-loops (*right*).
The peptidase and the TPR domains of the BepA crystal structure (PDB code: 6AIT) are shown in gray and orange, respectively. The α6-loop, the α9-loop, the proteolytic active site (the HExxH motif and the third zinc ligand, Glu-201) and the edge strand in the peptidase domain are shown in magenta, green, blue and red, respectively.

Table S1. Strains used in this study

Strain	Genotype	References
AD16	Δ*pro-lac thi*/F' *lacI^q Z*ΔM15 Y⁺ pro⁺	(Kihara et al., 1995)
SN56	AD16, Δ*bepA*	(Narita et al., 2013)
SN259	AD16, Δ*bepA* Δ*surA*::*kan*	(Narita et al., 2013)
RM2243	AD16, *bepA(E137Q) purC80*::Tn*10*	This study
RM3654	AD16, Δ*bepA bamA*⁺ *zae502*::Tn*10*	This study
RM3655	AD16, Δ*bepA bamA(S439C) zae502*::Tn*10*	This study
JE6631	*Hfr:str thi polA1*	(Mori et al., 2002)
RM2091	JE6631, *purC80*::Tn*10*	This study
YH188	JE6631, *zae502*::Tn*10*	(Hizukuri and Akiyama, 2012)
MC4100	F⁻ *araD139* Δ*(argF-lac)U169 rpsL150 relA1 flbB5301 deoC1 ptsF25 rbsR*	(Casadaban, 1976)
CU141	MC4100/F' *lacI^q lacZ*⁺, Y⁺, A⁺	(Akiyama et al., 1994)
HM1742	CU141 *ara*⁺	(Mori and Ito, 2006)
RM3588	HM1742, *kan araC*-P*araBAD*-*lptD*	This study
DY330	W3110, Δ*lacU169 gal490 λcI857* Δ(*cro-bioA*)	(Yu et al., 2000)
RM3444	DY330, *kan araC*-P*araBAD*-*lptD*	This study

(References)

Akiyama Y, Ogura T, Ito K. 1994. Involvement of FtsH in protein assembly into and through the membrane. I. Mutations that reduce retention efficiency of a cytoplasmic reporter. *J Biol Chem* **269**:5218–5224. doi:10.1016/S0021-9258(17)37677-9

Casadaban MJ. 1976. Transposition and fusion of the *lac* genes to selected promoters in *Escherichia coli* using bacteriophage lambda and Mu. *J Mol Biol* **104**:541–555. doi:10.1016/0022-2836(76)90119-4

Hizukuri Y, Akiyama Y. 2012. PDZ domains of RseP are not essential for sequential cleavage of RseA or stress-induced σ^E activation *in vivo*. *Mol Microbiol* **86**:1232–1245. doi:10.1111/mmi.12053

Kihara A, Akiyama Y, Ito K. 1995. FtsH is required for proteolytic elimination of uncomplexed forms of SecY, an essential protein translocase subunit. *Proc Natl Acad Sci U S A* **92**:4532–4536. doi:10.1073/pnas.92.10.4532

Mori H, Ito K. 2006. The long α-helix of SecA is important for the ATPase coupling of translocation. *J Biol Chem* **281**:36249–36256. doi:10.1074/jbc.M606906200

Mori H, Shimizu Y, Ito K. 2002. Superactive SecY variants that fulfill the essential translocation function with a reduced cellular quantity. *J Biol Chem* **277**:48550–48557. doi:10.1074/jbc.M204436200

Narita S, Masui C, Suzuki T, Dohmae N, Akiyama Y. 2013. Protease homolog BepA (YfgC) promotes assembly and degradation of β-barrel membrane proteins in *Escherichia coli*. *Proc Natl Acad Sci U S A* **110**:E3612–E3621. doi:10.1073/pnas.1312012110

Yu D, Ellis HM, Lee EC, Jenkins NA, Copeland NG, Court DL. 2000. An efficient recombination system for chromosome engineering in *Escherichia coli*. *Proc Natl Acad Sci U S A* **97**:5978–5983. doi:10.1073/pnas.100127597

Table S2. Plasmids used in this study

Plasmid	Vector	Encoded protein and description	Reference or source
pEVOL-pBpF		p15A-derivative encoding an evolved *M. jannaschii* aminoacyl-tRNA synthetase/ suppressor tRNA pair for incorporation of *p*BPA; CmR	(Young et al., 2010)
pSTD689		pACYC184-derived expression vector; P$_{lac}$, Spc^R	(Kanehara et al., 2003)
pRM290	pSTD689	BepA	(Daimon et al., 2017)
pRM291	pSTD689	BepA(E137Q)	(Daimon et al., 2017)
pRM512	pSTD689	BepA(N105P)	This study
pRM513	pSTD689	BepA(A106P)	This study
pRM514	pSTD689	BepA(F107P)	This study
pRM515	pSTD689	BepA(A108P)	This study
pRM516	pSTD689	BepA(F109P)	This study
pRM517	pSTD689	BepA(F110P)	This study
pRM833	pSTD689	BepA(N105C)	This study
pRM786	pSTD689	BepA(N105C, E137Q)	This study
pRM834	pSTD689	BepA(A106C)	This study
pRM787	pSTD689	BepA(A106C, E137Q)	This study
pRM835	pSTD689	BepA(F107C)	This study
pRM788	pSTD689	BepA(F107C, E137Q)	This study
pUC18		Expression vector; P$_{lac}$, Amp^R	Takara Bio
pUC-bepA(E137Q)	pUC18	BepA(E137Q)	(Narita et al., 2013)
pUC-bepA-his$_{10}$	pUC18	BepA-His$_{10}$	(Narita et al., 2013)
pUC-bepA(E137Q)-his$_{10}$	pUC18	BepA(E137Q)-His$_{10}$	(Narita et al., 2013)
pYN333	pUC18	BepA(N105P)-His$_{10}$	This study
pYN334	pUC18	BepA(A106P)-His$_{10}$	This study
pYN335	pUC18	BepA(F107P)-His$_{10}$	This study
pYN336	pUC18	BepA(A108P)-His$_{10}$	This study
pYN337	pUC18	BepA(F109P)-His$_{10}$	This study
pYN338	pUC18	BepA(F110P)-His$_{10}$	This study
pYN339	pUC18	BepA(N105*amb*)-His$_{10}$	This study
pRM536	pUC18	BepA(N105*amb*, E137Q)-His$_{10}$	This study
pRM778	pUC18	BepA(N105*amb*, A106P, E137Q)-His$_{10}$	This study
pRM779	pUC18	BepA(N105*amb*, F107P, E137Q)-His$_{10}$	This study
pYN340	pUC18	BepA(A106*amb*)-His$_{10}$	This study
pRM537	pUC18	BepA(A106*amb*, E137Q)-His$_{10}$	This study
pYN345	pUC18	BepA(A106*amb*, F107P, E137Q)-His$_{10}$	This study
pYN341	pUC18	BepA(F107*amb*)-His$_{10}$	This study
pRM538	pUC18	BepA(F107*amb*, E137Q)-His$_{10}$	This study

pYN346	pUC18	BepA(F107*amb*, A106P, E137Q)-His$_{10}$	This study
pYN342	pUC18	BepA(A108*amb*)-His$_{10}$	This study
pRM539	pUC18	BepA(A108*amb*, E137Q)-His$_{10}$	This study
pYN343	pUC18	BepA(F109*amb*)-His$_{10}$	This study
pRM540	pUC18	BepA(F109*amb*, E137Q)-His$_{10}$	This study
pYN344	pUC18	BepA(F110*amb*)-His$_{10}$	This study
pRM541	pUC18	BepA(F110*amb*, E137Q)-His$_{10}$	This study
pMW118		pSC101-derived expression vector; P$_{lac}$, Amp^{R}	Nippon Gene
pNB91	pMW118	BepA(E137Q)	This study
pRM807	pMW118	BepA(A106P, E137Q)	This study
pRM808	pMW118	BepA(F107P, E137Q)	This study
pTWV228		pBR322-derived expression vector; P$_{lac}$, Amp^{R}	Takara Bio
pRM267	pTWV228	LptD-His$_{10}$ (using the native SD sequence)	(Daimon et al., 2017)
pRM816	pTWV228	LptD-His$_{10}$^{SD} (using a strong SD sequence)	This study
pRM810	pTWV228	LptD(M287C)-His$_{10}$^{SD}	This study
pRM811	pTWV228	LptD(Y331C)-His$_{10}$^{SD}	This study
pRM812	pTWV228	LptD(E391C)-His$_{10}$^{SD}	This study
pRM813	pTWV228	LptD(Q393C)-His$_{10}$^{SD}	This study
pRM814	pTWV228	LptD(H416C)-His$_{10}$^{SD}	This study
pRM815	pTWV228	LptD(V430C)-His$_{10}$^{SD}	This study
pRM294		pBR322-derived expression vector; P$_{lac}$, Spc^{R}	(Miyazaki et al., 2018)
pRM309	pRM294	LptD-His$_{10}$^{SD} (using a strong SD sequence)	(Miyazaki et al., 2018)
pNB22	pRM294	LptD(R40*amb*)-His$_{10}$^{SD}	This study
pNB23	pRM294	LptD(N55*amb*)-His$_{10}$^{SD}	This study
pNB24	pRM294	LptD(T70*amb*)-His$_{10}$^{SD}	This study
pNB25	pRM294	LptD(D85*amb*)-His$_{10}$^{SD}	This study
pNB26	pRM294	LptD(P100*amb*)-His$_{10}$^{SD}	This study
pNB27	pRM294	LptD(N115*amb*)-His$_{10}$^{SD}	This study
pNB28	pRM294	LptD(T130*amb*)-His$_{10}$^{SD}	This study
pNB29	pRM294	LptD(R145*amb*)-His$_{10}$^{SD}	This study
pNB30	pRM294	LptD(N160*amb*)-His$_{10}$^{SD}	This study
pNB31	pRM294	LptD(P175*amb*)-His$_{10}$^{SD}	This study
pNB49	pRM294	LptD(D190*amb*)-His$_{10}$^{SD}	This study
pNB50	pRM294	LptD(V205*amb*)-His$_{10}$^{SD}	This study
pNB51	pRM294	LptD(V220*amb*)-His$_{10}$^{SD}	This study
pNB52	pRM294	LptD(Y235*amb*)-His$_{10}$^{SD}	This study
pNB53	pRM294	LptD(N250*amb*)-His$_{10}$^{SD}	This study
pRM618	pRM294	LptD(T258*amb*)-His$_{10}$^{SD}	This study
pRM619	pRM294	LptD(I259*amb*)-His$_{10}$^{SD}	This study

pNB54	pRM294	LptD(H265amb)-His$_{10}$^{SD}	This study
pRM620	pRM294	LptD(E273amb)-His$_{10}$^{SD}	This study
pNB55	pRM294	LptD(S280amb)-His$_{10}$^{SD}	This study
pRM621	pRM294	LptD(L286amb)-His$_{10}$^{SD}	This study
pRM622	pRM294	LptD(M287amb)-His$_{10}$^{SD}	This study
pNB56	pRM294	LptD(S295amb)-His$_{10}$^{SD}	This study
pNB57	pRM294	LptD(R310amb)-His$_{10}$^{SD}	This study
pRM623	pRM294	LptD(Y314amb)-His$_{10}$^{SD}	This study
pNB58	pRM294	LptD(W325amb)-His$_{10}$^{SD}	This study
pRM624	pRM294	LptD(F327amb)-His$_{10}$^{SD}	This study
pRM625	pRM294	LptD(D330amb)-His$_{10}$^{SD}	This study
pRM626	pRM294	LptD(Y331amb)-His$_{10}$^{SD}	This study
pNB59	pRM294	LptD(F340amb)-His$_{10}$^{SD}	This study
pNB60	pRM294	LptD(A355amb)-His$_{10}$^{SD}	This study
pRM627	pRM294	LptD(K358amb)-His$_{10}$^{SD}	This study
pNB61	pRM294	LptD(A370amb)-His$_{10}$^{SD}	This study
pRM628	pRM294	LptD(T371amb)-His$_{10}$^{SD}	This study
pNB62	pRM294	LptD(T385amb)-His$_{10}$^{SD}	This study
pRM629	pRM294	LptD(E391amb)-His$_{10}$^{SD}	This study
pRM630	pRM294	LptD(Q393amb)-His$_{10}$^{SD}	This study
pNB63	pRM294	LptD(Q400amb)-His$_{10}$^{SD}	This study
pNB64	pRM294	LptD(V415amb)-His$_{10}$^{SD}	This study
pRM631	pRM294	LptD(H416amb)-His$_{10}$^{SD}	This study
pRM632	pRM294	LptD(R429amb)-His$_{10}$^{SD}	This study
pNB65	pRM294	LptD(V430amb)-His$_{10}$^{SD}	This study
pRM633	pRM294	LptD(H431amb)-His$_{10}$^{SD}	This study
pNB66	pRM294	LptD(G445amb)-His$_{10}$^{SD}	This study
pRM634	pRM294	LptD(H457amb)-His$_{10}$^{SD}	This study
pRM635	pRM294	LptD(Y458amb)-His$_{10}$^{SD}	This study
pNB67	pRM294	LptD(Q460amb)-His$_{10}$^{SD}	This study
pNB68	pRM294	LptD(D475amb)-His$_{10}$^{SD}	This study
pRM636	pRM294	LptD(V478amb)-His$_{10}$^{SD}	This study
pRM637	pRM294	LptD(N479amb)-His$_{10}$^{SD}	This study
pNB69	pRM294	LptD(K490amb)-His$_{10}$^{SD}	This study
pNB70	pRM294	LptD(T505amb)-His$_{10}$^{SD}	This study
pNB71	pRM294	LptD(R520amb)-His$_{10}$^{SD}	This study
pNB72	pRM294	LptD(S535amb)-His$_{10}$^{SD}	This study
pNB73	pRM294	LptD(D550amb)-His$_{10}$^{SD}	This study
pNB74	pRM294	LptD(R565amb)-His$_{10}$^{SD}	This study
pNB75	pRM294	LptD(G580amb)-His$_{10}$^{SD}	This study

pNB76	pRM294	LptD(I595amb)-His$_{10}^{SD}$	This study
pNB77	pRM294	LptD(G610amb)-His$_{10}^{SD}$	This study
pNB78	pRM294	LptD(G625amb)-His$_{10}^{SD}$	This study
pNB79	pRM294	LptD(S640amb)-His$_{10}^{SD}$	This study
pNB80	pRM294	LptD(N655amb)-His$_{10}^{SD}$	This study
pNB81	pRM294	LptD(K670amb)-His$_{10}^{SD}$	This study
pNB82	pRM294	LptD(V685amb)-His$_{10}^{SD}$	This study
pNB83	pRM294	LptD(V700amb)-His$_{10}^{SD}$	This study
pNB84	pRM294	LptD(S715amb)-His$_{10}^{SD}$	This study
pNB32	pRM294	LptD(Y726amb)-His$_{10}^{SD}$	This study
pNB85	pRM294	LptD(V730amb)-His$_{10}^{SD}$	This study
pNB86	pRM294	LptD(H745amb)-His$_{10}^{SD}$	This study
pRM809	pRM294	LptD(D749amb)-His$_{10}^{SD}$	This study
pNB87	pRM294	LptD(G760amb)-His$_{10}^{SD}$	This study
pNB88	pRM294	LptD(S775amb)-His$_{10}^{SD}$	This study
pRM821	pRM294	LptD(Y331amb, D749amb)-His$_{10}^{SD}$	This study
pRM822	pRM294	LptD(Y331amb, Y726amb)-His$_{10}^{SD}$	This study
pRM829	pRM294	LptD(E733C)-His$_{10}^{SD}$	This study
pRM831	pRM294	LptD(Y331amb, E733C)-His$_{10}^{SD}$	This study
pUC118		Expression vector; P$_{lac}$, Amp^R	Takara Bio
pRM320	pUC118	$bepA$-$yfgD$	This study
pRM324	pUC118	$bepA(E137Q)$-$yfgD$	This study
pRM823	pUC118	$bamA$	This study
pRM836	pUC118	$bamA(S439C)$	This study
pK18mobsacB		pBR322-derived vector carrying $sacB$ for chromosomal gene replacement; Km^R	(Schäfer et al., 1994)
pRM330	pK18mobsacB	$bepA(E137Q)$-$yfgD$	This study
pRM845	pK18mobsacB	$bamA(S439C)$	This study

(References)

Daimon Y, Iwama-Masui C, Tanaka Y, Shiota T, Suzuki T, Miyazaki R, Sakurada H, Lithgow T, Dohmae N, Mori H, Tsukazaki T, Narita S-I, Akiyama Y. 2017. The TPR domain of BepA is required for productive interaction with substrate proteins and the β-barrel assembly machinery complex. *Mol Microbiol* **106**:760–776. doi:10.1111/mmi.13844

Kanehara K, Ito K, Akiyama Y. 2003. YaeL proteolysis of RseA is controlled by the PDZ domain of YaeL and a Gln-rich region of RseA. *EMBO J* **22**:6389–6398. doi:10.1093/emboj/cdg602

Miyazaki R, Myougo N, Mori H, Akiyama Y. 2018. A photo-cross-linking approach to monitor folding and assembly of newly synthesized proteins in a living cell. *J Biol Chem* **293**:677–686. doi:10.1074/jbc.M117.817270

Narita S, Masui C, Suzuki T, Dohmae N, Akiyama Y. 2013. Protease homolog BepA (YfgC) promotes assembly and degradation of β-barrel membrane proteins in *Escherichia coli*. *Proc Natl Acad Sci U S A* **110**:E3612–E3621. doi:10.1073/pnas.1312012110

Schäfer A, Tauch A, Jäger W, Kalinowski J, Thierbach G, Pühler A. 1994. Small mobilizable multi-purpose cloning vectors derived from the *Escherichia coli* plasmids pK18 and pK19: selection of defined deletions in the chromosome of *Corynebacterium glutamicum*. *Gene* **145**:69–73. doi:10.1016/0378-1119(94)90324-7

Young TS, Ahmad I, Yin JA, Schultz PG. 2010. An enhanced system for unnatural amino acid mutagenesis in *E. coli*. *J Mol Biol* **395**:361–374. doi:10.1016/j.jmb.2009.10.030