Shared and unique properties of place cells in anterior cingulate cortex and hippocampus

Ayaka Bota[1, 2, 3], Akihiro Goto[2, 3], Suzune Tsukamoto[2], Alexander Schmidt[4,5,6,7], Fred Wolf[4,5,6,7], Alessandro Luchetti[3], Junichi Nakai[1, 8], Hajime Hirase[3, 8], and Yasunori Hayashi[1, 2, 3, 8]

1. Graduate School of Science and Engineering, Saitama University, Saitama 338-8570, Japan
2. Department of Pharmacology, Kyoto University Graduate School of Medicine, Kyoto 606-8501, Japan
3. Brain Science Institute, RIKEN, Wako, Saitama 351-0198, Japan
4. Max Planck Institute for Dynamics and Self-Organization, Göttingen 37077, Germany
5. Max Planck Institute for Experimental Medicine, Göttingen 37075, Germany
6. Campus Institute for Dynamics of Biological Networks, Göttingen 37075, Germany
7. Center for Biostructural Imaging of Neurodegeneration, Göttingen 37075, Germany
8. Brain and Body System Science Institute, Saitama University, Saitama 338-8570, Japan

[†] Corresponding author:

Yasunori Hayashi, MD, PhD

E-mail: yhayashi-tky@umin.ac.jp

Keywords: Anterior cingulate cortex, hippocampus, place cells, memory consolidation, Ca^{2+}-imaging

Conflict of interest statement:

YH was partly supported by Fujitsu Laboratories and Dwango.

Present address:

Alessandro Luchetti
University of California, Los Angeles, CA 90095, USA

Hajime Hirase
University of Copenhagen, Center for Translational Neuromedicine, 2200 Copenhagen N., Denmark

Abstract

In the brain, spatial information is represented by neurons that fire when an animal is at specific locations, including place cells in hippocampus and grid cells in entorhinal cortex. But how this information is processed in downstream brain regions still remains elusive. Using chronic Ca^{2+} imaging, we examined the activity of neurons in anterior cingulate cortex (ACC), a brain region implicated in memory consolidation, and found neurons that fire in a manner consistent with the properties of place cells. While the ACC place cells showed stability, location and context specificity similar to the hippocampal counterparts, they also have unique properties. Unlike hippocampal place cells that immediately formed upon exposure to a novel environment, ACC place cells increased over days. Also, ACC place cells tend to have additional place fields whereas typical hippocampal place cells have only one. Hippocampal activity is required for the formation of ACC place cells, but once they are established, hippocampal inactivation did not have any impact on ACC place cell firing. We thus identified features of ACC place cells that carry spatial information in a unique fashion.

Introduction

Spatial navigation is an essential element of animal behavior that allows animals to forage, return home and avoid dangers. The hippocampus plays a crucial role in this process. It bears neurons called place cells that fire when an animal is located in a particular position of an environment but not in others, providing an allocentric cognitive map of a space [1-4]. Multimodal inputs including visual, tactile, olfaction, auditory, proprioception, and other sensory information are integrated to form the place cells in hippocampus [1-4]. It was also shown that the spatial representation of hippocampal place cell is initially dynamic but gradually stabilized [5-8]. The stabilization selectively occurs at a location with motivational (reward) and environmental (landmark) salience and results in an over-representation in these locations, indicating that the place cells' activity is not merely a representation of the space but also its cognitive value. In this way, hippocampal place cells are well characterized from years of studies [1-4]. However, how spatial information is further processed and chronically represented in downstream brain regions has not been fully elucidated at cellular resolution [5].

We decided to focus on anterior cingulate cortex (ACC), a part of medial prefrontal cortex (mPFC) implicated in memory consolidation [9-11]. Suppression of ACC impaired the recall of the remote but not of the recent memory[11]. This is in contrast to hippocampus, which is required for the recall of only recent but not remote memory [11, 12]. ACC receives a direct input from hippocampus [13] and also from striatum, amygdala and retrosplenial cortex, thereby serving as an

integration center of wide variety of sensory and motivational events [14, 15]. Recent work has

revealed neurons with spatially specific firing in mPFC including ACC. This spatial firing is regulated

by the environmental and task context, consistent with the spatial coding [16-19]. Lesion of the

hippocampus abolishes the place code activity in the mPFC indicating that mPFC is situated in

downstream of hippocampus for spatial coding [20]. However, how the spatial encoding of ACC

neurons develops has not been fully examined.

We, therefore, chronically monitored the activity of neurons in ACC while animals navigate

through a space. To this end, we used a head-mount miniaturized fluorescence microscope [5] to

image Ca^{2+}-responses of excitatory neurons in ACC over days from freely moving mice. We found

neurons that fire in a manner consistent with the properties of place cells (ACC place cell). ACC

place cell showed similar properties with hippocampal place cell such as increase in stability,

reliability and context specificity after repeated exposure to the same environment over days.

Bayesian decoding verified that those cells indeed carry positional information. On the other hand,

we found several properties unique to ACC place cells. The fraction of ACC place cells increased as

the task was repeated. ACC neurons tend to have multiple place fields, which might represent

association of more than one location in a context. Finally, the formation of the ACC place cells

requires hippocampal activity but once established, the cells become independent of hippocampus.

Thus, while ACC place cells share properties with hippocampal place cells, they have unique features

and mechanism of generation.

RESULTS

ACC has place cells similarly to hippocampus

For Ca^{2+} imaging, we used TRE-G-CaMP7-T2A-DsRed2 × CaMKIIα-tTA double transgenic

mice, that coexpress G-CaMP7 and DsRed2 in excitatory neurons [8]. To image ACC, a gradient

reflective index (GRIN) lens with a microprism attached at its end was implanted between the two

cortical hemispheres so that the center of the tip of the microprism is at 0.5 mm anterior from bregma

and 1.4 mm from the cortical surface and the prism faces the left ACC [21] (Fig. 1a and

Supplementary Fig. 1a). The dorsal hippocampus was imaged by implanting a GRIN lens above the

right dorsal hippocampus after removing a part of overlaying cortex as previously reported [5] (Fig.

1b and Supplementary Fig. 1b). The activity of ACC layer 2/3 neurons or hippocampal CA1

pyramidal neurons was observed by using a head-mounted fluorescent microscope across multiple

days from awake and behaving animals [5] (Fig. 1c).

The animals were placed first in a square track (one edge 50 cm, context S1) with a wall

installed at one of the corners, then in an open arena (50 x 50 cm, context O) with different wall

patterns and scent, and finally in the original square track (context S1') for 10 min each with 5 min
intermissions (Fig. 1d). In context S1, the animals ran back-and-forth between the two ends for food
rewards, given when the animal reached one end. The animals moved faster and finished more laps
over days (Fig. 1e), indicating that the animal became familiar with the context[11]. In the context O,
the rewards were randomly thrown into the open arena to motivate animals to navigate around [22].

In both hippocampus and ACC, we found neurons that fire in a manner consistent with the
properties known for place cells, such as location specificity, reproducibility, and direction selectivity
across laps (Fig. 1f and g). The proportion of hippocampal place cells did not change over days,
consistent with previous results [5]. The proportion of ACC place cells in early phase (days 1-3) was
smaller than in hippocampus (Fig. 1h) but increased as the task was repeated and reached the level
comparable to hippocampus in the late phase (days 7-9). We also calculated the reliability index of
each cell, an index of how reproducibly a cell fires in a specific location across multiple laps in a
single session (Supplementary Fig. 2). In early sessions (days 1-3), the index was lower in ACC than
in hippocampus, but gradually increased in the late sessions (days 7-9) though it never reached the
level of hippocampus (Fig. 1i). This indicates that the firing reproducibility of ACC place cells is not
as high as that of hippocampus place cells. We noticed that nearly a half of ACC neurons had extra
place fields (Fig. 1j) whereas hippocampal neurons typically had only one place field. The number of
extra place fields remained constant in both ACC and hippocampus during repeated training,
indicating that this is an intrinsic difference between these two brain regions.

In order to test if these cells indeed encode spatial information, we used Bayesian decoding
(Fig. 1k) by selecting the same number of the cells with the highest activity level from both ACC or
hippocampus. The decoder often registered one end of the square track as the other end. This may
reflect the fact that the ends are adjacent to each other, separated just by a wall. Nevertheless, the
analysis confirmed that the ACC neurons carry positional information comparable to those in
hippocampus (hippocampus vs ACC; $p = 0.46$ (early), $p = 0.201$ (middle), $p = 0.79$ (late), One way
ANOVA. Fig.1k). Therefore, we called these cells ACC place cells.

____

Bota et al. Fig. 1

Figure 1. The basic properties of ACC and hippocampal place cells.

(a) Imaging of the ACC using a right angle microprism inserted into the fissure.

(b) Imaging of hippocampus by using a GRIN lens implanted above hippocampal CA1 layer.

(c) Example images of ACC neurons. Left, max intensity image. Blood vessels appear as shadows.
Middle, relative fluorescent change ($\Delta F/F$). Right, active cells (red circles) overlaid with the max
intensity image.

(d) Behavior paradigms. The square linear track (context S1 or S1') and the open arena (context O).
In one session, mice visited two distinct environments (S1$\rightarrow$ O $\rightarrow$S1') each for 10 min. Location of
reward in S1 was indicated by asterisks. In context O, the reward was randomly thrown into the
arena. One set of experiment was conducted per day for 9 days while monitoring neuronal activity
in hippocampus or ACC.

(e) Behavioral changes induced by repeated training with context S1. Total traveled distance is
shown (n = 9 mice). Day 1 vs day 9, p = 1.88 x 10^{-8}, one-way AVOVA. Graphs show means $\pm$
SEM.

(f) Example of place cells in hippocampus in ACC in context S1. Blue lines show the trajectory of
the mouse and red dots mark calcium events.

(g) Example of place cells in hippocampus and in ACC in context O. Left, blue lines show the
mouse's trajectory and red dots mark calcium events. Right, Gaussian-smoothed density maps of
calcium events, normalized by the mouse's occupancy time per unit area and the cell's maximum
response.

(h) The fraction of ACC and hippocampal place cells relative to the number of total identified cells
in context S1. Hippocampus early vs late p = 0.69, ACC early vs late p = 2.9 x 10^{-4}, hippocampus
early vs ACC early p = 5.5 x 10^{-10}, hippocampus late vs ACC late p = 0.079; one-way ANOVA. N
= 36 data (18 sessions x 2 running directions) in early, 46 data (23 sessions x 2 running directions)
in middle, 34 data (17 sessions x 2 running directions) in late from 5 mice for hippocampus. N = 32
data (16 sessions x 2 directions) in early, 38 data (19 sessions x 2 directions) in middle, 32 data (16
sessions x 2 directions) in late from 4 mice for ACC.

(i) Reliability of firing of ACC and hippocampus ('HPC') place cells in the early (days 1-3) and the
late sessions (days 7-9). Reliability represents how reproducible is the during multiple laps in the
same session (Fig. S2). Data were pooled from 3265 cells in the early and 3527 cells in the late
sessions from 5 mice for hippocampus and 1206 cells in the early and 2494 cells in the late sessions
from 4 mice for ACC. Hippocampus early vs late p = 1.61 x 10^{-12}; ACC early vs late p = 7.23 x 10^{-8}
and ACC early vs hippocampus early p = 1.403 x 10^{-114} and ACC late vs hippocampus late p = 3.68
x 10^{-171}, one-way ANOVA.

(j) Example of an ACC place cell with extra place fields in context S1. Left Top, mouse trajectory
(blue) and calcium events (red). Left middle, linearized heat map of the event rate normalized by
maximum activity. Left bottom, Normalized event rate in each spatial bin. Vertical line shows the
threshold for criteria of place field. This cell has 2 place fields. Right, number of place field of
hippocampus and ACC place cells in the early (days 1-3), middle (days 4-6) and late (days 7-9)

sessions. N = 5161 place cells in early, 5018 place cells in middle, 3842 place cells from 5
hippocampus group mice. N = 2453 place cells in early, 2812 place cells in middle, 2972 place cells
from 4 ACC group mice.

(k) Bayesian decoding of the mouse trajectory (red dots) and actual position (black curves) from
hippocampus and ACC place cells. Average median errors of the early (days 1-3), middle (days 4-6)
and late (days 7-9) sessions. Not significant by Wilcoxon rank sum test. P = 0.46 (early), p = 0.201
(middle), p = 0.79 (late). N = 28 (hippocampus) and 16 (ACC) sessions in early sessions; 27
(hippocampus) and 9 (ACC) sessions in the middle sessions, 21 (hippocampus) and 6 (ACC) sessions
in the late sessions.

──

Stability of hippocampus and ACC place cells between sessions

We then compared the activity of each cell between two sessions in identical or distinct
contexts to see whether they switched their encoding mode between sessions (Fig. 2a). When the S1
and S1' sessions interleaved by a session O were compared, 66.1 ± 5.5% of ACC place cells in context
S1 still behaved as place cells in the context S1' (Fig. 2b). In contrast, only 7.0 ± 1.4% of ACC place
cell in context S1 behaved as a place cell in context O. The rest of the S1 place cells were either
unclassified (84.7 ± 2.1%) or not active (8.3 ± 2.4%). This proportion of encoding mode was
comparable to the proportion of all cell types in context O (place cells, 6.1 ± 1.1 %; unclassified, 84.8
± 2.2 %; not active 9.0 ± 2.4%. p = 0.65, 0.96, 0.85, N = 4 mice; one-way ANOVA), indicating
random reassignment of the place cells when mice moved to the other context. Likewise, place cells
in context O converted their encoding in context S1', in a proportion not different from their
proportion among all cells (Supplementary Fig. 3. Place cells, 30.6 ± 7.0%; unclassified cells 25.5 ±
5.4%, not active 44.0 ± 8.1% versus all cells in context S1': place cells, 33.2 ± 4.6%, unclassified
cells, 29.0 ± 3.6%; not active, 37.8 ± 3.0%; p = 0.92, 0.29, 0.50; N = 4 mice; one-way ANOVA). The
same tendency was observed in hippocampal place cells. Therefore, this analysis indicates that the
place cells in one context are randomly assigned either as place or non-place cells in different contexts
and suggests that there is no particular subclass of neurons which preferentially become place cells
in different contexts in both brain regions.

Finally, in order to quantitatively assess the change in the activity level of each cell, we
defined the selectivity index by calculating the absolute of the difference between firing rate in two
contexts normalized by the sum of firing rate in both contexts (Fig. 2c). This analysis revealed that
both hippocampal and ACC place cells showed a significant increase in the selectivity over days,

indicating that there was a significant differentiation in neuronal activity to one of the contexts during
repeated exposure to two different contexts (Fig. 2c).
203

Bota et al. Fig. 2

Figure 2. Context specificity of ACC and hippocampal place cells.

(a) Representative activity of an ACC place cell during one session (context S1→context
O→context S1'). Top, heat map for event rate of the cell. Bottom, mouse's trajectory (blue) and
calcium events (red). Firing intensity is normalized by the maximum activity of the cell throughout
the session.

(b) Comparison of place cell activity between two identical or different contexts. Encoding mode of
ACC place cells in context S1 was examined in context S1' or context O. ACC place cells in
context S1 were classified in 3 groups "place cells (PC)", "other" and "not active" according to its
activity pattern in context S1' or context O. Figure shows the average fraction of cells in each
group. Dotted lines show proportion of each class of cells in S1 or O, that serve as chance level.
Compared with the chance level, hippocampus S1 to S1': place cells to place cells $p = 1.8 \times 10^{-8}$,
place cells to other cell type $p = 1.2 \times 10^{-9}$, place cells to not active cells $p = 0.031$. Hippocampus S1
vs O: in the same order, $p = 0.42, 0.58, 0.018$. ACC S1 to S1': $p = 0.00027, 1.3 \times 10^{-7}, 0.52$. ACC
S1 to O: $p = 0.65, 0.96, 0.85$. One-way ANOVA. N = 4 mice for ACC, n = 5 mice for hippocampus.

(c) Selectivity index between context S1 versus context O. The index was pooled from 2045

hippocampal place cells (from 5 mice) and 931 ACC place cells (from 4 mice) in early phase (day 1-

3), 1655 hippocampal place cells (from 5 mice) and 1121 ACC place cells (from 4 mice) in late phase

(day 7-9). Whiskers show maximum and minimum value in each data set.

Long-term stability of place cells

We next examined whether these cells stably retain the same positional information on the

track across sessions on the same day or different days. We first compared the place field in context

S1 and S1' interleaved by a session in context O on the same day by calculating the population vector

correlation. In hippocampus the correlation increased during the 9-day sessions (days 1-3 vs 7-9, $p =$

2.5×10^{-3}, one-way ANOVA). This is consistent with the previous results which showed hippocampal

place cells become stable after repeating the same task [8, 23]. Also in ACC, we saw an increase of

correlation between S1-S1' sessions over days (days 1-3 vs 7-9, $p = 1.6 \times 10^{-7}$, one-way ANOVA).

We obtained essentially consistent result by calculating the ratio of cells with stable place fields (less

than 12.5 cm shift between S1-S1' sessions) among place cells in both hippocampus ($44.05 \pm 6.1\%$

of cells exhibited stable place field in days 1-3 and $64.7 \pm 5.5\%$ in days 7-9, $p = 0.0015$, one-way

ANOVA. Supplementary Fig.4) and ACC ($28.7 \pm 3.8\%$ of cells exhibited stable place field in days

1-3 and $58.1 \pm 3.6\%$ in days 7-9, $p = 9.6 \times 10^{-6}$, one-way ANOVA).

We then attempted to obtain a more holistic view of stability over days by comparing S1

sessions across days (Fig. 3c-e). We also found that both the hippocampal and ACC place cells

showed a gradual stabilization during the sessions repeated over 9 days. These results showed that

the ACC place cells have similar properties to the hippocampal place cells with respect to the stability

(Hippocampus early vs late $p = 0.022$; ACC early vs late $p = 0.014$; ACC early vs hippocampus early

$p = 0.93$; ACC late vs hippocampus late $p = 0.69$, one-way ANOVA).

Bota et al. Fig. 3

Figure 3. Stability of ACC and hippocampal place cells across sessions.

(a) A comparison of place cell map between context S1 and S1' within the same session on day 1 and day 9. Place-field maps was ordered according to the place field centroid position in context S1. The maps were normalized by the maximum activity of each cell.

(b) Population vector correlation of place cell representation in early, middle, and late sessions. Hippocampus early vs late $p = 2.5 \times 10^{-3}$; ACC early vs late $p = 1.6 \times 10^{-7}$; ACC early vs hippocampus early $p = 0.21$; ACC late vs hippocampus late $p = 0.24$, one-way ANOVA. N = 18 session pairs in early, 20 session pairs in middle, 20 session pairs in late for hippocampus. N = 18 session pairs in early, 14 session pairs in middle, 14 session pairs in late for ACC.

(c) A comparison of place cell map between context S1 and S1' in adjacent sessions.

(d) Heat map for population vector correlation of place cells in hippocampus and ACC between all pairs of sessions.

(e) Population vector correlation across early (day 1 vs day2 to day 3 vs day4), and middle (day 4 vs day5 and day 5 vs day 6) and late (day 6 vs day 7 to day 8 vs day 9) sessions for both hippocampus and ACC. Hippocampus early vs late $p = 0.022$; ACC early vs late $p = 0.014$; ACC early vs hippocampus early $p = 0.93$; ACC late vs hippocampus late $p = 0.69$, one-way ANOVA. N = 11 session pairs in early, 7 session pairs in middle, 8 session pairs in late for hippocampus. N = 10 session pairs in early, 7 session pairs in middle, 5 session pairs in late for ACC.

ACC place cell activity requires hippocampus for formation but becomes independent after training

  Finally, we examined whether the formation of ACC place cells requires hippocampal activity. To this end, we inhibited the hippocampal neuronal activity by administering a Designer Receptor Exclusively Activated by Designer Drugs (DREADD) agonist 21 (D21) to a mouse expressing its cognate inhibitory receptor, hM4Di-mCherry, in dorsal hippocampus (Fig. 4a) [24]. The mice received D21 20 min before each behavior experiments for 9 days. Mice in saline hM4D(Gi) group received saline for 9 days and then received D21 on day 10 (Fig. 4b).

  In the naïve, D21 naive or saline hM4D(Gi) groups, there was an increase in fraction of ACC place cells during 9-day period (Fig. 4c). However, in the D21 hM4D(Gi) group, the fraction remained low compared with control animals (on day 9. Naive vs. D21 naive, $p = 0.83$; Naive vs. D21 hM4D(Gi), $p = 1.9 \times 10^{-4}$; Naïve vs saline hM4D(Gi), $p = 0.38$. One way ANOVA. Fig. 4c). The stability of the map as assessed by the population vector correlation analysis between days 8 and 9 was also low stability in D21 hM4D(Gi) group compared saline hM4D(Gi) group (Fig. 4d. D21 hM4D(Gi) day 8 and 9 vs Saline hM4D(Gi) day 8 and 9, $p = 0.023$; D21 hM4D(Gi) day 8 and 9 vs Saline hM4D(Gi) day 9 and 10, $p = 4.7 \times 10^{-6}$. One-way ANOVA). This indicates that the hippocampal activity is required for establishing and maintaining ACC place cell maps.

  In contrast, when D21 was administered on day 10 in the saline hM4D(Gi) group after the place cell map was already established, it did not have effect on the proportion of ACC place cells

284 (Fig. 4c. p = 0.96, one-way ANOVA). Also, the population vector correlation between days 9 and 10

285 in the presence of D21 was not significantly different with days 8 and 9 when the same animals

286 received saline (Fig. 4d. p = 0.53, one-way ANOVA). This indicates that ACC place cell map, once

287 established, no longer requires hippocampal neuronal activity for maintenance.

Bota et al. Fig.4

Figure 4. The effect of chemogenetic inhibition of hippocampal excitatory neurons on ACC place cell.

(a) Representative mCherry image in TRE-GaMP7 x CaMKIIα-tTA double transgenic mice infected with AAV8-CaMKIIα-hM4D(Gi)-mCherry virus in hippocampi bilaterally. Scale bar = 100 μm.

(b) Experimental schedule.

(c) Percentage of ACC place cell in each context S1 and context S1'. In the bar graphs, circles represent individual trials from S1 or S1'. (Naive n =16 data (8 sessions x 2 direction) from 4 mice, D21 naive n =12 data (6 sessions x 2 direction) from 3 mice, D21 hM4D (Gi) n = 20 data (10 sessions x 2 direction) from 5 mice, Saline hM4D(Gi) n =12 data (6 sessions x 2 direction) from 3 mice). Day 1; naïve vs D21 naïve p = 0.23, naïve vs D21 hM4D (Gi) p = 0.033, naïve vs saline hM4D Gi) p = 0.054. Day 9; naïve vs D21 naïve p = 0.83, naïve vs D21 hM4D (Gi) p = 1.9 x 10^{-4}, naïve vs saline hM4D (Gi) p = 0.38, saline hM4D (Gi) day 9 vs day 10 p = 0.96, one-way ANOVA.

(d) Stability of place cell representation calculated by population vector correlation between day 8 and day 9, day 9 and day 10 in D21 hM4D (Gi) or saline hM4D (Gi) animals. P = 0.023 D21 hM4D (Gi) day 8 and 9 vs Saline hM4D (Gi) day 8 and 9, p = 4.7 x10^{-6} D21 hM4D(Gi) day 8 and 9 vs Saline hM4D(Gi) day 9 and 10, p = 0.53 Saline hM4D(Gi) day 8 and 9 vs Saline hM4D(Gi) day 9 and 10, one-way ANOVA. N = 10 session pairs for D21 hM4D (Gi) day 8 and day9, 6 session pairs for saline hM4D (Gi) day 8 and day 9, 6 session pairs for saline hM4D (Gi) day 9 and day 10.

Discussion

mPFC, in particular ACC, has been implicated in the process of memory consolidation [9, 10]. During this process, coherent theta oscillations coupling between the hippocampus and mPFC facilitates the transfer of memory [25]. However, the exact content of memory carried by the neurons in ACC and whether it changes their properties during consolidation, have not been fully elucidated. In this study, we examined how representation of positional information is processed in the downstream of hippocampus. To this end, we recorded calcium activity over days from the ACC of freely moving mice. We found a population of neurons in ACC which shows location specific activity. It shares basic properties with hippocampal place cells such as location specific firing and directional selectivity [1-4]. However, unlike hippocampus place cells, the ACC place cells often have extra firing field. This may represent association of information from more than one location on the track. Indeed, mPFC including ACC is involved in the processing contextual information and such association of multiple locations is likely to play a role.

In addition, we found that ACC place cells gradually increased over days in a manner requiring hippocampal activity for the formation. However, once the ACC place map is formed after 9 days, ACC no longer required hippocampal activity to fire and to exhibit a place cell map. This property is reminiscent of systems consolidation process of episodic memory, where hippocampus is required for initial formation of memory but not for recall of remote memory [11, 12, 26]. The memory consolidation theory predicts hippocampal activity after the memory events induces cortical plasticity and consolidates memory in cortical circuit[11]. After memory consolidation, hippocampus is no longer required for recall of remote memory[11]. The same analogy can be applied to the place cells in both hippocampus and ACC. At this point, it is not clear from which brain regions ~~the~~ spatial information is generated under hippocampal inactivation. ACC is bidirectionally connected to other cortical regions such as retrosplenial and entorhinal cortexes, both of which are implicated in processing spatial information [14, 27, 28]. It is therefore possible that spatial information arriving from these cortexes as well other regions can bypass the requirement of hippocampus.

In conclusion, we found that ACC place cells have both shared and unique properties with hippocampal counterparts. Further study is required to elucidate how their formation depends on the hippocampus, but they are maintained without hippocampal activity once they are formed. It will be intriguing to test if these cells indeed overlap with engram cells as defined by *c-fos* expression [6].

ACKNOWLEDGEMENTS

We thank Dr. Adam Z. Weitemier for their comments on the manuscript. This work was supported

by RIKEN, Grant-in-Aid for Scientific Research A JP19H01010, Grant-in-Aid for Scientific

Research on Innovative Area "Foundation of Synapse and Neurocircuit Pathology" JP22110006,

"Principles of memory dynamism elucidated from a diversity of learning systems" JP90466037, and

"Constructive understanding of multi-scale dynamism of neuropsychiatric disorders" JP18H05434

from MEXT, and Human Frontier Science Program RGP0022/2013 to Y.H. and JP20K22686 from

MEXT and the Sasakawa Scientific Research Grant from The Japan Science Society to A.B.

Reference

1. Eichenbaum, H., et al., *The hippocampus, memory, and place cells: is it spatial memory or a memory space?* Neuron, 1999. **23**(2): p. 209-26.
2. O'Keefe, J.N., Lynn, *The Hippocampus as a Cognitive Map.* Oxford: Clarendon Press, 1978.
3. Moser, M.B., D.C. Rowland, and E.I. Moser, *Place cells, grid cells, and memory.* Cold Spring Harb Perspect Biol, 2015. **7**(2): p. a021808.
4. Cobar, L.F., L. Yuan, and A. Tashiro, *Place cells and long-term potentiation in the hippocampus.* Neurobiol Learn Mem, 2017. **138**: p. 206-214.
5. Ziv, Y., et al., *Long-term dynamics of CA1 hippocampal place codes.* Nat Neurosci, 2013. **16**(3): p. 264-6.
6. Ghandour, K., et al., *Orchestrated ensemble activities constitute a hippocampal memory engram.* Nat Commun, 2019. **10**(1): p. 2637.
7. Gauthier, J.L. and D.W. Tank, *A Dedicated Population for Reward Coding in the Hippocampus.* Neuron, 2018. **99**(1): p. 179-193 e7.
8. Sato, M., et al., *Distinct Mechanisms of Over-Representation of Landmarks and Rewards in the Hippocampus.* Cell Rep, 2020. **32**(1): p. 107864.
9. Bontempi, B., et al., *Time-dependent reorganization of brain circuitry underlying long-term memory storage.* Nature, 1999. **400**(6745): p. 671-5.
10. Teixeira, C.M., et al., *Involvement of the anterior cingulate cortex in the expression of remote spatial memory.* J Neurosci, 2006. **26**(29): p. 7555-64.
11. Frankland, P.W., et al., *The involvement of the anterior cingulate cortex in remote contextual fear memory.* Science, 2004. **304**(5672): p. 881-3.
12. Kitamura, T., et al., *Engrams and circuits crucial for systems consolidation of a memory.* Science, 2017. **356**(6333): p. 73-78.
13. Vogt, B.A. and G. Paxinos, *Cytoarchitecture of mouse and rat cingulate cortex with human homologies.* Brain Struct Funct, 2014. **219**(1): p. 185-92.
14. Hoover, W.B. and R.P. Vertes, *Anatomical analysis of afferent projections to the medial prefrontal cortex in the rat.* Brain Struct Funct, 2007. **212**(2): p. 149-79.
15. Laubach, M., et al., *What, If Anything, Is Rodent Prefrontal Cortex?* eNeuro, 2018. **5**(5).
16. Fujisawa, S., et al., *Behavior-dependent short-term assembly dynamics in the medial prefrontal cortex.* Nat Neurosci, 2008. **11**(7): p. 823-33.
17. Hyman, J.M., et al., *Contextual encoding by ensembles of medial prefrontal cortex neurons.* Proc Natl Acad Sci U S A, 2012. **109**(13): p. 5086-91.
18. Ma, L., et al., *A Quantitative Analysis of Context-Dependent Remapping of Medial Frontal Cortex Neurons and Ensembles.* J Neurosci, 2016. **36**(31): p. 8258-72.
19. Mashhoori, A., et al., *Rat anterior cingulate cortex recalls features of remote reward locations after disfavoured reinforcements.* Elife, 2018. **7**.
20. Burton, B.G., et al., *Lesion of the ventral and intermediate hippocampus abolishes anticipatory activity in the medial prefrontal cortex of the rat.* Behav Brain Res, 2009. **199**(2): p. 222-34.
21. Low, R.J., Y. Gu, and D.W. Tank, *Cellular resolution optical access to brain regions in fissures: imaging medial prefrontal cortex and grid cells in entorhinal cortex.* Proc Natl Acad Sci U S A, 2014. **111**(52): p. 18739-44.
22. Hafting, T., et al., *Microstructure of a spatial map in the entorhinal cortex.* Nature, 2005. **436**(7052): p. 801-6.
23. Rubin, A., et al., *Hippocampal ensemble dynamics timestamp events in long-term memory.* Elife, 2015. **4**.
24. Thompson, K.J., et al., *DREADD Agonist 21 Is an Effective Agonist for Muscarinic-Based DREADDs in Vitro and in Vivo.* ACS Pharmacol Transl Sci, 2018. **1**(1): p. 61-72.

25.   Xing, B., M.D. Morrissey, and K. Takehara-Nishiuchi, *Distributed representations of temporal stimulus*
associations across regular-firing and fast-spiking neurons in rat medial prefrontal cortex. J
Neurophysiol, 2020. **123**(1): p. 439-450.
26.   Restivo, L., et al., *The formation of recent and remote memory is associated with time-dependent*
formation of dendritic spines in the hippocampus and anterior cingulate cortex. J Neurosci, 2009.
29(25): p. 8206-14.
27.   Takehara-Nishiuchi, K., *Entorhinal cortex and consolidated memory.* Neurosci Res, 2014. **84**: p. 27-33.
28.   de Landeta, A.B., et al., *Anterior retrosplenial cortex is required for long-term object recognition*
memory. Sci Rep, 2020. **10**(1): p. 4002.
29.   Kinsky, N.R., et al., *Hippocampal Place Fields Maintain a Coherent and Flexible Map across Long*
Timescales. Curr Biol, 2018. **28**(22): p. 3578-3588 e6.
30.   Cai, D.J., et al., *A shared neural ensemble links distinct contextual memories encoded close in time.*
Nature, 2016. **534**(7605): p. 115-8.
31.   Markus, E.J., et al., *Spatial Information-Content and Reliability of Hippocampal Ca1 Neurons - Effects*
of Visual Input. Hippocampus, 1994. **4**(4): p. 410-421.
32.   Kazumasa Z. Tanaka, H.H., Anupratap Tomar, Kazue Niisato, Arthur J. Y. Huang, Thomas J. McHugh,
The hippocampal engram maps experience but not place. Science, 2018.
33.   Kinsky, N.R., et al., *Trajectory-modulated hippocampal neurons persist throughout memory-guided*
navigation. Nat Commun, 2020. **11**(1): p. 2443.
34.   Leutgeb, J.K., et al., *Progressive transformation of hippocampal neuronal representations in*
"morphed" environments. Neuron, 2005. **48**(2): p. 345-58.
35.   Zhang, K.C., et al., *Interpreting neuronal population activity by reconstruction: Unified framework*
with application to hippocampal place cells. Journal of Neurophysiology, 1998. **79**(2): p. 1017-1044.

Materials and methods

Subjects

All experiments were conducted in accordance with institutional guidelines and protocols approved by the RIKEN and Kyoto University Animal Experiments Committees. Experiments were conducted on 10–24 weeks old male TRE-G-CaMP7 x CaMKIIα-tTA double transgenic mice [8]. Mice were housed singly in a cage with 12 h-12h light-dark cycle (with lights on at 6 pm and off at 6 am on the next day). All experiments were performed between 6 am and 6 pm.

To remove crosstalk from neighboring ROIs, clustering score of each ROI is calculated as follows.

After ROIs (red) were identified using a custom MATLAB routine (see method), 5-time enlarged ROIs (by area, yellow) were made. For a given frame of Ca2+ images, the location of pixels with top 20% brightness were detected. The proportion of the pixels within the original ROI is defined as the clustering score for each frame. Low clustering score indicates high likeliness of crosstalk of neighboring ROI.

Histology

Mice were perfused transcardially with phosphate-buffered saline (PBS) followed by 4% paraformaldehyde   in PBS. Brains were extracted and post-fixed in formalin for an additional day, and were transferred to a PBS for an additional day. Brains were then sliced in 50 µm sections

using a microslicer (Dosaka). For immunohistochemistory, sliced tissue sections were incubated in
0.1 M Tris-HCl, 0.15 M NaCl, 0.5 % Triton-X and 5 % blocking reagent (Roche) and rabbit anti-
GFP antibody (A11122, Thermo Fisher Scientific, 1:500) and incubated overnight at 4 °C. After
rinsing with PBS 3 times for 15 min each, sliced tissue sections were subsequently incubated with
AlexaFluor 488 (A11008, Thermo Fisher Scientific, 1:500) conjugated secondary antibodies. Sliced
tissue sections were then washed in PBS 3 times for 15 min, mounted, and coverslipped with
mounting medium with Hoechst 33258 (#382061, Calbiochem). Fluorescence images were taken
by confocal microscopy (Olympus FLUOVIEW FV1500) [29].

Surgery for calcium imaging from Hippocampal CA1

Mice were anesthetized with isoflurane in ambient air (3% induction, 1.5% maintenance) and
placed in a stereotaxic frame. A stainless-steel head plate (25 mm length, 4 mm width, 1 mm
thickness) with a circular opening (7 mm inner diameter and 10 mm outer diameter, the center is 2.5
mm off relative to the middle of the long side of the plate) was affixed to the skull using dental
cement. The center of the opening was targeted at 2 mm posterior to the bregma and 2 mm lateral to
the left from the midline. The cement was placed onto the skull such that it covered the entire skull,
including the three anchor screws, except for the area of skull inside the opening of the head plate.

Several days later, we used a trephine drill to remove a circular part of the left skull centered 2.0
mm posterior, 2.0 mm lateral, and 2.0 mm deep from bregma. We removed the dura and cortex
above the CA1 by suction with a 25 or 27 gauge blunt needle while constantly washing the exposed
tissue with sterile cortex buffer (123 mM NaCl, 5 mM KCl, 10 mM glucose, 2 mM $CaCl_2$, 2 mM
$MgCl_2$, 10 mM HEPES, pH 7.4). We then implanted a metal guide tube with its window just dorsal
to CA1 region and sealed the space between the skull and guide tube using dental acrylic [23].

Surgery for calcium imaging from ACC

A stainless-steel head plate was affixed to the skull as described above at 1 mm anterior to the
bregma centered at midline. The dura was removed to provide access to the longitudinal fissure.
Several days later, a microprism comprised of a right angle microprism (1 mm x 1 mm x 1 mm)
bonded to GRIN lens (0.85-mm diameter, 3.3-mm length, Inscopix) was implanted. The face of the
prism was aligned parallel to the fissure and positioned over the midline at the target 1.0-mm
anterior to bregma. After gently nudging the sinus laterally (~150 μm), the tip of the prism was
positioned at the entrance to the fissure. Then the micromanipulator was used to vertically lower the
prism into the subdural space in the longitudinal fissure so that its front faces against the medial
surface of the left ACC. We then sealed the space between the skull and lens using dental cement
and adhesive bond. The exposed lens was sealed with Silicone adhesive (Kwik-Sil, World Precision
Instruments) and dental cement [21].

Calcium imaging

For hippocampal imaging, four weeks after the surgery described above, a gradient refractive index
lens (GRINtech GmbH, 0.44 pitch length, 0.47 NA) was fixed in the guide tube using ultraviolet-
curing adhesive (Norland, NOA 81). For ACC imaging, the silicone adhesive was removed to
expose prism lens. We lowered the integrated microscope (nVistaHD, Inscopix) toward the GRIN
lens until we saw G-CaMP7 fluorescence using the microscope's LED light source (0.12–0.24
mW). After finding a suitable imaging site, we attached the microscope's base plate to the skull
using dental acrylic darkened with black acrylic paint. This plate stayed with the mouse even after
the microscope was detached to ensure reproducibility of imaging site. Before each behavior
experiment, the microscope was attached to the base plate under light isoflurane anesthesia. The
imaging session was started at least after 15 min recovery in the home cage.

In each session of the behavioral experiment, the animals were placed first in a square track
(context S1), then an open arena (context O) with different wall patterns, and then again in the
square track (context S1') for 10 min each with 5 min interval. Context S1 consists of a square track
of four divisions (each 50 cm long x 5 cm wide) and side walls (8 cm tall). A wall was placed at one
of the corners to separate the ends. The animals were trained to run back-and-forth between the two
ends for a sucrose tablet (#1811251 Sucrose Rewards Tablet, Test Diet), given when the animal
reached one end. In the context O (50 x 50 cm with wall of 25 cm tall), the rewards were randomly
thrown into the open arena. Immediately before the beginning of each session, the equipment was
wiped with differently scented paper towels (80% ethanol for context S and 0.5% acetic acid for
context O). Before beginning the imaging, mice were food restricted and they were exposed to
context O and context S as pre-training for 3 days. After pre-training, one S1-O-S1' session per day
was performed during calcium imaging and repeated for 9 days. On day 1, mice were let freely
move for 10 min in each context without any food reward. After day 1 pre-training, foods in their
home cage are removed. In day 2 and 3, food reward was given as described above. After each day
2 day3 of pre-training and each daily imaging, mice got 1-3 g food to keep their weight. We
recorded a total 5 mice from hippocampus and a total 4 mice from ACC [5, 23].

DREADD

For DREADD experiments, DREADD agonist 21 (D21; Cayman Chemical Company, #18907), an
alternative to CNO, was made in a stock solution of 5 mg/ml in DMSO and then diluted in saline to
desired concentration (0.05 mg or 0.1 mg/ml). D 21 was injected intraperitoneally at 1 mg/kg 20
min before the behavioral experiment.[30]

Processing of Calcium imaging data

We processed imaging data using Mosaic (Inscopix). To increase computation speed, we spatially
down-sampled the data by a factor of four in each dimension. The down-sampled imaging data was
motion corrected and transformed to relative changes in fluorescence, $\Delta F(t)/F0 = (F(t)-F0)/F0$,
where F0 is the value for each pixel averaged over the entire time span of the calcium imaging
movie. Finally, the movie was smoothed by applying disk average filter (disk radius: 3x3 pixels).

We separately used un-filtered movie for calculation of each neuron's fluorescence intensity, and
filtered movie for neuron identification and calculating clustering score, respectively. (see 'Neuron
Identification' and 'Calcium Event Identification').

(Users Manual of Mosaic, Inscopix, 2014)

Neuron Identification

Regions-of-interest (ROIs) was identified using a custom MATLAB routine. First, intensity of each
pixel is normalized by the mean value of all pixel in that frame of the filtered calcium imaging
movie. Then, it adaptively thresholds each imaging frame to identify separable pockets of calcium
activity, designated as blobs, on each frame. Blobs of activity were accepted at this stage of
processing only if they approximate the size and shape of a neuron, as measured by their size (min
= 250 μm^2 (30 pixels), max = 1500 μm^2 (180 pixels)), and the ratio of long to short axes (max = 2),
a metric used by the regionprops function of MATLAB we employ to reject jagged/strange shaped
blobs. Signal traces of each listed blob are calculated over all frames of the movie. Finally, overlap
of region among all blobs and correlation of activity (mean gray value of pixels in the blob) among
all blobs were calculated respectively. Blob pairs with high spatial correlation (r > 0.4) and high
signal correlation (r > 0.9) are considered as the same neuron. Smaller one of the two was used as
ROI for further analysis [29].

Calcium Event Identification

We extracted $\Delta F/F$ traces of each ROIs from un-filtered calcium imaging movies. To remove
crosstalk from neighboring ROIs, clustering score of each ROI is calculated as follows. After ROIs
(red) were identified using a custom MATLAB routine, 5-time enlarged ROIs (by area, yellow)
were made (Fig. S5). For a given frame of Ca^{2+} images, the location of pixels with top 20%
brightness were detected. The proportion of the pixels within the original ROI is defined as the
clustering score for each frame. Low clustering score indicates high likeliness of crosstalk of
neighboring ROI. By excluding value with low clustering score from each ROI, crosstalk was
removed. Second and third steps are performed in each frame of movie. Time points with $\Delta F/F$
signal >1.5 and clustering score >0.4 are detected as calcium event for each ROI.

Neuron Registration

548 Neuron registration occurred in two steps: session registration and neuron registration.

Session registration

Prior to mapping neurons between sessions, we determined how much the imaging window shifted
between sessions. In order to isolate consistent features of the imaging plane for each mouse (such
as vasculature or coagulated blood), we created a median projection of all of the frames of the
imaging movie for each recording session. One session (''registered session'') was then registered
to a base session (first session) using the 'motion correction' function from the Mosaic and

calculated transformation object was saved. This process of transformation was applied to each
calcium imaging movie of all sessions, and saved for further analysis. [29]

Neuron Registration

Next, each ROI in the registered session was transformed to its corresponding location in the base
session. Each neuron in the base session was then mapped to the neuron with the closest center-of-
mass in the registered session, unless the closest neuron exceeded our maximum distance threshold
of 4 pixels (10 μm). The spatial correlation between each pair of ROIsd are also calculated. We
designated the base session ROI with the high spatial correlation ($r > 0.6$) as mapping to the
registered session ROI. [29]

Place cells and place field

We used the following criteria to identify place cells and place fields in the open field and the
square track, respectively.

Place cells in open field

To identify place cells in the open field, firing rate maps were obtained as follows. First, calcium
events from each ROI were assigned to 16 cm x 16 cm spatial bins. Second, the number of calcium
events in each bin was divided by the occupancy time of that bin.

Then we computed the spatial information using the firing rate maps of each cell, as previously
described [31]

$$Spatial\ information = \sum_i P_i(r_i/\bar{r})log_2(r_i/\bar{r})$$

Where r_i is the calcium event rate of the neuron in the i^{th} bin; P_i is the probability of the mouse
being in the i^{th} bin (time spent in i^{th} bin/total session time); $\bar{r}$ is the overall mean calcium event
rate; and i running over all the bins. We then performed 1000 permutation shuffles, and calculated
the spatial information for each shuffle. This yielded the p value of the measured information
relative to the shuffles. Cells with $p < 0.05$ and mean firing rates higher than 0.1 Hz were classified
as place cell.

To detect place fields in the open field, we make firing rate maps with 2.5 cm x 2.5 cm spatial
bins, normalized firing rate of each bin by maximum value in the map, and smoothed the
normalized map with 1 SD Gaussian kernel. Next, we made binary map based on the firing rate of
each bin (spatial bin with firing rate>0.2 is 1, other spatial bins are 0). Connected bins in this binary
map are detected by using MATLAB bwlabel function (Pixel connectivity = 8-connected). Each
connected bin is considered as individual place field. Place field which contains bin with maximum
firing rate is designated as main place field, and other place fields as extra place field. [23] [32]

Place cell in Square track

To identify place cells in the square track, we employed reliability, which measured the consistency of a cell to fire on its preferred location along the track (Supplementary Fig. 2) [33]. We calculated reliability in the following manner. First, we divided track into 72 spatial bins (2.8cm each), and exclude the first and the last 5 bins where food rewards were given. Spatial activity in each lap was transformed into binarized vectors, in which 1 and 0 represent the presence and absence of a calcium events, respectively. Third, we calculated the correlation value between each lap and the mean of all correlation values. Finally, the locations of calcium events in the binarized vector are shuffled randomly 1000 times, then reliability of each shuffled data was calculated. Cells with significant reliability ($p < 0.05$) are considered as place cell in the square track (Supplementary Fig. 2). Number of running lap with calcium event of all cells and population vector correlations between a session and all other sessions are calculated to exclude session without spatial activity. The sessions with mean number of running with calcium event lap < 1.5 or mean population vector correlation < 0.005 are excluded from analysis. We separately considered place cells for forward and backward running directions.

To detect place fields in the square track, we normalized the firing rate map by maximum value, smoothed the normalized map with 1 SD Gaussian kernel, and converted the smoothed map to binary map (spatial bin with firing rate > 0.2 is 1, other spatial bins are 0). Connected bins in this binary map are detected, and each connected bin is considered as individual place field. Place field which contains bin with maximum firing rate is designated as main place field, and other place fields as extra place field[23].

Selectivity index

We used the following calculation to determine the selectivity index of each cell: |(activity in context S1 – activity in context O) / (activity in context S1 + activity in context O)|.

Population vector correlation

To determine the level of similarity between representations of the different sessions, we calculated the mean population vector correlation between them [34]. For each spatial bin (excluding the last 5 bins at both ends of the tracks) we defined the population vector as the mean event rate for each neuron given that bin's firing rate. We computed the correlation between the population vector in one session with that of the matching location in the other session, and averaged the scores over all positions. [23]

Statistics

Data are expressed as means ± SEM unless stated otherwise. All statistical tests were performed using MATLAB. One-way ANOVA was used for group/pair-wise comparisons. Where appropriate, Student's paired t-tests or Wilcoxon rank sum tests was conducted.　The null hypothesis was rejected at the $P < 0.05$ level.

Bayesian decoder

629   We used a Naive Bayes Classifier [5, 23, 35] to estimate the mouse location based on neuron
630   calcium activity. The computation of the conditional probability for the subject to be at location x is
631   based on Bayes formula:

632
$$P(x|\vec{n}) = P(\vec{n}|x)\,P(x)\,/\,P(\vec{n}).$$

n is the population activity vector of length N containing 1 at index i, when neuron i is
considered to be active and 0 when it is not. $P(x)$, the probability for the subject to be at position x
was obtained from the dwell time distribution at each spatial bin (bin size = 2.8 cm). $P(\vec{n}/x)$, the
conditional probability to observe $\vec{n}$ given the subject is at position x was computed from the
spatial map $P(n_i|x)$ of individual neurons i assuming statistical independence between their activity
[5].

639
$$P(\vec{n}|x) = \prod_{i=1}^{N} p(n_i|x)$$

640   The overall probability to observe n is obtained from $P(\vec{n}) = \sum_{j \in X} P(\vec{n}/x_j)\,P(\vec{n}/x_j)\,P(x_j)$, with
641    $P(x_j)$ normalized along the spatial dimension X. A reconstructed position of the subject is obtained
642   from the peak position of the evaluated $P(x/\vec{n})$, Eq.1 with $\vec{n} = \vec{n}$ (t). This estimate is further refined
643   taking into account the data from $dT = 8$ frames before the current frame, assuming a negligible
644   change in position of the animal within dT:

645
$$x_{rec} = \mathrm{argmax}_j \prod_{\delta t=0}^{dT} P\;(\vec{n}(t-\delta t)|x_j)\,P(x_j)\,/\,P(\vec{n}(t-\delta t))$$

646   The estimation error was calculated as the absolute difference between the real and the
647   reconstructed positions. We trained the decoder with the subjects observed positions and activities
648   of all place cells (place cells with top 6-15% higher firing rate were used) during the first half time
649   of the running period and estimated the trajectory for the following half time. The running time of
650   the same sessions using the corresponding place cell activities.

SUPPLEMENTARY FIGURES

Bota et al. supple. 1

Figure S1 Histological confirmation of imaging location.

(a) Left, a coronal section stained with Hoechst 33258 (blue) and anti-GFP antibody (green) of
TRE-G-CaMP7 x CaMKIIα-tTA mouse after imaging from ACC. Right, A zoomed image from
rectangle area. Hoechst image was overlaid with G-CaMP7 immunofluorescence.

(b) A coronal section stained with Hoechst 33258 and anti-GFP antibody of TRE-G-CaMP7 x
CaMKIIα-tTA mouse after imaging from hippocampus.

Correlation between each lap ⇒ average (Reliability)

Bota et al. supple. 2

Figure S2. Calculation of reliability of place cell.

Examples of trajectory of mouse and calcium events of ACC place cells. Spatial activity in each lap
was transformed into binarized vectors, in which 1 and 0 represent the presence and absence of a
calcium events, respectively. We calculated the correlation value between each lap and the mean of
all correlation value. Examples of ACC place cells with high reliability and low reliability are
shown.

Bota et al. supple. 3

Figure S3. Conversion of encoding mode of hippocampal and ACC place cells in context O
and in context S1.

Encoding mode of hippocampal or ACC place cells in context O was examined in context S1'.
Compared with unbiased change, hippocampus: p = 0.082, 0.090, 0.78. ACC: p = 0.92, 0.29, 0.50.
One-way ANOVA. N = 5 mice for hippocampus, n = 4 mice for ACC.

____

Bota et al. supple. 4

Figure S4. The fraction of stable place cells on the same day

Place cell stability calculated as the fraction of stable place cells relative to the number of total
place cells identified in each session that were compared. Cells in early phase (day 1-3), middle
phase (day 4-6) and late phase (day 7-9) were pooled respectively. p = 9.6 x 10^{-6} (ACC early vs
ACC late), p = 0.0015 (hippocampus early vs hippocampus late), p = 0.017 (hippocampus early vs
ACC early), p = 0.203 one-way ANOVA. N = 18 session pairs in early, 20 session pairs in middle,
16 session pairs in late for hippocampus. N = 18 session pairs in early, 16 session pairs in middle,
14session pairs in late for ACC.

____

Bota et al. supple. 5

Figure S5. Remapping of place cells across days.

Place-field maps of hippocampal and ACC place cells ordered by their centroid positions on day 1 (top), day 5 (middle) or day 9 (bottom).

Original ROI

5-time enlarged ROI

Bota et al. supple. 6

Figure S6. Clustering score.

To remove crosstalk from neighboring ROIs, clustering score of each ROI is calculated as follows.
After ROIs (red) were identified using a custom MATLAB routine (see method), 5-time enlarged
ROIs (by area, yellow) were made. For a given frame of Ca^{2+} images, the location of pixels with top
20% brightness were detected. The proportion of the pixels within the original ROI is defined as the
clustering score for each frame. Low clustering score indicates high likeliness of crosstalk of
neighboring ROI.

702
703