Innate immune signaling in trophoblast and decidua organoids defines differential antiviral defenses at the maternal-fetal interface

Liheng Yang[1†], Christina Megli[2,3], Carolyn B. Coyne[1†*]

[1]Department of Pediatrics, University of Pittsburgh School of Medicine, [2]Division of Maternal-Fetal Medicine, Division of Reproductive Infectious Disease, Department of Obstetrics, Gynecology and Reproductive Sciences, University of Pittsburgh Medical Center (UPMC),[3]Magee Womens Research Institute, Pittsburgh, PA USA

[†]Current address: Department of Molecular Genetics and Microbiology, Duke University School of Medicine, Durham, NC USA

*Address correspondence
Carolyn Coyne, PhD
3130 Medical Sciences Research Building III
3 Genome Court
Durham, NC 27710
USA
carolyn.coyne@duke.edu

Abstract

Infections at the maternal-fetal interface can directly harm the developing fetus and compromise the health of the pregnant woman. Innate immune signaling by both fetal-derived placental trophoblasts and the maternal decidua must provide antimicrobial defenses at this critical interface without compromising its integrity. Here, we developed trophoblast and decidua organoids from matched human placentas to define their relative contributions to innate immune defenses at the maternal-fetal interface. We show that trophoblast and decidua organoids recapitulate some, but not all, of the basal cytokine release observed in matched tissue explants. We further show that trophoblast, but not decidua, organoids constitutively release antiviral type III interferons (IFNs) and that fetal and maternal organoids differentially respond to viral infections through the induction of organoid-specific cytokines. These findings define key differences in innate immune signaling generated by fetal-derived trophoblasts and the maternal decidua, which together must protect the fetus from viral infections.

Introduction

The maternal-fetal interface is comprised of both the maternal-derived decidua and the fetal-derived placenta, which together support fetal growth and provide immunologic defenses that protect the fetus from infections. The relative contributions of specific maternal- and fetal-derived cells to these critical functions remain largely undefined given the limitations of models that recapitulate the complex nature of this interface.

The placenta forms a key barrier to viral vertical transmission throughout pregnancy and is comprised in part by fetal-derived trophoblast cells. Trophoblast stem cells located in chorionic villi, the branch like projections that directly contact maternal blood, give rise to cytotrophoblasts (CTBs), the proliferative mononuclear cells of the placenta, extravillous trophoblasts (EVTs), and the syncytiotrophoblast (STB), a multinucleated contiguous cell layer that covers the villous surfaces. EVTs directly remodel the maternal microvasculature to form spiral arteries, which ultimately deliver maternal blood to the surfaces of chorionic villi. Decidualization of the endometrium is required for implantation and appropriate placentation. This process involves the differentiation of fibroblast-like mesenchymal cells into secretory decidua cells. Prior to the formation of chorionic villi and the development of a hemochorial placenta, these secretory cells provide necessary nutrients to the developing embryo (Burton et al., 2002). Together, the decidua and placenta thus provide the fetus with essential hormones and nutrients to support its growth.

While it is clear that the placenta and decidua play complementary roles in fetal development, their contributions to antimicrobial defenses are less well defined. Studies of tissue explants and histopathologic analysis of the placenta suggest that the decidua and chorionic villi have differing roles in the pathogenesis of relevant infections. The decidua may be a primary site of replication for several microorganisms associated with congenital disease, including cytomegalovirus (CMV) (McDonagh et al., 2006; Pereira et al., 2003; Weisblum et al., 2011), Zika virus (ZIKV) (Guzeloglu-Kayisli et al., 2020; Platt et al., 2018; Weisblum et al., 2017), and *Listeria monocytogenes* (Rizzuto et al., 2017), suggesting it may serve as a reservoir for these infections. By contrast, the placenta is

largely resistant to infections and possesses a number of intrinsic mechanisms of innate defense. In the case of viruses, human trophoblasts constitutively release antiviral interferons (IFNs) that restrict infection in both autocrine and paracrine manners (Bayer et al., 2016; Corry et al., 2017). These IFNs are of the type III class, which include IFNs-λ1-3 in humans (Lazear et al., 2019; Wells and Coyne, 2018). Constitutive release of IFNs is a unique feature of trophoblasts as IFNs are tightly regulated in most cell types and induced only after detection of the products of viral replication. In mice, type III IFNs also confer protection from Zika virus (ZIKV) vertical transmission and fetal disease, which occurs in a gestational-associated manner (Chen et al., 2017; Jagger et al., 2017), although it is not clear whether mice constitutively release these IFNs. Together, these findings suggest that maternal- and fetal-derived tissue differentially respond to and control viral infections. However, the lack of available models to fully interrogate the differences between maternal- and fetal-derived cells has limited a complete understanding of these processes, including how each tissue type responds to infection.

The recent development of trophoblast and decidua organoids from first trimester placental tissue has expanded the models available to study the maternal-fetal interface (Turco et al., 2017; Turco et al., 2018). Previous work has shown that isolation and propagation of trophoblast stem cells from first trimester placental chorionic villi leads to the development of three-dimensional organoids that differentiate to contain all lineages of trophoblast cells present in the human placenta and release well-known pregnancy hormones such as human chorionic gonadotropin (hCG) (Turco et al., 2018). Similarly, decidua organoids isolated from uterine glands recapitulate the transcriptional profile of their tissue of origin and respond to hormonal cues (Turco et al., 2017). These organoid models thus provide ideal systems to profile differences between maternal- and fetal-derived cells, however, these models have not been applied to pathogens infections.

To directly compare antiviral signaling pathways between the human placenta and decidua, we generated trophoblast and decidual organoids from matched human placentas and profiled the release of cytokines, chemokines, and other factors from these models under uninfected and virally

infected conditions. In parallel, we performed similar profiling in matched tissue explants from which these organoids were developed and found that organoids recapitulated some, but not all, of the cytokine release profiles of this tissue. To determine whether there are differences in the responses of maternal- and fetal-derived cells to viral infections, we also compared the immunological secretome from organoids treated with a ligand to stimulate toll-like receptor 3 (TLR3) signaling or after infection with ZIKV. These studies revealed that trophoblast organoids selectively secrete the type III IFN IFN-λ2, which does not occur in matched decidua organoids. Moreover, trophoblast and decidua organoids differentially respond to TLR3 activation and ZIKV infection through the secretion of organoid-specific cytokines. Our studies highlight the differential responses of key cells comprising the maternal-fetal interface and provide new models to define the impact of pathogen exposure at this key site.

Results

Establishment and characterization of human trophoblast and decidual organoids

To generate matched human trophoblast organoids (TOs) and decidual organoids (DOs), we used human placental tissue isolated from the second and third trimesters of human pregnancy and cultured stem cells isolated from chorionic villi or isolated decidua in growth factor-containing medium similar to that described previously (Haider et al., 2018; Okae et al., 2018; Sheridan et al., 2020; Turco et al., 2017; Turco et al., 2018) (**Figure 1A**). Isolated stem cells were embedded in Matrigel to allow for self-organization and propagated for ~3 weeks to generate organoid structures that were morphologically similar to those described previously that were isolated from the first trimester (Turco et al., 2017; Turco et al., 2018) (**Figure 1B**). Also similar those isolated from the first trimester (Turco et al., 2018), cultured TOs developed projections that resembled villous structures (**Figure 1B**, arrows at bottom). Confocal microscopy confirmed the three-dimensional architecture of established organoids and verified that TOs expressed markers such as cytokeratin-19 (**Figure 1C**) and DOs expressed markers including cytokeratin-7 and -19 and E-cadherin (**Figure 1D**), consistent with

previous reports (Turco et al., 2017; Turco et al., 2018). Also consistent with first trimester tissue-derived TOs (Turco et al., 2018), established mid-to-late gestation TOs differentiated to include various trophoblast lineages, including EVTs, which extended from the TOs to invade the surrounding Matrigel (**Supplemental Figure 1A, 1B**). We confirmed that TOs and DOs expressed markers consistent with their tissue of origin, including the trophoblast specific marker GATA3, which was enriched in TOs but not DOs, and MUC20, which was expressed in DOs but not TOs (**Figure 1E, 1F**). We also confirmed that TOs, but not DOs, secreted the trophoblast-specific hormone human chorionic gonadotropin (hCG) by both ELISA and rapid test strip assays (**Figure 1G**). Lastly, we also confirmed that TOs secreted pregnancy-specific beta-1 glycoprotein 1 (PSG1), a major product released from the STB, similar to chorionic villous explants (**Figure 1H**) . TOs and DOs could be propagated for at least three months and passaged approximately every 2-3 weeks with stable hCG release (**Supplemental Figure 1C, 1D**). Collectively, these data establish matched TOs and DOs as model systems of the maternal-fetal interface and that they can be isolated from tissue after the first trimester, including from mid-to-late in gestation.

Cytokine profiling of TOs and DOs compared to matched tissue explants

We have shown previously that human chorionic villous explants release a number of cytokines, chemokines, and growth factors that impact both maternal and fetal antimicrobial defenses (Megli et al., 2021). Similarly, we have shown that primary human trophoblasts (PHTs) isolated from mid-gestation and full-term placentas secrete some of these same immunoactive factors (Corry et al., 2017; Megli et al., 2021). We therefore profiled the immunological secretome of TOs isolated from three independent placentas and compared these profiles to chorionic villous explants isolated from the same placentas. In addition, we compared these profiles to PHT cells to determine whether there were differences between stem cell-derived trophoblasts and primary trophoblasts isolated directly from full-term tissue. To do this, we used multianalyte Luminex-based profiling for ~80 cytokines, chemokines, growth and other factors from conditioned media (CM) isolated from these models

under unstimulated conditions. Factors whose secretion was reliably measured over 50pg/mL were

identified as present. We found that TOs secreted two cytokines (IL-6 and IL-28A/IFN-λ2) and three

immune-regulated secreted soluble receptors (sTNF-R1, gp130/sIL-6Rβ, sTNF-R2) (**Figure 2A**). By

contrast, matched chorionic villous explants were immunologically more active and we reliably

detected 51 factors (**Figure 2B**). Similarly, PHT cells also released a greater number of factors, with

31 detected (**Figure 2C**). The five factors released from TOs were also released from chorionic

villous explants and PHT cells (**Figure 2D**). The most abundant cytokine present in CM isolated from

TOs was IL-6, which was released at levels similar to those observed in both chorionic villi and PHT

cells (**Figure 2E**). In contrast, whereas the most abundant factor in CM isolated from PHT cells was

IL-8, which was also highly abundant in CM from chorionic villi, this cytokine was not detected in CM

isolated from TOs, suggesting that there might be differential immune regulation between terminally

differentiated PHT cells and those that are stem-cell derived (**Figure 2F**).

To define the immunological secretome of DOs, we performed similar Luminex-based

profiling as the studies described above from CM isolated from DOs and decidua explants.

Compared to TOs, DOs released a larger number of factors overall, with 21 detected in DO-derived

CM (**Figure 3A**). Of these, 19 were also released from human decidua explants, which contained

50 total cytokines and other factors (**Figure 3B, 3C**). DOs and decidua explant tissue secreted similar

levels of factors such as MMP-1 (**Figure 3D**). In contrast, we found that DOs secreted high levels of

IL-10, which was not detected in CM from decidua explants (**Figure 3E**). Together, these data

suggest that TOs and DOs partially recapitulate the secretion profile of chorionic villi and the decidua.

Trophoblast organoids constitutively secrete antiviral IFN-λ2

We next directly compared the secretion profiles from CM collected from matched TOs and DOs to

identify factors enriched in either maternal- or fetal-derived cells. TOs selectively secreted two

cytokines that were not present in CM isolated from matched DOs, IL-6 and IFN-λ2 (**Figure 4A-B,**

D). In contrast, DOs selectively released high levels of other factors such as IL-8 (**Figure 4C**). We

showed previously that PHT cells isolated from full-term placentas and mid-gestation chorionic villi

constitutively release type III IFNs, which protect from viral infections (Bayer et al., 2016; Corry et al.,

2017). We found that TOs recapitulated this phenotype and released high levels of IFN-λ2, but not

type I IFNs or IFN-λ1 (**Figure 4E**). In contrast, DOs did not basally secrete any detectable IFNs

(**Figure 2A, 4D**). We monitored the release of IFN-λ2 over the growth period of TOs and found that

it's release was detectable by ~10-days post-plating and reached high levels by >14-days in culture,

which correlated with similar trends for IL-6 (**Figure 4F**). We confirmed that CM isolated from TOs,

but not DOs, induced interferon stimulated genes (ISGs) using a HEK293 based reporter assay

(**Figure 4G**). Consistent with this, we found that CM isolated from TOs was antiviral and reduced

ZIKV infections in non-placental cells treated with this CM (**Figure 4H**). PHT cells and villous explants

basally express high levels of ISGs, most likely from the autocrine effects of released IFN-λs (Corry

et al., 2017). Therefore, we compared the basal expression of two ISGs, IFI44L and OASL, in TOs

and DOs and found that TOs expressed significantly higher levels of these ISGs that do DOs (**Figure**

4I). These data show that TOs recapitulate the constitutive release of type III IFNs observed in

primary trophoblast cells and tissue and that these IFNs function in paracrine to restrict viral

infections.

Differential induction of antiviral signaling in TOs and DOs

Because we observed differences in the constitutive release of IFN-λ2 from TOs and DOs, we next

determined how these organoids responded to viral infections. To do this, we used poly (I:C), a

synthetic ligand of TLR3 and measured differences in cytokine induction using Luminex-based

profiling. We found that both TOs and DOs robustly responded to poly I:C treatment, which

corresponded with the induction of ISGs in both models, although TOs induced higher transcript

levels of these ISGs (**Supplemental Figure 2A**). Cytokine profiling of CM isolated from poly I:C-

treated TOs and DOs showed that TOs induced 15 factors (>5-fold change from mock) and DOs

induced 14 factors (>5-fold change from mock) in response to TLR3 activation (**Figure 5A**). Of the factors induced, 9 were shared between both TOs and DOs (**Figure 5B**), which included IFN-λ1, which was the most induced cytokine in both organoids (**Figure 5C**) and IFN-λ2 (**Figure 5D**). Several cytokines were differentially induced in TOs versus DOs, including IFN-γ, which was induced in DOs but not TOs (**Supplemental Figure 2B)** and IL-8**,** which was induced in TOs but not DOs (**Supplemental Figure 2C).** In addition, we found that whereas TOs induced the type I IFNs IFN-α2 and IFN-β in response to poly I:C, DOs induced no IFN-α2 and low levels of IFN-β (**Figure 5E, 5F, Supplemental Figure 2A**).

We next profiled the differential response to viral infections in TOs and DOs using ZIKV. We found that both TOs and DOs supported ZIKV replication as assessed by the presence of viral RNA by qPCR at 72hrs post-infection (p.i.) (**Figure 6A**). Luminex-based profiling of CM isolated from these cultures at 24hrs, 48hrs, and 72hrs p.i. showed that few cytokines were induced by ZIKV infection and no cytokines or other factors were induced >5-fold (**Figure 6B, 6C**). Consistent with poly I:C treatment, TOs induced very low levels of IFN-λ1 and IFN-β at 72hrs p.i., which was not observed in DOs (**Figure 6D**). Consistent with this, we observed low level induction of ISGs in TOs that was not observed in DOs (**Supplemental Figure 2D**). In contrast, DOs secreted low levels of the glycophosphoprotein osteopontin at 24hrs p.i., which decreased over the course of infection (**Figure 6F**). Taken together, these data show that TOs and DOs differentially respond to viral infections through the release of select cytokines and other factors.

Differential response to IFNs in TOs and DOs

Previous work has shown that chorionic villous and decidua explants respond to treatment with recombinant IFNs, which protect from ZIKV infection (Jagger et al., 2017), although villous explants appear to be less responsive to type III compared to type I IFNs (Yockey et al., 2018). However, given that the tissue used in these studies contains many different cell types, we investigated the

sensitivity of TOs and DOs to recombinant type I (IFN-β) or type III (IFN-λ) treatment. We found that

DOs were highly responsive to both type I and III IFN treatment and robustly induced ISGs including

IFI44L and OASL (**Figure 7A, 7B**). These ISGs were similarly induced by IFN treatment of decidua

explants (**Figure 7C**). Although TOs did respond to both type I and III IFN treatment, the magnitude

of these responses were lower than those in DOs (**Figure 7A, 7B**), which were consistent with the

responses of IFN-treated chorionic villous explants (**Figure 7C, 7D**). These data suggest that there

are differences in the sensitivity of maternal-and fetal-derived cells to type I and III IFNs.

Discussion

Our work presented here establishes fetal- and maternal-derived organoids as models of innate

immune signaling at the maternal-fetal interface. We show that TOs constitutively secrete only two

cytokines, including the pro-inflammatory cytokine IL-6 and antiviral IFN-λ2. In contrast, we show

that DOs are immunologically more active under resting conditions and basally secrete a number of

cytokines not observed in TOs, but do not release IFNs under unstimulated conditions. In addition,

we show that TOs and DOs differentially respond to TLR3 activation and ZIKV infection and induce

organoid-specific cytokines in response to activation of antiviral innate immune signaling. Collectively,

our work suggests that cells that comprise the maternal-fetal interface defend from viral infections in

a cell-type specific manner and highlight the unique contributions of fetal-derived trophoblasts in

antiviral immunity.

The previous establishment of trophoblast and decidua-derived organoids has significantly

expanded the *in vitro* systems that can model the complexity of the maternal-fetal interface (Turco

et al., 2017; Turco et al., 2018). These previous studies utilized tissue isolated from the first trimester

of pregnancy (6-9 weeks gestation for TOs and 8-12 weeks gestation for DOs) (Turco et al., 2017;

Turco et al., 2018). Here, we show that these organoids can also be isolated from later stages of

gestation, including from full-term placental tissue. Moreover, we show that the growth medium and

conditions used to grow organoids from first trimester tissue can be extrapolated to grow organoids from the second and third trimesters of pregnancy. Two-dimensional cultures of stem cell-derived primary trophoblasts are also restricted to tissue obtained in the first trimester (Okae et al., 2018). Given that access to first trimester tissue can be complicated by access an/or regulatory restrictions, our studies provide an additional source for the development of placental organoids from tissue obtained later in gestation. Moreover, the establishment of these organoids from tissues later in gestation may allow for elucidation of the role of maternal-fetal signaling in obstetric related pathologies.

Unlike many other barrier cell types such as the epithelium, human trophoblasts have the unique capacity to robustly secrete cytokines and other immunological factors without stimulation. Primary human trophoblasts from full-term placentas and mid-gestation chorionic villi secrete a number of cytokines that mediate innate defenses from bacteria, parasites, and viruses (Ander et al., 2018; Bayer et al., 2016; Megli et al., 2021). Consistent with this, serum isolated from pregnant women often contains elevated levels of these cytokines, indicating that at least some have reached the systemic circulation (Benyo et al., 2001; Liu et al., 2019; Megli et al., 2021; Vassiliadis et al., 1998). Similarly, the decidua releases specific cytokines and alterations in these profiles may be associated with pregnancy complications including miscarriage (Alam et al., 2008; Denison et al., 1998; Laham et al., 1997; Lash and Ernerudh, 2015; Pitman et al., 2013; von Rango et al., 2003). The decidua also responds to microbial infections via the release of additional cytokines that function in innate defense (Barber et al., 2005; Guzeloglu-Kayisli et al., 2020; Qiu et al., 2009). However, in many cases, these studies were performed on tissue explants that were multicellular in nature, thus preventing an assessment of the contributions of cytokine release from distinct cell types. Similarly, previous work utilizing tissue explants that are treated with recombinant cytokines, synthetic ligands of various innate immune pathways, or infected with pathogens are also unable to define the contributions of distinct cell types in these responses. The use of organoids therefore allows for analyses of cellular responses to infection or innate immune activation from trophoblasts and

decidual glands in primary cell models that contain isolated cell types enriched at the maternal-fetal

interface. An additional strength of the TO model is that trophoblast stem cells differentiate into all

three distinct trophoblast cell types comprising the human placenta (Turco et al., 2018), which is not

possible in many other existing models including isolation of CTBs from full-term placentas.

The constitutive release of IFN-λs from human trophoblasts is a highly unique phenomenon

that runs counter to much of what is understood about antiviral IFN signaling, which is centered on

a requirement for viral sensing to induce this pathway. Our findings from TOs support that IFN

release occurs soon after trophoblast differentiation and is likely an early mechanism used by the

placenta to defend from viral infections. However, our studies indicated that TOs and DOs may be

similarly permissive to ZIKV infection based on levels of vRNA. Previous work has shown that human

CTBs and EVTs are permissive to ZIKV whereas the STB is resistant (Platt et al., 2018; Tabata et

al., 2016). Given that TOs contain EVTs, CTBs, and that the STB faces inward in the organoid

structure, it is possible that these cells support ZIKV infections in TOs. It remains unknown whether

EVTs and CTBs exhibit differential responses to type III IFNs which might impact this differential

susceptibility to ZIKV infection. Interestingly, our data also show that TOs constitutively release IFN-

λ2 and that IFN-λ1 is only induced upon activation of antiviral signaling pathways. IFN-λs genes are

located in tandem on chromosome 19 in humans and the mechanisms that might control the

constitutive release of IFN-λ2 specifically in trophoblasts is unclear. Our studies also show that type

III IFNs are further induced by activation of antiviral innate immune signaling in both TOs and DOs,

with IFN-λ1 being the most induced cytokine detected in both organoid types. However, the type I

IFN IFN-β is induced to greater levels in TOs than DOs, suggesting that at least some of the cells

that comprise the maternal-fetal interface differentially rely on type I and III IFNs for antiviral defenses.

Previous work suggests that type I IFNs damage placental structure and function in mice and

induce alterations in placental structure in *ex vivo* human chorionic villi (Yockey et al., 2018). In

contrast, type III IFN signaling in mice is protective against viral infections and there are no

deleterious effects of treatment of pregnant animals with recombinant IFN-λs (Chen et al., 2017;

Jagger et al., 2017). While the mechanisms for the opposing effects of type I and III IFNs have yet

to be fully elucidated, at least one ISG, interferon-induced transmembrane (IFITM), inhibits fusion of

CTBs into the STB (Buchrieser et al., 2019; Zani et al., 2019), which may indicate that expression of

select ISGs is tightly regulated to prevent IFN-mediated placental damage. Our data in TOs and

DOs suggests that trophoblasts and decidual cells exhibit differential responsiveness to recombinant

IFN treatment, with treatment of TOs inducing lower levels of ISGs than those observed in DOs.

While this might be attributed to reduced expression of the receptors associated with these IFNs in

TOs, there might also be additional mechanisms employed by trophoblasts to protect from IFN- and

ISG-mediated damage.

Our work presented here develops TOs and DOs from mid-to-late gestation placental tissue

and demonstrates that these organoids recapitulate some of the immunologic secretory phenotypes

of placental tissue isolated from matched placental tissue. We further show that TOs and DOs

constitutively secrete unique cytokines and respond to viral infections through the release of

organoid-specific cytokines. Our work also highlights the prominent role of type III IFNs in antiviral

immunity at the maternal-fetal interface and demonstrates that TOs constitutively release IFN-λ2.

Collectively, these studies define the differential responses of key cells comprising the maternal-fetal

interface and suggest that organoid-based models can be used to define innate immune signaling

at this interface.

Materials and Methods

Human samples. All human tissue used in this study was obtained through the UPMC Magee-

Womens Hospital Obstetric Maternal & Infant Database and Biobank after approval was received

from the University of Pittsburgh Institutional Review Board (IRB) and in accordance with the

guidelines of University of Pittsburgh human tissue procurement. Placental tissue used in this study

was collected from the second (23[rd] and 27[th]) and third (38[th]) trimesters, including from dizygotic

twins. Placental tissue was excluded for diagnosis of PPROM, labor, or maternal infection

Derivation and culture of trophoblast organoid (TOs) cultures from human placental samples.

Villous trophoblast stem cells were isolated from fresh placental tissues as described previously

(Turco et al., 2018). Briefly, villi collected from fresh placental tissue were intensively washed, then

were sequentially digested with 0.2% trypsin-250/0.02% EDTA and 1.0 mg/mL collagenase V.

Pooled digests were washed with Advanced DMEM/F12 medium (Gibco 12634-010) and pelleted

by centrifugation, then resuspended in approximate 10 times volume of ice-cold growth-factor-

reduced Matrigel (Corning 356231). Matrigel "domes" (40 µl/well) were plated into 24-well tissue

culture plates (Corning 3526), placed in a 37°C incubator to pre-polymerize for ~2 min, turned upside

down to ensure equal distribution of the isolated cells in domes for another 10 min, then carefully

overlaid with 500 µL prewarmed trophoblast organoid medium (TOM) consisting of Advanced

DMEM/F12 (Life Technologies) supplemented with 1× B27 (Life Technologies), 1× N2 (), 2 mM

GlutaMAX[TM] supplement (), 100 µg/mL Primocin, 1.25 mM N-Acetyl-L-cysteine (Sigma), 500 nM

A83-01 (Tocris), 1.5 µM CHIR99021 (Tocris), 50 ng/mL recombinant human EGF (Gibco), 80 ng/mL

recombinant human R-spondin 1 (R & D systems), 100 ng/mL recombinant human FGF2

(Peprotech), 50 ng/mL recombinant human HGF (Peprotech), 10mM nicotinamide (Sigma), 2 µM Y-

27632 (Sigma), and 2.5 µM prostaglandin E2 (PGE2, Sigma). Cultures were maintained in a 37°C

humidified incubator with 5% CO_2. Medium was renewed every 2-3 days. Small trophoblast organoid

spheroids became visible by approximately day 12 post-isolation. Derived TOs were passaged every

10 to 14 days depending on their size and density. To passage, TOs were released from domes with

ice-cold cell recovery solution (Corning) and then were digested in prewarmed TrypLE Express

(Gibco) or Stem Pro Accutase (Gibco) at 37°C for 5-7 min. Further mechanical dissociation was

achieved with an Eppendorf Explorer Plus electronic pipettes at its "mix" function setting by pipetting

up and down around 400 times through small-bore pipette tips. Dissociated TOs fragment were

collected and washed by centrifuge, then resuspended in fresh ice-cold Matrigel and replated as

domes at the desired density for continuous culture. For freezing TOs, overlaid TOM was aspirated,

TOs with Matrigel were resuspended in CryoStor CS10 stem cell freezing medium (STEMCell

Technologies) frozen at -80 ℃ and then transferred to liquid nitrogen for long-term storage. For

thawing cryopreserved TOs, TOs were thawed as quickly as possible, diluted with 5 times volume of

basic TOM containing Advanced DMEM/F12, 2 mM GlutaMAX supplement, 10 mM HEPES, 1 ×

Penicillin/Streptomycin and centrifuged to pellet. Afterwards, TOs were resuspended in new ice-cold

Matrigel and replated for recovery culture.

Derivation and culture of decidual organoids (DOs) cultures from human decidua samples.

Decidual gland-enriched cell suspension were acquired from fresh decidual tissue as described

previously (Turco, 2017; Turco et al., 2017). Briefly, isolated decidual tissues were cut into small

pieces then digested in 1.25 U/mL Dispase II (Sigma, D4693)/0.4 mg/mL collagenase V (Sigma, C-

9263) solution. Glandular cells were filtered out from the digestion solution supernatant and

resuspended in ice-cold Matrigel (Corning, 356231), plated as domes in 24-well plates (Corning

3526), and finally overlaid with 500 μL prewarmed organoid Expansion Medium (ExM) with the same

composition as previously described(Turco et al., 2017). The growth medium was renewed every 2-

3 days. Mature organoids were passaged by mechanically disruption following TrypLE Express

(Gibco, 12605) digestion every 7-10 days. Other maintenance were similar to those as previously

described (Turco, 2017).

Generation of TO and DO conditioned media (CM). Overlaid TOM and ExM were collected every

2-4 days based on the growth status of TOs and DOs. Media from at least ten timepoints for each

established organoid lines were collected as TO-CM or DO-CM. Non-conditioned medium (NCM)

was TOM or ExM (described above) that had not been exposed to TOs or DOs, respectively.

Cells and Viruses. PHT cells were established and cultured as described previously (Bayer et al., 2016) . Human osteosarcoma U2OS (ATCC HTB-96) and Vero (ATCC CCL-81) cells were grown in Dulbecco's minimal essential medium (DMEM) with 10% FBS and 1% antibiotic. HEK-Blue IFN-λ reporter cells (hkb-ifnl) were purchased from InvivoGen. ZIKV (Paraiba/2015, provided by David Watkins, University of Miami) was propagated in Vero cells. Viral titers were determined by fluorescent focus assay as previously described (Payne et al., 2006) using recombinant anti-double-stranded RNA monoclonal antibody J2 (provided by Abraham Brass, University of Massachusetts). Infection was determined by qRT-PCR as described below. To assess antiviral activity, U2OS cells were exposed to TO or DO CM, or matched NCM, 24 hr prior to infection with ZIKV Paraiba/2015 for 72hrs. ZIKV infection of TOs and DOs was performed using 10^6 FFU virus per well for 24-72hrs, as indicated.

Immunofluorescence microscopy. TOs and DOs were released from Matrigel domes with 1 mL of cell recovery solution (Corning, 354253) per well without disrupting their 3D architecture, then were fixed in 4%PFA for 30 min at room temperature, followed by 0.5% Triton X-100/PBS to permeabilize for 30 min at 4°C. Organoids were washed and blocked in 5%(v/v) goat serum/0.1%(v/v) Tween-20 in PBS for 15 min at room temperature and then incubated with primary antibodies in the above-described blocking solution at 4°C overnight. Organoids were then washed with PBS and then incubated for 1-2 h at room temperature with Alexa Fluor-conjugated secondary antibodies (Invitrogen). Organoids were washed again with PBS and mounted in Vectashield (Vector Laboratories) containing 4′,6-diamidino- 2-phenylindole (DAPI) and were transferred onto microscope slides with large-orifice 200 µL-tips (Fisher Scientific, 02707134). The following antibodies or reagents were used: cytokeratin-7 (Abcam, ab9021), cytokeratin-19 (Abcam, ab9221), E-cadherin (Invitrogen, PA5-85088), Abcam), Alexa Fluor 594–conjugated phalloidin (Invitrogen, A12381), Alexa Fluor Plus 488 Goat anti-Mouse IgG secondary antibody (Invitrogen, A32723), Alexa Fluor 488 Goat anti-Rabbit IgG secondary antibody (Invitrogen, A11034). Images were captured

using a Zeiss LSM 710 inverted laser scanning confocal microscope and contrast-adjusted in

Photoshop or Fiji.

RNA extraction, cDNA synthesis and quantitative PCR. Total RNA was extracted with the Sigma

GenElute total mammalian RNA miniprep kit following the manufacturer's instruction and using the

supplementary Sigma DNase digestion. RNA quality and concentration were determined using a

Nanodrop ND-1000 Spectrophotometer. Total RNA was reverse-transcribed with the iScript cDNA

synthesis kit (Bio-Rad) following the manufacturer's instructions. Quantitative PCR was performed

using the iQ SYBR Green Supermix (Bio-Rad, 1708882) on a CFX96 Touch Real-Time PCR

Detection System (Bio-Rad). Gene expression was determined based on a ΔC_T method normalized

to actin. Primer sequences for actin, IFN-β, IL-28A/IFN-λ2, IFI44L, OASL, GATA3, MUC20, and ZKIV

vRNA have been previously described (Good et al., 2019; Megli et al., 2021). All primer sequences

used in this study can be found in Table S1.

hCG and PSG1 ELISAs and Luminex assays. Human hCG (Sigma, RAB0092) and PSG1 (R&D

Systems, DY6799) ELISAs were performed according to the manufacturer's instructions. Plates

were read at 450 and 520 mm using a Synergy H1 microplate reader. All sample processing was

performed in duplicate and each experiment was performed with at least three biological replicates.

Luminex assays were performed using the following kits according to the manufacturer's instructions:

Bio-Plex Pro Human Inflammation Panel 1, 37-Plex kit (171AL001M; Bio-Rad Laboratories) and Bio-

Plex Pro Human Cytokine Screening Panel, 48-Plex kit (12007283; Bio-Rad Laboratories). The

plates were read on a MAGPIX (EMD Millipore) Luminex machine and analyzed using xPONENT

software (EMD Millipore).

IFN-λ activity reporter assay. For IFN-λ activity detection from CM, HEK-Blue IFN-λ reporter cells

were used according to the manufacture's protocol. Briefly, IFN-λ reporter cells were plated into a

96-well plate, then CM was added and incubated at 37 °C for 24 hr. Supernatants (20 µl) were

collected and then transferred into a 96-well plate and activity measured using QUANTI-Blue

Solution (InvivoGen, rep-qbs) after incubating at 37 °C for 2 hr using a spectrophotometer at 655 nm.

Each CM preparation was run in triplicate.

Recombinant human IFN and poly(I:C) treatment. For Poly (I:C) treatment, TOs or DOs were

incubated with 10 µg/mL Poly (I:C) diluted in organoid complete growth media for ~24 h, then

supernatant was collected for Luminex-based multianalyte profiling or RNA isolated as described for

qRT-PCR. For recombinant human IFN treatment, TOs or DOs were treated with IFN-β (8499-IF-

010; R&D Systems) or IFN-λ3 (5259-IL-025, R&D Systems) for 24 hr, then total RNA was extracted

as described above for following qRT-PCR analysis.

Statistics and reproducibility. All experiments reported in this study have been reproduced using

independent samples (tissues and organoids) from multiple donors. All statistical analyses were

performed using Prism software (GraphPad Software). Data are presented as mean ± SD, unless

otherwise stated. Statistical significance was determined as described in the figure legends.

Parametric tests were applied when data were distributed normally on the basis of D'Agostino-

Pearson analyses; otherwise, nonparametric tests were applied. P values <0.05 were considered

statistically significant, with specific P values noted in the figure legends.

Author contributions LY designed and performed experiments and participated in manuscript

review and writing, CM performed tissue dissections and participated in manuscript review and

writing, and CC designed and performed experiments, participated in manuscript review and writing,

and secured funding.

Acknowledgements

We thank Jon Boyle (University of Pittsburgh) for careful review of the manuscript and the Magee

Obstetric Maternal & Infant (MOMI) Database and Biobank at the Magee Women's Research

Institute (MWRI) for providing placental tissue. This project was supported by NIH AI145828 (C.B.C.)

and the Children's Hospital of Pittsburgh of the UPMC Health System (C.B.C). This project also used

the UPMC Hillman Cancer Center and Tissue and Research Pathology/Pitt Biospecimen Core and

Animal Facility shared resources which are supported in part by award P30CA047904.

Literature Cited

Alam, S.M., Konno, T., Sahgal, N., Lu, L., and Soares, M.J. (2008). Decidual cells produce a heparin-
binding prolactin family cytokine with putative intrauterine regulatory actions. J Biol Chem *283*,
18957-18968.

Ander, S.E., Rudzki, E.N., Arora, N., Sadovsky, Y., Coyne, C.B., and Boyle, J.P. (2018). Human
Placental Syncytiotrophoblasts Restrict Toxoplasma gondii Attachment and Replication and
Respond to Infection by Producing Immunomodulatory Chemokines. mBio *9*.

Barber, E.M., Fazzari, M., and Pollard, J.W. (2005). Th1 cytokines are essential for placental
immunity to Listeria monocytogenes. Infect Immun *73*, 6322-6331.

Bayer, A., Lennemann, N.J., Ouyang, Y., Bramley, J.C., Morosky, S., Marques, E.T., Jr., Cherry, S.,
Sadovsky, Y., and Coyne, C.B. (2016). Type III Interferons Produced by Human Placental
Trophoblasts Confer Protection against Zika Virus Infection. Cell Host Microbe *19*, 705-712.

Benyo, D.F., Smarason, A., Redman, C.W., Sims, C., and Conrad, K.P. (2001). Expression of
inflammatory cytokines in placentas from women with preeclampsia. J Clin Endocrinol Metab *86*,
2505-2512.

Buchrieser, J., Degrelle, S.A., Couderc, T., Nevers, Q., Disson, O., Manet, C., Donahue, D.A., Porrot,
F., Hillion, K.H., Perthame, E.*, et al.* (2019). IFITM proteins inhibit placental syncytiotrophoblast
formation and promote fetal demise. Science *365*, 176-180.

Burton, G.J., Watson, A.L., Hempstock, J., Skepper, J.N., and Jauniaux, E. (2002). Uterine glands
provide histiotrophic nutrition for the human fetus during the first trimester of pregnancy. J Clin
Endocrinol Metab *87*, 2954-2959.

Chen, J., Liang, Y., Yi, P., Xu, L., Hawkins, H.K., Rossi, S.L., Soong, L., Cai, J., Menon, R., and Sun,
J. (2017). Outcomes of Congenital Zika Disease Depend on Timing of Infection and Maternal-Fetal
Interferon Action. Cell Rep *21*, 1588-1599.

Corry, J., Arora, N., Good, C.A., Sadovsky, Y., and Coyne, C.B. (2017). Organotypic models of type
III interferon-mediated protection from Zika virus infections at the maternal-fetal interface. Proc Natl
Acad Sci U S A *114*, 9433-9438.

Denison, F.C., Kelly, R.W., Calder, A.A., and Riley, S.C. (1998). Cytokine secretion by human fetal
membranes, decidua and placenta at term. Hum Reprod *13*, 3560-3565.

Good, C., Wells, A.I., and Coyne, C.B. (2019). Type III interferon signaling restricts enterovirus 71
infection of goblet cells. Sci Adv *5*, eaau4255.

Guzeloglu-Kayisli, O., Guo, X., Tang, Z., Semerci, N., Ozmen, A., Larsen, K., Mutluay, D., Guller, S.,
Schatz, F., Kayisli, U.A.*, et al.* (2020). Zika Virus-Infected Decidual Cells Elicit a Gestational Age-

Dependent Innate Immune Response and Exaggerate Trophoblast Zika Permissiveness: Implication
for Vertical Transmission. J Immunol *205*, 3083-3094.

Haider, S., Meinhardt, G., Saleh, L., Kunihs, V., Gamperl, M., Kaindl, U., Ellinger, A., Burkard, T.R.,
Fiala, C., Pollheimer, J*., et al.* (2018). Self-Renewing Trophoblast Organoids Recapitulate the
Developmental Program of the Early Human Placenta. Stem Cell Reports *11*, 537-551.

Jagger, B.W., Miner, J.J., Cao, B., Arora, N., Smith, A.M., Kovacs, A., Mysorekar, I.U., Coyne, C.B.,
and Diamond, M.S. (2017). Gestational Stage and IFN-lambda Signaling Regulate ZIKV Infection In
Utero. Cell Host Microbe *22*, 366-376 e363.

Laham, N., Brennecke, S.P., and Rice, G.E. (1997). Interleukin-8 release from human gestational
tissue explants: the effects of lipopolysaccharide and cytokines. Biol Reprod *57*, 616-620.

Lash, G.E., and Ernerudh, J. (2015). Decidual cytokines and pregnancy complications: focus on
spontaneous miscarriage. J Reprod Immunol *108*, 83-89.

Lazear, H.M., Schoggins, J.W., and Diamond, M.S. (2019). Shared and Distinct Functions of Type I
and Type III Interferons. Immunity *50*, 907-923.

Liu, J., Hao, S., Chen, X., Zhao, H., Du, L., Ren, H., Wang, C., and Mao, H. (2019). Human placental
trophoblast cells contribute to maternal-fetal tolerance through expressing IL-35 and mediating
iTR35 conversion. Nat Commun *10*, 4601.

McDonagh, S., Maidji, E., Chang, H.T., and Pereira, L. (2006). Patterns of human cytomegalovirus
infection in term placentas: a preliminary analysis. J Clin Virol *35*, 210-215.

Megli, C., Morosky, S., Rajasundaram, D., and Coyne, C.B. (2021). Inflammasome signaling in
human placental trophoblasts regulates immune defense against Listeria monocytogenes infection.
J Exp Med *218*.

Okae, H., Toh, H., Sato, T., Hiura, H., Takahashi, S., Shirane, K., Kabayama, Y., Suyama, M., Sasaki,
H., and Arima, T. (2018). Derivation of Human Trophoblast Stem Cells. Cell Stem Cell *22*, 50-63 e56.

Payne, A.F., Binduga-Gajewska, I., Kauffman, E.B., and Kramer, L.D. (2006). Quantitation of
flaviviruses by fluorescent focus assay. J Virol Methods *134*, 183-189.

Pereira, L., Maidji, E., McDonagh, S., Genbacev, O., and Fisher, S. (2003). Human cytomegalovirus
transmission from the uterus to the placenta correlates with the presence of pathogenic bacteria and
maternal immunity. J Virol *77*, 13301-13314.

Pitman, H., Innes, B.A., Robson, S.C., Bulmer, J.N., and Lash, G.E. (2013). Altered expression of
interleukin-6, interleukin-8 and their receptors in decidua of women with sporadic miscarriage. Hum
Reprod *28*, 2075-2086.

Platt, D.J., Smith, A.M., Arora, N., Diamond, M.S., Coyne, C.B., and Miner, J.J. (2018). Zika virus-
related neurotropic flaviviruses infect human placental explants and cause fetal demise in mice. Sci
Transl Med *10*.

Qiu, X., Zhu, L., and Pollard, J.W. (2009). Colony-stimulating factor-1-dependent macrophage
functions regulate the maternal decidua immune responses against Listeria monocytogenes
infections during early gestation in mice. Infect Immun *77*, 85-97.

Rizzuto, G., Tagliani, E., Manandhar, P., Erlebacher, A., and Bakardjiev, A.I. (2017). Limited
Colonization Undermined by Inadequate Early Immune Responses Defines the Dynamics of
Decidual Listeriosis. Infect Immun *85*.

Sheridan, M.A., Fernando, R.C., Gardner, L., Hollinshead, M.S., Burton, G.J., Moffett, A., and Turco,
M.Y. (2020). Establishment and differentiation of long-term trophoblast organoid cultures from the
human placenta. Nat Protoc *15*, 3441-3463.

Tabata, T., Petitt, M., Puerta-Guardo, H., Michlmayr, D., Wang, C., Fang-Hoover, J., Harris, E., and
Pereira, L. (2016). Zika Virus Targets Different Primary Human Placental Cells, Suggesting Two
Routes for Vertical Transmission. Cell Host Microbe *20*, 155-166.

Turco, M., Gardner, L., Koo, B., Moffett, A., & Burton, G. (2017). Derivation and long-term expansion
of human endometrial and decidual organoids. Protocol Exchange.

Turco, M.Y., Gardner, L., Hughes, J., Cindrova-Davies, T., Gomez, M.J., Farrell, L., Hollinshead, M.,
Marsh, S.G.E., Brosens, J.J., Critchley, H.O.*, et al.* (2017). Long-term, hormone-responsive organoid
cultures of human endometrium in a chemically defined medium. Nat Cell Biol *19*, 568-577.

Turco, M.Y., Gardner, L., Kay, R.G., Hamilton, R.S., Prater, M., Hollinshead, M.S., McWhinnie, A.,
Esposito, L., Fernando, R., Skelton, H.*, et al.* (2018). Trophoblast organoids as a model for maternal-
fetal interactions during human placentation. Nature *564*, 263-267.

Vassiliadis, S., Ranella, A., Papadimitriou, L., Makrygiannakis, A., and Athanassakis, I. (1998).
Serum levels of pro- and anti-inflammatory cytokines in non-pregnant women, during pregnancy,
labour and abortion. Mediators Inflamm *7*, 69-72.

von Rango, U., Classen-Linke, I., Raven, G., Bocken, F., and Beier, H.M. (2003). Cytokine
microenvironments in human first trimester decidua are dependent on trophoblast cells. Fertil Steril
79, 1176-1186.

Weisblum, Y., Oiknine-Djian, E., Vorontsov, O.M., Haimov-Kochman, R., Zakay-Rones, Z., Meir, K.,
Shveiky, D., Elgavish, S., Nevo, Y., Roseman, M.*, et al.* (2017). Zika Virus Infects Early- and
Midgestation Human Maternal Decidual Tissues, Inducing Distinct Innate Tissue Responses in the
Maternal-Fetal Interface. J Virol *91*.

Weisblum, Y., Panet, A., Zakay-Rones, Z., Haimov-Kochman, R., Goldman-Wohl, D., Ariel, I., Falk,
H., Natanson-Yaron, S., Goldberg, M.D., Gilad, R., *et al.* (2011). Modeling of human cytomegalovirus
maternal-fetal transmission in a novel decidual organ culture. J Virol *85*, 13204-13213.

Wells, A.I., and Coyne, C.B. (2018). Type III Interferons in Antiviral Defenses at Barrier Surfaces.
Trends Immunol *39*, 848-858.

Yockey, L.J., Jurado, K.A., Arora, N., Millet, A., Rakib, T., Milano, K.M., Hastings, A.K., Fikrig, E.,
Kong, Y., Horvath, T.L., *et al.* (2018). Type I interferons instigate fetal demise after Zika virus infection.
Sci Immunol *3*.

Zani, A., Zhang, L., McMichael, T.M., Kenney, A.D., Chemudupati, M., Kwiek, J.J., Liu, S.L., and
Yount, J.S. (2019). Interferon-induced transmembrane proteins inhibit cell fusion mediated by
trophoblast syncytins. J Biol Chem *294*, 19844-19851.

Figure 1

Figure 1. Long-term three-dimensional organoids cultures can be established from human mid- and late-gestation trophoblasts and decidua. (A) Schematic representation of the derivation of trophoblast organoids (TOs) and decidual organoids (DOs) from placental tissue. **(B)** Representative bright-field images of TOs and DOs after passaging in complete growth medium TOM and ExM, respectively at 15 days (DO, top) or 32 days (bottom) post-passaging . Scale bar, 50 μm. Black arrows denote placental-like villi extending from TOs (bottom). **(C)** Confocal microscopy for cytokeratin-19 (Cyt-19, green), and actin (red) in a TO. DAPI-stained nuclei are in blue. Scale bar, 50 μm. White arrows denote placental-like villi. **(D)** Confocal microscopy for cytokeratin-7 (cyt-7, green, left), cytokeratin-19 (Cyt-19, green, middle), or E-cadherin (E-cad, grren, right) and actin (red in all) in DOs. DAPI-stained nuclei are in blue. Scale bar, 50 μm. **(E, F)** RT-qPCR analysis for the trophoblast specific marker *GATA3* (E) or the decidual gland marker MUC20 (F) in matched TOs (blue) and DOs (orange). Shown are the mean ± SD levels of expression normalized to *ACTB* and normalized to DOs (E) or TOs (F). **(G)** Levels of hCG in conditioned medium (CM) isolated from TOs (blue) or DOs (orange) as determined by ELISA. Each symbol represents an individual CM

611 preparation. Upper right shows the OTC pregnancy test stripe denoting an hCG positive line in TOs,
612 but not DOs, after being placed into CM. **(H)** Levels of PSG-1 in conditioned medium (CM) isolated
613 from TOs (blue) or chorionic villous explants (orange) as determined by ELISA. In (G) and (H) each
614 symbol represents an individual CM preparation and significance was determined by Mann-Whitney
615 U test. ****, $P < 0.0001$.

Figure 2

658
659

Figure 2. Cytokine secretion profiles from trophoblast organoids, matched placenta tissue, and primary cells. **(A)** Heatmap of twelve conditioned medium (CM) preparations generated from established TO lines analyzed by Luminex-based multianalyte profiling for the indicated cytokines, chemokines, and growth factors. Scale is shown at right. **(B)** Heatmap of CM generated from placental chorionic villi explants matched to TOs shown in A. Luminex assays were performed from three independent CM preparations. Scale is shown at right. **(C)** Heatmap of CM generated from primary human trophoblast cells cultured from full-term placentas. Luminex assays were performed from five independent CM preparations. Scale is at right. **(D)** Venn diagram showing the overlap of detectable factors in TOs, matched placenta chorionic villi, and primary cells from A-C. **(E, F)** Concentrations and comparison of the levels of IL-6 (E) and IL-8 (F) in CM from TOs (grey), matched chorionic villi (dark red), and primary cells (blue) shown in A-C. Each symbol represents a unique

671   CM preparation. Significance was determined by Kruskal Wallace U test with multiple comparisons.
672   **, P < 0.01; ns, not significant.

Figure 3

Figure 3. Cytokine secretion profiles from decidua organoids and decidua tissue. **(A)** Heatmap of twelve CM preparations generated from established DO lines analyzed by Luminex-based multianalyte profiling. Scale is at right. **(B)** Heatmap of CM generated from three independent decidual explants (two of these, at top and middle, are matched to DOs shown in A). Luminex assays were performed from three independent CM preparations. Scale is shown at right. **(C)** Venn diagram showing the overlap of detectable factors in CM isolated from DOs and decidual tissues. **(D, E)** Concentrations and comparison of the levels of MMP-1 (D) and IL-10 (E) in CM from DOs (orange) and decidual tissues (grey) shown in A and B. Each symbol represents a unique CM preparation. Significance was determined by Mann-Whitney U test. *, $P < 0.05$; ns, not significant.

Figure 4

Figure 4. Trophoblast organoids preferentially secrete high levels of antiviral IFN-λ2. **(A)** Venn diagram comparing the factors identified in CM isolated from TOs and DOs. **(B, C)** Concentrations and comparison of the levels of IL-6 (B) and IL-8 (C) released by TOs (blue) and DOs (orange) determined by Luminex-based profiling. Each symbol represents a unique CM preparation. **(D)** Comparison of the levels of IFN-λ2 in CM isolated from TOs (blue) and DOs (orange) as determined by Luminex. **(E)** Comparison of the levels of the type I IFNs IFN-α2 and IFN-β and type III IFNs IFN-λ1 and IFN-λ2 in CM isolated from TOs. **(F),** Concentrations of IL-6 (blue) and IFN-λ2 (red) in CM generated from a single TO line throughout the first 25 days of culture. **(G)** IFN-λ activity of CM isolated from TOs (blue) and DOs (orange) as assessed using an IFN-λ reporter 293T cells line. NCM (non-conditioned medium, grey) was as negative control, NCM supplemented with recombinant human IFN-λ2 (100ng) was used as a positive control (red). **(H)** Human osteosarcoma

U2OS cells were treated with CM isolated from TOs (blue) or with recombinant IFN-λ2 (100ng) for 24hrs and then infected with ZIKV for 24hrs (in the presence of CM). Level of ZIKV replication was assessed by viral RNA as determined by qPCR. Data are shown as fold change from NCM and from at least three independent CM preparations. **(I)** RT-qPCR analyses for two ISGs, IFI44L and OASL in TOs (blue) and DOs (orange). Data are shown as fold change from DOs and from at least three independent replicates. In B–H, each symbol represents unique CM preparations. In I, each symbol represents individual replicates (technical and experimental). Significance was determined by Mann-Whitney U test or by Kruskal Wallace U test with multiple comparisons. *, P < 0.05; **, P < 0.01; ****, P < 0.0001.

801
802
Figure 5. Differential cytokine release in TOs and DOs treated with poly I:C. (A) Heatmaps
demonstrating the induction of factors at left (shown as fold change from mock-treated controls) from
TOs (left panel) and DOs (right panel) treated with 10μg poly (I:C) and analyzed by multiplex
Luminex-based profiling. AVG denotes the average change in concentration of factors over CMs
from individual preparations. Dark blue denotes significantly induced factors compared with
untreated controls. Gray or white denotes little to no change (scale at top left). The red arrow
demonstrates factors with induction greater than five-fold change observed in the average of
separate experiments. Data are from four individual CM preparations from at least two unique
matched organoid lines. **(B)** Venn diagram showing overlap of the factors released >5-fold by poly
I:C treatment in TOs and DOs. **(C-F)** Levels of IFN-λ1 (C), IFN-λ2 (D) IFN-α2 (E), and IFN-β (F) in
CM isolated from poly I:C-treated TOs (blue) and DOs (orange) or in mock-treated controls (grey).
In C-F, each symbol represents individual CM preparations. In C-F significance was determined
using a Mann-Whitney U test. *, $P < 0.05$; ns, not significant.
816
817
818

Figure 6

Figure 6. **Differential cytokine release in TOs and DOs infected with ZIKV**. **(A)** ZIKV replication in TOs (blue) and DOs (red) was assessed by the production of vRNA by RT-qPCR 72hrs post-infection. Data are shown as fold change from mock controls. **(B, C)** Heatmaps demonstrating the induction of cytokines in CM (shown as fold change from mock controls) collected from TOs (B) and DOs (C) infected with ZIKV for indicated hours (24 to 72 hours) analyzed by multiplex Luminex-based profiling. Scale is shown at right. **(D, E)** Concentrations of IFN-λ1 (D) and IFN-β (E) in CM generated from TOs and DOs infected with ZIKV for 72 hrs (dark red) or in mock-treated controls (grey). **(F)** Concentrations osteopontin in CM generated from DOs infected with ZIKV for the indicated time (dark red) or in mock infected controls (mock, grey). In D-F, each symbol represents individual CM preparation. Significance was determined with Mann-Whitney U test. ns, not significant.

846   **Figure 7**

848
849

Figure 7. **Responses of TOs and DOs and placental tissue to recombinant IFNs. (A, B)** RT-
qPCR analyses for two ISGs, IFI44L (A) and OASL(B) in TOs (blue) and DOs (orange) treated with
100ng recombinant human IFN-β or IFN-λ3 treatment for ~24 hours. Data are shown as fold change
from mock-treated controls two independent replicates performed in triplicate. **(C, D),** RT-qPCR
analyses for IFI44L (C) and OASL(D) in chorionic villous (blue) or decidua (orange) explants treated
100ng recombinant human IFN-β or IFN-λ3 treatment for ~24 hours. Significance was determined
by Kruskal Wallace U test with multiple comparisons. *, P < 0.05; ***;**, P < 0.01; P < 0.001; ns, not
significant.

Table S1. qPCR primers used in this study.

Target	Forward, 5'→3'	Reverse, 5'→3'
Actin	ACTGGGACGACATGGAGAAAAA	GCCACACGCAGCTC
IFN-β	GAGCTACAACTTGCTTGGATTC	CAAGCCTCCCATTCAATTGC
IFN-λ2	ACATAGCCCAGTTCAAGTC	GACTCTTCTAAGGCATCTTTG
IFI44L	AACCTAGACGACATAAAGAGG	CTGAAACCAAGTCTGCATAG
OASL	GTACCAGCAGTATGTGAAAG	ATGGTTAGAAGTTCAAGAGC
GATA3	AAAATGAACGGACAGAACC	GGGGTCTGTTAATATTGTGAAG
MUC20	CTCTTCCTTCTGTCCCTC	CCATTTTCACCTGCACTC
ZIKV	AGATGACTGCGTTGTGAAGC	GAGCAGAACGGGACTTCTTC

903
904 **Supplemental Figure 1**

906
Supplemental Figure 1. **(A),** Brightfield image of a TO at 7 or 14 days post-plating. Black arrows
denote EVTs that migrate from the organois to invade surrounding Matrigel. **(B),** Confocal
micrograph of EVTs growing from a TO and immunostained for cytokeratin-19 (green) and actin
(red). DAPI-stained nuclei are shown in blue. **(C, D)** hCG levels as assessed by ELISA in
conditioned medium isolated from two TOs lines isolated at the indicated weeks during culture.
.

927
928
929 **Supplemental Figure 2**
930
931

932
Supplemental Figure 2. (A), RT-qPCR for IFN-β, IFNλ-2, IFI44L, and OASL in poly I:C treated TOs
(blue) and DOs (orange). Data are shown as mean ± standard deviation as a fold change from mock-
treated controls. **(B, C),** Levels of IFN-γ (B) or IL-8 (C) as assessed by Luminex in TOs (blue) and
DOs (orange) treated with poly I:C or in mock-treated controls (grey). Data are shown as mean ±
standard deviation. Symbols represents unique organoid preparations. **(D),** RT-qPCR for IFN-β,
IFNλ-2, IFI44L, and OASL in ZIKV-infected TOs (blue) and DOs (orange) 72hrs post-infection. Data
are shown as mean ± standard deviation as a fold change from mock-infected controls. Significance
was determined with Mann-Whitney U test. *, p<0.05, ns, not significant.