Title: Interactions, information and emergence: Exploring task allocation in ant colonies

using network analysis

List of Authors: Anshuman Swain*[1#], Sara D. Williams*[2], Louisa J. Di Felice*[3] and

Elizabeth A. Hobson[4]

Institutional affiliations: [1]Department of Biology, University of Maryland, College Park,

MD, USA; [2]Mote Marine Laboratory, Sarasota, FL, USA; [3]Institut de Ciencia i

Tecnologia Ambientals, Universitat Autònoma de Barcelona, ES; [4]Department of

Biological Sciences, University of Cincinnati, OH, USA; *contributed equally*

Contact Information: [#]answain@terpmail.umd.edu

ABSTRACT

11
In animal societies, individuals may take on different roles to fulfil their own needs

and the needs of their groups. Ant colonies display high levels of organisational

complexity, with ants fulfilling different roles at different timescales (what is known as *task*

allocation). Factors affecting task allocation can be at the individual level (e.g.,

physiology), or at the group level (e.g., interaction histories). In this work, we focus on

group level processes by exploring the impact of the history of interaction networks on

task allocation and task switching using a previously published dataset (Mersch et al.,

2013) tracking the behaviour of six *Camponotus fellah* colonies over 41 days. First, we

investigated the architecture of interaction networks using node (individual) level network

measures and their relation to the individual's task – foraging, cleaning or nursing – and

whether or not the ant switched tasks. We then explored how noisy information

propagation is among ants, as a function of the colony composition (how many ants are

carrying out which tasks), through the information-theoretic metric of effective information.

Our results show that interaction history affected task allocation, with ants more likely to

switch to a task if they had interacted with other ants carrying out that task. The degree

to which interaction history affected task allocation, as well as the noise in their

interactions, depended on which groups of ants are interacting. Overall, we showed that

colony cohesion is stable even as ant-level network measures vary more for ants when

they switched functional groups; thus ant colonies maintain a high level of information

flow as determined by network analysis and ant functional groups play different roles in

maintaining colony cohesion.

Keywords: Ant behaviour, effective information, emergent behaviour, insect social

networks, task allocation, information flow

Highlights

• We analysed the interaction networks of six *Camponotus fellah* colonies

• We tested how centrality and information flow affected task switching

• Node-level network metrics and the information theoretic measure of effective

information explain differences among functional groups

• Interaction histories predicted task switching, but the strength of the effect

differed across functional groups

INTRODUCTION

In animal societies, individuals may carry out different tasks to fulfil their own needs and the needs of their group (Sumpter, 2006; Clutton-Brock, 2009; Jeanson & Weidenmuller, 2014). Larger and more complex societies can self-organise to fulfil tasks beyond basic sustenance and reproduction (Boomsma and Frank, 2006; Sumpter, 2010). Local exchange of information, between individuals of a group and between individuals and their environment, is key to self-organisation (Sumpter, 2006; Boomsma and Frank, 2006; Couzin, 2009; Cavagna et al., 2010; Swain and Fagan, 2019). Social insect colonies display high levels of organisational complexity (Lukas & Clutton-Brock, 2018), where individual tasks may include foraging, nest construction and caring for the young (Gordon, 2002). The assignment of tasks, also referred to as *task allocation*, is the result of patterns of factors that vary across different scales (Gordon, 2015). These tasks can be fixed throughout each individual's lifetime due to physiological reasons, for example when only a fertile subset of the population is responsible for reproduction, or when a subset is responsible for providing food (Sumpter, 2010; Clutton-Brock et al., 2001).

Task allocation can also result in individuals changing their main task over time. Factors affecting task changes can occur at the individual level or at the group level. Individual-level factors include age, corpulence or physiology (Anderson and Shea, 2001; Tripet and Nonacs, 2004; Robinson et al., 2009). Studying individual-level factors associated with task change is often simpler than studying group-level ones. For example, individual-level changes can be easier to track because their rate of change is slower and often follows a consistent and predictable pattern, as in the case of ageing. Individual-level factors can also be directly quantified, e.g., by measuring age, corpulence, or

physiological features, and traditional statistical approaches can be used to predict task

changes.

Changes in task allocation affected by group-level factors are currently not well

understood. Structural features or macro-level social properties of groups can affect

micro-level individual actions if the social system is affected by feedbacks (Flack, 2017;

Hobson et al., 2019). However, quantifying relevant macro-to-micro feedbacks can be

challenging and can require large amounts of data. The development of automated

tracking systems has made this level of data collection possible. While these systems

have improved researchers' ability to track detailed social behaviour (Robinson et al.,

2009; Smith and Pinter-Wollman, 2021), assigning quantitative metrics to group dynamics

is still a non-trivial task. In the case of interaction patterns, tracking physical interactions

among individuals does not necessarily map onto the amount of meaningful (predictive)

information exchanged with each interaction (Valentini et al., 2020). Although tracking

technologies can tell us how many times individuals in a social group interact with one

another, they cannot explain to what extent these interactions drive task allocation.

Task allocation in ants has been the subject of much previous work (Anderson and

Shea, 2001; Gordon, 2015). Across ant species, studies have shown that, depending on

the tasks and on the colony, ants may display varying degrees of task flexibility, from

small colonies of totipotent ants to larger ones with a structured division of labour

(Anderson and Shea, 2001). Individual-level factors include physiology (Anderson and

Shea, 2001), age (Tripet and Nonacs, 2004), corpulence (Robinson et al., 2009) and past

experience (Ravary et al., 2007), whereas group-level factors involve colony size (Ravary

et al., 2007) and short-term interactions (Gordon and Mehdiabadi, 1999). However, we

do not possess a good understanding of how the nature of past interactions and topology

of the social interaction network structure affect an ant's propensity to switch to a new

task.

In this paper, we leverage social network methods to gain new insight into task

allocation changes in an existing dataset of ant interactions (published by Mersch et al.

2013). Mersch et al. studied task switching in *Camponotus fellah* by tracking and

analyzing the movements and interactions of individually-identified ants. Worker ants

were categorized into three *functional groups* (nurse, cleaner or forager). Analyses

showed that the ants had more interactions with others in their same functional group.

Communities defining the functional groups exhibited distinct behavioural signatures and

were highly spatially divided. Nurses spent most of their time with the brood, while

foragers spent time at the nest entrance and cleaners were located between the other

two groups and the rubbish pile (Mersch et al. 2013). Mersch et al. also explored the

questions of task switching cost, i.e., a time and energy investment associated with

learning new tasks (Goldsby et al., 2012), and of age polytheism, i.e., the correlation

between the age of an ant and which task they perform. The original study identified

spatial fidelity as a key regulator of ant social organization and interaction frequency

(Mersch et al. 2013). They also found that task switches were present but uncommon

and that when a shift in functional group occurred, ants showed a preferred direction of

task transition, from nurses to cleaners to foragers, mostly based on age (Mersch et al.,

2013). Task changes were thus hypothezied to be driven by age polyethism, but the

patterns were fairly noisy.

In this new analysis we focus specifically on this noisy process of task switching.

A question not addressed in the previous study is whether the history of an ant's

interactions with others and the resulting information flow within the colony could be one

of the elements explaining task switching. In other species, information flow patterns have

been shown to affect task allocation, interactions, and overall colony behavior, such as in

the case of midden workers in red harvester ants (*Pogonomyrmex barbatus*; Gordon and

Mehdiabadi, 1999, Pinter-Wollman et al., 2018), tandem running recruitment (Franklin

and Franks, 2012) and consensus-forming in rock ants (*Temnothorax albipennis*; Sasaki

& Pratt, 2018). To test whether the history of interactions or information flow could explain

the noise seen in task switching dynamics that was not explained by age polyethism

alone, we evaluate several potential macro-scale predictors of task switching not

addressed in the original paper.

First, we described the architecture of the interaction networks by focusing on

information flow (which in our case refers to the possible information exchange due to

interactions among ants). We tested whether the role played by individual ants in

regulating information flow in the colony and the functional group that they belong to were

correlated. To do this, we quantified three network measures that are tied to the

architecture of information flows at the local level for ant-ant interactions, strength,

betweenness centrality, bridge betweenness centrality; and a new measure, *effective*

information (EI) at the global level for the whole colony. While strength, betweenness

centrality and bridge betweenness centrality are common node-level measures in network

science and have been applied to animal social networks in the past (Holme et al., 2002;

Lusseau and Newman, 2004; Krause et al., 2009; Farine and Whitehead, 2015), EI is

central to our analysis and is a novel information theoretic metric reflecting how noisy a

mechanism connecting nodes (ants, in our case) is within a system. It is calculated by

perturbing a system through its repertoire of possible states and evaluating the effects of

the perturbation on the system's mechanisms (Hoel, Albantakis, & Tononi, 2013; Klein

and Hoel 2020). In interaction networks, EI reflects the noisiness of the interactions

among individuals (Hoel et al., 2020): a higher EI means that a system is more

deterministic, with information spreading in a more effective way throughout the network.

Second, we tested whether these four measures of information flow in the

interaction network were correlated with task switching to better understand the noise in

task allocation not explained by age polyethism as determined by Mersch et al. (2013).

We hypothesized that an ant's previous interactions with other ants affect switching

behaviour and tested whether interacting with a certain functional group increased the

probability of an ant then switching to that group. The relationship between the

architecture of information flows and the different functional groups, as described by

network measures, inform our understanding of the varying correlations between

interaction history and switching behaviour during task allocation. Our use of network

metrics, including the novel effective information metric, allowed us to determine the

influence of interaction history on task allocation and information flow among functional

groups in *Camponotus fellah* colonies.

METHODS

Data, network construction, and functional group assignment

The published Mersch et al. (2013) dataset contains summaries of interactions

among a total of 985 individually-marked ants in six *Camponotus fellah* colonies. The

authors collected interaction data for every pair of ants at a daily resolution over the 41-

day monitoring period, and the published dataset contains data pooled at the number of

interactions per dyad per day per colony. We matched this published dataset with the

colony metadata to inform our analyses (Supplementary material 1).

Consistent with Mersch et al. (2013), we used the pairwise daily number of

interactions to construct separate weighted, undirected, unipartite networks for each

colony per day. Each ant in a colony was represented by an individual node. An edge

between two nodes represents an interaction between those two ants on a given day.

The edge weight is proportional to the number of pairwise interactions between them on

that particular day. We used the available published dataset to recreate the 246 networks

for the 6 colonies over 41 days used by Mersch et al. (2013) as well as the general pattern

of task switching across the length of the experiments.

Mersch et al. (2013) assessed each ant's functional group every 10 days to

categorize them as a nurse, cleaner, or forager, representing their main task in the colony.

They assigned functional groups based on what community an ant spent at least 70% of

their time in, using the 'infomap' community detection algorithm. They split the ants into

the functional groups foragers (F), cleaners (C), nurses (N), queen (Q), and NA for ants

who were counted as missing at a time point (e.g., if they were dead or had lost their

tags). In our analyses, we used a similar method to assess functional groups and split the

ants into the same five groups used in the original study via community detection.

However, unlike the Mersch et al. approach, we assigned community membership using

a Louvain community detection algorithm (Csardi and Nepusz, 2006). This algorithm was

used as it performs better than the infomap algorithm on multiple metric tests (Emmons

et al., 2016). This approach resulted in very high agreement with previous task

assignments, as we will show in the Results.

Mersch et al. (2013) reported that their ants mostly did not change their task affiliation

within the 10-day observation period between task assessment points. We used the same

10-day snapshot window in our analyses which resulted in three time points at which a

switch in task to a new category could be detected. We assess whether ants switched

tasks by comparing the functional group assignments for each ant in one time period to

its assigntment in the next time period. For our analyses, we categorized each ant as

"switched" or "consistent", depending on whether they remained within the same

functional group, or were categorized as part of a different functional group.

Quantifying individual network metrics for each ant

Node metrics and centralities define various types of influence that individual nodes

exert on network connectivity and dynamics. For each network, we used R (v 3.6.2) and

the packages *igraph* (Csardi and Nepusz, 2006) and *networktools* (Jones, 2020) to

calculate three node-level, local metrics: strength, betweenness centrality, and bridge

betweenness. These local measures were calculated for every ant on each day of the

experiment for all six colonies.

Node strength was calculated as the sum of the weights of a node's edges. Thus, in

our context, it is a measure of not only how many interactions (edges) an ant (node) had

to other ants, but also of how frequently those interactions occurred during a day. While

degree is an index of potential communication activity (Freeman, 1979), strength

improves upon this index by weighting degrees according to frequency of communication

to better inform total interaction and information flow potential.

Node betweenness, also known as betweenness centrality, is a measure of

importance in a network based on shortest paths between pairs of nodes. For a given

pair of nodes in a weighted network, there exists at least one path between them such

that the sum of the link weights is minimized, thus forming a shortest path. The

betweenness of a node is therefore defined as the number of shortest paths that pass

through it. Freeman (1979) identified high betweenness centrality as a key indicator of

whether a node occupies a central location in the network for information transmission.

An ant with a high betweenness is an ant that is centrally located in the network, serving

as a key connection for seemingly disparate ants. Individuals with high betweenness are

responsible for maintenance of communication, group coordination, and network stability

(Lusseau and Newman, 2004; Farine and Whitehead, 2015).

Bridge betweenness extends the betweenness centrality metric to the level of

communities and is defined as the number of times a node lies on the shortest path

between two nodes from different communities. In network science, a community is

defined as a group of nodes that have a higher likelihood of connecting to each other than

to nodes from other communities. Ants with a high bridge betweenness serve as key

connectors for different groups in the network. This means that they are more integral to

network cohesion and information flow across groups, thus they may play an important

role in driving switching dynamics. We calculated bridge betweenness for ants using the

community structure determined when assigning functional groups via the Louvain
community detection algorithm.

We generated randomized networks to serve as null models for the daily interaction
networks of the six colonies. In order to preserve the underlying structure of the networks,
a degree-based randomization was used (through the R package *VertexSort*; Abd-Rabbo,
2017) to randomize the ant-to-ant interactions. We created 500 seeded null networks for
each colony's daily interaction network, for a total of 123,000 null networks. All metrics
calculated for the empirical ant-to-ant interaction networks were then calculated for this
new ensemble of null networks. Daily node-level metrics were compared to the
distribution of values determined by the null networks. Values were then Z transformed
to be compared across groups and measures.

Quantifying global network measures for each colony

240
We used Effective Information (EI) and its normalized measure of effectiveness to
measure colony-level noisiness in the system, with respect to its underlying mechanisms
(Hoel et al., 2020). Since we are considering the mechanism of communication and
information flow among ants, EI measures the level of noisiness in ant-to-ant interactions.
To calculate effectiveness, we first characterized the weight of the edges connected to a
node. We defined this weight as a vector W_i of the same length as the total number of
nodes, and referred to each element as ω_{ij}, signifying the normalized value of edge
weight between nodes *i* and *j*, such that for any index *i*, $\sum_j \omega_{ij} = 1$. Here, each term ω_{ij}
can be seen as the probability of moving from *i* to *j*, if a random walker is on the node *i*.
Next, we characterized the uncertainty associated with each node *i*, calculated using

Shannon's entropy measure $H(W_i)$. The average of this value across all the nodes in the

network is $< H(W_i) >$. This means that, if $< H(W_i) >$ is equal to 0, the network is

deterministic. We then assessed the certainty of the network by calculating the term

$H(< W_i >)$, which is Shannon's entropy of the average out-weights from nodes. If this

expression is equal to 0, the network is degenerate, with all edges leading to the same

node. Finally, we calculated EI using the following equation:

$$EI = H(< W_i >) - < H(W_i) > \tag{1}$$

258 As the value of EI can depend on the size of the network (Klein and Hoel, 2020), we

259 calculated effectiveness, the normalized EI with respect to network size, where N is the

260 number of nodes in the network:

$$Effectiveness = \frac{EI}{log_2 N} \tag{2}$$

Effectiveness was calculated for each constructed network (i.e., for each day, for every

colony), using the R package *einet* (Byrum et al., 2020). Linear models were fit to the

effectiveness for each constructed network as a function of the proportion of each

functional group in the network to determine significant relationships between

effectiveness and a colony's functional group composition.

Task interaction matrix and task switching

269
To investigate patterns of ants switching between functional groups and taking on

new colony tasks, we first tested whether we could replicate Mersch et al.'s (2013) results

of age polyethism using our task assignment method, i.e., the Louvain community

detection algorithm. We determined the probability that an ant would switch tasks once,

twice, or three times, against the age of the ant (Supplementary material 2, Figure S1A).

We further explored the possibility of a switching cost by determining the likelihood that

an ant would stay in the same task throughout the experimental time versus performing

two or three tasks (Supplementary material 2, Figure S1B).

If functional group identity of ants affected how individuals contributed to information

flow within a colony, then the number of times an ant switched to a new functional group

also affected the flow of information. Here we asked a question not addressed in the

original paper: if the number of interactions per ant were kept constant, and the

interactions were redistributed among everyone, did the proportion of interactions with

different functional groups significantly affect an ant's final functional group, as opposed

to what we saw in the original network? To test this, we quantified the average proportion

of an ant's interactions with each functional group before switching from its original

functional group to another in both the observed dataset and in the ensemble of random

networks described above. We compared the distribution of values computed from the

actual networks against those given by the null network distribution using a chi-square

distribution for all possible types of transition (including non-transitions), and calculated

differences significant at the alpha level of 0.05.

RESULTS

Functional group assignment comparison

Comparisons of our functional group assignments (via the Louvain community

detection algorithm) with groups assigned by Mersh et al. (2013; via infomap) showed

high levels of overall agreement. Community membership assignments from infomap and

Louvain, compared at an individual node level for a given network, resulted in an average

90.13 ± 7.25% similarity between the two methods across all the networks in the dataset.

Figure 1 shows a summary of the tasks of ants within all six colonies and how those tasks

changed over time (Figure 1; for details, see Supplementary material 2, Table S1).

Figure 1: Dynamics of task allocation across the experimental time for all ants in all six colonies, with functional group assigned via Louvain community detection algorithm (~90% assignments consistent with Mersch et al. 2013). The alluvial diagram shows the number of ants per functional group and number of ants staying in the same group or transitioning to a new functional group between time periods as proportional to box and flow sizes, respectively.

Individual network centrality measures and task switching

We compared network measures (and the variance in the measures) across each of

the functional groups for three categories: overall across all ants, for just switching ants,

and for just ants that remained consistent in their tasks during the assessment periods

(summarized in Figure 2; all values listed in Supplementary material 2 Table S2).

When we compared strength across ants in each of the tasks, we found that foragers

had the highest mean strength of any of the groups across all three of the categories,

showing that they had the most frequent interactions over a day regardless of whether

they remained foragers or switched task at some point. When considering the overall

mean strength, foragers had the strongest interaction potential (2.96). The strength of

cleaners (2.35) and nurses (2.25) were not significantly different from each other but were

still significantly greater than the null networks' strength (Figure 2A1). Overall strength

variance did not significantly differ across functional groups or from the null networks

(Figure 2A2), indicating that strength values remained fairly stable through time at the

colony level. When we looked at strength just for switching ants, we found that the mean

strength differed significantly across functional groups and was significantly greater than

for the null networks (Figure 2A3). Out of these, foragers that switched had the highest

strength (3.17), while cleaners that switched had a higher strength (2.92) than nurses that

switched (2.35). Strength variance of switching ants did not vary significantly among

functional groups or from the null networks (Figure 2A4). When we looked at strength just

for ants that were consistent, we found that the mean strength and strength variance

followed the same pattern seen for ants that switch (Figure 2A5 and 2A6).

When we compared betweeness across ants in each of the tasks, we found that

cleaners (2.73) had significantly higher betweenness than nurses (2.25) who also had

significantly higher betweenness than foragers (2.01, no different from null) and the null

networks (Figure 2B1). Overall, variance of betweenness did not differ among functional

groups or from the null networks (Figure 2B2). At the colony level, the betweenness metric

was stable and the cleaners played the most important role in connecting individual ants

for flow of information. When we looked at betweenness just for ants that switched, we

found that mean betweenness centrality measures were significantly greater than those

for the null networks (Figure 2B3). Cleaners (2.33) and nurses (2.29) that switched had

higher betweenness than foragers that switched (1.83). Variance of betweenness was

greater for foragers (2.67) and cleaners (3.01), however the variance of nurses was still

greater than that of the null networks (2.14, Figure 2B4). Betweenness of foragers and

cleaners was more unstable before these ants switched tasks. When we looked at

betweenness just for ants that were consistent, we found that they had the same patterns

of mean betweenness centrality (Figure 2B5). However, the variance of betweenness

was no longer significantly different than the null networks, thus consistent ants

maintained a stable betweenness position through time (Figure 2B6).

Since the communities we detected mapped primarily onto the previously determined

functional groups, a high bridge betweenness indicated a high potential for connecting

functional groups in a colony. When we compared bridge betweenness across each of

the tasks at the colony level, we found that the overall mean bridge betweenness values

did not vary among the functional groups or from the null networks (Figure 2C1).

Additionally, the overall variance remained similar across the functional groups and did

not vary significantly from the null networks (Figure 2C2). When we looked at bridge

betweenness for ants that switched compared to those that were consistent, we found

that mean bridge betweenness was higher for the switching ants for all functional groups,

though only significantly higher for foragers (Figure 2C 1,3,5). All ants that switched had

significantly higher mean bridge betweenness than the overall colony values per

functional group, suggesting that ants that switched played an important role in

connecting communities for information flow in the colony. Within the ants that switched,

foragers had the highest mean bridge betweenness (2.97), though they were not

significantly different from cleaners (2.63). Both foragers and cleaners had higher mean

bridge betweenness than nurses (2.22). All mean bridge betweenness values of ants that

switched were higher than the null networks. The variances of foragers (2.59) and

cleaners (2.41) that switched were greater than nurses (1.99) that switched (Figure 2C4).

Only the variance of foragers was significantly greater than that of nurses, which did not

differ from the null networks. The mean bridge betweenness of consistent ants did not

vary significantly among the functional groups or from the null network distribution (Figure

2C5). The variances of consistent cleaners (2.91) and foragers (2.75) were significantly

higher than nurses (2.11) and the null networks (Figure 2C6). Although the variance of

cleaners and foragers for both ants that switched and consistent ants did vary, the overall

colony values remained stable; these results may indicate that these stuctures could be

important for colony cohesion at the community level.

Figure 2: Z transformed values of the mean and variances of the strength, betweenness and bridge betweenness (columns A, B and C respectively) determined for all ants across the six colonies. The first two rows (rows 1 and 2; A1-C2) are values for all the ants in the experiment (overall colony values). The next two rows (rows 3 and 4; A3-C4) show values

for ants that switched during the experiment, and the last two rows (rows 5 and 6; A5-C6)
show values for for consistent ants (no switching) during the experiment. Line colors
correspond to ant functional group type: nurses (red), cleaners (blue), and foragers
(green). Shaded areas are the 95% confidence intervals based on seeded random
networks. The null distribution created from the null network simulations is shown in black
with a marked confidence interval (dashed lines).
379

Global information flow and task switching

We measured effectiveness as a function of the proportion of nurses, cleaners or

foragers in each colony for each day (resulting in 246 effectiveness measures, Figure 3).

We found that the colony networks with high proportions of nurses and cleaners had lower

effectiveness, but that the dependencies based on the linear model were weak and non-

significant (adj. R^2=0.12, P = 0.063, Figure 3A for nurses; and adj. R^2=0.11, P = 0.052,

Figure 3B for cleaners). Effectiveness significantly decreased with increasing proportions

of foragers in a colony (adj. R^2=0.22, P = 0.037, Figure 3C). This negative relationship

between increased numbers of foragers and decreased colony-level effectiveness shows

that the interactions mediated by foragers were noisier than those of nurses or cleaners.

Figure 3: Effectiveness (normalized EI) of the interaction networks constructed for each colony and every day of the experiment as a function of the proportion of different functional groups in the networks. Data are stacked because the available granularity for task allocation was at a 10-day interval. Linear models fit to effectiveness as a function of the proportions of nurses (A) and cleaners (B) separately return a nominally positive dependence (adj. R^2=0.12, P = 0.063 for nurses; and adj. R^2=0.11, P = 0.052 for cleaners). Effectiveness as a function of the proportion of foragers (C) returns a strong negative dependence (R^2 =0.22, p-value = 0.037).

Task interaction matrix and task switching

We tested whether previous interaction patterns affected switching behaviour using a task interaction matrix. We found that ants that remained consistent in their tasks usually interacted most with other ants occupying their same task (Table 1, Consistent ants). For example, consistent nurses were significantly more likely to only have interacted with other nurses (90% of nurse interactions, P=0.0326). Although cleaners and foragers who stayed within their functional group also more commonly interacted with other cleaners or

409 foragers, this difference in interaction proportions was not significantly higher than

410 expected by chance.

	Original task	Final task	Proportion interactions with other ants by role			p-value (Original task vs final task)
			Nurse	Cleaner	Forager	
Consistent ants	Nurse	Nurse	**0.9**	0.09	0.01	**0.0326***
	Cleaner	Cleaner	0.23	0.4	0.37	0.2105
	Forager	Forager	0.2	0.4	0.4	0.6744
Switching ants	Nurse	Cleaner	0.22	**0.71**	0.07	**0.0489***
	Cleaner	Forager	0.04	0.41	**0.55**	**0.0310***
	Forager	Nurse	0.05	**0.65**	0.3	**0.0446***
	Nurse	Forager	0.27	0.35	0.38	0.3671
	Cleaner	Nurse	**0.71**	0.24	0.05	**0.0229***
	Forager	Cleaner	0.02	**0.64**	0.34	**0.0019****

Table 1: The task interaction matrix, showing the proportion of an ants' interactions with a specified functional group before switching from its original to final group. P-values were calculated using a chi-square test contrasting the observed interaction proportions with the null model results for each type of task transition; values significantly differing from random expectations are indicated with asterisks. Bold type indicates the task and proportion of interactions with ants of that task that were significantly higher than expected.

However, most ants that switched to a new task interacted with ants currently

occupying a different task prior to switching (Table 1, Switching ants). For example,

nurses who switched to cleaning had interacted more frequently with cleaners (71% of

nurse interactions) and this was significantly more likely to occur based on interaction

history than by random chance (P = 0.0489). The result that an ant would transition to a

group that it previously interacted with the most was significant for the following other

transitions: cleaner to nurse, cleaner to forager, and forager to cleaner. Interestingly,

foragers who switched to nursing were significantly more likely to have interacted more

with ants of a different functional group, the cleaners (65% of forager interactions).

DISCUSSION

We explored task allocation in ant colonies to determine whether we could explain

how ants switched tasks based on information flow among functional groups and the

interaction history of the individuals. Mersch et al. (2013) attributed the pattern of ants

switching tasks to in this dataset to age polyethism based on the spatial division of

workers mediating the structure of the interaction network. The original authors

determined that task switching was a noisy process with a lot of individual variation, but

that at least some of the task switching could be explained by age polyethism based on

the spatial division of workers mediating the structure of the interaction network. Through

our reanalysis, we focused specifically on this noisy process of task switching. Our

approach allowed us to determine that previous interaction history can explain much more

of the noise behind task switching in *Camponotus fellah* colonies.

Our results suggest that ants in different functional groups had varying levels of

importance for information flow between individuals and groups in a colony. Additionaly,

ants that switched tasks often occupied positions in the interaction network that had high

potential for supporting information flow. Network analyses, combined with the task

interaction matrix, allowed us to describe how the architecture of interactions influences

the distribution of and switching among tasks in an ant colony.

At the scale of ant-to-ant interactions, we found that ants classified into the three main

tasks (forager, cleaner, nurse) differed in how they interacted with each other. This

affected their role in information flow for the colony as described by network measures.

Foragers had the highest interaction strength – they interacted longer and more frequently

than cleaners and nurses. Cleaners, however, had higher betweenness and thus were

key connections between ants interacting in the colony.

Ants that switched tasks functioned as key connectors for information flow in the

colony, supporting colony cohesion. In general, betweenness was higher for ants that

switched than for ants that remained consistent in their task. This suggests that ants who

switched tasks throughout the course of the experiment, and particularly cleaners, played

an important role in connecting other ants through information flows. Their high

betweenness means that they occupied a key network position for receiving and

transmitting information before they switched. If learning is required when ants switch

tasks, this increased access to information may have allowed them to be able to learn

new behaviours more quickly, helping them transition to a new task. Bridge betweenness

indicates how ants connected different communities within the colony. Ants who switched

tasks had higher bridge betweenness than those who remained consistent in their task.

In particular, foragers and cleaners had higher bridge betweenness than nurses, showing

that they were key connectors among the different functional groups. Cleaners especially

were less likely to interact within their functional group (consistent with Mersch et al.'s

2013 results). The low group cohesion of cleaners may strengthen colony-wide cohesion.

The instability of the centrality metrics may be related to cleaners' and foragers' ability

to transition tasks. Cleaners and foragers who switched functional groups had

significantly higher variances of betweenness and bridge betweenness, showing that

these individual measures of social network connectivity changed more over time.

However, when all ants in a colony were grouped for calculating the node-level network

metrics, pooled variances were not higher than those for the null networks. So while these

metrics varied significantly among functional groups and when ants switched tasks,

overall information flow in a colony remained fairly stable and colony cohesion was

maintained.

At the group level, the operationalization of effective information as a measure of the

noisiness of network mechanisms is relatively new and under-explored. Our effectiveness

results for the six *Camponotus fellah* colonies show a correlation between variations in

effectiveness and the functional group composition of each colony. We found that a higher

proportion of foragers lead to noisier communication among ants. Paired with the results

on interaction strength, this means that foragers interacted more than ants in other tasks

and that their interactions were noisier than interaction patterns of ants performing

different tasks. Results on centrality measures and effectiveness can be linked with task

allocation through our task interaction matrix. The matrix shows how previous interactions

with ants in a given task lead to a higher probability of the ant switching to that task. These

results are consistent with previous work in another species: Gordon and Mehdiabadi

(1999) found that, in red harvester ants, ants switching from other tasks to midden work

were more likely to have interacted with midden workers, and that switching was more

likely to occur the more frequent those interactions. In our results, interactions with

foragers appeared to drive switches to foraging: both cleaners and nurses who switched

to foraging had a higher probability of having interacted with foragers. Switches from

foragers to other tasks, however, showed different dynamics. Both foragers who switched

to nursing and foragers who switched to cleaning had a higher probability of having

interacted with cleaners, although the experimental data only showed two cases of the

former switch. This is in line with betweenness results, showing that cleaners are central

in driving switching patterns. These patterns suggest that, while previous interaction

patterns do affect switching behaviour, they do so to varying degrees depending on the

role played by the interacting ants and on the overall information flow of the system.

In future research, it would be interesting to further explore task switching in systems

with a higher granularity of data collection across both behaviours and interactions. One

limitation to the Mersch et al. (2013) dataset and to the original and current analyses is

that the task each ant was assigned to is assessed based on the interaction patterns, not

the types of actions or tasks the ant completes in the colony. Assessing not just who an

ant interacts with, but what actions that ant is actually completing, would provide useful

additional insight into the timing of behavioral and social change. This kind of data would

allow researchers to determine whether an ant alters its behaviors first (for example,

decreasing cleaning behaviors and increasing nursing behaviors) which then results in a

change in the social interaction patterns or whether an ant first begins to change its social

interaction patterns (for example, interacting less with other cleaners and more with

nurses) and then altering its behavior from cleaning actions to nursing actions. Another

open question is how the content of information flows, paired with the architecture of

interaction netowrks, affects individual and group behaviour (e.g., under conditions of

emergency). Future targeted data collection, involving both social and behavioral

observations, paired with statistically robust network methods, could be used to further

explore the relationships between patterns of interactions, individual-level behaviour, and

group-level behaviour.

ACKNOWLEDGEMENTS

We sincerely thank the organizers of 2019 Complex Networks Winter Workshop

(CNWW) for bringing us together and Brennan Klein for helpful discussions. A.S.'s

contribution to this research was supported in part through training by NSF award DGE-

1632976. S.W.'s contribution was supported in part by a NSF-GRF.

Author contributions: All the authors conceived the idea and the methodology together;

AS and SW did the analysis, and all the authors wrote the manuscript.

Competing interests: The authors declare no competing interests.

Data availability: All scripts and data used in this project are available at

https://github.com/anshuman21111/ant-colony-networks. A reformatted version of the

Mersch et al. (2013) dataset can be found as Supplementary material 1.

REFERENCES

1. Abd-Rabbo, D., 2017. VertexSort: Network Hierarchical Structure and Randomization. R package version 0.1-1. https://CRAN.R-project.org/package=VertexSort

2. Anderson, C., Shea, D.W.M., 2001. Individual versus social complexity, with particular reference to ant colonies. Biol. Rev. 49. doi:10.1023/A:1022699322624

3. Barabási, A., 2016. Network science.

4. Blonder, B., Dornhaus, A., 2011. Time-ordered networks reveal limitations to information flow in ant colonies. PLoS One 6, 1–8. doi:10.1371/journal.pone.0020298

5. Boomsma, J.J. and Franks, N.R., 2006. Social insects: from selfish genes to self organisation and beyond. Trends in Ecology & Evolution, 21(6), pp.303-308.

6. Byrum, T., Swain, A., Klein, B., Fagan, W., 2020. einet: Effective Information and Causal Emergence. R package version 0.1.0. https://CRAN.R-project.org/package=einet

7. Cavagna, A., Cimarelli, A., Giardina, I., Parisi, G., Santagati, R., Stefanini, F., Tavarone, R., 2010. From empirical data to inter-individual interactions: Unveiling the rules of collective animal behavior. Math. Model. Methods Appl. Sci. 20, 1491–1510.

8. Clutton-Brock, T.H., Brotherton, P.N., Russell, A.F., O'riain, M.J., Gaynor, D., Kansky, R., Griffin, A., Manser, M., Sharpe, L., McIlrath, G.M. and Small, T., 2001. Cooperation, control, and concession in meerkat groups. Science, 291(5503), pp.478-481.

9. Couzin, I.D., 2009. Collective cognition in animal groups. Trends Cogn. Sci.

10. Couzin, I.D., Ioannou, C.C., Demirel, G., Gross, T., Torney, C.J., Hartnett, A.,

Conradt, L., Levin, S.A. and Leonard, N.E., 2011. Uninformed individuals

promote democratic consensus in animal groups. *Science*, 334(6062), pp.1578-

1580.

11. Croft, D.P., Madden, J.R., Franks, D.W., James, R., 2011. Hypothesis testing in

animal social networks. Trends Ecol. Evol.

12. Csardi G, Nepusz T (2006). "The igraph software package for complex network

research." *InterJournal*, Complex Systems, 1695. http://igraph.org.

13. Emmons, S., Kobourov, S., Gallant, M. and Börner, K., 2016. Analysis of network

clustering algorithms and cluster quality metrics at scale. PloS one, 11(7).

14. Farine, D.R., Whitehead, H., 2015. Constructing, conducting and interpreting

animal social network analysis. J. Anim. Ecol. 84, 1144–1163.

15. Fewell, J.H., 2003. Social insect networks. Science (80-.).

16. Flack, J. C. (2017). Coarse-graining as a downward causation

mechanism. *Philosophical Transactions of the Royal Society A: Mathematical,*

Physical and Engineering Sciences, *375*(2109), 20160338.

17. Franklin, E.L. and Franks, N.R., 2012. Individual and social learning in tandem-

running recruitment by ants. Animal behaviour, 84(2), pp.361-368.

18. Freeman, L.C., 1979. Centrality in Social Networks Conceptual Clarification.

Social Networks.

19. Goldsby, H. J., Dornhaus, A., Kerr, B., & Ofria, C. (2012). Task-switching costs

promote the evolution of division of labor and shifts in individuality. *Proceedings*

of the National Academy of Sciences, *109*(34), 13686-13691.

20. Gordon, D.M., Mehdiabadi, N.J., 1999. Encounter rate and task allocation in

harvester ants. Behav. Ecol. Sociobiol. 45, 370–377.

21. Gordon, D.M., 2016. From division of labor to the collective behavior of social

insects. Behavioral ecology and sociobiology, 70(7), pp.1101-1108.

22. Hoare, D.J., Couzin, I.D., Godin, J.G. and Krause, J., 2004. Context-dependent

group size choice in fish. Animal Behaviour, 67(1), pp.155-164.

23. Holldobler, B., Wilson, E.O., 1990. The Ants. Belknap Press of Harvard

University Press.

24. Hoel, E., Klein, B., Swain, A., Griebenow, R. and Levin, M., 2020. Evolution leads

to emergence: An analysis of protein interactomes across the tree of life. bioRxiv.

25. Hoel, E. P., Albantakis, L., & Tononi, G. (2013). Quantifying causal emergence

shows that macro can beat micro. *Proceedings of the National Academy of*

Sciences of the United States of America, *110*(49), 19790–19795.

26. Kafsi, M., Braunschweig, R., Mersch, D., Grossglauser, M., Keller, L., Thiran, P.,

2016. Uncovering latent behaviors in ant colonies. 16th SIAM Int. Conf. Data Min.

2016, SDM 2016 450–458. doi:10.1137/1.9781611974348.51

27. Klein, B., & Hoel, E. (2020). The emergence of informative higher scales in

complex networks. *Complexity*, *2020*.

28. Krause, J., Lusseau, D. and James, R., 2009. Animal social networks: an

introduction. Behavioral Ecology and Sociobiology, 63(7), pp.967-973.

29. Lukas, D., & Clutton-Brock, T. (2018). Social complexity and kinship in animal

societies. *Ecology letters*, *21*(8), 1129-1134.

30. Lusseau, D., Whitehead, H., Gero, S., 2009. Incorporating uncertainty into the

study of animal social networks. Anim. Behav. 75, 1809–1815.

31. Lusseau, D. and Newman, M.E., 2004. Identifying the role that animals play in

their social networks. Proceedings of the Royal Society of London. Series B:

Biological Sciences, 271(suppl_6), pp.S477-S481.

32. McKinney Jr, E. H., & Yoos, C. J. (2010). Information about information: A

taxonomy of views. *MIS quarterly*, 329-344.

33. Mersch, D.P., Crespi, A., Keller, L., 2013. Tracking individuals shows spatial

fidelity is a key regulator of ant social organization. Science (80-.). 340, 1090–

1093. doi:10.1126/science.1234316

34. Molontay, R., Nagy, M., 2019. Two decades of network science as seen through

the co-authorship network of network scientists, in: Proceedings of the 2019

IEEE/ACM International Conference on Advances in Social Networks Analysis

and Mining, ASONAM 2019. Association for Computing Machinery, Inc, New

York, NY, USA, pp. 578–583.

35. Naug, D., 2009. Structure and resilience of the social network in an insect colony

as a function of colony size. Behav. Ecol. Sociobiol. 63, 1023–1028.

36. Pinter-Wollman, N., Wollman, R., Guetz, A., Holmes, S. and Gordon, D.M., 2011.

The effect of individual variation on the structure and function of interaction

networks in harvester ants. Journal of the Royal Society Interface, 8(64),

pp.1562-1573.

37. Pinter-Wollman, N., Hobson, E.A., Smith, J.E., Edelman, A.J., Shizuka, D., De

Silva, S., Waters, J.S., Prager, S.D., Sasaki, T., Wittemyer, G., Fewell, J.,

McDonald, D.B., 2014. The dynamics of animal social networks: Analytical,

conceptual, and theoretical advances. Behav. Ecol.

38. Radeva, T., Dornhaus, A., Lynch, N., Nagpal, R., Su, H.H., 2017. Costs of task

allocation with local feedback: Effects of colony size and extra workers in social

insects and other multi-agent systems. PLoS Comput. Biol. 13.

39. Ravary, F., Lecoutey, E., Kaminski, G., Châline, N., Jaisson, P., 2007. Individual

Experience Alone Can Generate Lasting Division of Labor in Ants. Curr. Biol. 17,

1308–1312.

40. Richardson, T.O., Mullon, C., Marshall, J.A., Franks, N.R. and Schlegel, T., 2018.

The influence of the few: a stable 'oligarchy'controls information flow in house-

hunting ants. Proceedings of the Royal Society B: Biological Sciences,

285(1872), p.20172726.

41. Robinson, E., Richardson, T., Feinerman, O., 2009. Radio tagging reveals the

roles of corpulence, experience and social information in ant decision making

The Bayesian Superorganism View project PhD ant learning and communication

View project. Springer 63, 627–636.

42. Robinson, E.J.H., Richardson, T.O., Sendova-Franks, A.B., Feinerman, O.,

Franks, N.R., 2009. Radio tagging reveals the roles of corpulence, experience

and social information in ant decision making. Behav. Ecol. Sociobiol. 63, 627–

636.

43. Sasaki, T., & Pratt, S. C. (2018). The psychology of superorganisms: Collective

decision making by insect societies. *Annual Review of Entomology*, *63*, 259-275.

44. Smith, Jennifer E., and Noa Pinter-Wollman (2021). "Observing the unwatchable:

Integrating automated sensing, naturalistic observations and animal social

network analysis in the age of big data." *Journal of Animal Ecology* 90.1: 62-75.

45. Sueur, C., Deneubourg, J.L., Petit, O. and Couzin, I.D., 2011. Group size,

grooming and fission in primates: a modeling approach based on group structure.

Journal of Theoretical Biology, 273(1), pp.156-166.

46. Sumpter, D.J.T., 2006. The principles of collective animal behaviour. Philos.

Trans. R. Soc. B Biol. Sci.

47. Sumpter, D.J., 2010. Collective animal behavior. Princeton University Press.

48. Swain, A. and Fagan, W.F., 2019. Group size and decision making: experimental

evidence for minority games in fish behaviour. Animal Behaviour, 155, pp.9-19.

49. Tononi, G. and Sporns, O., 2003. Measuring information integration. BMC

neuroscience, 4(1), p.31.

50. Tripet, F., Nonacs, P., 2004. Foraging for work and age-based polyethism: The

roles of age and previous experience on task choice in ants. Ethology 110, 863–

877.