Title: Interactions and information: Exploring task allocation in ant colonies using

network analysis

List of Authors: Anshuman Swain*[1#], Sara D. Williams*[2], Louisa J. Di Felice*[3] and

Elizabeth A. Hobson[4]

Institutional affiliations: [1]Department of Biology, University of Maryland, College Park,

MD, USA; [2]Mote Marine Laboratory, Sarasota, FL, USA; [3]IT University of Copenhagen,

Copenhagen, DK; [4]Department of Biological Sciences, University of Cincinnati, OH,

USA; *contributed equally

Contact Information: [#]answain@terpmail.umd.edu

ABSTRACT

11
In animal societies, individuals may take on different roles to fulfil their own needs

and the needs of their groups. Ant colonies display high levels of organisational

complexity, with ants fulfilling different roles at different timescales (what is known as *task*

allocation). Factors affecting task allocation can be at the individual level (e.g.,

physiology), or at the group level (e.g., the network of interactions). We focus on group

level processes by exploring the relationship between interaction networks, task

allocation and task switching using a previously published dataset (Mersch et al., 2013)

tracking the behaviour of six *Camponotus fellah* colonies over 41 days. In our new

analyses, our goal was to better explain the noisy process of task switching beyond simple

age polyethism. First, we investigated the architecture of interaction networks using node

(individual) level network measures and their relation to the individual's task – foraging,

cleaning or nursing – and whether or not the ant switched tasks. We then explored how

noisy information propagation was among ants, as a function of the colony composition

(how many ants carried out which tasks), through the information-theoretic metric of

Effective Information. Our results show that interaction history was tied to task allocation:

ants who switched to a task are more likely to have interacted with other ants carrying out

that task. The degree to which interactions related to task allocation, as well as the noise

in those interactions, depended on which groups of ants were interacting. Overall, we

showed that colony cohesion was stable even as ant-level network measures varied more

for ants when they switched functional groups; thus, ant colonies maintained a high level

of information flow as determined by network analysis, and ant functional groups played

different roles in maintaining colony cohesion through varied information flows.

Keywords: Ant behaviour, Effective Information, emergent behaviour, insect social

networks, task allocation, information flow

Highlights

● We analysed the interaction networks of six *Camponotus fellah* colonies

● We tested how centrality and information flow were tied to task switching

● Node-level network metrics and the information theoretic measure of *Effective*

Information explained differences among functional groups

● Interactions were correlated with task switching, but the strength of the

correlation differed across functional groups

INTRODUCTION

In animal societies, individuals may carry out different tasks to fulfil their own needs

and the needs of their group (Sumpter, 2006; Clutton-Brock, 2009; Jeanson &

Weidenmuller, 2014). Larger and more complex societies can self-organise to fulfil tasks

beyond basic sustenance and reproduction (Boomsma and Frank, 2006; Sumpter, 2010).

Local exchange of information, between individuals of a group and between individuals

and their environment, is key to self-organisation (Sumpter, 2006; Boomsma and Frank,

2006; Couzin, 2009; Cavagna et al., 2010; Swain and Fagan, 2019). Social insect

colonies display high levels of organisational complexity (Lukas & Clutton-Brock, 2018),

where individual tasks may include foraging, nest construction, and caring for the young

(Gordon, 2002). The assignment of tasks, also referred to as *task allocation*, is the result

of patterns of factors that vary across different scales (Gordon, 2015). These tasks can

be fixed throughout each individual's lifetime due to physiological reasons, for example

when only a fertile subset of the population is responsible for reproduction, or when a

subset is responsible for providing food (Sumpter, 2010; Clutton-Brock et al., 2001).

Task allocation can also result in individuals changing their main task over time.

Task allocation in ants has been the subject of much previous work (Anderson and Shea,

2001; Gordon, 2015). Across ant species, studies have shown that, depending on the

tasks and on the colony, ants may display varying degrees of task flexibility, from small

colonies of totipotent ants to larger ones with a structured division of labour (Anderson

and Shea, 2001). Factors affecting task changes can occur at the individual level or at

the group level. Individual-level factors include physiology (Anderson and Shea, 2001),

age (Tripet and Nonacs, 2004), corpulence (Robinson et al., 2009) and past experience

(Ravary et al., 2007), whereas group-level factors involve colony size (Ravary et al., 2007)

and interaction rates at the colony level (Gordon and Mehdiabadi, 1999). Studying

individual-level factors associated with task change is often simpler than studying group-

level ones. For example, individual-level changes can be easier to track because their

rate of change often follows a consistent and predictable pattern, as in the case of ageing.

Individual-level factors can also be directly quantified, e.g., by measuring age,

corpulence, or physiological features, and traditional statistical approaches can be used

to predict task changes.

An ant's propensity to switch to a new task could also be linked to nature of its

interactions and the topology of the group's social interaction network structure. However,

changes in task allocation affected by group-level factors are currently not well

understood. Structural features or macro-level social properties of groups can affect

micro-level individual actions if the social system is affected by feedbacks (Flack, 2017;

Hobson et al., 2019). However, quantifying relevant macro-to-micro feedbacks can be

challenging and can require large amounts of data. The development of automated

tracking systems has made this level of data collection possible. While these systems

have improved researchers' ability to track detailed social behaviour (Robinson et al.,

2009; Smith and Pinter-Wollman, 2021), assigning quantitative metrics to group dynamics

is still a non-trivial task. In the case of interaction patterns, tracking physical interactions

among individuals does not necessarily map onto the amount of meaningful (predictive)

information exchanged with each interaction (Valentini et al., 2020). Although tracking

technologies can tell us how many times individuals in a social group interact with one

another, they cannot explain to what extent these interactions are tied to task allocation

without considering the structure of these interactions and without including behavioural

observations. Network methods and metrics allow us to explore the interaction structure.

In this paper, we leverage social network methods to gain new insight into task

allocation changes in an existing dataset of ant interactions (published by Mersch et al.

2013). Mersch et al. studied task switching in *Camponotus fellah* by tracking and

analysing the movements and interactions of individually-identified ants. Worker ants

were categorised into three *functional groups* (nurse, cleaner or forager). Analyses

showed that ants had more interactions with others in their same functional group.

Communities defining the functional groups exhibited distinct behavioural signatures and

were highly spatially divided. Nurses spent most of their time with the brood, while

foragers spent time at the nest entrance and cleaners were located between the other

two groups and the rubbish pile (Mersch et al. 2013). Mersch et al. also explored the

questions of task switching cost, i.e., a time and energy investment associated with

learning new tasks (Goldsby et al., 2012), and of age polytheism, i.e., the correlation

between the age of an ant and which task they perform. The original study identified

spatial fidelity as a key regulator of ant social organisation and interaction frequency

(Mersch et al. 2013). They also found that task switches were present but uncommon and

that when a shift in functional group occurred, ants showed a preferred direction of task

transition, from nurses to cleaners to foragers, mostly based on age (Mersch et al., 2013).

Task changes were thus hypothesised to be driven by age polyethism, but the patterns

were fairly noisy.

In this new analysis we focus specifically on this noisy process of task switching

and its predictability. A question not addressed in the previous study is whether the history

of an ant's interactions with others could be one of the elements explaining task switching. In other species, information flow patterns have been shown to affect task allocation and overall colony behaviour, such as in the case of midden workers in red harvester ants (*Pogonomyrmex barbatus*; Gordon and Mehdiabadi, 1999, Pinter-Wollman et al., 2018), tandem running recruitment (Franklin and Franks, 2012) and consensus-forming in rock ants (*Temnothorax albipennis*; Sasaki & Pratt, 2018). To test whether the history of interactions or information flow could explain the noise seen in task switching dynamics that was not explained by age polyethism alone, we evaluated several potential macro-scale predictors of task switching not addressed in the original paper.

First, we described the architecture of the interaction networks by focusing on information flow (which in our case refers to the possible information exchange due to interactions among ants). We tested whether the role played by individual ants in regulating information flow in the colony and the functional group that they belong to were correlated. To do this, we quantified three network measures that are tied to the architecture of information flows at the local level for ant-to-ant interactions: strength mode, betweenness centrality, and bridge betweenness centrality. We also quantified a network level measure, *Effective Information* (EI), for the whole colony. At the scale of ant-to-ant interactions, strength measures the quantity and frequency of interactions of an ant, and strength mode finds the value in the distribution of strengths most commonly observed across all the ants in the group. Betweenness centrality measures the number of shortest paths between pairs of ants that pass through it. Bridge betweenness centrality extends betweenness to measure the number of shortest paths that pass through a node and connect separate highly connected groups of nodes, or communities. While strength,

betweenness centrality, and bridge betweenness centrality are common node-level

measures in network science and have been applied to animal social networks in the past

(Holme et al., 2002; Lusseau and Newman, 2004; Krause et al., 2009; Farine and

Whitehead, 2015), Effective Information is a new information theoretic metric reflecting

how noisy a mechanism connecting nodes (ants, in our case) is within a system. It is

calculated as the difference between degeneracy and determinism of the network (Hoel,

Albantakis, & Tononi, 2013; Klein and Hoel 2020). In interaction networks, Effective

Information reflects the noisiness and predictability of the interactions among individuals

(Hoel et al., 2020): a higher Effective Information means that a system is more

deterministic, with information spreading in a more effective way throughout the network.

Second, we tested whether these four measures of information flow in the

interaction network were related to task switching, to better understand the noise in task

allocation not explained by age polyethism as determined by Mersch et al. (2013). We

hypothesised that an ant's previous interactions with other ants affect switching behaviour

and tested whether interacting with a certain functional group increased the probability of

an ant switching to that group. We found that the relationship between the structure of the

interaction network and the different functional groups, as described by network

measures at the node and the global level, could explain the varying correlations between

interaction history and switching behaviour during task allocation. Our use of network

metrics, including the Effective Information metric, allowed us to determine the

relationship between interaction history, task allocation and information flow among

functional groups in *Camponotus fellah* colonies.

METHODS

Data, network construction, and ant categorisation

The published Mersch et al. (2013) dataset contains summaries of interactions among a total of 985 individually-marked ants in six *Camponotus fellah* colonies. The authors collected interaction data for every pair of ants at a daily resolution over the 41-day monitoring period, and the published dataset contains data pooled at the number of interactions per dyad per day per colony. We matched this published dataset with the colony metadata to inform our analyses (Supplementary material 1).

Consistent with Mersch et al. (2013), we used the pairwise daily number of interactions to construct separate weighted, undirected, unipartite networks for each colony per day. Each ant in a colony was represented by an individual node. An edge between two nodes represents the interactions between those two ants on a given day. The edge weight is proportional to the number of pairwise interactions between them on that particular day. We used the available published dataset to recreate the 246 observed networks for the 6 colonies over 41 days used by Mersch et al. (2013) as well as the general pattern of task switching across the length of the experiments.

Mersch et al. (2013) assessed each ant's functional group every 10 days to categorise them as a nurse, cleaner, or forager, representing their main task in the colony. They assigned functional groups based on what community an ant spent at least 70% of their time in, using the 'infomap' community detection algorithm paired with behavioural observations. They split the ants into the functional groups foragers (F), cleaners (C), nurses (N), queen (Q), and NA for ants who were counted as missing at a time point (e.g., if they were dead or had lost their tags).

Mersch et al. (2013) reported that their ants mostly did not change their task affiliation

within the 10-day observation period between task assessment points. We used the same

10-day snapshot window in our analyses which resulted in three time points at which a

switch in task to a new category could be detected. Based on Mersch et al's (2013)

observational data, when an ant switches functional groups, it switches tasks to that of

the new group. For our analyses, we categorised each ant as "switched" or "consistent",

depending on whether they were categorised as part of a different functional group, or

remained within the same functional group after each task assessment point in the

original behavioural data. These labels were assigned for each 10-day observation

period, meaning that an ant could be labelled as "consistent" in one time period because

it did not change tasks from the previous period, and "switched" in the next if it then

changed tasks, and thus functional groups during that next period. We utilised these

labels and the functional groups set by Mersch et al. (2013), throughout our work.

Before performing new analyses, we first investigated whether we could replicate

Mersch et al.'s (2013) results of age polyethism. We also tested whether we could

recapitulate Mersch et al.'s results about task switching by determining the likelihood that

an ant would stay in the same task throughout the experimental time versus performing

two or three tasks.

Quantifying individual network metrics for each ant

Our new analyses focused first at the individual scale within the networks. Node

metrics and centralities define various types of influence that individual nodes exert on

network connectivity and dynamics. For each network, we used R (v 3.6.2) and the

packages *igraph* (Csardi and Nepusz, 2006) and *networktools* (Jones, 2020) to calculate

three node-level, local metrics: (1) strength, (2) betweenness centrality, and (3) bridge

betweenness. Since networks were constructed for each daily set of interaction

observations for each colony, these metrics were calculated for each ant in every colony,

every day. Differences in these metrics were then analysed as a function of functional

groups at the colony level, and for just ants that switched or ants that remained consistent.

First, we calculated each ant's node strength as the sum of the weights of its edges.

Thus, in our context, it is a measure of not only how many interactions (edges) an ant

(node) had with other ants, but also of how frequently those interactions occurred during

a day. While degree is an index of potential communication activity (Freeman, 1979),

strength improves upon this index by weighting degrees according to communication

frequency, to better inform total interaction and information flow potential. To measure the

structure of the distribution of this node level metric at the network level, we calculated

the maxima of the density distribution of strength of all ants (or all within a functional group

subset) in a given colony on a given day to find the *strength mode*. The mode was used

instead of the mean because the strength distributions were skewed. The strength mode

provides a summary of how these strengths are generally distributed across each

network.

Second, we calculated each ant's node betweenness. Also known as betweenness

centrality, this measure is another way to assess the influence of a node for the

connectivity of the network. For a given pair of nodes in a weighted network, there exists

at least one path between them such that the sum of the link weights is minimized, thus

forming a shortest path. The betweenness of a node is therefore defined as the number

of shortest paths that pass through it. Freeman (1979) identified high betweenness

centrality as a key indicator of whether a node occupies a central location in the network

for information transmission. Individuals with high betweenness are often responsible for

maintenance of communication, group coordination, and network stability (Lusseau and

Newman, 2004; Farine and Whitehead, 2015). An ant with a high betweenness is an ant

that is centrally located in the network, serving as a key connection for seemingly

disparate ants.

Third, we measured the bridge betweenness for each ant in the network. Bridge

betweenness extends the betweenness centrality metric to the level of communities and

is defined as the number of times a node lies on the shortest path between two nodes

from different communities. In network science, a community is defined as a group of

nodes that have a higher likelihood of connecting to each other than to nodes from other

communities. Ants with a high bridge betweenness serve as key connectors for different

communities in the network, where communities mostly overlap with functional groups.

This means that ants with high bridge betweenness would be more integral to network

cohesion and information flow across groups, thus they may play an important role in

driving switching dynamics.

To quantify each ant's bridge betweenness, we needed to assign ants to network

communities in both the observed networks but also in our 123,000 reference networks

(see below). Assigning ants to network communities using the original network community

detection algorithm Infomap (used in Mersch et al. 2013) was computationally prohibitive

when applied to our many reference networks. Because of the computational demands

of the bridge betweenness analysis, we used a Louvain community detection algorithm

(Csardi and Nepusz, 2006) which saved computational time and memory (Emmons et al.,

2016) compared to the Infomap algorithm. These new network community assignments

were solely used for computing bridge betweenness and did not change the functional

group assignments of the ants made by Mersch et al. (2013), which we use in all other

cases in our analyses. To check that the Louvain algorithm assigned ants to network

communities in ways consistent with the original community assignments from Infomap,

we compared our community assignments to those found by Mersch et al. (2013): as we

show below, our new assignments were similar enough to the original assignments that

we could use our new method to assess bridge betweenness and the likelihood ants

would be connected to others within different functional groups (see results, below). All

other analyses involving functional group assignments of ants use the functional groups

assigned in Mersch et al. (2013).

Quantifying global network measures for each colony

267
We used _Effective Information_ and its normalised measure, _Effectiveness_, to measure

colony-level noisiness in the system, with respect to its underlying mechanisms (Hoel et

al., 2020; Klein et al., 2022). Since we are considering the mechanism of communication

and information flow among ants, Effective Information measures the level of

predictability, or degeneracy, in ant-to-ant interactions. To calculate Effectiveness, we first

found the sum of weights of all edges connected to each node in the interaction network,

where edge weights correspond to the number of interactions between a pair of ants. An

ant who had no interactions in a given network would have a weight of 0. We defined this

weight as a vector W_i of the same length as the total number of nodes and referred to

each element as ω_{ij}, signifying the normalised value of edge weight between nodes *i* and

j, such that for any index *i,* $\sum_j \omega_{ij} = 1$. Here, each term ω_{ij} can be seen as the probability

of moving from *i* to *j*, if a random walker is on the node *i*. Next, we characterised the

uncertainty associated with each node *i*, calculated using Shannon's entropy measure

$H(W_i)$. As node *i* has more connections, and as the weights $\omega_{ij}s$ of those connections to

other nodes (*j*) become more equal, Shannon's entropy (i.e., the uncertainty about where

a random walker would go) increases. The average of this value across all the nodes in

the network is $< H(W_i) >$. When $< H(W_i) >$ is equal to 0, the network is deterministic

(e.g., in the case of a line network or a ring lattice, both in directed and undirected cases,

where information can only flow in one dimension). We then assessed the certainty of the

network by calculating the term $H(< W_i >)$, which is Shannon's entropy of the average

out-weights from all nodes. When this expression is equal to 0, the network is degenerate,

with all edges leading to the same node. Finally, we calculated Effective Information using

the following equation:

$$EI = H(< W_i >) - < H(W_i) > \tag{1}$$

Here, the first term of the equation is determinism, and the second is degeneracy. Thus,

the Effective Information for a line graph or a ring lattice, which are maximally

deterministic and minimally degenerate, is the maximum. For the cases of a star

network, which is both maximally deterministic and degenerate, and that of a complete

graph, which is both minimally deterministic and degenerate, the value of Effective

Information is zero. As the value of Effective Information can depend on the size of the

network (Klein and Hoel, 2020), we calculated Effectiveness, the normalised Effective

Information with respect to network size. Effective Information is normalized by *log₂N*,

which is the maximal possible value of the entropy, where N is the number of nodes in

the network. For comparison, this is akin to the normalisation of Shannon diversity to

Shannon equitability in ecological studies.

303
$$Effectiveness = \frac{EI}{log_2 N} \tag{2}$$

Effectiveness was calculated for each observed network (i.e., for each day, for every

colony), using the R package *einet* (Byrum et al., 2020; Klein et al., 2022).

If node level properties were affected by which functional group an ant was in, then

system level measures could be affected by the proportion of ants in each functional

group, so we tested how group composition affected Effective Information. We used the

functional group assignment from Mersch et al. (2013), then fit linear models to the

Effectiveness for each observed network as a function of the proportion of each functional

group in the network to determine significant relationships between Effectiveness and a

colony's functional group composition.

Building reference models to test interaction patterns and task switching

To test how interaction patterns resulted in different network metrics and/or task

switching patterns than expected, we constructed randomised networks that served as

null models, or reference networks, for the daily interaction networks of the six colonies.

Using randomised networks as reference networks is a common method for testing the

effect of interaction structure and significance on various network properties and

dynamics (Hobson et al. 2021; Farine 2017). Constructing a reference model allowed us

to randomise some aspects of the interaction patterns while preserving other relevant

structural features of the networks (Hobson et al. 2021). We used a degree-based

randomisation (through the R package *VertexSort*; Abd-Rabbo, 2017) to generate our

reference networks. This approach preserved the total number of interaction partners per

any one ant on a given day but changed (1) who they interacted with and (2) how many

times they interacted. This process distributed the total number of original interactions

among the newly constructed edges of the randomised network.

This reference model approach allowed us to test how specific ant interaction

patterns affected the node and network level properties, while preserving the distribution

of connections in relation to functional groups of the ants (functional group assignment

and the degree of individual ants remained unchanged). As an example, a nurse that had

20 interaction partners (degree = 20) would still have a degree of 20 in the reference

network but would be interacting with 20 different ants with different frequencies, as the

edge weights were also randomly assigned from the initial distribution for each reference

network. This hypothetical reference model ant would then have a different total

frequency of interactions while maintaining their original number of partners. For

reproducibility, we created 500 seeded reference networks for each colony's daily

interaction network, for a total of 123,000 reference networks.

To test how observed network measures differed from those expected with the

identity of interaction partners and the number of interactions randomised, we compared

the observed node-level network measures to the distribution of those measures in our

reference networks. We found the strength mode, mean betweenness, and mean bridge

betweenness for every observed ant network (each colony, for each day) and for each of

the reference networks. We also estimated the variance for each metric for every ant in

a given colony on each day for the observed network and all the reference networks

associated with the observed one (please note that in the case of strength, the variance

is the standard variance in the strength distribution). The metrics were investigated

separately for each functional group within the following subsets: overall (all ants),

switching and consistent ants. Variance measured the individual variation of metrics

among ants of one group in a colony. The distribution of variances, in conjunction with

those of the central tendencies, helps us to explore the variation of these metrics across

colonies and through time. Central tendencies (mode for strength, and mean for

betweenness and bridge betweenness) and variance values were Z-transformed,

separately for each metric and individual observed network with its respective reference

networks, to facilitate comparison across observed networks which can vary in size

(number of ants), allowing us to find each metric's value for a given group, colony and

day relative to its own reference models. The Z-transformation allowed us to combine

values across colonies and days, and to visualize the means of those metrics across all

samples (groups, days and colonies). We then calculated the 95% confidence intervals

of the Z-transformed values for each functional group and ant subset (switching,

consistent and overall ants) to determine differences in the network measures in relation

to ant role and task switching. Since the distribution of variances indicates the variation

of a network metric both across colonies and through time, the size of the 95% confidence

intervals of the variances provides a proxy metric for stability of the network metrics, when

interpreted in conjunction with the distribution of the central tendencies of those metrics.

To test if the frequency of interactions with different functional groups significantly

affected an ant's functional group membership and whether these interactions could

explain how ants switched tasks, we compared the observed patterns of interactions in

relation to switching behaviour to that of the reference models. If the functional group

identity of ants affected how individuals contributed to information flow within a colony,

then the number of times an ant switched to a new functional group should affect the flow

of information. We tested how observed patterns differed from reference networks that

preserved the number of interactions per ant but redistributed the interactions among

every ant. We also tested whether the frequency of interactions with different functional

groups significantly affected an ant's final functional group. At each task assessment

point, we quantified the frequency of interactions with each functional group before

switching from its original functional group to the final one in both the observed dataset

and in the ensemble of randomised reference networks. We compared the distribution of

values computed from the observed networks against those given by the reference

network distribution using a suite of chi-square (independence/homogeneity)

comparisons separately for each possible type of task transition (including non-

transitions) and each observed network, wherever the specific transition/non-transition

occurred. To use the chi-square test, we assume that each interaction is independent as

it involves transfer of new information between two individuals, even if it might be biased

by more contact with certain individuals by choice. In addition, individual recognition is

not particularly well established for ants, although they can recognize brood mates and

their colony queen (see Esponda and Gordon, 2015; Sprenger and Menzel, 2020),

lending more credibility to our assumption.

The significance results from each type of transition across all observed networks

were then combined using Stouffer's method (see Heard and Rubin-Delanchy, 2017) and

significant differences at the alpha level of 0.05 were noted after accounting for multiple

comparisons (across transitions) through the Benjamini-Hochberg (BH) correction. We

combined all the cases where specific transitions were present, across all colonies and

days. Importantly, we were comparing the total number of interactions in these tests

pooled across all ants for a given subset, and not the number of ants that switched or

remained consistent in their tasks. In all cases, the frequency of interactions with each

type of functional group exceeded the minimum number required for using chi-square

tests, even though in certain cases the number of ants who were interacting were less

than five – their frequency of interactions exceed that number by a several orders of

magnitude. This process allowed us to assess whether interactions within the previous

observation period predicted functional group in the next observation period.

RESULTS

Replication of original results and visualisation of task switching

Figure 1 shows a new visualisation that summarizes the tasks of ants within all six

colonies and how those tasks changed over time (Figure 1; for details, see

Supplementary material 2, Table S1). To ground our analyses, we first replicated the main

results from the original Mersch et al. (2013) paper. We were able to replicate the original

results of age polyethism and recapitulated the distribution of the age of an ant that would

switch tasks once, twice, or three times (results from Mersch et al., novel visualisation in

Supplementary material 2, Figure S1A). We were also able to replicate Mersch et al.'s

results about task switching by determining the likelihood that an ant would stay in the

same task throughout the experimental time versus performing two or three tasks

(Supplementary material 2, Figure S1B).

Figure 1: Dynamics of task allocation across the experimental time for all ants in all six colonies. The alluvial diagram shows the number of ants per functional group and number of ants staying in the same group or transitioning to a new functional group between time periods as proportional to box and flow sizes, respectively (functional groups as originally determined in Mersch et al. 2013 via infomap).

We also compared our interaction community assignments used in calculating bridge

betweenness (via the Louvain community detection algorithm) with those obtained

through the Infomap algorithm originally used by Mersch et al. (2013; for the task of

functional group allocation). Community membership assignments, compared at an

individual node level for a given network, resulted in an average 90.13 ± 7.25% similarity

between the two methods across all the networks in the dataset. While the functional

group assignments used in our analyses were taken directly from Mersch et al.'s analysis

(which were validated by behavioural observations), this similarity of community

assignment is important as we wanted to certify that the structure of communities

detected by both algorithms was not divergent. The bridge betweenness metric used the

communities from the Louvain algorithm as it provided a substantial reduction in

computational time, and was indicative of the potential for an ant to connect ants from

different functional groups because of the high similarity with the community assignments

used to determine functional groups (i.e. by infomap and Mersch et al., 2013).

Individual network centrality measures and task switching

We compared network measures and their variances across each of the functional

groups for three categories: overall across all ants, for just switching ants, and for just

ants that remained consistent in their tasks during the assessment periods (summarised

in Figure 2; all values listed in Supplementary material 2 Table S2 and additional

visualisation in Figure S2), relative to their respective reference networks. Variance was

assessed due to apparent substantial fluctuations in the metrics for ants that switched

within the 10-day period leading to a task assessment point (Figure S3). These

fluctuations are represented by the 95% confidence intervals of the variances which

indicate how stable the relative network measures were across colonies and over time

(i.e., a larger confidence interval represents more fluctuations and less stability).

When we compared the distribution of the strength metric using the mode across

each of the tasks, we found that foragers had the highest strength mode of any of the

groups across all three of the categories, showing that they had the most frequent

interactions over a day regardless of whether they remained foragers or switched task at

some point. Values of variance (relative to their respective reference networks) of the

overall strength did not significantly differ across functional groups or from the reference

networks (see the 95% confidence intervals in Figure 2B) and the mode of strength

remained fairly consistent across functional groups (see the 95% confidence intervals in

Figure 2A). When we looked at strength just for switching ants, we found that the mode

differed significantly across functional groups and was significantly greater than for the

reference networks. Out of these, foragers that switched had the highest strength mode.

Strength mode variance of switching ants did not vary significantly among functional

groups. However, variance of the strength mode of switching foragers was higher and

more variable (i.e. larger confidence interval) than the reference networks, indicating less

stability of this metric among these individuals and over time. When we looked at strength

just for ants that were consistent, we found that the mode and variance followed the same

pattern seen for ants that switched, i.e., consistent foragers had higher strengths and the

values for all groups were of the same magnitude as those for the ants that switched.

At the colony level, the betweenness metric was stable (i.e., confidence intervals were

small for the both mean and the variance relative to the reference networks) and cleaners

played the most important role in connecting individual ants for flow of information, as

they had significantly higher betweenness than nurses and foragers (Figure 2). When we

assessed betweenness just for ants that switched, we found that mean betweenness

centrality measures were significantly greater than those for the reference networks,

except for foragers. Betweenness of switching ants was more variable than for consistent

ants. Consistent ants had the same relative patterns and magnitude of mean

betweenness centrality as ants that switched: consistent cleaners and nurses had higher

mean betweenness than consistent foragers. However, the variance of betweenness was

no longer significantly different than the reference networks, thus consistent ants

maintained a less variable betweenness distribution among networks and through time

(i.e., the mean and variance remain within small confidence intervals, and that the

variance is not significantly different from reference networks) than switching ants (where

even if the mean has similar values, the variance is much higher than reference

networks).

Figure 2: Z transformed values of the central tendencies (A) (note that the central tendency for strength is mode while for the other two metrics is mean) and variances (B) of the strength mode, betweenness and bridge betweenness determined for all ants across the six colonies. For each functional group, values were determined for all ants (overall), just ants that changed functional group during assessment periods (switching), and those ants that performed the same tasks from one time period to the next (consistent). Values were determined for every observed ant network (each colony, for each day) and for each of the reference networks before being Z-transformed to facilitate comparisons. Points are the values for each functional group (points depict average value and bars represent 95% confidence intervals among the measured central tendencies and variances (network-level statistics) of observed networks across days and colonies): nurses (red squares), cleaners (green triangles) and foragers (blue circles). The dashed black lines represent the upper extent of 95% confidence interval of the same normalised metrics from reference network simulations. (See Figure S2 for a more complete representation).

Since the communities we detected mapped primarily onto the previously determined

functional groups (see results above), a high bridge betweenness indicated a high

potential for connecting functional groups in a colony. When we compared bridge

betweenness across each of the functional groups at the colony level, we found that the

overall mean bridge betweenness values and their variance did not vary among the

functional groups or from the reference networks, indicating that connections among

network group communities stayed stable among networks and through time at the colony

level. Mean bridge betweenness was higher for the switching ants than for the consistent

ants for all functional groups, though only significantly higher for foragers. All ants that

switched had significantly higher mean bridge betweenness than the overall colony

values per functional group, suggesting that ants that switched played an important role

in connecting communities for information flow in the colony. The mean bridge

betweenness of consistent ants did not vary significantly among the functional groups or

from the reference network distribution. Although the variance of cleaners and foragers

for both ants that switched and those that were consistent was significantly higher and

more variable than the reference networks, the overall colony variance values remained

stable with small confidence intervals; these results may indicate that these interaction

structures could be important for colony cohesion at the community level.

495
496
Global information flow and task switching

We measured Effectiveness, the normalised Effective Information (difference between

how deterministic and degenerate a network is), as a function of the proportion of nurses,

cleaners or foragers in each colony for each day (resulting in 246 Effectiveness

measures, Figure 3). We found that the colony networks with high proportions of nurses

and cleaners had higher Effectiveness, but that the dependencies based on the linear

model were weak and non-significant (adj. R^2=0.12, P = 0.063, Figure 3A for nurses; and

Figure 3: Effectiveness (normalised Effective Information) of the interaction networks constructed for each colony and every day of the experiment as a function of the proportion of different functional groups in the networks. Data are stacked because the available granularity for task allocation was at a 10-day interval. Linear models fit to Effectiveness as a function of the proportions of nurses (A) and cleaners (B) separately return a nominally positive dependence (adj. R^2=0.12, p = 0.063 for nurses; and adj. R^2=0.11, p = 0.052 for cleaners). Effectiveness as a function of the proportion of foragers (C) returns a strong negative dependence (R^2 =0.22, p = 0.037).

adj. R^2=0.11, P = 0.052, Figure 3B for cleaners). Effectiveness significantly decreased

with increasing proportions of foragers in a colony (adj. R^2=0.22, P = 0.037, Figure 3C).

This negative relationship between the proportion of foragers and colony-level

Effectiveness suggests that interactions involving foragers were noisier than those

involving only nurses or cleaners.

Task interaction matrix and task switching

We tested whether previous interaction patterns affected switching behaviour using a

task interaction matrix. We found that ants that remained consistent in their tasks usually

interacted most with other ants occupying their same task (Table 1, Consistent ants). For

example, consistent nurses were significantly more likely to only have interacted with

other nurses (90% of nurse interactions, p=0.0326). Although cleaners and foragers who

stayed within their functional group also more commonly interacted with other cleaners or

foragers, this difference in interaction frequency was not significantly higher than

expected by chance. For simplicity and comparison throughout networks, we present the

average of proportion of interactions in each case in Table 1.

However, most ants that switched to a new task interacted with ants occupying a

different task prior to switching (Table 1, Switching ants). For example, nurses who

switched to cleaning had interacted more frequently with cleaners (71% of nurse

interactions) and this was significantly more likely to occur based on interaction history

than by random chance (p = 0.0489). The result that an ant would transition to a group

that it previously interacted with the most was significant for the following other transitions:

cleaner to nurse, cleaner to forager, and forager to cleaner. Interestingly, foragers who

switched to nursing were significantly more likely to have interacted more with ants of a

different functional group, the cleaners (65% of forager interactions who then switched to

nursing). However, it is important to note that this forager to nurse transition only occurred

in a few cases in the experimental data, so these results should be interpreted with

caution (due to the low number of observed ants), even though the interaction frequency

data was sufficient for the statistical comparison (see supplementary files for sample size

information).

	Original task	Final task	Proportion interactions with other ants by role			Corrected p-value (Original task to final task)
			Nurse	Cleaner	Forager	
Consistent ants	Nurse	Nurse	**0.9**	0.09	0.01	**0.0326***
	Cleaner	Cleaner	0.23	0.4	0.37	0.2105
	Forager	Forager	0.2	0.4	0.4	0.6744
Switching ants	Nurse	Cleaner	0.22	**0.71**	0.07	**0.0489***
	Cleaner	Forager	0.04	0.41	**0.55**	**0.0310***
	Forager	Nurse	0.05	**0.65**	0.3	**0.0446***
	Nurse	Forager	0.27	0.35	0.38	0.3671
	Cleaner	Nurse	**0.71**	0.24	0.05	**0.0229***
	Forager	Cleaner	0.02	**0.64**	0.34	**0.0019****

Table 1: The task interaction matrix, showing the proportion of an ant's interactions with a specified functional group before switching from its original to final group. P-values were calculated using a chi-square test contrasting the observed interaction proportions with the reference model results for each type of task transition; values significantly different from those obtained from their reference networks (after multiple comparison corrections) are indicated with asterisks. Bold type indicates the task and proportion of interactions with ants of that task that were dominant in each category (and which were higher than expected by random chance).

DISCUSSION

We explored task allocation in ant colonies to determine whether we could explain

how ants switched tasks based on information flow among functional groups and the

interaction history of the individuals. Mersch et al. (2013) determined that task switching

was a noisy process with a lot of individual variation, but that at least some of the task

switching could be explained by age polyethism based on the spatial division of workers

mediating the structure of the interaction network. In our analyses, we focused specifically

on this noisy process of task switching. Our approach allowed us to determine that

previous interaction history can help explain some of the noise behind task switching in

Camponotus fellah colonies and provides novel insight into task switching behaviour in

these ants.

Our results suggest that ants in different functional groups had varying levels of

importance for information flow between individuals and groups in a colony, based on

their individual roles in network connectivity as determined by the node-level metrics. Ants

that switched tasks often occupied positions in the interaction network that had high

potential for supporting information flow between groups. Network analyses, combined

with the task interaction matrix, allowed us to describe how the architecture of interactions

was related with the distribution of and switching among tasks in an ant colony.

At the scale of ant-to-ant interactions, we found that ants classified into the three main

tasks (forager, cleaner, nurse) differed in how they interacted with each other, which

affected their roles in information flow for the colony. Foragers had the highest interaction

strength mode – they interacted more frequently than cleaners and nurses. Cleaners,

however, had higher betweenness and thus were key connections between ants

interacting in the colony.

Ants that switched tasks functioned as key connectors for information flow in the

colony, supporting colony cohesion. In general, mean betweenness was higher for ants

that switched than for ants that remained consistent in their task, although confidence

intervals overlapped. Bridge betweenness (which indicated how ants connected different

communities within the colony) was significantly higher for ants that switched. This

suggests that ants who switched tasks throughout the course of the experiment, and

particularly foragers, played an important role in connecting functional groups through

information flows. Their high bridge betweenness means that they occupied a key network position for receiving and transmitting information before they switched tasks. If learning is required when ants switch tasks, this increased access to information may have allowed them to learn new behaviours more quickly, helping them transition to a new task. In general, cleaners were less likely to interact within their functional group (which was consistent with Mersch et al.'s 2013 results). The low group cohesion of cleaners may strengthen colony-wide cohesion.

The variability of the centrality metrics may be related to cleaners' and foragers' ability to transition tasks. Cleaners and foragers who switched functional groups had significantly higher variances of betweenness and bridge betweenness within networks. Across networks and over time, these variances also had a larger range, showing that these individual measures of social network connectivity changed more and were overall less consistent among individuals. However, when all ants in a colony were grouped for calculating the node-level network metrics, pooled variances were not higher than those for the reference networks and had small confidence intervals. So, while these metrics varied significantly among functional groups and when ants switched tasks, overall information flow in a colony remained fairly stable and colony cohesion was maintained.

At the group level, the operationalisation of Effective Information as a measure of the noisiness of network mechanisms is relatively new and under-explored. Our results for the six *Camponotus fellah* colonies show a correlation between variations in Effectiveness and the functional group composition of each colony. We found that a higher proportion of foragers led to noisier potential communication among ants. Paired with the results on interaction strength, this means that foragers interacted more frequently than ants

performing other tasks and that they had more diverse interactions with ants at different

positions in the interaction network. Results on centrality measures and Effectiveness

can be linked with task allocation through our task interaction matrix. The matrix shows

how previous interactions with ants in a given task are associated with a higher probability

of the ant switching to that task. These results are consistent with previous work in another

species: Gordon and Mehdiabadi (1999) found that, in red harvester ants, ants switching

from other tasks to midden work were more likely to have interacted with midden workers,

and that switching was more likely to occur the more frequent those interactions. In our

results, interactions with foragers were correlated with switches to foraging: both cleaners

and nurses who switched to foraging had a higher probability of interacting with foragers.

Switches from foragers to other tasks, however, showed different dynamics. Both

foragers who switched to nursing and foragers who switched to cleaning had a higher

probability of interacting with cleaners. These results should be interpreted with some

caution because ants switching from foragers to nursing was only observed three times

(the interaction frequency data, however, was sufficient for the chi-square comparison).

Consistent with betweenness results, these switching results show that cleaners were

central in driving switching patterns by occupying key positions for information flow in the

networks. These patterns suggest that, while previous interaction patterns were

correlated to switching behaviour, the degree of correlation varied depending on the role

played by the interacting ants and on the overall information flow of the system. It is

important to note that without more detailed data we cannot determine whether a change

in task or change in interactions happened first, but these insights provide valuable

information about system dynamics and suggestions for future experiments.

In future research, it would be interesting to further explore task switching in systems

with a higher granularity of data collection across both behaviours and interactions. One

limitation to the current analysis is that the task each ant was assigned to is assessed

based on the interaction patterns, not the types of actions or tasks the ant completes in

the colony. Even though interaction community membership was paired with behavioural

observations by Mersch et al. (2013), it may not have been at the level of detail needed

to assess fine-grain interaction patterns and task performance. It would be interesting to

use the combination of network methods and behavioural observations to further explore

existing results on the relationship between repetition (Langridge et al., 2004) and the

existence of experienced individuals (Langridge et al., 2007) on task performance.

Assessing not just who an ant interacts with, but what actions that ant is actually

completing, would provide useful additional insight into the timing of behavioural and

social change. This kind of data would allow researchers to determine whether an ant

alters its behaviours first (for example, decreasing cleaning behaviours and increasing

nursing behaviours) which then results in a change in the social interaction patterns, or

whether an ant first begins to change its social interaction patterns (for example,

interacting less with other cleaners and more with nurses) and then alters its behaviour

from cleaning actions to nursing actions. Future targeted data collection, involving both

social and behavioural observations, paired with statistically robust network methods,

could be used to further explore the relationships between patterns of interactions,

individual-level behaviour, and group-level behaviour.

ACKNOWLEDGEMENTS

We sincerely thank the organizers of 2019 Complex Networks Winter Workshop

(CNWW) for bringing us together and Brennan Klein for helpful discussions. A.S.'s

contribution to this research was supported in part through training by NSF award DGE-

1632976. S.W.'s contribution was supported in part by an NSF-GRF. EAH was

supported by NSF IOS 2015932 during preparation of this work.

Author contributions: All the authors conceived the idea and the methodology together;

AS and SW did the analysis, and all the authors wrote the manuscript.

Competing interests: The authors declare no competing interests.

Data availability: All scripts and data used in this project are available at

https://github.com/anshuman21111/ant-colony-networks. A reformatted version of the

Mersch et al. (2013) dataset can be found as Supplementary material 1.

REFERENCES

1. Abd-Rabbo, D., 2017. VertexSort: Network Hierarchical Structure and Randomization. R package version 0.1-1. https://CRAN.R-project.org/package=VertexSort

2. Anderson, C., Shea, D.W.M., 2001. Individual versus social complexity, with particular reference to ant colonies. Biol. Rev. 49. doi:10.1023/A:1022699322624

3. Blonder, B., Dornhaus, A., 2011. Time-ordered networks reveal limitations to information flow in ant colonies. PLoS One 6, 1–8. doi:10.1371/journal.pone.0020298

4. Boomsma, J.J. and Franks, N.R., 2006. Social insects: from selfish genes to self organisation and beyond. Trends in Ecology & Evolution, 21(6), pp.303-308.

5. Byrum, T., Swain, A., Klein, B., Fagan, W., 2020. einet: Effective Information and Causal Emergence. R package version 0.1.0. https://CRAN.R-project.org/package=einet

6. Cavagna, A., Cimarelli, A., Giardina, I., Parisi, G., Santagati, R., Stefanini, F., Tavarone, R., 2010. From empirical data to inter-individual interactions: Unveiling the rules of collective animal behavior. Math. Model. Methods Appl. Sci. 20, 1491–1510.

7. Clutton-Brock, T.H., Brotherton, P.N., Russell, A.F., O'riain, M.J., Gaynor, D., Kansky, R., Griffin, A., Manser, M., Sharpe, L., McIlrath, G.M. and Small, T., 2001. Cooperation, control, and concession in meerkat groups. Science, 291(5503), pp.478-481.

8. Couzin, I.D., 2009. Collective cognition in animal groups. Trends Cogn. Sci.

9. Couzin, I.D., Ioannou, C.C., Demirel, G., Gross, T., Torney, C.J., Hartnett, A.,

Conradt, L., Levin, S.A. and Leonard, N.E., 2011. Uninformed individuals

promote democratic consensus in animal groups. *Science*, 334(6062), pp.1578-

1580.

10. Croft, D.P., Madden, J.R., Franks, D.W., James, R., 2011. Hypothesis testing in

animal social networks. Trends Ecol. Evol.

11. Csardi G, Nepusz T (2006). "The igraph software package for complex network

research." *InterJournal*, Complex Systems, 1695. http://igraph.org.

12. Emmons, S., Kobourov, S., Gallant, M. and Börner, K., 2016. Analysis of network

clustering algorithms and cluster quality metrics at scale. PloS one, 11(7).

13. Esponda, F. and Gordon, D.M., 2015. Distributed nestmate recognition in ants.

Proceedings of the Royal Society B: Biological Sciences, 282(1806),

p.20142838.

14. Farine, D.R., Whitehead, H., 2015. Constructing, conducting and interpreting

animal social network analysis. J. Anim. Ecol. 84, 1144–1163.

15. Farine, D.R., 2017. A guide to null models for animal social network analysis.

Methods in Ecology and Evolution, 8(10), pp.1309-1320.

16. Fewell, J.H., 2003. Social insect networks. Science (80-.).

17. Flack, J. C. (2017). Coarse-graining as a downward causation

mechanism. *Philosophical Transactions of the Royal Society A: Mathematical,*

Physical and Engineering Sciences, *375*(2109), 20160338.

18. Franklin, E.L. and Franks, N.R., 2012. Individual and social learning in tandem-

running recruitment by ants. Animal behaviour, 84(2), pp.361-368.

19. Freeman, L.C., 1979. Centrality in Social Networks Conceptual Clarification.

Social Networks.

20. Goldsby, H. J., Dornhaus, A., Kerr, B., & Ofria, C. (2012). Task-switching costs

promote the evolution of division of labor and shifts in individuality. *Proceedings*

of the National Academy of Sciences, *109*(34), 13686-13691.

21. Gordon, D.M., Mehdiabadi, N.J., 1999. Encounter rate and task allocation in

harvester ants. Behav. Ecol. Sociobiol. 45, 370–377.

22. Gordon, D.M., 2016. From division of labor to the collective behavior of social

insects. Behavioral ecology and sociobiology, 70(7), pp.1101-1108.

23. Heard, N. A., & Rubin-Delanchy, P. (2018). Choosing between methods of

combining-values. Biometrika, 105(1), 239-246.

24. Hoare, D.J., Couzin, I.D., Godin, J.G. and Krause, J., 2004. Context-dependent

group size choice in fish. Animal Behaviour, 67(1), pp.155-164.

25.   Hobson, E. A., Ferdinand, V., Kolchinsky, A., & Garland, J. (2019).

Rethinking animal social complexity measures with the help of complex systems

concepts. Animal Behaviour, 155, 287-296.

26. Hobson, E.A., Silk, M.J., Fefferman, N.H., Larremore, D.B., Rombach, P., Shai,

S. and Pinter-Wollman, N., 2021   . A guide to choosing and implementing

reference models for social network analysis. Biological Reviews, 000-000.

27. Holldobler, B., Wilson, E.O., 1990. The Ants. Belknap Press of Harvard

University Press.

28. Hoel, E., Klein, B., Swain, A., Griebenow, R. and Levin, M., 2020. Evolution leads

to emergence: An analysis of protein interactomes across the tree of life. bioRxiv.

29. Hoel, E. P., Albantakis, L., & Tononi, G. (2013). Quantifying causal emergence

shows that macro can beat micro. *Proceedings of the National Academy of*

Sciences of the United States of America, *110*(49), 19790–19795.

30. Kafsi, M., Braunschweig, R., Mersch, D., Grossglauser, M., Keller, L., Thiran, P.,

2016. Uncovering latent behaviors in ant colonies. 16th SIAM Int. Conf. Data Min.

2016, SDM 2016 450–458. doi:10.1137/1.9781611974348.51

31. Klein, B., & Hoel, E. (2020). The emergence of informative higher scales in

complex networks. *Complexity*, *2020*.

32. Klein, B., Swain, A., Byrum, T., Scarpino, S. V., & Fagan, W. F. (2022). Exploring

noise, degeneracy and determinism in biological networks with the einet

package. Methods in Ecology and Evolution, 00, 1–6.

https://doi.org/10.1111/2041-210X.13805

33. Krause, J., Lusseau, D. and James, R., 2009. Animal social networks: an

introduction. Behavioral Ecology and Sociobiology, 63(7), pp.967-973.

34. Langridge, E. A., Franks, N. R., & Sendova-Franks, A. B. (2004). Improvement in

collective performance with experience in ants. Behavioral Ecology and

Sociobiology, 56(6), 523-529.

35. Langridge, E. A., Sendova-Franks, A. B., & Franks, N. R. (2008). How

experienced individuals contribute to an improvement in collective performance in

ants. Behavioral Ecology and Sociobiology, 62(3), 447-456.

36. Lukas, D., & Clutton-Brock, T. (2018). Social complexity and kinship in animal

societies. *Ecology letters*, *21*(8), 1129-1134.

37. Lusseau, D., Whitehead, H., Gero, S., 2009. Incorporating uncertainty into the

study of animal social networks. Anim. Behav. 75, 1809–1815.

38. Lusseau, D. and Newman, M.E., 2004. Identifying the role that animals play in

their social networks. Proceedings of the Royal Society of London. Series B:

Biological Sciences, 271(suppl_6), pp.S477-S481.

39. McKinney Jr, E. H., & Yoos, C. J. (2010). Information about information: A

taxonomy of views. *MIS quarterly*, 329-344.

40. Mersch, D.P., Crespi, A., Keller, L., 2013. Tracking individuals shows spatial

fidelity is a key regulator of ant social organization. Science (80-.). 340, 1090–

1093. doi:10.1126/science.1234316

41. Molontay, R., Nagy, M., 2019. Two decades of network science as seen through

the co-authorship network of network scientists, in: Proceedings of the 2019

IEEE/ACM International Conference on Advances in Social Networks Analysis

and Mining, ASONAM 2019. Association for Computing Machinery, Inc, New

York, NY, USA, pp. 578–583.

42. Naug, D., 2009. Structure and resilience of the social network in an insect colony

as a function of colony size. Behav. Ecol. Sociobiol. 63, 1023–1028.

43. Pinter-Wollman, N., Wollman, R., Guetz, A., Holmes, S. and Gordon, D.M., 2011.

The effect of individual variation on the structure and function of interaction

networks in harvester ants. Journal of the Royal Society Interface, 8(64),

pp.1562-1573.

44. Pinter-Wollman, N., Hobson, E.A., Smith, J.E., Edelman, A.J., Shizuka, D., De

Silva, S., Waters, J.S., Prager, S.D., Sasaki, T., Wittemyer, G., Fewell, J.,

McDonald, D.B., 2014. The dynamics of animal social networks: Analytical, conceptual, and theoretical advances. Behav. Ecol.

45. Radeva, T., Dornhaus, A., Lynch, N., Nagpal, R., Su, H.H., 2017. Costs of task allocation with local feedback: Effects of colony size and extra workers in social insects and other multi-agent systems. PLoS Comput. Biol. 13.

46. Ravary, F., Lecoutey, E., Kaminski, G., Châline, N., Jaisson, P., 2007. Individual Experience Alone Can Generate Lasting Division of Labor in Ants. Curr. Biol. 17, 1308–1312.

47. Richardson, T.O., Mullon, C., Marshall, J.A., Franks, N.R. and Schlegel, T., 2018. The influence of the few: a stable 'oligarchy' controls information flow in house-hunting ants. Proceedings of the Royal Society B: Biological Sciences, 285(1872), p.20172726.

48. Robinson, E.J.H., Richardson, T.O., Sendova-Franks, A.B., Feinerman, O., Franks, N.R., 2009. Radio tagging reveals the roles of corpulence, experience and social information in ant decision making. Behav. Ecol. Sociobiol. 63, 627–636.

49. Sasaki, T., & Pratt, S. C. (2018). The psychology of superorganisms: Collective decision making by insect societies. *Annual Review of Entomology*, *63*, 259-275.

50. Smith, Jennifer E., and Noa Pinter-Wollman (2021). "Observing the unwatchable: Integrating automated sensing, naturalistic observations and animal social network analysis in the age of big data." *Journal of Animal Ecology* 90.1: 62-75.

51. Sprenger, P.P. and Menzel, F., 2020. Cuticular hydrocarbons in ants

(Hymenoptera: Formicidae) and other insects: how and why they differ among

individuals, colonies, and species. Myrmecological news, 30.

52. Sueur, C., Deneubourg, J.L., Petit, O. and Couzin, I.D., 2011. Group size,

grooming and fission in primates: a modeling approach based on group structure.

Journal of Theoretical Biology, 273(1), pp.156-166.

53. Sumpter, D.J.T., 2006. The principles of collective animal behaviour. Philos.

Trans. R. Soc. B Biol. Sci.

54. Sumpter, D.J., 2010. Collective animal behavior. Princeton University Press.

55. Swain, A. and Fagan, W.F., 2019. Group size and decision making: experimental

evidence for minority games in fish behaviour. Animal Behaviour, 155, pp.9-19.

56. Tononi, G. and Sporns, O., 2003. Measuring information integration. BMC

neuroscience, 4(1), p.31.

57. Tripet, F., Nonacs, P., 2004. Foraging for work and age-based polyethism: The

roles of age and previous experience on task choice in ants. Ethology 110, 863–

877.