Identifying and correcting for misspecifications in GWAS summary statistics and polygenic scores

Florian Privé,[1,*] Julyan Arbel,[2] Hugues Aschard,[3,4] and Bjarni J. Vilhjálmsson[1,5]

[1]National Centre for Register-Based Research, Aarhus University, Aarhus, 8210, Denmark.

[2]Univ. Grenoble Alpes, Inria, CNRS, Grenoble INP, LJK, Grenoble, 38000, France.

[3]Department of Computational Biology, Institut Pasteur, Paris, 75015, France.

[4]Program in Genetic Epidemiology and Statistical Genetics, Harvard T.H. Chan School of Public Health, Boston, MA, 02115, USA.

[5]Bioinformatics Research Centre, Aarhus University, Aarhus, 8000, Denmark.

[*]To whom correspondence should be addressed.

Contact: `florian.prive.21@gmail.com`

Abstract

Are genome-wide association studies (GWAS) summary statistics of good enough quality for performing follow-up analyses? Can we detect possible misspecifications in GWAS summary statistics and correct for them in order to improve predictive performance of polygenic scores?

1 Introduction

Contrary to individual-level genotypes and phenotypes, summary statistics resulting from genome-wide association studies (GWAS) are widely available, and very large sample sizes can be obtained through meta-analysis (Yengo *et al.*, 2022). GWAS summary statistics have been extensively used to derive polygenic scores (PGS), perform fine-mapping, and estimate a range of key genetic architecture parameters (Privé *et al.*, 2021). However, GWAS summary statistics come with uneven imputation accuracy. There is also heterogeneity in the information made available in these summary statistics; for example, per-variant imputation INFO scores, sample sizes and allele frequencies are often missing. Moreover, many methods based on summary statistics use Bayesian models, which can be sensitive to model misspecifications (Walker, 2013; Miller and Dunson, 2019).

We present an overview of possible misspecifications that come with GWAS summary statistics (Table 1). First, the total sample size used can be misestimated, which would result in a biased estimation of the SNP heritability. For example, the total effective sample size is overestimated when computed from the total number of cases and controls from a meta-analysis of binary outcomes (Grotzinger *et al.*, 2021); using BOLT-LMM summary statistics can result in an increased effective sample size (Loh *et al.*, 2018; Gazal *et al.*, 2018); using SAIGE on binary traits with a large prevalence can result in a reduced effective sample size (Zhou *et al.*, 2018; Mbatchou *et al.*, 2021). Second, per-variant sample sizes can vary substantially and be much smaller than the total sample size when meta-analyzing GWAS summary statistics from multiple cohorts with different sets of variants (Wang *et al.*, 2021). Using very different per-variant sample sizes can be problematic for models implicitly assuming that summary statistics have all been derived from the same individuals (Zhu and Stephens, 2017; Zhou and Zhao, 2021). As a quality control (QC) step, one can e.g. remove SNPs with a sample size less than 0.67 times the 90th percentile of all sample sizes (Zheng *et al.*, 2017). Third, many genetic variants are imputed, which can cause some bias in summary statistics. Fourth, the imputation quality of each variant (e.g. INFO scores), often used in a QC step in statistical genetics analyses, can be severely misestimated (often overestimated) when computed from multi-ancestry individuals instead of a more homogeneous subset. Fifth, errors such as allele inversions can be present in the summary statistics; QC is particularly important here. Sixth, there can be some mismatch between the GWAS summary statistics and the LD reference used.

Here we investigate some of these misspecifications, and propose adjustments to improve the predictive performance of polygenic scores derived from GWAS summary statistics. We approach this from three different angles. First, based on additional summary information such as the imputation INFO scores and allele frequencies from the GWAS summary statistics, we refine our previously proposed QC (Privé *et al.*, 2020b). This QC consists in comparing standard deviations (of genotypes) inferred from GWAS summary statistics with the ones computed from a reference panel. This is useful to check that the input parameters used are consistent with each other. This was particularly important for LDpred2-auto, which directly estimates key model parameters from the data (Privé *et al.*, 2020b). Here we further show that standard deviations of imputed genotypes (allele dosages) are lower than the expected values under Hardy–Weinberg equilibrium. Second, we investigate possible adjustments to apply to the input parameters of methods using summary statistics (focusing on PGS methods here), namely the reference LD (linkage disequilibrium) ma-

Misspecification	Possible consequence(s)	Possible correction
Misestimation of the total sample size[1]	Bias estimation of SNP heritability (Gazal *et al.*, 2018; Grotzinger *et al.*, 2021)	Estimation of the effective sample size
Differences in per-variant sample sizes	Possible violation of model assumptions resulting in poor model convergence	Imputation using equations 3 and 4, and filtering[2]
Using imputed dosages for GWAS	Standard deviations of dosages are too small, resulting in overestimated effect sizes and standard errors	See section 2.2
Using imputed dosages for LD computation		None required; LD seems correctly estimated (Figure S28)
Misestimation of imputation INFO scores	Misuse when filtering or adjusting input parameters based on these	Recomputing from a homogeneous subset
Error in summary statistics (e.g. allele inversions)	Strong violation of model assumptions, which can result in e.g. identifying false-positives in fine-mapping (Zou *et al.*, 2021)	Quality control
Rounding of summary statistics (e.g. effect sizes and SEs)	Add unnecessary noise to the data	None, but this has almost no impact on the predictive performance (Figure S11)
Ancestry mismatch between summary statistics and LD	Strong mismatch of LD and allele frequencies, often resulting in divergence of models	Checking ancestry proportions from GWAS summary statistics (Privé, 2021b)
Any	Possible violation of model assumptions resulting in poor model convergence	Quality control, using more regularization, and constraining LD to blocks

Table 1: Overview of possible misspecifications when using GWAS summary statistics, along with possible consequences and corrections.
[1] Examples: using the total number of cases and controls in a meta-analysis of binary traits, a larger effective sample size when using BOLT-LMM summary statistics (Loh *et al.*, 2018), and a reduced effective sample size when using SAIGE on binary traits with a large prevalence (Zhou *et al.*, 2018; Mbatchou *et al.*, 2021).
[2] E.g. removing SNPs with an effective sample size less than 0.67 times the 90th percentile of sample size (Zheng *et al.*, 2017).

trix, the GWAS effect sizes, their standard errors and corresponding sample sizes. For example, we show that GWAS effect sizes computed from imputed dosages are larger in magnitude compared to if computed from true genotypes. Third, we introduce two new optional parameters in LDpred2-auto to make it more robust to these types of misspecification. We focus our investigations on LDpred2 and lassosum for two reasons. First, multiple studies have shown that LDpred2 and lassosum rank among the best methods for single-trait polygenic prediction (Mak *et al.*, 2017; Privé *et al.*, 2020b; Pain *et al.*, 2021; Kulm *et al.*, 2021). Second, we reimplement and use a new version of lassosum, called lassosum2, that uses the exact same input parameters as LDpred2, which makes it easy for us to test the different QCs and adjustments presented here.

2 Results

2.1 Misspecification of per-variant GWAS sample sizes

We design GWAS simulations where variants have different sample sizes, which is often the case when meta-analyzing GWAS summary statistics from multiple cohorts with different sets of variants (Wang *et al.*, 2021). Using 40,000 variants from chromosome 22 (Methods), we simulate quantitative phenotypes with a heritability of 20% and 2000 causal variants. We then divide the 40,000 variants into three groups: for half of the variants, we use 100% of 300,000 individuals for GWAS, but use only 80% for one quarter of the variants and 60% for the remaining quarter. We then run C+T, lassosum, lassosum2 (Methods), LDpred2-inf, LDpred2(-grid), and LDpred2-auto (Privé *et al.*, 2019; Mak *et al.*, 2017; Privé *et al.*, 2020b) by using either the true per-variant GWAS sample sizes, the total sample size for all variants, or imputed (per-variant) sample sizes. Note that we initially included PRS-CS and SBayesR in our comparison (Ge *et al.*, 2019; Lloyd-Jones *et al.*, 2019). However, results for SBayesR always diverged (stopping with an error), and the overlap with the LD reference provided for PRS-CS was too small. When providing true per-variant GWAS sample sizes, squared correlations between the polygenic scores and the simulated phenotypes are of 0.123 for C+T, 0.161 for lassosum, 0.169 for lassosum2, 0.159 for LDpred2(-grid), 0.140 for LDpred2-auto, and 0.141 for LDpred2-inf (Figure 1, averaging over 10 simulations). Results when using imputed sample sizes are quite similar to when using the true ones. Note that C+T does not use this sample size information. When using the total GWAS sample size instead of the per-variant sample sizes, predictive performance slightly decreases to 0.157 for lassosum and to 0.163 for lassosum2, but dramatically decreases for LDpred2 with new values of 0.134 for LDpred2-grid, 0.119 for LDpred2-auto, and 0.123 for LDpred2-inf (Figure 1). This extreme simulation scenario shows that LDpred2 can be sensitive to the misspecification of per-variant GWAS sample sizes, whereas lassosum (and lassosum2) seems little affected by this.

We conduct further investigations to explain the results of figure 1. First, the results for LDpred2-auto are similar to LDpred2-inf because it always converges to an infinitesimal model ($p = 1$) in these simulations. To overcome this limitation, we introduce two new parameters to make LDpred2-auto more robust, and refer to this as "LDpred2-auto-rob" here (Methods). The first parameter prevents p from diverging to 1, while the second shrinks the off-diagonal elements of the LD matrix (a form of regularization). Second, for lassosum2, results for a grid of parameters (over λ and δ) are quite smooth compared to LDpred2 (Figures S2 and S3). In these simulations with misspecified per-variant sample sizes, it seems highly beneficial to use a small value for the SNP heritability hyper-parameter h^2 in LDpred2, e.g. a value of 0.02 or even 0.002 when the true value is 0.2 (Figure S3). Indeed, using a small value for this hyper-parameter induces a larger regularization (shrinkage) on the effect sizes. Here we call "LDpred2-low-h2" when running LDpred2(-grid) with a grid of hyper-parameters including these low values for h^2. Results with LDpred2-low-h2 improves from 0.159 to 0.169 when using true sample sizes and from 0.134 to 0.163 when using the maximum sample size. Finally, we introduce a last change for robustness here: we form independent LD blocks in the LD matrix to prevent small errors in the Gibbs sampler to propagate to too many variants (Methods). This change seems to solve convergence issues of LDpred2 in these simulations (Figure S3) and further improves predictive performance for all LDpred2 methods (Figure 1).

Figure 1: Results for the simulations with sample size misspecification, averaged over 10 simulations for each scenario. Reported 95% confidence intervals are computed from 10,000 non-parametric bootstrap replicates of the mean. The GWAS sample size is either "true" when providing the true per-variant sample size, "max" when providing instead the maximum sample size as a unique value to be used for all variants, "imputed" (Methods), or "any" when the method does not use this information (the case for C+T). Red bars correspond to using the LD with independent blocks (Methods).

2.2 When using allele dosages from imputation

Marchini and Howie (2010) showed that the IMPUTE INFO measure is highly concordant with the MACH measure $\hat{r}_j^2 = \frac{\text{var}(G_j)}{2\hat{\theta}_j(1-\hat{\theta}_j)}$, where $\hat{\theta}_j$ is the estimated allele frequency of G_j, the genotypes for variant j. Therefore, when using the expected genotypes from imputation (allele dosages), their standard deviations are often lower than the expected value under Hardy–Weinberg equilibrium, because $\text{INFO}_j \approx \frac{\text{var}(G_j)}{2\hat{\theta}_j(1-\hat{\theta}_j)}$. In simulations (cf. Methods section "Data for simulations"), we verify that $\text{sd}(G_j)^{\text{true}} \approx \text{sd}(G_j)^{\text{imp}}/\sqrt{\text{INFO}_j}$ (Figure S7). As a direct consequence of the lower standard deviation of dosages, we also show that GWAS effect sizes $\hat{\gamma}$ computed from imputed dosages are overestimated: $\hat{\gamma}_j^{\text{true}} \approx \hat{\gamma}_j^{\text{imp}} \cdot \sqrt{\text{INFO}_j}$ and $\text{se}(\hat{\gamma}_j)^{\text{true}} \approx \text{se}(\hat{\gamma}_j)^{\text{imp}} \cdot \sqrt{\text{INFO}_j}$ (Figures S9 and S8). This is the first correction of summary statistics we consider in the simulations below. Note that we recompute INFO scores for the subset of 362,307 European individuals used in this paper since they can differ substantially from the ones reported by the UK Biobank for the whole data (Figures S5 and S6). As a second option, instead of using dosages to compute the GWAS summary statistics, it has been argued that using multiple imputation (MI) would be more appropriate (Palmer and Pe'er, 2016). In simulations, we show that $\hat{\gamma}_j^{\text{MI}} \approx \hat{\gamma}_j^{\text{imp}} \cdot \text{INFO}_j$ and $Z_j^{\text{MI}} \approx Z_j^{\text{imp}} \cdot \text{INFO}_j$, where $Z = \hat{\gamma}/\text{se}(\hat{\gamma})$ (Figure S10). This is the second correction of summary statistics we implement in the simulations below, along with $n_j \cdot \text{INFO}_j$ as the new per-variant sample sizes. Finally, we consider an in-between solution as a third

correction, using $\hat{\gamma}_j = \hat{\gamma}_j^{\mathrm{imp}} \cdot \mathrm{INFO}_j$, $\mathrm{se}(\hat{\gamma}_j) = \mathrm{se}(\hat{\gamma}_j)^{\mathrm{imp}} \cdot \sqrt{\mathrm{INFO}_j}$, and $n_j \cdot \mathrm{INFO}_j$.

Using 40,000 variants from chromosome 22, we simulate quantitative phenotypes assuming a heritability of 20% and 2000 causal variants using the "true" dataset (cf. Methods section "Data for simulations"). We compute GWAS summary statistics from the dosage dataset and use these summary statistics to run LDpred2 and lassosum2 with either no correction of the summary statistics, or with one of the three corrections described above. The LD reference used by LDpred2 and lassosum2 is computed from the validation set using the dataset with the "true" genotypes. For lassosum2 and LDpred2(-grid), which tune parameters using the validation set, correcting for imputation quality slightly improves predictive performance in these simulations (Figure 2). However, correcting for imputation quality can dramatically improve predictive performance for LDpred2-auto, provided the imputation quality is well estimated. Moreover, new additions for robustness introduced before, namely LDpred2-low-h2, LDpred2-auto-rob, and forming independent blocks in the LD matrix also improve predictive performance for all corrections (Figure 2).

In the real data applications hereinafter, we choose to use the first correction, "sqrt_info", which is simple because it is equivalent to post-processing PGS effects by multiplying them by $\sqrt{\mathrm{INFO}}$.

Figure 2: Results of predictive performance for the simulations using GWAS summary statistics from imputed dosage data, averaged over 10 simulations for each scenario. Reported 95% confidence intervals are computed from 10,000 non-parametric bootstrap replicates of the mean. Correction "sqrt_info" corresponds to using $\hat{\gamma}_j^{\mathrm{imp}} \cdot \sqrt{\mathrm{INFO}_j}$ and $\mathrm{se}(\hat{\gamma}_j)^{\mathrm{imp}} \cdot \sqrt{\mathrm{INFO}_j}$. Correction "info" corresponds to using $\hat{\gamma}_j^{\mathrm{imp}} \cdot \mathrm{INFO}_j$ and $n_j \cdot \mathrm{INFO}_j$. Correction "in_between" corresponds to using $\hat{\gamma}_j^{\mathrm{imp}} \cdot \mathrm{INFO}_j$, $\mathrm{se}(\hat{\gamma}_j)^{\mathrm{imp}} \cdot \sqrt{\mathrm{INFO}_j}$, and $n_j \cdot \mathrm{INFO}_j$. Red bars correspond to using the LD with independent blocks (Methods).

2.3 Mismatch between LD reference and GWAS summary statistics

Here we design simulations to understand the impact of using a mismatched LD reference panel, e.g. that comes from a different population compared to the one used to compute the GWAS summary statistics. We use the same simulation setup as before (Methods). In addition, we design an alternative reference panel based on 10,000 individuals from South Europe by using the "Italy" center defined in Privé *et al.* (2022). When using this alternative LD reference panel instead of a well-matched one as in the previous sections, squared correlations between the polygenic scores and the simulated phenotypes drop from 0.174 to 0.167 for lassosum2, from 0.168 to 0.147 for LDpred2(-grid), from 0.174 to 0.169 for LDpred2-low-h2, from 0.143 to 0.139 for LDpred2-auto, from 0.162 to 0.146 for LDpred2-auto-rob, and from 0.144 to 0.140 for LDpred2-inf (Figure S12, averaging over 10 simulations). Forming independent LD blocks in these two LD matrices always helps, but do not improve performance by much when using the alternative LD.

2.4 Application to breast cancer summary statistics

Breast cancer summary statistics are interesting because they include results from two mega analyses (Michailidou *et al.*, 2013, 2015, 2017), which means that parameters reported in these GWAS summary statistics, such as INFO scores and sample sizes, are estimated with high precision. Imputation INFO scores for the OncoArray summary statistics are generally very good (mean of 0.968 after having restricted to HapMap3 variants, Figure S14) and better than the ones from iCOGS (mean of 0.841, Figure S13), probably because the iCOGS chip included around 200K variants only, compared to more than 500K variants for the OncoArray. For both summary statistics, we compare the standard deviations (of genotypes) inferred from the reported allele frequencies (i.e. $\sqrt{2f(1-f)}$ where f is the allele frequency, and denoted as sd_af) versus the ones inferred from the GWAS summary statistics (Equation (2), and denoted as sd_ss). As shown in figures 3 and S17, there is a clear trend with sd_ss being lower than sd_af as INFO decreases; indeed, using sd_ss/$\sqrt{\text{INFO}}$ provides a very good fit for sd_af, except for some variants of chromosome 6 and 8 for the OncoArray summary statistics. Most of these outlier variants are either in region 25-33 Mbp of chromosome 6 or in 8-12 Mbp of chromosome 8 (Figure S15), which are two known long-range LD regions (Price *et al.*, 2008). We hypothesize that this is due to using principal components (PCs) that capture LD structure instead of population structure, as covariates in the GWAS (Privé *et al.*, 2020a). To validate this hypothesis, we simulate a phenotype using HapMap3 variants of chromosome 6 for 10,000 individuals from the UK Biobank, then we run GWAS with or without PC19 as covariate. PC19 from the UK Biobank was previously reported to capture LD structure in region 70-91 Mbp of chromsome 6 (Privé *et al.*, 2020a). In these simulations, the same bias as in figure 3B is observed for the variants in this region (Figure S16), confirming our hypothesis.

Therefore, providing an accurate imputation INFO score is useful for two reasons. First, it allows for correcting for a reduced standard deviation when using imputed data in the QC step we propose, in order to better uncover problems with the summary statistics. Second, using one of the proposed corrections based on INFO scores may lead to an improved prediction when deriving polygenic scores. We apply these corrections to the two breast cancer summary statistics. In the comparison, we use the QC proposed in Privé *et al.* (2020b) first (which ends up filtering on MAF here, which we call "qc1"). We then also filter out the two long-range LD regions of chromosome 6 and 8 for the OncoArray summary statistics and remove around

Figure 3: Standard deviations inferred from the OncoArray breast cancer GWAS summary statistics using equation 2 (**A:** Raw or **B:** dividing them by $\sqrt{\text{INFO}}$) versus the ones inferred from the reported GWAS allele frequencies f_j (using $\sqrt{2f_j(1 - f_j)}$). Only 100,000 HapMap3 variants are represented, at random.

500 variants when filtering on differences of MAFs between summary statistics and the validation dataset ("qc2"). As for the correction using INFO scores, we use the first correction, "sqrt_info", which is simple because it is equivalent to post-processing PGS effects by multiplying them by $\sqrt{\text{INFO}}$. Although results for both QC used are very similar, correcting using INFO scores slightly improves predictive performance when deriving polygenic scores based on iCOGS summary statistics (Figure 4). All other improvements introduced before have little to no effect here, probably because misspecifications are much smaller than in the simulations.

2.5   Results for other phenotypes

We use other external GWAS summary statistics for which INFO scores are reported (Table 2); they all have a very high mean INFO score (larger than 0.94), except for T1D-affy (0.885). QC plots comparing standard deviations usually show little deviation from the identity line (after the INFO score correction), except for coronary artery disease (CAD) summary statistics (Figures S17-S22). In figures S23-S26, we then provide similar results as figure 4 for other phenotypes. Most changes introduced before have little to no impact on predictive performance, except for the second quality control performed for CAD when using LDpred2-auto (Figure S23).

Figure 4: Raw partial correlations (for *all* models) for predicting breast cancer in the UK Biobank when using either the OncoArray or the iCOGS summary statistics. These are computed using function `pcor` of R package bigstatsr where 95% confidence intervals are obtained through Fisher's Z-transformation.

3   Discussion

We have investigated misspecifications in GWAS summary statistics, focusing particularly on the impact of sample size heterogeneity and imputation quality, and the application to polygenic scores (PGS) methods. Previously, we proposed a quality control (QC) based on comparing standard deviations (of genotypes) inferred from GWAS summary statistics with the ones computed from a reference panel (Privé *et al.*, 2020b). Here we show that we can refine this QC by deriving the latter directly from the reported allele frequencies in the GWAS summary statistics, and by correcting the former using imputation INFO scores. Using this refined QC, we are able to identify a potential issue with how principal components were derived in a GWAS of breast cancer. Fortunately, this has practically no effect on the predictive performance of the derived polygenic scores. Additional QC can also be performed, e.g. comparing reported GWAS allele frequencies with the ones from the LD reference panel, e.g. to detect genotyping errors or allele inversions. We perform this additional QC as part of "qc2" here. One can also run other QC tools such as DENTIST (Chen *et al.*, 2021), and also infer ancestry proportions from summary statistics to make sure these are matching with the LD reference used (Privé, 2021b).

Note that, in this study, we use GWAS summary statistics that include extended information (e.g. INFO scores and allele frequencies), yet most GWAS summary statistics do not provide such exhaustive information (MacArthur *et al.*, 2021). We acknowledge that, in the case of a meta-analysis from multiple studies, providing a single INFO score per variant may not be possible. Solutions such as using a weighted averaged INFO score might be worth exploring in future studies. Nevertheless, this quality control could be performed within each study before meta-analyzing results, to make sure that resulting summary statistics have

the best possible quality for follow-up analyses such as deriving polygenic scores. Another information, the effective sample size per variant, is often missing from GWAS summary statistics. Sometimes, it can even be challenging to recover the total effective sample size from large meta-analyses. We recall that if some studies of a meta-analysis have an imbalanced number of cases and controls, the total effective sample size should not be computed from the total numbers of cases and controls overall, but instead from the sum of the effective sample sizes of each study. Indeed, take the extreme example of meta-analyzing two studies, one with 1000 cases and 0 controls, and another one with 0 cases and 1000 controls, then the effective sample size of the meta-analysis is 0, not 2000. Misspecifying the GWAS sample size can lead to serious issues such as misestimating the SNP heritability (Grotzinger *et al.*, 2021). Fortunately, an overestimated sample size can be detected from the QC plot we propose, where the slope is then less than 1 for case-control studies using logistic regression; otherwise the standard deviation of the phenotype is also needed (Equation 1), but can be estimated (Privé *et al.*, 2020b).

We have assessed the impact of these misspecifications in GWAS summary statistics on the predictive performance of some polygenic score methods. Using both the Bayesian LDpred2 models (Privé *et al.*, 2020b) and our reimplementation of the frequentist lassosum model (Mak *et al.*, 2017) for deriving polygenic scores, we have introduced and investigated some changes to possibly make these models more robust to misspecifications. Overall, these changes provided large improvements of predictive performance in the simulations with large misspecifications. However, they had little to no effect when using the real GWAS summary statistics we chose, except for breast cancer and coronary artery disease summary statistics. Although these results are somewhat unfortunate, they are reassuring because it means that the summary statistics we use in this study are usually of good quality for follow-up analyses such as deriving polygenic scores.

Nevertheless, we recommend to adopt these changes, i.e. performing the (refined) QC proposed here, forming independent LD blocks in the LD matrix, and using more regularization when needed. More regularization can be achieved by testing additional smaller values for the heritability parameter in LDpred2-grid, and using the two new parameters introduced here in LDpred2-auto (Section 4.6). Note that LD blocks are already widely used by several methods (such as lassosum and PRS-CS) because they allow for processing smaller matrices at once (Mak *et al.*, 2017; Ge *et al.*, 2019); here we have shown that these blocks are also useful to make methods more robust. PRS-CS is currently one of the most robust PGS methods; for example, it can use the (small) 1000 Genomes data as LD reference (Ge *et al.*, 2019), and can even use a European LD reference panel with multi-ancestry GWAS summary statistics (Wang *et al.*, 2021). We think this is made possible by the use of a (very) strong regularization in PRS-CS ($\phi^{-1}\psi_j^{-1} \geq 1$, which would approximately correspond to using $s = 0.5$ in lassosum, $\delta = 1$ in lassosum2, and `shrink_corr = 0.5` in LDpred2-auto). Using enough regularization is good for robustness, but note that using too much regularization can also damage predictive performance. To address this limitation, we recommend to rely on a proper QC and choice of ancestry-matched LD instead of on too much regularization. As a future research direction, we are interested in using multi-ancestry-matched LD matrices to improve polygenic prediction in all ancestries. It would also be useful to investigate the impact of the QC and adjustments we propose here on other (non-PGS) methods, for which consistency and robustness are likely to be very important as well (Chen *et al.*, 2021).

4 Materials and Methods

4.1 Data for simulations

We use the UK Biobank imputed (BGEN) data (Bycroft *et al.*, 2018). We restrict individuals to the ones used for computing the principal components (PCs) in the UK Biobank (Field 22020). These individuals are unrelated and have passed some quality control including removing samples with a missing rate on autosomes larger than 0.02, having a mismatch between inferred sex and self-reported sex, and outliers based on heterozygosity (more details can be found in section S3 of Bycroft *et al.* (2018)). To get a set of genetically homogeneous individuals, we compute a robust Mahalanobis distance based on the first 16 PCs and further restrict individuals to those within a log-distance of 5 (Privé *et al.*, 2020a). This results in 362,307 individuals. We randomly sample 300,000 individuals to form a training set (e.g. to run the GWAS), 10,000 individuals to form a validation set (to tune hyper-parameters), and use the remaining 52,307 individuals to form a test set (to evaluate final predictive models).

Among genetic variants on chromosome 22 and with a minor allele frequency larger than 0.01 and an imputation INFO score larger than 0.4 (as reported by the UK Biobank), we sample 40,000 of them according to the inverse of the INFO score density so that they have varying levels of imputation accuracy (Figure S4). We read the UK Biobank data into two different datasets using function `snp_readBGEN` from R package bigsnpr (Privé *et al.*, 2018), one by reading the BGEN data at random according to imputation probabilities, and another one reading it as dosages (i.e. expected values according to imputation probabilities). The first dataset is used as what could be the real genotype calls and the second dataset as what would be its imputed version; this design technique was used in Privé *et al.* (2019).

4.2 Data for real analyses

We also use the UK Biobank data, and use the same individuals as described in the previous section. We sample 10,000 individuals to form a validation set and use the remaining 352,307 individuals as test set. We restrict to the 1,054,315 HapMap3 variants used in the LD reference provided in Privé *et al.* (2020b).

To define phenotypes in the UK Biobank, we first map ICD10 and ICD9 codes (UKBB fields 40001, 40002, 40006, 40013, 41202, 41270 and 41271) to phecodes using R package PheWAS (Carroll *et al.*, 2014; Wu *et al.*, 2019). We use published GWAS summary statistics listed in table 2 to derive polygenic scores.

Trait	GWAS citation	Effective GWAS sample size	# GWAS variants	# matched variants with INFO > 0.4	Mean INFO
Breast cancer (BRCA) [iCOGS]	Michailidou *et al.* (2017)	87,037	11,792,542	1,051,242	0.841
Breast cancer (BRCA) [OncoArray]	Michailidou *et al.* (2017)	104,442	11,792,542	1,054,233	0.968
Type 1 diabetes (T1D) [Affymetrix]	Censin *et al.* (2017)	5516	8,996,866	934,712	0.885
Type 1 diabetes (T1D) [Illumina]	Censin *et al.* (2017)	7982	8,996,866	949,334	0.942
Prostate cancer (PRCA)	Schumacher *et al.* (2018)	135,316	20,370,946	818,400	0.969
Depression (MDD) [without UKBB]	Wray *et al.* (2018)	110,464	9,874,289	1,049,455	0.968
Coronary artery disease (CAD)	Nikpay *et al.* (2015)	129,014	9,455,778	1,052,200	0.982

Table 2: Summary of external GWAS summary statistics used. PRCA summary statistics have many variants with a missing INFO score, which we discard. We also restrict to variants with an INFO score larger than 0.4.

4.3 GWAS sample size imputation

In this paper, we extensively use the following formula

$$\text{sd}(G_j) \approx \frac{\text{sd}(y)}{\sqrt{n_j \, \text{se}(\hat{\gamma}_j)^2 + \hat{\gamma}_j^2}} \, , \tag{1}$$

where $\hat{\gamma}_j$ is the marginal GWAS effect size of variant j, n_j is the GWAS sample size associated with variant j, y is the vector of phenotypes and G_j is the vector of genotypes for variant j. This formula is used in LDpred2 Privé *et al.* (2020b, 2022). Note that, for a binary trait for which logistic regression is used, we have instead

$$\text{sd}(G_j) \approx \frac{2}{\sqrt{n_j^{\text{eff}} \, \text{se}(\hat{\gamma}_j)^2 + \hat{\gamma}_j^2}} \, , \tag{2}$$

where $n_j^{\text{eff}} = \frac{4}{1/n_j^{\text{cases}} + 1/n_j^{\text{controls}}}$.

We can then impute n_j from equation (1) using

$$n_j \approx \frac{\text{var}(y)/\text{var}(G_j) - \hat{\gamma}_j^2}{\text{se}(\hat{\gamma}_j)^2} \, , \tag{3}$$

and impute n_j^{eff} from equation (2) using

$$n_j^{\text{eff}} \approx \frac{4/\text{var}(G_j) - \hat{\gamma}_j^2}{\text{se}(\hat{\gamma}_j)^2} \, . \tag{4}$$

In practice, we also bound this estimate to be between $0.5 \cdot N$ and $1.1 \cdot N$, where N is the total (effective) sample size. Also note that the estimate of $\text{var}(G_j)$ has to account for the imputation accuracy (Section 2.2).

4.4 New implementation of lassosum in bigsnpr

Instead of using a regularized version of the correlation matrix R parameterized by s, $R_s = (1 - s)R + sI$ (where $0 < s \leq 1$), we use $R_\delta = R + \delta I$ (where $\delta > 0$), which makes it clearer that lassosum is also using L2-regularization (therefore elastic-net). Then, from Mak *et al.* (2017), the solution from lassosum can be obtained by iteratively updating

$$\beta_j^{(t)} = \begin{cases} \dfrac{\text{sign}\big(u_j^{(t)}\big)\big(\big|u_j^{(t)}\big| - \lambda\big)}{\widetilde{X}_j^T \widetilde{X}_j + \delta} & \text{if } \big|u_j^{(t)}\big| > \lambda \, , \\ 0 & \text{otherwise.} \end{cases}$$

where

$$u_j^{(t)} = r_j - \widetilde{X}_j^T \big(\widetilde{X}\beta^{(t-1)} - \widetilde{X}_j \beta_j^{(t-1)} \big) \, .$$

Following the notations from Privé *et al.* (2020b), denote $\widetilde{X} = \frac{1}{\sqrt{n-1}} C_n G S^{-1}$, where G is the genotype matrix, C_n is the centering matrix and S is the diagonal matrix of standard deviations of the columns of G. Then $\widetilde{X}_j^T \widetilde{X} = R_{j,.} = R_{.,j}^T$ and $\widetilde{X}_j^T \widetilde{X}_j = 1$. Moreover, using the notations from Privé *et al.* (2020b),

$$u_j^{(t)} = \widehat{\beta}_j - R_{\cdot,j}^T \beta^{(t-1)} + \beta_j^{(t-1)}, \text{ where } r_j = \widehat{\beta}_j = \frac{\widehat{\gamma}_j}{\sqrt{n_j \operatorname{se}(\widehat{\gamma}_j)^2 + \widehat{\gamma}_j^2}} \text{ and } \widehat{\gamma}_j \text{ is the GWAS effect of variant}$$

j and n is the GWAS sample size (Mak *et al.*, 2017; Privé *et al.*, 2022). Then computing $R_{\cdot,j}^T \beta^{(t-1)}$ is the most time-consuming part of each iteration. To make this faster, instead of computing $R_{\cdot,j}^T \beta^{(t-1)}$ at each iteration (j and t), we can start with an initial vector of 0s only (for all j) since $\beta^{(0)} \equiv 0$, then update this vector when $\beta_j^{(t)} \neq \beta_j^{(t-1)}$ only. Note that only positions k for which $R_{k,j} \neq 0$ must be updated in this vector $R_{\cdot,j}^T \beta^{(t-1)}$.

In this new implementation of the lassosum model, which we call lassosum2, the input parameters are the correlation matrix R, the GWAS summary statistics ($\widehat{\gamma}_j$, $\operatorname{se}(\widehat{\gamma}_j)$ and n_j), and the two hyper-parameters λ and δ. Therefore, except for the two hyper-parameters, lassosum2 uses the exact same input parameters as LDpred2 (Privé *et al.*, 2020b). We try $\delta \in \{0.001, 0.005, 0.02, 0.1, 0.6, 3\}$ by default in lassosum2, instead of $s \in \{0.2, 0.5, 0.8, 1.0\}$ in lassosum. For λ, the default in lassosum uses a sequence of 20 values equally spaced on a log scale between 0.1 and 0.001. By default in lassosum2, we use a similar sequence, but between λ_0 and $\lambda_0/100$ instead, where $\lambda_0 = \max_j |\widehat{\beta}_j|$ is the minimum λ for which no variable enters the model because the L1-regularization is too strong. Note that we do not provide an "auto" version using pseudo-validation (as in Mak *et al.* (2017)) as we have not found it to be robust enough (Figure S27). Also note that, as in LDpred2, we run lassosum2 genome-wide using a sparse correlation matrix which assumes that variants further away than 3 cM are not correlated. Although we do not require splitting the genome into independent LD blocks anymore (as done in lassosum), we recommend to do so for robustness in LDpred2, and for extra speed gain (cf. next section).

4.5 New LD reference

We form nearly independent LD blocks using the optimal algorithm developed in Privé (2021a). For different numbers of blocks and maximum number of variants in each block, we use the split with the minimum cost within the ones reducing the original number of non-zero values to less than 60% (70% for chromosome 6). Having a correlation matrix with independent blocks prevents small errors in the algorithm (e.g. the Gibbs sampler in LDpred2) from propagating to too many variants. It also makes running LDpred2 (and lassosum2) faster, taking about 60% of the initial time (since only 60% of the initial non-zero values of the correlation matrix are kept).

We have also developed a new "compact" format for the SFBMs (sparse matrices on disk). Instead of using something similar to the standard "compressed sparse column" format which stores all $\{i, x(i,j)\}$ for a given column j, we only store the first index i_0 and all the contiguous values $\{x(i_0, j), x(i_0 + 1, j), \dots\}$ up to the last non-zero value for this column j. This makes this format about twice as efficient for both LDpred2 and lassosum2 (in terms of both memory and speed). Therefore, using both this new format and the LD blocks should divide computation times by 3.

4.6 LDpred2-low-h2 and LDpred2-auto-rob

Here we introduce the small changes made to LDpred2 (-grid and -auto) in order to make them more robust. First, LDpred2-low-h2 simply consists in running LDpred2-grid by testing h^2 within $\{0.3, 0.7, 1, 1.4\} \cdot h_{\text{LDSC}}^2$

(note the added 0.3 compared to Privé *et al.* (2020b)), where h^2_{LDSC} is the heritability estimate from LD score regression. Indeed, we show in simulations here that using lower values for h^2 may provide higher predictive performance in the case of misspecifications (thanks to more shrinkage of the effects). In simulations, because of the large misspecifications, we use a larger grid over $\{0.01, 0.1, 0.3, 0.7, 1, 1.4\} \cdot h^2_{\text{LDSC}}$.

For LDpred2-auto, we introduce two new parameters. The first one, `shrink_corr`, allows for shrinking off-diagonal elements of the correlation matrix. This is similar to parameter 's' in lassosum (Section 4.4), and acts as a means of regularization. We use a value of 0.9 in simulations and 0.95 in real data when running "LDpred2-auto-rob" (and the default value of 1, with no effect, when running "LDpred2-auto"). The second new parameter, `allow_jump_sign`, controls whether variant effect sizes can change sign over two consecutive iterations of the Gibbs sampler. When setting this parameter to false (in the method we refer to as "LDpred2-auto-rob" here), this forces the effects to go through 0 before changing sign. This is useful to prevent instability (oscillation and ultimately divergence) of the Gibbs sampler under large misspecifications, and is also useful for accelerating convergence of chains with a large initial value for p (the proportion of causal variants).

Code and data availability

All code used for this paper is available at `https://github.com/privefl/paper-misspec/tree/master/code`. We have extensively used R packages bigstatsr and bigsnpr (Privé *et al.*, 2018) for analyzing large genetic data, packages from the future framework (Bengtsson, 2021) for easy scheduling and parallelization of analyses on the HPC cluster, and packages from the tidyverse suite (Wickham *et al.*, 2019) for shaping and visualizing results.

The latest version of R package bigsnpr can be installed from GitHub, and a recent enough version from CRAN. Two tutorials on running LDpred2 and lassosum2 using R package bigsnpr are available at `https://privefl.github.io/bigsnpr/articles/LDpred2.html` and `https://privefl.github.io/bigsnpr-extdoc/polygenic-scores-pgs.html`.

We have recomputed the allele frequencies and imputation INFO scores within the European subset used here and across all imputed variants in the UK Biobank, and made them available at `https://doi.org/10.6084/m9.figshare.16635388`. We have formed independent LD blocks within the European LD references provided in Privé *et al.* (2020b), and made these updated versions available at `https://doi.org/10.6084/m9.figshare.19213299`.

Acknowledgements

Authors thank Timothy Shin Heng Mak, Shing Wan Choi, and Matthew Stephens for helpful discussions. Authors also thank GenomeDK and Aarhus University for providing computational resources and support that contributed to these research results. This research has been conducted using the UK Biobank Resource under Application Number 58024.

Funding

F.P. and B.J.V. are supported by the Danish National Research Foundation (Niels Bohr Professorship to Prof. John McGrath) and by a Lundbeck Foundation Fellowship (R335-2019-2339 to B.J.V.).

Declaration of Interests

B.J.V. is on Allelica's international advisory board. The other authors declare no competing interests.

References

Bengtsson, H. (2021). A Unifying Framework for Parallel and Distributed Processing in R using Futures. *The R Journal*.

Bycroft, C., Freeman, C., Petkova, D., Band, G., Elliott, L. T., Sharp, K., Motyer, A., Vukcevic, D., Delaneau, O., O'Connell, J., *et al.* (2018). The UK Biobank resource with deep phenotyping and genomic data. *Nature*, **562**(7726), 203–209.

Carroll, R. J., Bastarache, L., and Denny, J. C. (2014). R PheWAS: data analysis and plotting tools for phenome-wide association studies in the R environment. *Bioinformatics*, **30**(16), 2375–2376.

Censin, J., Nowak, C., Cooper, N., Bergsten, P., Todd, J. A., and Fall, T. (2017). Childhood adiposity and risk of type 1 diabetes: A mendelian randomization study. *PLoS Medicine*, **14**(8), e1002362.

Chen, W., Wu, Y., Zheng, Z., Qi, T., Visscher, P. M., Zhu, Z., and Yang, J. (2021). Improved analyses of GWAS summary statistics by reducing data heterogeneity and errors. *Nature Communications*, **12**(1), 7117.

Gazal, S., Loh, P.-R., Finucane, H. K., Ganna, A., Schoech, A., Sunyaev, S., and Price, A. L. (2018). Functional architecture of low-frequency variants highlights strength of negative selection across coding and non-coding annotations. *Nature Genetics*, **50**(11), 1600–1607.

Ge, T., Chen, C.-Y., Ni, Y., Feng, Y.-C. A., and Smoller, J. W. (2019). Polygenic prediction via Bayesian regression and continuous shrinkage priors. *Nature Communications*, **10**(1), 1776.

Grotzinger, A. D., de la Fuente, J., Nivard, M. G., and Tucker-Drob, E. M. (2021). Pervasive downward bias in estimates of liability scale heritability in GWAS meta-analysis: A simple solution. *medRxiv*.

Kulm, S., Marderstein, A., Mezey, J., and Elemento, O. (2021). A systematic framework for assessing the clinical impact of polygenic risk scores. *medRxiv*, pages 2020–04.

Lloyd-Jones, L. R., Zeng, J., Sidorenko, J., Yengo, L., Moser, G., Kemper, K. E., Wang, H., Zheng, Z., Magi, R., Esko, T., *et al.* (2019). Improved polygenic prediction by Bayesian multiple regression on summary statistics. *Nature Communications*, **10**(1), 1–11.

Loh, P.-R., Kichaev, G., Gazal, S., Schoech, A. P., and Price, A. L. (2018). Mixed-model association for biobank-scale datasets. *Nature Genetics*, **50**(7), 906–908.

MacArthur, J. A., Buniello, A., Harris, L. W., Hayhurst, J., McMahon, A., Sollis, E., Cerezo, M., Hall, P., Lewis, E., Whetzel, P. L., *et al.* (2021). Workshop proceedings: GWAS summary statistics standards and sharing. *Cell Genomics*, **1**(1), 100004.

Mak, T. S. H., Porsch, R. M., Choi, S. W., Zhou, X., and Sham, P. C. (2017). Polygenic scores via penalized regression on summary statistics. *Genetic Epidemiology*, **41**(6), 469–480.

Marchini, J. and Howie, B. (2010). Genotype imputation for genome-wide association studies. *Nature Reviews Genetics*, **11**(7), 499–511.

Mbatchou, J., Barnard, L., Backman, J., Marcketta, A., Kosmicki, J. A., Ziyatdinov, A., Benner, C., O'Dushlaine, C., Barber, M., Boutkov, B., Habegger, L., Ferreira, M., Baras, A., Reid, J., Abecasis, G., Maxwell, E., and Marchini, J. (2021). Computationally efficient whole-genome regression for quantitative and binary traits. *Nature Genetics*, **53**(7), 1097–1103.

Michailidou, K., Hall, P., Gonzalez-Neira, A., Ghoussaini, M., Dennis, J., Milne, R. L., Schmidt, M. K., Chang-Claude, J., Bojesen, S. E., Bolla, M. K., *et al.* (2013). Large-scale genotyping identifies 41 new loci associated with breast cancer risk. *Nature Genetics*, **45**(4), 353–361.

Michailidou, K., Beesley, J., Lindstrom, S., Canisius, S., Dennis, J., Lush, M. J., Maranian, M. J., Bolla, M. K., Wang, Q., Shah, M., *et al.* (2015). Genome-wide association analysis of more than 120,000 individuals identifies 15 new susceptibility loci for breast cancer. *Nature Genetics*, **47**(4), 373–380.

Michailidou, K., Lindström, S., Dennis, J., Beesley, J., Hui, S., Kar, S., Lemaçon, A., Soucy, P., Glubb, D., Rostamianfar, A., *et al.* (2017). Association analysis identifies 65 new breast cancer risk loci. *Nature*, **551**(7678), 92–94.

Miller, J. W. and Dunson, D. B. (2019). Robust Bayesian inference via coarsening. *Journal of the American Statistical Association*, **114**(527), 1113–1125.

Nikpay, M., Goel, A., Won, H.-H., Hall, L. M., Willenborg, C., Kanoni, S., Saleheen, D., Kyriakou, T., Nelson, C. P., Hopewell, J. C., *et al.* (2015). A comprehensive 1000 genomes–based genome-wide association meta-analysis of coronary artery disease. *Nature Genetics*, **47**(10), 1121.

Pain, O., Glanville, K. P., Hagenaars, S. P., Selzam, S., Fürtjes, A. E., Gaspar, H. A., Coleman, J. R., Rimfeld, K., Breen, G., Plomin, R., *et al.* (2021). Evaluation of polygenic prediction methodology within a reference-standardized framework. *PLoS Genetics*, **17**(5), e1009021.

Palmer, C. and Pe'er, I. (2016). Bias Characterization in Probabilistic Genotype Data and Improved Signal Detection with Multiple Imputation. *PLoS Genetics*, **12**(6), e1006091.

Price, A. L., Weale, M. E., Patterson, N., Myers, S. R., Need, A. C., Shianna, K. V., Ge, D., Rotter, J. I., Torres, E., Taylor, K. D., *et al.* (2008). Long-range LD can confound genome scans in admixed populations. *The American Journal of Human Genetics*, **83**(1), 132–135.

Privé, F. (2021a). Optimal linkage disequilibrium splitting. *Bioinformatics*, **38**(1), 255–256.

Privé, F. (2021b). Using the UK Biobank as a global reference of worldwide populations: application to measuring ancestry diversity from GWAS summary statistics. *bioRxiv*.

Privé, F., Aschard, H., Ziyatdinov, A., and Blum, M. G. B. (2018). Efficient analysis of large-scale genome-wide data with two R packages: bigstatsr and bigsnpr. *Bioinformatics*, **34**(16), 2781–2787.

Privé, F., Vilhjálmsson, B. J., Aschard, H., and Blum, M. G. B. (2019). Making the most of clumping and thresholding for polygenic scores. *The American Journal of Human Genetics*, **105**(6), 1213–1221.

Privé, F., Luu, K., Blum, M. G., McGrath, J. J., and Vilhjálmsson, B. J. (2020a). Efficient toolkit implementing best practices for principal component analysis of population genetic data. *Bioinformatics*, **36**(16), 4449–4457.

Privé, F., Arbel, J., and Vilhjálmsson, B. J. (2020b). LDpred2: better, faster, stronger. *Bioinformatics*, **36**(22-23), 5424–5431.

Privé, F., Zhu, Z., and Vilhjalmsson, B. J. (2021). Finding hidden treasures in summary statistics from genome-wide association studies. *Nature Genetics*, **53**(4), 431–432.

Privé, F., Aschard, H., Carmi, S., Folkersen, L., Hoggart, C., O'Reilly, P. F., and Vilhjálmsson, B. J. (2022). Portability of 245 polygenic scores when derived from the uk biobank and applied to 9 ancestry groups from the same cohort. *The American Journal of Human Genetics*, **109**(1), 12–23.

Schumacher, F. R., Al Olama, A. A., Berndt, S. I., Benlloch, S., Ahmed, M., Saunders, E. J., Dadaev, T., Leongamornlert, D., Anokian, E., Cieza-Borrella, C., *et al.* (2018). Association analyses of more than 140,000 men identify 63 new prostate cancer susceptibility loci. *Nature Genetics*, **50**(7), 928.

Walker, S. G. (2013). Bayesian inference with misspecified models. *Journal of Statistical Planning and Inference*, **143**(10), 1621–1633.

Wang, Y., Namba, S., Lopera-Maya, E. A., Kerminen, S., Tsuo, K., Lall, K., Kanai, M., Zhou, W., Wu, K.-H. H., Fave, M.-J., *et al.* (2021). Global biobank analyses provide lessons for computing polygenic risk scores across diverse cohorts. *medRxiv*.

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., *et al.* (2019). Welcome to the tidyverse. *Journal of Open Source Software*, **4**(43), 1686.

Wray, N. R., Ripke, S., Mattheisen, M., Trzaskowski, M., Byrne, E. M., Abdellaoui, A., Adams, M. J., Agerbo, E., Air, T. M., Andlauer, T. M., *et al.* (2018). Genome-wide association analyses identify 44 risk variants and refine the genetic architecture of major depression. *Nature Genetics*, **50**(5), 668.

Wu, P., Gifford, A., Meng, X., Li, X., Campbell, H., Varley, T., Zhao, J., Carroll, R., Bastarache, L., Denny, J. C., *et al.* (2019). Mapping ICD-10 and ICD-10-CM codes to phecodes: workflow development and initial evaluation. *JMIR Medical Informatics*, **7**(4), e14325.

Yengo, L., Vedantam, S., Marouli, E., Sidorenko, J., Bartell, E., Sakaue, S., Graff, M., Eliasen, A. U., Jiang, Y., Raghavan, S., *et al.* (2022). A saturated map of common genetic variants associated with human height from 5.4 million individuals of diverse ancestries. *bioRxiv*.

Zheng, J., Erzurumluoglu, A. M., Elsworth, B. L., Kemp, J. P., Howe, L., Haycock, P. C., Hemani, G., Tansey, K., Laurin, C., Pourcain, B. S., *et al.* (2017). LD Hub: a centralized database and web interface to perform LD score regression that maximizes the potential of summary level GWAS data for SNP heritability and genetic correlation analysis. *Bioinformatics*, **33**(2), 272–279.

Zhou, G. and Zhao, H. (2021). A fast and robust Bayesian nonparametric method for prediction of complex traits using summary statistics. *PLoS Genetics*, **17**(7), e1009697.

Zhou, W., Nielsen, J. B., Fritsche, L. G., Dey, R., Gabrielsen, M. E., Wolford, B. N., LeFaive, J., VandeHaar, P., Gagliano, S. A., Gifford, A., *et al.* (2018). Efficiently controlling for case-control imbalance and sample relatedness in large-scale genetic association studies. *Nature Genetics*, **50**(9), 1335–1341.

Zhu, X. and Stephens, M. (2017). Bayesian large-scale multiple regression with summary statistics from genome-wide association studies. *The Annals of Applied Statistics*, **11**(3), 1561.

Zou, Y., Carbonetto, P., Wang, G., and Stephens, M. (2021). Fine-mapping from summary data with the "sum of single effects" model. *bioRxiv*.