Decomposing virulence to understand bacterial clearance in persistent infections

Beatriz Acuña Hidalgo [1¶], Luís M. Silva [1,2¶], Roland R. Regoes [3], Mathias Franz [1], Sophie A. O. Armitage [1*]

[1] Institute of Biology, Freie Universität Berlin, Berlin, Germany

[2] Institute of Biology, University of Neuchâtel, Neuchâtel, Switzerland

[3] Institute of Integrative Biology, ETH Zurich, Zurich, Switzerland

[*] Corresponding author

E-mail: sophie.armitage@fu-berlin.de (SAOA)

[¶] These authors contributed equally to this work.

Abstract

Hosts are not always successful at controlling and eliminating a pathogen and the factors

causing variation in pathogen clearance are not well understood. To address this problem, we

used *Drosophila melanogaster* to investigate how infections with different bacterial

pathogens affects virulence, clearance and persistence. In this context we developed novel

hypotheses that focus on how variation in clearance should be related to variation in different

components of virulence, where virulence is the infection-related reduction in host fitness. To

achieve this, virulence was decomposed into exploitation, i.e., how well bacteria can replicate

inside the host, and per parasite pathogenicity (PPP), i.e., the amount of damage per parasite

inflicted on the host. We used four bacterial species: *Enterobacter cloacae*, *Providencia*

burhodogranariea, Lactococcus lactis and *Pseudomonas entomophila*. The injection doses

spanned four orders of magnitude, and survival was followed to estimate virulence. Bacterial

load was quantified in live flies during the acute (1-4 days) and chronic (7-35 days) phases of

infection, and we tested infection status of flies that had died up to ten weeks post infection.

We show that sustained persistent infection and clearance are both possible outcomes for

bacterial species across a range of virulence. Bacteria of all species could persist inside the

host for at least 75 days, and injection dose partly predicted within-species variation in

clearance. Our decomposition of virulence showed that species differences in bacterial

virulence could be explained by a combination of variation in both exploitation and PPP. In

addition, we found that that higher exploitation leads to lower bacterial clearance, whereas we

could not detect any effect of PPP on clearance. The differing effects of exploitation and PPP

imply that there can be different means by which variation in virulence is related to clearance,

which could critically affect pathogen transmission and the evolution of pathogen virulence.

Author summary

Following an infection, hosts are not always able to quickly clear the pathogen, and they instead either die or survive with a persistent infection. Such variation is ecologically and evolutionarily important, because it can affect infection prevalence and transmission, and also virulence evolution. But what causes variation in infection outcomes? Here we contribute towards answering this question by investigating infection dynamics in flies infected with one of four bacterial species. We first establish that the bacterial species differ in virulence, *i.e.*, the host death rate after infection. We find that variation in virulence arises because the bacteria differ in the two components of virulence: bacterial growth inside the host (exploitation), and the amount of damage caused per bacterium (per parasite pathogenicity). Furthermore, as early-phase exploitation increases, bacterial clearance later in the infection decreases. This finding can be explained by increasing exploitation leading to increasing clearance costs for the host. Taken together we demonstrate that variation in infection outcomes can be partly explained by how virulence, and its components, relate to the rate of pathogen clearance. We propose that the decomposition of virulence is valuable for understanding variation in infection outcomes – potentially also beyond the interrelation between virulence and clearance.

1. Introduction

Once a host has become infected, the immune system will potentially limit pathogen growth, a response termed host resistance [1-3]. Resistance can therefore be quantified as the inverse of pathogen load [2]. Although there are clear benefits to the host of being able to mount an immune response that suppresses pathogen growth, resistance can come with evolutionary [4, 5] and usage costs [6] for the host [reviewed in 7]. During infection, hosts may re-allocate resources from other life history traits, such as reproduction [8] or development [9], into mounting an immune response. Furthermore, immune responses can lead to self-inflicted damage to the host, namely immunopathology [10-12]. Therefore, whether a pathogen is eliminated or not, *i.e.,* persists, is likely to depend upon the costs of infection *versus* the costs and effectiveness of the immune response against the infection, in addition to how well the pathogen can survive and replicate in the host environment.

Across host taxa, there is ample evidence of persistent bacterial infections. For example bacterial infections caused by *Escherichia coli* and *Staphylococcus aureus* can evade the human immune system and persist inside the host [13]. After injection with bacteria, insects can also sustain persistent systemic infections, for example in the mosquito *Anopheles gambiae* [14], the fruit fly *D. melanogaster* [15, 16] and the yellow mealworm beetle *Tenebrio molitor* [17]. These experimentally-induced infections can persist for at least 28 days in both *T. molitor* [17] and *D. melanogaster* [18], although longer term estimates are lacking.

Disparate bacterial species have been shown to be able to chronically infect (here defined as a minimum of seven days) the host species used in this study, *D. melanogaster* [15, 16, 18-23]. Persistent infections could be influential because the inability to clear an infection will result in more infected individuals in a population, thereby potentially increasing the probability of

pathogen transmission. In the chronic infection phase in *D. melanogaster,* the bacterial load

has been shown to stabilise around a relatively constant pathogen load over time [16, 21],

which has been termed the set point bacterial load [SPBL; 21], after the set point viral load

[*e.g.*, 24]. However a stable infection load over time is not necessarily always the case, as the

load for some bacterial species can gradually reduce in the days following infection, for

example, in *D. melanogaster* injected with *E. coli* [20] and *T. molitor* injected with *S. aureus*

[17, 25]. Alternatively, after an initial decline the infection load can start to increase again, as

seen in the burying beetle, *Nicrophorus vespilloides*, injected with *Photorhabdus luminescens*

[26].

Bacterial clearance during the chronic infection phase has not been commonly reported in

insects and it may be related to the costs and benefits of immune system activation, and the

degree of pathogen virulence. Virulence can be defined as disease severity, given as the

decrease in host fitness caused by a pathogen [27], and which we here measure as reduced

host survival. Virulence will be influenced by both host and parasite traits, *i.e.*, it depends on

resistance and tolerance from the host side, and the ability of the parasite to replicate and

cause damage to the host [28]. From the pathogen perspective, variation in virulence across

parasite strains could be due to differences in exploitation, that is an increase in virulence is a

side effect of an increase in pathogen load [28, 29] (Fig 1A). However, variation in virulence

could also be due to differences in per-parasite pathogenicity (PPP), *i.e.*, the damage inflicted

by each individual pathogen. PPP can be quantified by the slope of the reaction norm linking

infection intensity (pathogen load) and host fitness [28, 29] (Fig 1B). A parasite genotype

causing a steeper negative slope across a range of infection intensities, suggests higher PPP

compared to a parasite genotype infection resulting in a shallower slope (Fig 1B). Here we

use the concepts of exploitation and PPP to disentangle the causes of variation in virulence

caused by infection with different bacterial species. We then use this information to uncover

whether exploitation and PPP link to bacterial clearance probability.

Exploitation and PPP are conceptually and mechanistically distinct ways in which parasites

may harm their hosts. Accordingly, there could be distinct evolutionary consequences – both

for pathogen and host traits. In the pathogen, reduced PPP and thus reduced virulence may be

selected for, because this would result in a longer host infectious period without lowering the

transmission rate [28]. On the other hand, higher exploitation might be selected for given that

higher infection intensities are predicted to increase transmission rates and to take longer to

clear (described as the recovery rate). However, increased exploitation and longer clearance

will also mean increased virulence and hence trade-offs are expected between virulence,

transmission and clearance [28, 30].

We hypothesise that exploitation and PPP can affect clearance in different ways, which could

result in different, and potentially even opposing, patterns of how variation in virulence is

related to clearance. In the following we consider how hosts are expected to react to infections

with different levels of exploitation, or different levels of PPP. When the costs of mounting an

immune response exceed the benefits of clearing an infection, one might predict a host to

manage a persistent infection [31], and vice versa for clearing an infection. Therefore, we

assume that hosts are more likely to clear an infection when the ratio of benefits to costs of

clearance increases. In addition, as will be explained in more detail below, we assume that

exploitation and PPP have different effects on the costs and benefits of clearance.

We assume that PPP mainly affects the benefits of clearance. Higher PPP directly results in a

higher host death rate, which makes it more beneficial for the host to clear an infection.

Accordingly, we expect that an increase in PPP leads to an increased effort by the host to

clear the infection, which should result in a higher clearance rate (S1A Fig). In this context it

is important to consider that an increased clearance effort could also lead to increased

immunopathological effects, *i.e.*, increased clearance costs in the form of increased virulence.

Accordingly, such increased virulence would result in an increase in PPP (S1A Fig). This

would mean that the measured PPP could reflect both the pathogen contribution to virulence

and also the resulting host contribution to virulence. However, this potentially circular effect

does not change our expectation for how PPP should be qualitatively related to clearance.

Accordingly, we predict that an increase in measured PPP will be related to an increase in

measured clearance.

In contrast to PPP, we assume that exploitation, *i.e.*, infection intensity, affects both the

benefits and also the costs of clearance. Increasing exploitation results in increased virulence,

which increases clearance benefits. However, increasing exploitation additionally increases

the costs of clearance if we assume that it is more difficult, and thus costlier, to clear

infections with higher pathogen loads. Due to this dual increase in benefits and costs, two

opposing predictions can be derived. First, if benefits of clearance increase faster than costs,

we predict that increasing exploitation results in increasing clearance (S1B Fig). This

prediction is consistent with observations of viral infections in humans and non-human

primates where larger viral loads led to a faster decline in viral load or in shorter durations of

viremia [reviewed in 32; e.g., 33]. Second, if costs of clearance increase faster than benefits,

then we predict that increasing exploitation results in decreasing clearance (S1B Fig).

The initial exposure dose will also determine the outcome of infection, partly because

microbe density at the beginning of an infection can determine the strength of the immune

response [34]. Not only is dose-dependent survival frequently reported in response to

bacterial infections [19, 26, 35], but bacterial load later in the infection has been demonstrated

to correlate with the initial injection dose [19, 21]. We here test the generality of the latter

finding. Furthermore, we predict that lower injection doses are more likely to be cleared,

because a smaller population would be more susceptible to being killed by the host immune

defences. The latter has been referred to as the inoculum effect in the context of *in vitro*

antimicrobial activity studies [36-38]. However, the extent to which this kind of pattern is

generalisable to bacterial infections *in vivo* is poorly understood.

Here we injected flies with four candidate bacterial species at a range of infection doses to

address a number of different aims. First, we documented the variation in virulence and

clearance caused by different bacterial species. Specifically, we tested whether species varied

in virulence, which was measured via survival. We then decomposed virulence into its two

constituent parts, to ask whether the species-level differences in bacterial virulence that we

observed, were due to variation in parasite exploitation (infection intensity) or due to variation

in PPP. To investigate clearance, we tested whether all four bacterial species establish a

persistent infection by assessing infection status up to 35 days post injection, which is longer

than any studies that we are aware of. In the second part we then tested our predictions

regarding how variation in clearance is related to (1) exploitation and PPP, and (2) infection

dose. Lastly, we assessed the infection status of flies that had died up to ten weeks post

injection, which allowed us to assess whether long-term persistent bacterial infections occur

in insects.

2. Materials and Methods

2.1. Fly population and maintenance

We used an outbred population of *Drosophila melanogaster* established from 160 *Wolbachia-*

infected fertilised females collected in Azeitão, Portugal [39], and given to us by Élio Sucena.

For at least 13 generations prior to the start of the experiments the flies were maintained on

standard sugar yeast agar medium [SYA medium: 970 ml water, 100 g brewer's yeast, 50 g

sugar, 15 g agar, 30 ml 10 % Nipagin solution and 3 ml propionic acid; 40], in a population

cage containing at least 5,000 flies, with non-overlapping generations of 15 days. They were

maintained at $24.3 \pm 0.2°C$, on a 12:12 hours light-dark cycle, at 60-80 % relative humidity.

The experimental flies were kept under the same conditions.

2.2. Bacterial species

We used the Gram-positive *Lactococcus lactis* (gift from Brian Lazzaro), Gram negative

Enterobacter cloacae subsp. dissolvens (hereafter called *E. cloacae*; German collection of

microorganisms and cell cultures, DSMZ; type strain: DSM-16657), *Providencia*

burhodogranariea strain B (gift from Brian Lazzaro, DSMZ; type strain: DSM-19968) and

Pseudomonas entomophila (gift from Bruno Lemaitre). *L. lactis* [41], *Pr. burhodogranariea*

[42] and *Ps. entomophila* [43] were isolated from wild-collected *D. melanogaster* and can be

considered as opportunistic pathogens. *E. cloacae* was isolated from a maize plant, but has

been detected in the microbiota of *D. melanogaster* [44]. These bacterial species were chosen

based on various studies, which together suggest that they may be expected to show a range

of virulence [18, 20, 21, 45-47].

2.3. Experimental design

For each bacterial species, flies were exposed to one of seven treatments: no injection (naïve),

injection with *Drosophila* Ringer's (injection control) or injection with one of five

concentrations of bacteria ranging from 5×10^6 to 5×10^9 colony forming units (CFUs)/mL,

corresponding to doses of approximately 92, 920, 1,840, 9200 and 92,000 CFUs per fly. The

injections were done in a randomised block design by two people. Each bacterial species was

tested in three independent experimental replicates. Per experimental replicate we treated 252

flies, giving a total of 756 flies per bacterium (including naïve and Ringer's injection control

flies). Per experimental replicate and treatment, 36 flies were checked daily for survival until

all of the flies were dead. A sub-set of the dead flies were homogenised upon death to test

whether the infection had been cleared before death or not. To evaluate bacterial load in living

flies, per experimental replicate, four of the flies were homogenised per treatment, for each of

nine time points: one, two, three, four, seven, 14, 21, 28- and 35-days post-injection.

2.4. Infection assay

Bacterial preparation was performed as in Kutzer *et al.* [18], except that we grew two

overnight liquid cultures of bacteria per species, which were incubated overnight for

approximately 15 hours at 30 ºC and 200 rpm. The overnight cultures were centrifuged at

2880 rcf at 4 ºC for 10 minutes and the supernatant removed. The bacteria were washed twice

in 45 mL sterile *Drosophila* Ringer's solution [182 mmol·L-1 KCl; 46 mol·L-1 NaCl; 3

mmol·L-1 CaCl2; 10 mmol·L-1 Tris·HCl; 48] by centrifugation at 2880 rcf at 4°C for 10

minutes. The cultures from the two flasks were combined into a single bacterial solution and

the optical density (OD) of 500 µL of the solution was measured in a Ultrospec 10 classic

(Amersham) at 600 nm. The concentration of the solution was adjusted to that required for

each injection dose, based on preliminary experiments where a range of ODs between 0.1 and

0.7 were serially diluted and plated to estimate the number of CFUs. Additionally, to confirm

post hoc the concentration estimated by the OD, we serially diluted to $1:10^7$ and plated the

bacterial solution three times and counted the number of CFUs.

The experimental flies were reared at constant larval density for one generation prior to the

start of the experiments. Grape juice agar plates (50 g agar, 600 mL red grape juice, 42 mL

Nipagin [10 % w/v solution] and 1.1 L water) were smeared with a thin layer of active yeast

paste and placed inside the population cage for egg laying and removed 24 hours later. The

plates were incubated overnight then first instar larvae were collected and placed into plastic

vials (95 x 25 mm) containing 7 ml of SYA medium. Each vial contained 100 larvae to

maintain a constant density during development. One day after the start of adult eclosion, the

flies were placed in fresh food vials in groups of five males and five females, after four days

the females were randomly allocated to treatment groups.

Before injection, females were anesthetised with CO_2 for a maximum of five minutes and

injected in the lateral side of the thorax using a fine glass capillary (Ø 0.5 mm, Drummond),

pulled to a fine tip with a Narishige PC-10, and then connected to a Nanoject II™ injector

(Drummond). A volume of 18.4 nL of bacterial solution, or *Drosophila* Ringer's solution as a

control, was injected into each fly. Full controls, *i.e.*, naïve flies, underwent the same

procedure but without any injection. After being treated, flies were placed in groups of six

into new vials containing SYA medium, and then transferred into new vials every 2-5 days.

At the end of each experimental replicate, 50 µL of the aliquots of bacteria that had been used

for injections were plated on LB agar to check for potential contamination. No bacteria grew

from the Ringer's solution and there was no evidence of contamination in any of the bacterial

replicates. In addition, to confirm the concentration of the injected bacteria, serial dilutions

were prepared and plated before and after the injections for each experimental replicate, and

CFUs counted the following day.

2.5. Bacterial load of living flies

Flies were randomly allocated to the day at which they would be homogenised. Prior to

homogenisation, the flies were briefly anesthetised with CO_2 and removed from their vial.

Each individual was placed in a 1.5 mL microcentrifuge tube containing 100 µL of pre-chilled

LB media and one stainless steel bead (Ø 3 mm, Retsch) on ice. The microcentrifuge tubes

were placed in a holder that had previously been chilled in the fridge at 4 °C for at least 30

minutes to reduce further growth of the bacteria. The holders were placed in a Retsch Mill

(MM300) and the flies homogenised at a frequency of 20 Hz for 45 seconds. Then, the tubes

were centrifuged at 420 rcf for one minute at 4 ºC. After resuspending the solution, 80 μL of

the homogenate from each fly was pipetted into a 96-well plate and then serially diluted 1:10

until $1:10^5$. Per fly, three droplets of 5 μL of every dilution were plated onto LB agar. Our

lower detection limit with this method was around seven colony-forming units per fly. We

consider bacterial clearance by the host to be when no CFUs were visible in any of the

droplets. Although we note that clearance is indistinguishable from an infection that is below

the detection limit. The plates were incubated at 28 ºC and the numbers of CFUs were

counted after ~20 hours. Individual bacterial loads per fly were backcalculated using the

average of the three droplets from the lowest countable dilution in the plate, which was

usually between 10 and 60 CFUs per droplet.

D. melanogaster microbiota does not easily grow under the above culturing conditions [e.g.,

47 and personal observation]. Nonetheless we homogenised control flies (Ringer's injected

and naïve) as a control. We rarely retrieved foreign CFUs after homogenising Ringer's

injected or naïve flies (23 out of 642 cases, *i.e.*, 3.6 %). We also rarely observed

contamination in the bacteria-injected flies: except for homogenates from 27 out of 1223 flies

(2.2 %), colony morphology and colour were always consistent with the injected bacteria [see

methods of 49]. Twenty one of these 27 flies were excluded from further analyses given that

the contamination made counts of the injected bacteria unreliable; the remaining six flies had

only one or two foreign CFUs in the most concentrated homogenate dilution, therefore these

flies were included in further analyses. For *L. lactis* (70 out of 321 flies), *P.*

burhodogranaeria (7 out of 381 flies) and *Ps. entomophila* (1 out of 71 flies) there were too

many CFUs to count at the highest dilution. For these cases, we denoted the flies as having

the highest countable number of CFUs found in any fly for that bacterium and at the highest

dilution [20]. This will lead to an underestimate of the bacterial load in these flies.

2.6. Bacterial load of dead flies

2.6. Bacterial load of dead flies

For two periods of time in the chronic infection phase, *i.e.*, between 14 and 35 days and 56 to

78 days post injection, dead flies were retrieved from their vial at the daily survival checks

and homogenised in order to test whether they died whilst being infected, or whether they had

cleared the infection before death. The fly homogenate was produced in the same way as for

live flies, but we increased the dilution of the homogenate (1:1 to $1:10^{12}$) because we

anticipated higher bacterial loads in the dead compared to the live flies. The higher dilution

allowed us more easily to determine whether there was any obvious contamination from

foreign CFUs or not. Because the flies may have died at any point in the 24 hours preceding

the survival check, and the bacteria can potentially continue replicating after host death, we

evaluated the infection status (yes/no) of dead flies instead of the number of CFUs. Dead flies

were evaluated for two experimental replicates per bacteria, and 160 flies across the whole

experiment. Similar to homogenisation of live flies, we rarely observed contamination from

foreign CFUs in the homogenate of dead bacteria-injected flies (3 out of 160; 1.9 %); of these

three flies, one fly had only one foreign CFU, so it was included in the analyses. Dead

Ringer's injected and naïve flies were also homogenised and plated as controls, with 6 out of

68 flies (8.8 %) resulting in the growth of unidentified CFUs.

2.7. Statistical analyses

2.7. Statistical analyses

Statistical analyses were performed in RStudio version 1.3.1073 [50]. The following packages

were used for plotting the data: "ggpubr" [51], "grid", "gridExtra" [52], "ggplot2" [53],

"scales" [54], "survival" [55, 56] and "viridis" [57]. To include a factor as a random factor in

a model it has been suggested that there should be more than five to six random-effect levels

per random effect [58], so that there are sufficient levels to base an estimate of the variance of

the population of effects [59]. In our experimental designs, the low numbers of levels within

the factors 'experimental replicate' (two to three levels) and 'person' (two levels), meant that

we therefore fitted them as fixed, rather than random factors [59]. However, for the analysis

of clearance (see 2.7.7) we included species as a random effect because it was not possible to

include it as a fixed effect due to the fact that PPP is already a species-level predictor.

2.7.1. Do the bacterial species differ in virulence? To test whether the bacterial species

differed in virulence, we performed a linear model with the natural log of the maximum

hazard as the dependent variable and bacterial species as a factor. Post-hoc multiple

comparisons were performed using "emmeans". The hazard function in survival analyses

gives the instantaneous failure rate, and the maximum hazard gives the hazard at the point at

which this rate is highest. We extracted maximum hazard values from time of death data for

each bacterial species/dose/experimental replicate. Each maximum hazard per

species/dose/experimental replicate was estimated from an average of 33 flies (a few flies

were lost whilst being moved between vials etc.). To extract maximum hazard values we

defined a function that used the "muhaz" package [60] to generate a smooth hazard function

and then output the maximum hazard in a defined time window, as well as the time at which

this maximum is reached. To assess the appropriate amount of smoothing, we tested and

visualised results for four values (1, 2, 3 and 5) of the smoothing parameter, b, which was

specified using *bw.grid* [61]. We present the results from $b = 2$, but all of the other values

gave qualitatively similar results (see S2 Fig). We used *bw.method="global"* to allow a

constant smoothing parameter across all times. The defined time window was zero to 20 days

post injection. We removed one replicate (92 CFU for *E. cloacae* infection) because there was

no mortality in the first 20 days and therefore the maximum hazard could not be estimated.

This gave final sizes of n = 14 for *E. cloacae* and n = 15 for each of the other three species.

Model 1: log(maximum hazard) ~ bacterial species

2.7.2. Are virulence differences due to variation in parasite exploitation or PPP? To test whether the bacterial species vary in PPP, we performed a linear model with the natural log of the maximum hazard as the dependent variable, bacterial species as a factor, and the natural log of infection intensity as a covariate. We also included the interaction between bacterial species and infection intensity: a significant interaction would indicate variation in the reaction norms, *i.e.*, variation in PPP. The package "emmeans" [62] was used to test which of the reaction norms differed significantly from each other. We extracted maximum hazard values from time of death data for each bacterial species/dose/experimental replicate as described in 2.7.1. We also calculated the maximum hazard for the Ringer's control groups, which gives the maximum hazard in the absence of infection (the y-intercept). We present the results from $b = 2$, but all of the other values gave qualitatively similar results (see results). We wanted to infer the causal effect of bacterial load upon host survival (and not the reverse), therefore we reasoned that the bacterial load measures should derive from flies homogenised before the maximum hazard had been reached. For *E. cloacae*, *L. lactis*, and *Pr. burhodogranariea*, for all smoothing parameter values, the maximum hazard was reached after two days post injection, although for smoothing parameter value 1, there were four incidences where it was reached between 1.8- and 2-days post injection. Per species/dose/experimental replicate we therefore calculated the geometric mean of infection intensity combined for days 1 and 2 post injection. In order to include flies with zero load, we added one to all load values before calculating the geometric mean. This was done using the R packages "dplyr" [63], "plyr" [64] and "psych" [65]. Each mean was calculated from the bacterial load of eight flies, except for four mean values for *E. cloacae*, which derived from four flies each.

For *Ps. entomophila* the maximum hazard was consistently reached at around day one post injection, meaning that bacterial sampling happened at around the time of the maximum

hazard, and we therefore excluded this bacterial species from the analysis. We removed two

replicates (Ringer's and 92 CFU for *E. cloacae* infection) because there was no mortality in

the first 20 days and therefore the maximum hazard could not be estimated. One replicate was

removed because the maximum hazard occurred before day 1 for all *b* values (92,000 CFU for

E. cloacae) and six replicates were removed because there were no bacterial load data

available for day one (experimental replicate three of *L. lactis*). This gave final sample sizes

of n = 15 for *E. cloacae* and n = 12 for *L. lactis*, and n = 18 for *Pr. burhodogranariea*.

Model 2: log(maximum hazard) ~ log(geometric mean bacterial load) × bacterial

species

To test whether there is variation in parasite exploitation (infection intensity measured as

bacterial load), we performed a linear model with the natural log of infection intensity as the

dependent variable and bacterial species as a factor. Similar to the previous model, we used

the geometric mean of infection intensity combined for days 1 and 2 post injection, for each

bacterial species/dose/experimental replicate. The uninfected Ringer's replicates were not

included in this model. Post-hoc multiple comparisons were performed using "emmeans". *Ps.*

entomophila was excluded for the reason given above. The sample sizes per bacterial species

were: n = 13 for *E. cloacae*, n = 10 for *L. lactis* and n = 15 for *Pr. burhodogranariea*.

Model 3: log(geometric mean bacterial load) ~ bacterial species

2.7.3. Are persistent infection loads dose-dependent? We tested whether initial injection

dose is a predictor of bacterial load at seven days post injection [19, 21]. We removed all flies

that had 0 CFU as they are not informative for this analysis. The response variable was

natural log transformed bacterial load at seven days post-injection and the covariate was

natural log transformed injection dose. Separate models were carried out for each bacterial

species. Experimental replicate and person were fitted as fixed factors. By day seven none of

the flies injected with 92,000 CFU of *L. lactis* were alive. The analysis was not possible for

Ps. entomophila infected flies because all flies were dead by seven days post injection.

395   Model 4: $\log(\text{day 7 bacterial load}) \sim \log(\text{injection dose}) + \text{replicate} + \text{person}$

2.7.4. Calculation of clearance indices. To facilitate the analyses of clearance we calculated

clearance indices, which aggregate information about clearance into a single value for each

bacterial species/dose/experimental replicate. All indices were based on the estimated

proportion of cleared infections (defined as samples with measured zero bacterial load) of the

whole initial population. For this purpose, we first used data on bacterial load in living flies to

calculate the daily proportion of cleared infections in live flies for the days that we sampled.

Then we used the data on fly survival to calculate the daily proportion of flies that were still

alive. By multiplying the daily proportion of cleared flies in living flies with the proportion of

flies that were still alive, we obtained the proportion of cleared infections of the whole initial

population – for each day on which bacterial load was measured. We then used these data to

calculate three different clearance indices, which we used for different analyses. For each

index we calculated the mean clearance across several days. Specifically, the first index was

calculated across days one and two post injection (clearance index$_{1,2}$); the second index was

calculated across days three and four (clearance index$_{3,4}$); and the third index was calculated

from days seven, 14 and 21 (clearance index$_{7,14,21}$).

2.7.5. Does injection dose affect bacterial clearance? For this purpose, we aimed to assess

clearance that occurs shortly after injection. Accordingly, we used the clearance index that

was calculated for days one and two post injection. We suspected that the effect of dose on

clearance might differ between bacterial species. Therefore, we ran separate tests for each

species. The distribution of clearance values did not conform to the assumptions of linear

models, therefore we used Spearman rank correlations to test whether clearance changes with

injection dose.

420   Model 5: clearance index$_{1,2}$ ~ injection dose

2.7.6. Do the bacterial species differ in clearance? To test whether the bacterial species

differed in clearance, we used clearance index$_{3,4}$, which is the latest timeframe for which we

could calculate this index for all four species: due to the high virulence of *Ps. entomophila* we

were not able to assess bacterial load and thus clearance for later days. The distribution of

clearance values did not conform to the assumptions of a linear model. We therefore used a

Kruskal-Wallis test with pairwise Mann-Whitney-U post hoc tests. To control for multiple

testing we corrected the p-values of the post hoc tests using the method proposed by

Benjamini and Hochberg [66] that is implemented in the R function *pairwise.wilcox.test*.

430   Model 6: clearance index$_{3,4}$ ~ bacterial species

2.7.7. Do exploitation or PPP predict variation in clearance? To assess whether

exploitation or PPP predict variation in clearance we performed separate analyses for

clearance index$_{3,4}$ and clearance index$_{7,14,21}$. As discussed above, this precluded analysing *Ps.*

entomophila. Exploitation and PPP were calculated based on bacterial load from days 1 and 2

(see section 2.7.2), therefore we did not perform an analysis for clearance index$_{1,2}$. For each of

the two indices we fitted a linear mixed effects model with the clearance index as the response

variable. As fixed effects predictors we used the replicate-specific geometric mean log

bacterial load (see section 2.7.2) and the species-specific PPP (see section 3.2). In addition,

we included species as a random effect.

In our analysis we faced the challenge that many measured clearance values were at, or very

close to zero. In addition, clearance values below zero do not make conceptual sense. To

appropriately account for this issue, we used a logit link function (with Gaussian errors) in our

model, which restricts the predicted clearance values to an interval between zero and one.

Initial inspections of residuals indicated violations of the model assumption of homogenously

distributed errors. To account for this problem, we included the log bacterial load and PPP as

predictors of the error variance, which means that we used a model in which we relaxed the

standard assumption of homogenous errors and account for heterogenous errors by fitting a

function of how errors vary. For this purpose, we used the option *dispformula* when fitting the

models with the function *glmmTMB*.

Model 7: clearance index$_{3,4}$ or clearance index$_{7,14,21}$ ~ log(geometric mean bacterial

load) + PPP + bacterial species$_{random}$

2.7.8. Does longer-term clearance depend upon the injection dose? In contrast to the

analyses described above, we additionally aimed to assess the long-term dynamics of

clearance based on the infection status of dead flies collected between 14 and 35 days and 56

to 78 days after injection. Using binomial logistic regressions, we tested whether initial

injection dose affected the propensity for flies to clear an infection with *E. cloacae* or *Pr.*

burhodogranariea before they died. The response variable was binary whereby 0 denoted that

no CFUs grew from the homogenate and 1 denoted that CFUs did grow from the homogenate.

Natural log transformed injection dose was included as a covariate as well as its interaction

with day post injection, and person was fitted as a fixed factor. Replicate was included in the

Pr. burhodogranariea analysis only, because of unequal sampling across replicates for *E.*

cloacae. L. lactis injected flies were not analysed because only 4 out of 39 (10.3 %) cleared

the infection. *Ps. entomophila* infected flies were not statistically analysed because of a low

sample size (n = 12). The two bacterial species were analysed separately.

Model 8: CFU presence/absence$_{dead}$ ~ log(injection dose) × day post injection +

replicate + person

To test whether the patterns of clearance were similar for live and dead flies we tested

whether the proportions of dead and live uninfected flies correlated with each other (S1 File).

Results

3.1. Bacterial species vary in virulence

Fly survival after infection with five doses of the four different bacterial species is shown in

Fig 2A-D. As predicted, the bacterial species differed significantly in virulence, given as the

maximum hazard ($F_{3,55}$ = 193.05, p < 0.0001; Fig 2E). All species differed significantly from

each other (p < 0.0017 in all cases; S1 Table): *E. cloacae* was the least virulent and *Ps.*

entomophila the most virulent bacterium. *Pr. burhodogranariea* and *L. lactis* were

intermediate, with the former being less virulent than the latter. In all figures, the bacterial

species are thus presented in order of virulence.

3.2. Differences in virulence are due to variation in parasite exploitation and PPP

We assessed infection intensity over time post injection (Fig 3) and used the geometric mean

of the values for the first two days post injection, as a proxy for parasite exploitation (see

methods for rationale). Bacterial species varied significantly in exploitation of their hosts

during this early infection phase ($F_{2,35}$ = 35.90; p < 0.0001; Fig 4A). The least virulent

bacterium, *E. cloacae*, had a significantly lower infection intensity, and thereby lower parasite

exploitation, compared to either of the other species (Tukey contrasts: *E. cloacae vs. Pr.*

burhodogranariea: t = -5.24, p < 0.0001; *E. cloacae vs. L. lactis*: t = -8.36, p < 0.0001). The

more virulent bacterium, *L. lactis*, had the highest infection intensity, and differed

significantly compared to the less virulent *Pr. burhodogranariea* (*L. lactis vs. Pr.*

burhodogranariea: t = 3.50, p = 0.0018).

The slopes of the relationship between infection intensity and maximum hazard differed

significantly across bacterial species, suggesting that the bacterial species differed in their

PPP (infection intensity × bacterial species: $F_{2,39} = 7.35$, p = 0.0020; Fig 4B). *E. cloacae* had a

relatively flat reaction norm, indicating a minimal increase in hazard with an increase in

bacterial load, and thus a significantly lower PPP compared to both *Pr. burhodogranariea*

(Tukey contrast: t = -3.74; p = 0.0017) and *L. lactis* (t = -3.34; p = 0.0052). In contrast, the

latter two species had similar PPP to each other (t = -0.68; p = 0.78); both species had

negative reaction norms, indicating an increase in hazard with an increase in bacterial load.

There was no significant effect of bacterial load ($F_{1,39} = 0.19$, p = 0.67) or bacterial species

$F_{2,39} = 0.50$, p = 0.61) on the maximum hazard. Qualitatively similar results were obtained

using the three alternative smoothing parameters (S2 Fig). *Ps. entomophila* was not included

in the above analyses because the maximum hazard was consistently reached at around day

one post injection, meaning that we could not infer the causal effect of bacterial load upon

survival.

3.3. All bacterial species established persistent infections

By homogenising living flies, we found that the two bacterial species with lower virulence, *E.*

cloacae and *Pr. burhodogranariea* were able to persist inside the fly until at least 35 days

post injection (Figs 2A and 2B respectively). The persistence estimates for *L. lactis* (28 days;

Fig 3C) and *Ps. entomophila* (four days; Fig 3D) were both shorter, because the high

mortality caused by these bacterial species meant that we could not test later time points.

However, by testing for the presence or absence of bacteria in homogenised dead flies, we

found that infections could persist for considerably longer, *i.e.,* around two and a half months:

E. cloacae = 77 days, *Pr. burhodogranariea* = 78 days, *L. lactis* = 76 days and *Ps.*

entomophila = 75 days (Fig 5A-D).

3.4. Injection dose correlates with persistent infection loads

At seven days post injection, *E. cloacae* (Fig 6A) and *Pr. burhodogranariea* (Fig 6B) loads were significantly positively correlated with the initial injection dose (Table 1). *L. lactis* loads showed no significant correlation with the initial injection dose (Fig 6C; Table 1). We hypothesised that the lack of significant relationship between *L. lactis* load and injection dose might be due to our underestimation of the load of some flies injected with this bacterial species: this is because some flies had too many CFUs to count even at the lowest dilution, and they were therefore assigned a maximum bacterial load value (see methods), which was necessarily lower than their actual load. When we excluded the two flies at day seven that had been assigned the maximum value, the relationship became statistically significant (log(injection dose): $F_{1,35} = 4.59$; $p = 0.039$).

Table 1. The effect of initial injection dose on bacterial load at seven days post injection (Model 4). Experimental replicate and the person performing the injection were also included as factors in the models. *Ps. entomophila* was not analysed because it caused high fly mortality. Statistically significant factors are in bold.

Injected bacterium	Tested effect	df	F	P
E. cloacae	Log(Injection dose)	1,25	26.41	**<0.0001**
	Person	1,25	0.16	0.69
	Replicate	2,25	1.78	0.19
Pr. burhodogranariea	Log(Injection dose)	1,47	37.33	**<0.0001**
	Person	1,47	0.23	0.63
	Replicate	2,47	2.11	0.13
L. lactis	Log(Injection dose)	1,37	3.81	0.058
	Person	1,37	0.71	0.40

	Replicate	2,37	1.98	0.15

3.5. Lower doses of E. cloacae are cleared more quickly than higher doses

Summing up across all doses and days, 39.4 % (177 of 449) of *E. cloacae*-injected flies, 11.8 % (45 of 381) of *Pr. burhodogranariea*-injected flies, 3.7 % (11 of 301) of *L. lactis*-injected flies, and 21.4 % (15 of 70) of *Ps. entomophila*-injected flies cleared the infections (Fig 7). To account for mortality in our estimates of bacterial clearance we calculated clearance indices. Using the clearance index for the early infection phase, i.e., days one and two post injection (clearance index$_{1,2}$), we found that flies injected with lower doses of *E. cloacae* were more likely to clear the infection compared to flies injected with higher doses (Spearman rank correlation: $\rho = -0.86$, $p < 0.001$, Fig 8A). However, the other three bacterial species did not show dose-dependent clearance (Fig 8B-D).

3.6. Bacterial species vary in clearance

The four bacterial species used in this study covered a broad spectrum of clearance on days three and four post injection (Fig 8E). There was a statistically significant difference among species ($p = 0.002$, Chisq = 15.309, df = 3), and after p-value correction, the post hoc tests indicated statistically significant differences among the following species pairs: *E. cloacae* and *Pr. burhodogranariea* ($p = 0.024$), *E. cloacae* and *L. lactis* ($p = 0.011$), *Ps. entomophila* and *Pr. burhodogranariea* ($p = 0.048$), *Ps. entomophila* and *L. lactis* ($p = 0.011$). Rather than matching the virulence gradient across species (Fig 2E), clearance formed a U-shaped pattern with the species with the highest virulence (*Ps. entomophila*) and lowest virulence (*E. cloacae*) showing higher levels of clearance compared to the two species of intermediate virulence (*Pr. burhodogranariea* and *L. lactis*).

3.7. Exploitation but not PPP predict clearance

Our analyses of clearance index$_{3,4}$ and clearance index$_{7,14,21}$ showed similar results. In both

cases we found no statistically significant effect of PPP, but a significant negative effect of

exploitation, such that as the mean bacterial load for days one and two increased, clearance

decreased (Fig 8F, G Table 2). Similar results were obtained using the three alternative

smoothing parameters for calculating PPP (S3 Fig).

Table 2. The effect of log bacterial load (exploitation) and PPP on two different clearance

indices. Statistically significant factors are in bold.

Response variable	Tested effect	df	Chisq	P
Clearance index$_{3,4}$	Log (geometric mean bacterial load)	1	14.16	**< 0.001**
	PPP	1	0.38	0.535
Clearance index$_{7,14,21}$	Log (geometric mean bacterial load)	1	36.34	**< 0.001**
	PPP	1	0.35	0.557

3.8. Bacterial clearance before death is dose dependent

We homogenised flies that died during the chronic phase of the infection (between 14 and 35

days and between 56- and 78-days post injection) to test whether they died whilst being

infected, or whether they were able to clear the infection before death. Flies were indeed able

to clear the infection before death, but the degree to which this occurred varied across

bacterial species (Fig 5). Furthermore, for all bacterial species in both homogenisation phases

there were flies where the infection persisted until death, and flies that were uninfected at

death (Fig 5A-D). Lower injection doses of *E. cloacae* were more likely to be cleared before

death than higher injection doses (Fig 5E; Table 3), but there was no significant effect for *Pr.*

burhodogranariea (p = 0.051; Fig 5F; Table 3). Summing up across all doses and days, 29.8

% (14 out of 47) of *E. cloacae*-injected flies, 33.3 % (20 out of 60) of *Pr. burhodogranariea*-

injected flies, 10.3 % (4 out of 39) of *L. lactis*-injected flies, and 66.7 % (8 out of 12) of *Ps.*

entomophila-injected flies cleared the infection before death.

Table 3. The effect of injection dose on presence/absence of infection in dead flies (Model 8).

Person performing the injection was also included as a factor in the models, and replicate was

included for the analysis for *Pr. burhodogranariea* infections. Statistically significant factors

are in bold.

Injected bacterium	Tested effect	df	LR Chisq	P
E. cloacae	Log(Injection dose)	1	3.82	0.051
	Day post injection	1	0.0037	0.95
	Person	1	1.22	0.27
	Log(Injection dose) × Day post injection	1	0.013	0.91
Pr. burhodogranariea	Log(Injection dose)	1	13.26	**0.00027**
	Day post injection	1	2.99	0.084
	Person	1	4.33	**0.037**
	Replicate	1	0.038	0.54
	Log(Injection dose) × Day post injection	1	1.75	0.19

Discussion

In this study we demonstrate that sustained persistent infection and clearance are both

possible outcomes for bacteria showing a range of virulence when they infect female *D.*

melanogaster. We show that bacteria of all species can persist inside the host for at least 75

days and that bacterial virulence differences can be explained by a combination of variation in

exploitation and PPP. We also show that lower doses of the bacterium with the lowest

virulence, *E. cloacae*, are cleared more quickly than higher doses, and that clearance rates are

species specific. Furthermore, higher exploitation of the host leads to lower bacterial

clearance. Finally, we found some support for our hypothesis that exploitation and PPP imply

different costs and benefits of clearance and thus both virulence components have different

effects on clearance.

4.1. Differences in virulence are due to variation in exploitation and PPP

The infecting bacterial species showed pronounced differences in virulence. To understand

why this was the case, we decomposed virulence into its two components: exploitation and

PPP [28, 29]. Exploitation, given as infection intensity or bacterial load, is the more

frequently tested explanation for variation in virulence [28]. There is ample evidence that

exploitation varies across parasite genotypes [67, e.g., monarch butterflies and their protozoan

parasites: 68, *Daphnia magna* infected with the bacterium *Pasteuria ramosa*: 69], and also,

unsurprisingly, that it varies across parasite species infecting the same host genotype [19-21].

Indeed, in the current study, all bacterial species tested showed significant differences in

exploitation, where bacterial load increased as virulence increased. Chambers et al. [19]

observed that the two bacterial species in their study that caused lower mortality showed little

initial proliferation inside the host, but that the species causing more mortality showed an

initial increase in the bacterial load: these results support our findings.

However, variation in virulence might not only be determined by the load that a pathogen

attains: Råberg & Stjernman [28] proposed that pathogen genotypes may also vary in PPP *i.e.*,

the harm or damage caused per parasite [28, 29]. Some pathogens may cause more damage to

the host independently of their density, through mechanisms that directly affect the host

homeostasis. For example, *Bacillus anthracis* produces two anthrax toxins which are

responsible for impairing the host immune system and disrupting basic cellular functions,

ultimately killing the host [70]. By comparison, the genetically similar *Bacillus cereus* usually

only produces a mildly virulent gastro-intestinal infection [71-73]. Variation in PPP can be

observed when different parasite genotypes show different reaction norms for the relationship

625 between host health and infection intensity, when infecting the same host genotype [29].

626 Variation in PPP has been demonstrated for rats infected with different clones of *Plasmodium*

627 *chabaudii,* the agent of rodent malaria [28], for different strains of protozoan parasites

628 infecting monarch butterflies [67], and for humans infected with different HIV-1 genotypes

629 [74]. Here we found a significant overall effect of PPP across bacterial species, whereby *Pr.*

630 *burhodogranariea* and *L. lactis* had significantly more negative slopes compared to *E.*

631 *cloacae*. This finding, combined with the exploitation results, implies that *E. cloacae* is less

632 virulent towards its host compared to the other two species, because of a combination of

633 lower PPP and less exploitation. On the other hand, given that *Pr. burhodogranariea* and *L.*

634 *lactis* both showed similar levels of PPP, it suggests that the variation in virulence between

635 these two species is due to higher exploitation by *L. lactis*, rather than differences in PPP.

636 Some of our *L. lactis* counts were underestimates because there were too many CFUs to

637 count, so it is possible that the slope of the reaction norm might have differed slightly had we

638 not encountered this issue. Nonetheless, had we only examined exploitation as a source of

639 variation, we might have concluded that load alone explains the differences that we found in

640 virulence. These two sources of variation, exploitation and PPP, have not frequently been

641 explored in the same study, so it is generally difficult to ascertain the relative importance of

642 the two sources of variation. However, variation in PPP was demonstrated to explain more of

643 the variance in virulence across HIV-1 genotypes than did set point viral load [74].

645 **4.2. All bacterial species established persistent infections**

646 All four bacterial species were able to establish persistent infections in *D. melanogaster*. *E.*

647 *cloacae* and *Pr. burhodogranariea* could be retrieved from live homogenised flies up to 35

648 days, *L. lactis* up to 28 days, and *Ps. entomophila* up to four days post injection. The reduced

649 estimates for the latter two species are due to higher mortality, meaning that no flies were

650 alive to sample at later time points. However, by homogenising flies that had died, we show

that all bacterial species can persist inside the host for at least 75 days. To the best of our

knowledge these estimates are far beyond the currently known length of persistent infections

after injection in insects [28 days: 17, 18]. The duration of infection can be of key ecological

and potentially also evolutionary importance, because persistence determines the prevalence

of infection in a population, and therefore could affect transmission.

It is unclear how these bacteria are able to persist for so long inside the host, although there

are a number of theoretical possibilities, for example, through surviving inside host tissue,

forming biofilms or existing as persister or tolerant cells [75]. *Salmonella typhimurium* [76]

and *S. aureaus* [77] can survive inside insect haemocytes. The bacterial species that we used

are not known to be intracellularly replicating, *e.g.*, some *Providencia* strains were able to

survive, but not replicate, at low numbers 24 hours after infecting a *D. melanogaster* S2 cell

line [45]. Both *E. cloacae* and *L. lactis* are able to produce biofilms *in vitro* [78, 79,

respectively], but it is unknown whether this is the case inside an insect host. *Pr.*

burhodogranariea is not able to form biofilms *in vitro* [45] although it is unknown if it might

still be possible *in vivo*. Biofilms can cause chronic infections such as *Pseudomonas*

aeruginosa in cystic fibrosis patients [80], and oral infection of *D. melanogaster* with *Ps.*

aeruginosa resulted in biofilm production in the crop [81], but it is unknown whether *Ps.*

entomophila forms biofilms *in vivo*. Lastly, bacteria could potentially survive inside the host

in persistent or tolerant cell states, as is discussed in relation to the failure of antibiotic

treatments [82]. However, persistent cells typically make up a small proportion (< 1%) of the

bacterial population [82], so this might not explain the high numbers of CFUs that are

retrieved from the flies. Future research will test the likelihood of these and other

mechanisms.

D. melanogaster that are able to control a bacterial infection during the acute infection phase

have been shown to have a relatively constant bacterial load in the chronic infection phase,

which Duneau *et al.* [21] found remains stable until at least ten days post injection for *Pr.*

rettgeri. Our bacterial load data (Fig 3) suggests that *E. cloacae*, *Pr. burhodogranariea* and *L.*

lactis, show relatively stable loads from around day three to four post infection, lending

support to the SPBL concept. In addition, Duneau *et al.* [21] observed that per host, bacteria

with low virulence had a SPBL of a few hundred bacteria, whereas bacteria of intermediate

virulence had a SPBL of a few thousand bacteria. Our data also lend support to the idea that

virulence relates to SPBL, given that low virulence *E. cloacae* had a persistent load of tens to

hundreds of bacteria, and high virulence *L. lactis* had a load of tens of thousands of bacteria.

This finding is supported by the virulence decomposition analysis (see section 4.1), which

shows that as virulence increases, so does exploitation of the host over the first couple of days

post infection. Therefore, more virulent bacteria have higher initial proliferation rates as

shown by exploitation. Given that the infection load stays relatively constant in the longer

term, the initial proliferation differences likely explain the relationship between SPBL and

virulence.

4.3. Injection dose correlates with persistent infection loads

The bacterial load at day seven post injection, positively correlated with the initial injection

dose for *E. cloacae, Pr. burhodogranariea* and *L. lactis* (but see results section for the latter).

Our results expand the known bacterial species for which this relationship exists, and they

lend weight to the idea that this may be a more general phenomenon in *D. melanogaster*

bacterial infections. Previous studies found that this relationship held for bacterial load at

seven- and fourteen-days post injection [E. faecalis, Pr. rettgeri and S. marcescens: 19, Pr.

rettgeri: 21;]. It has been suggested that the SPBL will remain at around the bacterial load at

which the infection was controlled [19, 21]. Given that insects can show dose dependent

inducible immune activation [34], and given that the antimicrobial peptide Drosocin has been

shown to control *E. cloacae* infections and that a combination of Drosocin, Attacins and

Diptericins control *Pr. burhodogranariea* infections [47], one could hypothesise that these

AMPs are to some degree involved. However, the mechanisms that allow a dose-dependent

persistent infection, remain to be uncovered. It was not possible to test *Ps. entomophila* given

its high mortality during the acute infection phase.

709   4.4. Lower doses of E. cloacae are cleared more quickly than higher doses

The likelihood of clearing *E. cloacae* was dose dependent, although we did not find a dose

threshold below which there was complete clearance in all flies. The finding of dose-

dependent clearance, whilst maybe not surprising, could explain some discrepancies across

studies in terms of whether evidence of persistent infections is found. Just as stochastic

variation explains variation in the outcome of the early infection phase [21], perhaps

stochasticity plays a role in the clearance of bacteria [83], particularly where infection loads

are low such as in *E. cloacae*, for example through variation in expression of Drosocin. In

comparison to *E. cloacae*, most replicates of the other species showed no clearance in the

early infection phase, i.e., one- and two-days post infection, thus dose did not influence

clearance for these species.

721   4.5. Bacterial species vary in clearance

Across species, we uncovered a U-shaped pattern in bacterial clearance, where the species

with the lowest and highest virulence had higher levels of clearance compared to the two

species of intermediate virulence. Our finding that the low virulent species could be cleared,

is supported by evidence from infections with low virulence *E. coli* and *Erwinia carotovora*

Ecc15, where an injection dose of 30,000 bacteria was cleared in 22 % and 8 % of flies,

respectively [21]. The clearance of intermediate and high virulence pathogens in *D.*

melanogaster has been described as being rare, because no bacteria were cleared from any of

the previously infected hosts over the seven-days post injection [21]. *Pr. rettgeri* and

Enterococcus faecalis were described as having intermediate virulence in that study, with a

survival of around 50-60% seven days post infection [21], which is in the region of the

survival that we found for our intermediate virulence species, *Pr. burhodogranariea* and *L.*

lactis. Although clearance in our intermediately virulent bacteria was lower than that of the

low and high virulent bacteria, our results challenge the finding that clearance is rare because

our three more virulent bacteria all appear to be clearable to differing degrees, including

within the first seven days of infection. Our results thereby show that persistent infections are

not inevitable. This finding is supported by the observation that *Pr. burhodogranariea* was

cleared in flies seven- to ten-days post injection [45; although low sample sizes]. Similarly to

the current study, Kutzer & Armitage [20] also found that a few female flies, inoculated with

a dose of *L. lactis* in common with this study (1,840 CFUs), cleared the infection (3 out of

141; 2.1 %). Lastly, we expected that there may be selection for a fast and efficient early

clearance of infection by *Ps. entomophila*, because of its high virulence. Clearance of *Ps.*

entomophila was indeed higher than for the intermediate bacteria, although mortality was too

high to assess clearance in living flies for longer than four days post injection. Nonetheless,

there is evidence from other studies that *Ps. entomophila* has high virulence and can be

cleared from other *D. melanogaster* populations/genotypes [18, 39, 46]. The ability to resist

Ps. entomophila infection varied across host genotypes; similarly, five out of the ten tested

genotypes contained some individuals who cleared this pathogen [46]. Host genotypic

variation in clearance may thereby more generally explain why some studies find clearance

and others do not.

Even though dead flies were sampled for a longer period post-injection (up to 78 days)

compared to live flies (up to 35 days), the patterns of bacterial clearance in dead flies largely

reflected the results for live flies, given that dead flies that had been infected with *E. cloacae*

showed dose dependency in clearance. Once again, comparatively few dead individuals had

cleared *L. lactis* infections, whereas proportionally more had cleared *Ps. entomophila.*

Because we processed dead flies up to 24 hours post-death we did not analyse the number of

CFUs, however it would be interesting to test whether the bacterial load upon death [21]

remains constant even after many weeks of a potentially costly infection.

4.6. Exploitation but not PPP predicts clearance

We next sought to understand how the two components that determine variation in virulence

affect clearance of the infection, whilst framing our argumentation in the context of the

benefits to costs ratio to the host of clearing an infection. Our results provide some support for

our hypothesis that exploitation and PPP can affect clearance in different ways, which could

result in different, and potentially even opposing, patterns of how variation in virulence is

related to clearance. As predicted, we found that across species, increased host exploitation

early in the infection, i.e., days one and two post infection, was associated with decreased

clearance at days three and four post infection, and also at days seven to 21. This finding is

consistent with the expectation that the host immune response is sensitive to clearance costs

associated with exploitation. We did not find support for our prediction that increasing PPP

would be associated with increased clearance, which would be expected if the host immune

response is sensitive to clearance benefits. Taken together, our results demonstrate that for the

investigated pathogens, variation in clearance is mainly explained by variation in clearance

costs whereas clearance benefits seem to have no or only little effect. Further studies are

necessary to assess whether this is a general pattern.

Our finding of a negative relationships between exploitation and clearance is in contrast to

data from vertebrate viral infections [reviewed in 32; , e.g., 33], where larger viral loads led to

a faster decline in viral load or in shorter durations of viremia. Although these studies did not

directly assess clearance as defined in our study, their results are consistent with the idea that

in these systems increasing exploitation might lead to a stronger increase in the benefits of

clearance compared to the costs of clearance. In this case one would expect a positive

relationship between exploitation and clearance.

In contrast to exploitation, by definition, PPP cannot vary within replicates of a species,

because each species has just one PPP slope. In addition, only three species could be included

in this analysis due to high mortality after injection with *Ps. entomophila*. Therefore, we note

that the statistical power of detecting an effect of PPP on clearance might have been lower

than the power to detect an effect of exploitation. Nonetheless, it is possible that analyses with

more species will reveal an effect of PPP on clearance. Based on existing information on the

biology of *Ps. entomophila* we propose that the increased levels of clearance for this species

(Fig 8E) might have been caused by particularly high levels of PPP. *Ps. entomophila* produces

the virulence factor AprA [84], and a pore-forming toxin called Monalysin [85] in association

with the activation of stress-induced pathways and an increase in oxidative stress [86].

Ultimately this leads to a lack of tissue repair in the gut, and in most cases fly death [reviewed

in 87]. If similar pathologies are induced in the haemocoel after infection, in contrast to other

bacterial species, sustaining a persistent bacterial load in the face of high levels of tissue

damage might rarely be a viable option. Instead, the fly host might activate a stronger immune

response to attempt to clear the infection [31, 88]. However, a few flies in ours and other

studies, did survive into the chronic phase whilst testing positive for a *Ps. entomophila*

infection. In *D. melanogaster* Drosocrystallin expression in the gut has been shown to confer

protection to Monalysin and lead to higher individual survival [89, 90]; perhaps individuals

surviving into the chronic infection phase exhibit higher expression of such mechanisms,

allowing them to tolerate the damage caused by infection. Nonetheless, we suggest that the

higher virulence of *Ps. entomophila* could have been driven by an increased PPP because of

the above-mentioned toxins, but additional experiments would be required to test this idea.

4.7. Conclusions

To conclude, our results strongly support that PPP is an important component driving

variation in virulence, and that disentangling its contribution towards virulence, in

combination with the contribution of exploitation, will undoubtedly help our mechanistic and

evolutionary understanding of host-pathogen interactions. We suggest that such a

decomposition of virulence can be used to better understand how virulence relates to other

infection processes such as clearance during persistent infections. Future research will be

needed to test the generality of the relationships we have uncovered between virulence

decomposed as exploitation and PPP, and the persistence and clearance of infections. Here we

have taken a pathogen perspective on virulence, however a more complete understanding of

how virulence relates to clearance will be achieved by simultaneously considering the host

factors that contribute to variation in virulence, i.e., resistance and tolerance. Such an

approach may contribute towards expanding the theory on virulence evolution.

Author contributions

SAOA conceived the overall idea. BAH, LS and SAOA designed the experiments and

collected the data. BAH, LS, MF and SAOA wrote the manuscript. MF & RRR conceived the

virulence decomposition and clearance analyses and MF, RRR & SAOA analysed the data.

All authors contributed critically to the drafts.

Acknowledgements

830 We thank the Hiesinger group, the Rolff group, Luisa Linke, Alexandro Rodríguez-Rojas,

831 Jens Rolff and Seulkee Yang for advice and/or technical support. We thank Jens Rolff for

832 feedback on an earlier draft of the manuscript.

Data Accessibility

835 Data will be made publicly available upon acceptance of this article.

References

838 1. Best A, White A, Boots M. Maintenance of host variation in tolerance to pathogens and
839   parasites. Proceedings of the National Academy of Sciences. 2008;105:20786-91. doi:
840   10.1073/pnas.0809558105.
841 2. Råberg L, Graham AL, Read AF. Decomposing health: tolerance and resistance to
842   parasites in animals. Phil Trans R Soc B. 2009;364:37-49. doi: 10.1098/rstb.2008.0184.
843 3. Schmid-Hempel P. Evolutionary Parasitology: The Integrated Study of Infections,
844   Immunology, Ecology, and Genetics: Oxford University Press; 2011.
845 4. Boots M, Begon M. Trade-offs with resistance to a granulosis virus in the indian meal
846   moth, examined by a laboratory evolution experiment. Functional Ecology. 1993;7:528-
847   34. doi: 10.2307/2390128.
848 5. Kraaijeveld AR, Godfray HCJ. Trade-off between parasitoid resistance and larval
849   competitive ability in *Drosophila melanogaster*. Nature. 1997;389:278-80. doi:
850   10.1038/38483.
851 6. Armitage SAO, Thompson JJW, Rolff J, Siva-Jothy MT. Examining costs of induced and
852   constitutive immune investment in *Tenebrio molitor*. Journal of Evolutionary Biology.
853   2003;16:1038-44. doi: 10.1046/j.1420-9101.2003.00551.x.
854 7. Schmid-Hempel P. Variation in immune defence as a question of evolutionary ecology.
855   Proc R Soc Lond B. 2003;270:357-66. doi: 10.1098/rspb.2002.2265.
856 8. Nystrand M, Dowling DK. Effects of immune challenge on expression of life-history and
857   immune trait expression in sexually reproducing metazoans-a meta-analysis. BMC
858   Biology. 2020;18. Epub 2020/10/09. doi: 10.1186/s12915-020-00856-7. PubMed PMID:
859   33028304; PubMed Central PMCID: PMCPMC7541220.
860 9. Bajgar A, Kucerova K, Jonatova L, Tomcala A, Schneedorferova I, Okrouhlik J, et al.
861   Extracellular adenosine mediates a systemic metabolic switch during immune response.
862   PLoS Biol. 2015;13.
863 10. Graham AL, Allen JE, Read AF. Evolutionary causes and consequences of
864   immunopathology. Annu Rev Ecol Evol Syst. 2005;36:373-97.
865 11. Sadd BM, Siva-Jothy MT. Self-harm caused by an insect's innate immunity. Proc R Soc
866   B. 2006;273:2571-4.
867 12. Khan I, Agashe D, Rolff J. Early-life inflammation, immune response and ageing. Proc R
868   Soc B. 2017;284.
869 13. Grant SS, Hung D, T. Persistent bacterial infections, antibiotic tolerance, and the
870   oxidative stress response. Virulence. 2013;4:273-83.

14. Gorman MJ, Paskewitz SM. Persistence of infection in mosquitoes injected with bacteria.
Journal of Invertebrate Pathology. 2000;75:296-7.
15. Boman HG, Ingrid N, Bertil R. Inducible antibacterial defence system in *Drosophila*.
Nature. 1972;237:232-5.
16. Hotson AG, Schneider DS. *Drosophila melanogaster* natural variation affects growth
dynamics of infecting *Listeria monocytogenes*. G3. 2015;5:2593-600. doi:
10.1534/g3.115.022558.
17. Haine ER, Moret Y, Siva-Jothy MT, Rolff J. Antimicrobial defense and persistent
infection in insects. Science. 2008;322:1257-9. doi: 10.1126/science.1165265.
18. Kutzer MAM, Kurtz J, Armitage SAO. A multi‑faceted approach testing the effects of
previous bacterial exposure on resistance and tolerance. J Anim Ecol. 2019;88:566-78.
doi: 10.1111/1365-2656.12953.
19. Chambers MC, Jacobson E, Khalil S, Lazzaro BP. Consequences of chronic bacterial
infection in *Drosophila melanogaster*. PLoS ONE. 2019;14(10):e0224440. doi:
10.1371/journal.pone.0224440.
20. Kutzer MAM, Armitage SAO. The effect of diet and time after bacterial infection on
fecundity, resistance, and tolerance in *Drosophila melanogaster*. Ecol Evol.
2016;6(13):4229-42. doi: 10.1002/ece3.2185.
21. Duneau D, Ferdy J-B, Revah J, Kondolf H, Ortiz GA, Lazzaro BP, et al. Stochastic
variation in the initial phase of bacterial infection predicts the probability of survival in
D. melanogaster. eLife. 2017;6:e28298. doi: 10.7554/eLife.28298.
22. Dionne MS, Pham LN, Shirasu-Hiza MM, Schneider DS. *Akt* and *foxo* dysregulation
contribute to infection-induced wasting in *Drosophila*. Current Biology. 2006;16:1977-
85.
23. Brandt SM, Dionne MS, Khush RS, Pham LN, Vigdal TJ, Schneider DS. Secreted
bacterial effectors and host-produced eiger/TNF drive death in a *Salmonella*-infected fruit
fly. PLoS Biol. 2004;2.
24. Regoes RR, McLaren PJ, Battegay M, Bernasconi E, Calmy A, F. GH, et al.
Disentangling human tolerance and resistance against HIV. PLoS Biol. 2014;12.
25. Zanchi C, Johnston PR, Rolff J. Evolution of defence cocktails: Antimicrobial peptide
combinations reduce mortality and persistent infection. Mol Ecol. 2017;26:5334-43. doi:
10.1111/mec.14267.
26. Miller CVL, Cotter SC. Resistance and tolerance: The role of nutrients on pathogen
dynamics and infection outcomes in an insect host. J Anim Ecol. 2017;87:500-10.
27. Read AF. The evolution of virulence. Trends in Microbiology. 1994;2(3):73-6. doi:
10.1016/0966-842X(94)90537-1.
28. Råberg L, Stjernman M. The evolutionary ecology of infectious disease virulence. In:
Demas G, Nelson R, editors. Ecological Immunology: Oxford University Press; 2012. p.
548-78.
29. Råberg L. How to live with the enemy: Understanding tolerance to parasites. PLoS Biol.
2014;12.
30. Frank SA, Schmid-Hempel P. Mechanisms of pathogenesis and the evolution of parasite
virulence. J Evol Biol. 2008;21(2):396-404. Epub 2008/01/09. doi: 10.1111/j.1420-
9101.2007.01480.x. PubMed PMID: 18179516.
31. Lazzaro BP, Rolff J. Danger, microbes, and homeostasis. Science. 2011;332:43-4.
32. Althouse BM, Durbin AP, Hanley KA, Halstead SB, Weaver SC, Cummings DA. Viral
kinetics of primary dengue virus infection in non-human primates: a systematic review
and individual pooled analysis. Virology. 2014;452-453:237-46. Epub 2014/03/13. doi:
10.1016/j.virol.2014.01.015. PubMed PMID: 24606701; PubMed Central PMCID:
PMCPMC4578724.

33. Ben-Shachar R, Koelle K. Transmission-clearance trade-offs indicate that dengue
virulence evolution depends on epidemiological context. Nat Commun. 2018;9(1):2355.
Epub 2018/06/17. doi: 10.1038/s41467-018-04595-w. PubMed PMID: 29907741;
PubMed Central PMCID: PMCPMC6003961.

34. Jent D, Perry A, Critchlow J, Tate AT. Natural variation in the contribution of microbial
density to inducible immune dynamics. Mol Ecol. 2019;28:5360-72. doi:
10.1111/mec.15293.

35. Louie A, Song KH, Hotson A, Tate AT, Schneider DS. How many parameters does it
take to describe disease tolerance? PLoS Biol. 2016;14(4):e1002435. doi:
10.1371/journal.pbio.1002435.

36. Brook I. Inoculum effect. Rev Infect Dis. 1989;11(3):361-8. doi:
10.1093/clinids/11.3.361. PubMed PMID: 2664999.

37. Chin J, Rybak M, Cheung C, Savage P. Antimicrobial activities of ceragenins against
clinical isolates of resistant *Staphylococcus aureus*. Antimicrobial Agents and
Chemotherapy. 2007;51(4):1268-73. doi: 10.1128/AAC.01325-06. PubMed PMID:
WOS:000245416500021.

38. Savini F, Luca V, Bocedi A, Massoud R, Park Y, Mangoni ML, et al. Cell-density
dependence of host-defense peptide activity and selectivity in the presence of host cells.
ACS Chem Biol. 2017;12(1):52-6. Epub 2016/12/09. doi: 10.1021/acschembio.6b00910.
PubMed PMID: 27935673.

39. Martins NE, Faria VG, Teixeira L, Magalhães S, Sucena É. Host adaptation is contingent
upon the infection route taken by pathogens. PLoS Pathog. 2013;9(9):e1003601. doi:
10.1371/journal.ppat.1003601.

40. Bass TM, Grandison RC, Wong R, Martinez P, Partridge L, Piper MDW. Optimization of
dietary restriction protocols in *Drosophila*. The Journals of Gerontology: Series A.
2007;62:1071-81. doi: 10.1093/gerona/62.10.1071.

41. Lazzaro BP. A population and quantitative genetic analysis of the *Drosophila*
melanogaster antibacterial immune response: The Pennsylvania State University; 2002.

42. Juneja P, Lazzaro BP. *Providencia sneebia* sp. nov. and *Providencia burhodogranariea*
sp. nov., isolated from wild *Drosophila melanogaster*. INTERNATIONAL JOURNAL
OF SYSTEMATIC AND EVOLUTIONARY MICROBIOLOGY. 2009;59:1108-11. doi:
10.1099/ijs.0.000117-0.

43. Vodovar N, Vinals M, Liehl P, Basset A, Degrouard J, Spellman P, et al. *Drosophila* host
defense after oral infection by an entomopathogenic *Pseudomonas* species. Proceedings
of the National Academy of Sciences. 2005;102(32):11414-9. doi:
10.1073/pnas.0502240102.

44. Cox CR, Gilmore MS. Native microbial colonization of *Drosophila* melanogaster and its
use as a model of *Enterococcus faecalis* pathogenesis. IAI. 2007;75(4):1565-76. doi:
10.1128/IAI.01496-06.

45. Galac MR, Lazzaro BP. Comparative pathology of bacteria in the genus *Providencia* to a
natural host, *Drosophila melanogaster*. Microbes and Infection. 2011;13:673-83.

46. Kutzer MAM, Kurtz J, Armitage SAO. Genotype and diet affect resistance, survival, and
fecundity but not fecundity tolerance. Journal of Evolutionary Biology. 2018;31:159-71.
doi: 10.1111/jeb.13211.

47. Hanson MA, Dostálová A, Ceroni C, Poidevin M, Kondo S, Lemaitre B. Synergy and
remarkable specificity of antimicrobial peptides in vivo using a systematic knockout
approach. eLife. 2019;8:e44341. doi: 10.7554/eLife.44341.

48. Werner T, Liu G, Kang D, Ekengren S, Steiner H, Hultmark D. A family of
peptidoglycan recognition proteins in the fruit fly *Drosophila melanogaster*. Proceedings
of the National Academy of Sciences. 2000;97:13772–7.

49. Lazzaro BP, Sackton TB, Clark AG. Genetic variation in *Drosophila melanogaster*
resistance to infection: A comparison across bacteria. Genetics. 2006;174:1539-54. doi:
10.1534/genetics.105.054593.
50. R Core Team. R: A language and environment for statistical computing. R Foundation
for Statistical Computing. 2020.
51. Kassambara A. ggpubr: 'ggplot2' based publication ready plots. 2020.
52. Baptiste A. gridExtra: Miscellaneous functions for "Grid" graphics. 2017.
53. Wickham H. ggplot2: Elegant graphics for data analysis. New York: Springer-Verlag
2016.
54. Wickham H, Seidel D. scales: Scale functions for visualization. 2020.
55. Therneau TM. A package for survival analysis in R. 2020.
56. Therneau TM, Grambsch PM. Modeling survival data: Extending the Cox model. New
York: Springer; 2000.
57. Garnier S. viridis: default color maps from 'matplotlib'. 2018.
58. Bolker BM, Brooks ME, Clark CJ, Geange SW, Poulsen JR, Stevens MHH, et al.
Generalized linear mixed models: a practical guide for ecology and evolution. Trends in
Ecology and Evolution. 2008;24:127-35.
59. Crawley MJ. The R book. Chichester, England ; Hoboken, N.J: Wiley; 2007.
60. S original by Kenneth Hess and R port by R. Gentleman. muhaz: Hazard function
estimation in survival analysis. 2019.
61. Moore DF. Applied survival analysis using R. Switzerland: Springer International
Publishing; 2016.
62. Lenth R. emmeans: Estimated marginal means, aka least-square means. 2020.
63. Wickham H, François R, Henry L, Müller K. dplyr: A grammar of data manipulation.
2020.
64. Wickham H. The split-apply-combine strategy for data analysis. Journal of Statistical
Software. 2011;40:1-29.
65. Revelle W. psych: Procedures for psychological, psychometric, and personality research.
2020.
66. Benjamini Y, Hochberg Y. Controlling the false discovery rate: A practical and powerful
approach to multiple testing. Journal of the Royal Statistical Society: Series B
(Methodological). 1995;57:289-300.
67. de Roode JC, Altizer S. Host-parasite genetic interactions and virulence-transmission
relationships in natural populations of monarch butterflies. Evolution. 2010;64:502-14.
68. de Roode JC, Pedersen AB, Hunter MD, Altizer S. Host plant species affects virulence in
monarch butterfly parasites. J Anim Ecol. 2008;77:120-6.
69. Clerc M, Ebert D, Hall MD. Expression of parasite genetic variation changes over the
course of infection: implications of within-host dynamics for the evolution of virulence.
Proc R Soc B. 2015;282.
70. Liu S, Moayeri M, Leppla S. Anthrax lethal and edema toxins in anthrax pathogenesis.
Trends in Microbiology. 2014;22(6):317-25. doi: 10.1016/j.tim.2014.02.012. PubMed
PMID: WOS:000337873500005.
71. Helgason E, Okstad OA, Caugant DA, Johansen HA, Fouet A, Mock M, et al. *Bacillus*
anthracis, *Bacillus cereus*, and *Bacillus thuringiensis*--one species on the basis of genetic
evidence. Appl Environ Microbiol. 2000;66(6):2627-30. doi: 10.1128/aem.66.6.2627-
2630.2000. PubMed PMID: 10831447; PubMed Central PMCID: PMCPMC110590.
72. Radnedge L, Agron PG, Hill KK, Jackson PJ, Ticknor LO, Keim P, et al. Genome
differences that distinguish *Bacillus anthracis* from *Bacillus cereus* and *Bacillus*
thuringiensis. Appl Environ Microbiol. 2003;69(5):2755-64. doi:
10.1128/aem.69.5.2755-2764.2003. PubMed PMID: 12732546; PubMed Central PMCID:
PMCPMC154536.

73. Kotiranta A, Lounatmaa K, Haapasalo M. Epidemiology and pathogenesis of *Bacillus*
cereus infections. Microbes and Infection. 2000;2(2):189-98. doi: 10.1016/S1286-
4579(00)00269-0. PubMed PMID: WOS:000086217900010.
74. Bertels F, Marzel A, Leventhal G, Mitov V, Fellay J, Gunthard HF, et al. Dissecting HIV
virulence: Heritability of setpoint viral load, CD4+ T-cell decline, and per-parasite
pathogenicity. Molecular Biology and Evolution. 2018;35(1):27-37. Epub 2017/10/14.
doi: 10.1093/molbev/msx246. PubMed PMID: 29029206; PubMed Central PMCID:
PMCPMC5850767.
75. Ellner SP, Buchon N, Dörr T, Lazzaro BP. Host-pathogen immune feedbacks can explain
widely divergent outcomes from similar infections. bioRxiv. 2021.
76. Shinzawa N, Nelson B, Aonuma H, Okado K, Fukumoto S, Miura M, et al. p38 MAPK-
dependent phagocytic encapsulation confers infection tolerance in *Drosophila*. Cell Host
Microbe. 2009;6:244-52. Epub 2009/09/15. doi: 10.1016/j.chom.2009.07.010. PubMed
PMID: 19748466.
77. McGonigle JE, Purves J, Rolff J. Intracellular survival of *Staphylococcus aureus* during
persistent infection in the insect *Tenebrio molitor*. Dev Comp Immunol. 2016;59:34-8.
Epub 2016/01/19. doi: 10.1016/j.dci.2016.01.002. PubMed PMID: 26778297.
78. Nyenje ME, Green E, Ndip RN. Evaluation of the effect of different growth media and
temperature on the suitability of biofilm formation by *Enterobacter cloacae* strains
isolated from food samples in South Africa. Molecules. 2013;18:9582-93. Epub
2013/08/24. doi: 10.3390/molecules18089582. PubMed PMID: 23966079; PubMed
Central PMCID: PMCPMC6270404.
79. Chodorski J, Hauth J, Strieth D, Wirsen A. Diffusion profiles in *L. lactis* biofilms under
different conditions. Engineering in Life Sciences. 2020.
80. Høiby N, Bjarnsholt T, Givskov M, Molin S, Ciofu O. Antibiotic resistance of bacterial
biofilms. Int J Antimicrob Agents. 2010;35:322-32. Epub 2010/02/13. doi:
10.1016/j.ijantimicag.2009.12.011. PubMed PMID: 20149602.
81. Mulcahy H, Sibley CD, Surette MG, Lewenza S. *Drosophila melanogaster* as an animal
model for the study of *Pseudomonas aeruginosa* biofilm infection *in vivo*. PLoS Pathog.
2011;7.
82. Brauner A, Fridman O, Gefen O, Balaban N. Distinguishing between resistance,
tolerance and persistence to antibiotic treatment. Nat Rev Microbiol. 2016;14(5):320-30.
doi: 10.1038/nrmicro.2016.34. PubMed PMID: WOS:000374147900010.
83. Coates J, Park BR, Le D, Simsek E, Chaudhry W, Kim M. Antibiotic-induced population
fluctuations and stochastic clearance of bacteria. Elife. 2018;7:e32976. doi: ARTN
e32976
10.7554/eLife.32976. PubMed PMID: WOS:000427105100001.
84. Liehl P, Blight M, Vodovar N, Boccard F, Lemaitre B. Prevalence of local immune
response against oral infection in a *Drosophila/Pseudomonas* infection model. PLoS
Pathog. 2006;2(6):e56. doi: 10.1371/journal.ppat.0020056.
85. Opota O, Vallet-Gély I, Vincentelli R, Kellenberger C, Iacovache I, Gonzalez MR, et al.
Monalysin, a novel ß-pore-forming toxin from the *Drosophila* pathogen *Pseudomonas*
entomophila, contributes to host intestinal damage and lethality. PLoS Pathog. 2011;7.
86. Chakrabarti S, Liehl P, Buchon N, Lemaitre B. Infection-induced host translational
blockage inhibits immune responses and epithelial renewal in the *Drosophila* gut. Cell
Host & Microbe. 2012;12:60-70.
87. Buchon N, Broderick NA, Lemaitre B. Gut homeostasis in a microbial world: insights
from *Drosophila melanogaster*. Nat Rev Microbiol. 2013;11:615-26. doi:
10.1038/nrmicro3074.

88.   Moreno-García M, Condé R, Bello-Bedoy R, Lanz-Mendoza H. The damage threshold
1072   hypothesis and the immune strategies of insects. Infection, Genetics and Evolution.
1073   2014;24:25-33. doi: 10.1016/j.meegid.2014.02.010.
1074   89.   Kuraishi T, Binggeli O, Opota O, Buchon N, Lemaitre B. Genetic evidence for a
1075   protective role of the peritrophic matrix against intestinal bacterial infection in
1076   *Drosophila melanogaster*. PNAS. 2011:1-6. doi: 10.1073/pnas.1105994108.
1077   90.   Shibata T, Maki K, Hadano J, Fujikawa T, Kitazaki K, Koshiba T, et al. Crosslinking of a
1078   peritrophic matrix protein protects gut epithelia from bacterial exotoxins. PLoS Pathog.
1079   2015;11(10):e1005244. Epub 2015/10/27. doi: 10.1371/journal.ppat.1005244. PubMed
1080   PMID: 26506243; PubMed Central PMCID: PMCPMC4646701.
1081

 Figure legends

 Fig 1. Decomposing virulence. Both exploitation and per parasite pathogenicity (PPP) can

 harm the host and thereby contribute towards virulence. Exploitation describes the infection

 intensity or bacterial load inside the host. PPP describes the damage per parasite that an

 infection does to the host. In this figure we show variation among pathogen species in

 exploitation and PPP, as illustrated by hypothetical relationships between host fitness and

 infection intensity for two species of parasite infecting the same host genetic background. **A**.

 In this example, the two parasite species have the same PPP but differ in exploitation. Parasite

 1 has lower exploitation compared with parasite 2, because it causes a lower infection

 intensity. **B**. In contrast with A, in this example, parasite species have the same average

 exploitation but species 1 has lower PPP because its reaction norm has a shallower slope. This

 means that compared with species 2, species 1 causes less damage to the host with increasing

 parasite load. Figure modified from [29].

 Fig 2. Fly survival after injection with one of four bacterial species. A − D. Survival

 curves after injection with four bacterial species, each at one of five doses. Controls were

 either injected with Ringer's solution or received no injection (naïve). Each survival curve is

 from n = 79 to 108 flies. The legend in panel **A**, shows the treatments for all survival curves.

 E. The natural log of maximum hazard for all bacterial species, where each data point is the

 maximum hazard calculated from one replicate per dose. Ringer's injected and naïve flies are

 not included. Black lines show means. Different letters denote means that are significantly

 different from one another (Tukey multiple comparison test).

 Fig 3. Bacterial load per living fly after injection with one of four bacterial species (A −

 D). Flies were homogenised at between 1- and 35-days post-injection. The injection dose

 legend for all panels is shown in **D**. The arrows on the y-axis indicate the approximate

1108 injection doses. Missing data are due to increasing fly death over time. Black lines show

1109 medians.

1110

Fig 4. Virulence decomposition. A. Parasite exploitation given as infection intensity/bacteria

load across bacterial species. Each data point is from one injection dose per bacteria, per

experimental replicate, and gives the geometric mean of bacterial load for days one and two

post injection (denoted as $_{1,2}$). The circles are jittered along the x-axis to aid visualisation of

overlapping data points. Black lines show means. Different letters denote means that are

significantly different from one another (Tukey multiple comparison test). **B**. PPP given as

the relationship between bacterial load and maximum hazard. The bacterial load data is the

same as that given in **A** but with the addition of the Ringer's control group. To allow

inclusion of the uninfected Ringer's control group to the figure we added one CFU to all

mean bacterial load values. The natural log of maximum hazard data is estimated from

survival data for the corresponding injection doses and experimental replicates. Maximum

hazard is plotted as the inverse, such that the hazard (virulence) increases with proximity to

the x-axis. Lines show linear regressions.

1124

Fig 5. Bacterial clearance in dead flies. Each row shows flies that had been injected with

one of four bacterial species. **A-D** The proportion of dead flies that were infected and

uninfected according to the day post injection at which they died and were homogenised.

Dead flies were homogenised at between 14 and 35, and 56 and 78, days post injection. **E-H**

The same data as shown in the left-hand panels but graphed by injection dose. Numbers above

the bars indicate the total numbers of flies from which the proportions were calculated, *i.e.*,

the total numbers of flies homogenised. Note that we cannot distinguish between flies that had

cleared the infection and those where the bacterial load was below our detection limit.

1133

Fig 6. The relationship between bacterial load at seven days post injection, and the initial injection doses. Each row shows data from one bacterial species. Panel **A** contains no flies injected with 92 CFUs because all flies had a bacterial load of zero at day seven; **C** contains no flies injected with 92,000 CFUs because all flies had died by this time point. Each circle is the bacterial load of one fly, they are jittered along the x-axis to aid visualisation of overlapping data points, and they are coloured according to the injection dose. Flies with zero bacterial load are not shown (see methods). Linear regression lines are shown in black with 95 % confidence intervals. Asterisks denote significant correlations, where $p < 0.0001$.

Fig 7. Bacterial clearance by living flies. Each row shows flies that had been injected with one of four bacterial species. **A-D** The proportion of live flies that were uninfected. Each column shows a different injection dose. Numbers above the bars indicate the total numbers of flies from which the proportions were calculated, *i.e.,* the total numbers of flies homogenised. Note that we cannot distinguish between flies that had cleared the infection and those where the bacterial load was below our detection limit.

Fig 8. Effects of injection dose, species and exploitation on bacterial clearance. For all figures, each data point is from one injection dose per bacteria, per experimental replicate, and gives the mean proportion of cleared infections (out of the initial infected population) on days one and two (clearance index$_{1,2}$), days three and four (clearance index$_{3,4}$) or days seven, 14 and 21 (clearance index$_{7,14,21}$). **A-D** Effect of injection dose on clearance index$_{1,2}$ for each bacterial species. There was a statistically significant negative correlation for **A**. *E. cloacae* ($\rho = -0.86$, $p < 0.001$), but not for **B**. *Pr. burhodogranariea* ($\rho = -0.36$, $p = 0.182$), **C**. *L. lactis* ($\rho = 0.13$, $p = 0.657$) or **D**. *Ps. entomophila* ($p = 0.982$, $\rho = -0.01$). **E**. Mean species differences in clearance index$_{3,4}$. The circles are jittered along the x-axis to aid visualisation of overlapping data points. Black lines show means. Different letters denote means that are

significantly different from one another (Mann-Whitney-U post hoc tests). The effect of

parasite exploitation, given as bacterial load, upon **F**. mean clearance index$_{3,4}$ and **G**. mean

clearance index$_{7,14,21}$. The geometric mean of bacterial load was calculated from days 1 and 2

post injection (denoted as $_{1,2}$), *i.e.*, the same values as in figure 3. There was a negative

relationship between the two variables. Statistics are given in the main text and the species

legend for both panels is shown in G.

Fig 1

Fig 2

Fig 3

A

B

1169   **Fig 4**

Fig 5

1172

Fig 6

1173

 Fig 7

1176   **Fig 8**

Supporting Information

S1 Fig. Illustration of the predictions for how clearance is expected to change with A. per parasite pathogenicity (PPP) and B. exploitation.

S2 Fig. Per parasite pathogenicity using different smoothing parameter values to estimate the maximum hazard.

S3 Fig. Effect of exploitation (bacterial load) on clearance for different values of the smoothing parameter.

S1 File. Statistical methods and results for the comparison between the proportions of live and dead uninfected flies.

S1 Table. Tukey multiple comparisons between bacterial species for differences in virulence.