1 **Invertebrate methylomes provide insight into mechanisms of environmental**
2 **tolerance and reveal methodological biases**
3

Shelly A. Trigg*[1], Yaamini R. Venkataraman*[1], Mackenzie R. Gavery[2], Steven B.

Roberts[1], Debashish Bhattacharya[3], Alan Downey-Wall[4], Jose M. Eirin-Lopez[5], Kevin M.

Johnson[6,7], Katie E. Lotterhos[4], Jonathan B. Puritz[8] and Hollie M. Putnam[8+]

[1] University of Washington, School of Aquatic and Fishery Sciences 1122 NE Boat St.
Seattle, WA, 98195, USA
[2] Environmental and Fisheries Sciences Division, Northwest Fisheries Science Center,
National Marine Fisheries Service, National Oceanic and Atmospheric Administration,
2725 Montlake Blvd E, Seattle, WA, 98112, USA
[3] Department of Biochemistry and Microbiology, Rutgers University, New Brunswick, NJ
08901 USA
[4] Department of Marine and Environmental Sciences, Northeastern University, 430 Nahant
Road, Nahant, MA 01908
[5] Florida International University, Environmental Epigenetics Laboratory, Institute of
Environment 3000 NE 151 St. North Miami, FL, 33181, USA
[6] Center for Coastal Marine Sciences, California Polytechnic State University, San Luis
Obispo, CA, 93407, USA
[7] California Sea Grant, University of California San Diego, La Jolla, CA, 92093
[8] Department of Biological Sciences, University of Rhode Island, Kingston, RI 02881, USA
23
24 + corresponding author: hputnam@uri.edu
25
26 * joint first authors/equal contribution

27
28 Keywords: bisulfite sequencing, coral, epigenetics, marine invertebrate
29

Abstract

There is a growing focus on the role of DNA methylation in the ability of marine

invertebrates to rapidly respond to changing environmental factors and anthropogenic

impacts. However, genome-wide DNA methylation studies in non-model organisms are

currently hampered by limited understanding of methodological biases. Here we compare

three methods for quantifying DNA methylation at single base pair resolution — Whole

Genome Bisulfite Sequencing (WGBS), Reduced Representation Bisulfite Sequencing

(RRBS), and Methyl-CpG Binding Domain Bisulfite Sequencing (MBDBS) — using

multiple individuals from two reef-building coral species with contrasting environmental

sensitivity. All methods reveal substantially greater methylation in *Montipora capitata*

(11.4%) than the more sensitive *Pocillopora acuta* (2.9%). The majority of CpG

methylation in both species occurs in gene bodies and flanking regions. In both species,

MBDBS has the greatest capacity for detecting CpGs in coding regions at our sequencing

depth, however MBDBS may be limited by intra-sample methylation heterogeneity. RRBS

yields robust information for specific loci albeit without enrichment of any particular

genome feature and with significantly reduced genome coverage. Relative genome size

strongly influences the number and location of CpGs detected by each method when

sequencing depth is limited, illuminating nuances in cross-species comparisons. These

findings reinforce the role and importance of DNA methylation underlying environmental

sensitivity in critical marine invertebrate taxa, and provide a genomic resource for

investigating the functional role of DNA methylation in environmental tolerance.

Introduction

Environmental stimuli interact with genomic content to drive variation in gene and

protein expression, resulting in phenotypic plasticity. This plasticity has the potential to

buffer against mortality under environmental change (Baldwin, 1902), or conversely be

maladaptive (Velotta et al., 2018). Furthermore, plasticity may enhance or diminish

evolutionary rates (Ghalambor et al., 2007), which is particularly relevant to plasticity-

evolution feedbacks (Ghalambor et al., 2007, 2015; Kronholm & Collins, 2016). This is of

particular concern in the Anthropocene (Lewis & Maslin, 2015), as global change

exacerbates the mismatch between phenotype and a rapidly changing environment.

The increase in negative global climate change consequences have prompted an

intensification of research into phenotypic plasticity, gene regulation, and epigenetic

mechanisms in non-model marine invertebrates (Eirin-Lopez & Putnam, 2019; Hofmann,

2017; Roberts & Gavery, 2012). Specifically, carryover effects and cross and multi-

generational plasticity in response to climate change (Byrne et al., 2020) may be

generated by epigenetic regulation of gene expression (Dixon et al., 2018; Liew et al.,

2018, 2020). As epigenetic research has increased, there has been a focus on DNA

methylation, or the addition of a methyl group on the Cytosine residues in the genome,

often in the Cytosine phosphate Guanine, or CpG context (Zemach et al., 2010). DNA

methylation has gene expression regulation capacity through the interaction of base

modification with transcriptional elements. Early bulk enzyme-based and fingerprinting

methods for quantifying DNA methylation in marine invertebrates provided initial insights

into DNA methylation and organismal phenotypic plasticity in response to environmental

changes (Dimond et al., 2017; Gavery & Roberts, 2010; Gonzalez-Romero et al., 2017;

Putnam et al., 2016; Rodriguez-Casariego et al., 2018; Suarez-Ulloa et al., 2018).

Non-sequencing approaches that quantify global or bulk methylation [e.g.,

colorimetric or fluorescent ELISAs (Dimond et al., 2017; Gavery & Roberts, 2010; Putnam

et al., 2016; Rodriguez-Casariego et al., 2018)] are low-cost, rapidly applied, and do not

require genomic resources to generate information on the responsiveness of the

methylome. These global estimates do not, however, fully capture local changes in DNA

methylation across different genome regions. Specifically, differences in the location and

amount of methylation in two samples or treatments could lead to an incorrect conclusion

when based on average percent methylation at the bulk level. Consequently, non-

sequencing methods are limited in their ability to elucidate specific mechanisms of

expression regulation and thus are unable to fully address the functional implications of

methylation-driven regulation within the genome. In contrast, the use of genome-wide

approaches that provide single base pair resolution allow the testing of hypotheses

regarding spurious transcription, alternative splicing, and exon skipping (Roberts &

Gavery, 2012). For example, the use of Whole Genome Bisulfite Sequencing (WGBS) to

investigate the role of DNA methylation in regulating genes involved in caste specification

in honeybees identified differential methylation in an exon of the *anaplastic lymphoma*

kinase (ALK) gene; this exon was differentially retained in a splice variant between queens

and workers (Foret et al., 2012). Thus there is a clear need for single base pair

assessment of DNA methylomes facilitated by next generation sequencing to more fully

elucidate the relationship of DNA methylation and gene expression in non-model

invertebrates.

Genome-wide levels of DNA methylation can be estimated by several bisulfite

conversion and sequencing approaches. Bisulfite conversion of DNA results in the

deamination of unmethylated Cytosine to Uracil, which leaves a base change signature in

the DNA that can be tracked via comparison of sequence between bisulfite-converted

samples and reference genomes. While the number of bisulfite sequencing approaches

are expanding [e.g., epiGBS (van Gurp et al., 2016)], the widely-used approaches are

WGBS, Reduced Representation Bisulfite Sequencing (RRBS), and more recently,

Methyl-CpG Binding Domain Bisulfite Sequencing (MBDBS). WGBS is considered to be

the gold-standard of bisulfite sequencing because it provides full coverage of the genome

and the capacity to detect the entire methylome at single base pair resolution.

While providing a comprehensive approach, the high cost of WGBS is juxtaposed

against the often very small fraction of methylated DNA common in invertebrate genomes

(Tweedie et al., 1997). Alternatively, approaches such as RRBS also use bisulfite

conversion to quantitatively assess DNA methylation with base pair resolution. RRBS

incorporates a restriction digestion of the genome to enrich for CpG rich regions, and is

designed to capture the majority of promoters and other genomic regions containing CpG

islands because they have important regulatory functions in vertebrates (Meissner et al.,

2008). This is a more cost-effective approach provided by sequencing only a small portion

of the genome, but requires restriction enzyme recognition sites near other CpGs to gather

high resolution data.Since DNA methylation in invertebrates is primarily limited to coding

regions (Dixon et al., 2018; Flores et al., 2012; Roberts & Gavery, 2012), it is less clear

whether enrichment of CG-rich DNA using RRBS will capture informative or regulatory

regions of invertebrate genomes, making the cost savings moot in the absence of

informative data.

In contrast to the CpG-rich, region-specific targeting of RRBS, MBDBS uses Methyl

Binding Domain Proteins to target and enrich methylated CpGs, then employs bisulfite

conversion to provide single base pair resolution of DNA fragments with methylated

regions. Many marine invertebrate genomes consist of highly methylated regions that are

distributed in predominantly unmethylated DNA in a mosaic pattern (Suzuki et al., 2007).

When invertebrate methylomes have been characterized, these highly methylated regions

overlap with gene bodies and have been shown to play a role in gene expression activity

(Roberts & Gavery, 2012). Therefore, using an enrichment approach such as MBDBS to

isolate gene body methylation can be a cost-effective and gene-body focused alternative

to WGBS or RRBS (Gavery & Roberts, 2013; Venkataraman et al., 2020). The base-pair

resolution offered by the combination of MBD enrichment and BS is an advantage

compared to MBD-seq alone (Dixon & Matz, 2020), as the latter assumes that methylation

level is proportional to read depth. In contrast to WGBS or RRBS, the quantification and

interpretation of MBDBS data can be complicated by individual variation in methylation

levels (e.g., one individual who has high methylation in a particular region would have

reads, whereas another individual who lacks methylation in that region would have

missing data).

Given the need to assess plasticity mechanisms and the acclimatization potential

of a variety of marine taxa, it is critical to compare the potential of different approaches to

to detect, quantify, and assess DNA methylation with respect to specific biological

hypotheses of interest. To this end, we studied three DNA methylation quantification

approaches that provide single base pair resolution data using bisulfite conversion and

sequencing: WGBS, RRBS, and MBDBS. We applied these methods to two reef building

corals, *Montipora capitata* and *Pocillopora acuta*, which have different environmental

sensitivity, phenotypic plasticity, inducible DNA methylation (Putnam et al., 2016), and

genome sizes (Shumaker et al., 2019; Vidal-Dupiol et al., 2019). We assessed species-

specific differences in genome-wide methylation and contrasted percent methylation of

common loci and orthologous genes across methods. Then, we compared the coverage

and genomic location of CpG data generated from the three methods. Compared to

WGBS, both MBDBS and RRBS have advantages and potential limitations associated

with biology, genome characteristics, and experimental design, highlighting the need to

fully consider these aspects when evaluating DNA methylation for particular hypotheses

of function in invertebrates. As part of this effort, we described DNA methylation

differences in two coral species, providing valuable insights into the epigenetic

underpinnings of phenotypic plasticity.

Materials and Methods

Sample collection

The reef-building scleractinian coral species *Montipora capitata* and *Pocillopora*

acuta were collected from the reefs of Kaneʻohe Bay Hawaiʻi under SAP 2019-60 between

4 - 7 September 2018. Corals were transported to the Hawaiʻi Institute of Marine Biology

where they were held in tanks under ambient conditions for 15 days and then snap frozen

in liquid nitrogen and stored at -80°C until nucleic acid extraction was performed. For each

of the two coral species, fragments were collected from three different individuals collected

from ambient conditions.

Nucleic Acid Extraction

Samples were removed from -80°C and small tissue fragments were clipped

directly into a tube containing RNA/DNA shield (1 ml) and glass beads (0.5 mm). The

tissue clippings consisted of all coral cell types and their symbionts. Samples were

homogenized on a vortexer for 1 minute for the thin tissue imperforate coral *Pocillopora*

acuta and 2 minutes for the thick tissue perforate coral *Montipora capitata* at maximum

speed to ensure tissue extraction of all cell types. The supernatant was removed and DNA

was extracted using the Zymo Quick-DNA/RNA™ Miniprep Plus Kit and subsequently

checked for quality using gel electrophoresis on an Agilent 4200 TapeStation and

quantified using a Qubit. In summary, one DNA preparation was made from each of the

three individuals per coral species and was subsequently divided into three aliquots for

each of the three bisulfite sequencing methods (WGBS, MBDBS, and RRBS) to yield a

total of 18 libraries (**Figure 1**).

Genome Information

Previously sequenced and assembled coral genomes were used for mapping of

DNA methylation data. These include *Montipora capitata* (Shumaker et al., 2019) and

Pocillopora acuta (Vidal-Dupiol et al., 2019). Both of the coral genomes have a high and

similar number of predicted genes (63,227 in *M.capitata* and 64,558 in *P. acuta*). However,

P. acuta is much smaller in size (~352MB vs. ~886 MB in *M. capitata*), has less repetition,

a greater number of scaffolds (25,553 in *P. acuta* vs. 3,043 in *M.capitata*), and lower

genome assembly continuity (N50 is 171,375 in *P. acuta* and 540,623 in *M. capitata*).

192   Genome feature tracks for *M. capitata* and *P. acuta* were derived directly from the

published genomes for use in DNA methylation analyses. The *M. capitata* genome

annotation yielded gene (a combination of AUGUSTUS and GeMoMa predictions), coding

sequence, and intron tracks (Shumaker et al., 2019). Similarly, gene (AUGUSTUS

predictions), coding sequence, and intron information was obtained from the *P. acuta*

genome (Vidal-Dupiol et al., 2019). Flanking regions upstream and downstream of genes

were generated with BEDtools v2.29.2 for each genome separately (Quinlan & Hall,

2010); `flankBED` was used to generate 1000 bp flanks upstream and downstream of

annotated genes from each genome. Overlaps between genes and flanks were removed

from up- or down-stream flanking region tracks using `subtractBED`. Similarly, an

intergenic region track was created by finding the complement of genes with

`complementBED`, then removing any overlaps with flanking regions using `subtractBED`.

All tracks were verified with the Integrative Genomics Viewer (Thorvaldsdóttir et al., 2013).

Feature track files generated for both species are available in the project large file

repository (Putnam et al., 2020)

MBD Enrichment

209   Before enrichment, DNA (1 μg) in 80 μL Tris HCl was sheared to 500 bp using a

QSonica Q800R3. Samples were sonicated for 90 sec, with 15 sec on and 15 sec off

intervals at 25% amplitude. Fragment length was checked using a D5000 TapeStation

System (Agilent Technologies) and samples were sonicated an extra 15 sec to shear DNA

from 600 bp to 500 bp as needed.

The MethylMiner kit (Invitrogen; Cat. #ME10025) was used to enrich for methylated

DNA prior to MBDBS library generation, with 1µg of input DNA. Manufacturer's

instructions were adhered to with the following modifications: The capture reaction

containing the fragmented DNA and MBD beads was incubated with mixing at 4ºC

overnight, and enriched DNA was obtained with a single fraction elution using 2,000 mM

NaCl. Following ethanol addition, samples were centrifuged at 14,000 rcf at 1ºC for five

min. Pellets were resuspended in 25 µL ultra-pure water. Captured DNA was quantified

using a Qubit dsDNA HS Kit (Invitrogen).

MBDBS and WGBS Library Preparation

WGBS and MBDBS libraries were prepared using the Pico Methyl-Seq Library Prep

Kit (ZymoResearch Cat. # D5456). Manufacturer's instructions were followed with the

following modifications: For each sample, 1 ng of coral DNA and 0.05 ng of lambda phage

DNA (ZymoResearch Cat. # D5016) were used. Samples were always centrifuged at

12,000 rcf for 30 sec, however, samples were centrifuged for 12,000 rcf for 90 sec after

the second 200 µL addition of M Wash Buffer. Warmed elution buffer (56°C) was added

to each sample to increase DNA elution yield. During the second amplification cycle, 0.5

µL of PreAmp Polymerase was added. After initial clean-up with the DNA Clean-up and

Concentrator, the first amplification step was run for eight cycles. For amplification with i5

and i7 index primers, 1 µL of each primer was used to improve amplification. The volume

of the 2X LibraryAmp Master Mix was increased to 14 µL to match the increase in index

primers.

To remove excess primers from WGBS and MBDBS preparations, samples were

cleaned with 11 µL of KAPA pure beads (1X) (KAPA Cat # KK8000) and 80% ethanol.

Cleaned samples were resuspended in 12 µL of room-temperature DNA elution buffer

from the Pico Methyl-Seq Library Prep Kit. Samples were re-amplified with either two or

four cycles, depending on DNA concentration. Re-amplification was conducted with only

0.5 µL of each i5 and i7 index primer. After re-amplification, 26 µL of KAPA pure beads

(1X) and 80% ethanol were used for clean-up. Final samples were resuspended in 14 µL

of room-temperature elution buffer. Primer removal was confirmed by running samples on

a D5000 TapeStation System.

RRBS Library Prep

RRBS libraries were prepared with the EZ DNA RRBS Library Prep Kit

(ZymoResearch Cat. # D5460). Manufacturer's instructions were used with the following

modifications: For MspI digestion, 300 ng of input DNA and 15 ng of lambda phage DNA

were used. Digestions were carried out at 37°C for 4 hours. Adapter ligation was

performed overnight, with samples held at 4°C once cycling was completed. Similar to

WGBS and MBDBS library preparation, samples were always centrifuged at 12,000 rcf for

30 sec, with the exception of 90 sec centrifugation after the second 200 µL addition of M

Wash Buffer. Warmed elution buffer (56°C) was added to each sample to increase DNA

elution yield. Index primers were ligated using eleven cycles of the recommended

thermocycling protocol. Samples were cleaned using 50 µL of KAPA pure beads (1X) and

80% ethanol, then resuspended in 16 µL of the elution buffer. Primer removal was

confirmed by running samples on a D5000 TapeStation System.

DNA Sequence Alignment

All libraries (n = 18) were pooled in equimolar amounts and loaded at 250 pM onto

a single Illumina NovaSeq S4 flow cell lane for 2x150 bp sequencing at Genewiz (South

Plainfield, NJ). This was estimated to give 111-138 M reads per library and 99-123x

coverage of the *P. acuta* genome (3.3 M bp) and 38-47x coverage of the *M. capitata*

genome (8.8 M bp), assuming 100% even coverage (e.g., 150 bp read * 2 pairs * 111 M

reads/336,684,533 bp for *P. acuta*).

Sequence quality was checked by FastQC v0.11.8 and adapters from paired-end

sequences were trimmed using TrimGalore! version 0.4.5 (Krueger, 2012). Following

recommendations for methylation sequence analysis from the manufacturer's protocol

and from the Bismark User Guide, 10 bp were hard trimmed from the 5' and 3' end of each

read for WGBS and MBDBS samples, and RRBS samples were trimmed with `--`

`non_directional` and `--rrbs` options. Bisulfite-converted genomes were created in-

silico with Bowtie 2-2.3.4 [Linux x84_64 version; (Langmead & Salzberg, 2012)) using

`bismark_genome_preparation` through Bismark v0.21.0 (Krueger & Andrews, 2011).

Trimmed reads were aligned to the BS-converted *P. acuta* genome (Vidal-Dupiol et al.,

2019) and the BS-converted *M. capitata* genome (Shumaker et al., 2019) with Bismark

v0.21.0 with alignment stringency set by `-score_min L,0,-0.6` and the default MAPQ

score threshold of 20. To check mapping rates for endosymbionts and quantify percent

methylation, trimmed reads from *P. actua* libraries were also aligned to the *Cladicopium*

goreaui genome [type C1, previously *Symbiodinium goreaui* (Liu et al., 2018)] using the

same settings as specified above. Reads that mapped ambiguously were excluded and

alignment files containing uniquely mapped reads were deduplicated with

`deduplicate_bismark` for WGBS and MBDBS samples only. Methylation calls were

extracted from sorted deduplicated alignment files using

`bismark_methylation_extractor`. Cytosine coverage reports were generated using

`coverage2cytosine` with the `--merge_CpG` option to combine data from both strand

methylation. Resulting files include bedgraphs and Bismark coverage files (Putnam et al.,

2020). MultiQC v1.8 (Ewels et al., 2016) was run on the trimmed reads, FastQC output,

and Bismark reports to assess quality and summarize results.

Bisulfite conversion efficiency assessment

Trimmed sequence reads were aligned to the genome of *E. coli* strain K-12

MG1655 (Riley et al., 2006) using Bismark v0.21.0 with the `-non_directional` option

and alignment stringency set by `-score_min L,0,-0.6`. Bisulfite conversion

efficiency was also estimated from coral alignments as the ratio of the sum of

unmethylated cytosines in CHG and CHH context to the sum of methylated and

unmethylated cytosines in CHG and CHH. ANOVA was used to test for an effect of

library preparation method on conversion efficiency within each species (conversion

efficiency ~ library preparation method) for both estimated and lambda alignment

calculated conversion efficiencies. A two-sample t-test was used to test if conversion

efficiency calculated from lambda alignments was the same as estimated conversion

efficiency for each library preparation method within each species.

Genome-Wide Methylation

General *M. capitata* and *P. acuta* methylation was characterized to describe any

species specific patterns. This was carried out by combining BEDgraphs derived from all

methods for each species using `unionBedGraphs`. Percent methylation for every CpG

locus with at least 5x coverage was averaged, irrespective of how many samples had

coverage for that locus. Loci with no data within a method were excluded from downstream

analysis. CpGs were classified as being either highly methylated (≥ 50% methylation),

moderately methylated (>10% and <50%), or lowly methylated (≤10% methylation).

Percent Methylation of Shared CpG Loci

Comparisons of percent DNA methylation at CpG loci analyzed by more than one

method were performed using the R-package `methylKit` (Akalin et al., 2012). A

minimum of 5x coverage was required across all samples for a CpG locus to be

considered in the analyses. The `unite` function in methylKit was used to identify CpG loci

that were covered across all 9 samples (3 individuals per method) per species.

Scatterplots and Pearson correlation coefficients were calculated using the function

`getCorrelation`.   Additionally, differential methylation tests were performed on

pairwise comparisons between methods (WGBS versus RRBS, WGBS versus MBDBS,

and RRBS versus MBDBS). Discordant methylation was quantified using a logistic

regression model on CpG loci that were covered across all 6 samples (3 samples from

each method compared) in each pairwise comparison using the `calculateDiffMeth`

function with default parameters.

CpG Coverage

To assess average genome-wide CpG coverage, the number of Cytosines passing

different read depth thresholds (5x, 10x, 15x, 20x, 25x, 30x, 40x, and 50x) were totaled

from the CpG coverage reports output by the Bismark `coverage2cytosine` function

(detailed above) for each sample. These totaled CpGs were then relativized to the number

of CpGs in their respective genomes (*M. capitata,* 28,684,519 CpGs; *P. acuta,* 9,155,620

CpGs). Next, average and standard deviation of genome-wide CpG fractions were

calculated for each method within each species (n = 3), and these were plotted across

different read depth thresholds using ggplot2 (Gómez-Rubio, 2017).

To estimate overall genome-wide CpG coverage, a downsampling analysis was

performed by pooling all sample reads within a method and species. Briefly, trimmed fastq

files were concatenated for each method and species then randomly subsampled to 50,

100, 150, and 200 million reads. Next, alignment and methylation calling were carried out

as described above on each subset, and the number of cytosines passing with 5 or more

reads were totaled from CpG coverage reports from each subset. Sequencing saturation

was estimated from a Michaelis-Menten model with the 'drm' function from the R package

drc (Ritz et al., 2015) using CpG coverage reports from subsampled data as input. Both

observed CpG coverage from subsampled data and estimated CpG coverage were

plotted using the R package ggplot2 (Gómez-Rubio, 2017).

Proportion of Detected CpGs for Orthologs

To describe the differences in DNA methylation detected by each method at a more

functional level, and given the connection of gene body methylation and gene expression

in invertebrates (Roberts & Gavery, 2012) and corals specifically (Liew et al., 2018), the

presence of CpG data within all genes was calculated for each species, by method. First,

a CpG gff track was generated using EMBOSS (Rice et al., 2011) with the `fuznuc`

command searching for the pattern CG. For each sample, `intersectBED` was used to

identify CpGs with 5x coverage that intersected with gene regions. This was also done for

the reference genome CpG gff track. CpG counts per gene were compiled for each sample

and the mean taken per method. The proportion of CpGs per orthologous gene was

calculated by dividing the mean number of CpGs with 5x coverage from the three samples

per method and dividing that by the number of CpG possible summed per gene from the

reference genome CpG gff track. The proportion of CpG data in a gene was then

visualized in heatmaps for all genes of *M. capitata* and *P. acuta*.

Genomic Location of CpGs

For both *M. capitata* and *P. acuta*, the overlap between genome feature tracks and

species-specific CpG data at 5x coverage was characterized with BEDtools v2.29.2 to

assess the presence CpGs in various regions by method (Quinlan & Hall, 2010). Since

only gene, coding sequence, intron, flanking regions, and intergenic region tracks were

common between species, these were the tracks used in downstream analyses. A

combination of PCoA, PERMANOVA and beta-dispersion tests, and chi-squared

contingency tests were used to determine if the library preparation method influenced the

proportion of CpGs detected in a specific genomic feature. A separate contingency test

was used for each genomic feature.

Results

To compare the performance of bisulfite sequencing methods in the reef-building

scleractinian corals *Montipora capitata* and *Pocillopora acuta*, we isolated DNA and

generated WGBS, RRBS, and MBDBS libraries for three individuals from each species to

yield a total of 18 libraries (**Figure 1**).

Sequencing of all 18 libraries resulted in 1.82×10^9 read pairs, of which 99.1%

remained after QC and trimming (Additional file 1: **Table ST1**). Individual libraries were

generally sequenced to the same depth (~7.5×10^7 reads) across library preparation

methods and species, with the exception of *P. acuta* RRBS libraries, which were

sequenced 2-4-fold deeper. The average mapping efficiencies for all *P. acuta* and all *M.*

capitata libraries were 45% and 39% read alignments, respectively (Additional file 2: **Table**

ST2). In comparison to other methods, MBDBS libraries had a larger proportion of reads

(73.1% ± 9.9%) that did not align to the coral genomes (Additional file 3: **Figure SF1**). To

investigate this we aligned *P. acuta* libraries to a known symbiont *Cladocopium goreaui*

genome (C1 (Liu et al., 2018)) for which the genome sequence was available at the time

of analysis. We found a sizable proportion of the MBDBS reads mapped to the symbiont

genome (23.6 ± 10.6%), while a much smaller proportion of RRBS and WGBS reads

mapped to the symbiont genome (5.04 ± 0.22% and 1.92 ± 0.3% respectively) (Additional

file 4: **Table ST3**).

Bisulfite conversion efficiency calculated from alignments of the unmethylated

lambda DNA spike-in ranged from 98.6 to 99.3% in *M. capitata* and from 98.3 to 99.1% in

P. acuta (Additional file 5: **Table ST4**), and this differed by library preparation method for

both *M. capitata* (F_{df}= X, *P* = 1.676e^{-05}) and *P. acuta* (F_{df}= X, *P* = 0.025) libraries. In

general, conversion efficiency calculated from the lambda alignments did not differ from

conversion efficiency estimates from CHG and CHH methylation (under the assumption

that non-CpG methylation does not occur in corals, see also (Liew et al., 2018)) from coral

alignments in *M. capitata* and *P. acuta*. (Additional file 6: **Figure SF2** and Additional file

7: **Table ST5**).

For each species, the general methylation landscape was characterized for CpG

loci with 5x coverage identified in any method. The *M. capitata* genome was more

methylated than *P. acuta* (**Figure 2**). Using a cutoff of ≥ 50% methylation to define

methylated CpGs, of the 13,340,268 CpGs covered by the *M. capitata* data, 11.4% were

methylated. In contrast, only 2.9% of the 7,326,297 CpGs in *P. acuta* were methylated.

Both genomes were predominantly lowly methylated (≤ 10% methylated): 79.6% CpGs in

M. capitata and 91.3% CpGs in *P. acuta* were lowly methylated. The remaining 9.0% of

CpGs in *M. capitata* and 5.8% of CpGs in *P. acuta* were moderately methylated (10-50%

methylation). The different methods captured varying proportions of highly, moderately,

and lowly methylated CpGs (Additional file 8: **Figure SF3**).

For quantitative comparison of method performance, we reduced the dataset to loci

covered at 5x read depth across all methods and samples for each species, referred to

here as 'shared loci'. The number of shared loci was 4,666 CpG for *M. capitata* and 93,714

CpG for *P. acuta*. A PCA of CpG methylation for loci covered at 5x read depth showed

that libraries tended to cluster in PC space by preparation method, rather than by

individual (Additional file 9: **Figure SF4**). Variation in methylation levels of the shared loci

across all *M. capitata* samples was lower within a method than between methods

(Additional file 9: **Figure SF4A**). In contrast, for *P. acuta*, RRBS and WGBS methods

showed similar methylation levels of shared loci but these were different from the

methylation level of loci identified in MBDBS (Additional file 9: **Figure SF4B**). To further

explore the variation in methylation observed by method, we directly correlated

quantitative methylation calls for the shared loci (**Figure 3**). For *M. capitata*, correlations

among biological replicates within a method were higher, whereas for *P. acuta*,

correlations were lower and more variable. Correlations between pairs of methods for *M.*

capitata ranged on average from 0.75-0.82, whereas correlations for *P. acuta* ranged from

0.40-0.64. For *M. capitata*, WGBS versus MBDBS had the highest correlation. For *P.*

acuta, WGBS versus RRBS had the highest correlation.

Discordance in methylation quantification between methods was evaluated by

identifying the number of CpG loci with large differences (>50%) in methylation for each

species. WGBS versus RRBS showed the lowest discordance in both species (0.4% for

M. capitata and 0.5% for *P. acuta*). The highest discordance in methylation was found in

comparisons with MBDBS for *P. acuta*, with 11% and 15% of CpG sites being called at

least 50% different for comparisons with WGBS and RRBS, respectively. In contrast, only

0.4% and 5% of common CpG sites were at least 50% difference between MBDBS versus

WGBS and MBDBS versus RRBS, respectively, for *M. capitata*. A majority of the

discordance was due to higher methylation calls in MBDBS compared to WGBS or RRBS

(**Figure 3B**).

Consistent with what would be expected based on genome size, *P. acuta* libraries

have higher genome-wide CpG coverage than *M. capitata* regardless of library

preparation method (**Figure 4A-C**). For both species, WGBS and MBDBS libraries

covered more CpGs than RRBS libraries, whereas RRBS libraries tended to show greater

read depth for the CpGs that it did cover. In other words, at >20x read depth, RRBS

libraries covered more CpGs than either WGBS or MBDBS (**Figure 4** insets). Modelling

increased sequencing depth for RRBS or MBDBS libraries showed little impact on the

fraction of genome-wide CpGs covered in *M. capitata*, while increasing sequencing depth

from 50 M to 200 M for WGBS libraries in both species and for MBDBS in *P. acuta* showed

a substantially larger fraction of CpGs covered (Additional file 10: **Figure SF5**).

In order to assess the potential for cross-species comparisons using an equivalent

dataset we quantified CpG data available across one-to-one orthologous genes. For *M.*

capitata, WGBS yielded the highest proportion of CpGs, followed by RRBS, and then

MBDBS (Additional file 11: **Figure SF6A**). This differed in *P. acuta* with WGBS yielding

the highest proportion of CpGs on average across orthologs, followed by MBDBS, and

then RRBS (Additional file 11: **Figure SF6B**).

In order to compare locations of CpG data between genomic features for each

species and method, all CpGs with 5x coverage were characterized based on genomic

feature location (**Figure 5)**. Global PERMANOVA tests found significant differences

between library preparation methods for CpG coverage in various genome features for *M.*

capitata and *P. acuta* (Additional file 12: **Table ST6**). Although post-hoc pairwise

PERMANOVA tests did not reveal differences between sequencing methods, power for

these was probably low (due to low sample size). Pairwise chi-squared tests indicated

there are differences in CpG location for both species. In particular, CpGs in gene bodies

were significantly enriched over other genomic features with MBDBS (**Figure 5**; Additional

file 13: **Table ST7**). Visual inspection of PCoA also revealed the proportion of CpGs

captured in coding sequences (CDS) drove differences between MBDBS and the other

methods in both species (**Figure 5C-D**).

Discussion

We evaluated the performance of three approaches that use bisulfite-treated library

preparation to enable single base pair resolution quantification of DNA methylation in

corals. Our results demonstrate that the methylation landscape can vary significantly

across species, which is a critical consideration for both interpreting environmental

response capacity, and therefore for experimental design. We report significant

differences in DNA methylation in two coral species that may contribute to their differential

environmental sensitivity of these organisms (Gibbin et al., 2015; Putnam et al., 2016).

Whereas WGBS is the gold standard for studying methylation, it comes at a high cost.

MBDBS enriches for gene regions, which may be useful for taxa with gene body

methylation. On the other hand, RRBS provides greater coverage depth for a smaller

fraction of the genome, but lacks specificity for genomic features, or DNA methylation.

Taken together, our findings indicate biology, genome architecture, regions of interest,

and depth of coverage are critical considerations when choosing methods for high

resolution quantification of DNA methylation profiles in invertebrates.

M. capitata has a relatively high environmental tolerance (Bahr et al., 2016; Gibbin

et al., 2015; Putnam et al., 2016), which has previously been attributed to its symbiont

composition (Cunning et al., 2016), genome characteristics (Shumaker et al., 2019),

perforate tissue-skeletal architecture and tissue thickness, and heterotrophic capacity

(Rodrigues & Grottoli, 2007). Of particular relevance to DNA methylation are genomic

aspects such as gene family duplication and high repeat content in *M. capitata* (Shumaker

et al., 2019). Given genetic-epigenetic correlations, particularly in the case of DNA

methylation and the requirement for a CpG sequence target site (Dimond & Roberts, 2020;

Johnson et al., 2020), variation in genome architecture, gene number, and content will

impact the presence and use of DNA methylation as a mechanism of gene expression

regulation. We found overall DNA methylation was higher in *M. capitata* than in *P. acuta*,

supporting early bulk analyses of DNA methylation in these species (Putnam et al., 2016).

While the predicted number of genes is similar, the genome size of *M. capitata* is over

twice that of *P. acuta* (Shumaker et al., 2019; Vidal-Dupiol et al., 2019). One explanation

for the higher methylation in *M. capitata* is that with greater energy availability — through

translocation from high density Symbiodiniaceae populations and energy stores in

perforate tissues — there is greater capacity for maintenance methyltransferase to

maintain high methylation, and thus reduce gene expression variability and spurious

expression (Liew et al., 2018; Li et al., 2018). High constitutive methylation could allow

"frontloading" of stress response genes (e.g., (Barshis et al., 2013)), providing greater

stress tolerance. Another possible explanation is that the higher level of methylation

contributes to the silencing of repeated genetic elements. In contrast, with a small and

non-repetitive genome, imperforate thin tissues, and low energy reserves, *P. acut*a may

be more energetically limited. Thus, *P. acuta* may be expected to show lower DNA

methylation across the genome as we demonstrate here, as well as a higher propensity

for inducible methylation in the presence of stressors (Putnam et al., 2016).

Another striking contrast in DNA methylation in these species is the lack of

concordance in the percent methylation values for *P. acuta* among methods compared to

M. capitata (**Figure 3**). The potential for chimerism in corals (Oury et al., 2020;

Schweinsberg et al., 2015) and differences in tissue structure (e.g., perforate or

imperforate) between species could contribute to differences in concordance across

methods for quantifying DNA methylation. One possibility is that Pocilloporids are chimeric

and multiple genotypes exist (Oury et al., 2020; Schweinsberg et al., 2015). Although

percent DNA methylation concordance across methods was generally high, in *P. acuta*

there was approximately a 10% higher level of discordance in percent methylation

quantification when compared WGBS to RRBS or MBDBS (**Figure 3**). This discordance

could have resulted from differences in *P. acuta* and *M. capitata* tissue structure. There is

the potential to homogenize and extract DNA from all cell types from the thin, imperforate

tissues of *P. acuta*, as opposed to the thick, perforate tissues in *M. capitata* (Putnam et

al., 2017), likely contributing to a greater number of cell types, and thus methylation

differences, captured in our *P. acuta* samples. Furthermore, the microhabitats created in

the tissues of these two species likely differ substantially spatially (Putnam et al., 2017),

creating cell-to-cell variability in methylation content. Since the likelihood of capturing

multiple cell types in bulk DNA extractions varies with tissue structure, future studies

should consider methods such as fluorescence-activated cell sorting (Hu et al., 2020;

Rosental et al., 2017), or laser microdissection (e.g., (Liew et al., 2018)), to target specific

tissues or cell types and reduce cell-to-cell methylation variability. Whereas this does not

necessarily indicate a bias in our methods, it highlights the needs to account for the

biological characteristics of a species when designing an experiment and evaluating

results.

The gold standard for bisulfite sequencing, WGBS, can be cost prohibitive

particularly if comparing multiple species and treatments. As expected, we found that

WGBS performed well, particularly for *P. acuta,* which has a smaller genome. Focusing

on gene orthologs, WGBS performed the best in terms of data for CpGs per gene. Based

on the gene ortholog comparisons, MBDBS provided more information than RRBS for *P.*

acuta, however the opposite held true for *M. capitata.* This is likely attributable to the

different genome size and inherent differences in methylation that result in differential

enrichment.

For both species, WGBS and MBDBS libraries covered more CpGs than RRBS

libraries; however, RRBS libraries showed greater read depth for CpGs. This is because

RRBS subsampled a specific, smaller portion of the genome than MBDBS or WGBS,

allowing more read coverage. Hence, CpG coverage did not largely increase when deeper

sequencing was modeled using RRBS data (Additional file 10: **Figure SF5**). RRBS was

designed to enrich for CpG islands, short stretches of DNA with higher levels of CpGs (~1

CpG per 10bp), that are typically found in mammalian promoters and enhancer regions

and thought to play a role in gene regulation (Gu et al., 2011). We found RRBS yielded a

well-covered reduced representation of the genome, which is important for bisulfite data

where high read depth is desired, and locus methylation levels were concordant with

WGBS for both species. However, RRBS did not enrich for promoters or other particular

genomic regions compared to the other bisulfite sequencing methods (**Figure 5**), and in

fact tended to identify unmethylated regions. For this reason, RRBS is not the best choice

for gene-focused methylation studies in corals and other invertebrates.

A critical consideration in deciding to perform MBDBS in corals is the amount of

DNA methylation present in any symbiont. If any non-target organisms have substantially

more DNA methylation than the target organism, MBDBS data could become saturated

by methylated DNA from non-target organisms, lowering sampling of the target species.

We observed this in *P. acuta*, for which we had the genome of its Symbiodiniaceae which

has ~90% genome-wide methylation (de Mendoza et al., 2018; Lohuis & Miller, 1998).

When compared to RRBS and WGBS data, we found a 4 to 10-fold enrichment of

Symbiodiniaceae DNA in *P. acuta* MBDBS data. Separation of host and symbionts is

therefore recommended to obtain the greatest read counts for the organism of interest,

but this comes at the cost of not being able to obtain RNA from the same nucleic acid

pool. For example, physical separation of the host and symbiont in living cells impacts

gene expression, and attempts at physical separation after freezing can degrade the host

RNA. Simultaneous extraction of holobiont RNA and DNA from the same nucleic acid pool

provides the optimal approach for detecting interactions between DNA methylation and

epigenetic regulation of gene expression. This comes at the cost of generating excess

reads to overcome highly methylated Symbiodiniaceae DNA.

MBDBS can enrich for gene regions in species where methylation is primarily found

in gene bodies such as in corals (reviewed in (Eirin-Lopez & Putnam, 2019)), and can thus

provide insight into mechanisms underlying physiological or organismal responses. We

found that MBDBS significantly enriched for gene bodies, specifically CDS and introns,

when compared to RRBS and WGBS in both *M. capitata* and *P. acuta* (**Figure 5**). While

MBDBS may be a good choice to examine gene body methylation at a reduced cost,

species differences in CpG coverage within orthologous genes with MBDBS (Additional

file 10: **Figure SF6**) may complicate cross-species comparisons by reducing the amount

of data available for analysis. Additionally, we found high discordance between MBDBS

and non-enrichment methods, WGBS and RRBS, for *P. acuta*. MBDBS is the only method

we evaluated that can non-randomly sub-sample genomes present in a DNA sample

through preferential pull-down of methylated DNA. Differences in methylation across the

sampled genomes could result from cell-to-cell heterogeneity in methylation or cell-type

(e.g., methylation of calcifying cells may differ from symbiont hosting cells). In other words,

MBDBS data may represent only a subpopulation of highly methylated cells, while WGBS

and RRBS represent the average methylation across all cells in the sample. Using a

consistent tissue type is important to limit potential methylation heterogeneity, and caution

should be taken when comparing MBDBS data directly to that of non-enrichment bisulfite

sequencing approaches.

Although MBDBS did enrich for methylated regions of the genome, 80% of CpGs

in *M. capitata* and 82% of CpGs in *P. acuta* interrogated with MBDBS were lowly

methylated (<10% methylated) (Additional file 8: **Figure SF3**). This is expected and is

consistent with previous reports applying MBDBS in other marine invertebrates where

unmethylated CpGs actually represent the highest proportion of loci in the data,

attributable to the nature of the methylation landscape and enrichment   protocol (e.g.

(Gavery & Roberts, 2013; Venkataraman et al., 2020)). The base pair resolution of

methylation revealed by MBDBS is a benefit over MBD-Seq alone because it enables a

fine-scale examination of specific genomic features (e.g. exon-intron boundaries) that may

not be possible with the regional resolution of MBD-Seq. Without complete knowledge of

the relative importance of a single loci compared to a region, it is difficult to compare trade-

offs between MBDBS and MBD-seq. However, bisulfite sequencing requires significant

coverage to quantify DNA methylation.

MBDBS may have potential biases that should be considered when interpreting

results. If a treatment, population comparison, or developmental change results in a given

region (~500bp) going from being highly methylated to fully unmethylated, then it is likely

that this region would not be interrogated by MBDBS, due to an absence of data in the

unmethylated condition. This is a potential source of bias in MBDBS data and may

contribute to important differentially methylated regions being overlooked: for example if

one treatment results in high methylation and is captured by MBDBS and another

treatment results in no methylation and is not captured by MBDBS, this region would be

filtered out of the analysis because of missing data in some individuals. Further, the

potential of MBDBS to provide limited information for unmethylated genes may introduce

bias in studies that seek to draw relationships between methylation level and gene

expression. Just as with many interpretations of key findings we present, a more complete

understanding of the mechanistic functional role DNA methylation plays in genome

regulation in the species of interest is needed.

There is a greater capacity to gain mechanistic insight when using methods that

have single base pair resolution compared to methylation enrichment without bisulfite

treatment or bulk percent methylation approaches. For example, hypotheses such as the

linkage between DNA methylation and alternative splicing (Roberts & Gavery, 2012) are

more accurately tested with bisulfite sequencing approaches. We acknowledge the cost

of generating genomic resources and bisulfite sequencing data can be higher than other

approaches. While WGBS is supported here as the gold standard for DNA methylation

quantification, consideration should be given to specific study hypotheses in light of the

pros and cons of the enrichment or reduced representation approaches presented here

and in other comparative works (Dixon & Matz, 2020). Our results suggest that it would

be unwise to use multiple different library preparation methods for comparing individuals

within a study, especially for studies in which familial relationships are to be compared. As

technology advances, it would be ideal to move away from harsh bisulfite conversion to

assess DNA methylation with single base pair resolution across whole genomes in the

absence of DNA treatment (e.g., Oxford Nanopore).

Our results provide a quantitative comparative assessment that can be used to

inform the choice of sequencing DNA methylation in corals and other non-model

invertebrates. Together these metrics enable comparative capacity for three common

methods in two coral taxa that vary in their phylogeny, genome size, symbiotic unions,

and environmental performance, and thus provide the community with a more

comprehensive foundation upon which to build laboratory and statistical analyses of DNA

methylation, plasticity, and acclimatization.

Abbreviations

CpG: cytosine and guanine separated by a phosphate group

ELISA: Enzyme-Linked Immunosorbent Assay

MBD-seq: Methyl-CpG Binding Domain Sequencing

MBDBS: Methyl-CpG Binding Domain Bisulfite Sequencing

RRBS: Reduced representation bisulfite sequencing

WGBS: Whole genome bisulfite sequencing

Acknowledgements

We thank Putnam Lab members E. Strand and M. E. Schedl for their technical support.

Data analyses were facilitated through the use of advanced computational, storage, and

networking infrastructure provided by the Hyak supercomputer system at the University of

Washington.

Funding

This work was partially supported by the USDA: National Institute of Food and Agriculture,

Hatch projects RI0019-H020 to HMP, NJ01180 to DB, and the NRSP-8 National Animal

Genome Research Program to HMP and SBR. This work was also partially supported by

funding from the National Science Foundation: 1756623 to HMP, 1756616 to DB, 1921149

to SBR, 1921402 to JEL, and 1635423 to KEL.

References

Akalin, A., Kormaksson, M., Li, S., Garrett-Bakelman, F. E., Figueroa, M. E., Melnick, A.,
& Mason, C. E. (2012). methylKit: a comprehensive R package for the analysis of
genome-wide DNA methylation profiles. *Genome Biology*, *13*(10), R87.
https://doi.org/10.1186/gb-2012-13-10-r87
Bahr, K. D., Jokiel, P. L., & Rodgers, K. S. (2016). Relative sensitivity of five Hawaiian

coral species to high temperature under high-pCO2 conditions. *Coral Reefs* , *35*(2),
729–738. https://doi.org/10.1007/s00338-016-1405-4
Baldwin, J. M. (1902). Development and evolution, including psychophysical evolution,
evolution by orthoplasy, and the theory of genetic modes.
https://doi.org/10.1037/13707-000
Barshis, D. J., Ladner, J. T., Oliver, T. A., Seneca, F. O., Traylor-Knowles, N., &
Palumbi, S. R. (2013). Genomic basis for coral resilience to climate change.
Proceedings of the National Academy of Sciences of the United States of America,
110(4), 1387–1392. https://doi.org/10.1073/pnas.1210224110
Byrne, M., Foo, S. A., Ross, P. M., & Putnam, H. M. (2020). Limitations of cross-and
multigenerational plasticity for marine invertebrates faced with global climate
change. *Global Change Biology*, *26*(1), 80–102. https://doi.org/10.1111/gcb.14882
Cunning, R., Ritson-Williams, R., & Gates, R. D. (2016). Patterns of bleaching and
recovery of *Montipora capitata* in Kāne 'ohe Bay, Hawai 'i, USA. *Marine Ecology*
Progress Series, *551*, 131–139. https://www.int-res.com/abstracts/meps/v551/p131-
139/
de Mendoza, A., Bonnet, A., Vargas-Landin, D. B., Ji, N., Li, H., Yang, F., Li, L., Hori, K.,
Pflueger, J., Buckberry, S., Ohta, H., Rosic, N., Lesage, P., Lin, S., & Lister, R.
(2018). Recurrent acquisition of cytosine methyltransferases into eukaryotic
retrotransposons. *Nature Communications*, *9*(1), 1–11.
https://doi.org/10.1038/s41467-018-03724-9
Dimond, J. L., Gamblewood, S. K., & Roberts, S. B. (2017). Genetic and epigenetic
insight into morphospecies in a reef coral. *Molecular Ecology*, *26*(19), 5031–5042.
https://doi.org/10.1111/mec.14252
Dimond, J. L., & Roberts, S. B. (2020). Convergence of DNA Methylation Profiles of the
Reef Coral *Porites astreoides* in a Novel Environment. *Frontiers in Marine Science*,
6, 792. https://doi.org/10.3389/fmars.2019.00792
Dixon, G., Liao, Y., Bay, L. K., & Matz, M. V. (2018). Role of gene body methylation in
acclimatization and adaptation in a basal metazoan. *Proceedings of the National*
Academy of Sciences of the United States of America, *115*(52), 13342–13346.
https://doi.org/10.1073/pnas.1813749115
Dixon, G., & Matz, M. (2020). Benchmarking DNA methylation assays in a reef-building
coral. *Molecular Ecology Resources*. https://doi.org/10.1111/1755-0998.13282
Eirin-Lopez, J. M., & Putnam, H. M. (2019). Marine Environmental Epigenetics. *Annual*
Review of Marine Science, *11*, 335–368. https://doi.org/10.1146/annurev-marine-
010318-095114
Ewels, P., Magnusson, M., Lundin, S., & Käller, M. (2016). MultiQC: summarize analysis
results for multiple tools and samples in a single report. *Bioinformatics* , *32*(19),
3047–3048. https://doi.org/10.1093/bioinformatics/btw354
Flores, K., Wolschin, F., Corneveaux, J. J., Allen, A. N., Huentelman, M. J., & Amdam,
G. V. (2012). Genome-wide association between DNA methylation and alternative
splicing in an invertebrate. *BMC Genomics*, *13*, 480. https://doi.org/10.1186/1471-
2164-13-480
Foret, S., Kucharski, R., Pellegrini, M., Feng, S., Jacobsen, S. E., Robinson, G. E., &
Maleszka, R. (2012). DNA methylation dynamics, metabolic fluxes, gene splicing,

and alternative phenotypes in honey bees. *Proceedings of the National Academy of*
Sciences of the United States of America, *109*(13), 4968–4973.
https://doi.org/10.1073/pnas.1202392109
Gavery, M. R., & Roberts, S. B. (2010). DNA methylation patterns provide insight into
epigenetic regulation in the Pacific oyster (*Crassostrea gigas*). *BMC Genomics*,
11(1), 483. https://doi.org/10.1186/1471-2164-11-483
Gavery, M. R., & Roberts, S. B. (2013). Predominant intragenic methylation is
associated with gene expression characteristics in a bivalve mollusc. *PeerJ*, *1*,
e215. https://doi.org/10.7717/peerj.215
Ghalambor, C. K., Hoke, K. L., Ruell, E. W., Fischer, E. K., Reznick, D. N., & Hughes, K.
A. (2015). Non-adaptive plasticity potentiates rapid adaptive evolution of gene
expression in nature. *Nature*, *525*(7569), 372–375.
https://doi.org/10.1038/nature15256
Ghalambor, C. K., McKay, J. K., Carroll, S. P., & Reznick, D. N. (2007). Adaptive versus
non-adaptive phenotypic plasticity and the potential for contemporary adaptation in
new environments. *Functional Ecology*, *21*(3), 394–407.
https://besjournals.onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-
2435.2007.01283.x
Gibbin, E. M., Putnam, H. M., Gates, R. D., Nitschke, M. R., & Davy, S. K. (2015).
Species-specific differences in thermal tolerance may define susceptibility to
intracellular acidosis in reef corals. *Marine Biology*, *162*(3), 717–723.
https://doi.org/10.1007/s00227-015-2617-9
Gómez-Rubio, V. (2017). ggplot2 - Elegant Graphics for Data Analysis (2nd Edition). In
Journal of Statistical Software (Vol. 77, Issue Book Review 2).
https://doi.org/10.18637/jss.v077.b02
Gonzalez-Romero, R., Suarez-Ulloa, V., Rodriguez-Casariego, J., Garcia-Souto, D.,
Diaz, G., Smith, A., Pasantes, J. J., Rand, G., & Eirin-Lopez, J. M. (2017). Effects of
Florida Red Tides on histone variant expression and DNA methylation in the
Eastern oyster *Crassostrea virginica*. *Aquatic Toxicology* , *186*, 196–204.
https://doi.org/10.1016/j.aquatox.2017.03.006
Gu, H., Smith, Z. D., Bock, C., Boyle, P., Gnirke, A., & Meissner, A. (2011). Preparation
of reduced representation bisulfite sequencing libraries for genome-scale DNA
methylation profiling. *Nature Protocols*, *6*(4), 468–481.
https://doi.org/10.1038/nprot.2010.190
Hofmann, G. E. (2017). Ecological Epigenetics in Marine Metazoans. *Frontiers in Marine*
Science, *4*, 4. https://doi.org/10.3389/fmars.2017.00004
Hu, M., Zheng, X., Fan, C.-M., & Zheng, Y. (2020). Lineage dynamics of the
endosymbiotic cell type in the soft coral *Xenia*. In *Nature* (Vol. 582, Issue 7813, pp.
534–538). https://doi.org/10.1038/s41586-020-2385-7
Johnson, K. M., Sirovy, K. A., Casas, S. M., La Peyre, J. F., & Kelly, M. W. (2020).
Characterizing the Epigenetic and Transcriptomic Responses to *Perkinsus marinus*
Infection in the Eastern Oyster *Crassostrea virginica*. *Frontiers in Marine Science*, *7*.
https://doi.org/10.3389/fmars.2020.00598
Kronholm, I., & Collins, S. (2016). Epigenetic mutations can both help and hinder
adaptive evolution. *Molecular Ecology*, *25*(8), 1856–1868.

https://doi.org/10.1111/mec.13296

Krueger, F. (2012). Trim Galore: a wrapper tool around Cutadapt and FastQC to
consistently apply quality and adapter trimming to FastQ files, with some extra
functionality for MspI-digested RRBS-type (Reduced Representation Bisufite-Seq)
libraries. *URL Http://www. Bioinformatics. Babraham. Ac. Uk/projects/trim_galore/.*
(Date of Access: 28/04/2016).

Krueger, F., & Andrews, S. R. (2011). Bismark: a flexible aligner and methylation caller
for Bisulfite-Seq applications. *Bioinformatics* , *27*(11), 1571–1572.
https://doi.org/10.1093/bioinformatics/btr167

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2.
Nature Methods, *9*(4), 357–359. https://doi.org/10.1038/nmeth.1923

Lewis, S. L., & Maslin, M. A. (2015). Defining the anthropocene. *Nature*, *519*(7542),
171–180. https://doi.org/10.1038/nature14258

Liew, Y. J., Howells, E. J., Wang, X., Michell, C. T., Burt, J. A., Idaghdour, Y., & Aranda,
M. (2020). Intergenerational epigenetic inheritance in reef-building corals. *Nature*
Climate Change, *10*(3), 254–259. https://doi.org/10.1038/s41558-019-0687-2

Liew, Y. J., Zoccola, D., Li, Y., Tambutté, E., Venn, A. A., Michell, C. T., Cui, G.,
Deutekom, E. S., Kaandorp, J. A., Voolstra, C. R., Forêt, S., Allemand, D.,
Tambutté, S., & Aranda, M. (2018). Epigenome-associated phenotypic
acclimatization to ocean acidification in a reef-building coral. *Science Advances*,
4(6), 188227. https://doi.org/10.1126/sciadv.aar8028

Liu, H., Stephens, T. G., González-Pech, R. A., Beltran, V. H., Lapeyre, B., Bongaerts,
P., Cooke, I., Aranda, M., Bourne, D. G., Forêt, S., Miller, D. J., van Oppen, M. J.
H., Voolstra, C. R., Ragan, M. A., & Chan, C. X. (2018). Symbiodinium genomes
reveal adaptive evolution of functions related to coral-dinoflagellate symbiosis.
Communications Biology, *1*, 95. https://doi.org/10.1038/s42003-018-0098-3

Li, Y., Liew, Y. J., Cui, G., Cziesielski, M. J., Zahran, N., Michell, C. T., Voolstra, C. R., &
Aranda Lastra, M. (2018). DNA methylation regulates transcriptional homeostasis of
algal endosymbiosis in the coral model *Aiptasia. Sci Adv, 4*.

Lohuis, M. R., & Miller, D. J. (1998). Hypermethylation at CpG-Motifs in the
dinoflagellates *Amphidinium carterae* (*Dinophyceae*): Evidence from restriction
analyzes, 5-azacytidine and ethionine treatment. In *Journal of Phycology* (Vol. 34,
Issue 1, pp. 152–159). https://doi.org/10.1046/j.1529-8817.1998.340152.x

Meissner, A., Mikkelsen, T. S., Gu, H., Wernig, M., Hanna, J., Sivachenko, A., Zhang,
X., Bernstein, B. E., Nusbaum, C., Jaffe, D. B., Gnirke, A., Jaenisch, R., & Lander,
E. S. (2008). Genome-scale DNA methylation maps of pluripotent and differentiated
cells. *Nature*, *454*(7205), 766–770. https://doi.org/10.1038/nature07107

Oury, N., Gélin, P., & Magalon, H. (2020). Together stronger: Intracolonial genetic
variability occurrence in *Pocillopora* corals suggests potential benefits. *Ecology and*
Evolution, *10*(12), 5208–5218. https://doi.org/10.1002/ece3.5807

Putnam, H. M., Barott, K. L., Ainsworth, T. D., & Gates, R. D. (2017). The Vulnerability
and Resilience of Reef-Building Corals. *Current Biology: CB*, *27*(11), R528–R540.
https://doi.org/10.1016/j.cub.2017.04.047

Putnam, H. M., Davidson, J. M., & Gates, R. D. (2016). Ocean acidification influences
host DNA methylation and phenotypic plasticity in environmentally susceptible

corals. *Evolutionary Applications*, *9*(9), 1165–1178.
https://doi.org/10.1111/eva.12408
Putnam, H. M., Roberts, S. B., & Venkataraman, Y. (2020). *Coral Methylation Methods*
Comparison. Open Science Framework. https://doi.org/10.17605/OSF.IO/X5WAZ
Quinlan, A. R., & Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics* , *26*(6), 841–842.
https://doi.org/10.1093/bioinformatics/btq033
Rice, P. M., Bleasby, A. J., & Ison, J. C. (2011). *EMBOSS User's Guide: Practical*
Bioinformatics with EMBOSS. Cambridge University Press.
https://play.google.com/store/books/details?id=zC5WGJxDNtcC
Riley, M., Abe, T., Arnaud, M. B., Berlyn, M. K. B., Blattner, F. R., Chaudhuri, R. R.,
Glasner, J. D., Horiuchi, T., Keseler, I. M., Kosuge, T., Mori, H., Perna, N. T.,
Plunkett, G., 3rd, Rudd, K. E., Serres, M. H., Thomas, G. H., Thomson, N. R.,
Wishart, D., & Wanner, B. L. (2006). *Escherichia coli* K-12: a cooperatively
developed annotation snapshot--2005. *Nucleic Acids Research*, *34*(1), 1–9.
https://doi.org/10.1093/nar/gkj405
Ritz, C., Baty, F., Streibig, J. C., & Gerhard, D. (2015). Dose-Response Analysis Using
R. *PloS One*, *10*(12), e0146021. https://doi.org/10.1371/journal.pone.0146021
Roberts, S. B., & Gavery, M. R. (2012). Is There a Relationship between DNA
Methylation and Phenotypic Plasticity in Invertebrates? *Frontiers in Physiology*, *2*,
116. https://doi.org/10.3389/fphys.2011.00116
Rodrigues, L. J., & Grottoli, A. G. (2007). Energy reserves and metabolism as indicators
of coral recovery from bleaching. *Limnology and Oceanography*, *52*(5), 1874–1882.
https://doi.org/10.4319/lo.2007.52.5.1874
Rodriguez-Casariego, J. A., Ladd, M. C., Shantz, A. A., Lopes, C., Cheema, M. S., Kim,
B., Roberts, S. B., Fourqurean, J. W., Ausio, J., Burkepile, D. E., & Eirin-Lopez, J.
M. (2018). Coral epigenetic responses to nutrient stress: Histone H2A.X
phosphorylation dynamics and DNA methylation in the staghorn coral *Acropora*
cervicornis. *Ecology and Evolution*, *8*(23), 12193–12207.
https://doi.org/10.1002/ece3.4678
Rosental, B., Kozhekbaeva, Z., Fernhoff, N., Tsai, J. M., & Traylor-Knowles, N. (2017).
Coral cell separation and isolation by fluorescence-activated cell sorting (FACS). In
BMC Cell Biology (Vol. 18, Issue 1). https://doi.org/10.1186/s12860-017-0146-8
Schweinsberg, M., Weiss, L. C., Striewski, S., Tollrian, R., & Lampert, K. P. (2015). More
than one genotype: how common is intracolonial genetic variability in scleractinian
corals? *Molecular Ecology*, *24*(11), 2673–2685. https://doi.org/10.1111/mec.13200
Shumaker, A., Putnam, H. M., Qiu, H., Price, D. C., Ehud, Z., Harel, A., Wagner, N. E.,
Gates, R. D., Yoon, H. S., & Bhattacharya, D. (2019). Genome analysis of the rice
coral *Montipora capitata*. *Scientific Reports*, *9*(2571).
https://doi.org/10.1038/s41598-019-39274-3
Suarez-Ulloa, V., Rivera-Casas, C., Michel, M., & Eirin-Lopez, J. M. (2018). Seasonal
variation and epigenetic changes in the flat tree oyster *Isognomon alatus* during a 2-
year study of a mangrove ecosystem in North Biscayne Bay, Florida. *Journal of*
Shellfish Research, 38(1):79-88 (2019). https://doi.org/10.2983/035.038.0108
Suzuki, M. M., Kerr, A. R. W., De Sousa, D., & Bird, A. (2007). CpG methylation is

targeted to transcription units in an invertebrate genome. *Genome Research, 17*(5),
625–631. https://doi.org/10.1101/gr.6163007
Thorvaldsdóttir, H., Robinson, J. T., & Mesirov, J. P. (2013). Integrative Genomics
Viewer (IGV): high-performance genomics data visualization and exploration.
Briefings in Bioinformatics, 14(2), 178–192. https://doi.org/10.1093/bib/bbs017
Tweedie, S., Charlton, J., Clark, V., & Bird, A. (1997). Methylation of genomes and
genes at the invertebrate-vertebrate boundary. *Molecular and Cellular Biology*,
17(3), 1469–1475. https://www.ncbi.nlm.nih.gov/pubmed/9032274
van Gurp, T. P., Wagemaker, N. C. A. M., Wouters, B., Vergeer, P., Ouborg, J. N. J., &
Verhoeven, K. J. F. (2016). epiGBS: reference-free reduced representation bisulfite
sequencing. *Nature Methods, 13*(4), 322–324. https://doi.org/10.1038/nmeth.3763
Velotta, J. P., Ivy, C. M., Wolf, C. J., Scott, G. R., & Cheviron, Z. A. (2018). Maladaptive
phenotypic plasticity in cardiac muscle growth is suppressed in high-altitude deer
mice. *Evolution; International Journal of Organic Evolution, 72*(12), 2712–2727.
https://doi.org/10.1111/evo.13626
Venkataraman, Y. R., Downey-Wall, A. M., Ries, J., Westfield, I., White, S. J., Roberts,
S. B., & Lotterhos, K. E. (2020). General DNA methylation patterns and
environmentally-induced differential methylation in the eastern oyster (*Crassostrea*
virginica). *Frontiers in Marine Science*. https://doi.org/10.3389/fmars.2020.00225
Vidal-Dupiol, J., Chaparro, C., Pratlong, M., Pontaroti, P., Grunau, C., & Mitta, G. (2019).
Sequencing, de novo assembly and annotation of the genome of the scleractinian
coral, *Pocillopora acuta*. In *bioRxiv* (p. 698688). https://doi.org/10.1101/698688
Zemach, A., McDaniel, I. E., Silva, P., & Zilberman, D. (2010). Genome-wide
evolutionary analysis of eukaryotic DNA methylation. *Science, 328*(5980), 916–919.
https://doi.org/10.1126/science.1186366

Data accessibility

The datasets supporting the conclusions of this article are available in the Coral Methylation Methods Comparison repository, http://doi.org/10.17605/OSF.IO/X5WAZ, and included included within the article (and its additional file(s)). All raw data can be accessed under NCBI Bioproject PRJNA691891.

Author contributions

HMP designed the study and collected the samples. YRV, SAT, MRG, HMP, and SBR performed data analyses and drafted the manuscript. DB, AD-W, JME-L, KMJ, KEL, and JBP provided analytical insight and manuscript revisions. All authors read and approved the final manuscript.

Ethics declarations

Ethics approval and consent to participate

The coral samples were collected under Hawai'i Department of Aquatic Resources Special Activity Permit SAP 2019-60.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Figures and Tables

Figure 1. Experimental design. Three biological replicate coral samples were obtained from both coral species. A) *M. capitata*, where B) a cross section of a decalcified fragment reveals thick tissue, and C) a perforate tissue skeletal interaction. In contrast in D) *P. acuta*, a E) a cross section of a decalcified fragment reveals thin tissue, and F) an imperforate tissue skeletal interaction. DNA was extracted from each coral sample and split for use in Whole Genome Bisulfite Sequencing (WGBS), Reduced Representation Bisulfite Sequencing (RRBS), and Methyl-CpG Binding Domain Bisulfite Sequencing (MBDBS) library preparation methods. Three libraries were generated for each of the three methods, yielding nine libraries for each species and 18 libraries total.

Figure 2. **Mean percent methylation of CpGs**. Data is presented for CpGs with 5x coverage for each method on the largest scaffolds of each genome. The outer track shows the scaffold locations and dots indicate the percent methylation as indicated by the y-axes from 0-100% for each of the inner tracks.

Figure 3. Matrix of pairwise scatter plots for shared CpG loci. Data is presented for CpG covered at ≥ 5x across all samples) for A) *M. capitata* (n=4,666 common loci) and B) *P. acuta* (n=93,714 common loci). The red lines represent linear regression fits and the

green lines are polynomial regression fits. Pearson correlation coefficients for each pairwise comparison are presented in the upper right boxes. Methods are color coded on the X and Y axes (WGBS = green, MBDBS = purple, and RRBS = orange) and replicate samples are indicated on the diagonal along with histograms of % CpG methylation.

Figure 4. CpG site coverage across library preparation methods. Mean fraction of CpG sites in the genome covered at different sequencing depths (read depths) by (**A**) MBDBS libraries, (**B**) RRBS libraries, and (**C**) WGBS libraries with standard deviations shown by shaded areas (see Additional file 2: **Table ST2** for number of reads in each sample).

Figure 5. Percent of CpGs detected by sequencing methods in genome features A) for *M. capitata* and B) *P. acuta*. Genome features considered were coding sequences

(CDS), introns, 1 Kb flanking regions upstream (Upstream Flank) or downstream of genes (Downstream Flank), and intergenic regions Principal Coordinate Analyses associated with PERMANOVA and beta-dispersion tests related to Additional file 12: **Table ST6** that show differences in proportion of CpGs in various genomic locations (CDS, introns, upstream flanks, downstream flanks, and intergenic regions) for **C**) M. capitata and **D**) P. acuta. WGBS is represented by green circles, RRBS by purple triangles, and MBDBS by orange diamonds. Percent variation explained by each PCoA axis is included in the axis label. Ellipses depict 95% confidence intervals for each sequencing method. All eigenvectors are significant at the $\alpha = 0.05$ level.

Supplementary Information