A MALDI-TOF assay identifies nilotinib as an inhibitor of inflammation in acute myeloid

leukemia and multiple myeloma.

Rachel E. Heap[1,#], José Luis Marín-Rubio[1,#,*], Tiaan Heunis[1,2], Maria Emilia Dueñas[1], Joseph

Inns[1], Barbara Saxty[3] and Matthias Trost[1,*].

[1] Laboratory for Biological Mass Spectrometry, Biosciences Institute, Newcastle University,

Newcastle-upon-Tyne, NE2 4HH, UK

[2]Sir William Dunn School of Pathology, University of Oxford, South Parks Road, Oxford OX1

3RE, UK

[3]LifeArc, SBC Open Innovation Campus, Stevenage, SG1 2FX, UK

[#]These authors contributed equally to this work.

* Corresponding author:

Matthias Trost, Tel: +44 191 2087009, E-mail: matthias.trost@ncl.ac.uk

José Luis Marín-Rubio, Tel +44 191 2088983 E-mail: jose.marin-rubio@newcastle.ac.uk

Key Points

Label-free cell-based assay identifies new anti-inflammatory drugs using MALDI-TOF MS.

Nilotinib reduces inflammation by inhibition of MAPK14 in AML and MM.

Topics: MALDI-TOF, nilotinib, inflammation, myeloma, leukemia, high-throughput drug

discovery

ABSTRACT

Inflammatory responses are important in cancer, particularly in the context of monocyte-rich aggressive myeloid neoplasm. We developed a label-free cellular phenotypic drug discovery assay to identify anti-inflammatory drugs in human monocytes derived from acute myeloid leukemia (AML), by tracking several biological features ionizing from only 2,500 cells using matrix-assisted laser desorption/ionization-time of flight (MALDI-TOF) mass spectrometry. A proof-of-concept screen showed that the BCR-ABL inhibitor nilotinib, but not the structurally similar imatinib, blocks inflammatory responses. In order to identify the cellular (off-)targets of nilotinib, we performed thermal proteome profiling (TPP) using TMTpro 16plex. Unlike imatinib, nilotinib inhibits p38-alpha (MAPK14) signalling. This inhibition suppressed the expression of inflammatory cytokines, cell adhesion and innate immunity markers in activated human monocytes derived from AML. Thus, our study provides a tool for the discovery of new anti-inflammatory drugs, which could contribute to the treatment of inflammation in myeloid neoplasms and other diseases.

INTRODUCTION

Monocyte-derived macrophages play important roles in both physiological and pathological processes. The activation of these cells during pathogen infection or other insults leads, in part, to the pathophysiology of inflammation.[1] Monocytes can recognize various damage-associated molecular (DAMPs) and pathogen-associated molecular patterns (PAMPs) through specific receptors, such as Toll-like receptors (TLRs). Activation of these receptors leads to the production of cytokines, chemokines, and mediators that are involved in inflammation in myeloid leukemia.[2,3] Several hematopoietic disorders, including lymphoproliferative disorders and myelodysplastic syndromes, which possess a high risk of transformation to leukemia, have been linked to aberrant TLR signalling.[4,5] Moreover, TLR2 and TLR4 show significantly higher expression in the bone marrow of patients with myeloid leukemia.[5] While TLRs are specific for exogenous PAMPs such as lipopolysaccharide (LPS) which triggers TLR4 signalling in monocytes and macrophages, they are also activated by specific endogenous DAMPs such as HSP60 or biglycan.[6,7] TLR activation leads to intracellular signalling cascades including the mitogen-activated kinase (MAP) pathways,[8,9] increasing the expression and secretion of inflammatory cytokines and chemokines including interleukins (IL-6, IL-8 and IL-10), interferons (IFN), and tumor necrosis factor alpha (TNF-α).[10]

The discovery of drugs that prevent chronic or acute inflammation is a major goal of the pharmaceutical industry for the treatment of infectious diseases, autoimmune diseases, and cancer.[11] The discovery and design of new compounds for inhibiting inflammation are typically achieved through high-throughput screening (HTS) approaches, however, fluorescence and chemiluminescence methodologies cannot be used in primary cells and

other read-outs, including antibody-based assays such as ELISA, are very expensive.[12,13] As

such, there is a growing need to develop physiological phenotypic HTS assays that can be

used in monocytic cells to accelerate the discovery of inhibitors.[14,10z5] Matrix-assisted laser

desorption/ionization-time of flight (MALDI-TOF) mass spectrometry (MS) is a versatile,

label-free technique that has the potential to accelerate HTS of promising drug

candidates.[16-18] In this study, we developed a MALDI-TOF MS cellular assay to perform a

blind screen of 96 compounds to assess their potential anti-inflammatory effects on human

monocytes derived from acute myeloid leukemia (AML). We discovered that nilotinib, but

not the structurally related imatinib, inhibits inflammatory signalling. Using Thermal

Proteome Profiling (TPP), we can show that nilotinib blocks the p38α MAP kinase pathway.

Overall, our data show that nilotinib is a drug capable of preventing inflammation in AML

and multiple myeloma (MM).

RESULTS

A novel MALDI-TOF MS cellular assay to characterize inflammatory phenotypes

Recently, high-throughput MALDI-TOF MS has been used successfully for *in vitro* assays of

specific enzymes. Here, we tested if MALDI-TOF MS can be used for phenotypic assays to

monitor inflammatory responses by identifying differences in the "fingerprint" of

biomolecules ionized when whole cells are spotted onto the MALDI target (**Figure 1A**). As a

proof of concept, we tested if we could distinguish phenotypes of THP-1 cells, a model of

human monocytes derived from AML,[19] upon stimulation with a number of pro-

inflammatory stimuli, namely LPS, Pam2CSK4, Pam3CSK4; which activate TLR4, TLR1/2 and

TLR2/6, respectively, as well as interferon-γ (IFN-γ). Principal component analysis (PCA) of all

biomolecules detected in the range of *m/z* 2,000-20,000 showed clear separation of

monocytes treated with TLR-agonists, while IFN-γ treatment did not separate from

untreated, suggesting that the features are not interferon-stimulation dependent but rather

TLR dependent (**Figure 1B**). A loading plot analysis of the PCA results identified the two

features at *m/z* 4632 and 4964 as the main drivers for the difference in the phenotypes

(**Figure 1C**). The feature *m/z* 4632 was significantly reduced while *m/z* 4964 increased in

LPS-stimulated cells (**Figure 1D-E**). We confirmed that other TLR ligands such as Pam2CSK4

and Pam3CSK4, which activate TLR1/2 and TLR2/6, respectively, did also increase this ratio

and in a dose-dependent manner, while interferon-gamma (IFN-γ) did not affect it (**Figure**

1F-G). Thus, we used the feature at *m/z* 4632 as a resting phenotype biomarker and feature

m/z 4964 as an LPS-stimulated monocyte phenotype biomarker. To verify that this

phenotype was not due to cell death, we treated the cells with 0.5 µM staurosporine for

24h before analysis with our MALDI-TOF assay. Our results showed that apoptosis leads to a

distinct cluster in the PCA compared to control or LPS-treated cells (**Figure 1H**), suggesting

that this assay can be multiplexed to identify other phenotypes such as cell toxicity.

Together, these results indicate that biomarkers identified by MALDI-TOF MS can be used to

detect changes in the inflammatory phenotype downstream of TLRs in AML cells.

Figure 1. Identification of features associated with the monocyte inflammatory phenotype by MALDI-TOF MS. A) Workflow of the MALDI-TOF MS assay: (1) AML cells were pre-treated with a drug for 1h before adding a pro-inflammatory stimulus for 24h. (2) Cells were frozen on dry ice, thawed and (3) washed with 100 mM Tris-HCl buffer at 4°C. (4) Cells were mixed with a matrix (10 mg/mL α-cyano-4-cinnamic acid in 50% acetonitrile, 0.1% trifluoroacetic acid). (5) Cells were analysed in a RapifleX PharmaPulse MALDI TOF mass spectrometer. (6 In the ionization chamber a laser is used to produce ions in the gas phase. These ions are separated according to their time-of-free flight (TOF: Time-Of-Flight). The smaller molecules reach the detector first, followed by the bigger, according to the *m/z* ratio. (7) The detector converts the received ions into electrical current which is amplified and digitized in the m/z spectra. **B)** Unsupervised PCA plot of LPS, Pam2CSK4, Pam3CSK4 and IFN-γ-treated cells showing separation of cells treated with bacterial ligands treated and resting monocytes. **C)** Loading plot derived from PCA in B) showing that m/z 4964 and 4632 contribute predominantly to the separation of the two clusters in component 1 and component 2. **D)** Unsupervised heat map of the relative intensities of three biological replicates of THP-1 cells treated with 100 ng/mL LPS, Pam2CSK3, Pam2CSK4 and 100 U/mL IFN-γ for 24h compared with resting cells **E)** Box plots of significantly changing intensities between resting and LPS-treated monocytes identified at *m/z* 4632 and 4964. **F)** Relative quantitation from three biological replicates of THP-1 cells treated with 100 ng/mL LPS, Pam2CSK3, Pam2CSK4 and 100 U/mL IFN-γ for 24h compared with resting cells. **G)** Titration of LPS, Pam2CSK4 and Pam3CSK4-treated cells from 10 - 100 ng/mL of stimulus. **H)** PCA plot with three dimensions represents resting, LPS-, and 0.5 μM staurosporine-treated cells. Significant differences between two groups were determined by a Students t-test. The statistical significance of the comparisons with resting is indicated as follows: ns, not significant; ***, $P \leq 0.001$. Error bars represent the SEM.

The MALDI-TOF MS cellular assay can detect inhibitors of inflammation

Next, we used the TAK1 inhibitor NG-25, which blocks TNF-α production, the TBK1 inhibitor

MRT67303, which blocks production of type I interferons (IFNs),[20,21] as well as BI2536, a

PLK1 inhibitor that has been show to inhibit pro-inflammatory gene transcription due to

inhibition of BET proteins[22], to test if pharmacological inhibition of these pathways can be

detected in our assay (**Figure 2A**). THP-1 cells were pre-treated with the inhibitors for one

hour before LPS-stimulation. NG-25 and BI2536 treatment, but not MRT67303, reduced

TNF-α secretion (**Figure 2B**). Comparably, our MALDI-TOF MS assay showed that treatment

with NG-25 or BI2536 blocked the inflammatory phenotype and was indistinguishable to

vehicle control cells not treated with LPS, while MRT67307-treated cells still showed an

increased ratio at *m/z* 4964/4632, similar to LPS-treated cells (**Figure 2C**). Moreover, using

all biomolecule features, PCA analysis showed that treatment with NG-25 and BI2536

reverted the features back close to control, while MRT67307 did not affect LPS-induced

features significantly (**Figure 2D**). This data shows that pharmacological intervention of

inflammatory pathways can be detected by our assay and that NG-25 and MRT67307 can be

used as positive and negative control compounds in high-throughput screens, respectively.

Label-free MALDI-TOF MS identifies nilotinib as an anti-inflammatory compound

Once the analytical workflow was established to identify an inflammatory phenotype in

THP1 cells, we evaluated the potential of the assay to discriminate positive and negative

inhibitors of inflammation in a proof-of-concept blind screen of 96 selected kinase inhibitors

(**Supplementary Table 1).** The assay showed very good sensitivity (z'=~0.8) and

reproducibility (R^2 > 0.8**) (Supplementary Figure 1A-B**). Only two positive hits were observed

when applying a cut-off of 40% to the ratio *m/z* 4964/4632. Unblinding revealed that these

compounds were NG-25 and nilotinib (Tasigna®) (**Figure 2E**). Nilotinib is a BCR-ABL tyrosine

kinase inhibitor[23,24] and is a second-generation derivative of imatinib (Gleevec®),[24] which

was also included in the panel screened. However, imatinib did not affect the phenotype of

inflammatory monocytes, while nilotinib-treated monocytes did not display an LPS-induced

inflammatory phenotype and showed no differences compared to non-activated monocytes

in the ratio of *m/z* 4964/4632 (**Figure 2F**).

In order to validate the MALDI-TOF MS results, we measured secreted TNF-α levels in THP-1

cells treated with each inhibitor. Nilotinib treatment reduced expression and secretion

levels of TNF-α, while imatinib did not affect TNF-α levels upon LPS treatment (**Figure 2G-H**).

Both MALDI-TOF MS and TNF-α ELISA data provided a similar IC50 of approximately 400-500

nM for the inhibitory effect of Nilotinib (**Supplementary Figure 1C-D**). This data indicates

that nilotinib is able to block inflammatory responses downstream of TLR activation, while

the structurally similar imatinib cannot.

Figure 2. Nilotinib is a positive hit in a cellular MALDI-TOF MS drug discovery screen and inhibits

inflammation in response to LPS. A) Inhibition of the TLR2 and TLR4 signalling pathways. MRT67307

inhibits TBK1 (TANK Binding Kinase 1) which blocks production of type I interferons (IFNs). NG-25

inhibits TAK1 (TGF-Beta-Activated Kinase 1) which induces TNF-α and IL-6 production. BI2536 inhibits

BET proteins (Bromodomain and Extra-Terminal motif proteins) which are required for pro-

inflammatory gene transcription. **B)** TNF-α secretion of THP-1 cells treated in either vehicle control-

treated (DMSO), 100 ng/mL LPS-treated, or pre-treated with 5 µM NG-25, BI2536 or MRT67307 for

one hour before 100 ng/mL LPS-treatment for up to 24h measured by ELISA. **C)** MALDI-TOF MS

relative quantitation of the ratio *m/z* 4964/4632 in THP-1 cells as treated in B). **D)** PCA of all features

identified by the MALDI-TOF MS of experiment in B). **E)** Compound hit map of the mean of three

biological replicates with 40% effectiveness cut off showing nilotinib and NG-25 as positive hits and

imatinib and MRT67307 as negative hits (arrows). **F)** MALDI-TOF MS relative quantitation of the ratio

m/z 4964/4632 in THP-1 cells in vehicle control-treated (DMSO), 100 ng/mL LPS-treated, or pre-

treated with 5 µM nilotinib or imatinib for one hour before 100 ng/mL LPS-treatment for up to 24h.

G) *TNF-α* expression of cells treated as in F.) determined by RT-qPCR at 24 h. The results were

analysed using the $2^{-\Delta\Delta Ct}$ method and normalized using *GAPDH* and *TBP* as the reference genes and

vehicle control (DMSO) as the reference sample. Significant differences between two groups were

determined by a Students t-test. **H)** TNF-α secretion of cells treated as in F.) measured by ELISA after

24h. The statistical significance of the comparisons with LPS is indicated as follows: ns, not

significant; *, $P \leq 0.05$; **, $P \leq 0.01$; ***, $P \leq 0.001$. Error bars represent the SEM of three (**B, F**) or

four (**C, G, H**) biological replicates.

Nilotinib blocks the inflammatory response by inhibiting the p38α MAP kinase pathway.

Nilotinib is known to inhibit BCR-ABL and other tyrosine kinases.[24] A putative role of these

kinases in inflammatory signalling downstream of TLRs is unknown. Moreover, the kinase

potentially involved in this particular function must be a specific nilotinib target, since the

treatment with imatinib did not result in the same phenotype. In order to identify the off-

targets responsible for the nilotinib-specific anti-inflammatory phenotype, we performed

thermal proteome profiling (TPP) by multiplexed quantitative mass spectrometry using

tandem mass tags (TMT) (**Figure 3A, Supplementary Table 2, and Supplementary Table 3)**.

This method is based on determining the changes in the thermal stability of proteins, which

may be due to direct drug binding, drug-induced conformation changes, binding to other

cellular components, or post-translational modifications such as phosphorylation.[25-29] We

pre-treated THP1 cells for 1 h with nilotinib or imatinib before stimulation with LPS for 15

min. We reduced the LPS treatment to 15 min in order to avoid changes in protein

abundance due to transcriptional responses to LPS activation. Overall, we identified 5,565

proteins in our TPP analysis (**Supplementary Table 4**). Only seven proteins changed

significantly in all four replicates in melting temperature between nilotinib and imatinib with

a standard deviation below two degrees. For three of these proteins, we detected increased

ΔT_m in nilotinib-treated samples. The strongest hit was MAP kinase p38α (MAPK14) with an

increased ΔT_m of 6°C in nilotinib-treated samples (**Figure 3B**). The closely related isoform

p38γ (MAPK12) did not show significant differences (**Supplementary Figure 2A**). Also, MK3,

a downstream protein of p38, showed an increased ΔT_m of 2.5°C in nilotinib-treated samples

(**Figure 3C**). The remainder of significant proteins are shown in the **Supplementary Figure**

2B-F. To validate p38 alpha, we performed western blots of lysates from a cellular thermal

shift assay. The thermal denaturation temperatures for p38 alpha in DMSO and LPS-treated

monocytes were 50.3°C and 50.2°C, respectively. In accordance with the TPP data, the

thermal denaturation temperatures for p38 with nilotinib was 60.7°C, while with imatinib it

was 50.8°C (**Figure 3D-E**). We further performed a concentration compound range

experiment at 56ºC to determine the amounts of nilotinib required to alter protein thermal

stability of p38. We found that nilotinib was able to stabilize p38 at an EC50 concentration

of 4.6 µM (**Supplementary Figure 2G-H**).

Figure 3. Thermal proteome profiling (TPP) reveals stabilization of p38 alpha and MK3 in nilotinib-
treated compared to imatinib-treated cells. A) Workflow of thermal proteome profiling (TPP):
(1) AML cells were pre-treated with drug for 1h before adding LPS for 15 min. (2) Cells were
split in 8 different tubes and heated for 3 min at a range of temperatures (40, 44, 48, 52, 56,
60, 64 and 68°C). Cells were lysed by freeze-thawing and precipitates removed by
ultracentrifugation. (3) Proteins in the supernatants were digested with trypsin, labelled
with 16-plex tandem mass tags (TMT) and (4) mixed. TMT-labelled peptides were
fractionated using high-performance liquid chromatography (HPLC). (5) Peptides were
analysed in an Orbitrap Fusion Lumos Tribrid mass spectrometer. (6) TPP data analysis was
performed using MaxQuant and the computing environment R. (7) To validate the results,
target proteins were analysed by western blot and melting temperatures (Tm) were
calculated. **B)** Determination of the thermostability of p38 alpha (MAPK14), and **C)** MK3
(MAPKAPK3) at the indicated temperatures with 5 µM nilotinib or imatinib for one hour
before 100 ng/mL LPS-treatment for up to 15 min by TPP analysis. Table insert shows
melting temperature (Tm ºC) and R^2. **D)** Determination of the thermostability of p38 alpha
at the indicated temperatures in vehicle control (DMSO), 100 ng/mL LPS and pre-treated
with 5 µM nilotinib or imatinib for one hour before 100 ng/mL LPS-treatment for up to 15
min. GAPDH served as a loading control. **E)** Quantification of thermostability of p38 MAPK
western blots from four independent experiments. Table insert shows melting temperature
(Tm ºC) and R^2. Error bars represent the SEM of four biological replicates. A representative
image of four replicates is shown. Relative mobilities of reference proteins (masses in kilo
Daltons) are shown on the left of each blot.
240

In order to further understand the effect of nilotinib in preventing the inflammatory
response by TLR activation in human monocytes, we used *E. coli* to stimulate and induce an
inflammatory response in THP-1 cells. We used live *E. coli* rather than LPS to activate
multiple inflammatory pathways and not only the pathways downstream of TLR4. Similar to
LPS treatment, we were able to identify the inflammatory phenotype after exposure to *E.*
coli for 24h using the MALDI-TOF MS assay (**Figure 4A**). We observed a reduction in p38

phosphorylation in stimulated monocytes treated with nilotinib similar to the specific p38

inhibitor SB202190, which eventually impacted on phosphorylation and activation of the

downstream MAPK-activated protein kinase 2 (MK2) (**Figure 4B**). As Nilotinib did not affect

the upstream activity of MKK3 (**Figure 4B**), it is likely that nilotinib binds p38α specifically. In

order to demonstrate that the suppressive effects of nilotinib on the pro-inflammatory

phenotype were mediated by inhibition of the p38 MAPK pathway in THP-1 cells, we

analyzed the cytokine production mediated by p38 MAPK. In contrast to imatinib, nilotinib

significantly reduced TNF-α, IL-6 and IL-1β levels and expression in monocytes stimulated

with *E. coli* (**Figure 4C-E, Supplementary Figure 3**). Altogether, our results indicate that

nilotinib targets p38 alpha, which inhibits its activity and affects the production of pro-

inflammatory cytokines, thereby reducing inflammation.

259   18

Figure 4. Nilotinib inhibits p38 MAPK signalling. A) MALDI-TOF MS relative quantitation of the ratio

m/z 4964 / 4632 in THP-1 cells pre-treated with DMSO, 5 µM nilotinib or imatinib for one hour

before stimulation with live *E. coli* for up to 24h. **B)** Western blot analysis of the p38 MAPK pathway

in THP-1 cells pre-treated with DMSO, 5 µM nilotinib, 5 µM imatinib or 10 µM SB202190 for one

hour before stimulation with live *E. coli* for up to 15 min shows loss of p38 phosphorylation in

response to nilotinib. Nilotinib and SB202190, but not imatinib, block phosphorylation of the down-

stream MK2. α-tubulin serves as a loading control. A representative image with two biological

replicates of four replicates is shown. Relative mobilities of reference proteins (masses in kilo

Daltons) are shown on the left of each blot. **C)** TNF-α, **D)** IL-6, and **E)** IL-1β secretion was measured

by ELISA at 24h. ELISA IC50 curves are shown below. Error bars represent the SEM of four biological

replicates. Significant differences between two groups were determined by a Students t-test. The

statistical significance of the comparisons with *E. coli* is indicated as follows: ns, not significant; **, *P*

≤ 0.01; ***, *P* ≤ 0.001.

Nilotinib prevents monocyte activation.

In order to better understand differences between nilotinib and imatinib treatments on

monocytes, we performed quantitative proteomics of THP-1 cells pre-treated with DMSO, 5

µM nilotinib or imatinib for one hour before stimulation with live *E. coli* for up to 24h. We

identified and quantified a total of 3570 proteins of which 242 proteins showed significant

differences between nilotinib and imatinib treatments (**Supplementary Table 5** and

Supplementary Figure). Comparing differences between nilotinib and imatinib, we

identified 30 significant proteins (log2 fold-change ≤ -0.6; adjusted p-value <0.05)

downregulated with nilotinib (**Supplementary Table 6** and **Supplementary Figure 4C**). We

found decreased levels of proteins related with cell-cell adhesion and leukocyte migration

between DMSO- and nilotinib-treated cells compared to *E. coli*-stimulated and imatinib-

treated samples. Monocyte activation leads to differentiation into macrophages and also

leads to an increased expression of cell adhesion proteins.[30-33] To further understand how

nilotinib affected activation of monocytes and their differentiation into macrophages upon

stimulation with *E. coli,* we studied cell surface differentiation and cell adhesion markers.

We validated our proteomics data by showing that MMP9 and CD44 upregulation induced

by *E. coli* was abolished with nilotinib but not with imatinib (**Figure 5A-B**). Furthermore,

monocyte activation markers CD11c, CD14, CD49a (integrin α1) and CD54 (ICAM-1) were

suppressed with nilotinib treatment but not with imatinib (**Figure 5C-F**). Also, we observed

less cell adhesion with nilotinib treatment than with imatinib (**Figure 5G**). These results

indicate that nilotinib inhibits the inflammatory response of monocytes under a pro-

inflammatory stimulus, inhibiting their differentiation into macrophages.

Figure 5. Nilotinib inhibits cell adhesion and monocyte activation. A) MMP9 and p38 levels in THP-1

pre-treated with 5 µM nilotinib or imatinib for one hour before stimulation with live *E. coli* for up to

24h. α-tubulin serves as a loading control. A representative image with two biological replicates of

four replicates is shown. Relative mobilities of reference proteins (masses in kilo Daltons) are shown

on the left of each blot. **B)** Expression levels of CD44, **C)** CD11c, **D)** CD14, **E)** CD49a, and **F)** CD54 on

the cell surface were measured by flow cytometry. **G)** Representative images of cell adhesion assay

and quantification shows reduced cell adhesion upon treatment with nilotinib. Cells were stained

with crystal violet and then observed by light microscopy. Optical density of dissolved crystal violet

was used to evaluate the adhesion rate. Significant differences between two groups were

determined by a Students t-test. The statistical significance of the comparisons with *E. coli* is

indicated as follows: ns, not significant; *, $P \leq 0.05$; **, $P \leq 0.01$; ***, $P \leq 0.001$. Error bars represent

the SEM of four biological replicates.

Nilotinib prevents an inflammatory response in AML and MM.

Nilotinib is a second-line therapy for CML patients who failed or were intolerant to

imatinib,[23,34] but little is known about its use in other types of cancer. We performed an *in-*

silico analysis of data from the Genomics of Drug Sensitivity in Cancer (GDSC) database to

identify nilotinib and imatinib sensitivity in different cancer types including hematological

and solid tumors. The sensitivity to nilotinib was significantly higher than that of imatinib in

hematological malignancies and gastrointestinal tumors (**Supplementary Figure 5**). In

particular, AML, MM and diffuse large B cell lymphomas (DLBCL) showed significant

differences between the sensitivities to nilotinib and imatinib treatments (**Figure 6A**). In

order to test whether nilotinib may be an effective treatment in hematological

malignancies, we validated the anti-inflammatory effects of nilotinib in another monocytic

AML cell line, OCI-AML2, and in a MM cell line, NCI-H929. We also observed a significant

reduction in the ratio *m/z* 4964/4632 by MALDI-TOF MS in both OCI-AML2 and NCI-H929

cell lines treated with nilotinib (**Figure 6B**). Furthermore, we confirmed that nilotinib inhibits

325 p38 MAPK phosphorylation and subsequent MK2 phosphorylation in OCI-AML2 and NCI-

326 H929 cell lines after exposure to *E. coli* (**Figure 6C**). Moreover, TNF-α, IL-6, and IL-1β

327 secretion were reduced with nilotinib treatment in OCI-AML2 (**Supplementary Figure 6**).

328 Taken together, the results show that nilotinib inhibits p38 phosphorylation during the

329 inflammatory response, reducing the levels of cytokines such as TNF-α, IL-6 or IL-1 β (**Figure**

330 **6D**).

Figure 6. Nilotinib reduces the inflammation phenotype in AML and MM. A) IC50 values of nilotinib

and imatinib in hematological malignancies according to TCGA classification from the Genomics of

Drug Sensitivity in Cancer (GDSC) database. The statistical significance of the comparisons is

indicated as follows: ns, not significant; *, $P \leq 0.05$; ***, $P \leq 0.001$. **B)** MALDI-TOF MS relative

quantitation of the ratio *m/z* 4964/4632 in OCI-AML2 and NCI-H929 cells pre-treated with DMSO, 5

μM nilotinib or imatinib for one hour before stimulation with live *E. coli* for up to 24h. Error bars

represent the SEM of three biological replicates. Significant differences between two groups were

determined by a Students t-test. The statistical significance of the comparisons with *E. coli* is

indicated as follows: ns, not significant; *, $P \leq 0.05$; **, $P \leq 0.01$; ***, $P \leq 0.001$. **C)** p38 MAPK

pathway status in OCI-AML2 and NCI-H929 cells. α-tubulin was detected as a loading control. A

representative image of four replicates is shown. Relative mobilities of reference proteins (masses in

kilo Daltons) are shown on the left of each blot. **D)** Effect of nilotinib during pro-inflammatory

stimulus. Activation of TLR4 by a pro-inflammatory stimulus induces p38 MAPK signalling.

Nilotinib inhibits p38 phosphorylation and MK2 phosphorylation reducing the levels of

cytokines such as TNF-α, IL-6 or IL-1 β, and cell adhesion molecules (CAMs).

DISCUSSION

Monocytes and macrophages are major contributors to host immunity and immune

surveillance against inflammation.[1] Development of new high-throughput phenotypic drug

screening approaches is important for the identification of novel drugs for the treatment of

inflammation in autoimmune diseases and cancer. Here, we present a rapid and label-free

MALDI-TOF MS assay able to identify an inflammatory phenotype in monocytic cells, and to

screen for anti-inflammatory drugs. In recent years, MALDI-TOF MS has become an exciting

new tool for drug discovery for *in vitro* assays.[35-37] Here, we applied a screen in whole cells,

allowing the identification of biomolecules that provide a specific fingerprint for the

phenotype of the analyzed cells.[38] Two features at *m/z* 4632 and 4964 were significantly

altered upon inflammatory stimuli. The feature *m/z* 4964 has been previously described as a

biomarker to identify patients with Crohn's Disease, intestinal tuberculosis or rheumatoid

arthritis using MALDI MS imaging.[39,40] It has been identified as thymosin beta 4 (Tb4), a

small protein involved in binding and promoting actin polymerization, however it has also

been described as having anti-inflammatory properties.[41] Using our MALDI-TOF MS assay,

we identified nilotinib as a drug that prevents the inflammatory phenotype in stimulated

AML cells. Imatinib and nilotinib, BRC-ABL tyrosine kinase inhibitors (TKIs), are the standard

first-line therapy for CML.[24] However, nilotinib possesses higher potency than imatinib

against BCR-ABL and is active against most imatinib-resistant BCR-ABL mutations.[23] Our *in*

silico data suggest that a higher concentration of imatinib is necessary to achieve the same

effect elicited by nilotinib treatment in myeloid malignancies and gastrointestinal cancer;

such high doses could contribute to resistance or intolerance to imatinib in CML[23,34] and in

gastrointestinal stromal tumors as it has been observed in several clinical trials such as

NCT00633295, NCT00718562 or NCT00756509. Our results indicate that a lower nilotinib

dose already exerts anti-inflammatory effects on AML and MM. It would be interesting to

explore whether nilotinib has the same effect in chronic gastrointestinal inflammatory

diseases such as inflammatory bowel disease or Crohn's disease, which are the primary risk

factor for the development of gastrointestinal malignancies.[42]

Stimulation with TLR ligands activates monocytes, which leads to the production of

cytokines, chemokines, and mediators that are involved in inflammation.[2,43] Dysregulation

of TLR signalling has been linked with myeloproliferative disorders.[4,5] The bacterial surface

molecule LPS activates TLR4 and CD14 in monocytes, inducing pro-inflammatory (IL-1, IL-6

and TNF-α) and then anti-inflammatory (IL-10, soluble TNF receptor, and IL-1 receptor

antagonist) cytokines.[6,10] Nilotinib treatment was able to decrease the levels of IL-1β, IL-6,

and TNF-α cytokines secretion upon stimulation with *E. coli* in AML cells. In line with our

results, pre-incubation with nilotinib in murine macrophages derived from bone marrow led

to a decreased LPS-induced *IL-6* expression.[44] Furthermore, monocyte differentiation into

macrophages induces an up-regulation of CD11c and CD14 surface markers,[30,31] as well as

cell adhesion molecules.[32,33] Interestingly, nilotinib was able to block the upregulation of

CD11c, CD14, CD44, CD49a, and CD54 markers, thus reducing immune response and cell

adhesion of monocytes. We further found MMP9, which cleaves CD44 and induces cell

migration and cell-cell and cell-matrix adhesion,[45] downregulated upon nilotinib treatment.

Furthermore, we observed a reduction in cell adhesion in activated monocytes upon

nilotinib treatment, compared with imatinib. Our data therefore suggests that nilotinib may

not only affect inflammatory signalling and immune cell adhesion, but also differentiation

from monocytes into macrophages.

TLR ligation induces the formation of a signalling complex that includes IL-1R-associated

kinases (IRAKs) and TNFR-associated factor 6 (TRAF6), which is mediated by K63-linked

polyubiquitylation. This complex interacts with and activates TGFβ-activated kinase 1

(TAK1), a MAPK kinase kinase (MAP3K) upstream of p38α and Jun N-terminal kinases (JNKs).

These events in turn lead to activation of transcription factors, triggering the production of

cytokines and chemokines.[9,46] TPP analysis identified p38α as a main off-target of nilotinib,

when compared with imatinib. Nilotinib has been reported to inhibit p38α and p38γ, in an *in*

vitro kinase assay and may stabilize p38α through direct interaction, as it has been observed

in vascular endothelial cells.[47] Furthermore, it was shown that nilotinib was able to block

p38 and JNK activation in an *in vivo* model of ischemic reperfusion injury, leading to

decreased cytokine production and modulated leukocyte recruitment.[48] However, we did

not find any evidence for p38γ or JNK inhibition in our experiments, suggesting remarkable

specificity for the 38α isoform. Furthermore, our results showed that nilotinib interferes

with the downstream MAPK signalling pathway comparable to a specific p38 inhibitor. The

potential of p38 MAPK inhibitors was initially explored within inflammatory conditions such

as rheumatoid arthritis and Crohn's disease, but early studies demonstrated poor clinical

efficacy and unacceptable side effects.[49] This suggests that nilotinib may be considered for

additional applications outside of cancer, particularly as it is well tolerated by patients and

has few side effects. [23,34] Within the cancer field, it appears that the anti-inflammatory off-

target effects from nilotinib may be beneficial as seen in tumors co-treated with doxorubicin

and vincristine.[50,51] Consistent with these results, nilotinib – but not imatinib – can reduce

the inflammation in human monocytes-derived from AML and other myeloid cells.

Our study provides a valuable tool to discover new anti-inflammatory drugs using MALDI-

TOF MS, as we demonstrate here for nilotinib. We show that nilotinib inhibits p38α,

prevents its phosphorylation and subsequent activation, affecting the phosphorylation of

downstream signalling targets and, ultimately preventing the transcription of pro-

inflammatory genes, cell adhesion markers and innate immunity markers. In consequence,

nilotinib may have therapeutic potential through the inhibition of p38 MAPK pathway in

inflammatory diseases as well as in myeloid malignancies such as myeloid leukemia and

multiple myeloma.

ACKNOWLEDGEMENTS

We would like to thank the animal staff at Newcastle University's Comparative Biology Centre, and Akshada Gajbhiye, Abeer Dannoura, Julien Peltier, and Anetta Härtlova for their technical support. This research was partly funded by a Wellcome Trust Investigator Award (215542/Z/19/Z) and generous start-up funding of Newcastle University to M.T. R.E.H. was funded by an iCASE studentship by the BBSRC and Bruker Daltonics. M.E.D. is a Marie Sklodowska Curie Fellow within the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 890296. For the purpose of Open Access, the author has applied a CC BY public copyright license to any Author Accepted Manuscript version arising from this submission.

AUTHORSHIP CONTRIBUTIONS

R.E.H. and J.L.M.-R. are co-first authors. J.L.M.-R. performed cellular thermal shift assay, ELISA and flow cytometry. R.E.H. and M.E.D. performed MALDI-TOF MS experiments. J.L.M.-R. and T.H. performed LFQ and TMT mass spectrometry-based proteomics experiments. T.H. analyzed TPP experiments. J.L.M.-R. and J.I. performed and analyzed RTqPCR assays. B.S. provided drug panel. M.T. conceived the original idea and provided supervision. J.L.M.-R., R.E.H., and M.T. wrote the manuscript. All authors provided critical feedback and helped shape the research, analysis and manuscript.

SUPPLEMENTAL INFORMATION

The online version of this article contains supplementary data.

452   **REFERENCES**

453   1   Gordon, S. & Taylor, P. R. Monocyte and macrophage heterogeneity. *Nat Rev*
454   *Immunol* **5**, 953-964, doi:10.1038/nri1733 (2005).
455   2   Stuhlmuller, B. *et al.* Identification of known and novel genes in activated
456   monocytes from patients with rheumatoid arthritis. *Arthritis Rheum* **43**, 775-
457   790, doi:10.1002/1529-0131(200004)43:4<775::AID-ANR8>3.0.CO;2-7 (2000).
458   3   Binder, S., Luciano, M. & Horejs-Hoeck, J. The cytokine network in acute
459   myeloid leukemia (AML): A focus on pro- and anti-inflammatory mediators.
460   *Cytokine Growth Factor Rev* **43**, 8-15, doi:10.1016/j.cytogfr.2018.08.004
461   (2018).
462   4   Hemmati, S., Haque, T. & Gritsman, K. Inflammatory Signaling Pathways in
463   Preleukemic and Leukemic Stem Cells. *Front Oncol* **7**, 265,
464   doi:10.3389/fonc.2017.00265 (2017).
465   5   Monlish, D. A., Bhatt, S. T. & Schuettpelz, L. G. The Role of Toll-Like Receptors
466   in Hematopoietic Malignancies. *Front Immunol* **7**, 390,
467   doi:10.3389/fimmu.2016.00390 (2016).
468   6   Takeuchi, O. *et al.* Differential roles of TLR2 and TLR4 in recognition of gram-
469   negative and gram-positive bacterial cell wall components. *Immunity* **11**, 443-
470   451, doi:10.1016/s1074-7613(00)80119-3 (1999).
471   7   Takeda, K. & Akira, S. Toll-like receptors in innate immunity. *Int Immunol* **17**,
472   1-14, doi:10.1093/intimm/dxh186 (2005).
473   8   Kim, E. K. & Choi, E. J. Pathological roles of MAPK signaling pathways in human
474   diseases. *Biochim Biophys Acta* **1802**, 396-405,
475   doi:10.1016/j.bbadis.2009.12.009 (2010).
476   9   Suzuki, T. *et al.* Different involvement of the MAPK family in inflammatory
477   regulation in human pulmonary microvascular endothelial cells stimulated
478   with LPS and IFN-gamma. *Immunobiology* **223**, 777-785,
479   doi:10.1016/j.imbio.2018.08.003 (2018).
480   10   Mogensen, T. H. Pathogen recognition and inflammatory signaling in innate
481   immune defenses. *Clin Microbiol Rev* **22**, 240-273, Table of Contents,
482   doi:10.1128/CMR.00046-08 (2009).
483   11   Inglese, J. *et al.* High-throughput screening assays for the identification of
484   chemical probes. *Nat Chem Biol* **3**, 466-479, doi:10.1038/nchembio.2007.17
485   (2007).
486   12   Neefjes, J. & Dantuma, N. P. Fluorescent probes for proteolysis: tools for drug
487   discovery. *Nat Rev Drug Discov* **3**, 58-69, doi:10.1038/nrd1282 (2004).
488   13   Adegbola, S. O., Sahnan, K., Warusavitarne, J., Hart, A. & Tozer, P. Anti-TNF
489   Therapy in Crohn's Disease. *Int J Mol Sci* **19**, doi:10.3390/ijms19082244
490   (2018).
491   14   Dulai, P. S. & Sandborn, W. J. Next-Generation Therapeutics for Inflammatory
492   Bowel Disease. *Curr Gastroenterol Rep* **18**, 51, doi:10.1007/s11894-016-0522-
493   0 (2016).

15   Li, P., Zheng, Y. & Chen, X. Drugs for Autoimmune Inflammatory Diseases:
From Small Molecule Compounds to Anti-TNF Biologics. *Front Pharmacol* **8**,
460, doi:10.3389/fphar.2017.00460 (2017).

16   Guitot, K. *et al.* A direct label-free MALDI-TOF mass spectrometry based assay
for the characterization of inhibitors of protein lysine methyltransferases. *Anal*
Bioanal Chem **409**, 3767-3777, doi:10.1007/s00216-017-0319-5 (2017).

17   Heap, R. E. *et al.* Identifying Inhibitors of Inflammation: A Novel High-
Throughput MALDI-TOF Screening Assay for Salt-Inducible Kinases (SIKs). *SLAS*
Discov **22**, 1193-1202, doi:10.1177/2472555217717473 (2017).

18   De Cesare, V. *et al.* The MALDI-TOF E2/E3 Ligase Assay as Universal Tool for
Drug Discovery in the Ubiquitin Pathway. *Cell Chem Biol* **25**, 1117-1127 e1114,
doi:10.1016/j.chembiol.2018.06.004 (2018).

19   Bosshart, H. & Heinzelmann, M. THP-1 cells as a model for human monocytes.
Ann Transl Med **4**, 438, doi:10.21037/atm.2016.08.53 (2016).

20   Clark, K. *et al.* Novel cross-talk within the IKK family controls innate immunity.
Biochem J **434**, 93-104, doi:10.1042/BJ20101701 (2011).

21   Tan, L. *et al.* Discovery of type II inhibitors of TGFbeta-activated kinase 1
(TAK1) and mitogen-activated protein kinase kinase kinase kinase 2 (MAP4K2).
J. Med. Chem. **58**, 183-196, doi:10.1021/jm500480k (2015).

22   Malik, N. *et al.* Suppression of interferon beta gene transcription by inhibitors
of bromodomain and extra-terminal (BET) family members. *Biochem. J.* **468**,
363-372, doi:10.1042/BJ20141523 (2015).

23   Wei, G., Rafiyath, S. & Liu, D. First-line treatment for chronic myeloid
leukemia: dasatinib, nilotinib, or imatinib. *J Hematol Oncol* **3**, 47,
doi:10.1186/1756-8722-3-47 (2010).

24   Pan, P. *et al.* Systematic Review and Meta-Analysis of -New-Generation
Tyrosine Kinase Inhibitors versus Imatinib for Newly Diagnosed Chronic
Myeloid Leukemia. *Acta Haematol*, 1-13, doi:10.1159/000501537 (2019).

25   Franken, H. *et al.* Thermal proteome profiling for unbiased identification of
direct and indirect drug targets using multiplexed quantitative mass
spectrometry. *Nat Protoc* **10**, 1567-1593, doi:10.1038/nprot.2015.101 (2015).

26   Mateus, A. *et al.* Thermal proteome profiling for interrogating protein
interactions. *Mol Syst Biol* **16**, e9232, doi:10.15252/msb.20199232 (2020).

27   Miettinen, T. P. *et al.* Thermal proteome profiling of breast cancer cells reveals
proteasomal activation by CDK4/6 inhibitor palbociclib. *EMBO J* **37**,
doi:10.15252/embj.201798359 (2018).

28   Becher, I. *et al.* Thermal profiling reveals phenylalanine hydroxylase as an off-
target of panobinostat. *Nat Chem Biol* **12**, 908-910,
doi:10.1038/nchembio.2185 (2016).

29   Savitski, M. M. *et al.* Tracking cancer drugs in living cells by thermal profiling of
the proteome. *Science* **346**, 1255784, doi:10.1126/science.1255784 (2014).

30   Ma, W. *et al.* The p38 mitogen-activated kinase pathway regulates the human
interleukin-10 promoter via the activation of Sp1 transcription factor in

lipopolysaccharide-stimulated human macrophages. *J Biol Chem* **276**, 13664-
13674, doi:10.1074/jbc.M011157200 (2001).

31   Chen, R. F., Wang, L., Cheng, J. T. & Yang, K. D. Induction of IFNalpha or IL-12
depends on differentiation of THP-1 cells in dengue infections without and
with antibody enhancement. *BMC Infect Dis* **12**, 340, doi:10.1186/1471-2334-
12-340 (2012).

32   Kounalakis, N. S. & Corbett, S. A. Lipopolysaccharide transiently activates THP-
1 cell adhesion. *J Surg Res* **135**, 137-143, doi:10.1016/j.jss.2005.12.018 (2006).

33   Park, G. S. & Kim, J. H. LPS Up-Regulates ICAM-1 Expression in Breast Cancer
Cells by Stimulating a MyD88-BLT2-ERK-Linked Cascade, Which Promotes
Adhesion to Monocytes. *Mol Cells* **38**, 821-828,
doi:10.14348/molcells.2015.0174 (2015).

34   Sasaki, K. *et al.* Frontline therapy with high-dose imatinib versus second
generation tyrosine kinase inhibitor in patients with chronic-phase chronic
myeloid leukemia - a propensity score analysis. *Haematologica* **101**, e324-327,
doi:10.3324/haematol.2015.139089 (2016).

35   De Cesare, V. *et al.* High-throughput matrix-assisted laser
desorption/ionization time-of-flight (MALDI-TOF) mass spectrometry-based
deubiquitylating enzyme assay for drug discovery. *Nat. Protoc.* **15**, 4034-4057,
doi:10.1038/s41596-020-00405-0 (2020).

36   Simon, R. P. *et al.* MALDI-TOF-Based Affinity Selection Mass Spectrometry for
Automated Screening of Protein-Ligand Interactions at High Throughput. *SLAS*
Discov **26**, 44-57, doi:10.1177/2472555220959266 (2021).

37   Winter, M. *et al.* Automated MALDI Target Preparation Concept: Providing
Ultra-High-Throughput Mass Spectrometry-Based Screening for Drug
Discovery. *SLAS Technol* **24**, 209-221, doi:10.1177/2472630318791981 (2019).

38   Heap, R. E., Segarra-Fas, A., Blain, A. P., Findlay, G. M. & Trost, M. Profiling
embryonic stem cell differentiation by MALDI TOF mass spectrometry:
development of a reproducible and robust sample preparation workflow.
Analyst **144**, 6371-6381, doi:10.1039/c9an00771g (2019).

39   Kriegsmann, M. *et al.* MALDI MS imaging as a powerful tool for investigating
synovial tissue. *Scand J Rheumatol* **41**, 305-309,
doi:10.3109/03009742.2011.647925 (2012).

40   Zhang, F. *et al.* Exploration of Serum Proteomic Profiling and Diagnostic Model
That Differentiate Crohn's Disease and Intestinal Tuberculosis. *PLoS One* **11**,
e0167109, doi:10.1371/journal.pone.0167109 (2016).

41   Holzlechner, M. *et al.* In Situ Characterization of Tissue-Resident Immune Cells
by MALDI Mass Spectrometry Imaging. *J. Proteome Res.* **16**, 65-76,
doi:10.1021/acs.jproteome.6b00610 (2017).

42   Boyapati, R. K., Rossi, A. G., Satsangi, J. & Ho, G. T. Gut mucosal DAMPs in IBD:
from mechanisms to therapeutic implications. *Mucosal Immunol* **9**, 567-582,
doi:10.1038/mi.2016.14 (2016).

43   Yang, R. B. *et al.* Toll-like receptor-2 mediates lipopolysaccharide-induced
cellular signalling. *Nature* **395**, 284-288, doi:10.1038/26239 (1998).
44   Wong, J. *et al.* Small molecule kinase inhibitors block the ZAK-dependent
inflammatory effects of doxorubicin. *Cancer Biol Ther* **14**, 56-63,
doi:10.4161/cbt.22628 (2013).
45   Chetty, C. *et al.* MMP-9 induces CD44 cleavage and CD44 mediated cell
migration in glioblastoma xenograft cells. *Cell Signal* **24**, 549-559,
doi:10.1016/j.cellsig.2011.10.008 (2012).
46   Arthur, J. S. & Ley, S. C. Mitogen-activated protein kinases in innate immunity.
Nat Rev Immunol **13**, 679-692, doi:10.1038/nri3495 (2013).
47   Hadzijusufovic, E. *et al.* Nilotinib-induced vasculopathy: identification of
vascular endothelial cells as a primary target site. *Leukemia* **31**, 2388-2397,
doi:10.1038/leu.2017.245 (2017).
48   Ocuin, L. M. *et al.* Nilotinib protects the murine liver from
ischemia/reperfusion injury. *J Hepatol* **57**, 766-773,
doi:10.1016/j.jhep.2012.05.012 (2012).
49   Fisk, M., Gajendragadkar, P. R., Maki-Petaja, K. M., Wilkinson, I. B. & Cheriyan,
J. Therapeutic potential of p38 MAP kinase inhibition in the management of
cardiovascular disease. *Am J Cardiovasc Drugs* **14**, 155-165,
doi:10.1007/s40256-014-0063-6 (2014).
50   Dent, P. The flip side of doxorubicin: Inflammatory and tumor promoting
cytokines. *Cancer Biol Ther* **14**, 774-775, doi:10.4161/cbt.26125 (2013).
51   Wong, J., Tran, L. T., Magun, E. A., Magun, B. E. & Wood, L. J. Production of IL-
1beta by bone marrow-derived macrophages in response to
chemotherapeutic drugs: synergistic effects of doxorubicin and vincristine.
Cancer Biol Ther **15**, 1395-1403, doi:10.4161/cbt.29922 (2014).

METHODS
Compounds

LPS and SB202190 were purchased from Sigma-Aldrich; Pam2CSK4 and Pam3CSK4 from

Invitrogen; IFN-γ from PeproTech; and nilotinib, imatinib, NG-25, MRT67307, and the other

compounds from Supplementary Table 1 were provided by LifeArc.

Cell culture.

Acute myeloid leukemia-derived cell lines THP-1 (ATCC) and OCI-AML2 (DSMZ), and multiple

myeloma-derived cell lines NCI-H929 (ATCC) were cultured at 37°C in a humidified 5% CO_2

atmosphere. ATCC and DSMZ routinely perform cell line authentication, using short tandem

repeat profiling as a procedure. Cell experimentation was always performed within a period

not exceeding six months after resuscitation in mycoplasma-free culture conditions.

Blind drug screening.

One million THP-1 cells/mL were incubated with each compound (**Supplementary Table 1**)

at 5 µM for one hour in technical triplicates before stimulation with 100 ng/mL LPS for 24h.

Compounds were staggered into 8 sets each with individual positive (NG-25) and negative

controls (MRT67307) to enable maximum relative peak area calculations. This screening

process was repeated in three biological replicates.

Cell infection with *E. coli*.

The DH5α strain of *E. coli* bacteria (Invitrogen) was harvested during mid-log phase.

Mammalian cells were infected with a multiplicity of infection of 2. The cells with bacteria

were centrifuged at 500 xg for 5 min at 37°C and incubated at 37°C in 5% CO_2 for bacterial

uptake for 15 min or 30 min. Thirty min post-infection, cells were washed once with PBS and

incubated for 1h with 100 µg/mL gentamicin to kill extracellular bacteria. Then, the cells

were washed with PBS twice and the media replaced with 20 µg/mL gentamicin for the

remainder of the experiment.

Sample preparation for MALDI-TOF MS and data analysis.

Cell pellets were frozen on dry ice, then thawed and washed with 100 mM Tris-HCl, pH 7.5

and centrifuged at 1,000 xg for 10 min at 4°C. 2,500 cells were spotted on target with 10

mg/mL α-cyano-4-cinnamic acid in 50% acetonitrile, 0.1% trifluoroacetic acid. Automated

target spotting was performed using a Mosquito liquid handling robot (TTP Labtech). A

RapifleX PharmaPulse MALDI TOF/TOF mass spectrometer (Bruker Daltonics) equipped with

a Smartbeam 3D laser was used in positive ion mode. Samples were acquired in automatic

mode (AutoXecute; Bruker Daltonics), totaling 10,000 shots at a 10 kHz frequency per spot.

A random walk pattern (complete sample) on spot laser ablation pattern was used with an

M5 Smart beam Parameter at a 45 μm × 45 μm scan range. Spot diameter was limited to

2,000 μm and a random walk pattern movement enabled at 1000 shots per raster position.

Ionization was achieved using a laser power of 75% (laser attenuator offset 14%, range 30%)

with a detector gain of ×6.8 in the mass range of *m/z* 2,000-20,000 with a mass suppression

up to m/z 1,600. Samples were analyzed in a linear geometry with optimized voltages for

ion sources (ion source 1, 20 kV; pulsed ion extraction 1.3 kV), lens (8.6 kV), and a pulsed ion

extraction of 180 ns. A novel 10-bit digitizer was used at a sampling rate of 1.25 GS/s. Raw

data were processed first by a TopHat baseline subtraction followed by smoothing with a

SavitzkyGolay algorithm. MALDI-TOF spectra were processed by FlexAnalysis Batch Process

(Compass 2.0). Spectra-based PCA plots were generated using ClinPro Tools (Bruker

Daltonics). A normalized relative peak area was used to measure a pro-inflammatory

response between *m/z* 4632 and 4963 and was calculated with the following equation:

$$Normalised\ peak\ area\ ratio = \left(\left(\frac{4963\ peak\ area}{4632\ peak\ area} \right) / (LPS\ treated) \right) \times 100$$

This method of intra-spectra quantification[1] was robust over ten different passages, where

we were able to quantify the inflammatory response by MALDI-TOF MS with Z'>0.5 and

$P \leq 0.001$.

Quantitative RT-PCR.

RNA was extracted using Trizol (Thermo Fisher Scientific). Real-time RT-qPCR from total RNA

in two steps, was performed with a QuantiTect reverse transcription kit (Qiagen), and

QuantiTect SYBR green kit (Qiagen) using a StepOne Applied Biosystems real-time PCR

system (Thermo Fisher Scientific). Expression values of *GAPDH* and *TBP* genes in the same

samples were used for normalization, using the $2^{-\Delta\Delta CT}$ method. The following primers were

used: *GAPDH*,   forward: GTCTCCTCTGACTTCAACAGCG   and,   reverse:

ACCACCCTGTTGCTGTAGCCAA; *IL-6*, forward: GCCCAGCTATGAACTCCTTCT and, reverse:

CTTCTCCTGGGGGTACTGG; *TBP*, forward: GAGTTCCAGCGCAAGGGTTT, and, reverse:

GGGTCAGTCCAGTGCCATA; *TNF-α*, forward: AAACTCATGAGCAGTCTGCA, and reverse:

AGGAGATCTTCAGTTTCGGAGG.

ELISA.

Cell culture supernatant was collected at 24h post-treatment. TNF-a, IL-6, and IL-1β were

measured by DuoSet ELISA kits (R&D Systems). Absorbance from four biological replicates at

450 nm was measured with the correction wavelength set at 540 nm using SpectraMax® M3

microplate reader (Molecular Devices).

Western blot.

Cells were lysed using 5% SDS supplemented with Protease Inhibitor Cocktail and

Phosphatase Inhibitor Cocktail 2 (Sigma-Aldrich). The following antibodies were purchased

from Cell Signaling Technology (Danvers): MK2 (#3042), Thr-334-P-MK2 (#3041), MK3

(#7421), MKK3 (#8535), Ser189-P-MKK3/Ser207-P-MKK6 (#12280), p38 MAPK (#9212), p38α

MAPK (#9218), p38γ MAPK (#2307) Thr180/Tyr182-P-p38 MAPK (#4511), MPP9 (#13667),

anti-rabbit IgG-HRP (#7074), and anti-mouse IgG (#7076). GAPDH (sc-47724) was from Santa

Cruz, and α-Tubulin (T9026) was from Sigma-Aldrich. Amersham Imager 600 digital imaging

system (GE Healthcare) was used for image acquisition.

Flow cytometry.

Cell surface staining was performed by direct immunofluorescent assay with fluorescent-

conjugated antibodies: CD11c-APC (#17-0114-82), CD14-AF700 (#56-0149-42), and CD54-

FITC (#11-0541-82) from Thermo Fisher; CD49a-PE (#562115) and CD44-FITC (#338803) from

BD Biosciences; and corresponding isotype control antibodies, for 30 min at 4°C in PBS with

1% FBS, 1% BSA and 1% human serum (Sigma) to block Fc receptors. Cells were analyzed in a

FACSCanto II flow cytometer (Becton-Dickinson). The results were analyzed using FlowJo.

Adhesion assay.

Cells were washed twice with PBS before being fixed with 4% paraformaldehyde for 15 min.

Then, cells were incubated with 5 mg/ml Crystal violet for 10 min before lysis with 2% SDS

for 30 min. Absorbance from four biological replicates at 550 nm was measured using a

SpectraMax® M3 microplate reader (Molecular Devices). Micrographs were acquired with an

inverted microscope Axio Vert.A1 FL LED (Zeiss, Cambridge).

Cellular thermal shift assay.

Cells were treated with 5 µM of nilotinib or imatinib for one hour, then 100 ng/mL LPS was

added for 15 min. Cells were washed with PBS supplemented with cOmplete, EDTA-free

Protease Inhibitor Cocktail (Sigma-Aldrich), 1.2 mM sodium molybdate, 1 mM sodium

orthovanadate, 4 mM sodium tartrate dihydrate, 5 mM glycerophosphate, and 20 mM N-

ethylmaleimide; then separated into 8 fractions for thermal profiling. Fractions were heated

at 40, 44, 48, 52, 56, 60, 64 and 68°C for 3 min. For concentration compound range

experiments, cells were treated with 0.1, 0.5, 1, 5 or 10 µM of nilotinib or imatinib for 1h

before treatment with 100 ng/mL LPS for 15 min. Cells were heated at 56°C for 3 min.

Samples were lysed with four freeze-thaw cycles using dry ice and a thermo block at 35°C.

Cell lysates were centrifuged at 20,000 xg for 2h at 4°C to separate protein aggregates from

soluble proteins. Supernatants were collected and used for western blot and mass

spectrometry.

Sample preparation for LC-MS/MS and label-free quantification (LFQ) data analysis.

Samples were prepared for mass spectrometry using S-Trap micro columns (Protifi)

according to the manufacturer's recommended protocol. Proteins were reduced with 20

mM TCEP at 37°C for 30 min and alkylated with 20 mM N-ethylmaleimide in the dark for 30

min. A ratio of 1:10 wt:wt Trypsin TPCK Treated (Worthington-Biochem) was used to digest

the samples for 2h at 47°C. Eluted peptides were dried down and resuspended in loading

buffer (2% acetonitrile, 0.1% trifluoroacetic acid). Peptide samples were injected on a

Dionex Ultimate 3000 RSLC (Thermo Fisher Scientific) connected to an Orbitrap Fusion

Lumos Tribrid mass spectrometer (Thermo Fisher Scientific). Samples were injected on a

PepMap 100 C18 LC trap column (300 µm ID x 5 mm, 5 µm, 100 Å) followed by separation

on an EASY-Spray column (50 cm x 75 μm ID, PepMap C18, 2 μm, 100 Å) (Thermo Fisher

Scientific). Buffer A consisted of water containing 0.1% FA and Buffer B of 80% acetonitrile

containing 0.1% FA. Peptides were separated with a linear gradient of 3-35% Buffer B over

180 min followed by a step from 35-90% Buffer B in 0.5 min at 250 nL/min and held at 90%

for 4 min. The gradient was then decreased to 3% Buffer B in 0.5 min at 250 nL/min, and the

column equilibrated for 10 min before the next injection. Column temperature was

controlled at 45°C. The Orbitrap Fusion Lumos Tribrid mass spectrometer was operated in

data-dependent, positive ion mode. Full scan spectra were acquired in the range of m/z 400

to 1,600, at a resolution of 120,000, with an automatic gain control (AGC) target of $4x10^5$

and a maximum injection time of 50 ms. The Orbitrap was operated in 'stop speed' to

maintain a 3 sec fixed duty cycle. The most intense precursor ions were isolated with a

quadrupole mass filter width of 1.6 m/z and higher-energy collision-induced dissociation

(HCD) fragmentation was performed in one-step collision energy of 30%. Detection of

MS/MS fragments was acquired in the linear ion trap in rapid scan mode with an AGC target

of $1x10^4$ ions and a maximum injection time of 45 ms. An electrospray voltage of 2.0 kV and

capillary temperature of 275°C, with no sheath and auxiliary gas flow, was used.

All discovery proteomics RAW mass spectra were analyzed using MaxQuant (version

1.6.10.43)[2] and searched against a SwissProt Homo sapiens database (containing 42,347

database entries with isoforms; downloaded on 21 April 2020). Trypsin/P was set as

proteolytic enzyme, N-ethylmaleimide on cysteine was set as fixed modification, and

methionine oxidation and acetylation of protein N-termini was set as variable modifications.

Two missed cleavages were allowed. A protein and peptide false discovery rate (FDR) of less

than 1% was employed in MaxQuant. LFQ data analysis was performed using Perseus

(version 1.6.2.3).[3] Reverse hits, contaminants and proteins only identified by site were

removed before downstream statistical and bioinformatics analysis. LFQ intensity data were

transformed (log 2) and filtered to contain at least two unique peptides and at least three

valid values in one group for comparisons. Significance testing was carried out using a two-

tailed unpaired Student's t-test and multiple hypothesis testing was controlled using the

Benjamini-Hochberg FDR threshold of p-value <0.05. Gene ontology term enrichments were

performed within DAVID (version 6.8).[4]

Thermal proteome profiling (TPP) and data analysis.

Isobaric labeling of peptides was performed using the 16-plex tandem mass tag (TMT)

reagents (Thermo Fisher Scientific) (**Supplementary Table 2**), according to the manufacturer

recommended protocol. According to the different temperature points, labeled peptides

were combined and desalted with C18 Macro Spin Columns (Harvard Apparatus, Holliston,

MA, USA). The TMT-labeled samples were subjected to fractionation using basic-pH

reversed-phase liquid chromatography on a Dionex Ultimate 3000 HPLC system (Thermo

Fisher Scientific) using a Gemini C18 column (250 mm × 3 mm, 3 µm, 110 Å; Phenomenex).

Buffer A consisted of 20 mM ammonium formate pH 8.0 and Buffer B of 100% acetonitrile.

Peptides were fractionated using a linear gradient of 1-49% buffer B over 49 min at 250

nl/min. Twelve fractions were dried under vacuum centrifugation and resuspended in 2%

acetonitrile with 0.1% TFA for LC-MS/MS analysis. Labeled peptide samples were injected on

a Dionex Ultimate 3000 RSLC (Thermo Fisher Scientific) connected to an Orbitrap Fusion

Lumos Tribrid mass spectrometer (Thermo Fisher Scientific). Buffer A consisted of water

containing 0.1% FA and Buffer B of 80% acetonitrile containing 0.1% FA. Peptides were

separated with a linear gradient of 3-35% Buffer B over 180 min followed by a step from 35-

90% Buffer B in 0.5 min at 250 nL/min and held at 90% for 4 min. The gradient was then

decreased to 3% Buffer B in 0.5 min at 250 nL/min, and the column equilibrated for 10 min

before the next injection. The scan sequence began with an MS1 spectrum (Orbitrap

analysis; resolution 120,000; mass range *m/z* 375-1500; automatic gain control (AGC) target

4x10E5; maximum injection time 50 ms). MS2 analysis consisted of collision-induced

dissociation (CID); AGC 1x10E4; normalized collision energy (NCE) of 30%; maximum

injection time 50 ms; and isolation window of 0.7. Following acquisition of each MS2

spectrum, MS3 precursors were fragmented by high energy collision-induced dissociation

(HCD) and analyzed using the Orbitrap (NCE 55%; AGC 5x10E4; maximum injection time 86

ms, resolution was 50,000) and synchronous precursor selection (SPS) was enabled to

include 10 MS/MS fragment ions in the FTMS3 scan.

Mass spectrometry data analysis was performed similarly as described above, but

TMT modification on the peptide N-termini or lysine residues were enabled for the 16-plex

TMT reagents, and deamidation of asparagine and glutamine was included as variable

modification in MaxQuant. TPP data analysis was performed using the TPP packages

(https://www.bioconductor.org/packages/release/bioc/html/TPP.html) in the R statistical

programming language. The data were normalized as detailed in Miettinen et al.[5]

(**Supplementary Table 3**).

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange

Consortium[6] via the PRIDE partner repository[7] with the data set identifier: PXD025020.

Reviewer account details: Username: reviewer_pxd025020@ebi.ac.uk   Password:

H0T2rCOL

References

1   Chen, X. *et al.* Ratiometric Mass Spectrometry for Cell Identification and
802   Quantitation Using Intracellular "Dual-Biomarkers". *Sci Rep* **7**, 17432,
803   doi:10.1038/s41598-017-17812-1 (2017).
804   2   Cox, J. & Mann, M. MaxQuant enables high peptide identification rates,
805   individualized p.p.b.-range mass accuracies and proteome-wide protein
806   quantification. *Nat Biotechnol* **26**, 1367-1372, doi:10.1038/nbt.1511 (2008).
807   3   Tyanova, S. *et al.* The Perseus computational platform for comprehensive
808   analysis of (prote)omics data. *Nat Methods* **13**, 731-740,
809   doi:10.1038/nmeth.3901 (2016).
810   4   Huang da, W., Sherman, B. T. & Lempicki, R. A. Systematic and integrative
811   analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* **4**,
812   44-57, doi:10.1038/nprot.2008.211 (2009).
813   5   Miettinen, T. P. *et al.* Thermal proteome profiling of breast cancer cells reveals
814   proteasomal activation by CDK4/6 inhibitor palbociclib. *EMBO J* **37**,
815   doi:10.15252/embj.201798359 (2018).
816   6   Deutsch, E. W. *et al.* The ProteomeXchange consortium in 2017: supporting
817   the cultural change in proteomics public data deposition. *Nucleic Acids Res* **45**,
818   D1100-D1106, doi:10.1093/nar/gkw936 (2017).
819   7   Perez-Riverol, Y. *et al.* The PRIDE database and related tools and resources in
820   2019: improving support for quantification data. *Nucleic Acids Res* **47**, D442-
821   D450, doi:10.1093/nar/gky1106 (2019).
822
823