1

Extensive aquatic subsidies lead to territorial breakdown and high

density of an apex predator

Charlotte E. Eriksson[1], Daniel L.Z. Kantek[2], Selma S. Miyazaki[2], Ronaldo G. Morato[3], Manoel dos Santos-Filho[4], Joel S. Ruprecht[1], Carlos A. Peres[5], Taal Levi[1]

[1]Department of Fisheries and Wildlife, Oregon State University, Corvallis, OR, United States of America

[2]Instituto Chico Mendes de Conservação da Biodiversidade (ICMBIO), Estação Ecológica de Taiamã (EET). Cáceres, MT, Brasil

[3]Instituto Chico Mendes de Conservação da Biodiversidade (ICMBIO), Centro Nacional de Pesquisa e Conservação de Mamíferos Carnívoros (CENAP). Atibaia, SP, Brasil

[4]Departamento de Ciências Biológicas, Universidade do Estado de Mato Grosso (UNEMAT). Cáceres-MT CEP: 78.200-000 Brasil

[5]Centre for Ecology, Evolution and Conservation, School of Environmental Sciences, University of East Anglia, Norfolk, Norwich, UK,

Abstract

Energetic subsidies between terrestrial and aquatic ecosystems can strongly influence food webs and

population dynamics. Our objective was to study how aquatic subsidies affected jaguar (*Panthera onca*)

diet, sociality, and population density in a seasonally flooded protected area in the Brazilian Pantanal. The

diet (n = 138 scats) was dominated by fish (46%) and aquatic reptiles (55%), representing the first jaguar

population known to feed extensively on fish and to minimally consume mammals (11%). These aquatic

subsidies supported the highest jaguar population density estimate to date (12.4 per 100 km²) derived

from camera traps (8,065 trap nights) and GPS collars (n = 13). Contrary to their mostly solitary behavior

elsewhere, we documented social interactions previously unobserved between same-sex adults including

cooperative fishing, co-traveling, and play. Our research demonstrates that aquatic subsidies seen in

omnivores can be highly influential to obligate carnivores leading to high population density and altered

social structure.

Introduction

Energetic subsidies through the transfer of resources between terrestrial and aquatic ecosystems

can strongly influence food webs and population dynamics (Polis, Anderson, & Holt, 1997). The transfer

of energy from large emergences of aquatic insects to terrestrial systems is the canonical example of this

phenomenon (Nakano & Murakami, 2001), but fish-consuming terrestrial vertebrates can also play an

important role. Salmon systems famously support hyperabundant bear (*Ursus arctos*) populations with

modified social dynamics due to the need to tolerate conspecifics at the point sources (e.g., waterfalls)

and linear features (e.g., streams) where salmon become accessible (Egbert & Stokes, 1974; Gende &

Quinn, 2004). Similarly, Rose and Polis (2016) found coastal coyote (*Canis latrans*) populations

subsidized by marine resources to be on average 4-5 times more abundant than inland populations.

Several other omnivorous terrestrial carnivores forage in marine or freshwater systems such as gray

wolves (*Canis lupus*) (Darimont, Paquet, & Reimchen, 2008), raccoons (*Procyon lotor*) (Carlton &

Hodder, 2003) and arctic foxes (*Vulpes lagopus*) (Roth, 2002), but aquatic subsidies in obligate terrestrial

carnivores are thought to be rare. Of the obligate carnivores, jaguars (*Panthera onca*) have been most

frequently linked to aquatic resources such as caiman and chelonians (Escobar-Lasso et al., 2016;

Guilder, Barca, Arroyo-Arce, Gramajo, & Salom-Pérez, 2015; Silveira, Ramalho, Thorbjarnarson, &

Magnusson, 2010), although terrestrial mammals still dominate their diet.

A resource-rich environment may enable multiple individuals to share a common resource, which

typically results in higher animal densities (Newsome, Ballard, Dickman, Fleming, & Ven, 2013), smaller

home-ranges and increased social interactions among individuals according to the resource dispersion

hypothesis (Macdonald, 1983). Most large felids, including the jaguar, are considered solitary species

with social interactions generally limited to courtship and territorial disputes (Sunquist and Sunquist

2002; Kleinman & Eisenberg 1973). However, the degree of intraspecific tolerance and sociality can

change with resource availability (Elbroch, 2017; Elbroch & Quigley, 2017). The Pantanal wetlands in

central South America harbors abundant terrestrial and aquatic jaguar prey resources. Here, jaguars reach

high population densities (Soisalo & Cavalcanti, 2006) and use smaller home-ranges than elsewhere in

their range (Morato et al., 2016), but spatial tolerance of conspecifics, defined by overlapping home-

ranges, seem to vary. Azevedo and Murray (2007) found that jaguars maintained exclusive territories

while Cavalcanti and Gese (2009) found extensive overlap among males but not among females.

Consumption of aquatic prey, mainly caiman (*Caiman yacare*), varies among study areas from

approximately 20% (Azevedo and Murray 2007; Cavalcanti and Gese 2010) to 30% (Perilli, Lima,

Rodrigues, & Cavalcanti, 2016) with terrestrial mammals still constituting the majority of the diet in all

prior research. In addition, domestic cattle are abundant and make up a considerable portion of the jaguar

diet in the southern Pantanal where most research has been conducted to date. Thus, it is currently unclear

whether aquatic subsidies have altered the abundance or social structure of these jaguar populations, as

has been demonstrated with other omnivorous carnivores.

In contrast to the southern Pantanal, the ecology of jaguars in the wetter northern portion (i.e.,

within the Brazilian state of Mato Grosso) remains largely unstudied. Taiamã Ecological Station

(hereafter, Taiamã), a small and seasonally flooded protected area (115.55 km^2), was suggested to have an

unusually high jaguar density based on frequent jaguar sightings (Kantek & Onuma, 2013). A large

presence of jaguars in this area is further made obvious by abundant jaguar footprints, scats, scratch

marks on trees and large caiman carcasses scattered along the river margins (this study). Taiamã (16°50'

S and 57°35' W, Figure 1) was established in 1981 to protect this unique wetland ecosystem and its high

diversity and abundance of fish species (ICMBIO 2017). Importantly, there are no human settlements or

cattle ranching within or in the immediate vicinities of Taiamã and fishing is strictly prohibited (ICMBIO

2017). Our objectives were to study the ecology of jaguars at Taiamã to determine if the presence of

abundant aquatic resources affect their diet, social behavior and ultimately population density, and to

place these findings in the context of previous literature on jaguar ecology. To accomplish these

objectives, we collected jaguar scats for dietary analysis, estimated jaguar density and documented social

interactions using camera traps and telemetry data from 13 jaguars. This research details a rare example

of how allochthonous resource subsidies can be so prolific that they influence the behavior and social

organization of an apex predator.

Figure 1 Study area. Map of study area at Taiamã Ecological Station within the Pantanal wetlands in Mato Grosso, Brazil.

96
97

Results

In total we operated 59 camera stations for 8,065 days between 2014 and 2018 (Table 1; Figure 2).

Jaguars were detected in 95% of the cameras (n = 56) and were the most frequently detected mammal

(Figure 3). We obtained 1,594 videos of jaguars (excluding the photos from the paired cameras in 2018),

representing 385 individual weekly detections with the number of detections per individual ranging from

one to 34 (mean 5.6 ± 5.9 SD). We detected 12 out of the 13 telemetered jaguars with our camera traps.

We detected an additional 29 females, 21 males, two individuals of unknown sex, and four cubs,

representing a total of 69 unique individuals. The maximum number of unique jaguars captured by one

camera was 15 across all 3 years including 9 just in 2015. Both right and left flanks were obtained for

most of the individuals, despite our unpaired camera setup in 2014 and 2015. Only one jaguar was

captured only on her left side but was still included in the dataset since the closest female with only a

right-side capture was detected 20 km away and had a completely different rosette pattern type. We were

therefore able to retain all identified jaguars in the analyses. The highest minimum number of jaguars

known to be present within the study area within one camera session was 40 (Table 1).

Table 1. Camera trapping summary statistics. M = male, F= female, U = unknown sex, C = cub.
[1]Capture defined as an individual per camera-trap per occasion. [2]Number of jaguars known to be present
in the study area based on telemetry or camera detections.

Camera session	1 (Aug-Oct 2014)	2 (Mar-Jun 2015)	3 (Jul-Nov 2015)	4 (Aug-Oct 2018)
Length	56	126	126	65
Season	Dry	Wet	Dry	Dry
Camera days	2069	2125	2366	1271
Camera stations	39	20	24	20
Captures[1]	108	165	183	135
Unique individuals	26 (13M, 12F, 1U)	33 (15M, 16F, 1U, 1C)	28 (17M, 10F, 1C)	25 (8M, 14F, 3C)
GPS-collared	8 (5M, 3F)	6 (4M, 2F)	4 (2M, 2F)	0
Min known to be alive[2]	31	40	31	27

Figure 2 and Supplemental Animation. Camera trap locations and jaguar movement at Taiamã Ecological Station. A) Camera trap placement along the Paraguay and Bracinho rivers. B) GPS locations of six resident males and one transient. C) GPS locations of five resident females and one transient

Figure 3 Total number of mammal detections. Number of detections of medium and large mammal species at Taiamã Ecological Station using camera traps from 2014-2018.

The telemetered jaguars spent on average 96% of the time within the study area (Table 2; Figure

2). Estimated jaguar density was 12.4 per 100 km² and mean capture probability per weekly occasion was

0.18 (0.15 - 0.21 Bayesian Credible Interval; BCI) for females and 0.27 (0.24 - 0.31 BCI) for males.

Density was similar across the three dry season sessions at 12.3 (10.1 - 14.4), 11.9 (10.1 - 12.9) and 11.3

(9.4 - 13.1) jaguars per 100 km², in session 1, 3, and 4, respectively. The wet season estimate from session

2 was 14.3 (12.2 - 15.4) jaguars per 100 km² (Table 2), even though some 40% of the Taiamã land area

becomes seasonally flooded.

Table 2 Model parameter estimates of the posterior distributions for jaguar density per 100 km²
and detection parameters in Taiamã Ecological Station across four camera trapping sessions. $\tilde{p}$ is
the fraction of time jaguars spent within the bounds of the trapping area as estimated by telemetry data
and p is the probability of detection within a weekly occasion.

Parameter	Mean	2.50%	50%	97.50%
Session 1 (Aug-Oct 2014)	12.28	10.13	12.26	14.43
Session 2 (Mar-Jun 2015)	14.30	12.20	14.47	15.41
Session 3 (Jul-Nov 2015)	11.90	10.12	12.02	12.95
Session 4 (Aug-Oct 2018)	11.26	9.36	11.26	13.05
$\tilde{p}$	0.96	0.82	0.97	1.00
p-females	0.18	0.15	0.18	0.21
p-males	0.27	0.24	0.27	0.31

Jaguar diet

We identified nine prey items in 138 jaguar scats collected from 2013-2018. The jaguar diet was dominated by three taxonomic groups: reptiles (55%), fish (46%) and mammals (11%) (Figure 4A). These findings were corroborated by camera data showing jaguars feeding on fish and caimans, and other field observations such as large caiman skulls and fish carcasses scattered along the rivers. Fish remains found in the scats could not be identified to species, however, the camera data revealed jaguars capturing several types of fish such as thorny catfish (*Doradidae*) (Figure 4D), pacu (*Piaractus mesopotamicus*), red-bellied piranha (*Pygocentrus nattereri*), and large tiger catfish (*Pseudoplatystoma fasciatum*) (Figure 4C). Most mammals consumed were semi-aquatic (i.e., capybaras; 7%) (Figure 4B). Rare prey items included green iguana (*Iguana iguana*), Brazilian porcupine (*Coendou prehensilis*), freshwater crab, and gray four-eyed opossum (*Philander opossum*) (Figure 4B). The high frequency of occurrence of fish and reptiles in the scats found a Taiamã was unparalleled with other studies throughout the jaguar's range (Figure 5). Jaguars foraged for fish mainly by searching while walking slowly through the water (Figure 6 A, B) and attacking rapidly when spotting a fish (Figure 6, C). Jaguars also looked for fish while sitting still in the water (Figure 6, D).

Figure 4 Relative percentage of occurrence of prey items found in jaguar scats (n = 138) in Taiamã

Ecological Station between 2013 and 2018. (A) Main food groups: reptiles, fish, and mammals. (B) All

identified prey items. (C) F8 carrying a tiger catfish (*Pseudoplatystoma fasciatum*). (D) M4 after catching

a thorny catfish (*Doradidae* spp.).

Figure 5. Frequency of occurrence of major vertebrate prey taxa (mammals, reptiles, fish, and birds) found in jaguar diets documented by different studies throughout the geographic range of jaguars. Only peer-reviewed studies with a sample size of at least 20 scats or kill sites were included. See Supplementary file 3 for more information on sample sizes and study locations.

Figure 6. Videos of jaguar fishing behavior. A) Mother and cub searching for fish together, B) Jaguar looking for fish by walking slowly through water, C) Failed capturing attempt, D) 'Sit and wait' foraging technique.

Social interactions

All resident GPS-collared males had overlapping 95% AKDE home ranges, except for M3 and M4 (Figure 7A). M1 overlapped from 17.6 to 25.5% with M2 and M3, respectively, while M2 and M3's home ranges overlapped by 64.7%. M5 overlapped with M1 by 7.6% and 16.1% with M3. The 95% AKDE home range also overlapped extensively between GPS-collared females (Figure 7B). F2's home range overlapped by 58.6% with F1 and the home ranges of F3 and F4 overlapped to 21.6%. F1 and F2 were unrelated while F3 and F4 were mother and daughter (D.L.Z Kantek, personal communication, January 2021). Since these home range overlap estimates were exclusively derived from telemetered individuals, they fail to account for spatial overlap with uncollared jaguars. Approximately 4-15 jaguars (including collared and uncollared) were detected with the cameras overlapping each home range.

Figure 7. 95% AKDE home ranges for resident jaguars GPS-collared at Taiamã Ecological Station between 2014-2015. A) Five male jaguars, and B) 4 female jaguars.

We documented 80 independent social interactions between adult jaguars based on the camera (n=40) and GPS data (n=40). The majority of the interactions were between males and females (M-F 85%), often displaying courtship behavior (Figure 8B and C) throughout the wet and dry seasons. The M-F interactions involved 29 individual jaguars (14 males and 15 females). One male and female mated with each other both in 2015 and 2018. We documented 12 interactions of the same sex (one F-F and 11 M-M) consisting of 10 individuals (two females and eight males). Two GPS-collared males, M2 and M1, moved continuously within approximately 41-182 m of each other for two days. Eight months later the same two males were captured with the same camera 5 min apart. M2 moved continuously with another GPS-collared male (M3) for 10 hours within 4-38 m of each other, again two days later (2 hours; 14 m apart) and were close again two weeks later (1 hour; 159 m). None of the three males were related (D.L.Z Kantek, personal communication, January 2021). Two other males (M7 and M21) were seen fishing together (Figure 8D) and walking past another camera together one month later. Two males (M29 and

197   M27) spent 30 min in front of a camera "playing" (Figure 8E and F). Two females (F2 and F4) were

198   documented in the same location 12 min apart (Figure 8G). We lack information on potential relatedness

199   among the interacting uncollared jaguars.

A

Type of interaction	Number of camera interactions	Number of GPS interactions	Photo
Female + male	33	35	B, C
Male + male	6	5	D, E, F
Female + female	1	0	G

200   **Figure 8. (A) Number and type of social interactions among adult jaguars determined by GPS data**
201   **and camera trap data.** (B-C) Male (M5) and female (F4) mating behavior, (D) Two adult males (M7 and
202   M21) fishing together, (E-F) two adult males (M29 and M27) playing, (G) Two females (F2 and F4)
203   captured with the same camera 12 minutes apart.

Discussion

With a density of 12.4 jaguars per 100 km², our results indicate that Taiamã is home to the

highest density jaguar population described to date. Other high jaguar density estimates using spatial-

capture recapture analysis include 4.4-4.5 jaguars/100 km² from the Peruvian Amazon (Tobler et al.,

2013; Tobler et al., 2018) and 4.4 jaguars/100 km² in the Venezuelan Llanos (Jędrzejewski et al., 2017).

Our density estimate using SCR was 6.0 jaguars per 100 km² (Supplementary file 2), but as discussed

previously, this is an underestimate due to the elongated home ranges of jaguars at Taiamã (Figure 2). As

a very minimum estimate, using just our 13 *telemetered* individuals, which we knew were all present in

2015, with on average 96% of GPS locations contained within the boundaries of our study area (236.7

km²), would suggest a density of approximately 5.4 jaguars per 100 km², without considering the

additional 56 individuals detected with cameras.

Reptile consumption by jaguars at Taiamã was also the highest observed to date (Figure 5)

although they can be a common component of jaguar diets in seasonally flooded habitats (Cavalcanti and

Gese 2010; Azevedo and Verdade 2012; Silveira et al. 2010). This is particularly the case in the flooded

várzea forests of the Amazon where common terrestrial mammalian prey are absent during the flood

pulse (Silveira et al., 2010). However, the heavy reliance on fish appears unique to the jaguars at Taiamã

and has been rarely documented in the literature (Figure 5) despite jaguars being commonly associated

with rivers and waterbodies (Morato et al. 2018; Kanda et al. 2019). Prior to the extensive fish

consumption observed here, the only records include anecdotal observations of jaguars fishing

summarized in Gudger (1946) and more recently Emmons (1987) found fish in two jaguar scats. Other

felids living in semi-aquatic habitats such as tigers (*Panthera tigris*) in the Sundarbans still consume

mostly terrestrial mammals (Aziz, Islam, & Groombridge, 2020). The fishing cat (*Prionailurus*

viverrinus) may be the most similar example of a piscivorous felid akin to the jaguars at Taiamã (Ganguly

& Adhya, 2020).

Ecological theory would predict that a high density of an apex predator, subsidized by

allochthonous resources, would have consequences for the surrounding community through modification

of top-down forces (Leroux & Loreau, 2008). It is unclear whether this exceptionally high abundance of

jaguars, an opportunist predator supported by abundant aquatic resources, has had a negative impact on

the terrestrial mammal community at Taiamã. Common terrestrial mammalian prey for jaguars such as

capybara, tapirs, peccaries (*Tayassuidae*), giant anteater, and marsh deer (Emmons, 1987; Novack, Main,

Sunquist, & Labisky, 2005; Weckel, Giuliano, & Silver, 2006) are present at Taiamã (De Lázari et al.

2013; Mauro et al. 1998), yet mammals were only found in 11% of the scats. The seasonally flooded

landscape at Taiamã may play a stronger role in determining the abundance and distribution of terrestrial

species (De Lázari et al., 2013). For example, mammals that are suited to aquatic life such as capybaras

are common, but rarely feature in the diet of these jaguars. It is possible that the abundance of aquatic

resources has resulted in a highly specialized jaguar diet, potentially benefitting the terrestrial community

through apparent mutualism (Holt & Bonsall, 2017) and provisioning of carcasses (Escobar-Lasso et al.,

2016). However, even if jaguars only occasionally take certain terrestrial prey, they may still suppress

those species due to the large number of jaguars.

Jaguars at Taiamã may have adapted their social system in response to abundant prey resources,

similar to how brown bears are more tolerant of conspecifics during periods of high salmon abundance

(Egbert & Stokes, 1974). According to optimal foraging theory, individuals widen their dietary niche by

switching to alternative food sources to avoid competition when food is limited. When food is abundant,

individual prey selectivity increases and the diversity of prey taken decreases (MacArthur and Pianka

1966). The almost complete dominance of just two vertebrate taxa — fish and caiman — in the diet of

jaguars at Taiamã further suggests food is abundant and intraspecific competition is relaxed. Home-range

overlap among jaguars, in particular males, has been documented to varying degrees in other areas, but

may not be an indication of social tolerance since individuals may still avoid each other temporally

through the use of scent and scrape marking (Harmsen, Foster, Gutierrez, Marin, & Doncaster, 2010) or

vocalizations. Outside of male and female courtship behavior, camera-trapping observations of adult

jaguars together, simultaneous GPS locations and/or direct observations are exceedingly rare in the

literature. Schaller and Crawshaw (1980) described an incident where a jaguar hunter saw three males

courting one female at the same time. Cavalcanti and Gese (2009) found two males sharing a carcass

together and two other males moving within 200 m of each other on three occasions. Our GPS data

revealed one male spending several hours, sometimes days within 200 m of two other males and our

cameras detected another two males walking and fishing together. In addition, the overlapping home-

ranges of a pair of females and the detection of a female followed by another female 12 min later, suggest

that jaguars can be more social than previously thought. Jaguars may share kills with offspring, mates or

even same-sex individuals (Cavalcanti & Gese, 2009), but observations of jaguars hunting together have

not, to our knowledge, been published previously. In addition to the two adult males mentioned earlier, a

female and her older cub were frequently observed fishing together (Figure 6, B). We cannot determine

whether this form of cooperative fishing improved fish capture efficiency, or whether the jaguars just

tolerated the close proximity of each other without deriving any prey-acquisition benefits.

Further indication that high resource abundance at Taiamã affects the social organization of

jaguars comes from clear evidence of social play which in animals is thought to occur when all biological

needs have been met (Bekoff, 1972). The incident of play between M27 and M29 started with passive

submissive patterns such as rolling over, facial rub, and face-paws (Bekoff, 1972). The play then

transitioned to play-fighting with wrestling, soft bites, and mounting. Play in juvenile mammals is

essential for behavioral development but play among adults is rarely described and its motivations are

more difficult to interpret. Latour (1981) suggested adult play in male polar bears was used to practice

and enhance social skills for future agonistic interactions. While our results indicate an increased social

tolerance among jaguars at Taiamã, some males displayed typical territorial behavior such as urine

spraying, scraping and roaring, suggesting that intraspecific social behavior is a complex phenomenon

that demands further investigation.

Conclusion

Our study demonstrates that aquatic subsidies, frequently described in omnivores, can be common in

obligate carnivores as well. Whether our results are generalizable to other jaguar populations in

seasonally flooded habitats is confounded by the presence of domestic cattle, another subsidy, in other

areas. The flooded nature of the landscape surrounding Taiamã has largely prevented the expansion of

cattle farming, yet this region faces many threats and efforts to increase protected areas are critically

needed to ensure future persistence of jaguars in the area (Cardoso et al., 2019; Quigley & Crawshaw,

1992). In an effort to prioritize conservation efforts throughout the jaguar's distribution, Sanderson et al.

(2002) stated "the goal is not to determine the most important site for jaguar conservation overall, or the

most important site in a given country, but rather to find the most important sites for ecologically distinct

populations of jaguars". In this study we have provided evidence that Taiamã is a stronghold for jaguars,

but also revealed aspects of jaguar ecology that are rare elsewhere; an extensive piscivorous diet

facilitating an extremely dense population with a more fluid social structure characterized by high levels

of individual tolerance. This work further sheds new light on the relationship between carnivore

socioecology and prey abundance and distribution, showing that the modal pattern of solitary hunting

observed in the vast majority of hyper-carnivores can be broken given the nature, productivity and

distribution of the resource base. Future research is needed to understand the role jaguars play linking

aquatic and terrestrial food webs and the implications of this highly subsidized population on the

surrounding community.

Materials and methods

Study Area

The study area encompassed Taiamã Island and contiguous Sararé Island bounded by the Paraguay and

Bracinho rivers (Figure 1). The area is characterized by seasonally flooded grasslands with numerous

seasonal and permanent lakes and riparian gallery forests growing along the rivers. The climate is defined

by a dry (April-September) and wet season (October-March) with an average annual precipitation of

1,200-1,300 mm (Mamede & Alho, 2006). The maximum river level occurs between February-June and

the lowest in August-November. The fluctuating water levels substantially alter the availability of

terrestrial vs aquatic habitats (Figure 1) and determine local ecological patterns and processes, including

the distribution and abundance of wildlife (Alho, 2008; Desbiez, Bodmer, & Tomas, 2010). In addition to

jaguars, the reserve supports several other vertebrates well-adapted to water, most notably caimans

(*Caiman yacare*) and capybaras (*Hydrochoerus hydrochaeris*). Other known jaguar prey species present

include giant anteaters (*Myrmecophaga tridactyla*), Brazilian tapir (*Tapirus terrestris*), marsh deer

(*Blastocerus dichotomus*), and Brazilian porcupine (*Coendou prehensilis*) (De Lázari et al. 2013; Mauro

et al. 1998; this study).

Camera dataset

We carried out three camera trapping surveys between 2014 and 2018. The absence of roads or trails

coupled with the prolonged flood pulse of the landscape and the high jaguar density presented unique

logistical challenges for field work and prevented any movement by car or foot. All camera deployments

were therefore boat-based and restricted to the major waterways and, thus, prevented the use of a

systematic camera grid. We used unbaited Bushnell (Trophy Cam 119636, Overland Park, Kansas) and

Browning (Dark Ops BTC-6, Birmingham, AL) camera traps separated by approximately 2 km along the

Paraguay and Bracinho rivers. The two waterways were parallel to each other and separated by 4-6 km

(Figure 1). Camera availability varied across the years resulting in 39 and 27 single cameras used in 2014

and 2015, respectively. In 2018 each camera station consisted of two paired cameras (20 stations in total).

In 2014 and 2015 the cameras were set to video mode (20 sec/video) and operated for 56 (August-

October) and 252 days (March-November), respectively. In 2018 one camera was set to video and the

other paired camera was set to take photos (three per burst) for 65 days (August-October). Cameras were

set approximately 50 cm off the ground on trees with batteries and SD cards checked every three weeks.

The study area encompassed an area of approximately 236.7 km², which is larger than a jaguar home

range in this area (Morato et al., 2016) as recommended for accurate density estimation (Tobler & Powell,

2013). Individual jaguars were identified in each photo or video based on their unique coat patterns. The

camera detections were collapsed into weekly occasions for density estimation. The 252-day long camera

deployment in 2015 was split into two sessions to assume demographic closure resulting in four sessions

of 8, 18, 18 and 10 weekly occasions each. Camera trap sampling details are summarized in Table 1.

Telemetry dataset

Seven male and six female jaguars were captured between December 2011 and April 2016 as part of

concurrent studies (Morato et al. 2016; Miyazaki et al., 2016; Cardoso et al., 2020) approved by Instituto

Chico Mendes de Conservação da Biodiversidade (SISBIO number 30896-3). The jaguars were captured

using foot snares and immobilized with tiletamine and zolazepam (10 mg/kg). Two of the jaguars were

fitted with VHF radio-collars and the remaining (n = 11) were fitted with Lotek-Iridium GPS collars

(Lotek Wireless, Newmarket, Canada) programmed to record location data every hour and to

automatically drop-off the animal after 400 days.

Density estimation

A fundamental issue with density estimation is how to deal with a lack of geographic closure given that

some fraction of detected individuals primarily lives outside the study area. Capture-recapture models

(CR; Otis et al. 1978) estimate abundance and derive density by dividing the number of individuals by the

effective sampling area. However, the effective sampling area is imprecisely estimated using a buffer strip

around the study area of either the full Mean Maximum Distance Moved (MMDM) or ½ MMDM from

the capture data, which are considered as a surrogate for home range diameter or radius. Thus capture-

recapture models can be sensitive to arbitrarily defined effective sampling area. In the past decade,

Spatially explicit Capture-Recapture models (SCR; Borchers and Efford 2008; Royle and Young 2008)

have become a preferred method for density estimation over conventional non-spatial capture-recapture

techniques. SCR models solve the geographic closure problem by incorporating the spatial location of

detections to estimate the activity center of each animal, and the density of activity centers is then treated

as the population density.

While SCR models present a useful advance over conventional CR models, they assume isotropic

(i.e. circular) home ranges such that detection probability of each individual falls off equally with distance

from its activity center. Our telemetry data revealed that the direction of jaguar movement at Taiamã

follows the rivers closely and the home ranges are therefore elongated and highly anisotropic (Figure 2).

In such a situation, SCR models may still perform well if sufficient detectors are placed in two

dimensions capturing both the major axis of movement (along the rivers in our case) and the minor axis

(perpendicular to the rivers) (Efford, 2019). The bias can then be corrected by applying anisotropic

detection models (Murphy et al., 2016). Since our camera deployment was boat-based and therefore

restricted to the rivers surrounding Taiamã island, we were unable to effectively capture jaguar movement

in the minor axis. Further, the telemetered jaguars did not uniformly share the same home range

orientation (Figure 2) which is necessary to fit anisotropic detection models. Applying a SCR model to

this unusual dataset where the detectors are aligned with elongated animal home ranges would therefore

lead to an overestimation of the spatial scale parameter and consequently underestimate density (Efford,

2019).

For data-rich applications where a representative portion of the population has been telemetered,

the extent of geographic closure violations can be corrected for directly (Ivan et al. 2013a, 2013b) to

estimate population density without the restrictive assumptions of circular home ranges or the size of the

effective sampling area. This method uses a conventional Huggins closed-capture model augmented by

telemetry data. The study area is first determined by applying a minimum convex hull around the

detectors. Crucially, rather than assuming each captured individual is fully present within the study area

such as in a standard CR model (1 in the equation 1 numerator) (i.e., the study area is geographically

closed) (eq. 1), residency ($\tilde{p}$) is instead used by estimating the proportion of time the telemetered

individuals spend within the area (eq. 2). p_i is the detection probability for individual i, M_{t+1} is the total

number of animals detected, and t is the number of occasions.

383
$$N = \sum_{i=1}^{Mt+1} \left(\frac{1}{p_i}\right) \tag{1}$$

384
$$N = \sum_{i=1}^{Mt+1} \left(\frac{\tilde{p}}{p_i}\right) \tag{2}$$

We estimated jaguar abundance (N) allowing for variation in capture probability (p) between males and

females as recommended by Sollmann et al. (2011). We used telemetry data to estimate $\tilde{p}$ as the grand

mean among individuals of the Binomial probability parameter in a model for the number of telemetry

points inside the study area relative to outside. We included a random intercept in the logit-linear model

for $\tilde{p}$ so that one individual's data would not dominate the estimate. Density was then estimated by

correcting N with $\tilde{p}$ (Eq. 2) and then dividing by the study area. All models were implemented in JAGS

(Plummer, 2003) accessed through the R package JagsUI (Kellner, 2019). Code is available in

Supplementary file 1.

Diet analysis

Jaguar scats were collected opportunistically during field work. We did not genetically confirm

that the scats were from jaguars but there are only two camera detections to date of another large felid

species (*Puma concolor*) in the study area compared to more than 1,600 jaguar detections (this study). We

were therefore confident that the vast majority (if not all) of the scats we collected were jaguar scats.

Once collected, scat samples were stored in individually-labeled plastic bags and kept frozen until

laboratory analysis. Undigested remains found in the scats were first isolated by rinsing the scats using a

fine mesh, then dried and identified to the lowest taxon possible using a dissecting scope. We recorded

jaguar diet using relative frequency of occurrence (number of occurrences of each prey item divided by

the total number of scats). We compared our results to other published jaguar diet studies using either scat

or kill site analysis with a sample size of at least 20. For comparative purposes, we grouped prey items

into 'mammals', 'reptiles', 'fish' and 'birds' and standardized frequency of occurrence values to sum to

100%.

Social interactions among jaguars

We investigated shared space use among the resident jaguars collared in 2014-2015 (n=9), by

quantifying the proportion of overlap at the 95% individual utilization distribution using the

Autocorrelated Kernel Density Estimator (AKDE) in the R package 'CTMM' (Calabrese, Fleming, &

Gurarie, 2016). Because jaguars exhibiting home-range overlap can still be avoiding each other

temporally, we analyzed the frequency of social interactions by assessing simultaneous locations from

both GPS data and camera data. GPS interaction rates were calculated using the package wildlifeDI (Long

& Nelson, 2013) in R and defined as the number of times two GPS-collared jaguars were within 200 m of

each other (Cavalcanti & Gese, 2009; Elbroch & Quigley, 2017). We defined camera interaction rates as

the number of times two jaguars were observed together or within 30 min of each other. We collapsed

interactions between individuals per day such that multiple contacts within the same day were not counted

as new interactions for both the GPS and camera data.

Acknowledgments

This study was funded by Helge Ax:son Johnson Foundation and Oregon State University. We thank

Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio) and Universidade do Estado de

Mato Grosso (UNEMAT) for facilitating this research. We also thank Manaav Kamath, Aurea Ignácio,

Claumir Muniz, Amabile Furlan and Derrick Campos for logistical support and field work assistance, as

well as Gerlane Costa for help with prey remain identification.

References

Alho, C. (2008). Biodiversity of the Pantanal: response to seasonal flooding regime and to environmental
degradation. *Brazilian Journal of Biology*, *68*(4), 957–966.

Azevedo, F. C., & Murray, D. L. (2007). Spatial organization and food habits of jaguars (*Panthera onca*)
in a floodplain forest. *Biological Conservation*, *137*(3), 391–402.

Azevedo, F., & Verdade, L. (2012). Predator-prey interactions: Jaguar predation on caiman in a
floodplain forest. *Journal of Zoology*, *286*(3), 200–207. https://doi.org/10.1111/j.1469-
7998.2011.00867.x

Aziz, M. A., Islam, M. A., & Groombridge, J. (2020). Spatial differences in prey preference by tigers
across the Bangladesh Sundarbans reveal a need for customised strategies to protect prey
populations. *Endangered Species Research*, *43*, 65–74. https://doi.org/10.3354/esr01052

Bekoff, M. (1972). The development of social interaction, play , and metacommunication in mammals:
An ethological perspective. *The Quarterly Review of Biology*, *47*(4), 412–434.

Borchers, D., & Efford, M. (2008). Spatially explicit maximum likelihood methods for capture–recapture
studies. *Biometrics*, (64), 377–385. https://doi.org/10.1111/j.1541-0420.2007.00927.x

Calabrese, J. M., Fleming, C. H., & Gurarie, E. (2016). ctmm: an R package for analyzing animal
relocation data as a continuous-time stochastic process. *Methods in Ecology and Evolution*, *7*, 1124–
1132. https://doi.org/10.1111/2041-210X.12559

Cardoso, H. M., Morato, R. G., Miyazaki, S. S., Pereira, T. D. C., Araújo, G. R. De, & Kantek, D. L. Z.
(2019). Effectiveness of protected areas for jaguars: the case of the Taiamã Ecological Station in
Brazil, 0–4.

Carlton, J. T., & Hodder, J. (2003). Maritime mammals: terrestrial mammals as consumers in marine
intertidal communities. *Marine Ecology Progress Series*, *256*, 271–286.

Cavalcanti, S. M., & Gese, E. M. (2009). Spatial ecology and social interactions of jaguars (*Panthera
onca*) in the southern Pantanal, Brazil. *Journal of Mammalogy*, *90*(4), 935–945.
https://doi.org/10.1644/08-MAMM-A-188.1

Cavalcanti, S. M., & Gese, E. M. (2010). Predation patterns of jaguars (*Panthera onca*) in a seasonally

flooded forest in the southern region of Pantanal, Brazil. *Journal of Mammalogy*, *91*(3), 722–736. https://doi.org/10.1644/09-MAMM-A-171.1.Key

Darimont, C. T., Paquet, P. C., & Reimchen, T. E. (2008). Spawning salmon disrupt trophic coupling between wolves and ungulate prey in coastal British Columbia, *12*, 1–12. https://doi.org/10.1186/1472-6785-8-14

De Lázari, P. R., dos Santos-Filho, M., Canale, G. R., & Graipel, M. E. (2013). Flood-mediated use of habitat by large and midsized mammals in the Brazilian Pantanal. *Biota Neotropica*, *13*(2), 70–75.

Desbiez, A. L. J., Bodmer, R. E., & Tomas, M. (2010). Mammalian densities in a neotropical wetland subject to extreme climatic events. *Biotropica*, *42*(3), 372–378.

Efford, M. (2019). Non-circular home ranges and the estimation of population density. *Ecology*, *100*(2), e02580. https://doi.org/10.1002/ecy.2580

Egbert, A., & Stokes, A. (1974). The social behaviour of brown bears on an Alaskan salmon stream. *Biology and Management*, *3*, 40–56.

Elbroch, L. M. (2017). Pumas: solitary but social? *Frontiers in Ecology and the Environment*, *15*(3), 168–169. https://doi.org/10.1002/fee.1479

Elbroch, L. M., & Quigley, H. (2017). Social interactions in a solitary carnivore. *Current Zoology*, *63*(4), 357–362. https://doi.org/10.1093/cz/zow080

Emmons, L. H. (1987). Comparative feeding ecology of felids in a neotropical rainforest. *Behavioral Ecology and Sociobiology*, *20*(4), 271–283. https://doi.org/10.1007/BF00292180

Escobar-Lasso, S., Gil-fernandez, M., Sáenz, J., Carrillo-Jimenez, E., Wong, G., & Fonseca, L. G. (2016). Inter-trophic food provisioning between sea and land: The jaguar (*Panthera onca*) as provider of sea turtle carcasses to terrestrial scavengers. *International Journal of Conservation Science*, *7*(4), 1081–1094.

Ganguly, D., & Adhya, T. (2020). How fishing cats *Prionailurus viverrinus* fish: Describing a felid's strategy to hunt aquatic prey. *BioRxiv*, *[Preprint]*. https://doi.org/https://doi.org/10.1101/2020.04.24.058925

Gende, S. M., & Quinn, T. P. (2004). The relative importance of prey density and social dominance in determining energy intake by bears feeding on Pacific salmon. *Canadian Journal of Zoology*, *82*, 75–85. https://doi.org/10.1139/Z03-226

Guilder, J., Barca, B., Arroyo-Arce, S., Gramajo, R., & Salom-Pérez, R. (2015). Jaguars (*Panthera onca*) increase kill utilization rates and share prey in response to seasonal fluctuations in nesting green turtle (*Chelonia mydas mydas*) abundance in Tortuguero National Park, Costa Rica. *Mammalian Biology*, *80*(2), 65–72. https://doi.org/10.1016/j.mambio.2014.11.005

Harmsen, B. J., Foster, R. J., Gutierrez, S. M., Marin, S. Y., & Doncaster, C. P. (2010). Scrape-marking behavior of jaguars (*Panthera onca*) and pumas (*Puma concolor*). *Journal of Mammalogy*, *91*(5), 1225–1234.

Holt, R. D., & Bonsall, M. B. (2017). Apparent competition. *Annual Review of Ecology and Systematics*, *48*, 447–471.

ICMBio. 2018. *Livro Vermelho da Fauna Brasileira Ameaçada de Extinção: Volume II - Mamíferos*. Avaliable at: https://www.icmbio.gov.br/portal/images/stories/comunicacao/publicacoes/publicacoes-diversas/livro_vermelho_2018_vol2.pdf. Accessed: 22/10/2019.

ICMBio. 2017. *Plano de Manejo da Estação Ecológica de Taiamã.* Instituto Chico Mendes de
Conservação da Biodiversidade, Brasília, Brasil. Available at:
http://www.icmbio.gov.br/portal/images/stories/plano-de-
manejo/plano_de_manejo_esec_Taiamã_vs_070617.pdf. Accessed: 20/01/2020.

Jędrzejewski, W., Puerto, M. F., Goldberg, J. F., Hebblewhite, M., Abarca, M., Gamarra, G., … Schmidt,
K. (2017). Density and population structure of the jaguar (*Panthera onca*) in a protected area of Los
Llanos, Venezuela, from 1 year of camera trap monitoring. *Mammal Research*, *62*(1), 9–19.
https://doi.org/10.1007/s13364-016-0300-2

Kanda, C. Z., Morato, R. G., Paula, R. C. De, & Rampim, L. E. (2019). Spatiotemporal dynamics of
conspecific movement explain a solitary carnivore's space use. *Journal of Zoology*, *308*, 66–74.
https://doi.org/10.1111/jzo.12655

Kantek, D. L. Z., & Onuma, S. S. M. (2013). Jaguar conservation in the region of Taiama Ecological
Station, Northern Pantanal, Brazil. *Publication UEPG. Ciencias Biologicas e Da Saude*, *19*, 69–74.
https://doi.org/10.5212/Publ.Biologicas.v.19i1.0008

Kellner, K. (2019). jagsUI: A Wrapper Around "rjags" to Streamline "JAGS" Analyses.

Kleiman, D.G. & Eisenberg, J.F. (1973) Comparisons of canid and felid social systems from an
evolutionary perspective. Animal Behaviour, 21, 637–659.

Latour, P. B. (1981). Interactions between free-ranging, adult male polar bears (*Ursus maritimus* Phipps):
a case of adult social play. *Canadian Journal of Zoology*, *59*, 1775–1783.

Leroux, S., & Loreau, M. (2008). Subsidy hypothesis and strength of trophic cascades across ecosystems.
Ecology Letters, (11), 1147–1156. https://doi.org/10.1111/j.1461-0248.2008.01235.x

Long, J. A., & Nelson, T. A. (2013). Measuring dynamic interaction in movement data. *Transactions in*
GIS, *17*(1), 62–77. https://doi.org/10.1111/j.1467-9671.2012.01353.x

Macarthur, R. H., & Pianka, E. R. (1966). On optimal use of a patchy environment. *The American*
Naturalist, *100*(916), 603–609.

Macdonald, D. W. (1983). The ecology of carnivore social behaviour. *Nature*, *301*, 379–383.

Mamede, S. B., & Alho, C. J. R. (2006). Response of wild mammals to seasonal shrinking-and-expansion
of habitats due to flooding regime of the Pantanal, Brazil. *Brazilian Journal of Biology*, *66*(664),
991–998. https://doi.org/10.1590/S1519-69842006000600006

Mauro, R., Mourão, G., Coutinho, M., Silva, M., & Magnusson, W. (1998). Abundance and distribution
of marsh deer *Blastocerus dichotomus* (*Artiodactyla: Cervidae*) in the Pantanal, Brazil. *Rev. Ecol.*
Lat. Am., *5*(1–2), 13–20.

Miyazaki, S., Chaves, L., Lara, M. do C., May-Júnior, J., Taques, Ísis, Fritzen, J., Alfieri, A., Ometto, T.,
Durigon, E. L., Araújo, J., Kantek, D. L., & Aguiar, D. (2016). Serological and molecular
investigation of viral agents in free-living jaguars of the Pantanal wetlands, state of Mato Grosso,
Brazil. *Brazilian Journal of Veterinary Research and Animal Science*, *53*(3), 270–279.
https://doi.org/10.11606/issn.1678-4456.bjvras.2016.108947

Morato, R., Connette, G., Stabach, J., Paula, R. De, Ferraz, K., Kantek, D. L. Z., … Leimgruber, P.
(2018). Resource selection in an apex predator and variation in response to local landscape

characteristics. *Biological Conservation*, *228*(October), 233–240.
https://doi.org/10.1016/j.biocon.2018.10.022

Morato, R. G., Stabach, J. A., Fleming, C. H., Calabrese, J. M., Paula, R. C. De, Ferraz, K. M. P. M., …
Leimgruber, P. (2016). Space use and movement of a neotropical top predator: the endangered
jaguar. *Plos One*, *11*(12), e0168176. https://doi.org/10.1371/JOURNAL.PONE.0168176

Murphy, S. M., Cox, J. J., Augustine, B. C., Hast, J. T., Guthrie, J. M., Wright, J., … Plaxico, J. H.
(2016). Characterizing recolonization by a reintroduced bear population using genetic spatial
capture–recapture. *Journal of Wildlife Management*, *80*(8), 1390–1407.
https://doi.org/10.1002/jwmg.21144

Nakano, S., & Murakami, M. (2001). Reciprocal subsidies: Dynamic interdependence between terrestrial
and aquatic food webs. *PNAS*, *98*(1), 166–170.

Newsome, T. M., Ballard, G., Dickman, C. R., Fleming, P. J. S., & Ven, R. Van De. (2013). Home range,
activity and sociality of a top predator, the dingo: a test of the Resource Dispersion Hypothesis.
Ecography, *36*, 914–925. https://doi.org/10.1111/j.1600-0587.2013.00056.x

Novack, A. J., Main, M. B., Sunquist, M. E., & Labisky, R. F. (2005). Foraging ecology of jaguar
(Panthera onca) and puma (*Puma concolor*) in hunted and non-hunted sites within the Maya
Biosphere Reserve, Guatemala. *Journal of Zoology*, *267*(2), 167–178.
https://doi.org/10.1017/S0952836905007338

Otis, D. L., Burnham, K. P., White, G. C., & Anderson, D. R. (1978). Statistical inference from capture
data on closed animal populations. *Wildlife Monographs*, (62), 3–135.

Perilli, M. L. L., Lima, F., Rodrigues, F. H. G., & Cavalcanti, S. M. C. (2016). Can scat analysis describe
the feeding habits of big cats? A case study with jaguars (*Panthera onca*) in southern Pantanal.
PLoS ONE, *11*(3), e0151814. https://doi.org/10.1371/journal.pone.0151814

Plummer, M. (2003). JAGS: A program for analysis of Bayesian graphical models using Gibbs sampling.

Polis, G. A., Anderson, W. B., & Holt, R. D. (1997). Toward an integration of landscape and food web
ecology: The dynamics of spatially subsidized food webs. *Annual Review of Ecology and*
Systematics, *28*, 289–316.

Quigley, H., Foster, R., Petracca, L., Payan, E., Salom, R. & Harmsen, B. 2017. *Panthera onca* (errata
version published in 2018). *The IUCN Red List of Threatened Species* 2017: e.T15953A123791436.
https://dx.doi.org/10.2305/IUCN.UK.2017-3.RLTS.T15953A50658693.en. Accessed on 26 August 2020.

Quigley, H. ., & Crawshaw, P. . (1992). A conservation plan for the jaguar *Panthera onca* in the Pantanal
region of Brazil. *Biological Conservation*, *61*, 149–157.

Rose, M. D., & Polis, G. A. (1998). The distribution and abundance of coyotes: The effects of
allochthonous food subsidies from the sea. *Ecology*, *79*(3), 998–1007.

Roth, J. D. (2002). Temporal variability in arctic fox diet as reflected in stable-carbon isotopes; the
importance of sea ice. *Oecologia*, *133*, 70–77. https://doi.org/10.1007/s00442-002-1004-7

Royle, J. A., & Young, K. V. (2008). A hierarchical model for spatial capture-recapture data. *Ecology*,
89(8), 2281–2289.

Sanderson, E. W., Redford, K. H., Chetkiewicz, C. L. B., Medellin, R. A., Rabinowitz, A. R., Robinson,
J. G., & Taber, A. B. (2002). Planning to save a species: The jaguar as a model. *Conservation*
Biology, *16*(1), 58–72. https://doi.org/10.1046/j.1523-1739.2002.00352.x

Schaller, G. B., & Crawshaw-Junior, P. G. (1980). Movement patterns of jaguar. *Biotropica*, *12*(3), 161–
168.

Silveira, R. Da, Ramalho, E. E., Thorbjarnarson, J. E., & Magnusson, W. E. (2010). Depredation by
jaguars on caimans and importance of reptiles in the diet of jaguar. *Journal of Herpetology*, *44*(3),
418–424.

Soisalo, M. K., & Cavalcanti, S. M. C. (2006). Estimating the density of a jaguar population in the
Brazilian Pantanal using camera-traps and capture-recapture sampling in combination with GPS
radio-telemetry. *Biological Conservation*, *129*(4), 487–496.
https://doi.org/10.1016/j.biocon.2005.11.023

Sollmann, R., Furtado, M. M., Gardner, B., Hofer, H., Jácomo, A. T. A., Tôrres, N. M., & Silveira, L.
(2011). Improving density estimates for elusive carnivores: Accounting for sex-specific detection
and movements using spatial capture-recapture models for jaguars in central Brazil. *Biological
Conservation*, *144*(3), 1017–1024. https://doi.org/10.1016/j.biocon.2010.12.011

Sunquist, M., & Sunquist, F. (2002). *Wild cats of the world*. Chicago: University of Chicago Press.

Tobler, M. W., Carrillo-Percastegui, S. E., Zúñiga Hartley, A., & Powell, G. V. N. (2013). High jaguar
densities and large population sizes in the core habitat of the southwestern Amazon. *Biological
Conservation*, *159*, 375–381. https://doi.org/10.1016/j.biocon.2012.12.012

Tobler, M. W., Garcia, R., Carrillo-percastegui, S. E., Ponce, G., Polisar, J., Zuñiga, A., & Goldstein, I.
(2018). Do responsibly managed logging concessions adequately protect jaguars and other large and
medium-sized mammals? Two case studies from Guatemala and Peru. *Biological Conservation*,
220(January), 245–253. https://doi.org/10.1016/j.biocon.2018.02.015

Tobler, M. W., & Powell, G. V. N. (2013). Estimating jaguar densities with camera traps: Problems with
current designs and recommendations for future studies. *Biological Conservation*, *159*, 109–118.
https://doi.org/10.1016/j.biocon.2012.12.009

Weckel, M., Giuliano, W., & Silver, S. (2006). Jaguar (*Panthera onca*) feeding ecology: Distribution of
predator and prey through time and space. *Journal of Zoology*, *270*(1), 25–30.
https://doi.org/10.1111/j.1469-7998.2006.00106.x