1 **Title: Occipital and parietal cortex participate in a cortical network for**
2 **transsaccadic discrimination of object shape and orientation**
3
4 **Abbreviated Title**: Dissociating feature-dependent transsaccadic mechanisms
5
6

7 **Author(s) and Affiliations:** B. R. Baltaretu[1-3], W. Dale Stevens[1,4], E. Freud[1,4] and J.
8 D. Crawford*[1,2,4,5]
9
10 [1] Centre for Vision Research and Vision: Science to Applications (VISTA) program,
11 York University, Toronto, Ontario M3J 1P3, Canada
12
13 [2] Department of Biology, York University, Toronto, Ontario M3J 1P3, Canada
14
15 [3] Department of Psychology, Justus-Liebig University Giessen, Giessen, Hesse
16 35394, Germany
17
18 [4] Department of Psychology and Neuroscience Graduate Diploma Program, York
19 University, Toronto, Ontario M3J 1P3, Canada
20
21 [5] School of Kinesiology and Health Sciences, York University, Toronto, Ontario
22 M3J 1P3, Canada
23
24
25
26
27 **Corresponding Author:** Bianca R. Baltaretu
28 b.baltaretu@gmail.com
29
30 Otto-Behaghel-Strasse 10F
31 Giessen, Hesse
32 35394
33 Germany

Number of Figures: 3
Number of Supplementary Figures: 0
Number of Tables: 2
Number of Words (Abstract): 197
Number of Words (Introduction): 607
Number of Words (Discussion): 1285
Declarations of Interest: none.

Abstract

Saccades change eye position and interrupt vision several times per second,

necessitating neural mechanisms for continuous perception of object identity, orientation,

and location. Neuroimaging studies suggest that occipital and parietal cortex play

complementary roles for transsaccadic perception of intrinsic versus extrinsic spatial

properties, e.g., dorsomedial occipital cortex (cuneus) is sensitive to changes in spatial

frequency, whereas the supramarginal gyrus (SMG) is modulated by changes in object

orientation. Based on this, we hypothesized that both structures would be recruited to

simultaneously monitor identity *and* orientation across saccades. To test this, we merged

two previous neuroimaging protocols: 21 participants viewed a 2D object and then, after

sustained fixation or a saccade, judged whether the shape *or* orientation of the re-

presented object changed. We, then, performed a bilateral region-of-interest analysis on

identified cuneus and SMG sites. As hypothesized, cuneus showed both saccade and

feature (i.e., orientation vs. shape change) modulations, and right SMG showed saccade-

feature interactions. Further, the cuneus activity time course correlated with several other

cortical saccade / visual areas, suggesting a 'functional network' for feature

discrimination. These results confirm the involvement of occipital / parietal cortex in

transsaccadic vision and support complementary roles in spatial versus identity updating.

Keywords

Saccades, transsaccadic perception, orientation, shape, posterior parietal cortex,
occipital cortex, fMRI, functional connectivity
69
70
71

Introduction

The ability to extract pertinent visual information from our surroundings depends on the brain's ability to aim, and account for, eye position [1-3]. Rapid eye movements (saccades) help gather new visual information by aligning the fovea with objects of interest, at the cost of disrupting visual stability and memory by displacing the retinal image relative to the world [4,5]. This necessitates some neural mechanism to continuously track both intrinsic object properties – cues to identity – and extrinsic spatial properties, like object location and orientation [5-8]. It is thought that during saccades, oculomotor signals are used to 'update' retinal location [9-15] and other visual features [5,7]. Recently, it has been shown that parietal and occipital cortex are involved in transsaccadic memory of spatial orientation and frequency, respectively [16-18]. However, in real world conditions, the brain must track intrinsic and extrinsic object properties simultaneously [19,20].

Various structures in the occipito-parieto-frontal stream have been implicated in the spatial aspects of transsaccadic vision, notably the parietal and frontal eye fields [9,10,21-26]. Likewise, transcranial magnetic stimulation (TMS) over various occipital, parietal, and frontal sites disrupts transsaccadic updating of object orientation [8,27-29]. More recently, we employed functional magnetic resonance imaging (fMRI) adaptation [30,31] to identify areas that are specifically involved in transsaccadic comparisons of object orientation. These experiments identified an area in the right supramarginal gyrus (SMG), anterolateral to the parietal eye fields, that showed transsaccadic orientation processing for both perception and grasp [16,17].

It has proven more difficult to localize the cortical mechanisms that update features

related to object identity [32]. Modest amounts of feature remapping have been observed

in monkey parietal eye fields [33]. A recent human study has shown that stimulus features

(specifically, spatial frequency) can be decoded from whole-brain

magnetoencephalography signals after an intervening saccade [34]. When we adapted

our transsaccadic fMRI adaption paradigm to test spatial frequency, the main source of

these modulations appeared to be superior-medial occipital cortex, i.e., cuneus [18].

Based on these results and our earlier experiments [16,17], we speculated that parietal

and occipital cortex show a relative specialization for updating extrinsic spatial attributes

of objects versus cues to intrinsic object identity, respectively.

In our previous adaption experiments [16-18], we only examined transsaccadic changes

in one feature at a time. However, as noted above, in real world conditions, the brain must

update multiple object features across saccades in order to track both object identity and

its location / orientation. This, in turn, may require the 'binding' of multiple object features

across saccades [35-37]. There is evidence that comparison between features evokes

different mechanisms than the sum of the parts [38], but it is not known how multiple

features are retained, compared, and discriminated across saccades.

Here, we investigated the cortical mechanisms for transsaccadic perception of multiple

object features, combining intrinsic (identity-related) and extrinsic (spatial) features. To

do this, we combined elements of our previous experiments [16-18]. Specifically, we used

an fMRI paradigm that required participants to remember both object shape and

orientation across a saccade and then discriminate which one of these had changed (Fig.

1). Our hypotheses were that 1) *both* SMG and cuneus would be modulated in this task,

2) SMG would show the saccade-orientation interactions that we observed previously [17]

and 3) if cuneus is involved in object identity updating, our previous spatial frequency

results should extend to the transsaccadic shape manipulation in the current task. To test

this, we performed a bilateral region-of-interest (ROI) analysis, based on the sites

obtained from our previous studies [17,18]. Our results confirmed the expected

transsaccadic feature modulations in both SMG and cuneus, and suggest that the latter

participates in a saccade-dependent network spanning several cortical lobes.

124   **-Figure 1 here-**

Results

To identify areas involved in transsaccadic discrimination of object orientation versus

shape, we employed a task (Fig. 1) where 21 participants briefly viewed an object with

one of three shapes (a rectangle, a convex 'barrel', or a concave 'hourglass') at one of

two orientations (45° clockwise or counterclockwise). They either maintained fixation to

the left or right of this central object or made a saccade to the opposite side. Then, they

re-viewed an object with either a different shape or different orientation from among the

same options, and then indicated which feature had changed. 17 participants met our

behavioural inclusion criteria for analysis of fMRI data collected during this task.

Based on previous experiments [16-18], we expected both SMG and cuneus to be

modulated in this task. To test this, we performed a hypothesis-driven region of interest

(ROI) analysis on these sites (see next sections for details), and then performed a

functional connectivity analysis to identify how saccade-related activity (if any) in these

regions correlates with activity in other cortical areas.

Region-of-interest analysis (I): Supramarginal Gyrus.

In a previous experiment, we found that an area in right SMG, slightly anterior and lateral

to the classic parietal eye field (revealed by a saccade-fixation subtraction) showed

transsaccadic repetition modulations for object orientation, i.e., different activation when

saccades were preceded and then followed by the same or different object orientations

[16]. More recently, we showed that right SMG shows an interaction between this effect

and saccades, i.e., significantly more orientation modulation during saccades compared

to fixation [17]. Here, we predicted the same result in an ROI analysis based on the

Talairach coordinates of peak right SMG activation in our previous study (Table 1) and

the mirror left hemisphere site, forming a bilateral pair.

Figure 2a, left panel, shows the location of the right SMG site relative to visual field-

specific areas (Left Visual Field – Right Visual Field Stimulation across all (fixation and

saccade) trials; yellow) and saccade-specific areas (Saccade – Fixation trials, all

conditions; magenta), superimposed on an 'inflated brain'. (Note that this rendering is

useful for showing full brain data, but causes some spatial distortions and cluster

separation.) Our right SMG site was anterior to two magenta clusters that might be part

of the parietal eye fields, but did not itself show significant independent modulations for

saccade or visual field. A whole-brain analysis of feature (i.e., shape) sensitivity at the

initial fixation point (e.g., barrel versus hourglass, clockwise versus counterclockwise

orientation) did not reveal significant clusters.

To test our specific hypothesis, we created spheres (radius = 5 mm) around our two SMG

coordinates, and then extracted β-weights from these ROIs. We then performed the same

interaction test as in our previous study on each ROI, i.e., Saccade (Same – different

orientation) – Fixation (same – different orientation) [18], except in this case 'same

orientation' corresponded to a shape change, thus: Saccade (Orientation change – Shape

change) – Fixation (Orientation change – Shape change). SMG showed a significant

hemispheric difference ($F_{1,16}$ = 54.84, p = 1.5 x 10-6, η^2 = 0.72), so we analyzed the left

and right ROIs separately. RM-ANOVAs were then used to test our specific criteria in

each of the ROIs. As predicted, right SMG passed this test (Fig. 2a, *right panel*) with a

significant saccade – feature change interaction ($F_{1,16}$ = 6.84, p = 0.019, η^2 = 0.076). It is

noteworthy, however, that the interaction was opposite to what we observed with

orientation alone. Left SMG (not shown) did not show significant modulation for saccades

($F_{1,16}$ = 0.39, p = 0.54, η^2 = 0.010), feature ($F_{1,16}$ = 0.048, p = 0.83, η^2 = 8.5 x 10^{-4}), or

saccade – feature interactions ($F_{1,16}$ = 1.46, p = 0.24, η^2 = 0.025).

173   **-Figure 2 here-**

Region-of-interest analysis (II): Cuneus.

In a previous experiment [18], we found that right and left cuneus showed significant

saccade modulations and right cuneus showed a significant transsaccadic feature

modulation for *spatial frequency* (with left cuneus showing a similar trend). Neither area

showed an interaction effect (perhaps because that occurs at some higher level). If this

area plays a more general role in transsaccadic perception of object identity, we expected

to see the same modulations for shape change in the current task.

To test this, we followed similar procedures and statistics as used for SMG. Figure 2b,

left and middle panels, show locations of the cuneus ROIs relative to saccade and field-

specific activity, respectively, superimposed on coronal slices. Both sites fall within a zone

that contains both significant saccade and field-specific modulations. The right panel of

Figure 2b shows the results of our specific hypotheses tests. In this case, cuneus did not

show significant hemispheric differences (p > 0.05), so data from the bilateral pair were

pooled. As predicted, this bilateral pair showed a significant main effect of eye movement

(| S-F | $F_{1,16}$ = 32.55, p = 3.25 x 10^{-5}, η^2 = 0.386) and feature (| O-S | $F_{1,16}$ = 11.68, p =

0.015, η^2 = 0.011). In this case, the directionality (analogous to same-different shape) was

the same as what we observed for spatial frequency in a previous study [18]. Further, as

in that previous study, we did not find a saccade-feature interaction ($F_{1,16}$ = 0.54, p = 0.47,

η^2 = 0.003).

Functional connectivity of cuneus with visual and sensorimotor regions.

In the final step of our analysis, we performed a functional connectivity analysis based

on our right cuneus ROI. To be consistent with our previous studies (and be feature

agnostic), we based this analysis on the Saccade-Fixation subtraction [17,18]. This

excluded SMG (since it did not show significant saccade modulations in this dataset).

And, to be consistent with our previous study [18], we chose the right cuneus as the

seed region. We then defined a sphere (radius = 5 mm) around this site and performed

a psychophysiological interaction (PPI) analysis. This should identify a 'network' of

cortical regions with similar saccade-related signals in this task.

This analysis revealed significant connectivity with (right) occipital, occipitotemporal,

parietal, and frontal cortex (Fig. 3, Table 2). Specifically, we observed significant

functional connectivity with early-to-intermediate visual occipital (right calcarine sulcus,

CalcS, and superior occipital gyrus, SOG, as well as bilateral lingual gyrus, LG),

sensorimotor parietal (right superior parietal lobule, SPL), oculomotor (left medial

superior frontal gyrus, SFG, and frontal eye field, FEF) and spatial coding (right inferior

frontal sulcus, IFS) frontal cortex.

209   **-Figure 3 here-**

Discussion

Our aim here was to investigate whether transsaccadic discrimination of multiple features

relies on the same or different mechanisms as those for single features. Specifically, we

tested if transsaccadic mechanisms related to intrinsic (spatial frequency) and extrinsic

(orientation) object properties would combine and generalize to a task that involves

discriminating object orientation versus shape changes. To do this, we tested for saccade

and feature sensitivity in four regions that showed both these properties in previous

transsaccadic tasks involving single features (SMG, cuneus) [17,18]. Of these regions,

right SMG and bilateral cuneus showed the predicted patterns of saccade and feature

sensitivity. Of these, cuneus showed main effects (of eye movement and feature change)

and visual field sensitivity, whereas only SMG showed interactions, perhaps suggesting

lower-level versus higher-level mechanisms. Further, saccade-related activity in cuneus

correlated with activity throughout visual, control, and saccade-related regions of cerebral

cortex. Overall, these analyses support the notion that SMG and cuneus participate in a

complex network for transsaccadic retention of object identity and orientation.

Monolithic versus Feature-specific transsaccadic mechanisms.

Although some theoretical discussions of transsaccadic feature perception have

suggested a role for occipital cortex [6,7,39,40], initial experimental studies emphasized

the parietal cortex as the key hub [8,16,17,33]. However, considering the diversity and

regionalization of visual functions [30,41,42], and the prevalence of saccades in normal

vision [43,44], the notion that the combination of saccades and vision would be handled

by one specialized mechanism seems naïve. As noted in the introduction, transsaccadic

updating of object location is found throughout visual and visuomotor cortex [5,9,34,45],

whereas transsaccadic perception of object orientation, spatial frequency, and shape

appears to involve parietal, medial occipital, and in some cases, lateral occipital cortex

[16-18]. The current results are also consistent with a more distributed system, with

feature-specific modules.

Role of SMG in transsaccadic perception.

Although the orientation of local features (such as the lines that make up curves and

shapes) can be a cue to identity [46-48], in our experiments, all lines in the stimuli have

the same orientation (Fig. 1), and the task instructions strongly reinforced the

interpretation that this was a cue to the orientation of the entire object. Given the

consistency of our results across various experiments [6,16,17] (and the current

experiment), it appears that right SMG plays a specific role in updating this property

across saccades. (See those previous papers and 'networks' below for discussions of the

hemispheric specialization of this structure). Taken together with the many studies of

spatial updating (e.g., [10,22]), this suggests a general role for parietal cortex in the

transsaccadic spatial updating of the fundamental extrinsic spatial properties of objects:

location, orientation, and the combination of these two properties [6,7].

Possible role of medial occipital cortex in transsaccadic perception.

In this and our previous study on spatial frequency [18], dorsomedial occipital cortex

appears to play a role in the postsaccadic integration of spatial frequency and object

shape. The cuneus sites described here likely fall within a region spanning visual areas

V2 and V3 [49]. These areas are ideally suited to provide the rudimentary machinery for

transsaccadic feature analysis [6,7,50,51].

First, they are well-positioned to provide rapid, bottom-up operations on their anatomic

inputs from V1 [52-54]. Second, they include the requisite machinery for feature

orientation, spatial frequency, and early shape processing [55-57]. Third, there are

indications that even these areas are able to store information in some sense [58]. Fourth,

these areas – V1, V2, and V3 – are known to show saccade modulations consistent with

remapping signals [23,29,50,51]. Finally, these areas have reciprocal connections with

both 'ventral stream' object recognition areas [58,59,60], and dorsal stream saccade and

attention areas [61-63]. This feedback, in combination with V1/V2/V3 retinotopic

topography (absent at higher levels), makes this area also ideally suited for integrating

location and identity information across saccades [6,7,39].

Thus, situated at both early and intermediate stages of object feature processing, with the

capability to monitor several object feature changes and receive feedback signals for

attention and saccades, medial occipital cortex is an excellent, additional candidate for

transsaccadic feature perception. Our experiments suggest that this area might play a

specific role in transsaccadic processing of lower-level intrinsic object features related to

identity, presumably working in concert with higher level 'ventral stream' mechanisms like

lateral occipital cortex and inferior parietal cortex [20,64,65].

Combining mechanisms for extrinsic and intrinsic feature updating.

As noted above, transsaccadic perception would be somewhat useless if we could not

update and link both object identity and location / orientation across saccades [43,66].

There is evidence that comparison between features evokes different mechanisms than

the sum of the parts [38]. Based on the common properties of spatial frequency and shape

(as intrinsic object features related to object *identity)*, we speculated that both cuneus and

SMG would be modulated by transsaccadic shape changes. In the experiment, both

cuneus and SMG showed the expected modulations, although the saccade-feature

interaction had the opposite directionality [17]. This suggests that the individual

mechanisms for transsaccadic orientation and shape updating combined, although not

necessarily in a linear fashion. What remains to be understood, is how these various

signals are combined – the classic 'binding problem' [67,68]. While cuneus and SMG

were not linked in the same functional network in the current study (likely because SMG

did not show strong enough saccade modulations), there is enough overlap between their

functional networks in different studies (see next section) to suggest that this is a possible

linking mechanism.

Transsaccadic networks: Task-specificity.

Considering the complexity of transsaccadic signals discussed above, it seems likely that

these mechanisms can be better understood at a network level. In our previous study,

where participants were required to update grasp orientation across saccades [17], they

recruited a cortical network that included each element required for the task, including a

transsaccadic orientation updater (SMG), saccade signals (FEF), and grasp motor areas

(anterior intraparietal and superior parietal cortex). Likewise, in the current study, we

identified a network that seems eminently suitable for our feature discrimination task (Fig.

3): calcarine sulcus for early feature processing [49,57,69], lingual gyrus and superior

occipital gyrus for mid-level shape recognition [70,71], posterior parietal cortex for

saccade/reach updating signals [9,10,22,72], and oculomotor superior frontal gyrus (likely

pre-supplementary motor area)/frontal eye field [27] and spatial coding in inferior frontal

sulcus [73]. The differences in these studies might thus simply be due to task and feature

details.

Finally, although we used right cuneus as a seed region here, it is noteworthy that its

activity predominantly correlated with activity in right occipital and parietal cortex, with

oculomotor activation in left hemisphere. Right cortex also predominated in our previous

regional and network analyses [16-18]. Consistent with this, transsaccadic memory is

most sensitive to right parietal stimulation and brain damage [6,45,74]. This is generally

consistent with the role of parietal cortex in spatial attention and cognition and suggests

that impaired transsaccadic perception might contribute to better-known disorders like

hemispatial neglect [75].

Conclusion

Here, we used fMRI to determine the cortical correlates of transsaccadic perception in a

task where participants had to remember and discriminate changes in object orientation

versus shape after a saccade. Our current findings support specific roles of two regions

in transsaccadic feature discrimination: SMG for orientation and medial occipital cortex

for shape. Taken together with the previous literature, these results suggest that the

cortical mechanisms for transsaccadic perception are both feature- and task-dependent.

Specifically, there seems to be relative specialization for updating of object location and

orientation in parietal cortex [9,16-18,50,51] versus transsaccadic perception of intrinsic

object features in medial occipital cortex (here and [18]). These areas appear to work in

conjunction for more complex tasks, apparently forming ad-hoc functional networks with

saccade centers (providing the updating signal) and higher order recognition and/or motor

areas, as required for the task.

Materials and Methods

Participants.

To determine the minimum number of participants required to attain adequate statistical

power, we used the effect size from right SMG, which is closest to predicted SMG here,

found in the most recent of a series of transsaccadic perception studies [17]. We

performed a power analysis (G*Power) using this effect size (0.887), with the following

parameters: 1) two-tailed t-tests for the difference of matched pair means, 2) a desired

power value of 0.90, and 3) an alpha-value of 0.05. On this basis, the suggested number

of participants was 13, though we wanted to ensure full desired power, so we recruited

21 participants for this study, of whom we analyzed data from 17 participants (see below).

The expected power was 0.914.

Twenty-one participants with normal or corrected-to-normal vision from York University,

Toronto, Canada took part in this study. One participant's data were excluded, as this

person did not complete the experiment, and three participants' data were excluded due

to poor behavioural performance (i.e., less than 80% behavioural accuracy across all

runs), which left 17 participants' data for all further analyses. Participants were

compensated monetarily for their time. All participants were right-handed (mean age: 28.7

+/- 5.2, 12 females). All participants provided written informed consent and did not have

any neurological disorders. This experiment was approved by the York University Human

Participants Review Subcommittee.

Experimental set-up and stimuli.

Participants were first required to pass MRI screening. Upon successful completion of

this step, participants were introduced to the task. Once they felt comfortable with the

requirements of the task, they were asked to assume a supine position on the MRI table.

Their head lay in a 64-channel head coil. A mount on the head coil was used to support

an infrared eye-tracker (for the right eye) to ensure appropriate fixation/eye movements,

as well as a mirror that reflected the image of the screen from inside the MRI bore. iViewX

software (SensoMotoric Instruments) was used to record eye movements for offline

verification of correct behavior (see exclusion criteria above). Participants were provided

with a button box that was held with the right hand (index and middle fingers positioned

over the first and second buttons, respectively). Button press responses were analyzed

offline.

The experiment was conducted under conditions of complete darkness. A fixation cross

was always visible (except during the presentation of the mask) to direct participants'

gaze as required by the task. The fixation cross could appear approximately 7° from either

the left or right edge of the gray screen (45.1° x 25.1°) along the horizontal meridian. The

black stimulus always appeared in the center of the screen and was either a: 1) rectangle,

2) barrel-shaped object, or 3) hourglass-shaped object (Fig. 1a). The dimensions of the

rectangle were 12° x 6°; the other two objects had the same area as the rectangle.

General paradigm/procedure.

Experiment.

To determine the cortical correlates of transsaccadic perception of object orientation

versus shape, we used an event-related fMRI design (Fig. 1). Each trial commenced with

a fixation cross, presented either to the left or right center, for 7.5 s. Then, the central

object (rectangle, barrel-shaped, or hourglass-shaped object) appeared for 3 s at +/- 45°

from vertical. This was followed by a static noise mask for 0.75 s to avoid an afterimage.

The fixation cross would then appear either at the same position as before the mask

(*Fixation* condition) or at the other location (*Saccade* condition) for 0.75 s. The same

object presented in a different orientation (*Orientation change* condition) or one of the

other two different objects presented in the same orientation (*Shape change* condition)

appeared in the center for 3 s. The object disappeared and the fixation cross was replaced

by an instruction ("I or O?") for 3 s, indicating that participants should indicate if the two

objects presented in the trial changed in shape (first button, 'I') or in orientation (second

button, 'O'). Thus, there were four main conditions: 1) Fixation, Orientation change, 2)

Fixation, Shape change, 3) Saccade, Orientation change, or 4) Saccade, Shape change.

These trial types were randomly intermingled within a run (24 trials); there were eight runs

in total. Each run began and ended with central fixation for 18 s to establish baseline

measures.

Imaging parameters.

We used a 3T Siemens Magnetom Prisma Fit magnetic resonance imaging (MRI)

scanner. Functional experimental data were acquired with an echo-planar imaging (EPI)

sequence (repetition time [TR]= 1500 ms; echo time [TE]= 30 ms; flip angle [FA]= 78

degrees; field of view [FOV]= 220 mm x 220 mm, matrix size= 110 x 110 with an in-slice

resolution of 2 mm x 2 mm; slice thickness= 2 mm, no gap). A total of 312 volumes of

functional data (72 slices) were acquired in an ascending, interleaved manner for each of

the eight runs. In addition to the functional scans, a T1-weighted anatomical reference

volume was obtained using an MPRAGE sequence (TR= 2300 ms, TE= 2.26 ms; FA= 8

degrees; FOV= 256 mm x 256 mm; matrix size= 256 x 256; voxel size= 1 x 1 x 1 mm^3).

192 slices were acquired per volume of anatomical data.

Analysis.

Behavioural data.

Eye movements and button presses were recorded throughout the experiment; the former

were monitored during the experiment and analyzed offline to ensure appropriate fixation

and saccade production. If participants made eye movements when not required to, the

data associated with those trials were excluded from further analysis. In addition, data for

trials where incorrect button presses were made were excluded from any additional

analysis. On these bases, 148 trials were removed (out of 3264 trials, representing 4.5%;

maximum four trials were removed across all participants, with a mode of four) for the 17

participants.

Functional imaging data.

Functional data from each run for each participant were preprocessed (slice time

correction: cubic spline, temporal filtering: <2 cycles/run, and 3D motion correction:

trilinear/sinc). Raw anatomical data were transformed to a Talairach template [76].

Functional data were co-registered via gradient-based affine alignment (translation,

rotation, scale affine transformation). Finally, data were smoothed using a Gaussian

kernel of 8 mm full width at half maximum. On the basis of missing predictors (more than

50% of trials with incorrect behavioral response), nine runs across all participants were

removed from the overall random effects general linear model (RFX GLM; i.e., 216 trials

out of 3116 trials; 6.9%).

Four separate GLMs were generated for each participant for every run. The first set of

GLMs contained 4 predictors: 1) Fixation, Orientation change, for an entire trial during

which participants maintained fixation throughout while only the orientation of the same

object changed; 2) Fixation, Shape change, for a trial during which participants

maintained fixation while only the shape of the object changed; 3) Saccade, Orientation

change, for a trial during which participants had to make a saccade while only the

orientation of the object changed; and 4) Saccade, Shape change, for a trial where

participants made a saccade while only the shape of the object changed. The first

presentation of the stimulus was referred to as the *Sensory/Encoding* phase and the

second presentation period was referred to as the *Change Detection* phase. These GLMs

were created in BrainVoyager QX 20.6 (Brain Innovation), where each of the predictors

was convolved with a haemodynamic response function (standard two-gamma function

model) [77]. GLMs were additionally adjusted via the addition of a last, confound predictor

("Error") to account for any errors made during trials due to unwanted eye movements or

incorrect button responses.

The second set of GLMs were structured such that the *Sensory/Encoding* phase was

modeled with two predictors to ultimately determine visual field effects: 1) "LVF" for when

the object appeared in the left visual field relative to fixation and 2) "RVF" for trials where

the object appeared in the right visual field relative to fixation. The rest of the trial was

modeled with "Fix" or "Sacc" predictors to indicate if it was a Fixate or Saccade trial,

respectively. An "Error" confound predictor was included to model trials where eye

movement or button press errors had been made.

The third set of GLMs were modified to test for the presence of orientation-sensitive bias

during the first stimulus presentation. As such, two predictors were modeled (as before)

for the *Sensory/Encoding* phase: 1) "45°" for objects that were initially presented at 45°

and 2) "135°" for objects that were initially presented at 135° from horizontal. The

remainder of the trial was simply coded according to whether it was a Fixate or Saccade

trial with "Fix" or "Sacc" predictors. A confound predictor ("Error") was also included for

eye movement or button press errors.

The last set of GLMs were modified to determine whether there was an initial shape-

related bias upon first stimulus presentation. Three predictors, one for each of the three

object shapes, were used to model the *Sensory/Encoding* phase: 1) "Rect" for when the

rectangle was presented during this phase; 2) "Barrel" for the barrel-shaped object; and

3) "Hourglass" for the hourglass-shaped object. Similar to the two previous GLMs, the

remainder of the trial was modeled according to whether it was a Fixate or Saccade trial

(i.e., "Fix" or "Sacc" predictors, respectively). Finally, the "Error" predictor was also

included for erroneous eye movements and/or button presses.

Region-of-interest (ROI) analysis.

A hypothesis-driven region-of-interest (ROI) analysis was performed based on Talairach

coordinates obtained our previous studies [17,18]. Specifically, we used the previously

identified coordinates of peak modulation for transsaccadic orientation changes (right

SMG) and spatial frequency changes (right cuneus), which we hypothesized would

generalize to transsaccadic shape detection if cuneus plays a more general role in object

identity. We then mirrored these coordinates laterally to obtain four ROIs (i.e., bilateral

SMG and cuneus; Table 1). ROIs were created around these coordinates as spheres with

5-mm radius using BrainVoyager QX v2.8 (Brain Innovation). β-weights were then

extracted and inspected for hemispheric differences with a two-tailed repeated-measures

t-test. If a significant result was obtained, the β-weights from the left and right

hemispheres were analyzed separately for each of the two criteria (see Results for

details). On this basis, we pooled data only for cuneus. Within each of the ROIs, we ran

a repeated-measures analysis-of-variance (RM-ANOVA) with two factors: Eye movement

(2 levels: Fixation, Saccade) and Object feature (3 levels: Rectangle, Barrel, Hourglass).

See Results and Table 1 for further information.

Whole-brain univariate analysis for saccade and visual field sensitivity.

We performed a whole-brain analysis to determine the presence of saccade sensitivity

within the ROIs. For these univariate analyses, we applied four independent contrasts to

test for four effects. First, we performed a whole-brain voxelwise Saccade > Fixation

contrast to identify cortical regions that were modulated by saccades in this task (Fig. 2).

Second, we applied a Left > Right Visual Field contrast using the predictors for the

Sensory/Encoding phase to find regions that are sensitive to perception of objects within

the respective visual fields (Fig. 2). To each of the resulting volumetric maps, we applied

False Discovery Rate correction to account for multiple comparisons ($q < 0.05$)

BrainVoyager QX v2.8, Brain Innovation).

Psychophysiological interaction.

Our last goal was to identify the functional network for saccade-related updating during

transsaccadic perception. In order to do this, we conducted psychophysiological

interaction (PPI) analysis [17,77-79] on data extracted from the cuneus peak site

(Talairach coordinates: 6, -79, 15). For optimal signal processing, we used a sphere of

radius = 5 mm around this site. To perform this analysis, we used three predictors: 1) a

physiological component (which is accounted for by z-normalized time-courses obtained

from the seed region for each participant for each run); 2) a psychological component

(task predictors, i.e. for the Saccade and Fixation conditions, were convolved with a

hemodynamic response function), and 3) a psychophysiological interaction component

(multiplication of seed region z-normalized time-courses with task model in a volume-

by-volume manner). For the psychological component, the *Saccade* predictors were set

to a value of '+1', whereas the Fixation predictors were set to a value of '-1'; all other

predictors were set to a value of '0' in order to determine the responses to Saccades as

compared to Fixation. (For greater detail, please see O'Reilly et al. [79] and [17]). We

created single design matrices for each run for each of the 17 participants, which were

ultimately included in an RFX GLM. Then, we used the third, psychophysiological

interaction predictor to determine the regions that constitute the functional network that

shows saccade-related modulations for feature updating. To the resulting volumetric

map, we applied FDR correction (q < 0.05).

Acknowledgements

Contributions: We thank Saihong Sun for technical support and Joy Williams and Dr.
Diana Gorbet for help with data collection.
Funding: This work was supported by a grant from the Natural Sciences and
Engineering Council (NSERC) of Canada and Canada Research Chair Program for Dr.

References

1. Javal, E., 1879. Essai sur la physiologie de la lecture. *Ann. Ocul. (Paris)*. **82**, 242-253 (1992).
2. Rayner, K. & Pollatsek, A. Eye movements and scene perception. *Can. J. Psychol.* **46**, 342-376 (1992).
3. Wade, N.J., Tatler, B.W. & Heller, D. Dodge-ing the issue: Dodge, Javal, Hering, and the measurement of saccades in eye-movement research. *Perception* **32**, 793-804 (2003).
4. Irwin, D.E. Integrating information across saccadic eye movements. *Curr. Dir. Psychol. Sci.* **5**, 94–100 (1996).
5. Melcher, D. & Colby, C.L. Trans-saccadic perception. *Trends Cogn. Sci.* **12**, 466-473 (2008).
6. Prime, S.L., Vesia, M. & Crawford, J.D. Transcranial magnetic stimulation over posterior parietal cortex disrupts transsaccadic memory of multiple objects. *J. Neurosci.* **28**, 6938–6949 (2008).
7. Prime, S.L., Vesia, M. & Crawford, J.D.. Cortical mechanisms for trans-saccadic memory and integration of multiple object features. *Philos. Trans. R. Soc. B. Biol. Sci.* **366**, 540–553 (2011).
8. Hamker, F. H., Zirnsak, M., Ziesche, A. & Lappe, M. Computational models of spatial updating in peri-saccadic perception. *Philos. Trans. R. Soc. B. Biol. Sci.* **366**, 554-571 (2011).
9. Duhamel, J.R., Colby, C.L. & Goldberg, M.E. The updating of the representation of visual space in parietal cortex neurons. *Science* **255**, 90-92 (1992).
10. Merriam, E., Genovese, C.R. & Colby, C.L. Spatial updating in parietal cortex. *Neuron* **39**, 361-373 (2003).
11. Klier, E.M. & Angelaki, D.E. Spatial updating and the maintenance of visual constancy. *Neuroscience* **156**, 801–818 (2008).
12. Mathôt, S. & Theeuwes, J. Evidence for the predictive remapping of visual attention. *Exp. Brain Res.* **200**, 117-122 (2010).
13. Burr, D. C. & Morrone, M. C. Spatiotopic coding and remapping in humans. *Philos. Trans. R. Soc. Lond. B, Biol. Sci.* **366**, 504-515 (2011).
14. Rao, H.M., Mayo, J.P. & Sommer, M.A. Circuits for presaccadic remapping. *J. Neurophysiol.* **116**, 2624-2636 (2016).
15. Bisley, J.W., Mirpour, K. & Alkan, Y. The functional roles of neural remapping in cortex. *J. Vis.* **20**, 1-15 (2020).
16. Dunkley, B.T., Baltaretu, B. & Crawford, J.D. Trans-saccadic interactions in human parietal and occipital cortex during the retention and comparison of object orientation. *Cortex* **82**, 263-276 (2016).

17. Baltaretu, B.R., Monaco, S., Velji-Ibrahim, J., Luabeya, G.N. & Crawford, J.D. Parietal cortex integrates saccade and object orientation signals to update grasp plans. *J. Neurosci.* **40**, 4525-4535 (2020).

18. Baltaretu, B.R., Dunkley, B.T., Stevens, W.D. & Crawford, J.D. Occipital cortex is modulated by transsaccadic changes in spatial frequency: an fMRI study. *Sci. Rep.* **11**, 8611 (2021).

19. Oksama, L. & Hyönä, J. Position tracking and identity tracking are separate systems: Evidence from eye movements. *Cognition* **146**, 393-409 (2016).

20. Nummenmaa, L., Oksama, L., Glerean, E. & Hyönä, J. Cortical circuit for binding object identity and location during multiple-object tracking. *Cereb. Cortex* **27**, 162-172 (2017).

21. Umeno, M.M. & Goldberg, M.E. Spatial processing in the monkey frontal eye field. I. Predictive visual responses. *J. Neurophysiol.* **78**, 1373-1383 (1997).

22. Medendorp, W.P., Goltz, H.C., Vilis, T. & Crawford, J.D. Gaze-centered updating of visual space in human parietal cortex. *J. Neurosci.* **23**, 6209-6214 (2003).

23. Merriam, E., Genovese, C.R. & Colby, C.L. Remapping in human visual cortex. *J. Neurophysiol.* **97**, 1738-1755 (2007).

24. Zirnsak, M., Steinmetz, N.A., Noudoost, B., Xu, K.Z. & Moore, T. Visual space is compressed in prefrontal cortex before eye movements. *Nature* **507**, 504-507 (2014).

25. Neupane, S., Guitton, S., & Pack, C.C. Two distinct types of remapping in primate cortical area V4. *Nat. Commun.* **7**, 1-11 (2016).

26. Hartmann, T.S., Zirnsak, M., Marquis, M., Hamker, F.H. & Moore, T. Two types of receptive field dynamics in area V4 at the time of eye movements? *Front. Syst. Neurosci.* **11**, 1-7 (2017).

27. Prime, S.L., Vesia, M. & Crawford, J.D. TMS over human frontal eye fields disrupts trans-saccadic memory of multiple objects. *Cereb. Cortex* **20**, 759-772 (2010).

28. Tanaka, L.L., Dessing, J.C., Malik, P., Prime, S.L. & Crawford, J.D. The effects of TMS over dorsolateral prefrontal cortex on trans-saccadic memory of multiple objects. *Neuropsychologia* **63**, 185-193 (2014).

29. Malik, P., Dessing, J. & Crawford, J.D. Role of early visual cortex in trans-saccadic memory of object features. *J. Vis.* **15**, 1-17 (2015).

30. Grill-Spector, K. & Malach, R. fMR-adaptation: a tool for studying the functional properties of human cortical neurons. *Acta Psychol.* **107**, 293-321 (2001).

31. Segaert, K., Weber, K., de Lange, F.P., Petersson, K.M. & Hagoort, P. The suppression of repetition enhancement: a review of fMRI studies. *Neuropsychologia* **51**, 59-66 (2013).

32. Lescroart, M.D., Kanwisher, N. & Golomb, J.D. No evidence for automatic remapping of stimulus features or location found within fMRI. *Front. Syst. Neurosci.* **10**, 53 (2016).

33. Subramanian, J. & Colby, C.L. Shape selectivity and remapping in dorsal stream visual area LIP. *J. Neurophysiol.* **111**, 613-627 (2014).

34. Fabius, J. H., Fracasso, A., Acunzo, D. J., Van der Stigchel, S. & Melcher, D. Low-level visual information is maintained across saccades, allowing for a postsaccadic handoff between visual areas. *J. Neurosci.* **40**, 9476-9486 (2020).

35. Treisman, A. Feature binding, attention and object perception. *Phil. Trans. R. Soc. Lond. B.* **353**, 1295-1306 (1998).

36. Colzato, L.S., Raffone, A. & Hommel, B. What do we learn from binding features? Evidence from multilevel feature integration. *J. Exp. Psychol. Hum. Percept. Perform.* **32** 705-716 (2006).

37. Utochkin, I.S. & Brady, T.F. Independent storage of different features of real-world objects in long-term memory. *J. Exp. Psychol. Gen.* **149**, 530-549 (2020).

38. Valyear, K.F., Culham, J.C., Sharif, N., Westwood, D. & Goodale, M.A. A double dissociation between sensitivity to changes in object identity and object orientation in the ventral and dorsal visual streams: a human fMRI study. *Neuropsychologia* **44**, 218-228 (2006).

39. Prime, S.L., Niemeier, M. & Crawford, J.D. Transsaccadic integration of visual features in a line intersection task. *Exp. Brain Res.* **169**, 532-548 (2006).

40. Hamker, F.H. & Zirnsak, M. V4 receptive field dynamics as predicted by a systems-level model of visual attention using feedback from the frontal eye field. *Neural Netw.* **19**, 1371-1382 (2006).

41. James, T.W., Culham, J., Humphrey, G.K., Milner, A.D. & Goodale, M.A. Ventral occipital lesions impair object recognition but not object-directed grasping: an fMRI study. *Brain* **126**, 2463-2475 (2003).

42. Grill-Spector, K. & Sayres, R. Object recognition: Insights from advances in fMRI Methods. *Curr. Dir. Psychol. Sci.* **17**, 73-79 (2006).

43. Land, M.F. & Hayhoe, M. In what ways do eye movements contribute to everyday activities? *Vis. Res.* **41**, 3559-3565 (2001).

44. Land, M.F. Eye movements and the control of actions in everyday life. *Prog. Retin. Eye Res.* **25**, 296-234 (2006).

45. Ten Brink, A.F., Fabius, J.H., Weaver, N.A., Nijboer, T.C.W. & Van der Stigchel, S. Trans-saccadic memory after right parietal brain damage. *Cortex* **120**, 284-297 (2019).

46. Kourtzi, Z., Tolias, A.S., Altmann, C.F., Augath, M. & Logothetis, N.K. Integration of local features into global shapes: Monkey and human fMRI studies. *Neuron* **37**, 333-346 (2003).

47. Wagemans, J., De Winter, J., Op de Beeck, H., Ploeger, A, Beckers, T., ... Identification of everyday objects on the basis of silhouette and outline versions. *Perception* **37**, 207-244 (2008).

48. Elder, J.H. Shape from contour: Computation and representation. *Ann. Rev. Vis. Sci.* **4**, 423-450 (2018).

49. McKeefry, D., Watson, J., Frackowiak, R., Fong, K. & Zeki, S. The activity in human areas V1/V2, V3, and V5 during perception of coherent and incoherent motion. *Neuroimage* **5**, 1-12 (1997).

50. Nakamura, K. & Colby, C.L. Visual, saccade-related, and cognitive activation of single neurons in monkey extrastriate V3A. *J. Neurophysiol.* **84**, 677-692 (2000).

51. Nakamura, K. & Colby, C.L. Updating of the visual representation in monkey striate and extrastriate cortex during saccades. *Proc. Natl. Acad. Sci.* **99**, 4026-5031 (2002).

52. Livingstone, M. & Hubel, D. Segregation of form, color, movement, and depth: Anatomy, physiology, and perception. *Science* **240**, 740-749 (1988).

53. Sincich, L.C. & Horton, J.C. Divided by cytochrome oxidase: a map of the
projections from V1 to V2 in macaques. *Science* **295**, 1734-1737 (2002).
54. Sincich, L.C. & Horton, J.C. The circuitry of V1 and V2: integration of color, form,
and motion. *Annu. Rev. Neurosci.* **28**, 303-326 (2005).
55. Tootell, R., Silverman, M., Hamilton, S., Switkes, E. & De Valois, R. Functional
anatomy of macaque striate cortex. V. Spatial frequency. *J. Neurosci.* **8,** 1610-
1624 (1988).
56. Gegenfurtner, K.R., Kiper, D.C. & Fenstemaker, S.B. Processing of color, form,
and motion in macaque area V2. *Vis. Neurosci.* **13**, 161-172 (1996).
57. Fang, F., Murray, S.O., Kersten, D. & He, S. Orientation-tuned fMRI adaptation in
human visual cortex. *J. Neurophysiol.* **94**, 4188-4195 (2005).
58. Tong, F. Primary visual cortex and visual awareness. *Nat. Rev. Neurosci.* **4**, 219-
229 (2003).
59. Salin, P.-A. & Bullier, J. Corticocortical connections in the visual system:
structure and function. *Physiol. Rev.* **75,** 107-154 (1995).
60. Lamme, V.A.F. & Roelfsema, P.R. The distinct modes of vision offered by
feedforward and recurrent processing. *Trends Neurosci.* **23**, 571-579 (2000).
61. Corbetta, M., Akbudak, E., Conturo, T.E., Snyder, A.Z., Ollinger, J.M., Drury,
H.A., … A common network of functional areas for attention and eye movements.
Neuron **21**, 761-773 (1998).
62. Yantis, S., Schwarzbach, J., Serences, J.T., Carlson, R.L., Steinmetz, M.A., …
Transient neural activity in human parietal cortex during spatial attention shifts.
Nat. Neurosci. **5**, 995-1002 (2002).
63. Behrmann, M., Geng, J.J. & Shomstein, S. Parietal cortex and attention. *Curr.
Opin. Neurobiol.* **14**, 212-217 (2004).
64. Kourtzi, Z. & Kanwisher, N. Cortical regions involved in perceiving object shape.
J. Neurosci. **20**, 3310-3318 (2000).
65. Denys, K., Vanduffel, W., Fize, D., Nelisson, K., Peuskens, H., … The processing
of visual shape in the cerebral cortex of human and nonhuman primates: A
functional magnetic resonance imaging study. *J. Neurosci.* **24**, 2551-2565
(2004).
66. Land, M., Mennie, N. & Rusted, J. The roles of vision and eye movements in the
control of activities of daily living. *Perception* **28**, 1311-1328 (1999).
67. Milner, P.M. A model for visual shape recognition. *Psychol. Rev.* **81**,
521 (1974).
68. Treisman, A. & Gelade, G. A feature integration theory of attention. *Cogn.
Psychol.* **12**, 97-136 (1980).
69. Cavina-Pratesi, C., Kentridge, R.W., Heywood, C.A. & Milner, A.D. Separate
channels for processing form, texture, and color: Evidence from fMRI adaptation
and visual object agnosia. *Cereb. Cortex* **20**, 2319-2332 (2010).
70. Lerner, Y., Hendler, T., Ben-Bashat, D., Harel, M. & Malach, R. A hierarchical
axis of object processing stages in the human visual cortex. *Cereb. Cortex* **11**,
287-297 (2001).
71. Ben-Shachar, M., Dougherty, R.F., Deutsch, G.K. & Wandell, B.A. Differential
sensitivity to words and shapes in ventral occipito-temporal cortex. *Cereb. Cortex*
17, 1604-1611 (2007).

72. Rossit, S., McAdam, T., Mclean, D.A., Goodale, M.A. & Culham, J.C. fMRI
reveals a lower visual field preference for hand actions in human superior
parieto-occipital cortex (SPOC) and precuneus. *Cortex* **49**, 2525-2541 (2013).
73. Hansen, K.A., Chu, C., Dickinson, A., Pye, B., Weller, J.P., … Spatial selectivity
in the temporoparietal junction, inferior frontal sulcus, and inferior parietal lobule.
J. Vis. **15**, https://doi.org/10.1167/15.13.15 (2015).
74. Sapir, A., Hayes, A., Henik, A., Danziger, S. & Rafal, R. Parietal lobe lesions
disrupt saccadic remapping of inhibition location tagging. *J. Cogn. Neurosci.* **16**,
503-509 (2004).
75. Ptak, R. & Müri, R. The parietal cortex and saccade planning: lessons from
human lesion studies. *Front. Hum. Neurosci.* **7**, 254 (2013).
76. Talairach, J. & Tournoux, P. Stereotaxic Atlas of the Human Brain (1988).
77. Friston, K. J., Buechel, C., Fink, G. R., Morris, J., Rolls, E., …
Psychophysiological and modulatory interactions in
neuroimaging. *Neuroimage* **6**, 218-229 (1997).
78. McLaren, D.G., Ries, M.L., Xu, G. & Johnson, S.C. A generalized form of
context-dependent psychophysiological interactions (gPPI): A comparison to
standard approaches. *Neuroimage* **61**, 1277–1286 (2012).
79. O'Reilly, J.X., Woolrich, M.W., Behrens, T.E.J., Smith, S.M. & Johansen-Berg, H.
Tools of the trade: Psychophysiological interactions and functional connectivity.
Soc. Cogn. Affect. Neurosci. **7**, 604–609 (2012).

Figure Legends

Figure 1. Experimental paradigm and criteria for transsaccadic feature modulation. An

example trial is presented (initial left fixation, rectangle, 45º) with the four possible main

conditions: 1) Fixation, Different Orientation, 2) Fixation, Different Shape, 3) Saccade,

Different Orientation, or 4) Saccade, Different Shape. Each of the 19.5 s trials had three

main phases: 1) *Sensory/Encoding*, during which the stimulus (one of the three possible

objects: rectangle, barrel, or hourglass) was first presented at 45º to the left or right of

vertical, while the participant was looking to the left or right of centre; 2) *Change Detection*,

wherein the same object was presented at the orthogonal orientation (*Different*

Orientation condition) or a different object at the same initial orientation (*Different Shape*

condition) while participants fixated on the initial fixation cross (*Fixation* condition) or after

they made a saccade (*Saccade* condition); and 3) *Response*, where participants made a

button press response to make a judgment about whether the shape ('I') or the orientation

('O') of the object has changed across the two stimulus presentations.

745
746
Figure 2. Region-of-interest (ROI) analysis results for tests of eye movement and

feature effects. (**a**) Voxelwise maps (n=17) from two contrasts used to determine simple

eye movement (Saccade > Fixation; "Sacc > Fix"; fuchsia) and visual field (Left Visual

Field > Right Visual Field; "LVF > RVF" and "RVF > LVF"; lime green and cornflower

blue) effects are overlaid onto inflated brain rendering of example participant's right

hemisphere (lateral view). Saccade sensitivity is shown parietally in angular gyrus (AG),

whereas visual field effects can be observed in occipital cortex (extending from inferior

through middle to superior occipital gyrus; IOG, MOG, and SOG). These regions are

posterior/medial to right SMG (orange). B-weight differences depicted for right SMG in

the bar graph on the right (error bars: SEMs) which show the significant interaction

between eye movement- and feature-related changes across saccades (SO: Saccade,

Orientation change condition; SS: Saccade, Shape change condition; FO: Fixation,

Orientation change condition; FS: Fixation, Shape change condition). (**b**) Transverse

slices (Talairach X-coordinates provided) for left and right cuneus (yellow) that show

saccade (leftmost) and visual field (middle) sensitivity. B-weights differences (error bars:

SEMs) from left (LH) and right (RH) cuneus are plotted (*left bar:* S: Saccade condition;

F: Fixation condition; *right bar*: O: Orientation change condition; S: Shape change

condition) and show significant eye movement and feature effects, as was expected

from similar results in another transsaccadic perception study [18].

766
767
Figure 3. Functional connectivity network involved in transsaccadic updating of object

features (orientation, shape). Using a Saccade > Fixation contrast, a psychophysiological

interaction (PPI) contrast revealed statistically significant (RFX GLM; FDR (q < 0.05))

functional connectivity between right cuneus and frontal (right inferior frontal sulcus, IFS,

left superior frontal gyrus, SFG, and frontal eye field, FEF), parietal (bilateral superior

parietal lobule, SPL), and occipital (right calcarine sulcus, CalcS, left superior occipital

gyrus, SOG, and bilateral lingual gyrus, LG) regions. (Note that these correlations do not

necessarily imply direct anatomical connections.) Left: Transverse slice through the

average brain of all participants (n=17) shows activation in left frontal (SFG, FEF) and

right parietal (SPL) cortex in yellow which depicts correlation with the seed region, right

Cu.

Figure 1

Figure 2

Figure 3

816 **Table 1**. Regions-of-interest, Talairach coordinates, t-statistic (T), and p-values (P) for *a*

817 *priori* selected areas tested for saccade and feature modulations.

Region	Talairach coordinates			Criterion 1		Criterion 2		Ref's
	x	y	z	T	P	T	P	
Occipital Cortex								
Cuneus	±6	-79	15	5.8	0.00028	2.9	0.0085	Baltaretu et al., 2021
Parietal Cortex								
LH Supramarginal Gyrus	-49	-40	48	-0.6	0.87	0.2	0.93	Baltaretu et al., 2020
RH Supramarginal Gyrus	49	-40	48	-0.2	0.93	0.1	0.93	Baltaretu et al., 2020

831 **Table 2.** Functional connectivity network regions and Talairach coordinates resulting

832 from psychophysiological interaction (PPI) analysis with right cuneus as the seed

833 region.

Region	Talairach coordinates						Active Voxels	References
	x	y	z	Std x	Std y	Std z		
Seed Region								
RH Cuneus	6	-79	15					
Occipital Areas								
LH Lingual Gyrus	-11	-76	-6	2.8	2.8	2.9	958	Decety et al., 1994
LH Superior Occipital Gyrus	-28	-77	14	2.6	2.5	2.1	587	Salmon et al., 1996
RH Calcarine Sulcus	4	-82	-2	2.9	2.9	2.9	1000	Miki et al. 2001
RH Lingual Gyrus	3	-86	-14	2.9	2.8	2.8	980	Decety et al., 1994
Parietal Areas								
LH Superior Parietal Lobule	-19	-77	39	2.8	1.6	2.5	437	Cavina-Pratesi et al., 2006
RH Superior Parietal Lobule	9	-68	55	2.7	2.1	1.8	379	Tamber-Rosenau et al., 2011
Frontal Areas								
LH Frontal Eye Field	-25	-12	53	1.3	1.2	2.2	141	Luna et al., 1998
LH Superior Frontal Gyrus	-4	0	52	1.8	1.4	1.6	154	Luna et al., 1998
RH Inferior Frontal Sulcus	41	3	31	2.8	2.2	2.4	602	Hansen et al., 2015