1 **Hybrid histidine kinase BinK represses *Vibrio fischeri* biofilm signaling at multiple**
2 **developmental stages**
3

4 Denise A. Ludvik[1,2], Katherine M. Bultman[1,3], and Mark J. Mandel[1,3,*]
5
6 [1] Department of Medical Microbiology and Immunology, University of Wisconsin-Madison,
7 Madison, WI USA
8
9 [2] Department of Microbiology-Immunology, Northwestern University Feinberg School of
10 Medicine, Chicago, IL USA
11
12 [3] Microbiology Doctoral Training Program, University of Wisconsin-Madison, Madison, WI USA
13
14

15 Short title: Antagonistic regulation by *V. fischeri* RscS and BinK
16
17 Keywords: biofilms, phosphorelay, two-component systems, symbiosis, *Vibrio fischeri*, *Aliivibrio*
18 *fischeri*
19
20

21 * Correspondence to:
22
Mark J. Mandel
University of Wisconsin-Madison
Department of Medical Microbiology and Immunology
1550 Linden Drive
Madison, WI 53706
Phone: (608) 261-1170
Fax: (608) 262-8418
Email: mmandel@wisc.edu

31
32
33
34

ABSTRACT

The symbiosis between the Hawaiian bobtail squid, *Euprymna scolopes*, and its exclusive light-

organ symbiont, *Vibrio fischeri*, provides a natural system in which to study host-microbe

specificity and gene regulation during the establishment of a mutually-beneficial symbiosis.

Colonization of the host relies on bacterial biofilm-like aggregation in the squid mucus field.

Symbiotic biofilm formation is controlled by a two-component signaling (TCS) system consisting

of regulators RscS-SypF-SypG, which together direct transcription of the Syp symbiotic

polysaccharide. TCS systems are broadly important for bacteria to sense environmental cues

and then direct changes in behavior. Previously, we identified hybrid histidine kinase BinK as a

strong negative regulator of *V. fischeri* biofilm regulation, and here we further explore the

function of BinK. To inhibit biofilm formation, BinK requires the predicted phosphorylation sites

in both the histidine kinase (H362) and receiver (D794) domains. Furthermore, we show that

RscS is not essential for host colonization when *binK* is deleted from strain ES114, and imaging

of aggregate size revealed no benefit to the presence of RscS in a background lacking BinK.

Strains lacking RscS still suffered in competition. Finally, we show that BinK functions to inhibit

biofilm gene expression in the light organ crypts, providing evidence for biofilm gene regulation

at later stages of host colonization. Overall, this study provides direct evidence for opposing

activities of RscS and BinK and yields novel insights into biofilm regulation during the maturation

of a beneficial symbiosis.

IMPORTANCE

Bacteria are often in a biofilm state, and transitions between planktonic and biofilm

lifestyles are important for pathogenic, beneficial, and environmental microbes. The critical

nature of biofilm formation during *Vibrio fischeri* colonization of the Hawaiian bobtail squid light

organ provides an opportunity to study development of this process *in vivo* using a combination

of genetic and imaging approaches. The current work refines the signaling circuitry of the biofilm

pathway in *V. fischeri*, provides evidence that biofilm regulatory changes occur in the host, and

identifies BinK as one of the regulators of that process. This study provides information about

how bacteria regulate biofilm gene expression in an intact animal host.

INTRODUCTION

Animals are host to microbial partners that perform essential functions, including

promoting tissue and immune development, nutrient acquisition, and defense (1, 2). In many

cases, the hosts emerge aposymbiotic (i.e., lacking their symbiont) and must then recruit and

retain the correct symbiotic microbes from the environment. On the microbial side, these

horizontally-acquired symbionts often navigate multiple distinct lifestyles, including a free-living

environmental stage and a distinct host-associated phase (3). To understand how reproducible

symbiotic colonization occurs against this backdrop of distinct microbial life stages, we can use

a model system in which a host organ is colonized by only a single bacterial partner. Here, we

focus on marine *Vibrio fischeri* and their colonization of the light organ of the Hawaiian bobtail

squid, *Euprymna scolopes* (4, 5). *V. fischeri* gains exclusive access to the squid's light organ

niche and creates luminescence that the squid manipulates as a counter-illumination

camouflage strategy (6, 7). Within hours of the aposymbiotic squid hatching, *V. fischeri* colonize

the light organ (8, 9). In addition to the binary nature of the symbiosis and the ability to rear both

partners separately, the amenability of *V. fischeri* to sophisticated genetic manipulation and the

power to image at the direct site of infection provides a powerful tool box to study how genes

and their products influence bacterial colonization.

Biofilm formation by the symbiotic bacteria is fundamental to the colonization of the

squid host (3, 10–15). During the establishment of the symbiosis, squid recruit *V. fischeri* by

pumping seawater through the mantle cavity and over the light organ. Biofilm formation is

required for *V. fischeri* to aggregate in the host mucus, and *V. fischeri* mutants unable to

synthesize biofilm are unable to colonize the squid (10, 11). The polysaccharide component of

the biofilm is the symbiosis polysaccharide (Syp), whose production is encoded by the 18-gene

syp locus on the *V. fischeri* second chromosome (11). Expression of this locus is controlled by a

two-component phosphorelay (**Figure 1A**). In strain ES114, the current model posits that hybrid

sensor kinase RscS autophosphorylates at residue H412 in its dimerization and histidine

phosphotransferase (DHp) domain and then relays the phosphoryl group to D709 in its receiver

domain (REC) (16). The phosphoryl group is then relayed to residue H705 in the histidine

phosphotransfer (HPt) domain of a distinct hybrid sensor kinase, SypF (17, 18). SypF then

phosphorylates D53 in the REC domain of response regulator SypG (17, 19). Phospho-SypG

acts as a σ^{54}-dependent enhancer binding protein to promote transcription of the *syp* locus (11,

20). SypF has additional effects, including regulating the activity of serine kinase/phosphatase

SypE to inactivate/activate its target SypA, which influences Syp biofilm formation downstream

of *syp* transcription (21).

Previously, we identified the biofilm regulator, BinK, as a strong inhibitor of symbiotic

biofilm formation (22). BinK was identified in an insertion sequencing screen where a mutant

library of *V. fischeri* was analyzed before and after colonization of squid hatchlings to find genes

that influenced colonization (23). Mutants of *binK* were overrepresented in the output pool

compared to the input, suggesting that deletion of *binK* confers an advantage during the

colonization process. Upon further study, it was determined that deletion of *binK* confers a

competitive advantage against the wild-type strain during squid colonization, and the mutant

forms larger aggregates on the surface of the squid light organ (22). Overexpressing BinK leads

to a substantial reduction in symbiotic biofilm formation and prohibits bacteria from colonizing

the host. BinK, which is conserved across *V. fischeri* strains, was also identified to be an

important colonization regulator in a squid colonization experimental evolution study (14).

Natural *V. fischeri* fish and seawater isolates that were experimentally evolved to colonize squid

were able to do so as a result of spontaneous mutations in *binK* (14). Strain ES114 does not

form robust biofilms outside of the squid. However, we can use culture-based assays with an

RscS-overexpressing allele (termed, *rscS**) that approximates *in vivo* biofilm phenotypes to

study BinK function, and with those approaches we demonstrated that deletion of *binK* can lead

to wrinkled colony formation, higher transcription of the *syp* locus, and higher production of the

Syp exopolysaccharide (22). Overexpressing *sypG* is epistatic to inhibitory signaling from

overexpressing *binK* (22). The Δ*binK* strain is significantly derepressed for biofilm formation and

in that background, calcium stimulates colony biofilm formation without the need for induction by

rscS* alleles (24). The calcium induction system led to the discovery of biofilm regulator HahK,

which influences biofilm development in response to host nitric oxide (24, 25).

Given the prominence of BinK as a strong negative regulator of biofilm across various *V.*

fischeri natural isolates, here we pursued multiple questions regarding its function during

symbiosis. First, while BinK has the predicted structure of a hybrid histidine kinase, we asked

whether it requires its putative phosphorylation sites for function. Second, given the strong

phenotypes of BinK, we asked whether canonical squid isolate ES114 would be capable of

colonizing squid in the absence of the positive regulator RscS if the negative regulator BinK was

also removed. We found that RscS was dispensable for colonization in the absence of BinK,

and this result enabled us to investigate the relative role of each protein during symbiotic

colonization. Third, using direct imaging of a fluorescent *syp* transcriptional reporter, we asked

whether symbiosis polysaccharide gene expression is regulated in the host by comparing

regulation at two distinct time points (and therefore, distinct sites with the host) and by

comparing both wild type and various mutant strains. Overall, this work provides an intriguing

look into how biofilm signaling is regulated when a symbiotic microbe encounters its animal

host.

RESULTS

BinK requires its conserved two-component histidine and aspartate for function

We identified BinK as an orphan hybrid histidine kinase that inhibits biofilm formation

and is a negative regulator of squid colonization (22). Protein domain prediction indicated that

BinK has the conserved dimerization and histidine phosphotransferase (DHp) domain and

catalytic (CA) domain typical of a two-component sensor kinase, and it also contains an

additional receiver (REC) domain making it a hybrid sensor kinase (**Figure 1B**) (26–29). His362

in the DHp domain and Asp794 in the REC domain are the predicted sites for phosphorylation in

BinK (**Figure 1C**) (22). Phosphotransfer through such sites is typically required for signaling by

sensor kinases, but there are examples where this is not the case (17, 30, 31). Therefore, we

first asked whether His362 and Asp794 are necessary for BinK function.

To assess the function of individual alleles, we conducted colony biofilm assays. We

started with a Δ*binK* strain and then introduced the *binK* gene (including 300 bp of upstream

and downstream sequence) into the neutral chromosomal *att*Tn*7* site **(Figure 2A)**. This

approach enabled us to test the wild-type and mutant alleles in comparable isogenic

backgrounds. The strain background also has an allele to induce biofilm formation under culture

conditions through the overexpression of RscS. This allele, termed *rscS** is carried on the

chromosome at the native *rscS* locus (**Figure 2A**) (32). In this background, deletion of *binK*

results in a wrinkled colony morphology when grown at 28 $^{\circ}$C, while a strain with a functional

binK has a smooth colony morphology (22). To test whether the His362 and/or Asp794 are

required for BinK function, we constructed H362Q and D794A mutations, which have been

shown to mimic the unphosphorylated state when similarly introduced into homologous domains

(16, 33). In an otherwise Δ*binK* background, mutation in either individual residue or in both

residues in the same protein resulted in a non-functional BinK (**Figure 2B**). We next asked

whether a predicted phosphomimetic allele of the REC domain D794E is functional. This allele

was constructed and unable to complement the lack of BinK in the *rscS** biofilm induction model

(**Figure 2B**). Using western blot analysis with a polyclonal antibody raised against a BinK

cytoplasmic epitope, we demonstrated that the mutant proteins are expressed to levels

comparable to that of the wild-type BinK (**Figure 2C**). Together, these results provide genetic

evidence that phosphorylation of BinK residues is required for its function. Given that we did not

observe complementation with either the D794A that is predicted to be non-phosphorylatable or

with the putative phosphomimetic D794E allele, we expect that phosphoryl groups at this

residue are transferred to (or from) a downstream signaling partner, and that this signaling is

required for BinK to inhibit biofilm formation.

BinK merodiploid analysis reveals a dominant negative phenotype for the D794A allele

Our data above demonstrated that the H362Q and D794A alleles were individually

nonfunctional. Given that histidine kinases typically operate as homodimers, we next inquired

whether the BinK(H362Q) or BinK(D794A) had any effect when expressed in a cell that also

expressed wild-type BinK protein. We continued to express the test alleles from the *att*Tn*7* site,

but now did so in a strain expressing wild-type *binK* from the native locus. In the wrinkled colony

assay, the nonfunctional H362Q allele was recessive to the wild-type allele as expected, but the

D794A allele exhibited a dominant negative phenotype, displaying a lack of biofilm inhibition

even in the presence of the wild-type allele (**Figure 2B**). A similar dominance was observed for

the D794A allele even if the H362Q mutation was introduced on the same polypeptide,

expressed from the *att*Tn*7* site (**Figure 2B**). While either the H362Q or the D794A mutation

disable BinK's biofilm inhibitory function, the latter allele additionally interferes with the ability of

a wild-type copy of BinK to inhibit biofilm formation.

Our data above demonstrated that the putative phosphomimetic D794E allele of BinK

was similarly nonfunctional as the D794A allele when expressed as the only BinK allele in the

cell. Merodiploid analysis revealed, however, that the D794E allele did not interfere with BinK

activity *in trans* (**Figure 2B**). This was the case whether it was the only mutation in *binK* or in the

context of a double *binK*(H362Q, D794E) allele (**Figure 2B**). Therefore, inactivation of the REC

domain does not correlate directly with the dominant interfering phenotype. However, it may be

the case that an unphosphorylated D794 (and the D794A allele that cannot be phosphorylated)

leads to inhibition of BinK activity, whereas phosphorylated D794 (mimicked by the D794E

allele) does not.

Sensor kinases can perform both phosphatase and kinase activities (34). To ask

whether BinK kinase or phosphatase activity is used for biofilm inhibition, we created point

mutations in the DHp domain that are predicted to differentially affect kinase versus

phosphatase activities of the protein. BinK contains a conserved ExxT motif immediately after

the conserved His362 residue (**Figure 1C**). This region of the H-box (i.e., the conserved

phosphoryl group-binding His and surrounding region) is important for coordinating the

phosphotransfer reactions in two-component proteins (35). In BinK we constructed E363A,

T366Q, and T366A mutations that are predicted to eliminate kinase activity, reduce

phosphatase activity, and eliminate phosphatase activity (with possible effects on autokinase

activity), respectively (35–41). Colony biofilm assays revealed nonfunctional BinK in each case

(**Figure S1**). This result further supports a role for phosphotransfer through BinK in its functional

role.

Single-copy phosphomimetic SypG is epistatic to BinK overexpression

As an orphan histidine kinase, BinK has no known paired response regulator. In a

previous study, we demonstrated that a separate RscS-dependent pathway that relies on

signaling through SypE was insensitive to BinK activity (22). Therefore, we can proceed to study

BinK signaling architecture in a Δ*sypE sypF2* mutant background to isolate the core biofilm

phosphorelay. Note that in this strain background, wrinkled colony formation is observed upon

overexpression of SypG (without the requirement for *rscS**) (18, 42). In our previous study, we

demonstrated that overexpression of SypG led the cell to be insensitive to overexpression of

BinK in a wrinkled colony assay (22). This result suggested that SypG was epistatic to BinK. To

further probe the genetic relationship between BinK and SypG, we constructed a

phosphomimetic SypG allele at the native *sypG* locus. Hussa *et al.* demonstrated that a plasmid

containing the SypG(D53E) allele conferred increased *syp* transcription and wrinkled colony

formation in a Δ*sypE sypF2* strain (18). We introduced the same amino acid change into

chromosomal *sypG* within the Δ*sypE sypF2* genetic background. We proceeded to determine

that overexpression of BinK does not reduce the colony biofilm produced in the Δ*sypE sypF2*

sypG(D53E) background (**Figure S2**). Therefore, this experiment supports and extends our

previous work and provides strong evidence that BinK acts upstream of SypG in the control of

syp transcription and biofilm development. We proceeded then to examine the relative impacts

of RscS and BinK on colonization phenotypes *in vivo*.

RscS is not required for aggregation or squid colonization in a strain lacking BinK

In strain ES114, the positive regulator RscS and the negative regulator BinK both exert

strong impacts on colonization and both act through the response regulator SypG. We therefore

considered models in which RscS and BinK exert opposing influence on *syp* gene transcription

during squid colonization. Deletion of *rscS* alone leads to severe *in vivo* biofilm and colonization

defects, whereas deletion of *binK* alone leads to enhanced biofilm production and improved

colonization in a competitive assay (10, 22, 43). Given these opposing phenotypes, we asked

whether removal of both regulators--RscS and BinK--would enable the bacteria to colonize the

host. An examination of experiments in diverse *V. fischeri* strains provides some insight into this

question. Strains can improve their ability to colonize squid in the laboratory by mutation of *binK*,

and this includes strain MJ11 that naturally lacks RscS (14, 22). The same MJ11 strain can

colonize squid robustly if RscS from squid symbiont ES114 is introduced (13). Together, these

results support our model that RscS and BinK could exhibit opposing activities, but we remained

uncertain as to what we would observe in native squid symbiont ES114. If RscS is mainly

required to counteract the negative regulation of BinK, then elimination of both regulators should

allow the bacteria to colonize the squid host. However, if RscS is required to transduce a

specific signal from the host, then we predict that elimination of both regulators would not allow

for colonization.

To test these models, we conducted single-strain colonization assays. As has been

shown previously, a Δ*rscS* mutant exhibits a significant colonization defect (**Figure 3**) (43). The

Δ*binK* mutant is known to exhibit a competitive advantage over wild-type (22), yet in single-

strain colonization displays similar bacterial yields and luminescence (**Figure 3**). The Δ*binK*

Δ*rscS* double mutant strain was able to colonize up to levels indistinguishable from the wild-type

strain (**Figure 3**). This result therefore supports the model that in ES114 BinK and RscS

antagonize each others' activity, and that in the absence of the negative regulator BinK, the

positive regulator RscS is no longer required for single-strain colonization.

The above result was surprising in that we identified a condition in which RscS,

discovered twenty years ago as a strong colonization factor (43), was no longer required for

squid colonization in strain ES114. This prompted us to ask whether the key symbiotic behavior

regulated by RscS--*in vivo* aggregate formation--occurs in the Δ*binK* Δ*rscS* background. For this

experiment, we introduced a plasmid that constitutively expresses GFP into the colonizing

strains from Figure 3, and we asked whether these strains form biofilm aggregates in the squid

mucus field. Direct visualization of the bacterial cells revealed the presence of biofilm

aggregates in the host for the Δ*binK* Δ*rscS* cells (**Figure 4A**). Notably, the size of these

aggregates were comparable to those formed by Δ*binK* single mutant cells, which were larger

than the aggregates formed by wild-type *V. fischeri* (**Figure 4A,B**). This result therefore reveals

that RscS is not required for aggregate formation in a background lacking BinK, and RscS does

not contribute to aggregate size in this background.

The above results prompted us to ask whether RscS performs any detectable function in

a strain lacking BinK. We employed a sensitive competition assay to ask whether strains lacking

RscS exhibit a defect upon co-inoculation. In a competitive colonization assay in which the

Δ*binK* Δ*rscS* strain was co-inoculated with a LacZ-expressing Δ*binK* single mutant, the strain

lacking RscS exhibited an approximately 100-fold defect in the competition (**Figure 5**).

In the absence of BinK, *syp* transcription *in vitro* does not require RscS

271   Recent work demonstrated that in a strain lacking BinK, RscS is not required for colony

biofilm formation when the symbiotic biofilm is induced with an additional 10 mM calcium in the

medium (LBS-Ca) (24). We therefore examined a *sypA'-gfp*[+] transcriptional reporter for cells

grown on LBS-Ca agar, in strains that lack BinK, RscS, or both regulators. We determined that

GFP activity from the reporter is induced in the Δ*binK* background, and the presence or

absence of *rscS* in this background did not impact *syp* expression (**Figure 6A,C**). Similarly, the

wrinkled colony biofilm phenotype is observed in both the Δ*binK* and Δ*binK* Δ*rscS* strains. Next,

we examined the same reporter when the cells were grown on rich medium without the

additional calcium (LBS). On LBS agar, the overall induction is lower than on LBS-Ca and

strains do not wrinkle. However, we still detect induction of the reporter in the Δ*binK* strain

compared to the WT parent. The induction is unaffected by the absence of *rscS* from this

background (**Figure 6B,C**). In summary, we can readily detect *syp* gene transcription in the

Δ*binK* background when grown on different solid media, and this induction is independent of

RscS.

BinK represses syp transcription in the squid crypts

287   We next examined expression of the *sypA'-gfp*[+] transcriptional reporter in *V. fischeri*

cells that had aggregated in the host mucus (3-4 hpi). The results in **Figure 7A,C** reveal

indistinguishable overall levels of GFP activity in WT, Δ*binK,* and Δ*binK* Δ*rscS* cells in the

aggregates of each animal examined. We note some limitations to these data. Given that cells

spend a short period of time in the aggregate stage--on the order of 1-2 h--it is unclear whether

the reporter is revealing the steady state transcription levels from the aggregate or whether this

information integrates time prior to the aggregation stage (growth in liquid media and in

seawater). Nonetheless, we present these data for two reasons. First, any physiological

response that uses transcription to regulate symbiosis would be subject to similar constraints.

Second, the similarity of the data points provides a useful control for the data in the crypts that

will be described below. Even with these caveats, we can conclude that the absence of RscS

does not diminish the median level of *sypA* transcription in the Δ*binK* background. We note that

there was more heterogeneous *sypA'-gfp*[+] activity across the aggregate in wild type compared

to samples that lacked BinK (**Figure 7A**).

We proceeded to conduct a similar analysis of the transcriptional reporter in the light

organ crypts (48 hpi). At this point, we observed a notable difference between the Δ*binK* strain

and wild type, with substantially elevated *sypA* transcription in the cells lacking BinK (**Figure**

7B,D). The absence of RscS did not affect the *sypA* reporter. From this imaging, we conclude

that a normal function of BinK is to repress *syp* transcription in the crypts.

DISCUSSION

By studying the *V. fischeri*-squid symbiosis model, we have refined our understanding of

how bacterial biofilm signaling is regulated at the initiation of bacterial colonization. This study

provides evidence that BinK acts as a hybrid histidine kinase, defines novel biofilm regulation in

the host, describes a role for BinK in that regulation, and reveals that the key ES114

colonization factor RscS is dispensable in the absence of BinK. These major conclusions are

discussed in detail below.

BinK acts as a hybrid histidine kinase. In our previous study, we identified BinK as a

putative hybrid histidine kinase based on its predicted domain structure containing CA, DHp,

and REC domains (**Figure 1**) (22). Furthermore, experimental evolution studies to improve

squid colonization of other *V. fischeri* strains revealed spontaneous *binK* point mutations in its

CA and HAMP domains that approximated the phenotypes of Δ*binK* strains, suggesting key

roles for these domains in BinK function (14). In this work, we used a combination of targeted

mutants with a newly-developed anti-BinK peptide antibody to demonstrate that mutagenesis of

predicted phosphorylation sites creates BinK proteins that are nonfunctional.

In most cases, histidine kinases dimerize, and our results provide genetic support that

this occurs in the case of BinK. In particular, our finding that a BinK protein containing a REC

domain that is locked as non-phosphorylated with the D794A mutation is not only nonfunctional,

but is dominant negative as it interferes with signaling of a wild-type *binK* allele expressed in the

same cell. There are published cases in which one monomer's REC domain interacts with the

same monomer's DHp domain (*in cis*), and other cases in which it interacts with the partner

monomer's DHp domain (*in trans*) (44, 45). In both cases, however, dynamic movement of both

monomers is required across the dimer interface (35, 46). It seems likely that the BinK D794A

allele disrupts some aspect of this dynamic process that is still intact in heterodimers containing

the phosphomimetic D794E allele. Therefore, BinK REC domain phosphorylation may both

contribute to phosphoryl group flow and regulate the ability of BinK to interact in productive

dimers.

Our data support a model in which phosphoryl groups flow from/to a downstream

signaling partner through the BinK REC domain. The lack of complementation of a Δ*binK* allele

with either the *binK*(D794A) or *binK*(D794E) alleles provide support for this model, in spite of

differences in the merodiploid analysis described above. Additionally, the double *binK*(H362Q,

D794E) allele is nonfunctional but not disruptive in the presence of a wild-type allele, suggesting

that phosphoryl group flow can proceed *in cis* through a single BinK monomer even when

present in mixed dimers with the double mutant. We note that there are examples in the

literature where a REC domain regulates activity of the DHp domain and does not directly

transfer phosphoryl groups (30, 31, 47). While this is possible for BinK, it seems unlikely given

the genetic results here. Future biochemical analysis of BinK phosphotransfer will be necessary

to further advance these studies. In particular, such analysis may enable asking whether BinK

acts as a kinase or phosphatase when it inhibits biofilm formation. Without knowledge of the

direct downstream partner of BinK, this question is difficult to access at present.

347   Considering that a phosphomimetic SypG is epistatic to BinK, it is likely that BinK is

acting to control the phosphorylation of SypG. Our results indicate that the BinK His and Asp

residues are necessary for activity. Thus, we predict BinK acts indirectly on SypG through the

HPt domain in another regulator. Among known players, the HPt domain that has been shown

to mediate phosphorylation of SypG is that of SypF (17, 24), making SypF a likely candidate. It

is also possible that the relevant partner has yet to be identified. Genetic approaches to identify

and characterize relevant partners will provide insight into the pathway downstream of BinK.

354   **RscS is dispensable for colonization in strain ES114 lacking BinK**. We found that in

strain ES114, *rscS* is not required for squid colonization in a Δ*binK* background. Strains lacking

both regulators colonize to a level comparable to that of wild type, though we found through

competitive colonization analysis that there still remains a role for RscS in this background

(Figures 3, 5). There are three major phylogenetic groups of *V. fischeri* (15). Relevant for this

work, the ancestral Group C strains include Mediteranean squid symbionts such as strain SR5.

These strains encode functional BinK, but do not encode RscS, and it is unclear how they can

colonize squid without the biofilm-promoting activity from RscS (15). Group C strains also

include fish symbionts such as MJ11, which cannot colonize squid unless they gain RscS or

lose BinK (13, 14). Derived from this group is Group B, which includes strain ES114, which is

the focus of the present study. Squid symbionts in Group B typically encode both RscS and

BinK, and mutation of RscS leads to an inability to colonize the squid (13, 15). In this study, it

was found that when *binK* is deleted, strain ES114 no longer requires *rscS* for colonization

(**Figure 3**). This result mirrors a previous finding in Group C, which found that strains lacking

RscS (due to evolution) and also lacking BinK (due to directed mutation) could colonize squid

(14). Given the diversity of biofilm regulation across *V. fischeri*, this result was not expected and

highlights conserved aspects of regulation that are shared across much of the species (15). It

therefore seems that one of the main functions of RscS is to antagonize BinK's negative

regulation of biofilm: without the negative regulator, the positive regulator is no longer absolutely

required for host colonization. From the evolutionary tree, we know that *binK*--which is found

throughout the species--predates the horizontal gene transfer event that enabled acquisition of

rscS, which is only present in a derived group of *V. fischeri* (13, 15). Therefore, our work raises

the question of whether there are other factors that antagonize BinK activity in strains that

colonize squid independent of RscS (e.g., the Group C Mediterranean squid symbionts). It

seems likely that such activity would be sufficient to enable colonization, given that mutations in

binK facilitate colonization by strains that are otherwise unable to colonize well (14).

BinK is a key regulator across the symbiotic life cycle. In a previous study, we used

a fluorescent biofilm gene promoter fusion to examine gene expression in liquid medium (22). In

this work, we expanded on that approach to examine expression of *sypA'-gfp*[+] on solid medium

and *in vivo* during colonization. On solid agar, we observed high levels of reporter expression

under conditions known to induce Syp biofilm formation (LBS-Calcium (24)). We observed a

less dramatic yet significant induction on medium where the Syp biofilm is not visibly apparent

(LBS), revealing the sensitivity of this reporter. Our results provide evidence for expression of

the *syp* biofilm in the aggregates, when biofilm formation is known to be required for

colonization. We also demonstrate that in the absence of BinK, there is expression of *syp*

reporter in the crypts, which supports a role for BinK in repressing biofilm gene expression at

this later stage in the wild type strain. Building on our observation above that RscS is

dispensable for colonization in strains lacking BinK, we observed similar levels of *syp* reporter

expression in Δ*binK* and Δ*binK* Δ*rscS* strains. Our discovery that BinK functions to repress *syp*

expression in the crypts hints that BinK regulation may serve a role in the daily expulsion of

bacteria from squid at dawn. Little is known about how this process is regulated, yet it occurs

daily in the mature symbiosis for the duration of the host's lifetime. If this proves correct, it may

help to explain why *binK* genes are widely conserved in *V. fischeri* despite the mutant having a

competitive advantage (14, 15, 22). It seems likely that while the absence of BinK is beneficial

to enter the squid host, the absence of the regulator (and subsequent inappropriate biofilm

formation at later stages) may be detrimental to the daily homeostasis that is maintained long-

term in the squid. We also examined *sypA'-gfp⁺* reporter activity in biofilm aggregates. The

average expression level within aggregates was similar for wild type or Δ*binK* mutant cells, but

we observed greater heterogeneity in the expression in wild type. This suggests that in the

presence of BinK, there is more variability in biofilm expression, and this is worthy of further

study.

405   This work provides an exciting view into how biofilm gene expression is regulated *in*

vivo. We know that within a few hours, planktonic bacteria transition to a biofilm state in the host

mucus (3, 48). Despite a number of biofilm regulators being identified, how this process is

controlled at the host interface is not well understood. Our results provide evidence that this

regulation is dynamic over the course of colonization, as evidenced by the BinK-dependent

repression that occurs specifically in the crypts at 48 hpi but is not evident in the aggregates at

3-4 hpi. We propose that the interaction of BinK with host-derived compounds may lead to

downregulation of biofilm genes as bacteria transition from the biofilm aggregates during

initiation to cells in the crypts during the persistence stage. Finally, we observed similar sizes of

in vivo biofilm aggregates in the host in Δ*binK* and Δ*binK* Δ*rscS* strains, arguing that there is no

requirement for stimulation of biofilm through RscS to initiate a productive symbiosis. We

therefore posit that a key regulatory mechanism to control the planktonic-to-biofilm transition is

host inhibition of BinK.

418   In summary, this work provides novel insight into the function of hybrid histidine kinase

419   BinK, its relationship to RscS is regulating symbiotic biofilm formation, and the temporal control

420   of symbiotic biofilm gene expression. Future work will continue to examine the signaling

421   architecture downstream of BinK and host-derived molecules that may regulate BinK activity.

MATERIALS AND METHODS

Bacterial Strains, Plasmids, and Media

V. fischeri and *Escherichia coli* strains used in this study are listed in **Table 1**. Plasmids used in

this study are listed in **Table 2**. *V. fischeri* strains were grown at 25^{o}C or 28^{o}C in Luria-Bertani

salt (LBS) medium (25 g Difco LB broth [BD], 10 g NaCl, and 50 mL 1 M Tris buffer, pH 7.5, per

liter). *E. coli* strains, used for cloning and conjugation, were grown shaking at 37^{o}C in Luria-

Bertani (LB) medium (25 g Difco LB broth [BD] per liter). Growth media were solidified with 1.5%

agar (15 g Bacto agar (BD) per liter) as needed. When necessary, antibiotics were added to the

media at the following concentrations: erythromycin, 5 μg/ml for *V. fischeri*; kanamycin, 100

μg/ml for *V. fischeri* and 50 μg/ml for *E. coli*; and chloramphenicol, 5 μg/ml for *V. fischeri* and 25

μg/ml for *E. coli*. The *E. coli* strain π3813 containing pKV496 is a thymidine auxotroph and was

grown in LB with 50 μg/ml kanamycin supplemented with 0.3 mM thymidine (49, 50).

DNA synthesis and sequencing

Each of the primers listed in **Table 3** was synthesized by Integrated DNA Technologies

(Coralville, IA). Full inserts from all cloned constructs were verified by Sanger DNA sequencing

at Northwestern University Feinberg School of Medicine Center for Genetic Medicine,

Functional Biosciences via UW-Madison, or the UW-Madison Biotechnology Center. Sequence

data were analyzed with SeqMan Pro (DNAStar software) and Benchling. For cloning and

sequencing PCR reactions, we used Q5 High-Fidelity DNA polymerase (NEB). For diagnostic

PCR, we used GoTaq polymerase (Promega).

Construction of *att*Tn*7*::*binK* mutant alleles

The previously generated pTn*7*-*binK* plasmid, which uses a mini-Tn*7* delivery vector backbone

(pEVS107), was purified and used as a template. Point mutations to the *binK* sequence on the

plasmid were designed using the NEBaseChanger tool and constructed with the Q5 site-

directed mutagenesis kit (New England BioLabs, Inc.). The constructed plasmid was

transformed into either electrocompetent or chemically competent DH5α λpir *E. coli*. The entire

binK gene on the plasmid construct was sequenced (using pEVS107 F and R primers and *binK*

sequencing primers). BinK alleles generated in this manner were then introduced into *V. fischeri*

by tetraparental mating by mixing the pEVS104-containing helper, pUX-BF13-containing

transposase, pEVS107 mini-Tn*7* vector containing donor, and the *V. fischeri* recipient (51). PCR

verification by amplifying around the *att*Tn*7* site with primers Tn7 site F and Tn7 site R

confirmed transposon insertion at the *att*Tn*7* site.

Wrinkled Colony Assays

Cultures were grown overnight and 8 μL was spotted onto LBS plates or LBS-Calcium (10 mM

CaCl$_2$). Plates were incubated at 25^{o}C or 28^{o}C for 48 hours and imaged using Leica M60

stereomicroscope with Leica Firecam software. For assays done with the pM1422 reporter,

plates were also imaged on a Zeiss Axio Zoom.v16 large-field fluorescent stereo microscope

and analyzed with Zen Blue software.

BinK peptide antibody creation and purification

ProSci (Poway, CA) analyzed the sequence of BinK and chose the peptide

SEYGEMIDLPHKRKDNLDIIK as an epitope for polyclonal antibody production. This peptide

was synthesized by ProSci and used to inoculate rabbits. Serum was analyzed at specific

checkpoints to assess antibody production. Final bleed serum was then purified in our

laboratory using a Proteus Protein A Mini Purification Starter Kit and yielded approximately 1

mg/mL of the antibody. Purified antibody was diluted 1:1 in 50% glycerol, aliquoted into small

volumes, and stored at -20^{o}C. Single aliquots were thawed and used for each blot.

SDS-PAGE and Western Blots

One milliliter of overnight culture was pelleted, washed, and lysed in 1% SDS. The volume of

SDS used to lyse the sample was adjusted based on the OD_{600} of the overnight culture to

standardize the concentration of total protein in the samples. The solution was then pelleted to

remove cell debris and the supernatant was mixed 1:1 with 2x Laemli sample buffer (Bio-Rad)

and beta-mercaptoethanol. Samples were heated at $95^{o}C$ for 15 minutes and loaded onto a

10% Bio-Rad Mini-PROTEAN TGX Precast Gel. The gel was then transferred to a PVDF

membrane and blocked overnight in 5% non-fat milk. The purified anti-BinK-peptide antibody

was used as the primary antibody in a 1:100 dilution in 0.5% nonfat milk in 1X TBS-Tween20.

The secondary antibody was a 1:5,000 dilution of the Pierce goat anti-rabbit IgG (H+L) HRP-

conjugate (Lot UK293475). Washes were done in 1X TBS-Tween20. Blots were developed

using the ThermoScientific SuperSignal West Dura Extended Duration Substrate and analyzed

using a Licor Odyssey Fc machine.

Construction of *sypG*(D53E)

Site-direction mutation of *sypG* was done using an allelic exchange approach modified from the

laboratory's gene deletion protocol (https://doi.org/10.5281/zenodo.1470836). In brief,

approximately 1.6-kb upstream sequence and 1.6-kb downstream sequence of the desired point

mutation site in *sypG* was amplified from ES114 genomic DNA using primers designed using

the NEB site-directed mutagenesis approach. These two fragments were then cloned into

pEVS79 (which had been linearized by primers MRH001 and MRH002) using isothermal

assembly (NEBuilder HiFi DNA assembly cloning kit) with the primer combinations listed in

Table 3. The reaction was transformed into *E. coli* with selection for transformants on LB-

chloramphenicol. PCR around the insertion using primers M13 For (-41) and M13 Rev (-48) was

used to confirm plasmid candidates. The resulting candidates were then confirmed by

sequencing and conjugated into the *V. fischeri* recipient (KV3299/MJM2536) by triparental

mating with helper plasmid pEVS104, selecting for the chloramphenicol resistance of the

plasmid backbone. Single recombinants in *V. fischeri* were screened for maintaining

chloramphenicol resistance. To obtain double recombinants, single recombinants were then

grown without antibiotics and patched onto LBS and LBS-Cam to find isolates that lost the

antibiotic resistance cassette. These candidates were then verified with PCR and sequencing to

confirm loss of the *cam* cassette and mutation of *sypG* to *sypG*(D53E). Strain MJM2536

sypG(D53E) was saved as MJM3236.

Construction of Δ*rscS* and Δ*binK* Δ*rscS* strains

Deletion of *rscS* was performed following the barcode-tagged gene deletion protocol from

Burgos et al. (52). In brief, the US homology arm was amplified using primers KMB_082 and

KMB_083 and the DS homology arm was amplified using primers KMB_086 and KMB_087.

Homology arms were fused to either side of a third fragment containing an *erm* cassette using

SOE-PCR. Mutagenic DNA was purified using the Qiagen PCR purification kit and transformed

into ES114 via transformation using pLostfoX-Kan (MJM1538) (53, 54). Mutant candidates were

selected using erythromycin and screened by PCR using primer pairs KMB_081/KMB_088,

KMB_081/HB8, and KMB_084/KMB_085. Insertion of the Erm-bar scar was confirmed by

Sanger sequencing using primers KMB_081, KMB_082, KMB_083, HB8, HB9, KMB_086,

KMB_087, KMB_088 and the barcode sequence recorded. The final bar-scar strain (MJM3903)

was constructed via a triparental mating with donor MJM3478 (π3813/pKV496) (50) and helper

strain MJM534 (CC118 λpir/pEVS104) with MJM3775. Candidates were selected for using

kanamycin and screened by PCR using the primer pairs listed above. The deletion scar was

verified by Sanger sequencing using primers KMB_081, KMB_082, KMB_083, HB41, HB42,

KMB_086, KMB_087 and KMB_088.

To create the Δ*binK* Δ*rscS* strain, a Δ*binK* strain with the pLostfoX-Kan plasmid

(MJM2252) was cultured for transformation (54). Donor DNA was 2.4 ug of genomic DNA from

the Δ*rscS*::*erm-bar* strain that was purified using the Qiagen Blood and Tissue Kit. Transformed

cells were plated onto LBS containing erythromycin to select for transformants. Isolates were

then patched onto LBS containing kanamycin to ensure loss of the pLostfoX-Kan plasmid. PCR

was performed to ensure the strain was transformed with Δ*rscS*::*erm-bar* and maintained Δ*binK*

(primers DAT_011F BinK ext, DAT_012R BinK ext, KMB_081, KMB_088). The *erm* cassette

was then removed using a FLP recombinase as described with the Δ*rscS::bar* strain above to

create the final Δ*binK* Δ*rscS*::*bar* strain (MJM4018).

Squid single strain colonizations

V. fischeri strains were grown overnight with aeration at 25°C in LBS. Overnight cultures were

diluted 1:80 in LBS and grown to an OD_{600} of approximately 0.3. The OD_{600} was used to

normalize the amount of each strain used to inoculate *E. scolopes* hatchlings at concentrations

of approximately 6×10^3 CFU/mL in 40 mL seawater for 3 hours. Squid were then washed and

transferred to individual vials with 4 mL of bacteria-free filter-sterilized Instant Ocean (FSIO)

until approximately 48 hours post-inoculation (hpi) with a water change that occurred at 24 hpi.

At 48 hpi, squid were transferred to 1.5 mL microfuge tubes with 750 mL of water and each

animal's luminescence was measured using the Promega GloMax 20/20 luminometer.

CFU counts per light organ were conducted as we described previously: euthanized

squid were homogenized and plated, and colonies were counted to determine CFU per light

organ (55).

For crypt visualization, squid were anesthetized in FSIO with 2% ethanol. At this point,

hatchlings were either immediately dissected and imaged, or fixed in 4% paraformaldehyde in

1X mPBS (50 mM phosphate buffer, 0.45 M NaCl, pH 7.4) for approximately 36 hours. Fixed

squid were thoroughly washed in 1X mPBS before being dissected and imaged. All images

were acquired on the Zeiss Axio Zoom.v16 large-field stereo microscope. The Zen Blue

software polygon tool was used to select regions of interest and measure fluorescence intensity.

Squid aggregation assays

V. fischeri strains were grown overnight with aeration at 25°C in LBS. 40 mL of overnight culture

was used to inoculate *E. scolopes* hatchlings at concentrations of approximately 5.5×10^6

CFU/mL in 40 mL for 3-4 hours. Hatchlings were then anesthetized in FSIO with 2% ethanol. At

this point, hatchlings were either immediately dissected and imaged, or fixed in 4%

paraformaldehyde in 1X mPBS (50 mM phosphate buffer, 0.45 M NaCl, pH 7.4) for

approximately 36 hours. Fixed squid were thoroughly washed in 1X mPBS before being

dissected and imaged. All images were acquired on the Zeiss Axio Zoom.v16 large-field

fluorescent stereo microscope. The Zen Blue software polygon tool was used to select regions

of interest and measure both area and fluorescence intensity

Squid competition assays

Strains were grown overnight with aeration at 25°C in LBS and LBS containing kanamycin to

maintain plasmid pVSV103. Strains with pVSV103 constitutively express LacZ (β-

galactosidase). Overnight cultures were diluted 1:80 in LBS and grown to an OD_{600} of

approximately 0.3. Using optical density to normalize the strains, the two strains were mixed in a

1:1 ratio. This mixed culture was used to inoculate *E. scolopes* hatchlings at concentrations of

approximately 7.6×10^3 bacteria for 3 hours. Squid were then washed and transferred to 40 ml

of bacteria-free filter-sterilized Instant Ocean (FSIO) until approximately 48 hours post-

inoculation (water was changed at 24 h post-inoculation), at which point they were euthanized

by storage at −80°C. Each squid was homogenized and plated on LBS-Xgal, and the blue/white

colony ratios were used to score these competitions as described previously (23, 55).

Data analysis and graphing

Data analysis was conducted using Python, including the pandas library. For fluorescence of

colonies, aggregates, and crypts, the mean GFP and mCherry for the region of interest and a

nearby background region was acquired using Zen Blue Software. The background for each

channel was subtracted from the region of interest. To normalize GFP to plasmid copy number,

GFP was divided by mCherry. This resulted in the reported mean GFP/mCherry reading for

each individual colony, aggregate, or crypt space. Graphpad Prism was used to construct

graphs and perform statistical analyses.

ACKNOWLEDGMENTS

We thank John F. Brooks II for contributing strains for this study and Karen L. Visick for helpful

comments on the manuscript. This work was funded by NIGMS grant R35 GM119627 to M.J.M.

Support for trainees was provided by NIGMS T32 GM008061 (D.A.L), T32 GM008349 (K.M.B.),

and an NSF Graduate Research Fellowship to K.M.B.

Table 1: Strains used in this study

Strain	Genotype	Source or Reference(s)
V. fischeri		
MJM1100 = ES114	Natural isolate, squid light-organ	(56, 57)
MJM1107	MJM1100 pVSV102	(23)
MJM1198	MJM1100 *rscS**	(32)
MJM1438	MJM1100 pM1422	JF Brooks II
MJM1538	MJM1100 pLostfoX-Kan	(23)
MJM1575	MJM1100 pVSV103	(22)
MJM2251	MJM1100 Δ*binK*	(22)
MJM2252	MJM2251 pLostfoX-Kan	(22)

MJM2255	MJM1100 *rscS* ΔbinK*	(22)
MJM2265	MJM2251 pVSV102	(22)
MJM2475	MJM1198 *att*Tn*7::binK*	(22)
MJM2476	MJM2255 *att*Tn*7::binK*	(22)
MJM2479	MJM1198 *att*Tn*7::erm*	(22)
MJM2480	MJM2255 *att*Tn*7::erm*	(22)
MJM2484	MJM2255 *att*Tn*7::binK*(H362Q)	JF Brooks II
MJM2493	MJM2251 pM1422	JF Brooks II
MJM2536 = KV3299	ES114 Δ*sypE sypF**	(18, 42)
MJM3210	MJM1198 *att*Tn*7::binK*(H362Q)	This study
MJM3213	MJM1198 *att*Tn*7::binK*(D794A)	This study
MJM3215	MJM2255 *att*Tn*7::binK*(D794A)	This study
MJM3218	MJM1198 *att*Tn*7::binK*(H362Q,D794A)	This study
MJM3220	MJM2255 *att*Tn*7::binK*(H362Q,D794A)	This study
MJM3221	MJM2251 pVSV103	This study
MJM3236	MJM2536 *sypG*(D53E)	This study
MJM3251	MJM1198 *att*Tn*7::binK*(D794E)	This study
MJM3252	MJM2255 *att*Tn*7::binK*(D794E)	This study
MJM3256	MJM1198 *att*Tn*7::binK*(E366A)	This study
MJM3257	MJM2255 Δ*binK att*Tn*7::binK*(E363A)	This study
MJM3775	MJM1100 Δ*rscS::erm-bar*	This study
MJM3903	MJM1100 Δ*rscS::bar*	This study
MJM4017	MJM1100 Δ*binK* Δ*rscS::erm-bar*	This study
MJM4018	MJM1100 Δ*binK* Δ*rscS::bar*	This study
MJM4071	MJM4018 pVSV102	This study
MJM4240	MJM3236 pVSV104	This study
MJM4241	MJM3236 pBinK	This study
MJM4242	KV3299 pVSV104	This study
MJM4243	KV3299 pBinK	This study
MJM4251	MJM1198 *att*Tn*7::binK*(H362Q,D794E)	This study
MJM4252	MJM2255 *att*Tn*7::binK*(H362Q,D794E)	This study

MJM4253	MJM1198 *att*Tn*7*::*binK*(T366Q)	This study
MJM4254	MJM2255 Δ*binK att*Tn*7*::*binK*(T366Q)	This study
MJM4255	MJM1198 *att*Tn*7*::*binK*(T366A)	This study
MJM4256	MJM2255 Δ*binK att*Tn*7*::*binK*(T366A)	This study
MJM4257	MJM3903 pM1422	This study
MJM4258	MJM4018 pM1422	This study
E. coli		
MJM534	CC118 λpir/pEVS104	(51)
MJM537	DH5α λpir	Laboratory Stock
MJM542	DH5α λpir/pVSV102	(58)
MJM570	DH5α/pEVS79	(51)
MJM580	DH5α λpir/pVSV104	(58)
MJM637	S17-1 λpir/pUX-BF13	(59)
MJM658	DH5α λpir/pEVS107	(60)
MJM1422	DH5α λpir/pM1422	(22)
MJM2090	DH5α λpir/pLostfoX-Kan	(23)
MJM2384	DH5α λpir/pBinK	(22)
MJM2474	DH5α λpir/pTn7BinK	(22)
MJM2482	DH5α λpir/pTn7BinK(H362Q)	JF Brooks II
MJM3211	DH5α λpir/pDAT01	This study
MJM3216	DH5α λpir/pDAT02	This study
MJM3234	DH5α λpir/pDAT05	This study
MJM3241	DH5α λpir/pDAT10	This study
MJM3255	DH5α λpir/pDAT12	This study
MJM3478	π3813/pKV496	(50)
MJM4248	DH5α λpir/pDAT13	This study
MJM4249	DH5α λpir/pDAT14	This study
MJM4250	DH5α λpir/pDAT15	This study

596 **Table 2: Plasmid List**

Plasmid	Description	Source or Reference(s)
pEVS107	Mini-Tn7 mobilizable vector (Kan^R,Erm^R)	(60)
pTn7BinK	pEVS107 carrying wild-type *binK* (Kan^R,Erm^R)	(22)
pTn7BinK(H362Q)	pEVS107 carrying *binK* with H362Q mutation (Kan^R,Erm^R)	This study
pDAT01	pEVS107 carrying *binK* with D794A mutation (Kan^R,Erm^R)	This study
pDAT02	pEVS107 carrying *binK* with H362Q,D794A mutation (Kan^R,Erm^R)	This study
pDAT10	pEVS107 carrying *binK* with D794E mutation (Kan^R,Erm^R)	This study
pDAT13	pEVS107 carrying *binK* with H362Q,D794E mutation (Kan^R,Erm^R)	This study
pDAT12	pEVS107 carrying *binK* with E363A mutation (Kan^R,Erm^R)	This study
pDAT14	pEVS107 carrying *binK* with T366Q mutation (Kan^R,Erm^R)	This study
pDAT15	pEVS107 carrying *binK* with T366A mutation (Kan^R,Erm^R)	This study
pEVS104	Conjugal helper plasmid (Kan^R)	(51)
pUX-BF13	Tn7 transposition helper (Amp^R)	(59)
pVSV104	Vector backbone for complementation (Kan^R)	(58)
pBinK	pVSV104 carrying wild-type *binK* (Kan^R)	(22)
pEVS79	Vector backbone for allelic exchange (Cam^R)	(51)
pDAT05	pEVS79 carrying *sypG* with D53E mutation (Cam^R)	This study
pM1422	pTM267 sypA'-gfp+ (Cam^R)	(22)
pVSV102	Constitutive GFP (Kan^R)	(58)
pVSV103	Constitutive LacZ (Kan^R)	(58)
pLostfoX-Kan	Arabinose-inducible TfoX for transformation (Kan^R)	(23)
pKV496	pEVS79 containing the FLP recombinase (Kan^R)	(50)

601   **Table 3: Primer List**

Primer Name	Sequence (5'-3')	Notes
DAT_011F BinK ext	CTTATGGGATAGCTTTAGCTCGAA	External primer to sequence around BinK gene
DAT_012R BinK ext	TTCTTTATTAAGTGAAGGCGAACC	External primer to sequence around BinK gene
DAT_018F tfox	TAAATAAATCCTGGTGTCCCTGTT	Screen for pLostfoX-Kan
DAT_019R tfox	TCGCTGTTAAAAGGACAATTACAA	Screen for pLostfoX-Kan
DAT_064 D794A F	AGTTTTAATGGcgTGTATGATGCCGATAATG	Site directed mutagenesis to make binK(D794A), lowercase letters represent mutated nucleotides
DAT_065 D794A R	AAACTATAAGGTGACGTTTTAC	Site directed mutagenesis to make binK(D794A)
DAT_066 D794A Fwd	GAATCGTCCAAAAATTTGTTCCGC	Screen for D794A mutation
DAT_067 D794A RevM	CATCCATTATCGGCATCATACAcg	Screen for D794A mutation, lowercase letters represent nucleotides specific to mutant
DAT_068 D794A RevWT	CATCCATTATCGGCATCATACAgt	Screen for D794A mutation, lowercase letters represent nucleotides specific to wild-type
DAT_071 sypG_US F	ccggtcgacggtatcgataaGATATTCTCGACTTCTCACGTATG	sypG-D53E cloning, capital letters indicate homology to template
DAT_072 sypG_US R	tcgagaatcaccaaatgTGGTGGATTCTTTTCCATAAATGC	sypG-D53E cloning, capital letters indicate homology to template
DAT_074 sypG_DS R	gcaggaattcgatatcaagcCTGGAATATAATGCCGCTTTGTAG	sypG-D53E cloning, capital letters indicate homology to template
DAT_077 sypG_DS F	CATTTGGTGATTCTCGAaTTGAAACTGCC	sypG-D53E cloning, lowercase letter represents mutated nucleotide
DAT_078 sypG seq1	CATGAAGGTTCTGGTTTAGGG	pEVS79 sypG-D53E sequencing
DAT_079 sypG seq2	CCGCTCTCTATCTGATAC	pEVS79 sypG-D53E sequencing
DAT_080 sypG seq3	GCATTAGAGTTTGAAGCG	pEVS79 sypG-D53E sequencing
DAT_081 sypG seq4	GATATGTCAGGGCAGGATG	pEVS79 sypG-D53E sequencing
DAT_082 sypG seq5	GTAAACGCAGCAATAAACC	pEVS79 sypG-D53E sequencing
DAT_083 sypG	CGCAAAAGAAGATCATAA	pEVS79 sypG-D53E sequencing

seq6		
DAT_084 sypG seq7	GATGGTAATGTTCTTAATG	pEVS79 sypG-D53E sequencing
DAT_085 sypG seq8	CCATTGGGCGAATTACAC	pEVS79 sypG-D53E sequencing
DAT_086 sypG seq9	GAATAGGCACAACATGGAC	pEVS79 sypG-D53E sequencing
DAT_087 sypG seq10	CCTGTACCGCTTTCACCA	pEVS79 sypG-D53E sequencing
DAT_088 sypG seq11	CATAAATGCCTTCGCTTG	pEVS79 sypG-D53E sequencing
DAT_089 sypG seq12	GATGGTTTCTACTCGCTC	pEVS79 sypG-D53E sequencing
DAT_090 sypG seq13	CATCCATTTCAGGCATAG	pEVS79 sypG-D53E sequencing
DAT_091 sypG seq14	GCTGTTATCTGCCATAGT	pEVS79 sypG-D53E sequencing
DAT_092 D53E Fwd	GAGTTTAATGAATCAACAGAAGAGC	sypG-D53E screen
DAT_093 D53E RevMT	CCCTGACATATCTGGCAGTTTCAAt	sypG-D53E screen, lowercase letters represent nucleotides specific to mutant
DAT_094 D53E RevWT	CCCTGACATATCTGGCAGTTTCAAa	sypG-D53E screen, lowercase letters represent nucleotides specific to wild-type
DAT_095 sypG fwd	CTACAGCAAGCCAGAAATGAAGCAG	Amplify around *sypG* insert area on pEVS79
DAT_096 sypG rev	GGGTGCCTTTTGATTGAATTAAGTTC	Amplify around *sypG* insert area on pEVS79
DAT_106 D794E F	TTTTAATGGAgTGTATGATGCC	Site directed mutagenesis to make binK(D794E), lowercase letters represent mutated nucleotides
DAT_107 D794E R	CTAAACTATAAGGTGACGTTTTAC	Site directed mutagenesis to make binK(D794E)
DAT_109 E363A_F	ATGTCACACGctATTCGAACAC	Site directed mutagenesis to make binK(E363A), lowercase letters represent mutated nucleotides
DAT_110 E363_R	ATTAGCTAAAAATGAACTTTTGG	Site directed mutagenesis to make binK(E363A)
DAT_111 T366A_F	CGAAATTCGAgcACCTCTAAATG	Site directed mutagenesis to make binK(T366A), lowercase letters represent mutated nucleotides

DAT_112 T366_R	TGTGACATATTAGCTAAAAATG	Site directed mutagenesis to make binK T366 mutations
DAT_114 T366Q_F	CGAAATTCGAcaACCTCTAAATGGCATC	Site directed mutagenesis to make binK(T366Q), lowercase letters represent mutated nucleotides
DAT_118 D794E_RevWT	CATCCATTATCGGCATCATACAg	Screen for D794E mutation, lowercase letters represent nucleotides specific to wild-type
DAT_119 D794E_RevM	CATCCATTATCGGCATCATACAc	Screen for D794E mutation, lowercase letters represent nucleotides specific to mutant
DAT_120 E363A_RevM	CCATTTAGAGGTGTTCGAATag	Screen for E363A mutation, lowercase letters represent nucleotides specific to mutant
DAT_121 E363A_RevWT	CCATTTAGAGGTGTTCGAATtt	Screen for E363A mutation, lowercase letters represent nucleotides specific to wild-type
DAT_122 E363A_Fwd	GGTTGATCGGTGTTATTGAATC	Screen for E363A mutation
DAT_257 T336 RevWT	CATACCAATGATGCCATTTAGAGGTgt	Screen for T366 mutations, lowercase letters represent nucleotides specific to wild-type
DAT_259 T366 RevQ	CATACCAATGATGCCATTTAGAGGTtg	Screen for T366 mutations, lowercase letters represent nucleotides specific to T366Q mutant
DAT_260 T366 RevA	CATACCAATGATGCCATTTAGAGGTgc	Screen for T366 mutations, lowercase letters represent nucleotides specific to T366A mutant
DAT_261 T366 Fwd	CGCTGTTGATGAAAGTATGTATGTTGTAG	Screen for T366 mutations
Tn7 site F	TGTTGATGATACCATTGAAGCTAAA	Amplify around *att*Tn*7* site
Tn7 site R	CTTGCTGTATGTATTTGCTGATGA	Amplify around *att*Tn*7* site
pEVS107 F	ACCTATCAAGGTGTACTGCCTTCC	Sequences around pEVS107 multiple cloning site
pEVS107 R	GTCGTTAAATGCCCTTTACCTGT	Sequences around pEVS107 multiple cloning site
MRH001	TTATCGATACCGTCGACC	Linearizes pEVS79 for cloning
MRH002	GCTTGATATCGAATTCCTG	Linearizes pEVS79 for cloning
M13 Rev (-48)	AGCGGATAACAATTTCACACAGG	Amplifies around plasmid insertions
M13 For (-41)	CGCCAGGGTTTTCCCAGTCACGAC	Amplifies around plasmid insertions

JFB_359	AAATGATAATCGCTGGTC	Sequencing of *binK*
JFB_361	GATGTTCATCAAGCATTA	Sequencing of *binK*
JFB_362	GAGGTGTTCGAATTTCGT	Sequencing of *binK*
JFB_363	GAGCGAAAGTCTCATCAG	Sequencing of *binK*
JFB_364	AAACCTCAGACCATGAAA	Sequencing of *binK*
JFB_365	GGAAAGAGAATGATTAAG	Sequencing of *binK*
JFB_366	ATTCAAAGAATATGGTGC	Sequencing of *binK*
JFB_371	CTATTTTATTGGCTTGTG	Sequencing of *binK*
JFB_372	AACTGAAACCGATTTAAC	Sequencing of *binK*
JFB_373	ATGCCGTTAAATTTACTC	Sequencing of *binK*
JFB_374	TTGAGGTGATTGAGCCAA	Sequencing of *binK*
JFB_375	TTGAACGTACAATTGAAG	Sequencing of *binK*
JFB_376	TAGATATGGTGATGAGTA	Sequencing of *binK*
JFB_377	ACTGAATTACGTTTAACG	Sequencing of *binK*
JFB_426	GCTAATATGTCACAAGAAATTCGAACACC	Site-directed mutagenesis to make pTn7BinK (H362Q)
JFB_427	GGTGTTCGAATTTCTTGTGACATATTAGC	Site-directed mutagenesis to make pTn7BinK (H362Q)
KMB_081	GTGTGATGCAGATATAAAAAATCCCTGATCTTAATC	F0 for ES114 *rscS* deletion
KMB_082	CGAACTTCCCCACCAGCTAAAATC	F1 for ES114 *rscS* deletion
KMB_083	CTGGCGAAGCATATATAAGAAGCTCGTCTCGTCATTGCATTAGCTCCTATAAAATAGTCTGTTTGG	R1-LL for ES114 *rscS* deletion
KMB_084	CACTACATTGCCGTAGAAAGAGACATCAC	FW for ES114 *rscS* deletion
KMB_085	CAGCTTGTTTACCTTTACCTGTTAGAGTATGG	RW for ES114 *rscS* deletion
KMB_086	GACTTGACCTGGATGTCTCTACCCACAAGATCGGAGAAGTATGAAACACAATAAACTTCGTCATAAAAAAGG	F2-RL for ES114 *rscS* deletion
KMB_087	CGTACTATGGTGTTTAATATCAATACACTTCAATGGG	R2 for ES114 *rscS* deletion
KMB_088	CTCTCGTGCTGTCATTCTTGACCAC	R0 for ES114 *rscS* deletion
HB8	ACAAAATTTTAAGATACTGCACTATCAACACACTCTTAAG	Sequencing of ES114 Δ*rscS*

HB9	GGGAGGAAATAAtCTAGAATGCGAGAGTAGG	Sequencing of ES114 ΔrscS
HB41	CGATCTTGTGGGTAGAGACATCCAGGTCAAGTCCAGCCCCGCTCTAGTTTGGGAATCAAGTGCATGAGCGCTGAAG	Sequencing of ES114 ΔrscS
HB42	ACGAGACGAGCTTCTTATATATGCTTCGCCAG	Sequencing of ES114 ΔrscS

REFERENCES

1.   Ruby E, Henderson B, McFall-Ngai M. 2004. Microbiology. We get by with a little help from
our (little) friends. Science 303:1305–1307.

2.   McFall-Ngai MJ. 2014. The importance of microbes in animal development: lessons from
the squid-Vibrio symbiosis. Annu Rev Microbiol 68:177–194.

3.   Nyholm SV, Stabb EV, Ruby EG, McFall-Ngai MJ. 2000. Establishment of an animal-
bacterial association: recruiting symbiotic vibrios from the environment. Proc Natl Acad Sci
U S A 97:10231–10235.

4.   Mandel MJ, Dunn AK. 2016. Impact and influence of the natural *Vibrio*-squid symbiosis in
understanding bacterial-animal interactions. Front Microbiol 7:1982.

5.   Visick KL, Ruby EG. 2006. *Vibrio fischeri* and its host: it takes two to tango. Curr Opin
Microbiol 9:632–638.

6.   Ruby EG. 1996. Lessons from a cooperative, bacterial-animal association: the *Vibrio*
fischeri-Euprymna scolopes light organ symbiosis. Annu Rev Microbiol 50:591–624.

7.   Ruby EG, McFall-Ngai MJ. 1992. A squid that glows in the night: development of an animal-
bacterial mutualism. J Bacteriol 174:4865–4870.

8.   Koch EJ, Miyashiro T, McFall-Ngai MJ, Ruby EG. 2014. Features governing symbiont
persistence in the squid-vibrio association. Mol Ecol 23:1624–1634.

9.   Nyholm SV, McFall-Ngai MJ. 2004. The winnowing: establishing the squid-*Vibrio* symbiosis.
Nat Rev Microbiol 2:632–642.

10. Yip ES, Geszvain K, DeLoney-Marino CR, Visick KL. 2006. The symbiosis regulator *rscS*

controls the *syp* gene locus, biofilm formation and symbiotic aggregation by *Vibrio fischeri*.

Mol Microbiol 62:1586–1600.

11. Yip ES, Grublesky BT, Hussa EA, Visick KL. 2005. A novel, conserved cluster of genes

promotes symbiotic colonization and σ^{54}-dependent biofilm formation by *Vibrio fischeri*. Mol

Microbiol 57:1485–1498.

12. Visick KL. 2009. An intricate network of regulators controls biofilm formation and

colonization by *Vibrio fischeri*. Mol Microbiol 74:782–789.

13. Mandel MJ, Wollenberg MS, Stabb EV, Visick KL, Ruby EG. 2009. A single regulatory gene

is sufficient to alter bacterial host range. Nature 458:215–218.

14. Pankey SM, Foxall RL, Ster IM, Perry LA, Schuster BM, Donner RA, Coyle M, Cooper VS,

Whistler CA. 2017. Host-selected mutations converging on a global regulator drive an

adaptive leap towards symbiosis in bacteria. Elife 6:e24414.

15. Rotman ER, Bultman KM, Brooks JF 2nd, Gyllborg MC, Burgos HL, Wollenberg MS,

Mandel MJ. 2019. Natural strain variation reveals diverse biofilm regulation in squid-

colonizing *Vibrio fischeri*. J Bacteriol 201:e00033–19.

16. Geszvain K, Visick KL. 2008. The hybrid sensor kinase RscS integrates positive and

negative signals to modulate biofilm formation in *Vibrio fischeri*. J Bacteriol 190:4437–4446.

17. Norsworthy AN, Visick KL. 2015. Signaling between two interacting sensor kinases

promotes biofilms and colonization by a bacterial symbiont. Mol Microbiol 96:233–248.

18. Hussa EA, Darnell CL, Visick KL. 2008. RscS functions upstream of SypG to control the

syp locus and biofilm formation in *Vibrio fischeri*. J Bacteriol 190:4576–4583.

19. Darnell CL, Hussa EA, Visick KL. 2008. The putative hybrid sensor kinase SypF

coordinates biofilm formation in *Vibrio fischeri* by acting upstream of two response

regulators, SypG and VpsR. J Bacteriol 190:4941–4950.

20. Ray VA, Eddy JL, Hussa EA, Misale M, Visick KL. 2013. The *syp* enhancer sequence plays

a key role in transcriptional activation by the σ^{54}-dependent response regulator SypG and in

biofilm formation and host colonization by *Vibrio fischeri*. J Bacteriol 195:5402–5412.

21. Morris AR, Visick KL. 2013. The response regulator SypE controls biofilm formation and

colonization through phosphorylation of the *syp*-encoded regulator SypA in *Vibrio fischeri*.

Mol Microbiol 87:509–525.

22. Brooks JF 2nd, Mandel MJ. 2016. The histidine kinase BinK Is a negative regulator of

biofilm formation and squid colonization. J Bacteriol 198:2596–2607.

23. Brooks JF 2nd, Gyllborg MC, Cronin DC, Quillin SJ, Mallama CA, Foxall R, Whistler C,

Goodman AL, Mandel MJ. 2014. Global discovery of colonization determinants in the squid

symbiont *Vibrio fischeri*. Proc Natl Acad Sci U S A 111:17284–17289.

24. Tischler AH, Lie L, Thompson CM, Visick KL. 2018. Discovery of calcium as a biofilm-

promoting signal for *Vibrio fischeri* reveals new phenotypes and underlying regulatory

complexity. J Bacteriol 200:e00016–18.

25. Thompson CM, Tischler AH, Tarnowski DA, Mandel MJ, Visick KL. 2019. Nitric oxide

inhibits biofilm formation by *Vibrio fischeri* via the nitric oxide sensor HnoX. Mol Microbiol

111:187–203.

26. Marchler-Bauer A, Bryant SH. 2004. CD-Search: protein domain annotations on the fly.

Nucleic Acids Res 32:W327–31.

27. Blum M, Chang H-Y, Chuguransky S, Grego T, Kandasaamy S, Mitchell A, Nuka G,

Paysan-Lafosse T, Qureshi M, Raj S, Richardson L, Salazar GA, Williams L, Bork P, Bridge A, Gough J, Haft DH, Letunic I, Marchler-Bauer A, Mi H, Natale DA, Necci M, Orengo CA, Pandurangan AP, Rivoire C, Sigrist CJA, Sillitoe I, Thanki N, Thomas PD, Tosatto SCE, Wu CH, Bateman A, Finn RD. 2021. The InterPro protein families and domains database: 20 years on. Nucleic Acids Res 49:D344–D354.

28. Letunic I, Khedkar S, Bork P. 2021. SMART: recent updates, new developments and status in 2020. Nucleic Acids Res 49:D458–D460.

29. Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, Tosatto SCE, Paladin L, Raj S, Richardson LJ, Finn RD, Bateman A. 2021. Pfam: The protein families database in 2021. Nucleic Acids Res 49:D412–D419.

30. Wise AA, Fang F, Lin Y-H, He F, Lynn DG, Binns AN. 2010. The receiver domain of hybrid histidine kinase VirA: an enhancing factor for *vir* gene expression in *Agrobacterium tumefaciens*. J Bacteriol 192:1534–1542.

31. Wise AA, Binns AN. 2015. The receiver of the *Agrobacterium tumefaciens* VirA histidine kinase forms a stable interaction with VirG to activate virulence gene expression. Front Microbiol 6:1546.

32. Singh P, Brooks JF 2nd, Ray VA, Mandel MJ, Visick KL. 2015. CysK plays a role in biofilm formation and colonization by *Vibrio fischeri*. Appl Environ Microbiol 81:5223–5234.

33. Freeman JA, Lilley BN, Bassler BL. 2000. A genetic analysis of the functions of LuxN: a two-component hybrid sensor kinase that regulates quorum sensing in *Vibrio harveyi*. Mol Microbiol 35:139–149.

34. Gao R, Stock AM. 2009. Biological insights from structures of two-component proteins. Annu Rev Microbiol 63:133–154.

35. Bhate MP, Molnar KS, Goulian M, DeGrado WF. 2015. Signal transduction in histidine

kinases: insights from new structures. Structure 23:981–994.

36. Atkinson MR, Ninfa AJ. 1993. Mutational analysis of the bacterial signal-transducing protein

kinase/phosphatase nitrogen regulator II (NRII or NtrB). J Bacteriol 175:7016–7023.

37. Willett JW, Kirby JR. 2012. Genetic and biochemical dissection of a HisKA domain

identifies residues required exclusively for kinase and phosphatase activities. PLoS Genet

8:e1003084.

38. Huynh TN, Noriega CE, Stewart V. 2010. Conserved mechanism for sensor phosphatase

control of two-component signaling revealed in the nitrate sensor NarX. Proc Natl Acad Sci

U S A 107:21140–21145.

39. Huynh TN, Stewart V. 2011. Negative control in two-component signal transduction by

transmitter phosphatase activity. Mol Microbiol 82:275–286.

40. Dutta R, Yoshida T, Inouye M. 2000. The critical role of the conserved Thr247 residue in

the functioning of the osmosensor EnvZ, a histidine kinase/phosphatase, in *Escherichia*

coli. J Biol Chem 275:38645–38653.

41. Wayne KJ, Li S, Kazmierczak KM, Tsui H-CT, Winkler ME. 2012. Involvement of WalK

(VicK) phosphatase activity in setting WalR (VicR) response regulator phosphorylation level

and limiting cross-talk in *Streptococcus pneumoniae* D39 cells. Mol Microbiol 86:645–660.

42. Thompson CM, Marsden AE, Tischler AH, Koo J, Visick KL. 2018. *Vibrio fischeri* biofilm

formation prevented by a trio of regulators. Appl Environ Microbiol 84:e01257–18.

43. Visick KL, Skoufos LM. 2001. Two-component sensor required for normal symbiotic

colonization of *Euprymna scolopes* by *Vibrio fischeri*. J Bacteriol 183:835–842.

44. Kinoshita-Kikuta E, Kinoshita E, Eguchi Y, Koike T. 2016. Validation of *cis* and *trans* modes

in multistep phosphotransfer signaling of bacterial tripartite sensor kinases by using Phos-

Tag SDS-PAGE. PLoS One 11:e0148294.

45. Teran-Melo JL, Peña-Sandoval GR, Silva-Jimenez H, Rodriguez C, Alvarez AF, Georgellis

D. 2018. Routes of phosphoryl group transfer during signal transmission and signal decay

in the dimeric sensor histidine kinase ArcB. J Biol Chem 293:13214–13223.

46. Jacob-Dubuisson F, Mechaly A, Betton J-M, Antoine R. 2018. Structural insights into the

signalling mechanisms of two-component systems. Nat Rev Microbiol 16:585–593.

47. Chang CH, Zhu J, Winans SC. 1996. Pleiotropic phenotypes caused by genetic ablation of

the receiver module of the *Agrobacterium tumefaciens* VirA protein. J Bacteriol 178:4710–

4716.

48. Altura MA, Heath-Heckman EAC, Gillette A, Kremer N, Krachler A-M, Brennan C, Ruby

EG, Orth K, McFall-Ngai MJ. 2013. The first engagement of partners in the *Euprymna*

scolopes-Vibrio fischeri symbiosis is a two-step process initiated by a few environmental

symbiont cells. Environ Microbiol 15:2937–2950.

49. Le Roux F, Binesse J, Saulnier D, Mazel D. 2007. Construction of a *Vibrio splendidus*

mutant lacking the metalloprotease gene *vsm* by use of a novel counterselectable suicide

vector. Appl Environ Microbiol 73:777–784.

50. Visick KL, Hodge-Hanson KM, Tischler AH, Bennett AK, Mastrodomenico V. 2018. Tools

for Rapid Genetic Engineering of *Vibrio fischeri*. Appl Environ Microbiol 84:e00850–18.

51. Stabb EV, Ruby EG. 2002. RP4-based plasmids for conjugation between *Escherichia coli*

and members of the *Vibrionaceae*. Methods Enzymol 358:413–426.

52. Burgos HL, Burgos EF, Steinberger AJ, Suen G, Mandel MJ. 2020. Multiplexed competition

in a synthetic squid light organ microbiome using barcode-tagged gene deletions.

mSystems 5:e00846–20.

53. Brooks JF 2nd, Gyllborg MC, Kocher AA, Markey LEH, Mandel MJ. 2015. TfoX-based

genetic mapping identifies *Vibrio fischeri* strain-level differences and reveals a common

lineage of laboratory strains. J Bacteriol 197:1065–1074.

54. Pollack-Berti A, Wollenberg MS, Ruby EG. 2010. Natural transformation of *Vibrio fischeri*

requires *tfoX* and *tfoY*. Environ Microbiol 12:2302–2311.

55. Naughton LM, Mandel MJ. 2012. Colonization of *Euprymna scolopes* squid by *Vibrio*

fischeri. J Vis Exp e3758.

56. Boettcher KJ, Ruby EG. 1990. Depressed light emission by symbiotic *Vibrio fischeri* of the

sepiolid squid *Euprymna scolopes*. J Bacteriol 172:3701–3706.

57. Mandel MJ, Stabb EV, Ruby EG. 2008. Comparative genomics-based investigation of

resequencing targets in *Vibrio fischeri*: focus on point miscalls and artefactual expansions.

BMC Genomics 9:138.

58. Dunn AK, Millikan DS, Adin DM, Bose JL, Stabb EV. 2006. New *rfp-* and pES213-derived

tools for analyzing symbiotic *Vibrio fischeri* reveal patterns of infection and *lux* expression *in*

situ. Appl Environ Microbiol 72:802–810.

59. Bao Y, Lies DP, Fu H, Roberts GP. 1991. An improved *Tn7*-based system for the single-

copy insertion of cloned genes into chromosomes of gram-negative bacteria. Gene

109:167–168.

60. McCann J, Stabb EV, Millikan DS, Ruby EG. 2003. Population dynamics of *Vibrio fischeri*

during infection of *Euprymna scolopes*. Appl Environ Microbiol 69:5928–5934.

61. Madeira F, Park YM, Lee J, Buso N, Gur T, Madhusoodanan N, Basutkar P, Tivey ARN,

Potter SC, Finn RD, Lopez R. 2019. The EMBL-EBI search and sequence analysis tools

APIs in 2019. Nucleic Acids Res 47:W636–W641.

FIGURE LEGENDS

Figure 1: BinK signaling and domain organization

A) Model of the biofilm pathway in strain ES114 as is relevant for this study. RscS signals

through SypF to SypG, which is a σ^{54}-dependent transcriptional activator of the *syp* locus.

Expression of the *syp* locus is required for biofilm formation. BinK inhibits biofilm production and

feeds into the pathway at an unknown location at or above SypG. (OM, outer membrane; IM,

inner membrane). **B)** Putative subcellular localization of a BinK homodimer **C)** BinK domain

diagram. Positions of key residues and the epitope used to generate the BinK peptide antibody

(722-742) are annotated. (TM, predicted transmembrane domains).

Figure 2: BinK requires H362 and D794 to inhibit colony biofilms.

A) Genome representations of the different strains used to assess *binK* alleles. The parent

ES114 *rscS** strain (MJM1198) was used to induce biofilm formation for wrinkled colony assays

on plates. The *att*Tn*7* site is located on chromosome I and the native locus of *binK* is on

chromosome II. *mut* designates a mutant allele of *binK,* while Δ indicates a clean deletion at that

locus. **B)** Wrinkled colony assay of the strains indicated, grown at 28°C for 48 hours. Mutations

(or WT control) expressed at the *att*Tn*7* site are listed beside each spot. In the left column, the

expressed allele is the only *binK* allele in the cell, while in the right column wild-type *binK* is

additionally present at its native locus. Scale bar is 2 mm. **C)** Western blot of whole cell lysates

assessed with a peptide antibody against BinK. Arrow indicates BinK, which is predicted to be

97 kDa (61), and which is absent in the strain that does not encode the protein.

Figure 3: RscS is not required for squid colonization in a strain lacking BinK

Bacteria were inoculated into FSIO containing squid and allowed to colonize for 3 hours. Squid

were washed and then maintained for two days to allow establishment of the symbiosis. Shown

are data on the luminescence (A) and CFU (B) per light organ (LO) at 48 hpi.Each dot

represents an individual animal. The dashed line indicates the limit of detection for CFU/LO. For

both graphs, data is pooled from three replicate experiments. Bars represent the median for

each strain with a 95% confidence interval. Statistical significance was calculated using Kruskal-

Wallis test with Dunn's multiple comparison tests. (ns, not significant; ****, $p < 0.0001$)

Figure 4: RscS does not impact aggregation in a strain lacking BinK

Imaging of bacterial aggregates in host mucus using fluorescence microscopy with the Zeiss

Axio Zoom fluorescent microscope. Squid were inoculated with *V. fischeri* cells that

constitutively express GFP from the pVSV102 plasmid and imaged at 3-4 hpi. **A)**

Representative images of aggregates of approximately median size for each strain. Scale bar is

200 µm with all panels at the same scale. **B)** Quantification of aggregate area. Each dot

represents one aggregate. A measure of zero indicates no aggregate was present. Significance

was determined with a Kruskal-Wallis test and Dunn's multiple comparison tests (ns, not

significant; **, $p < 0.01$; ***, $p < 0.001$). The median area with a 95% confidence interval is

displayed for each group. The data is pooled from two replicate experiments. WT = wild-type

ES114 (MJM1100).

Figure 5: RscS is required for competitive fitness in a Δ*binK* background. Competitive

fitness of the indicated strains (unlabeled) compared to a Δ*binK* strain (labeled with LacZ).

Squid were exposed to a mixed inoculum of the two strains for 3 hours, then assessed at 48 hpi.

Blue vs white CFU counts on LBS-Xgal were used to determine the representation of each

strain in the competition. The competitive index is equal to the Log_{10}-transformed value of the

ratio (indicated strain/Δ*binK* strain) after competition normalized to its ratio in the input inoculum.

Each point represents the competitive index from an individual squid. The median ratio with a

95% confidence interval is represented by the bar. Statistical significance was calculated with a

Mann-Whitney U-test (****, p<0.0001).

Figure 6: RscS does not alter *syp* production on agar in strains lacking BinK

syp'-gfp[+] transcriptional reporter activity normalized to constitutive mCherry from strains carrying

pM1422. **A, B)** Each strain was spotted onto LBS or LBS-Calcium and grown for 48 hours at

25^{o}C. Images were taken on both a Zeiss Axio Zoom fluorescent microscope and Leica

brightfield stereoscope. Scale bar in the bottom right of B is 2mm and all panels in A and B are

at the same scale. **C)** Quantification of the fluorescence intensity. GFP intensity of the entire

colony was normalized to mCherry intensity. Each bar represents the average GFP/mCherry of

five different colonies from replicate experiments on two distinct days. Error bars are standard

deviations. Statistical significance was determined with a two-way ANOVA analyzing the effect

of strain background and media on fluorescence with Tukey's test for multiple comparisons. (ns,

not significant; *, p<0.05; ****, p<0.0001)

Figure 7: BinK inhibits *sypA* transcription in the crypts

Imaging of *V. fischeri* containing the pM1422 *sypA'-gfp*[+] reporter plasmid (that encodes

constitutive mCherry) while aggregating in host mucus at 3-4 hpi (A, quantified in C) or in host

crypts at 48 hpi (B, quantified in D). For quantification, intensity of GFP and mCherry was

measured for each individual aggregate or crypt, background signal was subtracted, and GFP

was normalized to the mCherry level. Zen Blue software was used to collect signal and

background measurements. Each dot represents an individual aggregate or crypt. Bars

represent the median GFP/mCherry ratio with 95% confidence intervals. Statistical significance

within each location was determined by a Kruskal-Wallis test with Dunn's multiple comparison

tests (ns, not significant; **, p<0.01). Data for each bacterial location is pooled from at least 3

replicate experiments that each contained 4-8 squid per strain.

A

	Native Locus	*att*Tn7 Site	RscS
Parent	*binK⁺*	▪	*rscS**
Δ*binK*	Δ	▪	*rscS**
Δ*binK* *att*Tn7::*binK^mut*	Δ	*binK^mut*	*rscS**
binK⁺ *att*Tn7::*binK^mut*	*binK⁺*	*binK^mut*	*rscS**

B

binK expressed from *att*Tn7 site	Native locus Δ*binK*	*binK⁺*
(empty)		
WT		
H362Q		
D794A		
H362Q D794A		
D794E		
H362Q D794E		

C

rscS* Δ*binK* *att*Tn7::

(empty)   WT   H362Q   D794A   H362Q D794A   D794E   H362Q D794E   *rscS* binK⁺*

kDa
100
70
55
40

A Aggregates at 3-4 hpi

	Overlay	GFP	mCherry
WT			
Δ*binK*			
Δ*binK* Δ*rscS*			
Apo			

100 μm

B In the crypts at 48 hpi

	Overlay	GFP	mCherry
WT			
Δ*binK*			
Δ*binK* Δ*rscS*			
Apo			

100 μm

C

GFP/mCherry

ns
ns   ns

WT   Δ*binK*   Δ*binK* Δ*rscS*

D

GFP/mCherry

**   **
ns

WT   Δ*binK*   Δ*binK* Δ*rscS*